


The Rendering Equation

$$L_o = L_e + \int_{\Omega} f_r L_i \cos \theta d\omega$$

FUNDAMENTALS OF COMPUTER GRAPHICS
PHILIP DUTRÉ
DEPARTMENT OF COMPUTER SCIENCE

Overview Lecture

Radiometric accurate rendering



Light Energy

Counting number of "photons" incident or extant on a surface


Symbol: Q

Unit: Joule

Measurable with detector

Wavelength dependent

• Different counts for different "colors" of light.



Why is radiance important?

Radiance is invariant along a straight line

→ we can trace light energy along rays






Figure 15.17 Radiance along a Ray through Free Space is Invariant. To derive, we compare the radiant surface along a ray from point p in direction ω, we see that the ray within the hemisphere and parallel radiant surface in the direction ω. Then, the trace equals the radiant surface on the far surface that the ray goes through.

BRDF – Bidirectional Reflectance Distribution Function



$$f_r(p, \omega_i, \omega_o) = \frac{dL_o(p, \omega_o)}{L_i(p, \omega_i) \cos \theta_i d\omega_i}$$

Rendering Equation



$$f_r(p, \omega_i, \omega_o) = \frac{dL_o(p, \omega_o)}{L_i(p, \omega_i) \cos \theta_i d\omega_i}$$

$$dL_o(p, \omega_o) = f_r(p, \omega_i, \omega_o) L_i(p, \omega_i) \cos \theta_i d\omega_i$$

$$L_o(p, \omega_o) = L_e(p, \omega_o) + \int_{\text{hemisphere } \Omega} f_r(p, \omega_i, \omega_o) L_i(p, \omega_i) \cos \theta_i d\omega_i$$


Simple Stochastic Ray Tracing (aka Pure Path Tracing, Naïve Path Tracing, ...)

$$L_o(p, \omega_o) = ?$$

$$L_e(p, \omega_o)$$

+


$$\int_{\Omega} f_r(p, \omega_i, \omega_o) L_i(p, \omega_i) \cos \theta_i d\omega_i$$




Relevant sections in book: Chapter 13

(Illustrations from *Ray Tracing From The Ground Up, Physically-Based Rendering, Fundamentals of Computer Graphics*)
(Page numbering might skip some slides due to 'hidden' slides in my presentation.)

Radiometric accurate rendering




Visual comparison



Radiometric / Perceptual match (Meyer1986)

Visual comparison



Light Energy

Counting number of “photons” incident or exitant on a surface



Symbol: **Q**

Unit: **Joule**

Measurable with detector

Wavelength dependent

- Different counts for different ‘colors’ of light



Power




Energy per unit time (dQ/dt)

- “radiant flux”

Symbol: **P**

Unit: **Watt (Joule/sec)**

- “photons” per second
- measurable as current from a detector



Irradiance

Power per unit area (dP/dA)


- Defined with respect to a surface

Symbol **E**



Unit Watt/m²

- measurable as power on a small-area detector
- area density of power exiting a surface is also called **radiant exitance** (M) or **radiosity** (B)

Radiosity
Radiant exitance




Irradiance



Irradiance example

Uniform (point) light source with power P illuminates small surface A at distance r

- Think of A as a piece of the area of a sphere around the light
- P is uniformly spread over the area of this sphere




$$P_A = P \frac{A}{4\pi r^2}$$

$$E = \frac{P_A}{A} = \frac{P}{4\pi r^2}$$

Irradiance example

Uniform (point) light source with power P illuminates small surface A at distance r

- Think of A as a piece of the area of a sphere around the light
- P is uniformly spread over the area of this sphere



$$P_B = P_A = P \frac{A}{4\pi r^2}$$

$$E = \frac{P_B}{A} = \frac{P_B \cos \theta}{A} = \frac{P \cos \theta}{4\pi r^2}$$

Solid angle

Angle $\theta = s/r$

- subtended arc of circle
- s/r is unitless and called radians
- whole circle: 2π radians





Figure 5.8: Planar Angle. The planar angle of an object c as seen from a point p is equal to the angle it subtends as seen from p , or equivalently as the length of the arc s on the unit sphere.

Solid angle

Solid angle

- subtended area of sphere
- s/r^2 is unitless and called steradians
- whole sphere: 4π steradians




Solid angle and dodecahedron

Full sphere = 4π steradian = 12.566 sr

Dodecahedron


- 1 face = 1 steradian



Solid angle

Differential solid angle


$$d\omega = \frac{\cos \theta dA}{d^2}$$



Radiance

Power per unit solid angle, per unit area

- Irradiance per unit solid angle
- Measures light flow near a point x and direction ω
- “Counts photons” that
 - (a) go through a little area around x perpendicular to ω and
 - (b) are traveling in directions that fall in a little solid angle $d\omega$ around ω



$$L(x \rightarrow \omega) = \frac{d^2 P}{dA^\perp d\omega} = \frac{d^2 P}{dA \cos \theta d\omega}$$

◦

Symbol: L

Unit: $\text{Watt/m}^2 \text{ sr}$

Radiance


Power per unit solid angle, per unit area

- $L(x \rightarrow \omega)$
- x is position
- ω is direction

5D function

Defined for each wavelength

- (in practice r-g-b)



Some remarks w.r.t. notation

Outgoing radiance:

- $L_o(p, \omega)$ or $L(p \rightarrow \omega_o)$

Incoming radiance:

- $L_i(p, \omega)$ or $L(p \leftarrow \omega_o)$

All radiance is also wavelength dependent

- $L(p, \omega) \Rightarrow L(p, \omega, \lambda)$
- Often implicitly assumed as rgb-triplets

Integral over hemisphere of directions:

- 2D integral over hemispherical domain (see appendix)

Why is radiance important?

Radiance is invariant along a straight line

→ we can trace light energy along rays






Figure 15.3: Radiance along a Ray through Free Space Is Unchanged. Therefore, to compute the incident radiance along a ray from point p in direction ω , we can find the first surface the ray intersects and compute exitant radiance in the direction $-\omega$ there. The trace operator $t(p, \omega)$ gives the point p' on the first surface that the ray (p, ω) intersects.

Why is radiance important?




Why is radiance important?



Same radiance received from the Sun on the surface of Earth and Pluto?

BRDF – Bidirectional Reflectance Distribution Function



$$f_r(p, \omega_i, \omega_o) = \frac{dL_o(p, \omega_o)}{L_i(p, \omega_i) \cos \theta_i d\omega_i}$$

Properties of the BRDF

Reciprocity: $f_r(p, \omega_i, \omega_o) = f_r(p, \omega_o, \omega_i)$


BRDF usually does not depend on the intensity of light


BRDF is dependent on wavelength

- $f_r(p, \omega_i, \omega_o) \Rightarrow f_r(p, \omega_i, \omega_o, \lambda)$


Conservation of energy

- All reflected energy < incoming energy
- $\forall \omega_o: \int_{\Omega} f_r(p, \omega_o, \omega') \cos \theta' d\omega' \leq 1$





<https://cdfg.csail.mit.edu/wojciech/brdfdatabase>




<https://www1.cs.columbia.edu/CAVE/databases/tvbrdf/about.php>

Rendering Equation

$$f_r(p, \omega_i, \omega_o) = \frac{dL_o(p, \omega_o)}{L_i(p, \omega_i) \cos \theta_i d\omega_i}$$

$$dL_o(p, \omega_o) = f_r(p, \omega_i, \omega_o) L_i(p, \omega_i) \cos \theta_i d\omega_i$$

$$L_o(p, \omega_o) = L_e(p, \omega_o) + \int_{\text{hemisphere } \Omega} f_r(p, \omega_i, \omega_o) L_i(p, \omega_i) \cos \theta_i d\omega_i$$



Rendering Equation

Dallas, August 18-22

Volume 20, Number 4, 1986

THE RENDERING EQUATION

James T. Kajiya
California Institute of Technology
Pasadena, Ca. 91125

ABSTRACT. We present an integral equation which generalizes a variety of known rendering algorithms. In the course of discussing a monte carlo solution we also present a new form of variance reduction, called Hierarchical sampling and give a number of elaborations shows that it may be an efficient new technique for a wide variety of monte carlo procedures. The resulting rendering algorithm extends the range of optical phenomena which can be effectively simulated.

The equation is very much in the spirit of the radiosity equation, simply balancing the energy flows from one point of a surface to another. The equation states that the transport intensity of light from one surface point to another is simply the sum of the emitted light and the total light intensity which is scattered toward x from all other surface points. Equation (1) differs from the radiosity equation of course because, unlike the latter, no assumptions are made about reflectance characteristics of the surfaces involved.

Each of the quantities in the equation are new quantities which we call



Figure 6. A sample image. All objects are neutral grey. Color on the objects is due to caustics from the green glass balls and color bleeding from the base polygon.

Illuminated Mathematics

QUANTIFYING LIGHT IS THE KEY TO GREAT 3-D ANIMATION.

Light is one of the most challenging aspects of computer-generated imagery. It bounces around, reflects off surfaces, and changes color; if you don't get all those elements right, your animation will look no more real than Bugs Bunny—not necessarily bad, but not what the folks at places like Pixar are going for. In the mid-'80s, mathematicians quantified this process: If you know the intensity and color of the light arriving at a particular spot, you can figure out the same information for the light that bounces off. Here's the equation at the heart of every 3-D rendering package. —JULIE REHMEYER



$L_o(\lambda, x)$ The intensity in watts of a particular color (wavelength λ) of light from spot x to the viewer's eyeball.

L_e Light emitted from x . Unless some object—a lamp, the sun, a firefly, phosphorescent algae—is emitting a glow at x , this will be zero.

Ω A hemisphere around x . Each point on Ω represents one of the possible directions from which light could hit x .

\int Strain your brain back to calculus and you may remember that this is a fancy summation—adding up infinitely many infinitely tiny things. Here, you take Ω —every possible angle light may travel to x —compute how much each unit of that light bounces toward your eye, and then add 'em all up.

$$L_o(\lambda, x) = L_e + \int_{\Omega} f L_i \cos\theta d\omega$$

f This describes how much of the light that hits a material will reflect in a specific direction. A red surface, for example, will absorb every other color of light but red. A matte surface will spread light in all directions, reflecting only a bit toward your eye, while a mirrored surface will bounce all of the light in a single direction.

L_i The amount of incoming light of wavelength λ that strikes x from a particular direction.


$\cos\theta$ An adjustment for the angle of the light. Just as the low winter sun offers less heat than the high summer sun, less of the light coming from a low angle will bounce off x toward the eye than that arriving from a high angle.

$d\omega$ An infinitely small section of the hemisphere Ω around x —in other words, just one of the directions from which light could strike spot x .



Software for making animated films like *Despicable Me* relies on the equation above to model how light interacts with objects.

(WIRED magazine, July 2010)



directions for making pictures using numbers
(explained using only the ten hundred words people use most often)

the light that comes from an interesting direction towards the position on the stuff

direction towards the eye

position on the stuff

light made by the stuff (sometimes because it is very hot)

the answer to how much light from an interesting direction that will keep going in the direction towards the eye, after hitting stuff at the position (this is easy for mirrors, not so easy for everything else)

how much the light becomes less bright because the stuff leans away from the interesting direction

$$L_o(x, \omega_o) = L_e(x, \omega_o) + \int_{\Omega} f_r(x, \omega_i \rightarrow \omega_o) L_i(x, \omega_i) (\omega_i \cdot n) d\omega_i$$

light that leaves the position on the stuff and reaches the eye


light can be added said a man who sat under a tree many years ago

for lots of interesting directions inside half a ball facing up from the stuff, add up all the answers in between

this idea came from <http://xked.com/1133/>

@levork

Evaluating the Rendering Equation



https://www.youtube.com/watch?v=eo_MTI-d28s


Simple Stochastic Ray Tracing (aka Pure Path Tracing, Naïve Path Tracing, ...)

$$L_o(p, \omega_o) = ?$$

$$L_e(p, \omega_o)$$

+

$$\int_{\Omega} f_r(p, \omega_i, \omega_o) L_i(p, \omega_i) \cos \theta_i d\omega_i$$




(rendering L. Vanbesien)

Simple Stochastic Ray Tracing

Generate N random directions ω_j on hemisphere Ω around p , using probability density $p(\omega_j)$

$$\int_{\Omega} f_r(p, \omega_i, \omega_o) L_i(p, \omega_i) \cos \theta_i d\omega_i$$

$$\approx \frac{1}{N} \sum_{j=1}^N \frac{f_r(p, \omega_j, \omega_o) L_i(p, \omega_j) \cos \theta_j}{p(\omega_j)}$$



(rendering L. Vanbesien)


Simple Stochastic Ray Tracing

Generate N random directions ω_j on hemisphere Ω around p , using probability density $p(\omega_j)$

$$\int_{\Omega} f_r(p, \omega_i, \omega_o) L_i(p, \omega_i) \cos \theta_i d\omega_i \\ \approx \frac{1}{N} \sum_{j=1}^N \frac{f_r(p, \omega_j, \omega_o) L_i(p, \omega_j) \cos \theta_j}{p(\omega_j)}$$

Evaluate $L_i(p, \omega_j)$?







- Radiance is invariant along straight lines
- Trace ray to find closest intersection
- When path hits light source, we have a contribution
- Recursion



(rendering L. Vanbesien)

Simple Stochastic Ray Tracing

Reference - 1 - 4 - 16 - 64 - 256 viewing rays/pixel. No branching.



(rendering L. Vanbesien)

Simple Stochastic Ray Tracing

Choices?

- # viewing rays per pixel (anti-aliasing)
- # random rays over the hemisphere for each surface point: “branching factor”




Algorithm known as “path tracing”:

- Branching factor = 1
- Multiple viewing rays per pixel, but each ray branches only once at each reflection
- ‘path that is reflected through the scene’

Simple Stochastic Ray Tracing

Branching factor

Reference 4 viewing rays/pixel
Branching factor 1 - 2 - 4



(rendering L. Vanbesien)




Simple Stochastic Ray Tracing

Depth of Recursion

Reference



Recursion depth 1 - 2 - 3 is shown (same brightness)



Russian Roulette

Choose ‘**absorption probability**’ α

- probability α that ray will not be reflected → in this case, we stop the recursion
- probability $1 - \alpha$ that ray will be reflected → recursive evaluation is multiplied by $1/(1 - \alpha)$

E.g. $\alpha = 0.9$ ($1 - \alpha = 0.1$)

- 10% probability that ray is reflected
- If the ray is reflected, radiance along reflected ray ‘counts’ for $1 / 0.1 = 10$ times as much

E.g. $\alpha = 0.7$ (or $1 - \alpha = 0.3$)

- 30% probability that ray is reflected
- If the ray is reflected, radiance along reflected ray ‘counts’ for $1 / 0.3 = 3.3333$ times as much

Trivial stop-condition if $f_r = 0$ (set $\alpha = 1$)

- E.g. light sources are often modeled with $f_r = 0$

