



Review

Nigella sativa L as a potential phytotherapy for coronavirus disease

19: A mini review of in silico studies

Dr Abdulrahman E. Koshak^{1,*}, Prof Emad A. Koshak²¹ Department of Natural Products & Alternative Medicine, Faculty of Pharmacy, King Abdulaziz University, Jeddah, Saudi Arabia² Division of Allergy and Clinical Immunology, Department of Internal Medicine, Faculty of Medicine, King Abdulaziz University, Jeddah, Saudi Arabia

ARTICLE INFO

Article history:

Received 10 June 2020

Accepted 28 July 2020

Keywords:

Coronavirus

COVID-19

in silico

Nigella sativa

SARS-CoV-2

ABSTRACT

Background: Coronaviruses are responsible for several human diseases, such as the infectious novel coronavirus disease 2019 (COVID-19), which is caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). *Nigella sativa* is a natural food supplement with a known safety profile that may provide a wealth of documented antiviral compounds.

Objective: To explore the studies supporting the *N sativa* potential for hitting SARS-CoV-2 targets.

Methods: A literature search for published or preprint in silico studies between 1990 and 2020 in electronic databases (PubMed, Science Direct, Scopus, and Google Scholar) was performed for the terms *Nigella sativa*, *black seed*, *coronavirus*, *SARS-CoV-2*, and *COVID-19*.

Results: At least 8 in silico studies have shown that some compounds of *N sativa*, including nigellidine, α -hederin, hederagenin, thymohydroquinone, and thymoquinone, had high to moderate affinity with SARS-CoV-2 enzymes and proteins. These compounds may potentially inhibit SARS-CoV-2 replication and attachment to host cell receptors.

Conclusions: These preliminary data of in silico studies propose *N sativa* as a potential phytotherapy candidate for COVID-19. Further preclinical experimental evidence is required followed by a Phase I clinical trial. (*Curr Ther Res Clin Exp.* 2020; 81:XXX-XXX)

© 2020 The Author(s). Published by Elsevier Inc.

This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Introduction

Coronaviruses, enveloped RNA viruses, are characterized by spikes on their surface and belong to *Nidovirales* order.¹ They are responsible for a growing economic, social, and mortality burden in humans over the past decades. The spectrum of diseases associated with human coronaviruses range from the common cold to severe acute respiratory syndrome, and Middle East respiratory syndrome. Since December 2019, a newly discovered severe acute respiratory syndrome coronavirus (SARS-CoV-2) has been the causative agent of the current pandemic of infectious disease called coronavirus disease 2019 (COVID-19). Unfortunately, there are no effective approved antiviral agents for these coronavirus strains.^{2,3}

Natural products provide a wealth of biologically active molecules with antiviral activity, and thus may have utility as po-

tential therapeutic agents against coronavirus infections.⁴ Among these products is *Nigella sativa*, which has displayed several antiviral properties.⁵

N sativa is a well-known food supplement and medicinal plant in different cultures. The seeds of *N sativa* contain several active compounds in the classes of fixed oil, essential oil, saponins, and alkaloids. In the literature, *N sativa* exhibited several pharmacological properties including anti-inflammatory, antimicrobial, and immunostimulatory activities.^{5,6}

The safety and efficacy of *N sativa* used for many human diseases has been established in several randomized clinical studies.⁷ We also used *N sativa* oil in a randomized, double-blind placebo-controlled trial on asthmatic patients with acceptable safety and efficacy profile.⁸ Moreover, several meta-analyses have confirmed the beneficial effects and safety of *N sativa* on hyperlipidemia, type 2 diabetes, obesity, hypertension, and asthma.^{9–13} In a clinical study, Oral *N sativa* oil dosing of up to 5 g daily for up to 12 weeks is believed to be safe.¹⁴

In in vitro studies, the antiviral activities of *N sativa* on different viruses were documented in the literature.⁵ *N sativa* oil suppresses the viral load of murine cytomegalovirus in infected mice to an un-

* Address correspondence to: Dr Abdulrahman E. Koshak, 80260 Department of Natural Products & Alternative Medicine, Faculty of Pharmacy, King Abdulaziz University, Jeddah 21589, Saudi Arabia.

E-mail address: aekoshak@kau.edu.sa (D.A.E. Koshak).

Table 1A summary of effects of *Nigella sativa* compounds on severe acute respiratory syndrome-coronavirus disease 2 (SARS-CoV-2) targets.

Reference	<i>N sativa</i> material	SARS-CoV-2 targets	Control	Effects
16	Thymoquinone	6LU7	NA	-Thymoquinone had a moderate binding affinity with 6LU7
17	Nigellidine, α -Hederin	6LU7, 2GTB	-Chloroquine	-Nigellidine and α -hederin had the most binding affinity with 6LU7 and 2GTB
18	Hederagenin	6LU7, 6Y2E	-HCQ -Favipiravir Saquinavir	-Nigellidine was better than HCQ and favipiravir - α -Hederin better than chloroquine, HCQ, and favipiravir -Hederagenin had a high binding affinity with 6LU7 but less than saquinavir and 6Y2E close to saquinavir
19	Nigellidine	6LU7, NSP2, 6vsb, QHD43415_3, QHD43423, IL1R, TNFR1, TNFR2	NA	-Nigellidine had a high binding affinity with several SARS-CoV-2 and inflammatory molecular targets
20	Hederagenin	ACE2, GRP78	NA	-Hederagenin had the highest binding affinity with ACE2 and GRP78
21	Thymoquinone	6LU7, ACE2	HCQ	-Thymoquinone had a moderate binding affinity with 6LU7 and ACE2 1R42, but less than HCQ
22	Thymoquinone	HSPA5	NA	-Thymoquinone had a moderate binding affinity to HSPA5
23	Thymohydro-quinone	6LU7, Nsp15 / NendoU, ADRP, RdRp, rS, ACE2	NA	-Thymohydroquinone had a moderate binding affinity with several SARS-CoV-2 molecular targets

2GTB=main peptidase; 6LU7=main protease; 6vsb=spike glycoprotein; ACE2=angiotensin converting enzyme 2; ADRP=ADP-ribose-1''-phosphatase; HCQ=hydroxychloroquine; HSPA5=heat shock protein A5; IL1R=interleukin 1 receptor; NA=not available. NSP2=nonstructural protein 2; Nsp15/NendoU=endoribonuclease; QHD43415_3=N-terminus-proteinase; QHD43423=nucleocapsid; RdRp=RNA-dependent RNA polymerase; rS=binding domain of SARS-CoV-2 spike protein; TNFR1=tumor necrosis factor receptor 1; TNFR2=tumor necrosis factor receptor 2.

detectable level.¹⁵ *N sativa* honey was found to inhibit HIV-1 replication.¹⁶ *N sativa* had virucidal activity against herpes simplex and hepatitis A virus infections.¹⁶ *N sativa* decreased the coronavirus load in infected HeLa cells with stimulated interleukin 8 secretion and downregulation of transient receptor potential (TRP) genes expression such as TRPM6, TRPA1, TRPC4, and TRPM7.¹⁷ Hepatitis C virus replication was inhibited by *N sativa*.¹⁸ *N sativa* inhibited the growth of influenza virus H5N1 in vitro.¹⁹

In a human clinical study, patients with hepatitis C virus infection showed significant improvement in hepatitis C virus viral load after 3 months of *N sativa* treatment.²⁰ A case report of treatment with *N sativa* for 6 months showed a sustained seroreversion in a 46-year-old HIV patient and was also reported in an additional 6 HIV cases.^{21,22}

In recent years, in silico molecular docking studies on natural products enable computational screening approaches for assessing their therapeutic potential. These studies utilize bioinformatics techniques and can be used to discover how candidate drugs cause therapeutic activity by predicting interactions between drugs and proteins, and analysis of influence on biological pathways and functions.²³

The aim of this mini literature review was to explore any publication or preprint on in silico studies of the specific anticoronavirus potential of *N sativa*.

Methods

A literature search for scientific published manuscripts or preprint in silico studies found in electronic databases (PubMed, Science Direct, Scopus, and Google Scholar) was performed using the terms *Nigella sativa*, *black seed*, *coronavirus*, *SARS-CoV-2*, and *COVID-19*. Studies were searched for electronically between the years 1990 and 2020.

Results

In the literature review, there were at least 8 in silico studies that explored the effects of *N sativa* compounds on SARS-CoV-2. A summary of those studies is presented in the Table 1. However, there have been no reported clinical trials on *N sativa* in human coronavirus cases at this time.

Molecular docking of compounds from *N sativa* and some antiviral drugs was performed to determine their binding affinity

with SARS-CoV-2-related molecular targets such as main proteases (6LU7 and 6Y2E), main peptidase (2GTB), angiotensin converting enzyme 2 (ACE2), and heat shock protein A5. The binding of some natural compounds might prevent the adhesion of coronavirus to host epithelial cells. Nigellidine, an alkaloid in *N sativa*, docked with 6LU7 active sites showed an energy complex score close to chloroquine and better than hydroxychloroquine and favipiravir. α -Hederin, a saponin in *N sativa*, docked with 2GTB active sites showed an energy score better than chloroquine, hydroxychloroquine, and favipiravir.²⁴

Thymoquinone, the main essential oil constituent of *N sativa*, had a binding affinity with 6LU7, ACE2, and heat shock protein A5 active sites with a score less than hydroxychloroquine in 6LU7 and ACE2.^{25,26} Also, hederagenin, a saponin in *N sativa*, docked with 6LU7, 6Y2E, ACE2, and GRP78 active sites showed a binding score less than saquinavir in 6LU7 and 6Y2E.^{27,28} Thymohydroquinone showed moderate docking energy with SARS-CoV-2 6LU7, endoribonuclease, ADP-ribose-1''-phosphatase, RNA-dependent RNA polymerase, the binding domain of the SARS-CoV-2 spike protein, and human ACE2.²⁹ Nigellidine showed high binding affinity SARS-CoV-2 enzymes and proteins such as N-terminus-proteinase, 6LU7, nonstructural protein 2, spike-glycoprotein, and nucleocapsid. Nigellidine had high binding energy with human receptors, inflammatory signal molecules, and other proteins such as human IL1R (1itb), TNFR1 (1ncf), and TNFR2 (3alq).³⁰

Therefore, certain natural compounds found in *N sativa* such as nigellidine, α -hederin, hederagenin, thymohydroquinone, and thymoquinone were potentially active compounds that might inhibit coronavirus. Preclinical evidence is required to determine the activity of *N sativa* against coronavirus. If proven activity resulted from preclinical investigations, a clinical Phase I trial of *N sativa* in patients with COVID-19 is suggested to explore its clinical activity.

Conclusions

This mini literature review documented the inhibitory effects of some *N sativa* compounds against SARS-CoV-2 in several molecular docking studies. However, there is no reported clinical trial of *N sativa* in human coronavirus cases. Therefore, we propose *N sativa* as a potential phytotherapy candidate in further preclinical and clinical investigations in the treatment of coronavirus diseases such

as COVID-19. Also, further in silico investigation on other natural products from traditional medicines is suggested to apply them in the treatment of COVID-19.

Acknowledgments

All authors contributed equally in the literature search, study design, data collection, data interpretation, and writing.

Declaration of Competing Interest

None.

References

- Fehr AR, Perlman S. Coronaviruses: An overview of their replication and pathogenesis. *Coronaviruses: Methods and Protocols*, 1282, Springer; 2015:1–23.
- World Health Organization. Coronavirus; 2020 https://www.who.int/health-topics/coronavirus#tab=tab_1 Published 2020. Accessed May 12.
- Shereen MA, Khan S, Kazmi A, Bashir N, Siddique R. COVID-19 infection: Origin, transmission, and characteristics of human coronaviruses. *J Adv Res*. 2020;24:91–98. doi:10.1016/j.jare.2020.03.005.
- Lin LT, Hsu WC, Lin CC. Antiviral natural products and herbal medicines. *J Tradit Complement Med*. 2014;4(1):24–35. doi:10.4103/2225-4110.124335.
- Molla S, Abul M, Azad K, et al. A REVIEW ON ANTIVIRAL EFFECTS OF NIGELLA SATIVA L. *pharmacologyonline*, 2; 2019:47–53.
- Ahmad A, Husain A, Mujeeb M, et al. A review on therapeutic potential of *Nigella sativa*: A miracle herb. *Asian Pac J Trop Biomed*. 2013;3(5):337–352. doi:10.1016/S2221-1691(13)60075-1.
- Tavakkoli A, Mahdian V, Razavi BM, Hosseinzadeh H. Review on clinical trials of black seed (*Nigella sativa*) and its active constituent, thymoquinone. *J Pharmacopuncture*. 2017;20(3):179–193. doi:10.3831/KPI.2017.20.021.
- Koshak A, Wei L, Koshak E, et al. *Nigella sativa* Supplementation Improves Asthma Control and Biomarkers: A Randomized, Double-Blind, Placebo-Controlled Trial. *Phyther Res*. 2017;31(3). doi:10.1002/ptr.5761.
- He T, Xu X. The influence of *Nigella sativa* for asthma control: A meta-analysis. *Am J Emerg Med*. November 2019. doi:10.1016/j.ajem.2019.11.036.
- Sahebkar A, Soranna D, Liu X, et al. A systematic review and meta-analysis of randomized controlled trials investigating the effects of supplementation with *Nigella sativa* (black seed) on blood pressure. *J Hypertens*. 2016;34(11):2127–2135. doi:10.1097/HJH.0000000000001049.
- Namazi N, Larijani B, Ayati MH, Abdollahi M. The effects of *Nigella sativa* L. on obesity: A systematic review and meta-analysis. *J Ethnopharmacol*. 2018;219:173–181. doi:10.1016/j.jep.2018.03.001.
- Daryabeygi-Khotbehsara R, Golzarand M, Ghaffari MP, Djafarian K. *Nigella sativa* improves glucose homeostasis and serum lipids in type 2 diabetes: A systematic review and meta-analysis. *Complement Ther Med*. 2017;35:6–13. doi:10.1016/j.ctim.2017.08.016.
- Sahebkar A, Beccuti G, Simental-Mendía LE, Nobili V, Bo S. *Nigella sativa* (black seed) effects on plasma lipid concentrations in humans: A systematic review and meta-analysis of randomized placebo-controlled trials. *Pharmacol Res*. 2016;106:37–50. doi:10.1016/j.phrs.2016.02.008.
- Therapeutic Research Center. the Natural Medicines Research Collaboration. <https://naturalmedicines.therapeuticresearch.com/>. Published 2020. Accessed April 14, 2020.
- Salem ML, Hossain MS. Protective effect of black seed oil from *Nigella sativa* against murine cytomegalovirus infection. *Int J Immunopharmacol*. 2000;22(9):729–740. doi:10.1016/S0192-0561(00)00036-9.
- Barakat AB, Shoman SA, Dina N, Alfarouk OR. Antiviral activity and mode of action of *Dianthus caryophyllus* L. and *Lupinus termis* L. seed extracts against in vitro herpes simplex and hepatitis A viruses infection. *J Microbiol Antimicrob*. 2010;2(3):23–29. <http://www.academicjournals.org/JMA>. Accessed April 24, 2020.
- Ulasli M, Gurses SA, Bayraktar R, et al. The effects of *Nigella sativa* (Ns), *Anthemis hyalina* (Ah) and *Citrus sinensis* (Cs) extracts on the replication of coronavirus and the expression of TRP genes family. *Mol Biol Rep*. 2014;41(3):1703–1711. doi:10.1007/s11033-014-3019-7.
- Oyero OG, Toyama M, Mitsuhiro N, et al. Selective inhibition of hepatitis c virus replication by alpha-zam, a *Nigella sativa* seed formulation. *African J Tradit Complement Altern Med*. 2016;13(6):144–148. doi:10.21010/ajtcam.v13i6.20.
- Dorra N, El-Berrawy M, Sallam S, Mahmoud R. Evaluation of Antiviral and Antioxidant Activity of Selected Herbal Extracts. *J High Inst Public Heal*. 2019;49(1):36–40. doi:10.21608/jhiph.2019.29464.
- Barakat EMF, El Wakeel LM, Hagag RS. Effects of *Nigella sativa* on outcome of hepatitis C in Egypt. *World J Gastroenterol*. 2013;19(16):2529–2536. doi:10.3748/wjg.v19.i16.2529.
- Onifade AA, Jewell AP, Adediji WA. *Nigella sativa* concoction induced sustained seroreversion in HIV patient. *Afr J Tradit Complement Altern Med*. 2013;10(5):332–335.
- Onifade AA, Jewell AP, Ajadi TA, Rahamon SK, Ogunrin OO. Effectiveness of a herbal remedy in six HIV patients in Nigeria. *J Herb Med*. 2013;3(3):99–103. doi:10.1016/j.hermed.2013.04.006.
- Romano JD, Tatonetti NP. Informatics and computational methods in natural product drug discovery: A review and perspectives. *Front Genet*. 2019;10(APR):368. doi:10.3389/fgene.2019.00368.
- Bouchentouf S, Noureddine M. Identification of Compounds from *Nigella Sativa* as New Potential Inhibitors of 2019 Novel Coronavirus (COVID-19): Molecular Docking Study Elucidation of neurodegenerative pathologies processes by molecular modeling View project Molecular operating enviro. *ChemRxiv*. doi:10.26434/chemrxiv.12055716.v1.
- Sekiou O, Ismail B, Zihad B, Abdelhak D. In-Silico Identification of Potent Inhibitors of COVID-19 Main Protease (Mpro) and Angiotensin Converting Enzyme 2 (ACE2) from Natural Products: Quercetin, Hispidulin, and Cirsimaritin Exhibited Better Potential Inhibition than Hydroxy-Chloroquine Against. *chemRxiv*. April 2020. doi:10.26434/chemrxiv.12181404.v1.
- Elfiky AA. Natural products may interfere with SARS-CoV-2 attachment to the host cell. *J Biomol Struct Dyn*. April 2020:1–16. doi:10.1080/07391102.2020.1761881.
- Sampangi-ramaiah MH, Vishwakarma R, Shaanker RU. Molecular docking analysis of selected natural products from plants for inhibition of SARS-CoV-2 main protease. *Curr Sci*. 2020;118(7):1087–1092. <http://sts.bioe.uic.edu/castp/index.html?3igg>. Accessed May 3, 2020.
- Rajapaksa RMH, Perera BT, Nisansala MJ, Perera WPRT, Dissanayake KGC. POTENTIAL OF INHIBITING THE RECEPTOR BINDING MECHANISM OF SARS- COV-2 USING PHYTOCHEMICAL EXTRACTS OF MEDICINAL HERB; MOLECULAR DOCKING STUDY. *Glob J Eng Sci Res Manag*. 2020;7(4):51–61. doi:10.5281/zenodo.3766184.
- da Silva JKR, PLB Figueiredo, Byler KG, Setzer WN. Essential Oils as Antiviral Agents. Potential of Essential Oils to Treat SARS–CoV–2 Infection: An In–Silico Investigation. *Int J Mol Sci*. 2020;21(10):3426. doi:10.3390/ijms21103426.
- Maiti S, Banerjee A, Nazmeen A, Kanwar M, Das S. Active-site Molecular docking of Nigellidine to nucleocapsid / Nsp2 / Nsp3 / M Pro of COVID-19 and to human IL1R and TNFR1 / 2 may stop viral- growth / cytokine-flood , and the drug source *Nigella sativa* (black cumini) seeds show potent antioxidant role. *Research Square*. doi:10.21203/rs.3.rs-26464/v1.