



Inspire...Educate...Transform.


Unsupervised models

**Clustering- Understanding
Distances**

UNDERSTANDING DISTANCE



Numeric



Distance between categorical attributes

- For each categorical variable
 - Distance is zero if two records have same value
 - 1 if different
 - Sum the total distance across all attributes



Categorical attributes

$x_i \rightarrow$	1	1	1	0	1	0	0	1	0	1	0	1
$x_j \rightarrow$	1	1	0	1	1	0	1	1	0	0	1	1

For example, in above two feature vectors:

$a=5$
 $b=2$
 $c=3$
 $d=2$

(10)

		Data point j		
		1	0	
Data point i	1	a	b	$a+b$
	0	c	d	$c+d$
		$a+c$	$b+d$	$a+b+c+d$

- a : the number of attributes with the value of 1 for both data points.
- b : the number of attributes for which $x_{if} = 1$ and $x_{jf} = 0$, where x_{if} (x_{jf}) is the value of the f th attribute of the data point \mathbf{x}_i (\mathbf{x}_j).
- c : the number of attributes for which $x_{if} = 0$ and $x_{jf} = 1$.
- d : the number of attributes with the value of 0 for both data points.



Symmetric binary attributes

- Hamming distance function: Simple Matching Coefficient, proportion of mismatches of their values

$$\text{dist}(\mathbf{x}_i, \mathbf{x}_j) = \frac{b + c}{a + b + c + d}$$

$$\text{Dist} = \frac{\text{number of dissimilar attributes between the records}}{\text{number of dissimilar attributes} + \text{number of similar attributes}}$$



Asymmetric binary attributes

- **Asymmetric**: if one of the states is more important or more valuable than the other.
 - By convention, state 1 represents the more important state, which is typically the rare or infrequent state.
 - **Jaccard coefficient** is a popular measure

$$\text{Dist} = \frac{\text{number of dissimilar attributes between the records}}{\text{number of dissimilar attributes} + \text{number of similar attributes (excluding records with 0,0)}}$$

$$\text{dist}(\mathbf{x}_i, \mathbf{x}_j) = \frac{b + c}{a + b + c}$$

- We can have some variations, adding weights

Dissimilarity between Binary Variables

- Example

Name	Gender	Fever	Cough	Test-1	Test-2	Test-3	Test-4
Jack	M	Y	N	P	N	N	N
Mary	F	Y	N	P	N	P	N
Jim	M	Y	P	N	N	N	N

- gender is a symmetric attribute
- the remaining attributes are asymmetric binary
- let the values Y and P be set to 1, and the value N be set to 0

$$d(jack, mary) = \frac{0 + 1}{2 + 0 + 1} = 0.33$$

$$d(jack, jim) = \frac{1 + 1}{1 + 1 + 1} = 0.67$$

$$d(jim, mary) = \frac{1 + 2}{1 + 1 + 2} = 0.75$$



International School of Engineering

Plot 63/A, 1st Floor, Road # 13, Film Nagar, Jubilee Hills, Hyderabad - 500 033

For Individuals: +91-9502334561/63 or 040-65743991

For Corporates: +91-9618483483

Web: <http://www.insofe.edu.in>

Facebook: <https://www.facebook.com/insofe>

Twitter: <https://twitter.com/Insofeedu>

YouTube: <http://www.youtube.com/InsofeVideos>

SlideShare: <http://www.slideshare.net/INSOFE>

LinkedIn: <http://www.linkedin.com/company/international-school-of-engineering>

This presentation may contain references to findings of various reports available in the public domain. INSOF makes no representation as to their accuracy or that the organization subscribes to those findings.