

Root architecture: from a root picture to a working model

An original, passive-listening explanation of the OpenSimRoot methods paper: how root growth rules, carbon costs, soil resources, and measured root images can be connected, with careful limits and a sorghum modeling interpretation.

B1–B2 English | 1,771 words

Research source: Johannes A. Postma, Christian Kuppe, Markus R. Owen, Nathan Mellor, Marcus Griffiths, Malcolm J. Bennett, Jonathan P. Lynch, and Michelle Watt (2017), New Phytologist. [Read the original study.](#)

Full listening script

Highlighted words appear in the plain-English glossary after the script.

Welcome. Today we will explore a simple idea: a root system is more than a collection of lines in the soil. Its shape changes where the plant can find resources. Its living tissues also need resources to grow and work. A useful root model connects these two sides. It describes where roots go, and what those roots can do after they arrive.

Our paper is called “OpenSimRoot: widening the scope and application of root architectural models.” Johannes Postma and seven colleagues published it in New Phytologist in June twenty seventeen. This is an original methods paper. Its main contribution is a modeling framework and demonstrations of its use. It is not a new field trial proving that one root shape always produces the greatest yield.

The key term is **root system architecture**. Architecture means the arrangement of the root system in space and time. It includes the directions of roots, where branches appear, how fast they grow, and how different root types connect. Two plants can have the same total root length but very different architectures. One can place much of that length close to the surface. Another can place more of it deeper down.

A picture can show this difference. But a picture cannot directly tell us how much water enters each root, how much nitrogen the plant gains, or how much carbon the roots consume. Carbon here means the material and energy supplied largely through sugars made by the leaves. The central purpose of OpenSimRoot is to connect root structure with these functions. Scientists call this a **functional–structural model**.

Let us start with how the model grows a root. The root system is represented by connected points and **segments**. A segment is a short piece between two points. The growing tip changes position. As it moves, it leaves a record of the root behind it. For each new root, the model needs three kinds of information: when and where it begins, how quickly it grows, and which direction it takes.

Different root classes can have different rules. A main root and a small side root do not need to grow at the same speed. They can differ in diameter, branching pattern, and growth direction. This is important in grasses, where roots can emerge from the seed or later from the stem. A single average root may hide differences that matter for the whole plant.

The model also allows variation. Real roots of the same type are not perfectly identical. OpenSimRoot can give them different growth rates and can change their direction as they extend. One influence is **gravitropism**, which means a growth response to gravity. Other influences can represent local conditions or random changes in direction. These rules produce a three-dimensional network rather than one fixed root shape.

However, more root growth is not free. Every new segment requires carbon. Existing tissues also use carbon through **respiration**, the process that releases energy for living cells. The model compares the carbon available with the carbon needed for potential growth. Potential growth means the growth that could occur if resources were sufficient. If available carbon becomes too low, actual growth can fall below that potential.

This connection creates a useful way to think about branching. Imagine adding many more side roots. At first, this seems helpful because the plant gains more places to absorb resources. But those roots must share the available carbon. They may also compete with each other for the same nearby nutrients. More starting points do not automatically produce more useful exploration. Some branches may remain short because the budget is limited.

The twenty seventh paper explains these trade-offs using earlier SimRoot studies. We should keep that history clear. Results about the best branching density under low nitrogen were not all new experiments in this paper. They were examples of questions that the model family had already helped investigate. The new paper brought the framework together, opened the code, and expanded what researchers could connect and test.

Another important feature is **plasticity**. In plant science, plasticity means that a plant changes its development in response to its environment. In the model, a local nutrient level can change the rate of root extension or the number of branches. A soil region with abundant phosphorus can therefore support a different branching pattern from a region with little phosphorus, if the chosen response rule describes that behavior.

One demonstration placed more phosphorus near the soil surface. The plastic root system produced denser branching in this upper region. Deeper down, fewer branches allowed individual side roots to grow longer. This is a result of the specified model rules and resource interactions. It is not evidence that every crop uses exactly that response. The example shows how a local developmental rule can create a different whole-root pattern.

The distinction between phosphorus and nitrate is also useful. Nitrate is a form of nitrogen that can move with soil water. Phosphorus generally moves much less easily through soil. Because these resources behave differently, the paper describes different approaches for modeling their movement toward roots. A root arrangement that works well for one resource may work differently for another. The soil process and the root shape have to be considered together.

Water adds another connection. OpenSimRoot links movement through the soil, flow inside the root network, and water loss from the plant to the atmosphere. Xylem is the tissue that carries water through roots and shoots. A hydraulic network model represents the connected water pathways in the roots. Hydraulic simply means related to water movement. The soil model describes how water moves through the spaces between soil particles.

This coupling can produce **compensatory uptake**. That means roots in a wetter region take up more water when another region becomes less useful. The paper also shows that water can sometimes move through roots and return to drier soil. This is **hydraulic redistribution**. These behaviors depend on water conditions and the root network. They cannot be read directly from total root length alone.

Now we reach one of the paper's most interesting demonstrations: turning an observed root system into a functional model. The researchers used three-dimensional images from magnetic resonance imaging and X-ray computed tomography. These methods can reveal roots inside soil. The model can take measured positions as input, rather than creating every part of the architecture from general growth rules.

In one example, the authors reproduced an imaged maize root system. About eight percent of the roots visible in the original image were missing from the modeled reconstruction because of limits in image processing. This detail matters. A computer model is only as complete as the information supplied to it. Missing fine roots can affect estimated uptake, even when the larger visible roots look convincing.

The framework can also add fine roots that the imaging method cannot detect, using growth rules. That is useful, but the added roots are predictions. They are not newly observed roots. In another demonstration, an imaged wheat root system was used to simulate nitrate depletion in the surrounding soil. A **depletion zone** is a region where the nutrient concentration falls because roots remove nutrients. The image supplied structure; the model supplied an estimate of function.

This gives a careful way to interpret a colorful simulation. A realistic image of roots is one kind of evidence. A predicted pattern of nutrient removal is another. The second depends on the first, but also on assumptions about soil conditions, uptake behavior, and time. The paper is valuable partly because it makes these connections explicit. It helps researchers ask which missing measurement could most improve the prediction.

The model has important limits. In the version described in twenty seventeen, the water module did not simulate drought-related growth responses. It could calculate water movement without representing every way drought changes plant development. The shoot was also simplified, rather than represented as a detailed three-dimensional canopy. These are limits of the version reported in the paper. They should not be presented as a description of every later version.

Some carbon-allocation rules were practical choices rather than established physiological laws. **Carbon allocation** means how the plant shares carbon among its organs. For example, one rule gave different priorities to shoots, main roots, and branches when carbon was scarce. Changing that rule

can change the result. The authors also noted that their local plasticity responses did not include a mechanism for sensing environmental gradients.

These limits do not make the model useless. They tell us what kind of conclusion is justified. The authors emphasized the model as a tool for exploring mechanisms and testing ideas. A mechanism is an explanation of how a process works. A model can show that a proposed mechanism is sufficient to produce a pattern. It does not, by itself, prove that the same mechanism explains every observation in a real field.

For sorghum work using CPlantBox and DuMuX, the following connection is our interpretation, rather than a result tested in this paper. A good comparison should separate architecture from physiology. Architecture describes where the roots are. Physiology describes how the living roots function. If a simulated sorghum plant takes up more water, the cause might be deeper roots, different water conductance, or a change in plant demand. These possibilities should not be mixed together without checking.

One possible research design would begin with measured sorghum root growth and branching. CPlantBox could represent alternative architectures, while a coupled soil-water calculation could explore their behavior under the same soil and weather conditions. First, the comparison could keep root hydraulic properties fixed. Later, those properties could vary. This would help distinguish the effect of root placement from the effect of water transport inside the roots. It would be a new study, not a conclusion already established here.

There is also a lesson about measurement. A model comparison is stronger when it is checked against several observations. Root depth alone is not enough. Useful evidence might include root length at different depths, changes in soil water, and the total water lost by the plant. Agreement with one measurement can hide disagreement with another. Checking several parts of the system makes it harder for the right answer to arise for the wrong reason.

The overall message is that root architecture becomes more informative when it is connected to function and cost. More roots, deeper roots, or more branches are descriptions, not complete explanations. OpenSimRoot offered a transparent way to connect developmental rules, carbon supply, soil resources, and uptake. For a sorghum modeling project, the lasting lesson is to ask not only what root system appears, but why it appears and which evidence supports the function we assign to it.

Useful words

Plain-English meanings in the order supplied, with usage examples where available.

root system architecture: The arrangement and connections of roots in space and time.

Example: Two plants can have the same root length but different root system architecture.

functional–structural model: A model that connects the shape of a plant with processes such as growth and uptake.

Example: A functional–structural model links root position with water uptake.

segment: A short piece of a root represented between two points.

Example: Each root segment can have its own age and diameter.

gravitropism: A growth response to gravity.

Example: Gravitropism influences the direction in which a root grows.

respiration: The process through which living cells release energy for their activities.

Example: Root respiration is part of the carbon cost of soil exploration.

plasticity: A change in development in response to environmental conditions.

Example: Root plasticity can change branching in a nutrient-rich patch.

compensatory uptake: Greater uptake from one region when another region supplies less.

Example: Roots in wet soil may provide compensatory uptake.

hydraulic redistribution: Movement of water through roots from wetter soil into drier soil.

Example: A root network can sometimes enable hydraulic redistribution.

depletion zone: An area where a resource concentration falls because roots remove it.

Example: The simulation estimates a nitrate depletion zone around the roots.

carbon allocation: How a plant shares carbon among organs and processes.

Example: A carbon-allocation rule can influence the growth of side roots.

Sources and attribution

Original research

Johannes A. Postma, Christian Kuppe, Markus R. Owen, Nathan Mellor, Marcus Griffiths, Malcolm J. Bennett, Jonathan P. Lynch, and Michelle Watt (2017). OpenSimRoot: widening the scope and application of root architectural models. *New Phytologist*, 215. Published 27 June 2017.

[doi:10.1111/nph.14641](https://doi.org/10.1111/nph.14641)

<https://nph.onlinelibrary.wiley.com/doi/10.1111/nph.14641>

Source license: Creative Commons Attribution License (article license verified in PMC publication record; version not asserted).

Original listening lesson based on Postma et al. (2017), *New Phytologist* 215:1274–1286, DOI 10.1111/nph.14641. Source article distributed under a Creative Commons Attribution License. The lesson paraphrases and explains the paper; it is not the original article.

How to read the evidence

The primary source is an original methods/model-development paper, not a review or a new field-yield trial. Earlier experiments cited by that paper are explicitly identified as earlier work. Statements about the 2017 software version are time-bounded. Sorghum/CPlantBox/DuMuX applications are interpretation and proposed tests, not findings of the source paper.

Sources checked on 1 October 2026.