

Competing pathways in polaritonic chemistry and general harmonic environments

Jonas Vinther,^{1,2, a)} Leonardo A. Cunha,^{3,4} and Johannes Flick^{3,5,6, b)}

¹⁾ *Department of Computer Science, University of Copenhagen, DK*

²⁾ *Niels Bohr Institute, University of Copenhagen, DK*

³⁾ *Center for Computational Quantum Physics, Flatiron Institute, 162 5th Avenue, New York, New York 10010, USA*

⁴⁾ *Department of Chemistry and Biochemistry, Bates College, Lewiston, Maine 04240, USA*

⁵⁾ *Department of Physics, City College of New York, New York, New York 10031, USA*

⁶⁾ *Department of Physics, The Graduate Center, City University of New York, New York, New York 10016, USA*

Recent experiments suggest that strong light-matter coupling between molecules and cavity modes can significantly modify chemical reaction rates and yields. Despite multiple proposed mechanisms, a clear microscopic understanding of how vibrational strong coupling alters reaction kinetics remains incomplete. Here, we employ a general quantum-dynamical framework for analyzing competing reaction pathways in polaritonic chemistry by considering reaction coordinates with multiple relevant transition frequencies and general harmonic environments. We extend on a tensor-network formulation of the hierarchical equations of motion (HEOM) method to compute probability currents between vibrational states, leveraging a concise formulation for effective spectral densities and a controlled compression of Markovian bath modes into system-level jump operators. This allows us to decompose the resulting state-to-state probability currents, revealing how cavity-induced changes in individual transitions propagate through multistep reaction networks. We find that these local modifications can alter kinetic bottlenecks and selectively redistribute flux between competing pathways, changing branching ratios under irreversible conditions without substantially affecting the equilibrium yield, in the reversible regime. Finally, for coupled reaction-coordinate, spectator-mode, and cavity environments, we find structured, asymmetric rate profiles arising from interference between hybridized decay channels. Together, these results provide a mechanistic framework for understanding pathway-specific chemical reactivity under vibrational strong coupling.

I. INTRODUCTION

Hybrid light-matter states, known as polaritons, have received increased attention partly due to the novel phenomena that can emerge when molecules and materials are embedded in structured electromagnetic environments^{1,2}. In polaritonic chemistry³⁻⁵, for instance, an ensemble of molecules is strongly coupled to the harmonic modes of an optical cavity. Even in the dark, these strong light-matter interactions have been observed to alter the reaction rate and chemical yield of the molecular processes involved. The seminal discovery by Hutchison et al. in 2012, provided the first experimental observation that hybrid light-matter states could actively modify chemical landscapes⁶. This was followed by Shalabney et al.’s demonstration of vibrational strong coupling (VSC)⁷ and, in 2016, by Thomas et al.’s report of modified ground-state chemical reactivity by strongly coupling the vibrational modes of an ensemble of molecules to the vacuum electromagnetic field⁸. These developments paved the way to a flurry of subsequent experiments⁹⁻¹⁴ and form part of a broader effort to exploit polaritonic states to control molecular processes, including intermolecular energy transfer^{15,16}, long-range energy transport¹⁷, photochemical stability¹⁸, and selective emission from molecular isomers¹⁹. A recent review

surveys these emerging functionalities²⁰.

For ground-state chemical reactivity, experiments have reported a diverse range of responses to VSC. Reaction suppression has been reported by strongly coupling cavity modes to reactant vibrations^{8,21} or additionally to product or cooperative reactant-solvent modes¹². Reaction acceleration has been observed under cooperative coupling of overlapping vibrations of the reacting molecules and the surrounding solvent, both in Fabry-Pérot cavities^{10,13} and using resonances supported by non-local metasurfaces¹⁴. Temperature-dependent kinetic studies of both suppressed and accelerated reactions have further revealed compensating changes in apparent activation parameters under cavity coupling^{11,13}. It should be noted, as a testament to the difficulty of bridging the gap between theory and experiments, that the reproducibility of some reported rate enhancements remains contested²², with subsequent analyses identifying optical and kinetic-fitting artifacts that may contribute to discrepancies between studies²³.

In addition to modifying overall reaction rates, VSC has been reported to alter product selectivity. Ref. 9 observed a redistribution of products arising from competing pathways. Ref. 24 further showed that coupling different vibrations of the same molecule biased the product distribution in opposite directions, even though both coupling conditions slowed the overall reaction. Alongside these changes in kinetic selectivity, Ref. 25 reported shifts in chemical equilibrium, with the direction of the shift depending on the coupled vibrational mode.

Together, these findings motivate microscopic descrip-

^{a)} jonas.vinther@nbi.ku.dk

^{b)} jflick@ccny.cuny.edu

tions that resolve how coupling at distinct vibrational frequencies affects both molecular transitions and equilibrium populations, such that the conditions under which cavity coupling can control reaction rates and product distributions, can be identified.

In an attempt to interpret this range of observed cavity-modified reactivity, theoretical approaches have drawn on transition-state theory²⁶ and dynamical barrier-crossing theories²⁷, such as Grote-Hynes²⁸ and Pollak-Grabert-Hänggi theory^{29,30}. Despite the ability to account for environmental effects, these approaches are fundamentally concerned with the energy barrier between reactant and product populations and their classical implementation do not explicitly resolve transitions between quantized vibrational states. Consequently, some classical models predict rate-modification profiles that are substantially broader than reported experimentally, or whose extrema occur near the barrier frequency rather than the relevant molecular vibrational frequencies³¹. Comparisons with quantum simulations have further demonstrated limitations of approximate treatments of molecular and cavity dynamics in reproducing resonant rate modifications³².

Quantum dynamical treatments have demonstrated that the sharp resonance in VSC-modified reaction rates can be recovered when the molecular, cavity, and environmental degrees of freedom are treated quantum mechanically^{31,33–38}. Complementary theoretical analyses based on effective environmental descriptions and Fermi’s Golden Rule have provided microscopic interpretations of this resonance and its dependence on cavity properties^{33–36}. Tensor-network approaches have extended quantum simulations to models of collective effects^{37–39}, yet, establishing the connection between such microscopic models and collective experimental systems nevertheless remains an open challenge⁴⁰. Additional theoretical efforts seek to relate changes in the equilibrium chemical composition to the thermodynamics of the coupled light-matter system⁴¹.

An important next step is to determine how frequency-selective modifications of individual transitions propagate through reactions involving distinct reactant and product vibrational spectra or multiple accessible products. Furthermore, extending the mechanistic insights provided by Fermi’s golden rule through a nonperturbative analysis of state-to-state probability currents, resolved into system and environmental contributions, would further clarify how individual vibrational transitions translate into changes in overall reaction rates and product distributions.

Here, we extend a tensor-network formulation of the hierarchical equations of motion to analyze reaction coordinates with independently adjustable reactant and product well frequencies, including a model with two competing product wells. We combine quantum dynamical simulations with effective spectral densities for general harmonic environments to extract probability currents between vibrational states and decompose these into sys-

tem and environmental contributions. This allows us to identify how cavity-induced modifications of individual transitions alter kinetic bottlenecks and redistribute flux between competing pathways. By examining both reaction dynamics and equilibrium populations of the cavity- and bath-coupled system, we distinguish changes in kinetic selectivity from changes in thermodynamic preference.

This paper is organized as follows. Section II formulates the model of the reaction coordinate and its harmonic environment. It also presents the reaction rate, defined in terms of system observables, and the probability currents, which serves as a diagnostic tool for the reaction mechanism. Section III describes our method for performing the numerically exact simulations of the dynamics. The results are presented and discussed in section IV, where we address potential energy surfaces with multiple relevant transition frequencies and competing pathways and harmonic environments with interfering decay channels. Our conclusions are summarized in Sec. V.




FIG. 1. An example setup consists of a single molecule (red), comprising a reaction coordinate, and may also include spectator vibrational modes. The molecule is embedded in a solvent environment represented by a phonon bath (purple). The molecular and solvent degrees of freedom are confined within an optical cavity, represented by a single cavity mode (green).

II. THEORY

A. Model and effective harmonic environment

We employ a complete quantum description of a reaction coordinate (the system) on a one dimensional potential energy surface (PES) in a harmonic environment (the bath). A schematic for the physical setup is presented in Fig. 1.

The microscopic model that we consider, is given by the Pauli-Fierz Hamiltonian in the length gauge^{31,36,42}:

$$H = H_R + H_Q + H_c + H_R^{\text{CL}} + H_Q^{\text{CL}} + H_c^{\text{CL}} \quad (1)$$

$$H_R = \frac{\hat{P}^2}{2M} + V(\hat{R}) \quad (2)$$

$$H_Q = \frac{\hat{P}_Q^2}{2} + \frac{\omega_Q^2}{2} \left(\hat{Q} + \frac{c_Q}{\omega_Q^2} \hat{R} \right)^2 \quad (3)$$

$$H_c = \frac{\hat{p}_c^2}{2} + \frac{\omega_c^2}{2} \left(\hat{q}_c + \sqrt{\frac{2}{\omega_c}} \eta_c (\eta_R \hat{R} + \eta_Q \hat{Q}) \right)^2 \quad (4)$$

This Hamiltonian describes a reaction coordinate $\hat{R}$, coupled to the displacement $\hat{Q}$ of a spectator mode, both of which are coupled to a single cavity mode with displacement $\hat{q}_c$. Furthermore, each of these coordinates are assumed to experience their own harmonic environment, as described by the Caldeira-Leggett model of dissipation⁴³, via the inclusion of $H_{(R,Q,c)}^{\text{CL}}$ in Eq. 1. These are of the form:

$$H_R^{\text{CL}} = \sum_l \frac{\hat{p}_l^2}{2} + \frac{\omega_l^2}{2} \left(\hat{x}_l - \frac{c_l}{\omega_l^2} \hat{R} \right)^2 \quad (5)$$

The reaction coordinate (RC) is the only anharmonic degree of freedom in the Hamiltonian, Eq. 1, and all the other harmonic degrees of freedom may be referred to as the bath. Consequently, if we assume that the RC is initially in a product state with the bath and that the bath is in a thermal state, the influence of the bath is fully described by the bath correlation function (BCF)^{44,45}:

$$C(t) = \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} S_\beta(\omega) e^{-i\omega t} = \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} \frac{2J(\omega) e^{-i\omega t}}{1 - e^{-\beta\hbar\omega}} \quad (6)$$

where $S_\beta(\omega)$ is the noise spectrum, β is the inverse temperature and $J(\omega)$ is the spectral density of the full harmonic environment.

In Supplementary Section S2 we provide a succinct formula for $J(\omega)$ for arbitrary harmonic environments. It relies on the inversion of a matrix that can be formed straightforwardly from the parameters of the Hamiltonian, thus mitigating the algebraic labor that earlier methods required^{36,46}.

Figure 2 schematically shows the abstraction from the microscopic harmonic environment in question to the full spectral density $J(\omega)$. With our approach, Eq. 1 immediately leads to:

$$J(\omega) = J_R^{(DL)}(\omega) + \text{Im} [\mathbf{c}^T M^{-1}(\omega) \mathbf{c}] \quad (7)$$

$$\mathbf{c} = \begin{pmatrix} \sqrt{2\omega_c^3} \eta_R \eta_c \\ c_Q + 2\omega_c \eta_c^2 \eta_R \eta_Q \end{pmatrix} \quad (8)$$

$$M(\omega) = \begin{pmatrix} \Gamma_c(\omega) & -\sqrt{2\omega_c^3} \eta_Q \eta_c \\ -\sqrt{2\omega_c^3} \eta_Q \eta_c & \Gamma_Q(\omega) - 2\omega_c \eta_c^2 \eta_Q^2 \end{pmatrix} \quad (9)$$

$$\Gamma_{(c,Q)}(\omega) = \omega^2 - \omega_{(c,Q)}^2 - \left(i + \frac{\omega}{\gamma_{(c,Q)}} \right) J_{(c,Q)}^{(DL)}(\omega) \quad (10)$$

For simplicity, we have assumed that the spectral densities of the individual harmonic environments, described by $H_{(R,Q,c)}^{\text{CL}}$, are all of the Drude-Lorentz form:

$$J_n^{(DL)}(\omega) = \frac{2\lambda_n \gamma_n \omega}{\omega^2 + \gamma_n^2} \quad \text{with } n \in \{R, Q, c\} \quad (11)$$




FIG. 2. Schematic for the efficient modeling of the harmonic degrees of freedom in terms of the spectral density for the full harmonic environment. Supplementary Section S2 derives and presents the formula for arbitrary harmonic environments. The system refers to the reaction coordinate $\hat{R}$, here visualized in terms of the PES $V(\hat{R})$ for the triple well system treated in Sec. IV A 3. The spectator mode has coordinate $\hat{Q}$ and the cavity $\hat{q}_c$. The couplings between them are denoted by c_Q , η_R and η_Q and $J_{(R,Q,c)}^{(DL)}(\omega)$ refer to the individual baths which we choose to be of the Drude-Lorentz form.

Finally, we consider the cavity bath to be Markovian, and consider the limit $\gamma_c \rightarrow \infty$ with $\frac{2\lambda_c}{\gamma_c} = \alpha(\omega_c) = \frac{1 - e^{-\beta\hbar\omega_c}}{\tau_c}$ near constant ??




FIG. 3. The noise spectrum when there is no spectator mode ($c_Q = \eta_Q = 0$). The effect of the cavity is captured by the peak centered around ω_c (Eq. 12). The height of this peak scales with $\sim \omega_c^2$, as dictated by the Pauli-Fierz Hamiltonian. For visualizations of noise spectra with the full spectral density of Eq. 7, see Fig. 8

In the case of no spectator mode ($c_Q = \eta_Q = 0$), the harmonic environment reduces to that of the phonon

bath (modeled by the spectral density $J_R^{(DL)}(\omega)$) plus the lossy cavity³³:

$$J(\omega) = J_R^{(DL)}(\omega) + \frac{2\omega_c^3 \eta_R^2 \eta_c^2 \alpha(\omega_c) \omega}{(\omega^2 - \omega_c^2)^2 + \alpha^2(\omega_c) \omega^2}. \quad (12)$$

The resulting noise spectrum is visualized for three different cavity frequencies in Fig. 3.

As alluded to in Fig. 2, the cavity-modified kinetics are essentially governed by two scalar function: the potential energy surface $V(\hat{R})$ and the effective spectral density for the full harmonic environment $J(\omega)$. The former accounts for the specifics of the reaction while the latter contains all the harmonic degrees of freedom such as lossy cavity modes, spectator modes and the vibrations of the solvent. Thus, the simple spectral density (Eq. 12) will be used to isolate the effects of the cavity when studying PES driven effects (Sec. IV A). The full spectral density (Eq. 7) will be applied to study how the alignment between the reaction coordinate and spectator mode polarities, encoded in η_R and η_Q respectively, affects the reaction rate profile (Sec. IV B).

We parameterize the potential energy surface $V(\hat{R})$ in terms of piecewise parabolic functions, which provides a straightforward way to study independent well frequencies and competing pathways. Explicit expressions for $V(\hat{R})$ are found in Supplementary Section S3. While the simulations are carried out in the eigenbasis of the system Hamiltonian H_R , the analysis is performed in a basis where select eigenstates have been hybridized to construct localized states. Our approach for constructing the rotation between these two bases relies on diagonalizing $\hat{R}$ in the subspace of the selected states (see Supplementary Section S3.3).

B. Observables and Mechanistic Analysis

The reaction rate is defined in terms of expectation values of projection operators onto a reactant region and product regions^{47,48}. We define these explicitly, in terms of the eigenstates $\hat{R}|R\rangle = R|R\rangle$. The double well PESs are centered around $R = 0$ with a reactant region in the left well $\hat{P}_{\text{left}} = \int_{-\infty}^0 dR |R\rangle \langle R|$ and a product region in the right well $\hat{P}_{\text{right}} = \mathbb{1} - \hat{P}_{\text{left}}$. The forward reaction rate $k_{\rightarrow}$ is obtained under the ansatz of the rate equation:

$$\dot{P}_{\text{left}}(t) = -P_{\text{left}}(t)k_{\rightarrow} + (1 - P_{\text{left}}(t))k_{\leftarrow} \quad (13)$$

by obtaining the time-series $P_{\text{left}}(t) = \text{Tr}[\hat{P}_{\text{left}}\rho(t)]$ and solving for $k_{\rightarrow}$. The backwards reaction rate is denoted by $k_{\leftarrow}$ and its proportionality to $k_{\rightarrow}$ is linked to the equilibrium distribution of the reaction coordinate $\frac{k_{\rightarrow}}{k_{\leftarrow}} = \frac{P_{\text{right}}^{eq}}{P_{\text{left}}^{eq}}$, with e.g. $P_{\text{left}}^{eq} = \lim_{t \rightarrow \infty} P_{\text{left}}(t) = \frac{\text{Tr}[\hat{P}_{\text{left}} e^{-\beta \hat{H}}]}{\text{Tr}[e^{-\beta \hat{H}}]}$ ⁴⁹. In the case of PESs with more than two wells, we define a projection operator for each well and solve for the reaction rate between neighboring wells. Further details are found in Supplementary Section S5.

In order to study the dynamics in more detail, we decompose the rate of the population of the vibrational states in terms of probability currents associated with the transitions from state n to state m ^{50,51}:

$$\dot{\rho}_{nn}(t) = - \sum_m j_{n \rightarrow m}(t) \quad (14)$$

We further decompose each current $j_{n \rightarrow m}$ in terms of the contributions from the system Hamiltonian and contributions from the harmonic environment. We will mainly use this diagnostic tool in the case of the simple spectral density Eq. 12, such that we make the distinction between the following three contributions:

$$j_{n \rightarrow m}(t) = j_{n \rightarrow m}^{\text{Tunneling}}(t) + j_{n \rightarrow m}^{\text{Solvent}}(t) + j_{n \rightarrow m}^{\text{Cavity}}(t) \quad (15)$$

This allows us to directly interpret how the cavity affects the quantum dynamics and furthermore the role of the tunneling dynamics and the phonon bath. We also note that the reaction rate can be related to the probability currents that go from the reactant region to the product region (Supplementary Section S6.1)

$$k_{\rightarrow} \approx \sum_n^{\text{reactant}} \sum_m^{\text{product}} j_{n \rightarrow m}(t) \quad (16)$$

This applies when the chosen basis is such, that the sum n runs over states localized in the left well and m indexes states localized the right well and when considering times t for which e.g. $P_{\text{left}}(t)$ has not changed considerably.

In the perturbative limit, the bath contributions to Eq. 14 can be reduced to the Pauli-equation

$$j_{n \rightarrow m}(t) \sim \Gamma_{n \rightarrow m} \rho_{nn}(t) - \Gamma_{m \rightarrow n} \rho_{mm}(t) \quad (17)$$

with the transition rates $\Gamma_{n \leftarrow m}$ given by Fermi's Golden Rule (Supplementary Section S6.2). These two reductions (Eq. 16 & 17) provide a simple perspective on how to relate the cavity modified noise spectrum to the cavity modified reaction rates, in a perturbative setting. Such a perspective has already been successfully applied in Refs. 33–36, 52, 53. For the definition of the currents (Eq. 15) in terms of the master equation presented in the next section (Eq. 18), see Supplementary Section S6.

III. METHODOLOGY

The time-series for Eq. 13 is obtained from a numerically exact simulation of the open quantum system dynamics using the methodology known under the names of HEOM⁵⁴, pseudomode^{55,56} and QD-MESS^{57–59}, among others^{60–63}. The main principle behind these methods is to use Markovian embeddings to emulate the effect of a macroscopic bath through a reduced, few-body harmonic environment.

For the simulations in this study, the equations of motions for the few-body system $\varrho(t)$ is:

$$\begin{aligned} \dot{\varrho}(t) = & -i \left[\hat{H}_R + \lambda \hat{R}^2, \varrho(t) \right] + \hat{L} \varrho(t) \hat{L}^\dagger - \frac{1}{2} \left\{ \hat{L}^\dagger \hat{L}, \varrho(t) \right\} \\ & + \sum_{k \in K} \left\{ -z_k \hat{a}_k^\dagger \hat{a}_k \varrho(t) - i \sqrt{d_k^*} \hat{R} \varrho(t) \hat{a}_k^\dagger \right. \\ & \left. - i \sqrt{d_k} \left(\hat{a}_k + \hat{a}_k^\dagger \right) \hat{R} \varrho(t) + H.C. \right\} \end{aligned} \quad (18)$$

where $H.C.$ refers to the hermitian conjugate of everything inside the brackets. $\hat{a}_k$ is the lowering operator of the k 'th auxiliary mode and z_k and d_k are complex parameters that characterizes the effective few-body harmonic environment. We have introduced the system jump operator $\hat{L}$, in order to more efficiently emulate the environment, as will be elaborated upon in the following. The so-called reorganization energy λ is given by:

$$\lambda = -\text{Im} \sum_{k \in K} \frac{d_k}{z_k} \quad (19)$$

and the initial state and system density matrix is given by:

$$\varrho(0) = \rho_S(0) \otimes \overbrace{|0\rangle\langle 0| \otimes \dots \otimes |0\rangle\langle 0|}^{|K| \text{ times}} =: \rho_S(0) \otimes^K |0\rangle\langle 0| \quad (20)$$

$$\rho_S(t) = \text{Tr}_K [\mathbb{1} \otimes^K |0\rangle\langle 0| \varrho(t)] \quad (21)$$

That is, the auxiliary modes are initialized in the vacuum and to obtain the reduced density matrix of the system $\rho_S(t)$, the auxiliary modes should be projected onto their vacuum. While K is the set that labels the effective bath modes of Eq. 18, consider an effective harmonic environment of K_{fit} harmonic oscillators, similar to those in Eq. 18. These give rise to a bath correlation function (BCF) that is a sum of complex exponentials⁵⁷. If this effective BCF resembles the original BCF:

$$C(t) \approx \sum_{k \in K_{\text{fit}}} d_k e^{-z_k t} \quad d_k, z_k \in \mathbb{C} \quad (22)$$

the few-body system successfully emulates the macroscopic environment with deviations in observables being controlled by the decomposition in Eq. 22^{64–66}, barring errors from numerically solving the master equation. Thus, the starting point of our method is to decompose the original BCF to numerical precision, as a sum of complex exponentials. This is achieved using the AAA algorithm^{67,68}. Details and general considerations regarding the method or fitting procedure can be found in Supplementary Section S4. Once the K_{fit} pairs of z_k and d_k have been obtained, we reduce K_{fit} to the smaller set K in the following manner.

1. Controlled compression of Markovian bath modes

Modern introductions to HEOM-type master equations, such as Eq. 18, generally do not include system jump operators $\hat{L}$ ^{54–59}. The inclusion of $\hat{L}$ in Eq. 18 represent our novel approach to compress the number of effective bath modes needed to accurately simulate the open system dynamics, in a controllable fashion. Essentially, some of the K_{fit} harmonic oscillators resemble a Markovian bath and simulating them as an explicit degree of freedom is not necessary. We distinguish our approach from earlier proposals to utilize adiabatic elimination^{59,69–71} to address the short-comings of HEOM in computationally challenging regimes. Here, we utilize the rigorous analysis underlying the Universal Lindblad Equation⁷², to accurately and controllably account for the Markovian bath modes via the system jump operator:

$$\hat{L} = \sum_{n,m} \sqrt{S_M(E_m - E_n)} R_{nm} |n\rangle\langle m| \quad (23)$$

$$S_M(\omega) = \sum_{k \in K_M} 2 \text{Re} \left(\frac{d_k}{z_k - i\omega} \right) \quad (24)$$

In Supplementary Section S4.2, we show how to control any error introduced by this modeling technique by using the rigorous error bound provided in Ref. 72. For our simulations, the calculation of $\hat{L}$ is based on the eigen-decomposition of $\hat{H}_R + \lambda \hat{R}^2 = \sum_n E_n |n\rangle\langle n|$ and K_M denote the set of auxiliary bath modes that are deemed to be Markovian *a priori*. For brevity, we quote the necessary but insufficient condition, that serves as a guiding principle for assigning bath modes to K_M :

$$\sum_{k \in K_M} \frac{|d_k|}{\text{Re}(z_k)^2} \ll 1 \quad (25)$$

The set $K = K_{\text{fit}} \setminus K_M$, used in Eq. 18, contains the remaining elements of K_{fit} when the set K_M has been removed. In terms of the BCF:

$$C(t) = \sum_{k \in K_{\text{fit}}} d_k e^{-z_k t} = \overbrace{\sum_{k \in K_M} d_k e^{-z_k t}}^{\text{Markovian bath}} + \overbrace{\sum_{k \in K} d_k e^{-z_k t}}^{\text{Residual bath}} \quad (26)$$

This allows us to fit the BCF to numerical precision, while compressing the Liouville space in Eq. 18 for the ensuing numerical simulations. We present the results for compressing the phonon bath in Supplementary Table I, where $K_{\text{fit}} = 19$ is reduced to $K = 2$ (1) with a relative error in the extracted reaction rate, for the given example, of 0.1% (1.2%) as ensured by the small value of the numerically calculated error bound. The dynamical simulations are carried out using the tensor network algorithm TDVP⁷³ implemented in ITensor^{74,75}.




FIG. 4. (a) The wavefunction of the localized states of the reaction coordinate visualized alongside the potential with the y-axis providing the energy difference to the ground state. The transitions between these states are labeled and visualized by arrows. (b) Reaction rate profile when varying the cavity frequency compared to the out-of-cavity reaction rate (horizontal dashed line). Due to the symmetry of the PES, the forward and backward reaction rates are equal $k := k_{\rightarrow} = k_{\leftarrow}$. (c) The probability current at $t = 6 \text{ps}$ for the labeled transitions as a function of the cavity frequency. The probability currents are decomposed in terms of contributions from the system Hamiltonian (tunneling) and from coupling to bath degrees of freedom (solvent and cavity).

IV. RESULTS AND DISCUSSIONS

The model for the physical system is primarily characterized by the two scalar functions $V(\hat{R})$, for the PES, and $J(\omega)$, for the effective spectral density. Therefore, the results section is divided into two parts, which aim to elucidate the role of each in the resulting reaction kinetics.

A. PES-driven effects

In the following, the environment will be assumed to consist of a phonon bath and a lossy cavity mode (Eq. 12). We utilize our parameterization of the PES to control and add additional transition frequencies to the reaction pathway. The probability currents between vibrational states, decomposed and assigned to their different contributors (Eq. 15), will then be used to analyze the dynamics.

1. Symmetric double well potential

We will begin by considering the simple symmetric double well, whose PES and vibrational states are shown in Fig. 4a. Outside the cavity, the reaction coordinate only interacts with the phonon bath. The strength of the coupling between the reaction coordinate and the solvent puts the system in the beginning of the energy-diffusion limited regime (Supplementary Section S10). Tunneling between the wells has almost been minimized

by the solvent and the thermal fluctuations it provides to the reaction is minimal.

Still, as seen in Fig. 4c, the reaction outside the cavity is mainly driven by tunneling. Notice that the reaction rate, denoted by the horizontal line in Fig. 4b, is approximately given by the net current going across the barrier (Eq. 16), which in this case is the sum of the probability currents for the T_0 and T_1 transitions. The transitions labeled a, b, c and d are included to provide the complete picture, despite their small values. Aside from diminishing tunneling, the thermal fluctuations of the phonon bath help the reaction coordinate transition over the potential energy barrier, which is captured by the transitions denoted by $T_{\uparrow}$ and $T_{\downarrow}$.

The cavity modified reaction rate is shown in Fig. 4b and the related probability currents are shown in Fig. 4c at different values of the cavity frequency, spanning resonant and off-resonant tunings. When resonant with the transition frequency of the wells $\omega_c = 1200 \text{cm}^{-1}$, the cavity is seen to greatly enhance both the excitation $T_{\uparrow}$ and relaxation $T_{\downarrow}$ processes. Furthermore, the tunneling through the top of the potential barrier T_1 is accordingly increased while the tunneling among the ground states T_0 is largely unaffected. Finally, when the cavity is tuned off resonance, its effect on the dynamics, as diagnosed by the probability currents, vanishes.

To elucidate these observations, we consider the time-scales of the system. Such an approach will also be helpful when we interpret the results of the coming sections. First, consider the timescale of the transition labeled T_1 . It must be much faster than T_0 since it constitutes tunneling the barrier at a point where it is narrower. Con-

sidering the transition T_0 itself provides a sizable portion of the reaction dynamics, despite its longer timescale, the transition T_1 must be far from being a rate-limiting step. Conversely, the rate at which spectral weight reaches this excited state manifold is much slower. To leading order, we can estimate the rate by Fermi's Golden Rule (Eq. 17) such that intra-well decay, due to the phonon bath, occurs at a rate $\sim \frac{2\lambda_R\gamma_R}{M\omega^2}$, while the excitation rate is related by detailed balance $\frac{2\lambda_R\gamma_R}{M\omega^2}e^{-\beta\omega}$ where ω is the frequency of the well⁷⁶. Thus, excitation is the rate-limiting step, whose rate for the given values is of the order⁷⁷ 10^{-6}fs^{-1} . As a result, once the cavity is resonant with the molecular transition, the rate is strongly enhanced due to the increased value of the noise spectrum at the transition frequency³³. Effectively, the lossy cavity provides an additional thermalization channel, in close analogy with the Purcell effect⁷⁸.

Having established that the probability-current analysis is well aligned with our understanding of this simple scenario, we will now focus on understanding how the complexity and asymmetries in the PES can affect the overall rate.

2. Asymmetry in reactant and product well frequencies

Figure 5 shows both the forward and backward reaction rate profiles for four double well PESs, where the transition frequency of the right well has been lowered when compared to the symmetric PES. The reaction rate outside the cavity is decreasing with the decreasing well frequency because the overlap between the ground states of each well becomes smaller.




FIG. 5. The bottom row shows four different potential energy surfaces, with transition frequencies in the right-well of around $\omega_r \in \{950, 700, 570, 500\} \times \text{cm}^{-1}$. The top row shows the reaction rate from the left well to the right well in blue and the converse in red, for varying ω_c , when coupled to the spectral density in Eq. 12. The horizontal dashed lines correspond to the reaction rates outside the cavity. The probability currents of the system with $\omega_r \approx 950\text{cm}^{-1}$ are shown in Fig. 6.

This decreases the tunneling rate, which was the main reaction pathway for the symmetric PES (Fig. 4c). In the most asymmetric PES we consider, with a right well frequency of $\omega_r \approx 500\text{cm}^{-1}$, tunneling along T_0 is practically zero and the only reaction mechanism is the phonon bath inducing the transition over the barrier (see Supplementary Fig. S8).

For all of the PESs, we see an asymmetry in the backward ($k_{\leftarrow}$ in red) vs. forward ($k_{\rightarrow}$ in blue) reaction rates. Their ratio is linked to the equilibrium populations in each well $\frac{k_{\rightarrow}}{k_{\leftarrow}} = \frac{P_{\text{right}}^{eq}}{P_{\text{left}}^{eq}}$ (Sec. IIB). The ratio persists when changing ω_c , indicating that the cavity does not modify the equilibrium distribution. We can compare $\frac{k_{\rightarrow}}{k_{\leftarrow}}$ to the theoretical prediction for the equilibrium state given by the *mean force Gibbs state*⁴⁹. To leading order in the bath coupling, the reduced density matrix for the system at equilibrium is the Gibbs state $\rho_S^{eq} \propto e^{-\beta H_R}$. The ground states in both wells are only approximately degenerate, with a slight energetic preference for the left well ($\Delta E \approx 20\text{cm}^{-1}$) and at $T = 300\text{K}$ (corresponding to $\approx 208.5\text{cm}^{-1}$) the population of the left well will be larger in the steady-state distribution, which puts $k_{\leftarrow}$ above $k_{\rightarrow}$ in magnitude. When more low-energy states are introduced in the right well, by lowering its frequency, the asymmetry decreases as seen and eventually P_{right}^{eq} will dominate, for which we also expect $k_{\rightarrow} \gg k_{\leftarrow}$. Going beyond leading order in the calculation of ρ_S^{eq} ⁷⁹ (Supplementary Section S8), we see a small correction to the Gibbs state, which biases the left well. This can be understood in terms of the bath observing the reaction coordinate, which localizes the wavefunction and favors dispensing more spectral weight in the deeper (left) well. This effect is mainly due to the phonon bath. The presence of the cavity has nearly no effect on the equilibrium distribution and the minor dependence on ω_c does not exhibit resonant behavior.




FIG. 6. The associated probability currents for both the $k_{\rightarrow}$ (top) and $k_{\leftarrow}$ (bottom) reaction dynamics of the least asymmetric system in figure 5. Probability currents for the other systems can be found in Supplementary Fig. S8.

The frequency of the right well ω_r represent a secondary relevant transition frequency, different from the

left well's frequency at $\omega_l \approx 1200\text{cm}^{-1}$. This shows up as a secondary peak in the reaction rate profiles located at ω_r , whose magnitude, however, quickly vanishes when decreasing ω_r . To explain this, we consider the reaction mechanism which goes over the barrier. Once the excited states are reached, the population inside this manifold will oscillate back and forth between the two wells, before decaying into either. Whether the excitation decays to the opposite well or returns to its original well determines whether a reaction takes place. Thus, tuning the cavity to match the frequency of a well, which provides an additional thermalization channel, will only have an effect on the reaction rate if it changes the rate-limiting step of the overall process. Excitations to the top of the potential energy barrier are still the rate-limiting step, so once the cavity is in resonance with one well, it means that the phonon-induced excitations of the other well becomes the rate-limiting step. Resultingly, the ratio between the magnitude of the primary peak and the secondary peak, relative to the baseline k_0 in the reaction rate profiles, are related by the timescales provided in Sec. IV A 1, such that for both the forward and backward reaction rates: $\frac{k(\omega_c=\omega_r)-k_0}{k(\omega_c=\omega_l)-k_0} \approx \frac{k_{\leftarrow}}{k_{\rightarrow}} \left(\frac{\omega_r}{\omega_l}\right)^2 e^{-\beta(\omega_l-\omega_r)}$ ⁸⁰. This explains the observed vanishing magnitude of the peak in k at ω_r when decreasing ω_r .

We also observe that the difference between the greatest cavity-enhanced reaction rate and the reaction rate outside the cavity increases when widening the right well. This is attributable to the increased exposure to the phonon bath at lower frequencies. Once the cavity help facilitate the large frequency transition, the phonon bath can more easily assist the transitions of the lower frequency well, leading to a larger relative enhancement of the reaction rate for decreasing ω_r .

Figure 6 shows the probability currents of the transitions labeled in Fig. 5. Note that the probability currents are, by definition, anti-symmetric. Thus, if for a single dynamical simulation, we compared $T_{\downarrow}$ to its reversed partner, we would find that they only differ by a sign. However, here we compare two distinct dynamical simulations, that differ by which well the reaction coordinate wavefunction was initialized in. The currents for all systems shown in Fig. 5 are provided in Supplementary Fig. S8.

Compared to the currents for the symmetric double well (Fig. 4c), the ground state tunneling rate T_0 is suppressed, as already noted. From the currents, it is also apparent that the transition at 1200cm^{-1} , denoted $T_{\uparrow}$, is the rate-limiting step. Again, the tunneling along T_1 occurs at a faster time scale such that the net contribution simply follows what the thermodynamics of the wells dictates. At $\omega_c = \omega_r$ ($\approx 950\text{cm}^{-1}$ for this PES) the cavity simply assumes the role of the phonon bath in enabling both excitation and relaxation transitions in the right well, as captured by $T_{\downarrow}$. How large a contribution either environmental dof provide to these transitions, is determined by the relative magnitude of their noise spectra

(Fig. 3), in this perturbative regime. The more the cavity noise spectrum overlaps with the $\omega_l = 1200\text{cm}^{-1}$ transition, the more the reaction dynamics are limited by the transitions in the right well, which occur on a faster time scale due to the $\sim \frac{1}{\omega^2}$ scaling of the approximate phonon-induced downward transition rate. This is apparent from the $T_{\downarrow}$ transitions in Fig. 6, whose solvent-induced component grows significantly when $\omega_c = 1200\text{cm}^{-1}$.

3. Competing Pathways

Finally, we add a third relevant transition frequency to the dynamics. This will serve as our simplified model of a reaction coordinate containing two different product regions and thus two competing pathways.

Figure 7a shows the PES and the associated vibrational states. We define the middle well to be the reactant region and the left and right well to be two distinct product regions. The three near-degenerate states at frequency 1570cm^{-1} are not eigenstates of H_R , but instead have been mixed to construct the localized states seen (Supplementary Section S3.3).

Outside the cavity, the reaction rate from the reactant to the right product region $k_{\rightarrow}$ is about $3.36\times$ larger than the reaction rate to the left product well $k_{\leftarrow}$, as seen in Fig. 7b. Informed by the probability currents in Fig. 7c, this difference in time scales can mainly be attributed to two factors. The right well ($\omega_r \approx 530\text{cm}^{-1}$) samples the phonon bath at a favorable frequency, leading to the increase in $T_{\rightarrow}$ over $T_{\leftarrow}$. Additionally, the right product well has an excited state, located in frequency just above 2000cm^{-1} , that shares support with the reactant excited state leading to the phonon-assisted transition labeled R_0 , adding to the $k_{\rightarrow}$ reaction rate. Notice that $k_{\leftarrow}$ is approximately given by the probability current of the $T_{\leftarrow}$ transition and $k_{\rightarrow}$ by the $T_{\rightarrow}$ and R_0 transitions, in accordance with Eq. 16.

Inside the cavity, there exists three distinct scenarios depending on which of the three wells the cavity is resonant with. Similar to the examples with double-well PESs, we generally observe that the cavity accelerates the dynamics in the resonant well and that, due to the shape of the phonon bath, the higher frequency well generally hosts the rate-limiting steps.

When the cavity is resonant with the right product well $\omega_c = \omega_r$ the reaction rate $k_{\rightarrow}$ is slightly increased ($\sim 1.12\times$) whereas $k_{\leftarrow}$ is unaffected. The probability currents reflect this observation, where the cavity is seen to help thermalize the R_{0-4} transitions. Instead, when the cavity is resonant with the frequency of the reactant region $\omega_c = \omega_0 \approx 800\text{cm}^{-1}$ both reaction rates naturally increase, leading to an enhancement in $k_{\leftarrow}$ by $\sim 1.2\times$ and in $k_{\rightarrow}$ by $\sim 1.25\times$ in line with the cavity enhanced dynamics of the $M_{0,1}$ transitions.

More interestingly, when the cavity is resonant with the left product well $\omega_c = \omega_l$, the rate $k_{\leftarrow}$ is increased by $\sim 2.6\times$ whereas $k_{\rightarrow}$ is decreased by $\sim 0.86\times$, making




FIG. 7. (a) Visualization of the triple well PES and the vibrational states. The middle well is the reactant region and $k_{\leftarrow}$ and $k_{\rightarrow}$ denote the reaction rate to the left and right product wells, respectively. The black arrows label the state-to-state transitions. (b) Shows the reaction rates at different cavity frequencies. The horizontal dashed lines denote the rates outside the cavity. (c) Probability currents for the associated transitions, both outside and inside the cavity. (d) The inferred kinetic yield of the left product in red and the right product in blue for different cavity frequencies. This does not constitute the yield at equilibrium, which remains largely unaffected by the presence of the cavity.

the two reaction rates almost equal. The increased rate of equilibration between the left well and the reactant well is expected, since the left-well frequency is resonant with the lossy cavity. The resulting cavity-induced bias toward the left well, reflected in the $T_{\leftarrow}$ and L_0 transitions, is accompanied by an increased excitation rate within the reactant well, as seen in the $M_{0,1}$ transitions. However, these effects do not fully compensate, so the excited reactant state feeding both product pathways is more depleted than without the cavity. This leads to the decrease in the R_{0-4} transitions and ultimately the $k_{\rightarrow}$ reaction rate.

The fact that the cavity can modify the two reaction rates differently can lead to cavity modified yield of the reaction. However, whether this occurs depends on the reversibility of the reaction. The system we have considered is best described as a reversible parallel reaction. The population in the middle well can flow into both the left and right well and the population in these outer wells can also flow back into the middle well. This reversibility allows the system to equilibrate and as such, the yield of the reaction is determined by the equilibrium distribution. Using the mean force Gibbs state^{49,79} already employed for the double well PESs (Sec. IV A 2) we can study how the cavity affects the equilibrium distribution. Similar to the double well examples, the solvent itself induces a small bias towards the more narrow and deeper left well but the cavity provides no noticeable effect (Supplementary Section S8.1). Thus, for the setup in question, the presence of the cavity does not change the yield of left and right product but it affects the speeds at which the reactant equilibrates with the two products.

A change in reaction rates can still induce a change in the yield, if the reaction is more accurately described as

an irreversible parallel reaction. This is the case, for example, if the left and right product wells are much lower in energy than the reactant well, when compared to the temperature. Then the backwards flow from the outer wells to the middle well would be negligible and the yield would be given by the branching ratios of the two forward reaction rates $P_{\text{left}}^{eq} = \frac{k_{\leftarrow}}{k_{\leftarrow} + k_{\rightarrow}}$ and $P_{\text{right}}^{eq} = \frac{k_{\rightarrow}}{k_{\leftarrow} + k_{\rightarrow}}$ (see Supplementary Section S9 for a mathematical formulation of this discussion). Thomas et al.⁹ provided the first experimental observation of such cavity-modified yield under kinetic control. Our model reproduces this effect, as seen in Fig. 7d, where the branching ratio is changed from being around 1 : 3 to 1 : 1 when the cavity is resonant with the frequency of the left well⁸¹. Thus, the cavity can change product yields under kinetic control, by modifying the relative rates of competing pathways, but it does not significantly alter the equilibrium product distribution for the thermodynamically controlled reversible reaction considered here.

B. Environment-driven effects

All calculations in the preceding section employed the simple spectral density given in Eq. 12. We now turn our attention to the general spectral density, Eq. 7, which describes a reaction coordinate interacting with a spectator mode, both of which are coupled to the cavity mode. Simultaneously, we simplify the system by considering the symmetric double well seen in Fig. 4a.

How strongly the cavity couples to either the reaction coordinate η_R or the spectator mode η_Q depends on the alignment between the respective dipole moments and the cavity polarization. We parameterize these couplings

in terms of two angles:

$$\eta_Q \propto \cos(\phi) \quad \text{and} \quad \eta_R \propto \cos(\theta - \phi) \quad (27)$$

where ϕ is the angle between the cavity polarization and the spectator mode's dipole moment and θ is the angle between the reaction-coordinate and spectator-mode dipoles. This parameterization constitutes a reduced 2-dimensional orientational model in which the two molecular dipoles rotate rigidly, at a fixed relative angle θ , within a plane containing the cavity polarization. To approximate the response of an ensemble of molecular orientations, we first calculate the orientationally averaged spectral density $\langle J(\omega) \rangle_\phi := \frac{1}{2\pi} \int_0^{2\pi} J(\omega) d\phi$ and then use this averaged spectral density in the dynamical simulation.




FIG. 8. Visualization of the noise spectra with spectral density given by Eq. 7, without the phonon bath, which has been averaged over the orientation of the molecule. The spectator mode frequency is $\omega_Q = 1170\text{cm}^{-1}$ with a lifetime of about $\sim 500\text{fs}$. θ is the angle between the reaction coordinate and spectator mode dipole moments. Five different values of θ is shown vertically, while two different values of ω_c are shown horizontally.

These two simplifications allow us to isolate how the relative geometry of the two vibrational dipoles controls the effect of the cavity on the reaction rate. Their validity in the parameter regime considered here is assessed in Supplementary Section S7, by comparing the reduced orientational model with a 3-dimensional isotropic treatment and with an explicit account of orientation-resolved reaction rates. These comparisons show that the reduced model retains the qualitative features of the cavity-induced rate changes. In particular, the rate extracted from dynamics using $\langle J(\omega) \rangle_\phi$ closely approximates the

effective rate obtained by fitting the ensemble reactant decay to a single exponential. Likewise, the 2- and 3-dimensional orientational models yield similar results in this regime, supporting the use of the reduced model to examine the dependence on relative dipole geometry. The resulting noise spectra are visualized in Fig. 8, with explicit expressions found in Supplementary Section S2.2.

We consider a spectator mode with frequency $\omega_Q = 1170\text{cm}^{-1}$, slightly below the relevant system transition at around 1200cm^{-1} . Outside the cavity, the damped spectator mode produces a Lorentzian feature in the noise spectrum. Introducing the cavity hybridizes the cavity and spectator mode, giving rise to features associated with two Rabi-split damped resonances distanced $\Omega_R \sim 200\text{cm}^{-1}$ apart. Since the reaction coordinate also couples directly to the cavity, the resulting spectrum develops asymmetric and non-Lorentzian features that may be understood as reflecting interference between the two hybridized decay channels. This structure depends on θ , which controls the relative signs and magnitudes of the corresponding dipole couplings.




FIG. 9. Numerically exact reaction rates compared to predictions calculated via Eq. 28. The Lorentzian has parameters $\omega_0 = 1205\text{cm}^{-1}$ and $\Gamma = 30\text{cm}^{-1}$ and the approximate scaling factor was rescaled $\frac{1}{2} \rightarrow 0.38$ to best match the results from HEOM. The horizontal dashed line correspond to the reaction rate outside the cavity.

Using our method described in Sec. III, we decompose, to great accuracy, the noise spectra in Fig. 8, for different cavity frequencies, and perform simulations of the dynamics in order to extract the reaction rate $k := k_{\rightarrow} = k_{\leftarrow}$. Figure 9 shows k as a function of ω_c for different angles θ ⁸². The black horizontal line denotes the reaction rate outside the cavity, while the circles show the cavity-modified rates. The solid lines represent the perturbative predictions obtained from the probability currents. We now derive the corresponding closed expression for k , drawing on the analysis of Fig. 4.

According to Eq. 16, the reaction rate can be ap-

proximated by the total probability current across the potential barrier, $k \approx j_{T_0} + j_{T_1}$. The currents shown in Fig. 4c suggest that j_{T_0} is largely unaffected by the cavity, whereas the cavity-induced change in j_{T_1} originates from the cavity contribution to the $T_{\uparrow,\downarrow}$ transitions. We may therefore approximate the cavity-modified reaction rate as the rate independent of the cavity, k_0 , plus the cavity dependent contribution to these transitions, e.g. $k \approx k_0 + j_{T_{\uparrow}}^{(c)}$. In the perturbative limit, Eq. 17 allows the cavity contribution to be evaluated as $j_{T_{\uparrow}}^{(c)} \approx \Gamma_{\uparrow}\rho_{gg} - \Gamma_{\downarrow}\rho_{ee}$ where the rates $\Gamma_{\uparrow,\downarrow}$ are given by Fermi's Golden Rule and the subscripts in ρ_{gg} and ρ_{ee} refer to the reactant ground and excited state, respectively. Despite the small value of ρ_{ee} the product $\Gamma_{\downarrow}\rho_{ee}$ can be sizeable. Thus, we account for ρ_{ee} through adiabatic elimination. As alluded to in the earlier discussions, the dynamics among the excited vibrational states is fast, so an excitation coming from the reactant ground state can decay into either the product or reactant well. Schematically, $\dot{\rho}_{ee} \approx \Gamma_{\uparrow}\rho_{gg} - 2\Gamma_{\downarrow}\rho_{ee}$ and the intermediate steady state $\rho_{ee} \approx \frac{\Gamma_{\uparrow}}{2\Gamma_{\downarrow}}\rho_{gg}$ is quickly reached before back-excitations from the product ground state needs to be accounted for. Consequently, the apparent rate given by FGR is rescaled by a factor-half $j_{T_{\uparrow}}^{(c)} \approx \frac{\Gamma_{\uparrow}}{2}\rho_{gg}$. Together with $\rho_{gg} \approx 1$, we arrive at the following expression for the cavity modified reaction rate:

$$k \approx k_0 + \frac{1}{2}|\langle e|\hat{R}|g\rangle|^2 \int_{-\infty}^{\infty} e^{-\beta\omega} S_{\beta}^{(c)}(\omega) \mathcal{A}(\omega) d\omega \quad (28)$$

$S_{\beta}^{(c)}(\omega)$ refers to the cavity-modified component of the noise spectrum and $\mathcal{A}(\omega) = \frac{1}{\pi} \frac{\Gamma}{(\omega - \omega_0)^2 + \Gamma^2}$ encodes the broadening of the transition frequency due to the cavity-independent component, in this case the phonon bath³³ (see also Supplementary Section S6.2.1). $\langle e|\hat{R}|g\rangle$ is the matrix element that couples the ground and excited state of the reactant well and the factor $\frac{1}{2}$ is the approximate rescaling due to the decay of the excited states, as described above. Such expressions for the reaction rate were successfully employed in References 33,35, for the case of no spectator mode ($C_Q = \eta_Q = 0$) and in Ref. 36, for no direct coupling between the reaction coordinate and the cavity ($\eta_R = 0$), with empirical rescaling factors introduced to obtain quantitative agreement with numerically exact results. Here, we have formulated a rationale in terms of probability currents, including a possible explanation for the required rescaling coefficient, and applied it to the case where all couplings are non-zero.

As seen in Fig. 9, the FGR predictions agree well with the HEOM simulations. Such agreement is not expected in general, since the assumptions underlying FGR require sufficiently weak bath coupling and a sufficiently smooth noise spectrum for the Born–Markov approximation to be applicable. Furthermore, the agreement in Fig. 9 requires an empirical rescaling of the cavity-induced FGR contribution. For Eq. 28 to be applicable in the first place, the system must also contain a well-defined rate-

limiting step that can be identified.

Nonetheless, the qualitative form of the rate profiles in Fig. 9 is governed by which parts of the structured noise spectrum overlap with the broadened transition function $\mathcal{A}(\omega)$. If the spectator mode is far detuned from the relevant transition, $|\omega_Q - \omega_0| \gg \Gamma$, the two hybridized channels do not contribute comparably within the transition window, and the asymmetric interference feature becomes less pronounced. The rate profile then approaches a simpler resonant cavity response, although its amplitude and specific shape can still depend on θ when the cavity is tuned sufficiently close to the spectator mode to hybridize appreciably, $|\omega_c - \omega_Q| \sim \Omega_R \sim 200\text{cm}^{-1}$. This residual angular dependence reflects the way the dipole geometry redistributes spectral weight between the hybridized resonances.

When the spectator mode lies within the broadened transition window, $|\omega_Q - \omega_0| \lesssim \Gamma$, both cavity-spectator hybrid modes can contribute to the same rate-limiting transition, making the overlap with $\mathcal{A}(\omega)$ sensitive to interference between the two channels.

This interpretation is made more concrete by the fact that the rate curves in Fig. 9 permit a unified phenomenological description in terms of a Fano-plus-Lorentzian profile⁸³. The Fano component captures the interference between competing hybridized channels within the broadened transition window³⁸, while the Lorentzian component represents the remaining resonant contribution from the spectator mode's residual spectral weight. In this form, the angle θ tunes the interference term through the Fano-parameter in addition to the magnitude of the residual Lorentzian contribution.

The rate profiles in Fig. 9 are therefore organized by an effective Fano-plus-Lorentzian structure, reflecting the interplay between cavity-spectator hybridization, spectral overlap with the broadened transition, and geometry-dependent interference.

V. CONCLUSION

We have developed a general quantum-dynamical framework for understanding cavity-modified reaction kinetics in terms of the individual transitions that comprise a reaction pathway. The framework combines a concise expression for the effective spectral density of an arbitrary harmonic environment with a controlled compression of Markovian auxiliary modes into system-level jump operators, enabling efficient tensor-network simulations of HEOM dynamics while retaining a physically transparent description of the environment.

The probability-current decomposition provides the central diagnostic for connecting these dynamics to observable reaction rates. It shows that cavity resonance selectively modifies particular transitions and that the impact on the overall reaction depends on their role in the reaction network. In systems with multiple transition frequencies, this distinction explains why resonances can

differ greatly in their effect on the total rate and, in competing pathways, can redistribute flux between products.

For the competing pathways, the cavity modifies the rates of population transfer and hence the timescale of equilibration between the wells, but does not modify the equilibrium distribution itself in the parameter regime considered here. The latter is governed by the mean force Gibbs state, whereas the cavity-induced changes in the transition rates redistribute flux along the way to equilibrium. Under irreversible conditions, where the final product distribution is instead determined kinetically by the relative pathway rates, this redistribution can translate directly into a cavity-modified product yield.

Finally, when spectator modes are included, hybridization and interference between environmental decay channels produce asymmetric, structured rate profiles that can be described in terms of Fano-plus-Lorentzian features. Taken together, these results show that cavity-modified reactivity is governed by how the cavity reshapes transition-resolved flux within a reaction network, rather than by a uniform modification of the reaction as a whole.

An important direction for future work is to extend this picture beyond the perturbative regime explored here. In the present calculations, the probability currents are consistent with the transition-rate intuition provided by Fermi's Golden Rule, while being obtained directly from the full HEOM dynamics rather than relying on a perturbative rate description. This makes the current-based analysis particularly promising for regimes where such rate theories become inadequate, as it provides a direct way to resolve how environmental effects propagate through the underlying reaction pathways. Extending the approach to more complex potential-energy landscapes, stronger system–environment and light–matter couplings, and more realistic models of the electromagnetic environment will help establish which of the mechanisms identified here persist beyond the present model systems. Importantly, the combination of general harmonic–environment representations, Markovian-mode compression, and tensor-network HEOM dynamics developed here provides a practical foundation for carrying out this analysis in increasingly complex settings.

ACKNOWLEDGMENTS

JV gratefully acknowledges the support of the Flatiron Institute during the Pre-Doctoral Researcher Program. The Flatiron Institute is a division of the Simons Foundation. JV also acknowledge support from the Carlsberg Foundation, and the Novo Nordisk Foundation, Grant No. NNF22SA0081175, Quantum Computing Programme.

The data and the code for reproducing the figures can be accessed on GitHub⁸⁴

- ¹H. Xu, A. Baydin, Q. Yi, I.-T. Lu, N. Zhu, T. E. Kritzell, J. Doumani, D. Kim, F. Tay, A. Rubio, and J. Kono, "Vacuum-dressed superconductivity in NbN observed in a high- q terahertz cavity," (2026).
- ²I. Keren, T. A. Webb, S. Zhang, J. Xu, D. Sun, B. S. Y. Kim, D. Shin, S. S. Zhang, J. Zhang, G. Pereira, J. Yao, T. Okugawa, M. H. Michael, E. Viñas Boström, J. H. Edgar, S. Wolf, M. Julian, R. P. Prasankumar, K. Miyagawa, K. Kanoda, G. Gu, M. Cothrine, D. Mandrus, M. Buzzi, A. Cavalleri, C. R. Dean, D. M. Kennes, A. J. Millis, Q. Li, M. A. Sentef, A. Rubio, A. N. Pasupathy, and D. N. Basov, "Cavity-altered superconductivity," *Nature* **650**, 864–868 (2026).
- ³T. W. Ebbesen, A. Rubio, and G. D. Scholes, "Introduction: Polaritonic chemistry," *Chemical Reviews* **123**, 12037–12038 (2023).
- ⁴R. Bhuyan, J. Mony, O. Kotov, G. W. Castellanos, J. Gómez Rivas, T. O. Shegai, and K. Börjesson, "The rise and current status of polaritonic photochemistry and photophysics," *Chemical Reviews* **123**, 10877–10919 (2023), pMID: 37683254.
- ⁵B. Xiang and W. Xiong, "Molecular polaritons for chemistry, photonics and quantum technologies," *Chemical Reviews* **124**, 2512–2552 (2024).
- ⁶J. A. Hutchison, T. Schwartz, C. Genet, E. Devaux, and T. W. Ebbesen, "Modifying chemical landscapes by coupling to vacuum fields," *Angewandte Chemie International Edition* **51**, 1592–1596 (2012).
- ⁷A. Shalabney, J. George, J. Hutchison, G. Pupillo, C. Genet, and T. W. Ebbesen, "Coherent coupling of molecular resonators with a microcavity mode," *Nat. Commun.* **6**, 5981 (2015).
- ⁸A. Thomas, J. George, A. Shalabney, M. Dryzhakov, S. J. Varma, J. Moran, T. Chervy, X. Zhong, E. Devaux, C. Genet, J. A. Hutchison, and T. W. Ebbesen, "Ground-state chemical reactivity under vibrational coupling to the vacuum electromagnetic field," *Angewandte Chemie International Edition* **55**, 11462–11466 (2016).
- ⁹A. Thomas, L. Lethuillier-Karl, K. Nagarajan, R. M. A. Vergauwe, J. George, T. Chervy, A. Shalabney, E. Devaux, C. Genet, J. Moran, and T. W. Ebbesen, "Tilting a ground-state reactivity landscape by vibrational strong coupling," *Science* **363**, 615–619 (2019).
- ¹⁰J. Lather, P. Bhatt, A. Thomas, T. W. Ebbesen, and J. George, "Cavity catalysis by cooperative vibrational strong coupling of reactant and solvent molecules," *Angewandte Chemie International Edition* **58**, 10635–10638 (2019).
- ¹¹A. Thomas, A. Jayachandran, L. Lethuillier-Karl, R. M. Vergauwe, K. Nagarajan, E. Devaux, C. Genet, J. Moran, and T. W. Ebbesen, "Ground state chemistry under vibrational strong coupling: dependence of thermodynamic parameters on the rabi splitting energy," *Nanophotonics* **9**, 249–255 (2020).
- ¹²W. Ahn, J. F. Triana, F. Recabal, F. Herrera, and B. S. Simpkins, "Modification of ground-state chemical reactivity via light–matter coherence in infrared cavities," *Science* **380**, 1165–1168 (2023).
- ¹³J. Lather, A. N. K. Thabassum, J. Singh, and J. George, "Cavity catalysis: modifying linear free-energy relationship under cooperative vibrational strong coupling," *Chemical Science* **13**, 195–202 (2022).
- ¹⁴F. Verdelli, Y.-C. Wei, K. Joseph, M. S. Abdelkhalik, M. Goudarzi, S. H. C. Askes, A. Baldi, E. W. Meijer, and J. Gomez Rivas, "Polaritonic chemistry enabled by non-local metasurfaces," *Angewandte Chemie International Edition* **63**, e202409528 (2024).
- ¹⁵D. M. Coles, N. Somaschi, P. Michetti, C. Clark, P. G. Lagoudakis, P. G. Savvidis, and D. G. Lidzey, "Polariton-mediated energy transfer between organic dyes in a strongly coupled optical microcavity," *Nature Materials* **13**, 712–719 (2014).
- ¹⁶X. Zhong, T. Chervy, L. Zhang, A. Thomas, J. George, C. Genet, J. A. Hutchison, and T. W. Ebbesen, "Energy transfer between spatially separated entangled molecules," *Angewandte Chemie International Edition* **56**, 9034–9038 (2017).

- ¹⁷S. Hou, M. Khatoniar, K. Ding, Y. Qu, A. Napolov, V. M. Menon, and S. R. Forrest, "Ultralong-range energy transport in a disordered organic semiconductor at room temperature via coherent exciton-polariton propagation," *Advanced Materials* **32**, 2002127 (2020).
- ¹⁸B. Munkhbat, M. Wersäll, D. G. Baranov, T. J. Antosiewicz, and T. Shegai, "Suppression of photo-oxidation of organic chromophores by strong coupling to plasmonic nanoantennas," *Science Advances* **4**, eaas9552 (2018).
- ¹⁹S. Satapathy, M. Khatoniar, D. K. Parappuram, B. Liu, G. John, J. Feist, F. J. Garcia-Vidal, and V. M. Menon, "Selective isomer emission via funneling of exciton polaritons," *Science Advances* **7**, eabj0997 (2021).
- ²⁰J. Singh, P. Garg, J. Lather, and J. George, "Stirring chemical reaction landscapes through strong light-matter interactions," *Chemical Communications* **61**, 18289–18301 (2025).
- ²¹K. Hirai, R. Takeda, J. A. Hutchison, and H. Uji-i, "Modulation of Prins cyclization by vibrational strong coupling," *Angewandte Chemie International Edition* **59**, 5332–5335 (2020).
- ²²G. D. Wiesehan and W. Xiong, "Negligible rate enhancement from reported cooperative vibrational strong coupling catalysis," *The Journal of Chemical Physics* **155**, 241103 (2021).
- ²³R. M. A. Vergauwe, J. J. Toppari, and G. Groenhof, "Toward reliable spectroscopic analysis of reaction kinetics in polaritonic chemistry," *Nanophotonics* **15**, e70143 (2026).
- ²⁴A. Sau, K. Nagarajan, B. Patrahaui, L. Lethuillier-Karl, R. M. A. Vergauwe, A. Thomas, J. Moran, C. Genet, and T. W. Ebbesen, "Modifying Woodward-Hoffmann stereoselectivity under vibrational strong coupling," *Angewandte Chemie International Edition* **60**, 5712–5717 (2021).
- ²⁵Y. Pang, A. Thomas, K. Nagarajan, R. M. A. Vergauwe, K. Joseph, B. Patrahaui, K. Wang, C. Genet, and T. W. Ebbesen, "On the role of symmetry in vibrational strong coupling: The case of charge-transfer complexation," *Angewandte Chemie International Edition* **59**, 10436–10440 (2020).
- ²⁶D. G. Truhlar, B. C. Garrett, and S. J. Klippenstein, "Current status of transition-state theory," *J. Phys. Chem.* **100**, 12771–12800 (1996).
- ²⁷P. Hänggi, P. Talkner, and M. Borkovec, "Reaction-rate theory: fifty years after Kramers," *Rev. Mod. Phys.* **62**, 251–341 (1990).
- ²⁸X. Li, A. Mandal, and P. Huo, "Cavity frequency-dependent theory for vibrational polariton chemistry," *Nat. Commun.* **12**, 1315 (2021).
- ²⁹L. P. Lindoy, A. Mandal, and D. R. Reichman, "Resonant cavity modification of ground-state chemical kinetics," *J. Phys. Chem. Lett.* **13**, 6580–6586 (2022).
- ³⁰A. Mandal, M. A. Taylor, B. M. Weight, E. R. Koessler, X. Li, and P. Huo, "Theoretical advances in polariton chemistry and molecular cavity quantum electrodynamics," *Chemical Reviews* **123**, 9786–9879 (2023).
- ³¹L. P. Lindoy, A. Mandal, and D. R. Reichman, "Quantum dynamical effects of vibrational strong coupling in chemical reactivity," *Nature Communications* **14**, 2733 (2023).
- ³²Y. Ke and J. O. Richardson, "Quantum nature of reactivity modification in vibrational polariton chemistry," *The Journal of Chemical Physics* **161**, 054104 (2024).
- ³³W. Ying and P. Huo, "Resonance theory and quantum dynamics simulations of vibrational polariton chemistry," *The Journal of Chemical Physics* **159**, 084104 (2023).
- ³⁴W. Ying, M. A. D. Taylor, and P. Huo, "Resonance theory of vibrational polariton chemistry at the normal incidence," *Nanophotonics* **13**, 2601–2615 (2024).
- ³⁵W. Ying and P. Huo, "Resonance theory of vibrational strong coupling enhanced polariton chemistry and the role of photonic mode lifetime," *Communications Materials* **5**, 110 (2024).
- ³⁶S. M. Vega, W. Ying, and P. Huo, "Theoretical insights into the resonant suppression effect in vibrational polariton chemistry," *Journal of the American Chemical Society* **147**, 19727–19737 (2025).
- ³⁷Y. Ke, "Stochastic resonance in vibrational polariton chemistry," *The Journal of Chemical Physics* **162**, 064702 (2025).
- ³⁸Y. Ke, "Cooperative chemical reactions in optical cavities: A complex interplay of mode hybridization, timescale balance, and pathway interference," *The Journal of Chemical Physics* **164**, 174103 (2026).
- ³⁹L. P. Lindoy, A. Mandal, and D. R. Reichman, "Investigating the collective nature of cavity-modified chemical kinetics under vibrational strong coupling," *Nanophotonics* **13**, 2617–2633 (2024).
- ⁴⁰W. Ying, M. E. Mondal, E. R. Koessler, S. M. Vega, and P. Huo, "Collective effects in polariton chemistry and photophysics," *Annual Review of Physical Chemistry* **77**, 201–224 (2026).
- ⁴¹K. Sun and R. F. Ribeiro, "Theoretical formulation of chemical equilibrium under vibrational strong coupling," *Nat. Commun.* **15**, 2405 (2024).
- ⁴²M. Ruggenthaler, D. Sidler, and A. Rubio, "Understanding polaritonic chemistry from ab initio quantum electrodynamics," *Chemical Reviews* **123**, 11191–11229 (2023).
- ⁴³A. Caldeira and A. Leggett, "Quantum tunnelling in a dissipative system," *Annals of Physics* **149**, 374–456 (1983).
- ⁴⁴R. Feynman and F. Vernon, "The theory of a general quantum system interacting with a linear dissipative system," *Annals of Physics* **24**, 118–173 (1963).
- ⁴⁵H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, 2007).
- ⁴⁶A. Garg, J. N. Onuchic, and V. Ambegaokar, "Effect of friction on electron transfer in biomolecules," *The Journal of Chemical Physics* **83**, 4491–4503 (1985).
- ⁴⁷W. H. Miller, S. D. Schwartz, and J. W. Tromp, "Quantum mechanical rate constants for bimolecular reactions," *The Journal of Chemical Physics* **79**, 4889–4898 (1983).
- ⁴⁸I. R. Craig, M. Thoss, and H. Wang, "Proton transfer reactions in model condensed-phase environments: Accurate quantum dynamics using the multilayer multiconfiguration time-dependent hartree approach," *The Journal of Chemical Physics* **127**, 144503 (2007).
- ⁴⁹A. S. Trushechkin, M. Merkli, J. D. Cresser, and J. Anders, "Open quantum system dynamics and the mean force gibbs state," *AVS Quantum Science* **4**, 012301 (2022).
- ⁵⁰J. J. J. Roden and K. B. Whaley, "Probability-current analysis of energy transport in open quantum systems," *Phys. Rev. E* **93**, 012128 (2016).
- ⁵¹B. Schumacher, M. D. Westmoreland, A. New, and H. Qiao, "Probability current and thermodynamics of open quantum systems," (2016), arXiv:1607.01331.
- ⁵²J. E. Lawrence, T. Fletcher, L. P. Lindoy, and D. E. Manolopoulos, "On the calculation of quantum mechanical electron transfer rates," *The Journal of Chemical Physics* **151**, 114119 (2019).
- ⁵³Y. Lai, W. Ying, and P. Huo, "Non-equilibrium rate theory for polariton relaxation dynamics," *The Journal of Chemical Physics* **161**, 104109 (2024).
- ⁵⁴Y. Tanimura, "Numerically "exact" approach to open quantum dynamics: The hierarchical equations of motion (HEOM)," *The Journal of Chemical Physics* **153**, 020901 (2020).
- ⁵⁵G. Pleasance, B. M. Garraway, and F. Petruccione, "Generalized theory of pseudomodes for exact descriptions of non-markovian quantum processes," *Phys. Rev. Res.* **2**, 043058 (2020).
- ⁵⁶G. Park, Z. Huang, Y. Zhu, C. Yang, G. K.-L. Chan, and L. Lin, "Quasi-Lindblad pseudomode theory for open quantum systems," *Phys. Rev. B* **110**, 195148 (2024).
- ⁵⁷M. Xu, Y. Yan, Q. Shi, J. Ankerhold, and J. T. Stockburger, "Taming quantum noise for efficient low temperature simulations of open quantum systems," *Phys. Rev. Lett.* **129**, 230601 (2022).
- ⁵⁸M. Xu, V. Vadimov, M. Krug, J. T. Stockburger, and J. Ankerhold, "A universal framework for quantum dissipation: minimally extended state space and exact time-local dynamics," (2023), arXiv:2307.16790.
- ⁵⁹M. Xu, V. Vadimov, J. T. Stockburger, and J. Ankerhold, "Colloquium: Simulating non-markovian dynamics in open quantum

- systems,” *Rev. Mod. Phys.* **98**, 021002 (2026).
- ⁶⁰K. H. Hughes, C. D. Christ, and I. Burghardt, “Effective-mode representation of non-markovian dynamics: A hierarchical approximation of the spectral density. i. application to single surface dynamics,” *The Journal of Chemical Physics* **131**, 024109 (2009).
- ⁶¹A. W. Chin, Á. Rivas, S. F. Huelga, and M. B. Plenio, “Exact mapping between system-reservoir quantum models and semi-infinite discrete chains using orthogonal polynomials,” *Journal of Mathematical Physics* **51**, 092109 (2010).
- ⁶²M. P. Woods, R. Groux, A. W. Chin, S. F. Huelga, and M. B. Plenio, “Mappings of open quantum systems onto chain representations and markovian embeddings,” *Journal of Mathematical Physics* **55**, 032101 (2014).
- ⁶³A. Nazir and G. Schaller, “The reaction coordinate mapping in quantum thermodynamics,” in *Thermodynamics in the Quantum Regime: Fundamental Aspects and New Directions*, edited by F. Binder, L. A. Correa, C. Gogolin, J. Anders, and G. Adesso (Springer International Publishing, Cham, 2018) pp. 551–577.
- ⁶⁴F. Mascherpa, A. Smirne, S. F. Huelga, and M. B. Plenio, “Open systems with error bounds: Spin-boson model with spectral density variations,” *Phys. Rev. Lett.* **118**, 100401 (2017).
- ⁶⁵Z. Huang, Y. Zhu, G. Park, and L. Lin, “Unified error bounds for perturbations of non-markovian open quantum systems in gaussian environments,” (2025), arXiv:2411.08741v2.
- ⁶⁶K. Liu and J. Lu, “Error Bounds for Open Quantum Systems with Harmonic Bosonic Bath,” *Quantum* **9**, 1896 (2025).
- ⁶⁷T. A. Driscoll, Y. Nakatsukasa, and L. N. Trefethen, “AAA rational approximation on a continuum,” *SIAM Journal on Scientific Computing* **46**, A929–A952 (2024).
- ⁶⁸T. A. Driscoll, “Rational function approximation in julia,” (2025), doi: 10.5281/zenodo.8355791, version used: v0.1.4.
- ⁶⁹A. Ishizaki and Y. Tanimura, “Quantum dynamics of system strongly coupled to low-temperature colored noise bath: Reduced hierarchy equations approach,” *J. Phys. Soc. Jpn.* **74**, 3131–3134 (2005).
- ⁷⁰T. P. Fay, “A simple improved low temperature correction for the hierarchical equations of motion,” *J. Chem. Phys.* **157**, 054108 (2022).
- ⁷¹E. M. Kessler, “Generalized schrieffer-wolff formalism for dissipative systems,” *Phys. Rev. A* **86**, 012126 (2012).
- ⁷²F. Nathan and M. S. Rudner, “Universal Lindblad equation for open quantum systems,” *Phys. Rev. B* **102**, 115109 (2020).
- ⁷³J. Haegeman, J. I. Cirac, T. J. Osborne, I. Pizorn, H. Verschelde, and F. Verstraete, “Time-dependent variational principle for quantum lattices,” *Phys. Rev. Lett.* **107**, 070601 (2011).
- ⁷⁴M. Fishman, S. R. White, and E. M. Stoudenmire, “The ITensor Software Library for Tensor Network Calculations,” *SciPost Phys. Codebases*, 4 (2022).
- ⁷⁵M. Fishman, S. R. White, and E. M. Stoudenmire, “Codebase release 0.3 for ITensor,” *SciPost Phys. Codebases*, 4-r0.3 (2022).
- ⁷⁶At the relevant transition frequencies $\omega > \gamma_R = 200\text{cm}^{-1}$ the phonon bath spectral density becomes $\frac{2\lambda_R\gamma_R\omega}{\omega^2 + \gamma_R^2} \approx \frac{2\lambda_R\gamma_R}{\omega}$ and the transition matrix element within a well is approximated as $|\langle 1|\hat{R}|0\rangle|^2 \approx \frac{1}{2M\omega}$ with $|n\rangle$ the harmonic oscillator states. The noise spectrum evaluated at the transition frequency times the matrix element, yields the FGR rate which are the quoted expressions. Alternatively, see Supplementary Section S6.2.
- ⁷⁷ $\frac{\lambda\gamma}{\omega^2} e^{-\omega/T} \frac{2}{M} \approx \frac{83.6\text{cm}^{-1} \times 200\text{cm}^{-1}}{(1200\text{cm}^{-1})^2} e^{-1200\text{cm}^{-1}/208.5\text{cm}^{-1}} \times 0.045\text{fs}^{-1} \approx 1.66 \times 10^{-6}\text{fs}^{-1}$.
- ⁷⁸B. Metzger, E. Muller, J. Nishida, B. Pollard, M. Hentschel, and M. B. Raschke, “Purcell-enhanced spontaneous emission of molecular vibrations,” *Phys. Rev. Lett.* **123**, 153001 (2019).
- ⁷⁹J. D. Cresser and J. Anders, “Weak and ultrastrong coupling limits of the quantum mean force gibbs state,” *Phys. Rev. Lett.* **127**, 250601 (2021).
- ⁸⁰Using that $k_{\rightarrow} = \frac{P_{\text{right}}^{\text{eq}}}{P_{\text{left}}^{\text{eq}}} k_{\leftarrow}$, the relative increase in the rate $k_{\rightarrow}(\omega_c = \omega_l) - k_0$ can be related to the phonon-induced excitations in the right well $k_{\rightarrow}(\omega_c = \omega_l) - k_0 \sim \frac{P_{\text{right}}^{\text{eq}}}{P_{\text{left}}^{\text{eq}}} \frac{e^{-\beta\omega_r}}{\omega_l^2}$. Conversely, $k_{\rightarrow}(\omega_c = \omega_r) - k_0$ is related to the phonon-induced excitations in the left well, such that the ratio leads to the quoted result.
- ⁸¹Whether the cavity-modification is enhancive or suppressive in nature, is largely dependent on the harmonic environment³¹. Thus, if we instead considered the coupling between the reaction coordinate and the cavity to be mediated by a spectator mode³⁶, we would expect the presence of the cavity to have a suppressive effect, which could place our model more in line with the experimental study⁹. Additionally, the potential energy of the product regions could be lowered to incorporate the irreversible aspect of the reaction.
- ⁸²The HEOM results for some ω_c are omitted. This is because the automated AAA decomposition (Eq. 22) did not reach the specified accuracy of 10^{-10} . HEOM results for any ω_c and θ can still be obtained with our method, in some cases one would just have to tune the settings for the AAA-fit by hand or choose a different method to achieve the decomposition.
- ⁸³Specifically, we have in mind:
- $$\frac{k(\tilde{\omega}_c)}{k_0} \propto \frac{(a(\theta) + \tilde{\omega}_c)^2}{1 + \tilde{\omega}_c^2} + b(\theta) \frac{\tilde{\Gamma}}{(\tilde{\omega}_c + \tilde{\omega}_0)^2 + \tilde{\Gamma}^2}$$
- where $\tilde{\omega}_c$ is a dimensionless cavity frequency that has been shifted and scaled appropriately and $a(\theta)$ is the angle-dependent Fano-parameter. Reducing Eq. 28 to this form was not immediately successful because naive pole-expansion of the terms in Eq. S250 missed some of the observed features. Yet, this ansatz faithfully captures the rate profiles in Fig. 9.
- ⁸⁴GitHub repository: <https://github.com/Vinther901/Vinther2026dynamics>.
- ⁸⁵J. Williamson, “The exponential representation of canonical matrices,” *American Journal of Mathematics* **61**, 897–911 (1939).
- ⁸⁶F. Nicacio, “Williamson theorem in classical, quantum, and statistical physics,” *American Journal of Physics* **89**, 1139–1151 (2021).
- ⁸⁷D. T. Colbert and W. H. Miller, “A novel discrete variable representation for quantum mechanical reactive scattering via the s-matrix kohn method,” *The Journal of Chemical Physics* **96**, 1982–1991 (1992).
- ⁸⁸M. Krug and J. Stockburger, “On stability issues of the HEOM method,” *The European Physical Journal Special Topics* **232**, 3219–3226 (2023).
- ⁸⁹M. Sánchez-Barquilla and J. Feist, “Accurate truncations of chain mapping models for open quantum systems,” *Nanomaterials* **11** (2021), 10.3390/nano11082104.
- ⁹⁰L. K. Zhou, G. R. Jin, and W. Yang, “Systematic and efficient pseudomode method to simulate open quantum systems under a bosonic environment,” *Phys. Rev. A* **110**, 022221 (2024).
- ⁹¹Z. Huang, G. Park, G.-L. Chan, and L. Lin, “Coupled Lindblad pseudomode theory for simulating open quantum systems,” *Phys. Rev. Lett.* **136**, 090403 (2026).
- ⁹²F. Nathan and M. S. Rudner, “Erratum: Universal Lindblad equation for open quantum systems [phys. rev. b 102, 115109 (2020)],” *Phys. Rev. B* **104**, 119901 (2021).
- ⁹³J. Agerskov and F. Nathan, “Markovian quantum master equations are exponentially accurate in the weak coupling regime,” (2026), arXiv:2603.04504.
- ⁹⁴L. Chen and Q. Shi, “Quantum rate dynamics for proton transfer reactions in condensed phase: The exact hierarchical equations of motion approach,” *The Journal of Chemical Physics* **130**, 134505 (2009).
- ⁹⁵If the trace-preserving version of the master equation (Eq. S41) was instead used, then $\rho_k(t) = \text{Tr}_B[\hat{a}_k \rho(t)]$.
- ⁹⁶A. Plonka, *Dispersive Kinetics* (Springer Netherlands, 2001).
- ⁹⁷E. Anslyn and D. Dougherty, *Modern Physical Organic Chemistry*, G - Reference, Information and Interdisciplinary Subjects Series (University Science Books, 2006).

⁹⁸C. Zhang, E. Jeckelmann, and S. R. White, “Density matrix approach to local hilbert space reduction,” *Phys. Rev. Lett.* **80**, 2661–2664 (1998).

⁹⁹E. Jeckelmann and S. R. White, “Density-matrix renormalization-group study of the polaron problem in

the holstein model,” *Phys. Rev. B* **57**, 6376–6385 (1998).

¹⁰⁰M. S. Rudolph and J. Tindall, “Simulating and sampling from quantum circuits with 2d tensor networks,” (2025).

SUPPLEMENTARY INFORMATION

S1	Gaussian environments	15
S2	Basis-independent spectral density for general harmonic environments	16
S2.1	Spectral density of interacting lossy harmonic modes.	19
S2.2	Spectral density of interacting cavity and spectator mode, averaged over molecule orientation	21
S3	Reaction Coordinate Potentials & vibrational state details	21
S3.1	Double Well	22
S3.1.1	Derivation	22
S3.2	Triple Well	22
S3.3	Localized states	23
S3.4	Numerics	23
S4	Details on Hierarchical Equations Of Motion	24
S4.1	Decomposing the Bath Correlation Function.	25
S4.2	Effective Markovian correction	26
S5	Reaction Rate Analysis	27
S5.1	Triple well reaction rates	29
S6	Probability Currents: Transition Rate Between States	30
S6.1	Relating probability-currents to reaction rates.	31
S6.2	FGR as perturbative limit of probability currents.	31
S6.2.1	Broadening of transition frequencies due to phonon bath	32
S7	Distribution of reaction rates and comparison to isotropic averaging	33
S8	Equilibrium state of the reaction coordinate	33
S8.1	Mean force Gibbs state for the triple well, Fig. 7	35
S8.2	Mean force Gibbs state for the double wells, Fig. 5.	35
S9	Kinetic and thermodynamic control of the triple well system.	36
S10	The phonon bath and Kramers turnover	37
S10.1	Tensor network simulations in the strong coupling regime.	37
S11	Additional asymmetric double well probability currents	38
S12	Overview of parameter values	38

Supplemental Material S1: Gaussian environments

A generic starting point for considering a system coupled to a harmonic environment, is the mass-weighted Caldeira-Legget model of dissipation⁴³:

$$\hat{H} = \hat{H}_S + \frac{1}{2} \sum_j \left(\hat{p}_j^2 + \omega_j^2 \left(\hat{x}_j - \frac{c_j}{\omega_j^2} \hat{S} \right)^2 \right) \quad (\text{S11})$$

The coupling $\frac{\omega_j^2}{2} \left(\hat{x}_j - \frac{c_j}{\omega_j^2} \hat{S} \right)^2 = \frac{\omega_j^2}{2} \hat{x}_j^2 - c_j \hat{x}_j \hat{S} + \frac{c_j^2}{2\omega_j^2} \hat{S}^2$ gives rise to the reorganization term: $\sum_j \frac{c_j^2}{2\omega_j^2} \hat{S}^2 =: \lambda \hat{S}^2$. It is important, typically, to include this term as it counters unwarranted bath-induced renormalization of the system. For example, for two harmonic oscillators coupled bilinearly through their coordinates, omitting the reorganization term while holding the bare oscillator frequencies fixed causes one squared normal-mode frequency to become negative at

sufficiently strong coupling, signaling an instability. Including the reorganization term keeps the quadratic Hamiltonian positive definite for all finite coupling strengths. For brevity, we will often absorb the reorganization term into $\hat{H}_S$.

By assuming an initially factorized state and with the harmonic environment in a thermal equilibrium, one can formally write the system reduced density matrix⁴⁵:

$$\rho_S(t) = e^{-i[\hat{H}_S, \bullet]t} \mathcal{T}_S \exp \left[- \int_0^t dt_1 \int_0^{t_1} dt_2 \hat{\mathcal{K}}(t_1, t_2) \right] \rho_S(0) \quad (\text{S12})$$

with the influence superoperator:

$$\hat{\mathcal{K}}(t_1, t_2) \bullet = \left(\hat{S}(t_1) \bullet - \bullet \hat{S}(t_1) \right) \left(C(t_1 - t_2) \hat{S}(t_2) \bullet - \bullet \hat{S}(t_2) C^*(t_1 - t_2) \right) \quad (\text{S13})$$

where $C(t_1 - t_2)$ is the bath correlation function (BCF), $\hat{S}(t) = e^{i\hat{H}_S t} \hat{S} e^{-i\hat{H}_S t}$ the interaction picture system operator coupling to the bath and $\mathcal{T}_S$ the time-ordering operator. From the above, we see that the only information we need to model the environment is exactly the BCF. We may rewrite this in terms of the *spectral density* $J(\omega)$:

$$J(\omega) = \frac{\pi}{2} \sum_j \frac{c_j^2}{\omega_j} \delta(\omega - \omega_j) \quad (\text{S14})$$

such that:

$$\int_{-\infty}^{\infty} dt C(t) e^{i\omega t} = S_\beta(\omega) = \frac{2}{1 - e^{-\beta\omega}} (J(\omega) - J(-\omega)) \quad (\text{S15})$$

This expression can also be found in the literature, written with only one $J(\omega)$ with the understanding that $J(\omega)$ is already an odd function or else you employ the odd version $J(\omega) - J(-\omega)$.

Some remarks regarding the BCF and noise spectrum:

- $S_\beta(\omega)$ is real and positive and within Fermi's Golden Rule, the transition rate at frequency ω is proportional to $S_\beta(-\omega)$.
- $S_\beta(-\omega) = e^{-\beta\omega} S_\beta(\omega)$ known as the detailed balance condition.
- $C(t)$ is complex with:

$$C(t) = C^*(-t) \quad (\text{S16})$$

$$\text{Re}[C(t)] = \int_0^\infty \frac{d\omega}{\pi} \coth\left(\frac{\beta\omega}{2}\right) J(\omega) \cos(\omega t) \quad (\text{S17})$$

$$\text{Im}[C(t)] = - \int_0^\infty \frac{d\omega}{\pi} J(\omega) \sin(\omega t) \quad (\text{S18})$$

- The reorganization energy is given by $\sum_j \frac{c_j^2}{2\omega_j^2} = \int_0^\infty \frac{d\omega}{\pi} \frac{J(\omega)}{\omega} = -\text{Im} \int_0^\infty C(t) dt$.

Supplemental Material S2: Basis-independent spectral density for general harmonic environments

The above model considered a collection of uncoupled harmonic oscillators, each coupled to the system. However, it is of interest to consider more general microscopic models where the interactions can be arbitrary. This will allow us to describe the entire harmonic environment, including cavity modes, spectator modes and so on, in terms of a single spectral density (or more if more system operators are coupled to the environment). For now, we will absorb the reorganization term into $\hat{H}_S$ and write the system-bath Hamiltonian as:

$$\hat{\mathcal{H}} = \hat{H}_S + \hat{S} \mathbf{c}^T \hat{\mathbf{r}} + \frac{1}{2} \hat{\mathbf{r}}^T H \hat{\mathbf{r}} \quad (\text{S21})$$

where $\hat{\mathbf{r}}$ is a vector containing the $\hat{x}_j$'s and $\hat{p}_j$'s and H is a positive-definite symmetric matrix that contains the parameters for the bath modes. $\mathbf{c}$ is the vector of coupling constants. We will calculate the BCF, which in this notation, is:

$$C(t) = \text{Tr}[\mathbf{c}^T \hat{\mathbf{r}} e^{-\frac{it}{2} \hat{\mathbf{r}}^T H \hat{\mathbf{r}}} \mathbf{c}^T \hat{\mathbf{r}} e^{\frac{it}{2} \hat{\mathbf{r}}^T H \hat{\mathbf{r}}} \rho_B(0)] \quad (\text{S22})$$

$\rho_B(0) \propto e^{-\frac{\beta}{2}\hat{\mathbf{r}}^T H \hat{\mathbf{r}}}$ is the initial thermal state of the bath. The goal is to check whether the above permits the same expression for the noise spectrum in terms of a spectral density as the Caldeira-Leggett model and to find a practical expression in terms of H and $\mathbf{c}$. The time-dependence can be simplified by considering:

$$\frac{1}{2}[\mathbf{c}^T \hat{\mathbf{r}}, \hat{\mathbf{r}}^T H \hat{\mathbf{r}}] = \frac{1}{2}c_n H_{ml}(\hat{r}_n \hat{r}_m \hat{r}_l - \hat{r}_m \hat{r}_l \hat{r}_n) \quad (\text{S23})$$

$$= \frac{1}{2}c_n H_{ml}([\hat{r}_n, \hat{r}_m]\hat{r}_l - \hat{r}_m[\hat{r}_l, \hat{r}_n]) \quad (\text{S24})$$

$$= \frac{i}{2}\mathbf{c}^T \Omega H \hat{\mathbf{r}} - \frac{i}{2}\hat{\mathbf{r}}^T H \Omega \mathbf{c} = i\mathbf{c}^T \Omega H \hat{\mathbf{r}} \quad (\text{S25})$$

where we have defined the symplectic matrix $i\Omega_{nm} = [\hat{r}_n, \hat{r}_m]$. Then we can apply the Campbell identity:

$$e^{-\frac{i\mathbf{c}^T H \hat{\mathbf{r}}}{2}} \mathbf{c}^T \hat{\mathbf{r}} e^{\frac{i\mathbf{c}^T H \hat{\mathbf{r}}}{2}} = e^{\frac{i\mathbf{c}^T [\bullet, \hat{\mathbf{r}}^T H \hat{\mathbf{r}}]}{2}} \mathbf{c}^T \hat{\mathbf{r}} = \mathbf{c}^T e^{-t\Omega H \hat{\mathbf{r}}} \quad (\text{S26})$$

such that:

$$C(t) = \mathbf{c}^T \text{Tr}[\hat{\mathbf{r}} \hat{\mathbf{r}}^T \rho_B(0)] e^{tH\Omega} \mathbf{c} \quad (\text{S27})$$

where the trace is still in the Liouville-space sense and not in the matrix sense. Through the normal mode or Williamson decomposition^{85,86}, we may write $H = S^T D S$ where the diagonal matrix D contains the frequencies $\{\omega_j\}$ and S is a symplectic matrix ($S^T \Omega S = \Omega$). Then we can define the following Gaussian unitary:

$$\mathcal{U} \hat{\mathbf{r}} \mathcal{U}^\dagger = S \hat{\mathbf{r}} \quad (\text{S28})$$

such that:

$$\text{Tr}[\hat{\mathbf{r}} \hat{\mathbf{r}}^T \rho_B(0)] = \text{Tr}[\hat{\mathcal{U}}^\dagger \hat{\mathbf{r}} \hat{\mathbf{r}}^T \hat{\mathcal{U}} \rho_B^D(0)] = S^{-1} \text{Tr}[\hat{\mathbf{r}} \hat{\mathbf{r}}^T \rho_B^D(0)] S^{-T} \quad (\text{S29})$$

where $\rho_B^D(0)$ is the thermal state of the diagonal modes:

$$\rho_B^D(0) \propto \prod_j e^{-\frac{\beta\omega_j}{2}(\hat{x}_j^2 + \hat{p}_j^2)} \quad (\text{S210})$$

Then, we can calculate the trace using that:

$$\text{Tr} \left[\begin{pmatrix} \hat{x}_j \\ \hat{p}_j \end{pmatrix} \begin{pmatrix} \hat{x}_j & \hat{p}_j \end{pmatrix} \rho_B^D(0) \right] = \frac{1}{2} \begin{pmatrix} 2n(\omega_j) + 1 & i \\ -i & 2n(\omega_j) + 1 \end{pmatrix} \quad (\text{S211})$$

with $n(\omega_j) = 1/(e^{\beta\omega_j} - 1)$, such that we get:

$$\text{Tr}[\hat{\mathbf{r}} \hat{\mathbf{r}}^T \rho_B^D(0)] = \frac{1}{2} \left(i\Omega + \coth(\frac{\beta}{2}D) \right) \quad (\text{S212})$$

Putting everything together, we have:

$$C(t) = \frac{1}{2} \mathbf{c}^T S^{-1} \left(i\Omega + \coth(\frac{\beta}{2}D) \right) S^{-T} e^{tH\Omega} \mathbf{c} \quad (\text{S213})$$

and because of $S^{-1} \Omega S^{-T} = \Omega$ we can rewrite:

$$C(t) = \frac{1}{2} \mathbf{c}^T \left(i\Omega + S^{-1} \coth(\frac{\beta}{2}D) S^{-T} \right) e^{tH\Omega} \mathbf{c} \quad (\text{S214})$$

$$= \frac{1}{2} \mathbf{c}^T \left(i\Omega + S^{-1} \coth(\frac{\beta}{2}D) S^{-T} \right) S^T e^{tD\Omega} S^{-T} \mathbf{c} \quad (\text{S215})$$

$$= \frac{1}{2} \mathbf{c}^T S^{-1} \left(i\Omega + \coth(\frac{\beta}{2}D) \right) e^{tD\Omega} S^{-T} \mathbf{c} \quad (\text{S216})$$

Specifically for this kind of diagonal matrix D from the Williamson decomposition, where the diagonal entries come in pairs, we have $\Omega D \Omega = -D$ (equivalently $[\Omega, D] = 0$) such that:

$$e^{tD\Omega} = \sum_{n=0}^{\infty} \frac{t^n}{n!} (D\Omega)^n = \cos(Dt) + \sin(Dt)\Omega \quad (\text{S217})$$

Inserting this, we have:

$$C(t) = \frac{1}{2} \mathbf{c}^T S^{-1} \left(i\Omega + \coth\left(\frac{\beta}{2}D\right) \right) (\cos(Dt) + \sin(Dt)\Omega) S^{-T} \mathbf{c} \quad (\text{S218})$$

We may now realize that some of these terms are zero. To see this, we consider the transpose of a scalar, which must equal itself, and utilize the commutativity of Ω and D to show that it is equal to minus itself and thus equates to zero:

$$\mathbf{c}^T S^{-1} \Omega \cos(Dt) S^{-T} \mathbf{c} \stackrel{\text{transpose}}{=} \mathbf{c}^T S^{-1} \cos(Dt) (-\Omega) S^{-T} \mathbf{c} = 0 \quad (\text{S219})$$

$$\mathbf{c}^T S^{-1} \coth\left(\frac{\beta}{2}D\right) \sin(Dt) \Omega S^{-T} \mathbf{c} \stackrel{\text{transpose}}{=} \mathbf{c}^T S^{-1} (-\Omega) \sin(Dt) \coth\left(\frac{\beta}{2}D\right) S^{-T} \mathbf{c} = 0 \quad (\text{S220})$$

Together with $\Omega^2 = -\mathbb{1}$, we therefore end up with:

$$\boxed{C(t) = \frac{1}{2} \mathbf{c}^T S^{-1} \left(-i \sin(Dt) + \coth\left(\frac{\beta}{2}D\right) \cos(Dt) \right) S^{-T} \mathbf{c}} \quad (\text{S221})$$

We will now consider the noise spectrum, which is the Fourier transform of the bath correlation function:

$$S_\beta(\omega) = \int_{-\infty}^{\infty} C(t) e^{i\omega t} dt \quad (\text{S222})$$

By utilizing:

$$-i \int_{-\infty}^{\infty} \sin(Dt) e^{i\omega t} dt = \pi [\delta(\omega - D) - \delta(\omega + D)] \quad (\text{S223})$$

$$\int_{-\infty}^{\infty} \cos(Dt) e^{i\omega t} dt = \pi [\delta(\omega - D) + \delta(\omega + D)] \quad (\text{S224})$$

where scalars such as ω are understood to be proportional to the identity matrix $\omega \mathbb{1}$ and the delta function with a diagonal matrix as argument, is simply a diagonal matrix with elements $\delta(\omega - D)_{ii} = \delta(\omega - D_{ii})$, we recover:

$$S_\beta(\omega) = \frac{\pi}{2} \mathbf{c}^T S^{-1} \left[\delta(\omega - D) - \delta(\omega + D) + \coth\left(\frac{\beta}{2}D\right) [\delta(\omega - D) + \delta(\omega + D)] \right] S^{-T} \mathbf{c} \quad (\text{S225})$$

$$S_\beta(\omega) = \frac{\pi}{2} \mathbf{c}^T S^{-1} \left[\delta(\omega - D) - \delta(\omega + D) + \coth\left(\frac{\beta}{2}D\right) [\delta(\omega - D) - \delta(\omega + D)] \right] S^{-T} \mathbf{c} \quad (\text{S226})$$

where we have used that $\coth$ is an odd function. By defining the spectral density

$$J(\omega) = \frac{\pi}{2} \mathbf{c}^T S^{-1} \delta(\omega - D) S^{-T} \mathbf{c} \quad (\text{S227})$$

we can write the noise spectrum in the familiar form:

$$S_\beta(\omega) = \left(\coth\left(\frac{\beta}{2}\omega\right) + 1 \right) (J(\omega) - J(-\omega)) = \frac{2}{1 - e^{-\beta\omega}} (J(\omega) - J(-\omega)) \quad (\text{S228})$$

Exactly like Eq. S15, except, now we have a more general presentation of the spectral density.

We will now write the spectral density in a way that does not rely on the Williamson decomposition of the matrix H . Simply inserting $D = S^{-T} H S^{-1}$ does not perform the job. Instead, we will use the Sokhotski-Plemelj theorem to write the odd combination of delta functions as:

$$\pi [\delta(\omega - D) - \delta(\omega + D)] = \text{Im} \left[\frac{1}{\omega - D - i0^+} - \frac{1}{\omega + D - i0^+} \right] = \text{Im} \left[\frac{1}{\omega^2 - D^2 - \text{sgn}(\omega) i0^+} \right] D \quad (\text{S229})$$

We can then use that:

$$-D^2 = (-D)D = \Omega D \Omega D = \Omega S^{-T} H S^{-1} \Omega S^{-T} H S^{-1} = S \Omega H \Omega H S^{-1} \quad (\text{S230})$$

Thus, we can write the odd-combination of the spectral density $J^o(\omega)$ as:

$$J^o(\omega) := J(\omega) - J(-\omega) = \mathbf{c}^T S^{-1} S \operatorname{Im} \left[\frac{1}{\omega^2 + (\Omega H)^2 - \operatorname{sgn}(\omega) i 0^+} \right] S^{-1} D S^{-T} \mathbf{c} \quad (\text{S231})$$

Lastly, $S^{-1} D S^{-T} = -S^{-1} \Omega D \Omega S^{-T} = -\Omega H \Omega$ and we arrive at:

$$J^o(\omega) = -\mathbf{c}^T \operatorname{Im} \left[\frac{1}{\omega^2 + (\Omega H)^2 - \operatorname{sgn}(\omega) i 0^+} \right] \Omega H \Omega \mathbf{c} \quad (\text{S232})$$

The problem of finding the spectral density has now been turned into a matrix inversion problem. The practicality of this becomes apparent in the later examples.

We may simplify the expression further if the examples we consider are trivial in the momentum sector and only couple in the position sector. For example, in the basis $\mathbf{r} = (\hat{x}_1, \hat{x}_2, \dots, \hat{p}_1, \hat{p}_2, \dots)^T$, we consider:

$$\Omega = \begin{pmatrix} 0 & \mathbb{1} \\ -\mathbb{1} & 0 \end{pmatrix} \quad H = \begin{pmatrix} X & 0 \\ 0 & \mathbb{1} \end{pmatrix} \quad \mathbf{c} = \begin{pmatrix} \mathbf{c}_x \\ 0 \end{pmatrix} \quad (\text{S233})$$

Then $(\Omega H)^2 = \begin{pmatrix} -X & 0 \\ 0 & -X \end{pmatrix}$ and $\Omega H \Omega = -\begin{pmatrix} \mathbb{1} & 0 \\ 0 & X \end{pmatrix}$ which results in:

$$J^o(\omega) = \mathbf{c}_x^T \operatorname{Im} \left[\frac{1}{\omega^2 - X - \operatorname{sgn}(\omega) i 0^+} \right] \mathbf{c}_x \quad (\text{S234})$$

S2.1. Spectral density of interacting lossy harmonic modes

Consider N bosons that may couple directly to the system, as described by a vector $\mathbf{c}$, and interact among themselves through coupling constants contained in a matrix χ . χ should be positive-definite. In addition, each boson may be coupled to its own structured bath. Such a scenario is visualized in Fig. S1. The resulting Hamiltonian is:




FIG. S1. Schematic of the efficient modeling of an interacting harmonic environment in terms of an effective spectral density for the full environment (Eq. S238). The example used for ‘the system’ is the triple well PES from Sec. IV A 3.

$$H = H_R + \lambda \hat{R}^2 + \hat{R} \sum_n^N c_n \hat{x}_n + \sum_n^N \left[\frac{\hat{p}_n^2}{2} + \omega_n^2 \frac{\hat{x}_n^2}{2} \right] + \sum_{n \neq m}^N \frac{\chi_{nm}}{2} \hat{x}_n \hat{x}_m + \sum_n^N H_{\text{CL}}^{(n)} \quad (\text{S235})$$

H_R is the system Hamiltonian and $\hat{R}$ is the system operator that couples to the harmonic environment. The mass-weighted coordinates of the harmonic environment $\hat{x}_n$, are mutually interacting and they each experience their own

Harmonic environment through the inclusion of the $H_{\text{CL}}^{(n)}$

$$H_{\text{CL}}^{(n)} = \sum_l \frac{\hat{p}_{n,l}^2}{2} + \frac{\omega_{n,l}^2}{2} \left(\hat{q}_{n,l} - \frac{c_{n,l}}{\omega_{n,l}^2} \hat{x}_n \right)^2 \quad (\text{S236})$$

which gives rise to the spectral densities $J_n(\omega) = \frac{\pi}{2} \sum_l \frac{c_{n,l}^2}{\omega_{n,l}} \delta(\omega - \omega_{n,l})$ that characterize the local harmonic environment for the n 'th oscillator $\hat{x}_n$.

This Hamiltonian (Eq. S235), with formally infinitely many degrees of freedom, is of the form Eq. S21. In the appropriate basis, one can then write the coupling matrix of the full harmonic environment, X (Eq. S233) in the following block form:

$$X = \begin{pmatrix} \chi & C^T \\ C & W \end{pmatrix} \quad (\text{S237})$$

where W is diagonal and contains all the squared frequencies of the $J_i(\omega)$ dofs, C is a matrix with N columns that couples each boson i to its respective bath $J_i(\omega)$ and χ is the general symmetric $N \times N$ matrix that couples the lossy bosons. The system only couples to the χ part, such that Eq. S234 becomes:

$$J_{\text{eff}}(\omega) = \text{Im} \left(\begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_N \end{pmatrix}^T \begin{pmatrix} \omega^2 - \omega_1^2 - iJ_1^o(\omega) - \tilde{\mathcal{R}}_1(\omega) & -\chi_{12} & \dots & -\chi_{1N} \\ -\chi_{21} & \omega^2 - \omega_2^2 - iJ_2^o(\omega) - \tilde{\mathcal{R}}_2(\omega) & \dots & -\chi_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ -\chi_{N1} & -\chi_{N2} & \dots & \omega^2 - \omega_N^2 - iJ_N^o(\omega) - \tilde{\mathcal{R}}_N(\omega) \end{pmatrix}^{-1} \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_N \end{pmatrix} \right) \quad (\text{S238})$$

on account of the inversion of a block matrix $\begin{pmatrix} A & B \\ C & D \end{pmatrix}^{-1} = \begin{pmatrix} (A - BD^{-1}C)^{-1} & \dots \\ \dots & \dots \end{pmatrix}$. The Schur complement produced entries along the diagonal, which we rewrote by again applying the Sokhotski-Plemelj theorem:

$$\sum_j c_{n,j}^2 \frac{1}{\omega^2 - \omega_{n,j}^2 - \text{sgn}(\omega) i 0^+} = \sum_j c_{n,j}^2 \left[\text{sgn}(\omega) i \pi \delta(\omega^2 - \omega_{n,j}^2) + \mathcal{P} \left(\frac{1}{\omega^2 - \omega_{n,j}^2} \right) \right] = iJ_n^o(\omega) + \mathcal{R}_n(\omega) \quad (\text{S239})$$

We applied the identity: $\text{sgn}(\omega) \delta(\omega^2 - \nu^2) = \frac{1}{2\nu} [\delta(\omega - \nu) - \delta(\omega + \nu)]$ (for $\nu > 0$) and defined the principal value integral:

$$\mathcal{R}_n(\omega) = \frac{2}{\pi} \mathcal{P} \int_0^\infty \frac{\omega' J_n(\omega')}{\omega^2 - \omega'^2} d\omega' \quad (\text{S240})$$

$$= \mathcal{P} \int_{-\infty}^\infty \frac{d\omega'}{\pi} \frac{\omega' J_n^o(\omega')}{\omega^2 - \omega'^2} \quad (\text{S241})$$

$$= \mathcal{P} \int_{-\infty}^\infty \frac{d\omega'}{2\pi} J_n^o(\omega') \left(\frac{1}{\omega - \omega'} - \frac{1}{\omega + \omega'} \right) \quad (\text{S242})$$

$$= \mathcal{P} \int_{-\infty}^\infty \frac{d\omega'}{\pi} \frac{J_n^o(\omega')}{\omega - \omega'} \quad (\text{S243})$$

Lastly, we reintroduced the neglected reorganization energies by the following consideration:

$$\hat{H} = \dots + \frac{\hat{p}_n^2}{2} + \frac{\omega_n^2}{2} \hat{x}_n^2 + \hat{x}_n^2 \underbrace{\sum_j \frac{c_{n,j}^2}{2\omega_{n,j}^2}}_{\int_0^\infty \frac{d\omega}{\pi} \frac{J_n(\omega)}{\omega}} + \dots \Rightarrow \omega_n^2 \rightarrow \omega_n^2 + 2 \int_0^\infty \frac{d\omega}{\pi} \frac{J_n(\omega)}{\omega} \quad (\text{S244})$$

which means that we may incorporate them into $\mathcal{R}_n(\omega)$ in the following way:

$$\tilde{\mathcal{R}}_n(\omega) = \mathcal{R}_n(\omega) + 2 \int_0^\infty \frac{d\omega'}{\pi} \frac{J_n(\omega')}{\omega'} = \omega \mathcal{P} \int_{-\infty}^\infty \frac{d\omega'}{\pi} \frac{J_n^o(\omega')}{(\omega - \omega')\omega'} \quad (\text{S245})$$

It will be a helpful reminder, that if the pole structure of the bath spectral density is simple e.g. if $J_n^o(\omega) = \frac{f(\omega)}{g(\omega)}$ where $g(\omega)$ accounts for the simple poles, then:

$$\tilde{\mathcal{R}}(\omega) = -2\omega \operatorname{Im} \left(\sum_{\omega_p} \frac{f(\omega_p)}{(\omega - \omega_p)\omega_p g'(\omega_p)} \right) \quad (\text{S246})$$

where ω_p are the zeros of $g(\omega)$ in the upper-half of the complex plane. For the Drude-Lorentz spectral density

$$J^{(DL)}(\omega) = \frac{2\lambda\gamma\omega}{\omega^2 + \gamma^2} \quad \tilde{\mathcal{R}}^{(DL)}(\omega) = \frac{2\lambda\omega^2}{\omega^2 + \gamma^2} \quad (\text{S247})$$

For the following section, and in the main text in general, we use $J(\omega)$ and its odd-version $J^o(\omega) = J(\omega) - J(-\omega)$ interchangeably.

S2.2. Spectral density of interacting cavity and spectator mode, averaged over molecule orientation

We consider the spectral density in Eq. 7 while ignoring the phonon bath:

$$J(\omega) = \operatorname{Im} \frac{\Gamma_c(\omega)c_Q^2 + \eta_R^2\Gamma_Q(\omega)2\omega_c^3\eta_c^2 + 4\eta_Q\eta_R(\omega^2 - i\alpha\omega)c_Q\omega_c\eta_c^2 + 4\eta_Q^2\eta_R^2(\omega^2 - i\alpha\omega)\omega_c^2\eta_c^4}{\Gamma_c(\omega)\Gamma_Q(\omega) - 2(\omega^2 - i\alpha\omega)\omega_c\eta_c^2\eta_Q^2} \quad (\text{S248})$$

where $\Gamma_c(\omega) = \omega^2 - \omega_c^2 - i\alpha\omega$ and $\Gamma_Q(\omega) = \omega^2 - \omega_Q^2 - \left(i + \frac{\omega}{\gamma_Q}\right)J_Q^{(DL)}(\omega)$

We will now introduce $\eta_R = \tilde{\eta}_R \cos(\theta - \phi)$ and $\eta_Q = \tilde{\eta}_Q \cos(\phi)$ where θ is the angle between the reaction-coordinate and the spectator mode and ϕ is the angle between the cavity polarization and the spectator mode dipole moment. We are interested in averaging over ϕ

$$\langle J(\omega) \rangle_\phi = \frac{1}{2\pi} \int_0^{2\pi} J(\omega) d\phi \quad (\text{S249})$$

We apply $\cos(\theta - \phi) = \cos(\theta)\cos(\phi) + \sin(\theta)\sin(\phi)$ and utilize that the average over odd functions equals zero which together with polynomial division gives:

$$\left\langle \frac{1}{a - b \cos^2(\phi)} \right\rangle_\phi = \frac{1}{a\sqrt{1 - b/a}} \quad (\text{S250})$$

$$\left\langle \frac{\eta_R^2}{a - b \cos^2(\phi)} \right\rangle_\phi = \tilde{\eta}_R^2 \left(\frac{\sin^2(\theta)}{a\sqrt{1 - b/a}} + \frac{\cos(2\theta)}{b} \left(\frac{1}{\sqrt{1 - b/a}} - 1 \right) \right) \quad (\text{S251})$$

$$\left\langle \frac{\eta_R\eta_Q}{a - b \cos^2(\phi)} \right\rangle_\phi = \tilde{\eta}_R\tilde{\eta}_Q \frac{\cos(\theta)}{b} \left(\frac{1}{\sqrt{1 - b/a}} - 1 \right) \quad (\text{S252})$$

$$\left\langle \frac{\eta_R^2\eta_Q^2}{a - b \cos^2(\phi)} \right\rangle_\phi = \tilde{\eta}_R^2\tilde{\eta}_Q^2 \left(\frac{\sin^2(\theta)}{b} \left(\frac{1}{\sqrt{1 - b/a}} - 1 \right) + \cos(2\theta) \left(\frac{a}{b^2} \left(\frac{1}{\sqrt{1 - b/a}} - 1 \right) - \frac{1}{2b} \right) \right) \quad (\text{S253})$$

with $a = \Gamma_c(\omega)\Gamma_Q(\omega)$ and $b = 2\tilde{\eta}_Q^2(\omega^2 - i\alpha\omega)\omega_c\eta_c^2$ and a branch cut along the negative real axis.

Supplemental Material S3: Reaction Coordinate Potentials & vibrational state details

We consider a model system which consists of a reaction coordinate on a one dimensional potential energy surface.

$$H_R = \frac{\hat{P}^2}{2M} + V(\hat{R}) \quad (\text{S31})$$

S3.1. Double Well

One way to parameterize the symmetric double well potential is as a quartic: $V_{\text{sym}}(\hat{R}) = E_b \left(1 - \left(\frac{\hat{R}}{R_0}\right)^2\right)^2$. However, we are interested in a parameterization where we have control of the product-, reactant- and barrier frequencies independently. To this end, we consider a piecewise function of quadratic polynomials.

$$V(R) = \begin{cases} \frac{1}{2}M\omega_l^2(R + R_0^l)^2 - E_l & R < R_l \\ E_b - \frac{1}{2}M\omega_0^2R^2 & R_l \leq R \leq R_r \\ \frac{1}{2}M\omega_r^2(R + R_0^r)^2 - E_r & R_r < R \end{cases} \quad (\text{S32})$$

where we only have to specify the frequencies and offsets, with the location of the parabolas and the regions given by:

$$R_0^l = \sqrt{\frac{2(\omega_0^2 + \omega_l^2)(E_b + E_l)}{M\omega_0^2\omega_l^2}} \quad (\text{S33})$$

$$R_l = -\frac{\omega_l^2}{\omega_0^2 + \omega_l^2}R_0^l \quad (\text{S34})$$

$$R_0^r = -\sqrt{\frac{2(\omega_0^2 + \omega_r^2)(E_b + E_r)}{M\omega_0^2\omega_r^2}} \quad (\text{S35})$$

$$R_r = -\frac{\omega_r^2}{\omega_0^2 + \omega_r^2}R_0^r \quad (\text{S36})$$

The definition of ω_r and ω_l differ from that used in the main text, where they referred to the energy spacing between states in the right and left well, respectively.

S3.1.1. Derivation

We require any contiguous pair of parabolas to touch at a single point:

$$a(R + b)^2 - c = C - A(R + B)^2 \quad (\text{S37})$$

$$\Rightarrow 0 = (a + A)R^2 + 2(ab + AB)R + ab^2 + AB^2 - c - C \quad (\text{S38})$$

Where a,b,c and A,B,C denote the parameters of the well and barrier, respectively. We require a unique solution for R, which is the case when the discriminant is zero (for real parameters):

$$R = -\frac{ab + AB}{a + A} \Leftrightarrow (ab + AB)^2 = (a + A)(ab^2 + AB^2 - c - C) \quad (\text{S39})$$

Furthermore, we would like to solve for either b or B since the other parameters are fixed. To this end, the rightmost identity can be rewritten as:

$$(b - B)^2 = \frac{(a + A)(c + C)}{aA} \quad (\text{S310})$$

Associating a,b,c and A,B,C with the parameters of the potential results in the expressions above.

S3.2. Triple Well

In order to study competing pathways, we consider a potential energy surface with three wells. We can utilize the same parameterization as above, which results in:

$$V(R) = \begin{cases} \frac{1}{2}M\omega_{ll}^2(R + R_0^{ll})^2 - E_{ll} & -\infty < R \leq R_{ll} \\ E_l - \frac{1}{2}M\omega_l^2(R + R_0^l)^2 & R_{ll} < R \leq R_l \\ \frac{1}{2}M\omega_0^2R^2 - E_0 & R_l < R \leq R_r \\ E_r - \frac{1}{2}M\omega_r^2(R + R_0^r)^2 & R_r < R \leq R_{rr} \\ \frac{1}{2}M\omega_{rr}^2(R + R_0^{rr})^2 - E_{rr} & R_{rr} < R < \infty \end{cases} \quad (\text{S311})$$

with

$$R_0^l = \sqrt{\frac{2(\omega_0^2 + \omega_l^2)(E_0 + E_l)}{M\omega_0^2\omega_l^2}} \quad (\text{S312})$$

$$R_l = -\frac{\omega_l^2}{\omega_0^2 + \omega_l^2} R_0^l \quad (\text{S313})$$

$$R_0^l = R_0^l + \sqrt{\frac{2(\omega_{ll}^2 + \omega_l^2)(E_{ll} + E_l)}{M\omega_{ll}^2\omega_l^2}} \quad (\text{S314})$$

$$R_{ll} = -\frac{\omega_l^2 R_0^l + \omega_{ll}^2 R_0^l}{\omega_l^2 + \omega_{ll}^2} \quad (\text{S315})$$

$$R_0^r = -\sqrt{\frac{2(\omega_0^2 + \omega_r^2)(E_0 + E_r)}{M\omega_0^2\omega_r^2}} \quad (\text{S316})$$

$$R_r = -\frac{\omega_r^2}{\omega_0^2 + \omega_r^2} R_0^r \quad (\text{S317})$$

$$R_0^r = R_0^r - \sqrt{\frac{2(\omega_{rr}^2 + \omega_r^2)(E_{rr} + E_r)}{M\omega_{rr}^2\omega_r^2}} \quad (\text{S318})$$

$$R_{rr} = -\frac{\omega_r^2 R_0^r + \omega_{rr}^2 R_0^r}{\omega_r^2 + \omega_{rr}^2} \quad (\text{S319})$$

The ground state of the different wells can then be made approximately degenerate, by specifying the offset energy as $E = \frac{\omega_{\text{well}}}{2}$. This was done for triple well potential in Figure 7a, with some additional fine-tuning to get the excited-state degeneracy seen there. Alternatively, which was done for the double wells in figure 5, one can instead specify $E_{l,r} = 0$ and add a linear term $\frac{1}{2} \frac{\omega_l - \omega_r}{x_0^l + x_0^r} \hat{R}$ to the potential. Probably, these approaches are equivalent even though we do not attempt to show that here.

S3.3. Localized states

In order to analyze the dynamics in terms of intra- and inter-well dynamics, we consider localized states which are not eigenstates of the reaction coordinate Hamiltonian $H_R = \sum_n E_n |n\rangle \langle n|$. We obtain these localized states by diagonalizing the reaction coordinate operator in the sub-space of the relevant delocalized energy-states, e.g.:

$$\begin{pmatrix} \langle n_1 | \hat{R} | n_1 \rangle & \langle n_1 | \hat{R} | n_2 \rangle & \langle n_1 | \hat{R} | n_3 \rangle \\ \langle n_2 | \hat{R} | n_1 \rangle & \langle n_2 | \hat{R} | n_2 \rangle & \langle n_2 | \hat{R} | n_3 \rangle \\ \langle n_3 | \hat{R} | n_1 \rangle & \langle n_3 | \hat{R} | n_2 \rangle & \langle n_3 | \hat{R} | n_3 \rangle \end{pmatrix} = O \widehat{D}^{\text{diag}} O^T \quad (\text{S320})$$

The localized states $|l_{1,2,3}\rangle$ are then given by:

$$|l_i\rangle = \sum_{j=1,2,3} O_{ji} |n_j\rangle \quad (\text{S321})$$

In figure 4, the two ground states and the two excited states were hybridized to produce the localized states shown. In figure 5, even though the eigenstates were already approximately localized due to the broken symmetry of the potentials, we still hybridized the ground states and also the two states closest to Frequency = 1200cm⁻¹ to obtain the localized states shown. In figure 7a the three excited states around 1570cm⁻¹ were localized with this process.

S3.4. Numerics

For the simulations, the reaction coordinate was expressed in the eigen basis of the system Hamiltonian (without the reorganization term). The system Hamiltonian was numerically diagonalized after expressing it in the Discrete Variable Representation (DVR) based on the sinc-function⁸⁷. Sinc-DVR resembles that of discretizing the position basis at N points spaced by ΔR , where operators that are a function of $\hat{R}$ remain diagonal in this representation.

However, instead of a finite-difference representation of the momentum operator, in sinc-DVR, the momentum operator is instead represented as:

$$(\hat{P})_{nm} = \begin{cases} 0 & n = m \\ \frac{-i(-1)^{n-m}}{\Delta R(n-m)} & n \neq m \end{cases} \quad (\text{S322})$$

$$(\hat{P}^2)_{nm} = \begin{cases} \frac{\pi^2}{3\Delta R^2} & n = m \\ \frac{2(-1)^{n-m}}{\Delta R^2(n-m)^2} & n \neq m \end{cases} \quad (\text{S323})$$

N and ΔR was chosen such that convergence was observed in the eigenvalue of the relevant states.

Supplemental Material S4: Details on Hierarchical Equations Of Motion

We refer to our choice of method for simulating the dynamics as HEOM⁵⁴, but instead of viewing the equations of motion as a hierarchy of coupled differential equations, we treat it as the system coupled to a collection of bosons. See e.g. references^{55–59}. For example, the master equation^{56,58,59}:

$$\begin{aligned} \frac{d\varrho}{dt} = \mathcal{L}\varrho = & -i \left[\hat{H}_S + \sum_k \text{Im}(z_k) \hat{a}_k^\dagger \hat{a}_k + \sum_k \text{Re}(\sqrt{d_k}) \hat{S}(\hat{a}_k + \hat{a}_k^\dagger), \varrho \right] \\ & + 2 \sum_k \text{Re}(z_k) \left(\hat{a}_k \varrho \hat{a}_k^\dagger - \frac{1}{2} \{ \hat{a}_k^\dagger \hat{a}_k, \varrho \} \right) \\ & - 2 \sum_k \text{Im}(\sqrt{d_k}) \left(\hat{a}_k \varrho \hat{S} - \frac{1}{2} \{ \hat{S} \hat{a}_k, \varrho \} \right) \\ & - 2 \sum_k \text{Im}(\sqrt{d_k}) \left(\hat{S} \varrho \hat{a}_k^\dagger - \frac{1}{2} \{ \hat{a}_k^\dagger \hat{S}, \varrho \} \right) \end{aligned} \quad (\text{S41})$$

constitutes an open quantum system for the system S with a harmonic environment of K bosons with BCF:

$$C(t) = \sum_k^K d_k e^{-z_k t} \quad (\text{S42})$$

provided that the initial state factorizes as $\varrho(t=0) = \rho_S(0) \otimes^K |0\rangle\langle 0|$

The noise spectrum is consequently:

$$S_\beta(\omega) = \sum_k^K 2 \frac{\text{Re}(d_k z_k^*) - \text{Im}(d_k) \omega}{\text{Re}(z_k)^2 + (\omega - \text{Im}(z_k))^2} \quad (\text{S43})$$

In order to better understand the generator in Eq. S41, consider $\text{Im}(\sqrt{d_k}) = 0$. In such cases, the structure is equivalent to a Lindbladian (a generator for a completely positive, trace preserving dynamical semigroup). This would constitute a physical setup, where the system is coupled to bosons with Markovian loss and the noise spectrum is decomposed as a sum of Lorentzians. However, in general, this will not be an efficient decomposition and K would need to be too large to be practical. Allowing $\text{Im}(\sqrt{d_k}) \neq 0$ alleviates this problem, since the terms in Eq. S43 are now able to acquire negative values and may cancel eachother in a manner that assists the decomposition⁵⁷. Allowing this comes at the cost of the two terms in Eq. S41, which breaks the complete positivity of the generator. This means that, even though Eq. S41 is trace-preserving ($\text{Tr}[\dot{\varrho}] = 0$) the full density matrix $\varrho(t)$ may acquire negative eigenvalues and no longer corresponds to a physical setup. However, crucially, the reduced density matrix $\rho_S(t) = \text{Tr}_K[\varrho(t)]$ is still a physical density matrix.^{56,58,59}

Ref. 88 pointed to a potential problem. Namely, when cutoffs of the auxiliary modes are inevitably introduced for numerical implementation, the evolution can diverge in a way that is seemingly not just remedied by increasing the cutoff. However, Ref. 56 notes that the presence of the divergence depends on the choice of 'gauge' when decomposing $C(t-t')$. This gauge-freedom is made manifest in the master equation by applying a similarity transformation using $\kappa_k \hat{a}_k^\dagger \hat{a}_k$, which does not change the initial and final conditions for the evolution, but simply transforms $\hat{a}_k \rightarrow \hat{a}_k / \kappa_k$ and $\hat{a}_k^\dagger \rightarrow \kappa_k \hat{a}_k^\dagger$. The κ_k 's can then be chosen such that norm of the CP-breaking term is minimized (but non-zero), which,

in that case, remedied the divergence. In the case of a single system operator interacting with the environment, i.e. a scalar valued BCF, the optimal gauge is known and Eq. S41 is written in this gauge. The ubiquitous and successful application of these CP-breaking formulations suggests that these concerns are not dire and also, in our experience, Eq. S41 has yet to exhibit divergent behavior.

Nonetheless, a lack in understanding how and when such divergent behavior arises and because it may be related to the CP-breaking of these types of methods, could motivate the search for Markovian embeddings that preserves complete-positivity. There are already numerous formulations that achieve this (see refs. ^{61,62,89–91} to name a few). However, an overview of the drawbacks for each and every method, such as computational complexity and robustness, has yet to be developed. Thus, the tried and tested HEOM formulations remain our choice of method.

Another useful version of HEOM is obtained from Eq. S41 by applying a similarity transformation under $e^{\hat{a}_k \bullet \hat{a}_k^\dagger}$, which has the following effects^{58,59}:

$$\rho_S(t) = \text{Tr}_K [\varrho(t)] \rightarrow \langle \mathbf{0} | \varrho(t) | \mathbf{0} \rangle \quad (\text{S44})$$

$$\hat{a}_k^\dagger \bullet \rightarrow \hat{a}_k^\dagger \bullet + \bullet \hat{a}_k^\dagger \quad (\text{S45})$$

$$\bullet \hat{a}_k \rightarrow \bullet \hat{a}_k + \hat{a}_k \bullet \quad (\text{S46})$$

$$(\text{S47})$$

with $\hat{a}_k \bullet$ and $\bullet \hat{a}_k^\dagger$ unchanged. Now, the reduced density matrix for the system can be obtained by simply projecting the auxiliary modes onto the vacuum^{57–59}. This is helpful for both the hierarchical approaches, where you can now read-off the density matrix from the lowest tier and also tensor networks, where it allows you to place left and right dofs of the auxiliary modes independently of each other, since the trace could otherwise introduce loops that in the worst case could be difficult to contract. The price for this benefit, is that the master equation:

$$\dot{\varrho}(t) = -i \left[\hat{H}_S, \varrho(t) \right] + \sum_k^K \left\{ -z_k \hat{a}_k^\dagger \hat{a}_k \varrho(t) - i\sqrt{d_k} \left(\hat{a}_k + \hat{a}_k^\dagger \right) \hat{S} \varrho(t) - i\sqrt{d_k^*} \hat{S} \varrho(t) \hat{a}_k^\dagger + H.C. \right\} \quad (\text{S48})$$

Is no longer trace perserving:

$$\text{Tr} [\dot{\varrho}(t)] = \sum_k -2 \text{Re}(z_k) \text{Tr} \left[\hat{a}_k^\dagger \hat{a}_k \varrho(t) \right] - i \text{Tr} \left[\left(\sqrt{d_k} \hat{a}_k^\dagger - \sqrt{d_k^*} \hat{a}_k \right) \hat{S} \varrho(t) \right] \quad (\text{S49})$$

For weak to moderate coupling, this does not pose a problem and on the contrary, we experienced that simulating the dynamics with Eq. S48 required fewer computational resources than Eq. S41 (e.g. due to a lower bond-dimension). However, we did not carry out a precise comparison and it remains anecdotal information. Nonetheless, for strong coupling scenarios the trace can grow uncontrollably and one should resort to the trace preserving formulation, Eq. S41.

S4.1. Decomposing the Bath Correlation Function

We will now describe how to obtain the parameters d_k and z_k . There exists multiple methods to fit a given bath correlation function by a sum of complex exponentials. We rely on the AAA-algorithm^{67,68} to efficiently acquire the decomposition in Eq. 22. In essence, $S_\beta(\omega)$ is decomposed as a sum of simple poles p_n with residues r_n within a window of frequencies $\omega \in [\omega_l, \omega_h]$, such that:

$$d_k = -i\omega_- r_k \quad z_k = i(\omega_- p_k + \omega_+) \quad (\text{S410})$$

where $\omega_\pm = (\omega_h \pm \omega_l)/2$ and k labels the poles in the lower half of the complex plane. The decomposition can usually be carried out to numerical accuracy with K on the order of 10 – 100, depending on the intricacies of the noise spectrum.

The reorganization energy in terms of the effective spectral density can be written as:

$$\lambda = \frac{1}{\pi} \int_0^\infty d\omega \frac{J(\omega)}{\omega} \quad (\text{S411})$$

$$= \int_0^\infty \frac{d\omega}{2\pi} (1 - e^{-\beta\omega}) \frac{S_\beta(\omega)}{\omega} \quad (\text{S412})$$

$$= \mathcal{P} \int_{-\infty}^\infty \frac{d\omega}{2\pi} \frac{S_\beta(\omega)}{\omega} \quad (\text{S413})$$

$$= \mathcal{P} \int_{-\infty}^\infty \frac{d\omega}{2\pi} \int_{-\infty}^\infty dt C(t) \frac{e^{i\omega t}}{\omega} \quad (\text{S414})$$

$$= \frac{i}{2} \int_{-\infty}^\infty dt C(t) \text{sgn}(t) \quad (\text{S415})$$

$$= - \int_0^\infty dt \text{Im} [C(t)] \quad (\text{S416})$$

by using that $S_\beta(\omega) = \frac{2}{1-e^{-\beta\omega}} J(\omega) = e^{\beta\omega} S_\beta(-\omega) = \int_{-\infty}^\infty dt C(t) e^{i\omega t}$ and $C(t) = C^*(-t)$. Then the reorganization energy for our emulating bath is easily obtained as Eq. 19.

S4.2. Effective Markovian correction

As detailed in Sec. S4, the noise spectrum is decomposed into contributions from each auxiliary mode. We have found that some of these auxiliary modes constitute a Markovian bath. Thus, instead of treating these degrees of freedom explicitly, we can incorporate them as a jump operator on the system degree of freedom. This can be made precise using the Universal Lindblad Equation (ULE) by Nathan and Rudner^{72,92}. Their rigorous analysis proves that a Gaussian environment constitutes a Markovian bath once the characteristic correlation time of the bath τ remain small compared to the rate of bath-induced evolution Γ . Once $\Gamma\tau \ll 1$ the Markovian dynamics are accurately described by introducing the jump operator $\hat{L} = \sum_{n,m} \sqrt{S_\beta(E_m - E_n)} R_{nm} |n\rangle\langle m|$ for the system. They⁹² also provide an expression for the Hermitian Lamb shift, which for Markovian noise spectrums (e.g. $S_\beta(\omega) \sim \text{constant}$) cancel the reorganization term:

$$H_{\text{Lamb Shift}} = - \sum_{n,m,l} \mathcal{P} \int_{-\infty}^\infty \frac{d\omega}{2\pi} \frac{\sqrt{S_\beta(\omega + E_n - E_l)} S_\beta(\omega + E_m - E_l)}{\omega} R_{nl} R_{lm} |n\rangle\langle m| \approx - \int_{-\infty}^\infty \frac{d\omega}{2\pi} \frac{S_\beta(\omega)}{\omega} \hat{R}^2 = -\lambda \hat{R}^2 \quad (\text{S417})$$

Thus, for the auxiliary modes that are deemed Markovian, we simply remove their contribution to the reorganization energy.

In their exposition, they have scaled the system operator to have unit spectral norm, such that the complete error bound is $\|\hat{R}\|^2 \Gamma \tau$, which becomes trivial for the unbounded operator $\hat{R}$. However, the system Hilbert space is anyways truncated when carrying out the dynamical simulation, so applying their results to the system after it has been projected to a finite-dimensional Hilbert space $\hat{R} \rightarrow \hat{P} \hat{R} \hat{P}$ ensures that the error induced by compressing the Markovian auxiliary modes into the system jump operator $\hat{L}$ is bounded by $\|\hat{P} \hat{R} \hat{P}\|^2 \Gamma \tau$, when comparing to the uncompressed dynamical simulation. Assuming the dynamical simulation is converged with regards to the projection error, the error induced by the Markovian compression should not scale with increasing the projected space, as the formal error bound $\|\hat{P} \hat{R} \hat{P}\|^2 \Gamma \tau$ otherwise suggests. Instead, in the following discussion, we treat $\|\hat{P} \hat{R} \hat{P}\|$ as the characteristic length scale of the reaction coordinate under the dynamics, which is $\mathcal{O}(1) \times \text{a.u.}$.

It should be noted that E_n and $|n\rangle$ refer to the energies and eigenstates of the system dressed by the residual environment, which may be difficult to compute exactly. However, as discussed by Nathan and Rudner, the jump operator can be evaluated through an interaction-picture expansion. Writing $H = H_0 + V$, we treat the evolution generated by the tractable reference Hamiltonian H_0 exactly and incorporate the residual interaction V perturbatively. In our case, $H_0 = H_R + \lambda \hat{R}^2 + H_B$ contains the system Hamiltonian, the reorganization term, and the free Hamiltonian of the residual bath, while V contains the coupling to that bath. A low-order truncation is justified when the interaction-picture perturbation $e^{iH_0 t} V e^{-iH_0 t}$ changes the coupling operator ($\hat{R}$ in our case) only weakly over the timescale of the Markovian bath. In the numerical examples considered, we find that the zeroth-order construction of $\hat{L}$, which retains the evolution under H_0 exactly, yields quantitatively accurate results in the regime of validity of the ULE, $\Gamma\tau \ll 1$. Practically, this means that we may approximate the E_n and $|n\rangle$ that are used to compute $\hat{L}$ by $H_R + \lambda \hat{R}^2 = \sum_n E_n |n\rangle\langle n|$. This additional approximation is motivated by the perturbative argument above and

supported by our numerical results, but is not covered by the original ULE error bound. Although $\Gamma\tau \ll 1$ alone does not guarantee the accuracy of our zeroth-order truncation, the overall method remains controllable by the inclusion of higher orders.

While Nathan and Rudner⁷² control the Markovian approximation in terms of $\Gamma\tau$, they also define $\Gamma_0\tau_0 = 4 \int_0^\infty t|C(t)|dt$ which is more practical to work with than their expression for $\Gamma\tau$. However, as they explain, $\Gamma_0\tau_0 \leq \Gamma\tau$ and while $2\Gamma_0\tau_0$ can be used to bound the correction to the Bloch-Redfield equation⁹³ they are only able to bound the correction to their master equation using $\Gamma\tau$ ⁷². Interestingly, for a single exponential with a real prefactor $C(t) = |d|e^{-zt}$, we can compute $\Gamma\tau = \frac{2}{\pi} 8 \frac{|d|}{\text{Re}(z)^2}$ (using integrals of the zeroth modified Bessel function of the second kind) which compared to $2\Gamma_0\tau_0 = 8 \frac{|d|}{\text{Re}(z)^2}$ is smaller by a factor $\frac{2}{\pi}$. Thus, as a practical guiding principle, we may assess the Markovian-ness of the auxiliary modes in terms of:

$$2\Gamma_0\tau_0 = 8 \int_0^\infty t|C(t)|dt = 8 \int_0^\infty t \left| \sum_k d_k e^{-z_k t} \right| dt \leq 8 \sum_k \int_0^\infty t |d_k e^{-z_k t}| dt = 8 \sum_k \frac{|d_k|}{\text{Re}(z_k)^2} \quad (\text{S418})$$

and use this to supply candidates for the Markovian compression. However, for the approximation to be controlled, the final delegation of auxiliary modes to the Markovian compression should be informed by the exact error bound by numerically evaluating $\Gamma\tau$, as the two error bounds need not be comparable in general.

Lastly, we note that if the noise spectrum of the proposed Markovian bath $S_M(\omega) = \sum_{k \in K_M} 2 \text{Re} \left(\frac{d_k}{z_k - i\omega} \right)$ does not remain nonnegative, $S_M(\omega) \not\geq 0$, it is in conflict with the assumptions of Nathan and Rudner⁷² and their results do not immediately apply. Compression methods based on adiabatic elimination^{69–71} generally do not rely on such a semi-positivity requirement and possibly the results of Nathan and Rudner, could also be extended to such cases. However, developing such an extension and additionally deriving a bound for the error of approximating the residual-bath-dressed system, in order to elevate our compression method from controllable to generally rigorously bounded, remains outside the scope of this work.

We complement the discussion above, by exemplifying the validity of our approach, when simulating the symmetric double well interacting with the phonon bath, for different values of $2\Gamma_0\tau_0$. The resulting reaction rates are shown in Fig. S2. The phonon bath has been fitted to high-accuracy (tolerance of $\sim 10^{-14}$) and the number of auxiliary modes is $|K_{\text{fit}}| = 19$. However, many of these modes basically constitute a Markovian environment, so choosing K_M of these such that $2\Gamma_0\tau_0 \approx 0.01$, reduces the number of auxiliary modes that need to be simulated explicitly to $|K| = |K_{\text{fit}} \setminus K_M| = 2$. That is, the evolution induced by $|K_M| = 17$ of these auxiliary modes can be captured by the system jump operator $\hat{L}$, as seen by the resulting reaction rate $k(t) = \frac{P_r(t)}{1 - 2P_r(t)}$ (Eq. S53). The resulting evolution is nearly indistinguishable from the case where all the auxiliary modes have been simulated explicitly, $2\Gamma_0\tau_0 = 0$ and $|K| = 19$. However, increasing $|K_M| = 18$ and thus $2\Gamma_0\tau_0 \approx 0.06$, begins to introduce a deviation on the calculated reaction rate k . Based on this, we have chosen to truncate K_{fit} to K by choosing K_M such that $\Gamma\tau$ is at most 0.01 (in practice often smaller) for all our simulations. This allows us to substantially reduce the number of explicit auxiliary modes while staying true to the dynamics.

The results for this example are summarized in Table I

Number of explicit modes, $ K $	$ K_M $	$4 \sum_k^M \frac{ d_k }{\text{Re}(z_k)^2}$	$\Gamma\tau$	Absolute relative deviation in k
19	0	0	0	0%
2	17	0.00504	0.00598	0.102%
1	18	0.03045	0.03643	1.215%
0	19	7.3125	5.50957	25.8932%

TABLE I. Comparison of different levels of compression for the dynamics of the symmetric double well interacting with the phonon bath. The heuristic for "Markovianity" $4 \sum_k^M \frac{|d_k|}{\text{Re}(z_k)^2}$ is compared to the proper error bound $\Gamma\tau$ and the deviation in the reaction rate $|k - k_{\text{max}}|/k_{\text{max}}$ when compared to the fully explicit simulation k_{max} is presented.

Supplemental Material S5: Reaction Rate Analysis

We define the reaction rate constant by following e.g. references^{47,48,94} For the double well potential, we define the left well as the reactant region with partial concentration P_r and the right well as the product region with partial concentration $P_p = 1 - P_r$.




FIG. S2. Comparison of $k(t) = \frac{\dot{P}_r(t)}{1-2P_r(t)}$ (Eq. S53) for different levels of Markovian compression of the phonon bath. See Table I for an overview of the results. The observed oscillations correspond to the inherent time scale of the phonon bath $\gamma_R^{-1} \approx \beta \approx 25\text{fs}$ and originates from the quenched dynamics since the system is initialized in a product state with the phonon bath. The simulations were carried out using tensor networks, and for this environment, the number of physical indices needed to represent the environment could be reduced from $2 \times 19 = 38$ to $2 \times 2 = 4$ without having any effect on the observed reaction rate. This should be compared to reducing the number of indices by performing a less accurate decomposition of the bath correlation function in terms of only two exponentials, which would result in a larger deviation in the observed rate.

P_r is defined as an expectation value of the projection onto the left well:

$$P_r(t) = \text{Tr} [\hat{P}_r \rho(t)] \quad \text{with} \quad \hat{P}_r = \int_{-\infty}^0 dR |R\rangle \langle R| \quad (\text{S51})$$

Past some transient behavior, we can assume the expectation values follow the rate equation:

$$\dot{P}_r(t) = -kP_r(t) + k_bP_p(t) \quad (\text{S52})$$

where k is the forward reaction rate and k_b is the backward reaction rate, which determine the equilibrium distribution $\frac{k_b}{k} = \frac{P_r(t \rightarrow \infty)}{P_p(t \rightarrow \infty)}$. Using this, one can isolate k in Eq. S52:

$$k = \frac{\dot{P}_r(t)}{-P_r(t) + \frac{P_r^{eq}}{P_p^{eq}}(1 - P_r(t))} \quad (\text{S53})$$

The time derivative of $P_r(t)$ can be calculated either by finite difference:

$$\dot{P}_r(t) \equiv \frac{P_r(t + \Delta t) - P_r(t)}{\Delta t} \quad (\text{S54})$$

or by using the equations of motion, $\dot{\rho} = \mathcal{L}[\rho]$:

$$\dot{P}_r(t) = \text{Tr} [\hat{P}_r \dot{\rho}(t)] = \text{Tr} [\hat{P}_r \mathcal{L}[\rho(t)]] = \text{Tr} [\mathcal{L}^\dagger[\hat{P}_r] \rho(t)] \quad (\text{S55})$$

Where $\mathcal{L}^\dagger$ is the adjoint of the super-operator $\mathcal{L}$.

Furthermore, for symmetric potential energy surfaces, it is straight forward to show that the equilibrium distribution is uniform $P_r^{eq} = P_p^{eq} = \frac{1}{2}$. Consider e.g. the joint system-bath parity operation $\hat{S} : \begin{pmatrix} R \\ B \end{pmatrix} \rightarrow \begin{pmatrix} -R \\ -B \end{pmatrix}$, which leaves $H = H_R + \hat{R}\hat{B} + H_B$ invariant. Then:

$$\text{Tr} [e^{-\beta \hat{H}} \hat{P}_r] = \text{Tr} [\hat{S} e^{-\beta \hat{H}} \hat{S} \hat{P}_r] = \text{Tr} [e^{-\beta \hat{H}} \hat{S} \hat{P}_r \hat{S}] = \text{Tr} [e^{-\beta \hat{H}} (\mathbb{1} - \hat{P}_r)] \quad (\text{S56})$$

However, for general potential energy surfaces, it is helpful to consider alternatives to Eq. S53 for retrieving the forward reaction rate. One approach is to fit the solution of the rate equation to the expectation value of the simulation:

$$P_r(t) = \frac{k_b}{k + k_b} + \left(P_r(t_0) - \frac{k_b}{k + k_b} \right) e^{-(k+k_b)(t-t_0)} \quad (\text{S57})$$

where t_0 is typically chosen to be at a later time than the initialization of the simulation ($t = 0$) past the transients. Another approach is to consider a linear fit of Eq. S52 in terms of k and k_b e.g. through the pseudo-inverse of the time-series data:

$$\begin{pmatrix} \dot{P}_r(t_0) \\ \dot{P}_r(t_1) \\ \dot{P}_r(t_2) \\ \vdots \end{pmatrix} = \begin{pmatrix} -P_r(t_0) & 1 - P_r(t_0) \\ -P_r(t_1) & 1 - P_r(t_1) \\ -P_r(t_2) & 1 - P_r(t_2) \\ \vdots & \vdots \end{pmatrix} \begin{pmatrix} k \\ k_b \end{pmatrix} \quad (\text{S58})$$

$$\Rightarrow \begin{pmatrix} k \\ k_b \end{pmatrix} = \begin{pmatrix} -P_r(t_0) & 1 - P_r(t_0) \\ -P_r(t_1) & 1 - P_r(t_1) \\ -P_r(t_2) & 1 - P_r(t_2) \\ \vdots & \vdots \end{pmatrix}^{-1} \begin{pmatrix} \dot{P}_r(t_0) \\ \dot{P}_r(t_1) \\ \dot{P}_r(t_2) \\ \vdots \end{pmatrix} \quad (\text{S59})$$

However, the transients in $P_r(t)$ can become long lived in $\dot{P}_r(t)$ due to the time-derivative increasing the amplitude of the fluctuations. Therefore, it can be advantageous to "average out" the transients by considering the integral of Eq. S52 before performing the linear fit of k and k_b .

$$\begin{pmatrix} k \\ k_b \end{pmatrix} = \begin{pmatrix} -\int_{t_0}^{t_1} P_r(t) dt & \int_{t_0}^{t_1} (1 - P_r(t)) dt \\ -\int_{t_0}^{t_2} P_r(t) dt & \int_{t_0}^{t_2} (1 - P_r(t)) dt \\ -\int_{t_0}^{t_3} P_r(t) dt & \int_{t_0}^{t_3} (1 - P_r(t)) dt \\ \vdots & \vdots \end{pmatrix}^{-1} \begin{pmatrix} P_r(t_1) - P_r(t_0) \\ P_r(t_2) - P_r(t_0) \\ P_r(t_3) - P_r(t_0) \\ \vdots \end{pmatrix} \quad (\text{S510})$$

where the integrals can be performed as a Riemann sum. By simulating for long enough times and calculating the reaction rates for different t_0 , we make sure that k has converged upon a stationary value. By comparing the different approaches, we found Eq. S510 to be the most robust with fastest convergence, in the general (asymmetric potential) setting.

S5.1. Triple well reaction rates

In this setting, we consider the left and right well for distinct product regions with partial concentrations P_l and P_r , respectively. The middle well is the reactant region denoted by P_0 . Similar to the above, these are defined as expectation values of (notice the sign-convention of Eq. S311, e.g. $-x_0^l < 0 < -x_0^r$):

$$\hat{P}_l = \int_{-\infty}^{-x_0^l} dx |x\rangle \langle x| \quad (\text{S511})$$

$$\hat{P}_0 = \int_{-x_0^l}^{-x_0^r} dx |x\rangle \langle x| \quad (\text{S512})$$

$$\hat{P}_r = \int_{-x_0^r}^{\infty} dx |x\rangle \langle x| \quad (\text{S513})$$

for which the rate equation becomes:

$$\begin{pmatrix} \dot{P}_l(t) \\ \dot{P}_0(t) \\ \dot{P}_r(t) \end{pmatrix} = \begin{pmatrix} P_0(t) & 0 & -P_l(t) & 0 \\ -P_0(t) & -P_0(t) & P_l(t) & P_r(t) \\ 0 & P_0(t) & 0 & -P_r(t) \end{pmatrix} \begin{pmatrix} k_{0 \rightarrow l} \\ k_{0 \rightarrow r} \\ k_{l \rightarrow 0} \\ k_{r \rightarrow 0} \end{pmatrix} \quad (\text{S514})$$

Similar to Eq. S510, the reaction rates are calculated by forming a matrix from the discrete time-series and solving the linear problem.

Supplemental Material S6: Probability Currents: Transition Rate Between States

We are interested in how the rate of change in the population of the different system states depend on the environment. Using Eq. 18, we can consider the time derivative of $\hat{P}_n = |n\rangle\langle n|$, in terms of the reduced system density matrix $\rho_S(t) = \langle \mathbf{0} | \rho(t) | \mathbf{0} \rangle$. The instantaneous rate of change of $\rho_S(t)$ is (in our case $\hat{H}_S = \hat{H}_R$ and $\hat{S} = \hat{R}$):

$$\dot{\rho}_S(t) = -i \left[\hat{H}_S + \sum_k \lambda_k \hat{S}^2, \rho_S(t) \right] - i \sum_k \left[\hat{S}, \sqrt{d_k} \rho_k(t) + \sqrt{d_k^*} \rho_k^\dagger(t) \right] + \sum_i L_i \rho_S(t) L_i^\dagger - \frac{1}{2} \left\{ L_i^\dagger L_i, \rho_S(t) \right\} \quad (\text{S61})$$

with $\lambda_k = -\text{Im} \left(\frac{d_k}{z_k} \right)$ and $\rho_k(t) = \langle \mathbf{0} | \hat{a}_k \varrho(t) | \mathbf{0} \rangle$ ⁹⁵ and where we accommodate multiple markovian correctional terms L_i , which will be useful when we later subdivide the noise spectrum into distinct components. The Heisenberg equation of motion for $\hat{P}_n$ is then:

$$\begin{aligned} \dot{P}_n(t) = & i \text{Tr} \left[[\hat{H}_S, \hat{P}_n] \rho_S(t) \right] + i \sum_k \left(\lambda_k \text{Tr} \left[[\hat{S}^2, \hat{P}_n] \rho_S(t) \right] + \text{Tr} \left[[\hat{S}, \hat{P}_n] \left(\sqrt{d_k} \rho_k(t) + \sqrt{d_k^*} \rho_k^\dagger(t) \right) \right] \right) \\ & + \sum_i \left(\text{Tr} \left[L_i^\dagger \hat{P}_n L_i \rho_S(t) \right] - \frac{1}{2} \text{Tr} \left[\{ L_i^\dagger L_i, \hat{P}_n \} \rho_S(t) \right] \right) \end{aligned} \quad (\text{S62})$$

Utilizing References 50,51 one can imbue the interpretation of a probability current flowing from the state n to m :

$$\dot{P}_n(t) = - \sum_m j_{n \rightarrow m}(t) \quad (\text{S63})$$

We will expand on this interpretation by decomposing the probability current further. Specifically, we will assign the different terms in Eq. S62 to different contributions $j_{n \rightarrow m}^{(i)}(t)$ of the probability current $j_{n \rightarrow m}(t) = \sum_i j_{n \rightarrow m}^{(i)}(t)$. To do this, we consider a grouping of the transitions in terms of those induced by the system Hamiltonian and those induced by coupling to the harmonic environment. Furthermore, the harmonic environment is sub-divided into contributions from the distinct spectral densities:

$$J(\omega) = \sum_i J^{(i)}(\omega) \quad \Rightarrow \quad C(t) = \sum_i \sum_k d_k^{(i)} e^{-z_k^{(i)} t} \quad (\text{S64})$$

where the $d_k^{(i)}$ and $z_k^{(i)}$ parameters are obtained by decomposing the i 'th noise spectrum independently from the others. In our case of Eq. 12, we have two distinct spectral densities such that we define three different contributions to the probability currents (Eq. 15). We denote the probability current due to H_S as *Tunneling*, the contributions from the Drude-Lorentz spectral density of the phonons as *Solvent* and the contributions associated with the spectral density of the lossy cavity is simply referred to as *Cavity*. The probability currents are then defined as follows:

$$\partial_t \langle n | \rho_S(t) | n \rangle = - \sum_m j_{n \rightarrow m}(t) \quad (\text{S65})$$

$$j_{n \rightarrow m}(t) = j_{n \rightarrow m}^{\text{Tunneling}}(t) + j_{n \rightarrow m}^{\text{Solvent}}(t) + j_{n \rightarrow m}^{\text{Cavity}}(t) \quad (\text{S66})$$

$$j_{n \rightarrow m}^{\text{Tunneling}}(t) = 2 \text{Im} \left((H_S)_{mn} (\rho_S(t))_{nm} \right) \quad (\text{S67})$$

$$j_{n \rightarrow m}^{\text{Solvent}}(t) = \sum_{k \in \text{solvent}} 2 \text{Im} \left(S_{mn} \left[\sqrt{d_k} (\rho_k(t))_{nm} + \sqrt{d_k^*} (\rho_k(t))^*_{mn} \right] \right) + \lambda_k 2 \text{Im} \left((S^2)_{mn} (\rho_S(t))_{nm} \right) \quad (\text{S68})$$

$$+ \frac{1}{2} \left((L_s \rho_S(t))_{mn} (L_s^\dagger)_{nm} + (L_s)_{mn} (\rho_S(t) L_s^\dagger)_{nm} \right) \quad (\text{S69})$$

$$- \frac{1}{2} \left((\rho_S(t) L_s^\dagger)_{mn} (L_s)_{nm} + (L_s^\dagger)_{mn} (L_s \rho_S(t))_{nm} \right) \quad (\text{S610})$$

$$j_{n \rightarrow m}^{\text{Cavity}}(t) = \sum_{k \in \text{cavity}} 2 \text{Im} \left(S_{mn} \left[\sqrt{d_k} (\rho_k(t))_{nm} + \sqrt{d_k^*} (\rho_k(t))^*_{mn} \right] \right) + \lambda_k 2 \text{Im} \left((S^2)_{mn} (\rho_S(t))_{nm} \right) \quad (\text{S611})$$

$$+ \frac{1}{2} \left((L_c \rho_S(t))_{mn} (L_c^\dagger)_{nm} + (L_c)_{mn} (\rho_S(t) L_c^\dagger)_{nm} \right) \quad (\text{S612})$$

$$- \frac{1}{2} \left((\rho_S(t) L_c^\dagger)_{mn} (L_c)_{nm} + (L_c^\dagger)_{mn} (L_c \rho_S(t))_{nm} \right) \quad (\text{S613})$$

where L_s and L_c refer to the jump-operator for the Markovian correction (Sec. S4.2) of the solvent degrees of freedom and lossy cavity degrees of freedom, respectively.

In the main text, when we refer to the value of a probability current, it is the value of $j_{n \rightarrow m}(t)$ at $t = 6$ ps. At this time, the transients have died out and $j_{n \rightarrow m}(t)$ has become largely time-independent. However, $j_{n \rightarrow m}(t)$ still depends on the populations and coherences of $\rho(t)$, so it slowly decays to zero until the equilibrium distribution is reached.

S6.1. Relating probability-currents to reaction rates.

At early times, past the transients but before the population in the reactant well has decayed substantially, the reaction rate (Eq. S53) is approximated by:

$$k \approx -\dot{P}_r(t) \quad (\text{S614})$$

We can relate this basis-independent expectation value to the basis-dependent probability currents, because we have used the localized well states in the visualization and analysis of the kinetics. Most of the spectral weight is contained to these states during the dynamics and as such:

$$P_r(t) = \text{Tr} \left(\hat{P}_r \rho(t) \right) \approx \sum_n^{\text{reactant}} \rho_{nn}(t) \quad (\text{S615})$$

where the sum is only over the states in the reactant well. The approximation stems from the fact that $\hat{P}_r$ is not diagonal in the chosen basis for $\rho(t)$. However, as long as we use these localized states, we only neglect a very small contribution and the approximation remains good. Also, crucially, n index the same basis as in $\dot{\rho}_{nn} = -\sum_m j_{n \rightarrow m}(t)$ (Eq. 14) such that we can write:

$$k \approx \sum_n^{\text{reactant}} \sum_m j_{n \rightarrow m}(t) = \sum_n^{\text{reactant}} \sum_m^{\text{product}} j_{n \rightarrow m}(t) \quad (\text{S616})$$

where the last equality follows from the fact that $j_{n \rightarrow m}$ is anti-symmetric, i.e. the intra-reactant contributions cancel. These arguments can be extended to the case of multiple pathways. Thus, as long as the probability currents are calculated for states that are localized inside the distinct reactant and product regions and the currents are evaluated at a time t where $P_r(t) \approx 1$ has not depreciated too much, the reaction rates are well approximated by the sum of the currents that go across a dividing surface.

S6.2. FGR as perturbative limit of probability currents

Here, we will gain some intuition about the probability currents introduced in Sec. S6 by considering the Born-Markov-Secular approximation and relating the resulting expression to the well known Fermi's Golden Rule. We will begin by considering a generic system-bath setup:

$$H = H_S + \hat{S}\hat{B} + H_B \quad (\text{S617})$$

Since we consider the perturbative limit we have rightfully neglected the reorganization term in H_S . The resulting bath contribution to the probability current will be:

$$j_{n \rightarrow m}^B(t) = 2 \text{Im} \left(S_{mn} \langle n | \text{Tr}_B(\hat{B}\rho(t)) | m \rangle \right) \quad (\text{S618})$$

With $\rho(t)$ the full system plus bath density matrix. Thus, we seek to approximate $\text{Tr}_B(\hat{B}\rho(t))$. In the interaction picture $\rho_I(t) = e^{i(H_S+H_B)t} \rho(t) e^{-i(H_S+H_B)t}$ we may write:

$$\rho_I(t) = \rho_I(t_0) - i \int_{t_0}^t dt' \left[\hat{S}(t') \hat{B}(t'), \rho_I(t') \right] \quad (\text{S619})$$

Then, under the Born-Markov approximation $\rho_I(t') \approx \rho_I^S(t') \otimes \rho_B$ with $\rho_B \propto e^{-\beta H_B}$:

$$\text{Tr}_B(\hat{B}\rho(t)) \approx -i \int_{t_0}^t e^{-iH_S(t-t')} \hat{S} e^{iH_S(t-t')} \rho_S(t) C(t-t') dt' + H.C. \quad (\text{S620})$$

where we've used $\text{Tr}_B(\hat{B}(t)\rho_B) = 0$ and $C(t-t') = \text{Tr}_B(\hat{B}(t)\hat{B}(t')\rho_B)$. We can then write the probability current as:

$$j_{n \rightarrow m}^B(t) \approx 2 \text{Re} \left[- \int_0^{t-t_0} C_S^{nm}(t, \tau) C(\tau) d\tau + \int_0^{t-t_0} \left(C_S^{mn}(t, \tau) C(\tau) \right)^* d\tau \right] \quad (\text{S621})$$

$$C_S^{nm}(t, \tau) = \langle m | \hat{S} | n \rangle \langle n | e^{-iH_S \tau} \hat{S} e^{iH_S \tau} \rho_S(t) | m \rangle \quad (\text{S622})$$

When $|n\rangle$ and $|m\rangle$ are eigenstates of H_S , we get:

$$\langle n | e^{-iH_S(t-t')} \hat{S} e^{iH_S(t-t')} \rho_S(t) | m \rangle = \sum_l e^{-i(E_n - E_l)(t-t')} S_{nl} \rho_{lm}^S(t) \quad (\text{S623})$$

and we can invoke the secular approximation to argue that only the diagonal part of the density matrix contributes to the integral, such that $\sum_l S_{nl} \rho_{lm}^S(t) \approx S_{nm} \rho_{mm}^S(t)$. If we furthermore introduce the noise spectrum $C(\tau) = \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} e^{-i\omega\tau} S_\beta(\omega)$, we recover:

$$j_{n \rightarrow m}^B(t) = \int_{-\infty}^{\infty} S_\beta(\omega) [\mathcal{A}^{mn}(\omega, t) - \mathcal{A}^{nm}(\omega, t)] d\omega \quad (\text{S624})$$

$$\mathcal{A}^{nm}(\omega, t) = \frac{1}{\pi} \text{Re} \left[\int_0^{t-t_0} C_S^{nm}(t, \tau) e^{-i\omega\tau} d\tau \right] \quad (\text{S625})$$

$$= |\langle m | \hat{S} | n \rangle|^2 \rho_{mm}^S(t) \times \underbrace{\frac{1}{\pi} \text{Re} \left[\int_0^{t-t_0} e^{-i(E_n - E_m)\tau} e^{-i\omega\tau} d\tau \right]}_{\int_{-(t-t_0)}^{t-t_0} \frac{d\tau}{2\pi} e^{-i(\omega_{nm} + \omega)\tau}} \quad (\text{S626})$$

$$(\text{S627})$$

such that in the long time limit $t \gg t_0$:

$$j_{n \rightarrow m}^B \approx |S_{nm}|^2 S_\beta(\omega_{nm}) \rho_{nn}^S - |S_{nm}|^2 S_\beta(-\omega_{nm}) \rho_{mm}^S \quad (\text{S628})$$

That is, the probability current from n to m becomes the population in n times the FGR rate from n to m minus the population in m times the FGR rate from m to n (the two rates are related by $S_\beta(-\omega) = e^{-\beta\omega} S_\beta(\omega)$, see Sec. S1). Thus, when the probability current is subject to the same approximations as FGR, the resulting expression is consistent with FGR.

S6.2.1. Broadening of transition frequencies due to phonon bath

The derivation above (and FGR in general) relied on $|n\rangle$ and $|m\rangle$ being eigenstates of H_S . However, in the present case, we are interested in intra-well transitions between localized states, which are not eigenstates of H_S . The reason that FGR can still be applied as is, is because a secondary bath, in this case the solvent, renders these localized states metastable by suppressing the tunneling between the constituent eigenstates (this can be seen in Fig. S4 where the bath-dressed states look like the localized states in Fig. 7a, even though no such post-processing localization took place when visualizing the states in Fig. S4). Furthermore, the secondary bath broadens the transition frequencies of the system by inducing a finite lifetime of constituent states.

To incorporate these ideas into the result above, we may view H_S as an enlarged system that not only contains the system degrees of freedom, but now also incorporates e.g. the phonon degrees of freedom. In order to recover a succinct expression for $C_S^{nm}(t, \tau)$ for this enlarged system, we make the following considerations. The Markov approximation simply says that, when evaluating the integral S620, we may consider the dynamics of $\rho_S(t - \tau) \approx e^{iH_S \tau} \rho_S(t) e^{-iH_S \tau}$ to be closed under H_S (i.e. we disregard the perturbing bath when calculating the induced transition rates). In the current case where H_S contains the phonon bath, the dynamics are not only closed but it will evolve towards the thermal equilibrium distribution of the system plus phonon bath ρ_S^{eq} . Such that in the long time limit and within this approximation $C_S^{nm}(\tau) \approx \text{Tr} \left[e^{iH_S \tau} |m\rangle \langle m| \hat{S} | n \rangle \langle n | e^{-iH_S \tau} \hat{S} \rho_S^{eq} \right]$ which resembles the dipole-dipole correlation function $C_S(\tau)$, except it is restricted to the pathway between n and m : $C_S(\tau) = \text{Tr} \left[e^{iH_S \tau} \hat{S} e^{-iH_S \tau} \hat{S} \rho_S^{eq} \right] = \sum_{nm} C_S^{nm}(\tau)$. Thus, the simplest way to incorporate dissipation, is to broaden the delta function in Eq. S626 by including decay:

$$\mathcal{A}^{nm}(\omega, t) = |\langle m | \hat{S} | n \rangle|^2 \rho_{mm}^S(t) \text{Re} \underbrace{\int_0^{t-t_0} \frac{d\tau}{\pi} e^{-i(\omega_{nm} + \omega)\tau - \Gamma\tau}}_{\frac{1}{\pi} \frac{\Gamma}{(\omega_{nm} + \omega)^2 + \Gamma^2}} \quad (\text{S629})$$

where the inverse lifetime Γ and the possibly renormalized transition frequency ω_{nm} , can be read-off from the part of $C_S(\tau)$ that is due to the $n \rightarrow m$ transition. In our example, Fig. 9, we did not attempt to find Γ and ω_{nm} a priori, but simply found $\Gamma = 30\text{cm}^{-1}$ and $|\omega_{nm}| = 1205\text{cm}^{-1}$ to best explain the results across all θ and ω_c . The value for Γ align well with what has earlier been found for this system³⁵. However, the bare transition frequency is $\approx 1195\text{cm}^{-1}$ indicating that the phonon bath not only broadens the transition but also shifts it slightly.

Supplemental Material S7: Distribution of reaction rates and comparison to isotropic averaging

For the symmetric double well PES, we can write the difference of the population in the two wells (Eq. S57) as $\Delta P(t) = \Delta P_0 e^{-2kt}$. Based on the agreement seen in Fig. 9, we can use the FGR rate in Eq. 28 to study what the reaction rate profile would look like for a given molecule orientation. We can then imagine an ensemble of molecules with random orientations and since the orientation affects the reaction, this yields a distribution of reaction rates⁹⁶. Thus, instead of the average reaction rate $\langle k(\phi) \rangle_\phi$, which was studied in Fig. 9, the primary experimentally relevant entity is arguably:

$$\Delta P(t) = \langle \Delta P(t, \phi) \rangle_\phi = \Delta P_0 \int_{\Omega} p(\Omega) e^{-2k(\Omega)t} d\Omega \quad (\text{S71})$$

where $p(\Omega)$ is the distribution of the molecule orientation relative to the cavity polarization in terms of the solid angle Ω . In the 2-dimensional coplanar model (Sec. IV B), $\Omega = \phi \in [0, 2\pi)$ and $p(\phi) = \frac{1}{2\pi}$. For an isotropic distribution in three dimensional space $\eta_R/\tilde{\eta}_R = \cos(\theta)\cos(\varphi) + \sin(\theta)\sin(\varphi)\cos(\psi)$ and $\int p(\Omega)d\Omega = \frac{1}{4\pi} \int_0^\pi \sin(\varphi)d\varphi \int_0^{2\pi} d\psi$

Since the reaction rate depend on the orientation, the distribution of reaction rates $p(k)$ can be defined by a change of variables:

$$\Delta P(t) = \Delta P_0 \int_{k_{\min}}^{k_{\max}} p(k) e^{-2kt} dk \quad (\text{S72})$$

As an example, consider the harmonic environment without the spectator mode (Eq. 12) and an isotropically distributed orientation. Using Eq. 28, as was done in³³, we schematically recover: $k(\Omega) = k_0 + \cos^2(\varphi)k_1$, where k_0 is the reaction rate outside the cavity and k_1 is the maximal contribution from the cavity. Using the change-of-variables, one recovers the following distribution of reaction rates: $p(k) = \frac{1}{2\sqrt{k_1(k-k_0)}}$ which naturally reproduces the average

reaction rate: $\int_{k_0}^{k_0+k_1} k p(k) dk = k_0 + \frac{1}{3}k_1 = \langle k(\Omega) \rangle_\Omega$

In this kind of simplified scenario, where we consider an ensemble of molecules whose reaction rate is described by the distribution $p(k)$, we could imagine an observer fitting the time-series data $\Delta P(t)$ with a single exponential. The observer would find a rate k_{fit} that lies between the result at early times $\langle k \rangle$ and the result obtained from the long time asymptotics $\langle k^{-1} \rangle^{-1}$ ¹⁹⁶. The effect of a distribution of rates can perhaps be seen in the experimental data presented in e.g. Fig. 2E of Ref. 9. While the errorbars make it difficult to make a precise comparison, the mean-values suggests that there is an initial fast decay and a subsequent slower decay. A fit would then yield a value that lies in between these two rates.

Now, for the example in Sec. IV B, we would like to understand what the distribution of rates look like for both the 2- and 3-dimensional orientational distributions. Figure S3 show these distributions, based on Eq. 28, alongside the resulting fast rate $\langle k \rangle$ and the slow rate $\langle k^{-1} \rangle^{-1}$. We see that these two rates are comparable, so, in this example, $\langle k \rangle$ is representative of what one could expect to obtain from a single-exponential fit to the decay of reactant from an ensemble of molecules with random orientations. Furthermore, the angular distributions lead to qualitatively similar results, so even though $p(\phi) = \frac{1}{2\pi}$ was used in Sec. IV B, the discussion there also applies to the scenario where the molecule orientations are distributed isotropically in 3-dimensions.

Supplemental Material S8: Equilibrium state of the reaction coordinate

For an N-level system coupled to a dissipative environment, such as the one considered in this work and under fairly general conditions⁴⁹, it has been proven that the long-time asymptotics of system observables $\mathcal{O}_S$ is given by:

$$\lim_{t \rightarrow \infty} \text{Tr} \left[e^{-i(H_S + H_{SB} + H_B)t} \rho_S(0) \otimes \frac{e^{-\beta H_B}}{\text{Tr} e^{-\beta H_B}} e^{i(H_S + H_{SB} + H_B)t} \mathcal{O}_S \right] = \text{Tr}_S [\tau_{\text{MF}} \mathcal{O}_S] \quad (\text{S81})$$




FIG. S3. Distribution of reaction rates $p(k)$ for an ensemble of molecules with random orientations, visualized by shaded rate profiles. The rate is calculated from Eq. 28 and the distribution of the molecule orientations is either uniform in the plane containing the cavity polarization (top panel) or isotropic in 3-dimensions (lower panel). The dashed lines represent the minimum and maximum rates and the full colored line is the mean $\langle k \rangle$ while the black line is the asymptotic reaction rate $\langle k^{-1} \rangle^{-1} = \left(\int_0^\infty p(k) k^{-1} dk \right)^{-1}$. Fitting a single exponential to the decay curve of such an ensemble, would yield a reaction rate in between these two limits.

where τ_{MF} is the *mean force Gibbs state*:

$$\tau_{\text{MF}} = \text{Tr}_B \left[\frac{e^{-\beta(H_S + H_{SB} + H_B)}}{\text{Tr} [e^{-\beta(H_S + H_{SB} + H_B)}]} \right] \quad (\text{S82})$$

While proving the same for infinite-dimensional systems with discrete spectra remains an open problem, we have utilized a finite dimensional representation of our system in the dynamical simulation and may apply the above result to describe our observations. The mean force Gibbs state has been calculated perturbatively in the bath coupling⁷⁹:

$$\tau_{\text{MF}} = \tau + \tau_{\text{MF}}^{(2)} + \dots \quad (\text{S83})$$

where $\tau \propto e^{-\beta H_S}$ is the system Gibbs state and $\tau_{\text{MF}}^{(2)}$ is the leading correction. Whether this perturbative expansion applies is governed by:

$$\epsilon := \sqrt{\left| \beta \sum_{nm} |S_{nm}|^2 \frac{e^{-\beta E_n}}{\sum_l e^{-\beta E_l}} D_\beta(E_n - E_m) \right|} \ll 1 \quad (\text{S84})$$

where $\hat{S}$ is the system operator that couples to the bath and the function $D_\beta(\nu)$, which is also used in the expression for $\tau_{\text{MF}}^{(2)}$, is:

$$D_\beta(\nu) = \mathcal{P} \int_0^\infty \frac{d\omega}{\pi} J(\omega) \left(\frac{\nu \coth(\beta\omega/2) + \omega}{\omega^2 - \nu^2} - \frac{1}{\omega} \right) = \frac{\nu}{2} \mathcal{P} \int_{-\infty}^\infty \frac{d\omega}{\pi} \frac{S_\beta(\omega)}{\omega(\omega - \nu)} \quad (\text{S85})$$

which for the decomposed noise spectra (Eq. 22) reduces to:

$$D_\beta(\nu) = \nu \sum_k \text{Im} \left[\frac{d_k}{z_k(\nu + iz_k)} \right] \quad (\text{S86})$$

For the systems and noise spectra considered in this study $\epsilon \sim 0.12$. In case it is of interest to apply this perturbative methodology for stronger coupling, the method known as *reaction coordinate mapping* (see Ref. 63) could prove useful to reduce the perturbative error. The basic idea is to include strongly coupled degrees of freedom in H_S and perturbatively expand in the residual couplings.

By calculating τ_{MF} to second order using the results of Ref. 79, we evaluate the population inside each well $P_{\text{left/middle/right}}^{eq}$ which, for the double well PES, we can compare to the reaction rates through $\frac{k_{\rightarrow}}{k_{\leftarrow}} = \frac{P_{\text{right}}^{eq}}{P_{\text{left}}^{eq}}$. Furthermore, we can visualize τ_{MF} in terms of its eigendecomposition $\tau_{\text{MF}} = U D U^\dagger$. The eigenstates U correspond to the bath dressed vibrational states in thermal equilibrium, with the frequencies given by $\omega_{nm} = -\frac{1}{\beta} (\ln D_n - \ln D_m)$.

S8.1. Mean force Gibbs state for the triple well, Fig. 7

In Figure S4, the effective system (described above) is visualized for the triple well PES coupled to the phonon bath. Compared to Fig. 7a, we see that all the excitations have been somewhat lowered in energy. However, the left well has been favored due to a more significant lowering of the energy of its excited state. More importantly, this is also reflected in the energy of the left wells ground state which has been decreased, while the energy of the right well ground state has been increased, slightly.

An understanding that can explain these observations, is that the environment observes the position ($\hat{S} = \hat{R}$) of the reaction coordinate, which localizes the wavefunction. The reason this favors the left well is because it has lower potential energy. In other words, the kinetic energy is anyway increased by interacting with the environment, so it might as well save some potential energy by residing more in the left well.

We quantify this effect in the second part of figure S4, where we also include the effect of the cavity on the equilibrium population of each well. We see a minor correction from the phonon bath, to the prediction given by the Gibbs state τ (dotted lines). However, the presence of the cavity has nearly no effect on the equilibrium distribution. Especially, we do not see any resonant behavior. The cavity seems to only enhance the effect described above, where the localization of the reaction coordinate favors the deeper left well. This behavior seemed to persist when we considered non-linear dipole moments.




FIG. S4. (left) Visualization of the phonon-dressed vibrational states of the reaction coordinate according to the diagonalization of τ_{MF} described in Sec. S8. (right) Comparison of the population in the left (red), middle (black) and right (blue) wells at equilibrium. The dotted lines are the predictions from the Gibbs state $\tau \propto e^{-\beta H_R}$. The solid lines include the correction from the structured environment via Eq. S83, for which the horizontal line is for outside the cavity while the thicker line includes the cavity contribution.

S8.2. Mean force Gibbs state for the double wells, Fig. 5

For the double well systems, we can compare the predictions of the mean force Gibbs state to those obtained from the dynamical simulation $P^{eq}_{left/right} = \frac{k_{\leftarrow/\rightarrow}}{k_{\leftarrow} + k_{\rightarrow}}$. This is plotted in figure S5 alongside the distribution determined by the Gibbs state τ and its leading bath-induced correction $\tau_{MFG}^{(2)}$. The ω_r refers to the frequency of the right well and the ordering of the subfigures from left to right is exactly the same as in Fig. 5.

Again, we see the same behavior that the narrow well is favored because it has lower potential energy once the wavefunction is localized.

It is difficult to establish quantitative agreement to the numerical predictions via the reaction rates, because the numerical noise is magnified from the mutual division of $k_{\leftarrow}$ and $k_{\rightarrow}$. These numerical predictions display an ω_c dependence, which could be an artifact from this kind of short-time extrapolation to long-time asymptotics. For a proper comparison, one could consider obtaining the steady state directly from the dynamical equation, Eq. 18.




FIG. S5. Comparison of the population in the left (red) and right (blue) wells at equilibrium for the double wells in Fig. 5. The dotted lines are the predictions from the Gibbs state $\tau \propto e^{-\beta H_R}$. The solid lines include the correction from the structured environment via Eq. S83, for which the horizontal line is for outside the cavity while the thicker line includes the cavity contribution to the noise spectrum. The dashed lines represent the empirical prediction from the normalized reaction rates.

Supplemental Material S9: Kinetic and thermodynamic control of the triple well system

The triple well PES gives rise to reversible parallel reactions, which can be described by the rate equation:

$$\begin{pmatrix} \dot{P}_l \\ \dot{P}_0 \\ \dot{P}_r \end{pmatrix} = \begin{pmatrix} -k_{l0} & k_{0l} & 0 \\ k_{l0} & -(k_{0l} + k_{0r}) & k_{r0} \\ 0 & k_{0r} & -k_{r0} \end{pmatrix} \begin{pmatrix} P_l \\ P_0 \\ P_r \end{pmatrix} \quad (\text{S91})$$

with $P_0(t=0) = 1$ and $k_{ij} := k_{i \rightarrow j}$, i.e. this is the same rate equation as Eq. S514. In order to preserve probability, one of the eigenvalues of the matrix are zero and the associated eigenvector is the steady state distribution:

$$\lim_{t \rightarrow \infty} \begin{pmatrix} P_l(t) \\ P_0(t) \\ P_r(t) \end{pmatrix} \propto \begin{pmatrix} k_{0l}k_{r0} \\ k_{l0}k_{r0} \\ k_{l0}k_{0r} \end{pmatrix} \quad (\text{S92})$$

Here, we see why we should not expect a modification of the steady state distribution from simply manipulating the left or right reaction rates, within the linear rate equation model. Because, if either k_{0l} or k_{0r} is scaled by a factor α , the reverse rate k_{l0} , k_{r0} is also scaled by α since they are linked by detailed balance. Then, every well distribution acquires the same scaling, which disappears when accounting for normalization. A simple way to affect the equilibrium distribution, within this model, would be to change the temperature which affects the detailed balance condition and thus the equilibrium populations. The regime in which the product distribution is determined by these equilibrium populations, rather than by the rates of formation, is referred to as thermodynamic control⁹⁷. Sec. S8 describes a way to study cavity-modified thermodynamics of the reaction coordinate, that does not rely on the above simplified argument.

Nonetheless, the timescales of how the distribution reaches the above steady state is determined by the remaining two negative eigenvalues $-\gamma_{\pm}$ of the rate matrix:

$$\gamma_{\pm} = \frac{K_l + K_r}{2} \pm \frac{\Delta}{2} \quad (\text{S93})$$

$$\text{with } K_l = k_{0l} + k_{l0}, \quad K_r = k_{0r} + k_{r0} \quad (\text{S94})$$

$$\text{and } \Delta = \sqrt{(K_l + K_r)^2 - 4(K_l K_r - k_{0l} k_{0r})} \quad (\text{S95})$$

In the irreversible limit $k_{l0} \approx 0 \approx k_{r0}$ the two decay rates become a slow $\gamma_- \approx 0$ and a fast mode $\gamma_+ \approx k_{0l} + k_{0r}$. During the time $\mathcal{O}(\gamma_+^{-1})$ the middle well has depleted into both the left and the right well. The population in either well is determined by the branching ratio $P_l(t \sim \gamma_+^{-1}) \approx \frac{k_{0l}}{k_{0l} + k_{0r}}$, $P_r(t \sim \gamma_+^{-1}) \approx \frac{k_{0r}}{k_{0l} + k_{0r}}$ which constitutes the kinetic yield. The kinetic yield is straightforwardly affected by changes in the reaction rates. The regime in which the product distribution is governed by relative reaction rates rather than equilibrium populations, is generally referred to as kinetic control⁹⁷. Nonetheless, as long as the reaction is not completely irreversible ($k_{l0} \neq 0 \neq k_{r0}$), the distribution will eventually reach the steady state. Accounting for the non-zero backward rates, the kinetic yield becomes $P_l(t \sim \gamma_+^{-1}) \approx \frac{k_{0l}(\gamma_+ - k_{r0})}{\gamma_+(\gamma_+ - \gamma_-)}$, $P_r(t \sim \gamma_+^{-1}) \approx \frac{k_{0r}(\gamma_+ - k_{l0})}{\gamma_+(\gamma_+ - \gamma_-)}$ Figure S6 shows two examples of the dynamics




FIG. S6. Toy example of the population dynamics for the triple well rate equation $P_l \xrightleftharpoons[k_{l0}]{k_{0l}} P_0 \xrightleftharpoons[k_{r0}]{k_{0r}} P_r$. The dotted lines are the equilibrium distributions while the dashed lines are the kinetic yield.

of the population for the reversible parallel rate equation, Eq. S91. The horizontal lines represent the equilibrium distribution (dotted) and the kinetic yield (dashed). The kinetic yield best describe the distribution of the products in the intermediate time window $\mathcal{O}(\gamma_+^{-1}) \lesssim t \lesssim \mathcal{O}(\gamma_-^{-1})$ as defined in Eq. S93. The kinetic yield is affected by the doubling of the rates between the left well and the reactant well whereas the equilibrium distribution remains unaffected. In both examples, the populations slowly evolve towards the equilibrium distribution.

Supplemental Material S10: The phonon bath and Kramers turnover

The phonon spectral density is the same for all simulations. It is of the Drude Lorentz form with

$$J_R^{(DL)}(\omega) = \frac{2\lambda_R\gamma_R\omega}{\gamma_R^2 + \omega^2} \quad \lambda_R = \frac{\eta_\nu M \gamma_R}{2} \times 1000 \text{cm}^{-1} \approx \eta_\nu \times 0.00381157 \text{a.u.} \quad \gamma_R = 200 \text{cm}^{-1} \quad (\text{S101})$$

The reaction rate for the symmetric double well coupled to the phonon bath, were calculated for different coupling strengths η_ν , as seen in Fig. S7. In the limit of vanishing η_ν the state of the system would oscillate between the wells at a rate $\sim 600 \times 10^{-6} \text{fs}^{-1}$. By increasing $0 < \eta_\nu < 0.25$, the coupling to the bath greatly suppresses the tunneling. In this regime, the reaction is limited by the sparse energy transactions with the environment, so by increasing further $0.25 < \eta_\nu < 2$, the bath helps drive the transition until the crossover $\eta_\nu \approx 2$. From this point, the reaction rate decreases due to the increased localization from strongly coupling to the environment.

S10.1. Tensor network simulations in the strong coupling regime

In obtaining the above results, we looked into different tensor network techniques to alleviate the numerical hurdles that arise in the strong coupling regime $3 \lesssim \eta_\nu$. The primary obstacles are that the cutoff of the auxiliary modes need to be increased together with the bond-dimension of the tensor network ansatz.

In an attempt to compress the local Hilbert space dimensions, we applied the Optimized Boson Basis or Local Basis Optimization approach⁹⁸. In short, the three legged tensors of the MPS, which all have high local dimension in this scenario, are decomposed such that the physical index is separated from the two virtual bond indices. This introduces





FIG. S7. Reaction rates for the symmetric double well coupled to the phonon bath for different coupling strengths η_ν . For all other simulations $\eta_\nu = 0.1$ was used (third visible dot from the left).

an intermediate index that connects the two tensors, and the goal is to substantially compress this new bond, such that the computational cost of the MPS representation is reduced.



However, our experience is that such a compression is not achievable for this scenario, and that the bond that connects the two new tensors had to be too large to render the approach useful.

Another approach we tested is the pseudosite method⁹⁹. The bosonic occupation number is represented in binary form, such that a bosonic degree of freedom with a local cutoff of e.g. $64 = 2^6$ is mapped exactly onto six two-dimensional pseudosites. The joint state of these six pseudosites therefore encodes the Fock state of the original bosonic mode. Bosonic operators can be represented in terms of operators acting on the pseudosites, allowing the original bosonic site to be treated within a standard tensor-network framework. By distributing the original large local Hilbert space over several two-dimensional sites, standard tensor-network compression can be applied to the resulting pseudosite representation, potentially reducing the computational cost.



Doing this for each bosonic degree of freedom, we again experienced that this did not reduce the computational cost because the resulting tensor network could not compress enough while still keeping the necessary accuracy.

In the end, we simply simulated the strong-coupling scenarios using 1-site TDVP with GPU acceleration¹⁰⁰. Methods such as the reaction coordinate mapping (see Ref. 63) might also prove also for simulating the dynamics of such scenarios.

Supplemental Material S11: Additional asymmetric double well probability currents

Figure S8 provides a complete overview of the probability currents for the systems shown in Fig. 5.

Supplemental Material S12: Overview of parameter values




FIG. S8. Probability currents for the asymmetric double wells. A negative rate means that the current goes in the opposite direction of the arrow. All currents with magnitude larger than $0.05 \times 10^{-6} \text{fs}^{-1}$ are shown.

Symbol	Description	Value	Unit
Reaction Coordinate PES:			
Symmetric Double Well			
M	Effective mass	1836	a.u.
E_b	Barrier height	2120	cm^{-1}
ω_0	Barrier frequency	866.020	cm^{-1}
$\omega_l = \omega_r$	Reactant and product well frequency	1414.177	cm^{-1}
Asymmetric Double Well			
E_b	Barrier height	2120	cm^{-1}
ω_0	Barrier frequency	866.020	cm^{-1}
ω_l	Reactant well frequency	1414.177	cm^{-1}
ω_r	Product well frequency	{999.914, 707.046, 577.301, 500.077}	cm^{-1}
Triple Well			
E_{ll}	Left well offset	960.660	cm^{-1}
E_l	Left barrier height	2120	cm^{-1}
E_0	Middle well offset	426.963	cm^{-1}
E_r	Right barrier height	2120	cm^{-1}
E_{rr}	Right well offset	266.695	cm^{-1}
ω_{ll}	Left well frequency	2121.320	cm^{-1}
ω_l	Left barrier frequency	866.02	cm^{-1}
ω_0	Middle well frequency	853.927	cm^{-1}
ω_r	Right barrier frequency	866.02	cm^{-1}
ω_{rr}	Right well frequency	533.389	cm^{-1}
Harmonic Environment:			
T	Temperature	300	K
η_c	Light-matter interaction strength	0.1	—
$\tilde{\eta}_R$	Reaction coordinate - cavity coupling	1.0	—
$\tilde{\eta}_Q$	Spectator mode - cavity coupling	0.1	a.u.
c_Q	Reaction coordinate - Spectator mode coupling	2.2326×10^{-5}	a.u.
ω_Q	Spectator mode frequency	1170	cm^{-1}
Cavity loss parameters			
τ_c	Cavity lifetime	100	fs
α	Cavity Markovian parameter	$(1 - e^{-\beta\omega_c})/\tau_c$	fs^{-1}
Reaction coordinate Drude Lorentz bath			
γ_R	Characteristic frequency	200	cm^{-1}
λ_R	Reorganization energy	83.654	cm^{-1}
Spectator mode Drude Lorentz bath			
λ_Q	Reorganization energy	0.147	cm^{-1}
γ_Q	Characteristic frequency	6000	cm^{-1}
Bath decomposition hyper-parameters			
$[\omega_l, \omega_h]$	Frequency window	[-0.99,1]	a.u.
Tolerance	Criterion for convergence	below 10^{-10}	—
Lawson iterations	post-processing step	20	—

TABLE II. Overview of simulation parameters. Note that ω_r and ω_l in the main text refers to the energy spacing between states in the right and left well, respectively. Here, they refer to the curvature of the parabolas that define the potential energy surface in Sec. S3

Description	Value range
Simulation parameters:	
System Hilbert space dimension	12 to 30
Fock space cutoff	4 to 75
Maximum value of $\Gamma\tau$ (for the Markovian compression, Sec. S4.2)	0.01 with typical values of $\mathcal{O}(0.001)$
bond dimension	4 to 600
Tolerance of the Krylov update for the local propagation in TDVP	1e-8
Time step	0.25fs

TABLE III. Additonal simulation parameters. Convergence has been carefully checked for these. The extra large Fock space cutoff and bond dimension was mainly used for the strong coupling scenarios in Sec. S10. Additional information: The probability currents quoted throughout the figures are $j_{n\rightarrow m}(t)$ evaluated at $t = 6\text{ps}$. The initial state of the reaction coordinate is the localized ground state in the reactant well and the total simulation time was 10ps.