

CS215 Assignment-1


210050115 Patil Vipul Sudhir

210050119 Hari Prakash Reddy

1 Question 5.a

Uniform Distribution

- As mentioned in the question, we have to take elements in uniform distribution.
- Used `rand(1,1)` for getting random values with the above specification.
- Found all the errors by finding the sum and taking its average and subtracting the ideal mean(0.5).
- Repeated the above process for 100 times for each N.
- Stored errors for each N, in separate arrays of length 100.
- Combined all the 10 arrays of 100 length into a Matrix.
- Used `boxplot()` for plotting the above matrix in boxplot




Instructions to run the code

Run the WHISKERPLOT.m code in the codes section on MATLAB, to get the above boxplot.

2 Question 5.b

Gaussian Distribution

- As mentioned in the question, we have to take elements in Gaussian distribution.
- Used `normrnd(0,1)` for getting random values with the above specification.
- Found all the errors by finding the sum and taking its average and subtracting the ideal mean(0).
- Repeated the above process for 100 times for each N.
- Stored errors for each N, in separate arrays of length 100.
- Combined all the 10 arrays of 100 length into a Matrix.
- Used `boxplot()` for plotting the above matrix in boxplot



Instructions to run the code

Run the WHISKERPLOT2.m code in the codes section on MATLAB, to get the above boxplot.

3 Question 5.c

Interpretation

- From the graphs,
- When N is low, the error medians are high in both the graphs.
- When $N=5$ the median of errors is higher in Gaussian than those taken from Uniform Distribution.
- As N increases, this trend continued.
- As N increases the median of errors has gone down in both Uniform and Gaussian Distribution.
- As N increased the errors bar has grown narrower and at $N=10^4$ it is almost a single value.