


# SDMs as Geostatistical models

David V. Conesa Guillén




Universitat de València



## *Species distribution modelling (SDMs)*


- SDMs links spatially referenced records of species occurrence with maps of environmental variables in order to **create a statistical model of the relationship** between a **species and its environment**.



- Typical examples: diseases, fish species, plants, animals, etc.
- Typical covariates: elevation, climate, vegetation, human disturbance, temperature, chlorophyll-a, etc.
- Applications: climate change, conservation of species, etc.
- Many options to deal with them. See for instance among many other reviews Rodrigues et al (2023a, 2023b). Our focus here will be deal with them as **Geostatistical Bayesian Hierarchical models**.


# Occurrence of Mackerel in Gulf of Almería

The measures are presences (red) and absences (white)



- 1 Species Distribution Models as Geostatistical models
- 2 Modelling the Ocurrence of a species
- 3 Modelling the Abundance of a species
- 4 Dealing with proportions
- 5 Dealing with counts
- 6 References

# 1 Species Distribution Models as Geostatistical models



## *SDMs as Bayesian Hierarchical Geostatistical models.*

- Different response variables (normal, Poisson, binomial, beta, etc.) depending of the situation.
- **Data Model**  $y = (y_1, \dots, y_n)$  represents the observed values of the corresponding **variable of interest** with mean  $\mu = (\mu_1, \dots, \mu_n)$  at  $n$  locations.
- **Process model** each  $\mu_i$  can be easily linked to a **structured additive predictor**  $\eta_i$  and a **geostatistical spacial random effect**:

$$g(\mu_i) = X_i\beta + W_i, i = 1, \dots, n.$$

The **spatial effect**  $W$  is Gaussian: it is a **Latent Gaussian field**.

- This predictor can incorporate also smooth nonlinear effects of covariates, time trends and seasonal effects, random intercept and slopes as well as temporal random effects.
- **Prior distributions** for the parameters and hyperparameters need to be assigned.


- The resulting hierarchical spatial model **do not** yield to analytical expressions for the posterior distributions of the parameters.
- Posterior distributions of the parameters can be approached by means of MCMC but they are not computationally efficient in this context.
- The Integrated Nested Laplace Approximation (Rue et al., 2009), colloquially known as **INLA**, is a better and faster alternative to approximate the posterior marginals of interest.
- But geostatistics involves working with **continuously indexed Gaussian Fields** and INLA cannot be applied directly.
- Lindgren et al. (2011) proposed an explicit link between *Gaussian fields* and *Gaussian Markov random fields* (GMRF): the **Stochastic Partial Differential Equation approach (SPDE)**.
- GMRFs are discretely indexed → the Markov property makes the precision matrix involved sparse (allowing the use of faster numerical algorithms).

### Final modelling using SPDE

$$\begin{aligned}
 Z_i &\sim \text{Ber}(\pi_i), i = 1, \dots, n \\
 \text{logit}(\pi_i) &= X_i \beta + w_i \\
 \pi(\beta_j) &\sim N(\mu_{\beta_j}, p_{\beta_j}) \\
 w &\sim N(0, Q(\kappa, \tau)) \\
 2\log\kappa &\sim N(\mu_\kappa, p_\kappa) \\
 \log\tau &\sim N(\mu_\tau, p_\tau)
 \end{aligned}$$

- The spatial effect depends on now two different parameters:  $\kappa$  (the range of the effect) and  $\tau$  (the total variance). Prior information for them via pc-priors (Simpson et al., 2017).
- Once the inference is performed, the model is used to predict presence/abundance in unsampled places (providing a map of the species behaviour).
- As usual, model comparison can be done with criterions:
  - ▶ DIC (Spiegelhalter et al., 2002), WAIC (Watanabe, 2010)
  - ▶ LCPO (Gneiting and Raftery, 2007)

## 2 Modelling the Ocurrence of a species




### *Logistic regression models for presences and absences*

When the observed variable is the occurrence (presence/absence), we obtain maps of the posterior predictive distribution of the probability of finding the species (or disease):


- Distribution of Mediterranean horse mackerel in Gulf of Almería (Muñoz et al., 2013).
- Distribution of prevalence of Bovine paramphistomosis in Galicia (González-Warleta et al., 2013).
- Distribution of prevalence of Citrus Black Spot (Martínez-Minaya et al., 2018).
- Distribution of vulnerable (to fishing pressure) elasmobranch species (Pennino et al., 2013).
- Distribution of prevalence of *Xylella fastidiosa* in Alicante (Cendoya et al., 2020).

- In spite of its low commercial value, mackerel plays an important role in the observed transition zone between the Mediterranean and Atlantic sea.
- **Covariates:** geographical coordinates, bathymetry, monthly precipitation, sea surface temperature, chlorophyll-a concentration.
- We used DIC and CPO to compare models with different combinations of the available covariates.
- Final model includes **log(Depth)** (negative effect) and **chlorophyll-a** (positive effect).



## Distribution of Mediterranean horse mackerel in Gulf of Almería (2)

### Spatial effect




Posterior mean and standard deviation of the spatial effect

### Interpretation

- There seems to be a **east-west effect**.
- This was associated a posteriori by local experts with the fact that **the western area of the bay is a protected coastline** with favourable conditions for the species.
- The model provided (unexpectedly) a **quantification of the impact of this protective action** on the Mediterranean horse mackerel.

### Posterior probability of occurrence




Median (top) and quartiles 1 and 3 of  $\pi_i|\text{data}$

*Distribution of three elasmobranch species (Pennino et al., 2013).*


- There is an increasing concern over elasmobranch species because they are **highly vulnerable to fishing** pressure.
- Main predictors of elasmobranch habitats are depth, slope of seabed and type of substrate, followed by temperature and chlorophyll-a.

### Median of the posterior probability of the presence of elasmobranch species



- Species show **different optimum depths**: could indicate a sort of fine-tuned bathymetric segregation, though they coexist on shelf and slope bottoms.
- These maps can be used to **identify sensitive habitats**, with a final aim to improve management and conservation of these vulnerable species.

### 3 Modelling the Abundance of a species




#### *Dealing with positive continuous distributions*


The **observed variable** can be also abundances (normal, gamma, lognormal, etc.):

- **Abundance of hake recruitment** (Paradinas et al., 2015, 2017; Izquierdo et al., 2021), **amount of discards** (Pennino et al., 2014), **abundance of red mullet** (Paradinas et al, 2020), **CPUE biomass indices** (Fuster-Alonso et al. 2024).
- **Milk protein content** of cows with fasciolosis (González-Warleta et al., 2024).
- **Spatial distribution of bioclimatic indices** (Barber et al., 2017).

Mean (left) and standard deviation (right) of the posterior predictive distribution of the Ombrothermic Index



- Bayesian statistics allows us to go further with the bioclimatic indices. We can obtain maps of the **spatial bioclimatic probability distribution** by showing the posterior probability of an index belonging to each climate subtype.
- The map contains the posterior **probability of the nine possible ombootypes** (categories of the Ombothermic Index) that can be observed in the Mediterranean bioclimate at each location.




## 4 | Dealing with proportions

## Beta spatial regression for proportions

The observed variable can also be expressed in terms of proportions:

- Paradinas et al. (2016, 2018) have applied spatial Beta regression models to analyse the proportion of discards, and identified two main fishing-suitable areas based on the proportion of discarded regulated fish proportions.

Posterior mean (left) standard deviation (right) of the spatial component of the regulated species discard proportion




- Martínez-Minaya et al. (2019) have analysed the proportion of *Arabidopsis thaliana* to study the effect of climate change.
- Castilla et al. (2020) have also analysed the ecological, genetic and evolutionary drivers of regional genetic differentiation in *Arabidopsis thaliana*.

## Dirichlet regression for vectors of proportions

The observed variable can be expressed in terms of proportions of different groups:

- Compositional data  $\mathbf{Y} = \{y_{c \times i}\}$  are defined as a vector of values in the interval  $(0,1)$  that satisfies  $\sum_{c=1}^C y_{ci} = 1 : c \in \{1, \dots, C\}$  and  $n \in \{1, \dots, N\}$ .
- In other words, data consisting in proportions (of categories) adding up to 1.
- Martínez-Minaya and Rue (2024) have re-analysed the proportions of *Arabidopsis thaliana* in Peninsula Iberica to study the effect of climate change.
- Figueira et al (2025) have analysed proportions of land-use in Europe.

Posterior predictive distribution of proportions of land use.



## 5 | Dealing with counts




### Regression models for counts


The **observed variable** can also be expressed in terms of counts:

- Paradinas et al. (2020) have considered spatial Negative Binomial regression models to analyse the **number of catchs per 30 minutes** (NPUE) in order to understand the spatiotemporal persistence of fish distributions and so help to define fish hotspots and effective fisheries-restricted areas.

Average spatial distributions of *Mullus barbatus* and *M. surmuletus* occurrence, NPUE and median length (cm) between 2000 and 2016. Blue rectangles represent zoomed sections in next slide.



Zoomed spatial distribution of *Mullus barbatus* and *M. surmuletus* NPUE in Murcia (Zone A), Valencia (Zone B) and Catalonia (Zone C). Hotspots, identified as areas with NPUE above the 90 th percentile, are highlighted in blue



## 6 | References

# References

- ① X. Barber, J. Morales, A. López-Quílez, D. Conesa, A. Martínez-Mayoral and A. Barber (2017). Bayesian hierarchical models for analysing the spatial distribution of bioclimatic indices. *Statistics and Operations Research Transactions*, **41**(2), 277–296.
- ② A. Castilla, B. Méndez-Vigo, A. Marcer, J. Martínez-Minaya, D. Conesa, X. Picó and C. Alonso-Blanco (2020). Ecological, genetic and evolutionary drivers of regional genetic differentiation in *Arabidopsis thaliana*. *BMC Evolutionary Biology*, **20**:71.
- ③ M. Cendoya, J. Martínez-Minaya, V. Dalmau, A. Ferrer, M. Saponari, D. Conesa, A. López-Quílez, A. Vicent (2020). Spatial Bayesian modeling applied to the surveys of *Xylella fastidiosa* in Alicante (Spain) and Apulia (Italy). *Frontiers in Plant Science*, **11**:1204.
- ④ M. Figueira, C. Guarner, D. Conesa, A. López-Quílez, T. Krisztin (2025). Unveiling land use dynamics: Insights from a hierarchical Bayesian spatio-temporal modelling of Compositional Data. *Journal Agricultural, Biological and Env. Statistics*, **in press**.
- ⑤ T. Gneiting and A.E. Raftery (2007). Strictly proper scoring rules, prediction, and estimation. *J. Amer. Statist. Assoc.* **102** (477), 359–378.
- ⑥ M. González-Warleta, S. Lladosa, J.A. Castro-Hermida, A.M. Martínez-Ibeas, D. Conesa, F. Muñoz, A. López-Quílez, Y. Manga-González and M. Mezo, (2013). Bovine paramphistomosis in Galicia (Spain): Prevalence, intensity, aetiology and geospatial distribution of the infection. *Veterinary Parasitology*, **191**(3-4): 252-263.

- ⑦ M. González-Warleta, J.A. Castro-Hermida, M. Figueira, J. López, D. Conesa, A. López-Quílez, F.M. Ubeira, M. Mezo (2024). Bayesian hierarchical modelling of the geospatial distribution of fasciolosis in dairy cattle and the impact on production: Application to the main milk-producing region (Galicia) in Spain. *Veterinary Parasitology*, **325**, 110091.
- ⑧ F. Izquierdo, I. Paradinas, S. Cerviño, D. Conesa, A. Fernández, F. Velasco, I. Preciado, A. Punzón, F. Saborido-Rey and M. G. Pennino (2021). Spatio-temporal assessment of the European hake (*Merluccius merluccius*) recruits in the northern Iberian Peninsula. *Frontiers in Marine Science, section Marine Conservation and Sustainability*, **8**: 614675.
- ⑨ F. Lindgren, H. Rue and J. Lindstrom, (2011). An explicit link between Gaussian fields and Gaussian Markov random fields: the SPDE approach (with discussion). *Journal of the Royal Statistical Society, Series B*, **73**: 423-498.
- ⑩ J. Martínez-Minaya, D. Conesa, A. López-Quílez and A. Vicent (2018). Spatial and climatic factors associated with the geographical distribution of citrus black spot disease in South Africa. A Bayesian latent Gaussian model approach. *European Journal of Plant Pathology*, **151**(4), 991–1007.
- ⑪ J. Martínez-Minaya, D. Conesa, M.J. Fortin, C. Alonso-Blanco, F.X. Picó and A. Marcer (2019). A hierarchical Bayesian Beta regression approach to study the effects of geographical genetic structure and spatial autocorrelation on species distribution range shifts. *Molecular Ecology Resources*, **19**: 929–943.
- ⑫ J. Martínez-Minaya and H. Rue (2024). A flexible Bayesian tool for CoDa mixed models: logistic-normal distribution with Dirichlet covariance. *Statistics and Computing*, **34**, 116.
- ⑬ F. Muñoz, M. G. Pennino, D. Conesa, A. López-Quílez and J.M. Bellido (2013). Estimation and prediction of the spatial occurrence of fish species using Bayesian latent Gaussian models. *Stoch Environ Res Risk Assess*, **27**: 1171–1180.

- ⑯ I. Paradinas, D. Conesa, M.G. Pennino, F. Muñoz, A.M. Fernández, A. López-Quílez and J.M. Bellido (2015). Bayesian spatio-temporal approach to identifying fish nurseries by validating persistence areas. *Marine Ecology Progress Series*, **528**: 245–255.
- ⑰ I. Paradinas, M. Marín, M.G. Pennino, A. López-Quílez, D. Conesa, D. Barreda, M. González and J.M. Bellido (2016). Identifying the best fishing-suitable areas under the new European discard ban. **Selected as Editor's choice**, *ICES Journal of Marine Science*, **73**(10): 2479–2487.
- ⑱ I. Paradinas, D. Conesa, A. López-Quílez, J.M. Bellido (2017). Spatio-Temporal model structures with shared components for semi-continuous species distribution modelling. *Spatial Statistics*, **22**, 434-450.
- ⑲ I. Paradinas, M. G. Pennino, M. Marín, A. López-Quílez, J. M. Bellido and D. Conesa (2018). Modelling spatially sampled proportion processes. *RevStat*, **16**(1): 71–86.
- ⑳ I. Paradinas, D. Conesa, A. López-Quílez, A. Esteban, L. Martín, J.M. Bellido and M.G. Pennino (2020). Identifying Mullidae species (*M. surmuletus* and *M. barbatus*) persistent hot-spots in the Western Mediterranean Sea. *Marine Ecology Progress Series*, **644**: 173–185.
- ㉑ M.G. Pennino, F. Muñoz, D. Conesa, A. López-Quílez, J.M. Bellido (2013). Modelling sensitive elasmobranch habitats. *Journal of Sea Research*, **83**: 209–218.
- ㉒ M.G. Pennino, F. Muñoz, D. Conesa, A. López-Quílez, J. M. Bellido (2014). Bayesian spatio-temporal discard model in a demersal trawl fishery. *Journal of Sea Research*, **90**: 44–53.

- ㉓ L. Rodrigues, E. Kikuchi, P. Kinas, D. Conesa, M.G. Pennino, F. Barbosa and L. Cardoso (2023). Modelling the distribution of marine fishery resources: where are we? *Fish and Fisheries*, **24**(1), 159–175
- ㉔ L. Rodrigues, N. W. Daudt, L.G. Cardoso, P. G. Kinas, D. Conesa and M.G. Pennino (2023). Species distribution modelling in the Southwestern Atlantic Ocean: trends and perspectives. *Ecological Modelling*, **486**, 110514.
- ㉕ H. Rue, S. Martino and N. Chopin (2009). Approximate Bayesian inference for latent Gaussian models by using integrated nested Laplace approximations. *Journal of the Royal Statistical Society, Series B*, **71**(2): 319-392.
- ㉖ D. Simpson, H. Rue, A. Riebler, T.G. Martins and S.H. Sørbye (2017). Penalising model component complexity: A principled, practical approach to constructing priors. *Stat. Sci.* 2017, **32**, 1–28.
- ㉗ D.J. Spiegelhalter, N.G. Best, B.P. Carlin and A. van der Linde (2002). Bayesian measures of model complexity and fit. *Journal of the Royal Statistical Society: Series B*, **64**: 583–616.
- ㉘ S. Watanabe (2010). Asymptotic equivalence of Bayes cross validation and widely applicable information criterion in singular learning theory. *J. Mach. Learn. Res.*, **11**, 3571–3594.