

USEFUL FORMULAE & CONCEPT

ARITHMETIC



CAT QUANTITATIVE APTITUDE



About iQuanta?

Founded in 2015 by Indrajeet Singh, iQuanta began as a Facebook doubt-solving group from a small apartment. With strong expertise in Quants, he earned the title 'Wizard of Quant' by solving thousands of students' queries with clarity and speed. Noticing the lack of quality guidance for CAT aspirants, he officially launched iQuanta to bridge this gap.

Over the years, iQuanta has continued to support over 4 lakh students through its active online communities, offering resources, and peer-driven learning. In the last 5 years, more than 8,000 students have converted IIMs. In 2024, iQuanta produced 500+ 99 percentilers, many of whom made it to top IIMs. iQuanta continues to guide thousands of aspirants on their journey to top B-schools.



Indrajeet Singh
Founder & CEO, iQuanta
Forbes 30 Under 30

ARITHMETIC

Averages

Percentages

SI & CI

Profit & Loss

Mixtures & Alligation

Ratio & Proportion

Time, Speed & Distance

Races

Time & Work

www.iQuanta.in

AVERAGES



$$\text{Average} = \frac{\text{Sum of all the observations}}{\text{Number of observations}}$$

Note: Average of a set of values always lies between minimum to maximum.
i.e., Minimum $\leq$ Average $\leq$ Maximum

Remember the following points about averages:

- i. Average on 1st n natural number = $\frac{n+1}{2}$ (i.e., $n(n+1)/2/n = (n+1)/2$)
- ii. Average of 1st n odd numbers (1, 3, 5,, $2n-1$) = n (i.e., $n^2/n = n$)
- iii. Average of 1st n even numbers (2, 4, 6,, $2n$) = $n+1$
- iv. Average of n terms in Arithmetic Progression = $\frac{n+1}{2}$ th term = middle term, when n is odd.

For example: Average (3, 7, 11, 15, 19, 23, 27) = 15

- v. Average of n terms in Arithmetic Progression = $\frac{\frac{n}{2}\text{th term} + (\frac{n}{2}+1)\text{th term}}{2}$ = Average of middle terms, when n is even.

For example: Average of (6, 10, 14, 18, 22, 26, 30, 34) = $\frac{18+22}{2} = 19$

- v. Average of n terms of an Arithmetic Progression = $\frac{1\text{st term} + \text{last term}}{2}$

$$= \frac{2\text{nd term} + 2\text{nd last term}}{2}$$

$$= \frac{3\text{rd term} + 3\text{rd last term}}{2}$$

and so on.....

- vi. If each term in a set is increased by the value k, then their average is also increased by k.

- vii. If each term in a set is decreased by the value k then their average is also decreased by k.

iQuanta
Building Learning Networks

"iQuanta has been the best source for daily practice. The mentors here are always there to help you out with your queries."


HEENAL RAGHVANI
CAT 99.63%iler

www.iQuanta.in

viii. If each term in a set is multiplied by the value k then their average is also multiplied by k .

ix. If each term in a set is divided by the value k then their average is also divided by k .

Concept of assumed average:

Example: Find the average of 54, 55, 57, and 61

Solution: Let us assume average is 57 (We pick the assumed average in such a way that it lies around the middle of min and max values)

Next, we find deviation of the values from this assumed average.

$$54 - 57 = -3$$

$$55 - 57 = -2$$

$$57 - 57 = 0$$

$$61 - 57 = 4$$

$$\text{Average of all the deviations} = \frac{(-3) + (-2) + (0) + 4}{4} = -0.25$$

Hence, actual average will be $57 - 0.25 = \mathbf{56.75}$.

PERCENTAGES



The percentage equivalents of fractions:

Fraction	Percentage	Fraction	Percentage	Fraction	Percentage
1/2	50%	1/8	12.50%	1/14	7.14%
1/3	33.33%	1/9	11.11%	1/15	6.66%
1/4	25%	1/10	10%	1/16	6.25%
1/5	20%	1/11	9.09%	1/17	5.88%
1/6	16.16%	1/12	8.33%	1/18	5.56%
1/7	14.28%	1/13	7.69%	1/19	5.26%

Important points:

i. x is what % of $y = \frac{x}{y} * 100\%$

ii. $x\%$ of $y = y\%$ of x

iii. x is what % more than $y = \frac{(x-y)}{y} * 100$, $x > y$.

iv. x is what % less than $y = \frac{(y-x)}{y} * 100$, $x < y$.

v. Multiplication Factor (MF)

For example, increase x by 10%:

$$x + 10\% \text{ of } x = x(1+10\%) = x(1 + \frac{10}{100}) = x(1 + 0.1) = 1.1x$$

Here we are basically multiplying x by 1.1 which is known as multiplication factor for 10% increase.

So next time if we want to decrease a quantity by 10% we can directly multiply that quantity by $(1 - 10\%) = 0.9$ and get the desired value.


E.g. Find the resultant value when 100 is increased by 20%, decreased by 30% and then increased by 12%.

$$\text{Solution: } 100 \times 1.2 \times 0.7 \times 1.12 = 94.08$$

Here are a few multiplication factors for the respective % changes:

Percentage Change	Multiplication Factor	Percentage Change	Multiplication Factor
+5%	1.05	-5%	0.95
+10%	1.1	-10%	0.9
+15%	1.15	-15%	0.85
+20%	1.2	-20%	0.8
+25%	1.25	-25%	0.75
+30%	1.3	-30%	0.7
+40%	1.4	-40%	0.6
+50%	1.5	-50%	0.5
+60%	1.6	-60%	0.4

From 74%ile to 99.2%ile after joining iQuanta's CAT Course



Syed Abdus Sattar

Last Year in CAT 2019, I got 74 percentile. That year I started preparing in March and thought would do self preparation. Then I studied for 10-15 days with Arul Sharma but was unable to manage job and work. I got demotivated and left preparation. Bas exam ke din jake paper diya and got 74 percentile and did not even want to attempt the paper after 1 and half hours and then I thought I am not going to waste 1 more year and will prepare sincerely. 2020 me same cycle repeat hui. I started preparing in March with Arul Sharma and left after 10 days. Socha ki I am not giving test year. Then work from home ki wajah se I went to home and I was following iQuanta on Fb. Jab 4.5 months bache the tab my sister and mother asked me to join any coaching and then see whatever happens. I joined in the last batch of iQuanta and started preparing from scratch in the last batch. I attended all the events and practised as much I could after whatever time was left after job. Started giving mocks 2 weeks before CAT 2020. I think if I practised more mera LRD me confidence better hota but still I do not have any regrets. Mocks me marks kharab hi the but CAT tension free hokar diya and I think tough CAT examination helped my cause. I got 99.2 percentile this year. Want to thank Jeet Singh sir and iQuanta for teaching me but more importantly for getting me to start the preparation systematically and get into a schedule

51w Love Reply

8

700+ 99%ilers & 1000s of CAT Toppers




Rishi Mittal



Sabyasachi



Mridul Agarwal




Parikshit Tomar



Guru Charan



Manav Jain



Shikhar Sachdeva



Mihir Kapse



Satyajit Das



Swadesh Rath




Kislay Jha




Scan QR to watch
Topper's Interview

SIMPLE & COMPOUND INTEREST



Difference & Graphical Presentation of Simple and Compound Interest:

- Simple interest is always charged on a sum (original principal) at a particular rate and specified period of time.
- Simple interest for all years is the same.
- Compound interest is charged on principle and accumulated interest.
- Compound interest for all years is different.



SIMPLE INTEREST :

Present 0	R = 10% I = 100	End of 1 st year	R = 10% I = 100	End of 2 nd year	R = 10% I = 100	End of 3 rd year
P = 1000		P = 1000		P = 1000		P = 1000

COMPOUND INTEREST :

Present 0	R = 10% I = 100	End of 1 st year	R = 10% I = 110	End of 2 nd year	R = 10% I = 121	End of 3 rd year
P = 1000		P = 1000+100 P = 1100		P = 1100+110 P = 1210		P = 1210+121 P = 1331

Let P be the principal, R be the interest rate % Per annum, and T be the time period (in years)

Simple Interest :

$$S.I = \frac{P \times R \times T}{100}$$

Compound Interest:

$$C.I = P \left(1 + \frac{R}{100} \right)^T - P$$

$$\text{Amount, } A = P \left(1 + \frac{r}{100} \right)^T$$

If interest is being compounded n-times a year then $A = P \left(1 + \frac{r/n}{100} \right)^{n \times T}$

Difference between Compound Interest and Simple Interest

For Principal = P, Rate of interest per annum = R%

$$= \frac{R}{100}$$

i. Difference between C.I and S.I after 1 year = 0

ii. Difference between C.I and S.I after 2 years =

$$P \left(\frac{R}{100} \right)^2$$

iii. Difference between C.I and S.I after 3 years

$$= P \left(\frac{R}{100} \right)^2 \left(3 + \frac{R}{100} \right)$$

Non Engineer got 99.7%ile in CAT Quant with iQuanta



Aayush Agarwal

1 Jan 2019 · 🌐

Last year I scored 93 percentile in quants and the first thing I told sir before joining iquanta was sir non engineer hu quants me kam reh gaye. This year the only thing I concentrated in quants was sir's classes. For a non engineer like me i never imagined 99.69 in quants sir and this is all because of you. Log kehte h unka kaam h kehna ki iquanta sirf shortcuts pe focus karta h par har shortcut ke pehle basics bataya jaata h. I know I did not live up to your expectations with a overall 98.63 but Indrajeet Singh sir thank you for everything you did for us.



Attached topics: #cattopperssuccesstories



Sarah Singhal and 60 others

1 comment

PROFIT & LOSS



i. Keywords

a. Cost Price (CP): Price of an article at which it is bought.

b. Selling Price (SP): Price of an article at which it is sold.

c. Marked Price (MP): It is also known as listed price or printed price or maximum retail price of an article which is decided by the seller so that he can sell it to a customer and make profit.

d. A Profit is earned when selling price of an article is greater than the cost price of the same article. Profit % is calculated on the cost price.

$$\text{Profit} = \text{SP} - \text{CP}, \text{Profit \%} = \left(\frac{\text{SP} - \text{CP}}{\text{CP}}\right) * 100 = \left(\frac{\text{Profit}}{\text{CP}}\right) * 100.$$

e. A Loss occurs when selling price of an article is lesser than the cost price of the same article. Loss % is also calculated on the cost price.

$$\text{Loss} = \text{CP} - \text{SP}, \text{Loss \%} = \left(\frac{\text{CP} - \text{SP}}{\text{CP}}\right) * 100 = \left(\frac{\text{Loss}}{\text{CP}}\right) * 100.$$



f. Mark up is the difference between marked price and cost price of an article. Mark up % is also calculated on the cost price.

$$\text{Mark up value} = \text{MP} - \text{CP},$$

$$\text{Mark up \%} = \left(\frac{\text{MP} - \text{CP}}{\text{CP}}\right) * 100$$

g. Discount is the difference between marked price and selling price. Discount % is calculated on marked price.

$$\text{Discount} = \text{MP} - \text{SP}, \text{Discount \%} = \left(\frac{\text{MP} - \text{SP}}{\text{MP}}\right) * 100.$$



If there are two articles, one of them is sold at x% profit and other is sold at x% loss then overall profit OR loss % on overall transaction will be

1. 0, when both articles have same CP.

2. Loss = $\frac{x^2}{100}$ %, when both articles have same SP.

iii. Dishonest Seller & Faulty Measurements

These are cases when a shopkeeper uses lesser weights or measurements than what is promised to the customer while selling goods to get some profit out of it.

In these types of situations,

$$\text{Profit \%} = \frac{\text{Claimed weight} - \text{Actual weight}}{\text{Actual weight}} \times 100$$

E.g. A seller sells 800g for the price of 1000g.

$$\text{Profit \%} = (1000 - 800) / 800 \times 100 = 25\%$$

iv. Partnerships:

When n number of people invest $A_1, A_2, A_3, \dots, A_n$ amounts for $T_1, T_2, T_3, \dots, T_n$ time periods in the same business then profits are shared in the ratio $A_1T_1 : A_2T_2 : A_3T_3 : \dots : A_nT_n$.

ALLIGATION & MIXTURE



Alligation: The rule allows us to determine the ratio of quantity in which two or more ingredients at the given price must be combined to produce the desired price.

Rule of Alligation:

If two ingredients are mixed, then

Quantity of cheaper	=	C.P. of dearer - Mean Price
Quantity of dearer		Mean price - C.P. of cheaper

iQuanta CAT 2024 IIM Converts

IIM Ahmedabad Converts



Vedant Chandewar
99.98%iler



Swaraj Pal
99.95%iler



Raunak Das
99.55%iler



Siddhant Kumar
99.42%iler

IIM Bangalore Converts



Mridul Tiwari
99.97%iler



Shaivi Goyal
99.54%iler



Rahul Sinha
99.49%iler



Aman Ehtesham
99.94%iler

IIM Calcutta Converts



Mridul Tiwari
99.98%iler



Sanvie Singhal
99.92%iler




Chitturi Sakshvith
Sanjeev
99.86%iler



Madhumitha
Sriharsha
99.6%iler

And many more

Cross proportion representation:



$$\therefore (\text{Cheaper quantity}) : (\text{Dearer quantity}) = (d - m) : (m - c)$$

Shivam Kumar, IIM ABC Convert (99.95%ler) from iQuanta.




Shivam Kumar
1 hr · 🌐



Holy Trinity (IIM A,B,C) Converted

2020 was a year of uncertainty, CAT was no exception to it. With high hopes, I started my preparation. The initial few mocks were pretty similar to Team India's batting performance against Australia in the 2nd innings of 1st test (India Tour of Australia 2020-21). But initial push from the mentors and the peers helped me gain some momentum. I had been a silent observer but I closely followed the discussion which took place in the paid group. All the mentors were approachable 24*7 and that was the best part. I would get to learn unique approach to tackle the problems everytime I opened the group and the comments. As I said I was a silent spectator, most of my learnings happened through amazing peers and mentors. Every discussions led to new insights which proved to be very helpful. The mocks were pretty similar to the actual level. The analysis helped me reassess my strategies and increase my efficiency. iQuanta 250 & LRDI 70 series helped me buckle up for the final hustle. iQuanta 250 acted as a cherry on top, one place to revise the entire syllabus. The unparalleled hard work put by the team motivated me to take that extra step. And the cumulation of those steps mattered in the end. I would like to thank the entire iQuanta team for their relentless efforts and their selfless service. And to my peers: you are awesome, unknowingly all of you have taught me a lot. Hope all of you do well in your life.

FYI,
Converted IIM Ahmedabad, IIM Bangalore, IIM Calcutta, Kozhikode, Shillong, Indore, IIT B, NITIE !
CAT'20: 99.95%ile



www.iQuanta.in

www.iQuanta.in

Removal and Replacement

Let us say a container has x units of liquid, and y units are removed and replaced by water.


After n operations, the quantity of the pure liquid = $x \left(1 - \frac{y}{x}\right)^n$ units.

- General formula for n operations:**


Final or reduced concentration
= initial concentration $\left(1 - \frac{\text{amount being replaced in each operation}}{\text{total amount}}\right)^n$

- Weighted Average method:**

Two mixtures having weights f_1, f_2 and corresponding averages A_1, A_2 , their weighted average $A_w = \frac{A_1 * f_1 + A_2 * f_2}{f_1 + f_2}$



RATIO & PROPORTION



Properties of ratio:

i. If $0 < \frac{A}{B} < 1$, then $\frac{A}{B} < \frac{A+x}{B+x}$ where $x > 0$.

For example: $\frac{9}{11} < \frac{109}{111}$

ii. If $\frac{A}{B} > 1$, then $\frac{A}{B} > \frac{A+x}{B+x}$ where $x > 0$.

For example: $\frac{17}{13} > \frac{57}{53}$

iii. If $\frac{A}{B} = \frac{C}{D} = \frac{E}{F} = \dots = K$ then

a. $\frac{A+C+E+\dots}{B+D+F+\dots} = K$

b. $\frac{pA+qC+rE+\dots}{pB+qD+rF+\dots} = K$

c. $\frac{pA^n+qC^n+rE^n+\dots}{pB^n+qD^n+rF^n+\dots} = K^n$

Note: In points a, b, and c none of the denominators should be equal to zero.

Merging Ratios

Ratios are merged to compare all the quantities at once. Usually, ratios are merged by making the common term equal or multiplying all the ratios together.

E.g. If $a : b = 3 : 5$ and $b : c = 4 : 5$ then find $a : b : c$.

Solution:

Method – 1: firstly, it can be noticed that b is common in both so we can make it equal in both ratios, LCM (4, 5) = 20.

$a : b = 3 : 5 = 12 : 20$, $b : c = 4 : 5 = 20 : 25$

Hence, $a : b : c = 12 : 20 : 25$.

Method – 2:

$$\frac{a}{c} = \frac{a}{b} * \frac{b}{c}$$

$$\text{or, } \frac{a}{c} = \frac{3}{5} * \frac{4}{5} = \frac{12}{25},$$

when $a = 12$, $b = 20$ and $c = 25$.

Hence, $a : b : c = 12 : 20 : 25$.

This method is very useful when there are more than two ratios to be merged.

Proportion:

i. If a , b and c are in continued proportion, then $\frac{a}{b} = \frac{b}{c}$ and vice-versa is also true.

Here a is known as the first proportion, b is known as the mean proportion and c is known as the third proportion.

ii. If a , b , c and d are in continued proportion, then $\frac{a}{b} = \frac{c}{d}$ and vice-versa.

iii. If $\frac{a}{b} = \frac{c}{d}$, then
 $\frac{a+b}{a-b} = \frac{c+d}{c-d}$ which is
 also known as
 Componendo &
 Dividendo.



iQuanta
Building Learning Networks

CAT PREVIOUS YEAR PAPERS

iQuanta Website has all Past CAT Papers, with detailed solutions in all required format.


- ✓ Topic-wise Past CAT Questions
- ✓ Year-wise Past CAT Questions
- ✓ Take Past CAT as a Mock

Solving these along with **iQuanta CAT Classes** can make an aspirant geared upto at least 95% of his Preparation.

Click here!

www.iQuanta.in

TIME, SPEED & DISTANCE:



General formula: Distance = Speed $\times$ Time

Unit conversion:

Unit of speed = $\frac{\text{km}}{\text{hr}}$

$$1 \frac{\text{km}}{\text{hr}} = \frac{5}{18} \text{ m/sec}, 1 \frac{\text{m}}{\text{sec}} = \frac{18}{5} \text{ km/hr}$$

Proportionality:

1. When distance is constant: $s \propto \frac{1}{t}$ or $t \propto \frac{1}{s}$ so, $st = \text{constant}$, $s_1 t_1 = s_2 t_2$

	A		B		P		Q		R
Speed	3	:	4	:	3	:	4	:	6
Time	$\frac{1}{3}$	:	$\frac{1}{4}$	:	$\frac{1}{3}$	:	$\frac{1}{4}$	:	$\frac{1}{6}$
	4	:	3	:	4	:	3	:	2

Note: If speed is being multiplied by a factor x , then time will be multiplied by $\frac{1}{x}$.

2. When speed is constant: $D \propto t$ or $D_1 : D_2 = t_1 : t_2$

3. When time is constant: $D \propto s$ or $D_1 : D_2 = s_1 : s_2$


Average Speed: Total Distance/Total time taken

Relative speed:

The relative speed of two objects moving with speeds S_1 & S_2 when they are moving in:

1. Same direction, $S_R = S_1 - S_2$, $S_1 > S_2$

2. Opposite direction, $S_R = S_1 + S_2$



Special Cases for Trains:

- When a train passes a pole (or, any stationary object of negligible length), it covers a distance which is equal to its own length.
- When a train passes a platform, it covers a distance which is equal to the sum of the length of the platform and its own length.
- When a train A passes a moving train B, it covers a distance which is equal to the sum of the length of both the trains A and B with relative speed.
- When a train A crosses a stationary train B, it covers a distance which is equal to the sum of the length of both the trains.

Special Cases for Boats:

Speed of Boat = B, Speed of river or stream = R (in still water)

Speed in downstream (D) = B + R

Speed in upstream(U) = B - R

In the case of boats and streams, as the distance is constant in upstream and downstream movements, time taken is inversely proportional to the upstream and downstream speeds.

LINEAR & CIRCULAR RACES:



Linear races:

1. In a race of d meters A beats B by x metres:

$$S_A/S_B = d/(d-x)$$

2. In a race, A beats B by t seconds:

$$S_A/S_B = (t_A + t)/t_A$$



3. In a race, A beats B by x meters or t sec.

$$S_B = \frac{x}{t} \text{ m/sec}$$

Circular Races:

If two people are running on a circular track with speeds in ratio A: B **where A and B are co-prime**, then

- They will meet at A+B distinct points if they are running in opposite direction.
- They will meet at $|A-B|$ distinct points if they are running in same direction



If two people are running on a circular track having perimeter L, with speeds m and n,

- The time for their first meeting = $L/(m+n)$ (when they are running in opposite directions)
- The time for their first meeting = $L/(|m-n|)$ (when they are running in the same direction)
- Time when 3 objects meet for the first time: Find the time taken for any 2 pairs to meet for the first time and then take the LCM of those times.
- Time when 2 objects meet for the first time at the starting point after moving = $\text{LCM}(L/m, L/n)$

Note: LCM of Fractions $\text{LCM}(a/b, c/d) = \text{LCM}(a, c)/\text{HCF}(b, d)$

TIME & WORK



t – time, e – efficiency, a – amount of work

Case 1: When a is constant, $t \propto \frac{1}{e}$ or $e \propto \frac{1}{t}$

Case 2: When t is constant, $a \propto e$ or $e \propto a$

Case 3: When e is constant $t \propto a$ or $a \propto t$

Wages $\propto$ amount of work done by worker

- If A can do a piece of work in n days, then A's 1 day's work = $\frac{1}{n}$
- If A can do a piece of work in x days and B in y days, and together they take z days to complete the work. Then,

$$\frac{1}{z} = \frac{1}{x} + \frac{1}{y}$$

$$\Rightarrow z = \frac{xy}{x+y}$$

Concept of Man-days: If M_1 men can do W_1 work in D_1 days working H_1 hours per day and M_2 men can do W_2 work in D_2 days working H_2 hours per day (where all men work at the same rate), then

$$\frac{M_1 \times D_1 \times H_1}{W_1} = \frac{M_2 \times D_2 \times H_2}{W_2}$$

Points to Remember for Pipe & Cistern questions:

- If we are filling the tank, the Inlet pipes do positive work while the Outlet pipes do negative work.
- If the goal is to empty the tank, the Outlet Pipes do positive work while the Inlet Pipes perform negative work.

iQuanta **vs** OTHER COACHINGS

30,000 IIM Calls from iQuanta
CAT Course

COURSE FEATURES	CAT FULL COURSE	CAT FULL COURSE PRO	OTHER COACHINGS
Live Conceptual Classes	✓	✓	✓
Application Classes	✓	✓	✗
All Year Practice Sessions	✓	✓	✗
Assignments	✓	✓	✓
24*7 Doubt Solving	✓	✓	✗
Full Mocks with Video Solution	✓	✓	✓
Sectionals with Video Solution *	✓	✓	✗
Non-Engineer's Quant *	✓	✓	✗
Engineer's VARC *	✓	✓	✗
LRDI Inception *	✓	✓	✗
Ai Mock Analysis	✓	✓	✗
Special Initiatives * (QA250, LRDI 70)	✓	✓	✗
Peer to Peer Learning	✓	✓	✗
Free 2 Month Crash Course	✓	✓	✗
7500 CAT Qs in IIM ABC Course	✗	✓	✗
IIM ABC Practice Sessions (Daily 12 QA, 3RCs & 3 LRDI Sets)	✗	✓	✗
12 Sets: CAT Level Books *	✗	✓	✗

CHOOSE YOUR CAT PLAN

* New Additions this Year.

Enroll Now, www.iQuanta.in

For CAT Course, Visit: www.iQuanta.in

To Join CAT Preparation Group, [click here](#)

CAT 2025

NEW BATCH LAUNCHED

1300+ Hours
of Videos

500+ Students scored
99%ile in CAT 2024

35 Mocks & 45
Sectional Mocks

Kamal Lohia

CAT 99.99%iler
QA, LRDI Expert

Abhishek Leela Pandey

CAT 99.93%iler,
GMAT 770

Raj Kumar Jha

6 Times CAT
QA 100%iler

Indrajeet Singh

Founder & CEO
Wizard of Quant

Prashant Chadha

9 Times CAT,
VARC 99%iler

Shabana Ma'am

CAT VARC
99.5+%iler

