

1 Norm Approximation

Let $\mathbf{A} \in \mathbb{R}^{m \times n}$, $\mathbf{b} \in \mathbb{R}^m$. The objective is to minimize the residual vector $\mathbf{r} = \mathbf{Ax} - \mathbf{b}$.
The general problem is formulated as,

$$\min_{\mathbf{x} \in \mathbb{R}^n} \|\mathbf{Ax} - \mathbf{b}\|.$$

Depending on the norm chosen, this results in different convex optimization problems with unique statistical interpretations and solution characteristics.

1.0.1 ℓ_1 Norm Approximation (Sum of Absolute Residuals)

$$\|Ax - b\|_1 = \sum_{i=1}^m |(Ax - b)_i|$$

Also known as **Least Absolute Deviations (LAD)**.

- **Robustness:** Unlike the ℓ_2 norm, the ℓ_1 norm places less weight on large residuals (outliers). It yields a robust estimator where the solution tends to minimize the median of the residuals rather than the mean.
- **Sparsity:** In problems like Basis Pursuit (where we minimize $\|x\|_1$ subject to constraints), ℓ_1 induces sparsity. Here, in residual minimization, it allows many residual entries to be exactly zero.

Convex Formulation (Linear Program): The problem is non-differentiable at zero. We reformulate it as a Linear Program (LP) by introducing slack variables $t \in \mathbb{R}^m$:

$$\begin{aligned} & \text{minimize} && \mathbf{1}^T t \\ & \text{subject to} && -t \preceq Ax - b \preceq t \end{aligned}$$

where $\preceq$ denotes component-wise inequality.

1.0.2 ℓ_2 Norm Approximation (Least Squares)

$$\|Ax - b\|_2 = \left(\sum_{i=1}^m (Ax - b)_i^2 \right)^{1/2}$$

Minimizing the ℓ_2 norm is equivalent to minimizing the squared Euclidean norm $\|Ax - b\|_2^2$. This is the standard **Least Squares** problem.

- **Statistical Interpretation:** Corresponds to the Maximum Likelihood Estimator (MLE) under the assumption of Gaussian noise.
- **Sensitivity:** Highly sensitive to outliers because errors are squared.

Convex Formulation (Unconstrained Quadratic): This is a differentiable, unconstrained convex problem with a closed-form analytical solution (assuming A is full rank):

$$x^* = (A^T A)^{-1} A^T b$$

1.0.3 ℓ_∞ Norm Approximation (Chebyshev)

$$\|Ax - b\|_\infty = \max_{1 \leq i \leq m} |(Ax - b)_i|$$

Also known as **Minimax** or **Chebyshev approximation**.

- **Goal:** Minimizes the worst-case residual.
- **Applications:** Critical in control design and engineering where ensuring that error never exceeds a safety threshold is more important than the average error.

Convex Formulation (Linear Program): We introduce a scalar variable $t \in \mathbb{R}$ representing the maximum bound:

$$\begin{aligned} & \text{minimize} && t \\ & \text{subject to} && -t\mathbf{1} \preceq Ax - b \preceq t\mathbf{1} \end{aligned}$$

1.1 Penalty Functions

In many approximation problems, we minimize a sum of penalty functions applied to the residuals $r_i = (Ax - b)_i$. The problem takes the form:

$$\min_x \sum_{i=1}^m \phi(r_i)$$

where $\phi : \mathbb{R} \rightarrow \mathbb{R}$ is a convex penalty function.

1.1.1 Norm Based ($p = 1, 2, \infty$)

$$\|\cdot\|_1, \|\cdot\|_2, \|\cdot\|_\infty.$$

1.1.2 Dead Zone Linear

Functions of the form, for $r \in \mathbb{R}^n$,

$$\sum_i \phi(r_i),$$

where

$$\phi(r) = \max\{|r| - a, 0\}.$$

1.1.3 Log Barrier

$$\phi(u) = \begin{cases} -\log(1 - u^2), & |u| < 1 \\ \infty, & |u| \geq 1 \end{cases}$$

1.1.4 Huber Penalty

2 Moreau Envelope

Definition 2.1 (Moreau Envelope): Given $f : \mathbb{R}^n \rightarrow \mathbb{R}$, we have the Moreau Envelope as

$$f_u(\mathbf{x}) = \inf_{\mathbf{y} \in \mathbb{R}^n} \left\{ f(\mathbf{y}) + \frac{1}{2u} \|\mathbf{x} - \mathbf{y}\|^2 \right\}.$$

Remark 2.2: The above expression is essentially an infimal convolution of $uf : \mathbb{R}^n \rightarrow \mathbb{R}$ and $g : \mathbb{R}^n \rightarrow \mathbb{R}$ where $g = (1/2)\|\cdot\|_2^2$.

Definition 2.3 (Proximal Operator): The proximal operator $\text{prox}_u(\mathbf{x}) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ of f with parameter u is defined as,

$$\text{prox}_u(\mathbf{x}) = \arg \min_{\mathbf{y} \in \mathbb{R}^n} \left\{ f(\mathbf{y}) + \frac{1}{2u} \|\mathbf{x} - \mathbf{y}\|^2 \right\}.$$

2.0.1 Properties of Moreau Envelope

1. $f_u(\mathbf{x})$ is differentiable if f is continuous. Further, the derivative is continuous.
2. $f_u(\mathbf{x})$ is defined for all $\mathbf{x} \in \mathbb{R}^n$, even when f is not.
3. $f_u(\mathbf{x})$ is convex whenever f is convex.
4. If x^* is the unique minimizer of f , then it is the unique minimizer of $f_u(\mathbf{x})$

Remark 2.4: $f_u(\mathbf{x})$ is convex, since it is the min over one variable of a jointly convex function.

Remark 2.5: $f_u(\mathbf{x})$ is continuously smooth.

Remark 2.6:

$$\text{epi} f_u(x) = \text{epi} f(x) \oplus \text{epi} \left(\frac{1}{2u} \|x\|^2 \right)$$

Remark 2.7: x^* is a minima of f if and only if x^* is a minima of f_u .

Remark 2.8:

$$\nabla f_u(\mathbf{x}) = \frac{1}{u}(x - \text{prox}_u(x)).$$

3 Epigraphs

TODO: complete this

Duality

Supporting Hyperplane Theorem

Definition 3.1 (Conjugate of a function): Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$. The conjugate $f^* : \mathbb{R}^n \rightarrow \mathbb{R}$ of f is defined as,

$$f^*(\mathbf{y}) = \sup_{\mathbf{x} \in \text{dom} f} (\mathbf{y}^T \mathbf{x} - f(\mathbf{x})).$$

Remark 3.2: The conjugate of f is also called Legendre transform, polar of f , Fenchel dual.

Remark 3.3: f^* is convex, irrespective of f .

Remark 3.4:

$$-f^*(0) = \min_{\mathbf{x} \in \mathbb{R}^n} f(\mathbf{x})$$

Theorem 3.5 (Fenchel Young inequality):

$$f(\mathbf{x}) + f^*(\mathbf{m}) \geq \mathbf{m}^T \mathbf{x}$$

3.1 Recovering f from f^*

Definition 3.6 (Biconjugate): Let $f^* : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$ be the conjugate of a function $f : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$. The **biconjugate** $f^{**} : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$ is defined as the conjugate of f^* ,

$$f^{**}(\mathbf{x}) = (f^*)^*(\mathbf{x}) = \sup_{\mathbf{y} \in \mathbb{R}^n} (\mathbf{y}^T \mathbf{x} - f^*(\mathbf{y})).$$

Remark 3.7 (Geometric Interpretation): The biconjugate f^{**} constructs the pointwise supremum (upper envelope) of all affine minorants of f . Specifically, the epigraph of f^{**} is the **closed convex hull** of the epigraph of f ,

$$\text{epi } f^{**} = \text{cl}(\text{conv}(\text{epi } f)).$$

Thus, f^{**} represents the "tightest" closed convex function that minorizes f .

3.1.1 Properties of the Biconjugate

Corollary 3.8 (Weak Duality): For any proper function f , the inequality $f^{**}(\mathbf{x}) \leq f(\mathbf{x})$ holds for all $\mathbf{x} \in \mathbb{R}^n$.

Proof. Recall the **Fenchel-Young inequality**, which states that for any $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$:

$$f^*(\mathbf{y}) + f(\mathbf{x}) \geq \mathbf{x}^T \mathbf{y}.$$

Rearranging terms yields:

$$\mathbf{x}^T \mathbf{y} - f^*(\mathbf{y}) \leq f(\mathbf{x}).$$

Since this inequality holds for all $\mathbf{y} \in \mathbb{R}^n$, it must also hold for the supremum over $\mathbf{y}$:

$$\sup_{\mathbf{y} \in \mathbb{R}^n} (\mathbf{x}^T \mathbf{y} - f^*(\mathbf{y})) \leq f(\mathbf{x}).$$

By definition of the biconjugate, this is exactly $f^{**}(\mathbf{x}) \leq f(\mathbf{x})$. □

Theorem 3.9 (Fenchel-Moreau): Let $f : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$ be a proper function. Then $f = f^{**}$ if and only if f is **closed** (lower semi-continuous) and **convex**.

Proof. The forward implication is immediate: since f^{**} is defined as the pointwise supremum of a family of affine functions, it is necessarily closed and convex. Thus, if $f = f^{**}$, f must share these properties.

For the reverse implication, assume f is proper, closed, and convex. By the weak duality lemma, $f^{**}(\mathbf{x}) \leq f(\mathbf{x})$ for all $\mathbf{x} \in \mathbb{R}^n$. It remains to show that $f(\mathbf{x}) \leq f^{**}(\mathbf{x})$.

We proceed by contradiction. Suppose there exists a point $\mathbf{x}_0 \in \mathbb{R}^n$ such that $f^{**}(\mathbf{x}_0) < f(\mathbf{x}_0)$. This implies that the point $(\mathbf{x}_0, f^{**}(\mathbf{x}_0)) \in \mathbb{R}^{n+1}$ lies strictly below the epigraph of f , denoted $\text{epi}(f)$. Since f is closed and convex, $\text{epi}(f)$ is a closed convex set.

By the **Strict Separating Hyperplane Theorem**, there exists a non-zero vector $(\mathbf{y}, \mu) \in \mathbb{R}^n \times \mathbb{R}$ and a scalar $\alpha \in \mathbb{R}$ such that:

$$\mathbf{y}^T \mathbf{x}_0 + \mu f^{**}(\mathbf{x}_0) > \alpha > \mathbf{y}^T \mathbf{x} + \mu t, \quad \forall (\mathbf{x}, t) \in \text{epi}(f).$$

Since $(\mathbf{x}, t) \in \text{epi}(f)$ allows t to be arbitrarily large, we must have $\mu \leq 0$ (otherwise the RHS would be unbounded).

Case 1: Non-vertical separation ($\mu < 0$) Divide the inequality by $-\mu > 0$. Let $\mathbf{m} = \mathbf{y}/(-\mu)$ and $c = \alpha/(-\mu)$. The separation becomes:

$$\mathbf{m}^T \mathbf{x}_0 - f^{**}(\mathbf{x}_0) > c > \mathbf{m}^T \mathbf{x} - t, \quad \forall (\mathbf{x}, t) \in \text{epi}(f).$$

Setting $t = f(\mathbf{x})$ for $\mathbf{x} \in \text{dom}(f)$, we obtain $\mathbf{m}^T \mathbf{x} - f(\mathbf{x}) < c$. Taking the supremum over $\mathbf{x}$:

$$f^*(\mathbf{m}) = \sup_{\mathbf{x}} (\mathbf{m}^T \mathbf{x} - f(\mathbf{x})) \leq c.$$

However, the left side of the separation inequality yields $\mathbf{m}^T \mathbf{x}_0 - c > f^{**}(\mathbf{x}_0)$. This implies that the affine function $h(\mathbf{z}) = \mathbf{m}^T \mathbf{z} - f^*(\mathbf{m})$ satisfies:

$$h(\mathbf{x}_0) = \mathbf{m}^T \mathbf{x}_0 - f^*(\mathbf{m}) \geq \mathbf{m}^T \mathbf{x}_0 - c > f^{**}(\mathbf{x}_0).$$

This contradicts the definition of f^{**} as the supremum of *all* such affine minorants: $f^{**}(\mathbf{x}_0) \geq h(\mathbf{x}_0)$.

Case 2: Vertical separation ($\mu = 0$) If $\mu = 0$, the separation reduces to $\mathbf{y}^T \mathbf{x}_0 > \alpha > \mathbf{y}^T \mathbf{x}$ for all $\mathbf{x} \in \text{dom}(f)$. This means $\mathbf{x}_0$ is strictly separated from the domain of f in $\mathbb{R}^n$. Since f is proper and convex, it admits at least one affine minorant $g(\mathbf{z}) = \mathbf{a}^T \mathbf{z} - b \leq f(\mathbf{z})$. Consider the function $g_k(\mathbf{z}) = g(\mathbf{z}) + k(\mathbf{y}^T \mathbf{z} - \alpha)$. For $\mathbf{x} \in \text{dom}(f)$, since $\mathbf{y}^T \mathbf{x} < \alpha$, the term $k(\mathbf{y}^T \mathbf{x} - \alpha)$ is negative. Thus, $g_k(\mathbf{x}) \leq g(\mathbf{x}) \leq f(\mathbf{x})$ for all $k > 0$. This means g_k is an affine minorant of f for any k . At $\mathbf{x}_0$, however, $\mathbf{y}^T \mathbf{x}_0 > \alpha$, so the term $k(\mathbf{y}^T \mathbf{x}_0 - \alpha) \rightarrow +\infty$ as $k \rightarrow \infty$. Consequently, $f^{**}(\mathbf{x}_0) \geq g_k(\mathbf{x}_0) \rightarrow +\infty$. Since we assumed $f^{**}(\mathbf{x}_0) < f(\mathbf{x}_0) < \infty$ (if $f(\mathbf{x}_0) = \infty$ the inequality holds trivially), this yields a contradiction.

Thus, the assumption $f^{**}(\mathbf{x}_0) < f(\mathbf{x}_0)$ is false. We conclude $f = f^{**}$. $\square$

Remark 3.10: If f is convex but not closed, f^{**} is the lower semi-continuous closure of f . If f is not convex, f^{**} is the convex envelope of f .

4 Duality

Corollary 4.1 (Fenchel-Young-Corollary): Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex and differentiable function. For any $\mathbf{x}, \mathbf{m} \in \mathbb{R}^n$, the following statements are equivalent:

1. $\mathbf{m} = \nabla f(\mathbf{x})$
2. $f(\mathbf{x}) + f^*(\mathbf{m}) = \mathbf{m}^T \mathbf{x}$

Proof. Recall the Fenchel-Young inequality, which states $f(\mathbf{x}) + f^*(\mathbf{m}) \geq \mathbf{m}^T \mathbf{x}$ for all $\mathbf{x}, \mathbf{m}$. We prove the equivalence of the equality condition and the gradient condition.

(1 $\implies$ 2): Assume $\mathbf{m} = \nabla f(\mathbf{x})$. Since f is convex and differentiable, the first-order condition for convexity states that for all $\mathbf{y} \in \mathbb{R}^n$,

$$f(\mathbf{y}) \geq f(\mathbf{x}) + \nabla f(\mathbf{x})^T (\mathbf{y} - \mathbf{x}).$$

Substituting $\mathbf{m} = \nabla f(\mathbf{x})$,

$$f(\mathbf{y}) \geq f(\mathbf{x}) + \mathbf{m}^T \mathbf{y} - \mathbf{m}^T \mathbf{x}.$$

Rearranging terms to isolate $\mathbf{m}^T \mathbf{y} - f(\mathbf{y})$,

$$\mathbf{m}^T \mathbf{y} - f(\mathbf{y}) \leq \mathbf{m}^T \mathbf{x} - f(\mathbf{x}).$$

Taking the supremum of the left-hand side over all $\mathbf{y} \in \mathbb{R}^n$ yields the definition of the conjugate $f^*(\mathbf{m})$,

$$f^*(\mathbf{m}) = \sup_{\mathbf{y}} (\mathbf{m}^T \mathbf{y} - f(\mathbf{y})) \leq \mathbf{m}^T \mathbf{x} - f(\mathbf{x}).$$

This implies $f(\mathbf{x}) + f^*(\mathbf{m}) \leq \mathbf{m}^T \mathbf{x}$. Combining this with the Fenchel-Young inequality (which gives the reverse inequality $\geq$), we obtain equality,

$$f(\mathbf{x}) + f^*(\mathbf{m}) = \mathbf{m}^T \mathbf{x}.$$

(2 $\implies$ 1): Assume $f(\mathbf{x}) + f^*(\mathbf{m}) = \mathbf{m}^T \mathbf{x}$. Rearranging for $f(\mathbf{x})$,

$$f(\mathbf{x}) = \mathbf{m}^T \mathbf{x} - f^*(\mathbf{m}).$$

By definition of the conjugate, $f^*(\mathbf{m}) \geq \mathbf{m}^T \mathbf{y} - f(\mathbf{y})$ for any $\mathbf{y} \in \mathbb{R}^n$. Substituting this inequality into the equation above,

$$f(\mathbf{x}) \leq \mathbf{m}^T \mathbf{x} - (\mathbf{m}^T \mathbf{y} - f(\mathbf{y})).$$

Rearranging terms to compare $f(\mathbf{y})$ and $f(\mathbf{x})$,

$$f(\mathbf{y}) \geq f(\mathbf{x}) + \mathbf{m}^T (\mathbf{y} - \mathbf{x}).$$

This inequality holds for all $\mathbf{y} \in \mathbb{R}^n$. This is precisely the defining inequality for $\mathbf{m}$ being a subgradient of f at $\mathbf{x}$. Since f is differentiable, the subgradient is unique and equal to the gradient. Thus,

$$\mathbf{m} = \nabla f(\mathbf{x}).$$

□

Remark 4.2: Geometrically, this corollary asserts that the vector $\mathbf{m}$ defines a supporting hyperplane to the graph of f at $(\mathbf{x}, f(\mathbf{x}))$ if and only if Fenchel's inequality holds with equality.

4.1 Partial Minimization & Relation To Conjugate

Let $f : \mathbb{R}^n \times \mathbb{R}^k \rightarrow \mathbb{R} \cup \{+\infty\}$ be a function of two variables $(\mathbf{x}, \mathbf{y})$, where $\mathbf{x} \in \mathbb{R}^n$ and $\mathbf{y} \in \mathbb{R}^k$.

Definition 4.3 (Partial Minimization): The partial minimization of f with respect to $\mathbf{x}$ is the function $h : \mathbb{R}^k \rightarrow \mathbb{R} \cup \{+\infty\}$ defined as,

$$h(\mathbf{y}) = \inf_{\mathbf{x} \in \mathbb{R}^n} f(\mathbf{x}, \mathbf{y}).$$

Lemma 4.4 (Preservation of Convexity): If f is jointly convex in $(\mathbf{x}, \mathbf{y})$, then $h(\mathbf{y})$ is a convex function.

Remark 4.5:

$$f^*(\mathbf{m}_1, \mathbf{m}_2) = \sup_{\mathbf{x}, \mathbf{y}} (\mathbf{m}_1^T \mathbf{x} + \mathbf{m}_2^T \mathbf{y} - f(\mathbf{x}, \mathbf{y}))$$

Remark 4.6: $h^*(\mathbf{m}) = f^*(0, \mathbf{m})$

Proof:

$$\begin{aligned} h^*(\mathbf{m}) &= \sup_{\mathbf{y}} (\mathbf{m}^T \mathbf{y} - h(\mathbf{y})) \\ &= \sup_{\mathbf{y}} \left(\mathbf{m}^T \mathbf{y} - \min_{\mathbf{x}} f(\mathbf{x}, \mathbf{y}) \right) \\ &= \sup_{\mathbf{y}} \sup_{\mathbf{x}} (\mathbf{m}^T \mathbf{y} - f(\mathbf{x}, \mathbf{y})) = f^*(0, \mathbf{m}) \end{aligned}$$

Remark 4.7: $\sup [-f^*(0, \mathbf{m})] \leq \inf f(\mathbf{x}, 0)$ i.e. $h^{**}(0) \leq h(0)$

Remark 4.8: Partial minimization appears naturally when eliminating primal variables in dual constructions.

4.1.1 Perturbation Interpretation Of Duality

For any problem $p^* = \min_{\mathbf{x}} f(\mathbf{x})$, "the" dual is constructed by

1. Introduce perturbations $\mathbf{y}$, and extend f such that $f(\mathbf{x}, 0)$ returns the function to minimize
2. the dual is $d^* = \sup [-f^*(0, \mathbf{m})]$

where $\mathbf{m}$ is the dual variable and $\mathbf{x}$ is the primal variable.

We have

$$d^* \leq p^*,$$

under convexity + mild conditions, we have strong duality, i.e. $d^* = p^*$.

5 Lagrangian Duality

Given the optimization problem in standard form:

$$p^* = \inf \{f_0(x) \mid f_i(x) \leq 0, i = 1, \dots, m, x \in \mathcal{D}\}$$

where $\mathcal{D} \subseteq \mathbb{R}^n$ is the intersection of the domains of all functions f_i .

5.1 The Lagrangian and Dual Function

Define the **Lagrangian** $L : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}$ for $\lambda \succeq 0$ as,

$$L(x, \lambda) = f_0(x) + \sum_{i=1}^m \lambda_i f_i(x).$$

The **Lagrange dual function** $g : \mathbb{R}^m \rightarrow \mathbb{R}$ is defined as the pointwise infimum of the Lagrangian over x ,

$$g(\lambda) = \inf_{x \in \mathcal{D}} L(x, \lambda).$$

By the property of pointwise infima, g is a concave function, and it provides a lower bound on the primal optimal value: $g(\lambda) \leq p^*$ for all $\lambda \succeq 0$.

The Dual Problem

The **Lagrange dual problem** seeks the tightest lower bound,

$$\begin{aligned} d^* = \sup_{\lambda} \quad & g(\lambda) \\ \text{s.t.} \quad & \lambda \succeq 0 \end{aligned}$$

The relationship between the primal and dual optimal values is governed by **Weak Duality** ($d^* \leq p^*$). Under constraint qualifications (e.g., Slater's condition) and convexity, **Strong Duality** holds ($d^* = p^*$).

6 Shadow Price Interpretation of Duality

7 Game Theoretic Interpretation of Duality


8 KKT Conditions

Consider a standard optimization problem with strong duality and differentiable constraints. Let x^* be the primal optima and λ^* be the dual optima.

Definition 8.1 (KKT Conditions): *The following statements constitute the KKT conditions,*

1. $f_i(x^*) \leq 0$ & $h_j(x^*) = 0, \forall i, j$ (Primal feasibility).
2. $\lambda_i^* \geq 0, \forall i$ (Dual feasibility).
3. $\lambda_i^* f_i(x^*) = 0, \forall i$ (Complementary slackness).
4. $\nabla_x L(x^*, \lambda) = 0$.

9 Mechanistic Interpretation of KKT Conditions



There's maybe no need for this.

Yeah, there's no need for this.

10 Multicriterion Optimization

Some shit. Write it.

11 Moreau Envelope (Continued)

Continued from [here](#).

Example:

$$p^* = \min_{\mathbf{x}} \|\mathbf{Ax} - \mathbf{b}\|_2 \quad g(x) = p^*$$

Reframe as:

$$\begin{aligned} \min_{\mathbf{x}, \mathbf{y}} \quad & \|\mathbf{y}\|_2 \\ \text{s.t.} \quad & \mathbf{Ax} - \mathbf{b} = \mathbf{y} \end{aligned}$$

Lagrangian:

$$L(\mathbf{x}, \mathbf{y}, \mathbf{v}) = \|\mathbf{y}\|_2 + \mathbf{v}^T (\mathbf{Ax} - \mathbf{b} - \mathbf{y})$$

Dual Function:

$$\begin{aligned} g(\mathbf{v}) &= \inf_{\mathbf{x}, \mathbf{y}} [\|\mathbf{y}\|_2 + \mathbf{v}^T (\mathbf{Ax} - \mathbf{b} - \mathbf{y})] \\ &= -\mathbf{v}^T \mathbf{b} + \left[\inf_{\mathbf{x}, \mathbf{y}} \left(\underbrace{\|\mathbf{y}\|_2 - \mathbf{v}^T \mathbf{y}}_{\substack{-\infty \text{ if } \|\mathbf{v}\|_2 > 1 \\ 0 \text{ otherwise}}} \right) + \left(\underbrace{(\mathbf{A}^T \mathbf{v})^T \mathbf{x}}_{\substack{0 \text{ if } \mathbf{A}^T \mathbf{v} = 0 \\ -\infty \text{ otherwise}}} \right) \right] \end{aligned}$$

Result:

$$g(\mathbf{v}) = \begin{cases} -\mathbf{v}^T \mathbf{b} & \text{if } \mathbf{A}^T \mathbf{v} = 0 \text{ \& } \|\mathbf{v}\|_2 \leq 1 \\ -\infty & \text{otherwise} \end{cases}$$

11.1 Dual of Moreau envelope

Let $f : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$ be a proper, closed, convex function. Recall the Moreau envelope $f_u(x)$ with parameter $u > 0$ is defined as the infimal convolution of f and a quadratic kernel,

$$f_u(x) = \inf_y \left\{ f(y) + \frac{1}{2u} \|x - y\|^2 \right\}.$$

To derive its dual representation, we rewrite this as a constrained minimization problem by introducing an auxiliary variable z

$$\begin{aligned} \min_{y, z} \quad & f(y) + \frac{1}{2u} \|z\|^2 \\ \text{s.t.} \quad & x - y = z. \end{aligned}$$

The dual function $g(v)$ is given by

$$g(v) = \langle v, x \rangle - f^*(v) - \frac{u}{2} \|v\|^2,$$

which leads to the identity

$$f_u(x) = \left(f^* + \frac{u}{2} \|\cdot\|^2 \right)^*(x).$$

Proof. We form the Lagrangian $L(y, z, v)$ with dual variable $v \in \mathbb{R}^n$,

$$L(y, z, v) = f(y) + \frac{1}{2u} \|z\|^2 + \langle v, x - y - z \rangle.$$

The Lagrange dual function $g(v)$ is obtained by minimizing the Lagrangian with respect to the primal variables y and z ,

$$\begin{aligned} g(v) &= \inf_{y, z} L(y, z, v) \\ &= \inf_{y, z} \left\{ f(y) + \frac{1}{2u} \|z\|^2 + \langle v, x \rangle - \langle v, y \rangle - \langle v, z \rangle \right\} \\ &= \langle v, x \rangle + \inf_y \{ f(y) - \langle v, y \rangle \} + \inf_z \left\{ \frac{1}{2u} \|z\|^2 - \langle v, z \rangle \right\}. \end{aligned}$$

We evaluate the infima term-wise:

1. Term involving y : Using the definition of the Fenchel conjugate, $f^*(v) = \sup_y \{ \langle v, y \rangle - f(y) \}$, we have,

$$\inf_y \{ f(y) - \langle v, y \rangle \} = -\sup_y \{ \langle v, y \rangle - f(y) \} = -f^*(v).$$

2. Term involving z : This is the conjugate of the quadratic function $h(z) = \frac{1}{2u} \|z\|^2$. Setting the gradient with respect to z to zero,

$$\nabla_z \left(\frac{1}{2u} \|z\|^2 - \langle v, z \rangle \right) = \frac{z}{u} - v = 0 \implies z = uv.$$

Substituting $z = uv$ back into the expression,

$$\frac{1}{2u}\|uv\|^2 - \langle v, uv \rangle = \frac{u}{2}\|v\|^2 - u\|v\|^2 = -\frac{u}{2}\|v\|^2.$$

Combining these, the dual function

$$g(v) = \langle v, x \rangle - f^*(v) - \frac{u}{2}\|v\|^2.$$

The dual problem consists of maximizing $g(v)$. Since f is proper, closed, and convex, strong duality holds ($p^* = d^*$), and we can express the primal optimum as the supremum of the dual:

$$\begin{aligned} f_u(x) &= \sup_v g(v) \\ &= \sup_v \left\{ \langle v, x \rangle - \left(f^*(v) + \frac{u}{2}\|v\|^2 \right) \right\}. \end{aligned}$$

Recognizing the right-hand side as the Fenchel conjugate of the function $\phi(v) = f^*(v) + \frac{u}{2}\|v\|^2$ evaluated at x , we conclude:

$$f_u(x) = \left(f^* + \frac{u}{2}\|\cdot\|^2 \right)^* (x).$$

□

11.2 Moreau Decomposition