

Math 239

Vishesh Gupta

December 2, 2017

Contents

1	Combinatorial Analysis	5
1.1	Introduction	5
1.2	Sums and Products	5
1.3	Binomial Coefficients	5
1.4	Bijections(One-to-One Correspondence)	5
1.5	Combinatorial Proofs	5
1.6	Generating Series	5
1.7	Formal Power Series	5
1.8	The Sum and Product Lemmas	5
2	Compositions and Strings	7
2.1	Compositions of an Integer	7
2.2	Subsets with Restrictions	7
2.3	Binary Strings	7
2.4	Unambiguous Expressions	7
2.5	Some Decomposition Rules	7
2.6	Sum and Product Lemma Rules for Strings	7
2.7	Decomposition Using Blocks	7
2.8	Recursive Definition of Binary Strings	7
3	Recurrences, Binary Trees and Sorting	9
3.1	Coefficients of Rational Functions	9
3.2	Solutions to Recurrence Equations	9
4	Introduction to Graph Theory	11
4.1	Definitions	11
4.2	Isomorphism	11
4.3	Degree	12
4.4	Bipartite Graphs	12
4.4.1	N - Dimensional Cube (Q_n)	12
4.5	Connectness	13
4.6	Eulerian Circuits	13
4.7	Bridges	13

5	Trees	15
5.1	Trees	15
5.2	Spanning Trees	15
5.3	Characterizing Bipartite Graphs	15
5.4	Minimum Spanning Trees	15
6	Planar Graphs	17
6.1	Planarity	17
6.2	Euler's Formula	17
6.3	Platonic Solids	17
6.4	Colouring and Planar Graphs	18
6.4.1	Six Colour Theorem	19
6.4.2	Five Colour Theorem	19
6.4.3	$K_{3,3}$ is not planar	20
6.5	Kuratowski's Theorem:	20
7	Matchings	21
7.1	Matching or Matching in Graphs	21
7.1.1	Covers or Covering the edges of a graph	21
7.1.2	Augmented Paths	22
7.2	Konig's Theorem	24
7.3	Applications of Konig's Theorem	24
7.4	System of Distinct Representatives	24

Chapter 1

Combinatorial Analysis

1.1 Introduction

1.2 Sums and Products

1.3 Binomial Coefficients

1.4 Bijections(One-to-One Correspondence)

1.5 Combinatorial Proofs

1.6 Generating Series

1.7 Formal Power Series

1.8 The Sum and Product Lemmas

Chapter 2

Compositions and Strings

- 2.1 Compositions of an Integer
- 2.2 Subsets with Restrictions
- 2.3 Binary Strings
- 2.4 Unambiguous Expressions
- 2.5 Some Decomposition Rules
- 2.6 Sum and Product Lemma Rules for Strings
- 2.7 Decomposition Using Blocks
- 2.8 Recursive Definition of Binary Strings

Chapter 3

Recurrences, Binary Trees and Sorting

3.1 Coefficients of Rational Functions

3.2 Solutions to Recurrence Equations

Chapter 4

Introduction to Graph Theory

4.1 Definitions

A graph is an ordered pair $\{V(G), E(G)\}$ of finite sets $V(G)$ and $E(G)$ such that E is a subset of the set of unordered pairs of elements from $V(G)$.

If $\{V(G), E(G)\}$ is a graph then vertices v, w are adjacent in the graph G if $\{v, w\} \in E$.

The degree of a vertex, v , is the number of vertices that are adjacent to v , and is denoted by $d(v)$ or $\deg(v)$

A graph G is considered r -regular if every vertex has a degree r .

4.2 Isomorphism

Two graphs $\{V(G), E(G)\}$ and $\{V'(G), E'(G)\}$ are isomorphic if there is a bijection

$$\phi : V \rightarrow V'$$

such that $E'(G) = \{\{\phi(u), \phi(v) \mid \{u, v\} \in E\}$.

Examples

- Empty Graph $(V(G), \emptyset)$
- Complete Graph $(V, \binom{V}{2})$
- Path Graph
- Null Graph $\{\emptyset, \emptyset\}$
- Cycle Graph, denoted by C_n :

Theorem: Handshake Lemma

Suppose $\{V(G), E(G)\}$ is a graph, then $\sum_{v \in V(G)} \deg(v) = 2|E(G)|$.

$$\begin{aligned}
 \textbf{Proof: } \sum_{v \in V(G)} \deg(v) &= \sum_{v \in V(G)} |\{u \mid \{u, v\} \in E(G)\}| \\
 &= \sum_{v \in V(G)} |\{e \in E(G) \mid v \in e\}| \\
 &= \sum_{e \in E(G)} \sum_{v \in V(G)} 1 \\
 &= \sum_{e \in E(G)} 2 = 2|E(G)|
 \end{aligned}$$

4.3 Degree**4.4 Bipartite Graphs**

A graph G is bipartite if $V(G)$ can be partitioned into two parts X, Y such that each edge has one vertex in X and one vertex in Y .

$$\begin{aligned}
 V &= X \cup Y \\
 \phi &= X \cap Y
 \end{aligned}$$

Empty Graphs are bipartite.

FACT: Even cycles are bipartite.

FACT: If H is a subgraph of a bipartite graph G , then H is bipartite.

Implication

If G has a subgraph that is an odd cycle, then G is not a bipartite graph.

Two More examples of Bipartite Graph

We already showed all the paths and all even cycles are bipartite.

$K_{p,q}$ = Complete bipartite graphs of p and q vertices.

How many vertices? $p+q$

What are the degrees? q, p

How many edges? $p \cdot q$

4.4.1 N - Dimensional Cube (Q_n)

$V(G)$ = All binary strings of length n

$E(G)$ = Two strings that differ in exactly one place.

Q_n has 2^n vertices. They have degrees, all n , and number of edges is $n2^{n-1}$

FACT: Let G be a bipartite graph with bipartition (X,Y) , i.e, $V(G) = X \cup Y$ and $X \cap Y = \phi$, and every edge consists of vertex in X and a vertex in Y .

Then $|E(G)| = \sum_{v \in X} \deg(v) = \sum_{v \in Y} \deg(v)$.

X = Odd # 1's and Y = Even # 1's

4.5 Connectness

Last time, we agreed that K_n is connected for any empty graph ≥ 2 is disconnected.

- A graph G is connected if, for every two vertices v,u there is a path in G joining u & v .
- A graph is disconnected if it is not connected. That means there exists a pair of vertices, u,v such that no path in the graph joins u and v .
- How do we prove that Q_{758} is connected?
- How do we prove some path is disconnected?

4.6 Eulerian Circuits

4.7 Bridges

Chapter 5

Trees

5.1 Trees

5.2 Spanning Trees

5.3 Characterizing Bipartite Graphs

5.4 Minimum Spanning Trees

Chapter 6

Planar Graphs

6.1 Planarity

6.2 Euler's Formula

Theorem: **Proof:**

6.3 Platonic Solids

Images of Tetrahedron, octahedron, hexahedron, icosahedron, dodecahedron

Theorem: Let d and d^* be integers, both at least 3. If there is a planar graph such that all the vertices have degree d & all faces of length d^* , then $(d, d^*) \in \{(3, 3), (3, 4), (3, 5), (4, 3), (5, 3)\}$

Proof: Handshake Lemma says $d|V(G)| = 2|E(G)|$. For faces, let r be the number of faces. Then Handshaking lemma for faces says $d^*r = 2|E(G)|$. Also we have Euler's formula:

$$p - q + r = 2$$

Therefore,

$$\begin{aligned} \frac{2}{d}|E(G)| - |E(G)| + \frac{2}{d^*}|E(G)| &= 2. \\ \frac{E(G)}{dd^*}[2d^* - dd^* + 2d] &= 2. \\ dd^* - 2d - 2d^* &= \frac{-2dd^*}{E(G)} \\ (d-2)(d^*-2) - 4 &= \frac{-2dd^*}{E(G)} \\ (d-2)(d^*-2) &= 4 - \frac{2dd^*}{E(G)} \end{aligned}$$

What positive integer pairs a, b have $a, b < 4$? where $a = (d-2)$ and $b = (d^*-2)$
 $\rightarrow \{(1, 1), (1, 2), (1, 3), (2, 1), (3, 1)\}$

So now when $a, b \in A$, we get (d, d^*) such that $\{(3, 3), (3, 4), (3, 5), (4, 3), (5, 3)\}$

Let us take $d = 3$ and $d^* = 3$;

$\frac{E(G)}{9}(6-9+6) = 2$, simplifying and solving for $E(G)$ we get 6. Then $3V(G) = 2E(G)$ $V(G) = 4$ and hence $G = K_4$

Let us take $d = 3$ and $d^* = 5$;

$\frac{E(G)}{15}(6-15+10) = 2$, simplifying and solving for $E(G)$ we get 30. Then $3V(G) = 2E(G)$.
 $V(G) = 40$, $r = 12$

Let us take $d = 3$ and $d^* = 4$;

$V(G) = 8$, $E(G) = 12$ and $r = 6$ Should be the same as hexahedron

△ Joining the vertex of top triangle with bottom triangle

□□

△

Colouring Planar Graphs Adjacent vertices get different colors

How many colours for K_n ? n If G is connected planar graph then G has a vertex with degree ≤ 5

$$E(G) \leq 3V(G) - 6$$

Handshaking Lemma: $\sum \deg(V(G)) = 2E(G) \leq 6V(G) - 12$


6.4 Colouring and Planar Graphs

Colouring: assigns a colour to each vertex so that adjacent vertices get different colours.


1) Every bipartition in a bipartite graph can be coloured with two colours. Use different colour on each part of bipartition.

2) K_n can only be coloured with n colours.


Figures




Square




Pentagon



Some graph ———





K 100 K5

Peterson Graph Cross-Cover graph

6.4.1 Six Colour Theorem

Theorem: A planar graph can be coloured with six colours.

1) Every planar graph has a vertex with at most 5.

Pf: Let C be a component of a planar graph G . If $V(C) \leq 6$, then every vertex of G has a degree of at most 5. If $V(C) \geq 6$, then because it is a connected planar graph, we know that $E(C) \leq 3V(C) - 6$. Therefore the average degree is $\sum_{v \in V(C)} \frac{\deg(v)}{V(C)} \leq \frac{6V(C) - 12}{V(C)} = 6 - \frac{12}{V(C)} < 6$. There is a vertex v of C whose degree is at most the average. Thus $\deg(v) < 6$ and so $\deg(v) \leq 5$.

2) To colour G , we proceed with induction on $V(G)$

Pf: If $V(G) = 1$, then G can be coloured with 1 (and therefore 6) colours.

For the induction step, choose a vertex V of G with degree at most 5. Colour $G \setminus V$ with at most 6 colours (using inductive assumption). The neighbours are coloured by the colouring of $G \setminus V$

Proof: Because $\deg(v) \leq 5$, at most five of the 6 colours are used as colours for the neighbours of v . Thus there is one of the six colours not used to colour a neighbour of v , we can use this colour to colour v . Hence proved.

6.4.2 Five Colour Theorem

Theorem: A planar graph can be coloured with five colours.

1) Every planar graph has a vertex with at most 5.

Pf: Let C be a component of a planar graph G . If $V(C) \leq 6$, then every vertex of G has a degree of at most 5. If $V(C) \geq 6$, then because it is a connected planar graph, we know that $E(C) \leq 3V(C) - 6$. Therefore the average degree is $\sum_{v \in V(C)} \frac{\deg(v)}{V(C)} \leq \frac{6V(C) - 12}{V(C)} = 6 - \frac{12}{V(C)} < 6$. There is a vertex v of C whose degree is at most the average. Thus $\deg(v) < 6$ and so $\deg(v) \leq 5$.

2) To colour G , we proceed with induction on $V(G)$

Pf: If $V(G) = 1$, then G can be coloured with 1 (and therefore 6) colours.

For the induction step, choose a vertex V of G with degree at most 5. Colour $G \setminus V$ with at most 6 colours (using inductive assumption). The neighbours are coloured by the colouring of $G \setminus V$

Problem: $\text{Deg}(v) = 5$. Because K_5 is not planar, some two of the neighbours of v are not adjacent.

Solution: Let there be u and w . These two are not adjacent. Delete the edges from u to the other three neighbours of v & contract to v the edge uv and v as to create a smaller graph G' .

Proof: Continuing further from the solution, we see that by colouring G' with 5 colours by induction and colouring u and w , with the colour of the contracted vertex, every one else with their colour in G' . Since u, w have the same colour, neighbours of u and v have at most 4 colours in $G \setminus v$. This leaves the fifth colour for v .

6.4.3 $K_{3,3}$ is not planar

Lemma: If a planar graph has no cycle and has at least three vertices, then every face boundary has length ≥ 4 .

Corollary: If G is a connected planar graph with ≥ 3 vertices and no 3 cycle, then $E(G) \leq 2V(G)$.

So by using this above corollary $K_{3,3}$ is not planar

6.5 Kuratowski's Theorem:

A graph is not planar if and only if it contains a subdivision of either K_5 or $K_{3,3}$
Fig 1 and Fig 2 to be embedded here.

Chapter 7

Matchings

7.1 Matching or Matching in Graphs

A *matching* in a graph G is a set M of edges of G , no two of which are incident with a common vertex.

Fig 3 and Fig 4

For any matching M in a graph, $2|M| \leq |V(G)|$.

A *perfect Matching* in a graph G is a matching M in G such that $2 = |V(G)|$

The empty graph with at least one vertex has no perfect matching. Notice ϕ is a matching.

Note: Even cycles have perfect matching. If $|V(G)|$ is odd then G does not have a perfect matching

In a $K_{p,q}$ graph we get the $\min\{p, q\}$ giving us the largest matching size.

7.1.1 Covers or Covering the edges of a graph

A cover of a graph G is a set C of vertices such that every edge is incident with at least one vertex in C .

Fact: If M is a matching and C is a cover in G , then $|M| \leq |C|$.

In a *Cycle* of length n :

the size of a largest matching is $\lfloor \frac{n}{2} \rfloor$

the size of a smallest cover is $\lceil \frac{n}{2} \rceil$

In a K_n graph:

the size of the matching $\lfloor \frac{n}{2} \rfloor$

the cover for the cycle is: $n-1$

In a $K_{p,q}$ complete graph (with $p \leq q$):

the size of the matching is: p

the size of the cover is: p

Lemma: If M is a matching and C is a cover then $|M| \leq |C|$.

Proof: Let $e = uv$ be an edge in M . By definition, either u or v is in C (or both), without loss of generality, assume it is u . Let e' be a different edge in M . One of the endpoints of e' (w) is in C . Can $u = w$? By definition of the matching, e and e' share no endpoints, so $u \neq w$. There are $|M|$ edges in M . Each yields a unique vertex in $C \rightarrow |M| \leq |C|$

Maximum matching is a largest matching G . (No matching exist with more edges)

If the size of a maximum matching is $\frac{n}{2}$, then it is a perfect

Minimum cover is a smallest cover in G .

Lemma: If M is a matching and C is a $|M| = |C|$ then, M is a maximum matching and C is the minimum cover.

Proof: Let M^* be a maximum matching, then $|M^*| \leq |C| = |M|$. M has at least as many edges as M^* (which is a maximum matching). So M is also a maximum matching. Let C^* be a minimum cover. $|C^*| \geq |M| = |C|$. So C is also a minimum cover.

Lemma: If M has an augmenting path then it is not a maximum matching

Proof: Let P be the augmenting path. The edges alternate between being in M and not being in M . The endpoints are unsaturated. More edges of P are not in M than are in M (one more).

Replace the edges in $M \cap P$ with the edges in P set difference M . New matching M' . If $v \in P$ is saturated by M . We removing the edge in M is incident to v and replacing it with one in M' . If $v \in P$ is unsaturated, we only add one edge incident to v to form the matching M' . M' is a matching $|M'| = |M| + 1$ is not a maximum matching.

7.1.2 Augmented Paths

An alternating Path is a path in G where the edges alternate between being in M and not being in M . An augmented Path is an alternating path where its endpoints are unsaturated.

Odd length

First + Last edge not in the matching

A vertex is saturated by a matching M if it is incident to an edge in M . Otherwise it is unsaturated which is also called exposed. In a perfect matching, all vertices are saturated. If there's no perfect matching, at least one vertex is unsaturated in any matchings.

Lemma: Let G be a graph with matching M . If there is a M -augmented path in G , then M is not a maximum matching (swap the edges of the ones in M with the edges which are not in M)

Lemma: If M is not a max matching then it has an augmenting path. Will be done in tutorial
Let G be a graph with bipartition A, B and let M be a matching of G .

- Let X_0 be the set of vertices in A not saturated by M .
- Let Z be the set of vertices in G joined to a vertex in X_0 by an alternating path.
- If there is an augmenting path use it to update the matching.
- Otherwise: $X = A \cap Z$.
 $Y = B \cap Z$
Cover: $C = Y \cup (A \setminus X)$

Figure here

$X_0 = \{b, e\}$

Alternating Paths (Start from $b \setminus e$) $\{b, e, 4, 5, a, d, 1, 2, 3\}$

Figure 2 here

$X_0 = \{e\}$

$Z = \{e, 4, 5, b, d\}$

In order to have an augmenting path, Z must include an unsaturated vertex not in X_0 . Since Z does not contain 3 there is no augmenting path

$X = A \cap Z = \{b, d, e\}$

$Y = B \cap Z = \{4, 5\}$

$C = Y \cup (A \setminus X) = \{4, 5\} \cup \{a, c\} = \{4, 5, a, c\}$ is a min cover.

Lemma: Let M be a matching of a bipartite graph with partition A, B . Then

- a) There is no edge from X to $B \setminus Y$.
- b) $C = Y \cup (A \setminus X)$ is a cover.
- c) There is no edge of M from Y to $(A \setminus X)$.
- d) If u is the set of unsaturated vertices in Y then $|M| = |C| - |U|$.
- e) there is an augmenting path to each vertex in U .

Proof: a) Suppose there is an edge where $u \in X, v \in B \setminus Y$. There is an alternating path P from X_0 to u $P' = P + uv$. uv is not an edge of M . Alternating path to v . So v should be in Z .

b) Edges from:

$X \rightarrow Y$ - one end in Y and one end in C

$X \rightarrow B \setminus Y$ - don't exist by part a of the proof

$A \setminus X \rightarrow Y$ - both end in C

$A \setminus X \rightarrow B \setminus Y$ - one end in C

Hence C is cover

c) Suppose the edge $vw \in M$, $v \in A \setminus X$, $w \in Y$, $w \in Z$, so there is an alternating path from X_0 to w $w \in B$, so P has odd length (So to travel to the other set it should have an odd length length 1, 3, 5, ...). Starting from X_0 , the first edge is not in M , so the last edge (incident edge to w) is also not in the matching. $P' = P + vw$ is an alternating path and starts from X_0 to v . Thus v is in Z , hence it is in X . And since we said that $v \in A \setminus X$, it is a contradiction ($\longrightarrow! \longleftarrow$)

d) $X \rightarrow Y *$

$X \rightarrow B \setminus Y$ - does not \exists

$A \setminus X \rightarrow Y$ - not in M by part c

$A \setminus X \rightarrow B \setminus Y *$ - (2)

Number of edges X to Y in M . $|U|$ unsaturated vertices in Y . $|Y| - |U|$ saturated vertices in Y .

One edge in M for each in M for each of these. Number of edges from X to Y in $M = |Y| -$

$|U|$. From (2) now, 0 unsaturated vertices in $A \setminus X$. $|A \setminus X|$ saturated vertices in $A \setminus X$. $X_0 \subseteq Z$. So $X_0 \subseteq X$. Number is equal to $|A \setminus X|$. So $|M| = |Y| - |U| + |A \setminus X|$. So by using part b of the proof, $|M| = |C| - |U|$.

e) Let $v \in U$. V is unsaturated and $v \in Y$ and since $Y \in B \cap Z$ then $v \in Z$. Then there is an alternating path from X_0 to v then it is an augmenting path.

Hence proved the lemma.

7.2 König's Theorem

König's Theorem: In a bipartite graph, the maximum size of a matching is the minimum size of a cover.

Proof: Let M be a maximum matching. Then there is no augmenting path in G . Furthermore, $|U| = 0$. So by part b from the previous lemma, $C = Y \cup (A \setminus X)$ and then it is a cover. By part d) we get that $M = C$. \square .

D is a set of vertices.

Neighbour set, denoted by $N(D)$, is the set of vertices adjacent to a vertex in D

Halls' Theorem: G is a bipartite graph with partition on A, B . G has a matching saturating every vertex in A if and only if every subset D of A satisfies

$$|N(D)| \geq |D|$$

Picture: $D = \{1, 2, 3\}$

$N(D) = \{a, c\}$

$|N(D)| = 2 < 3 = |D|$

7.3 Applications of König's Theorem

7.4 System of Distinct Representatives