

ECLIPSE iGEM 2026

Thermodynamic Analysis — Complete Documentation

All thermodynamic scripts, equations, outputs, and interpretation for the lucCage-PfLB1 LOCKR biosensor (v1.0 and v2.2)

Denmark High School · iGEM 2026 · Computational Lead: Vishnu T.

Date: June 2026 · Constants: $RT = 0.592 \text{ kcal/mol}$ (37°C), $K_{\text{open}} = 1\text{e-}3$, $K_{\text{CK}} = 1\text{e-}8 \text{ M}$, $[\text{lucKey}] = 500 \text{ nM}$

Contents

1. System Overview and Thermodynamic Framework
2. Constants and Parameters
3. The K_{CK} Problem — Charge Optimization Thermodynamics
4. ADCP Docking Affinities and K_d Conversion
 - Script 1: Original binder docking (-26.3 kcal/mol)
 - Script 2: v1.0 optimized binder docking (-12.0 kcal/mol)
 - Script 3: v2.2 tandem binder docking (-16.6 kcal/mol)
 - Script 4: DDG and K_d improvement calculation
5. Avidity K_d Estimate (Pre-Docking Bounds)
 - Script 5: Simple and cooperative K_d estimates
6. Free Energy of Latch Opening
 - Script 6: Full latch opening and cascade analysis
7. LOCKR Fold-Change Model
 - Script 7: Thermodynamic model — all variants and pull strengths
 - Script 8: Updated model with ADCP K_d values
8. lucKey Competition Analysis — Why K_{open} Changes Don't Help
 - Script 9: v1.1 latch mutation thermodynamics
9. Complete Thermodynamic Results Summary
10. Wet Lab: Measuring Pull Strength from Luminescence Data

1. System Overview and Thermodynamic Framework

The ECLIPSE biosensor operates through a three-state thermodynamic cascade. Each transition has an associated free energy that determines whether the sensor will produce signal.

```

State 1: Cage closed (latch folded, SmBiT sequestered)
|
| Step 1: PflDH binds binder in latch
| dG_binding = -12.0 kcal/mol (v1.0) / -16.6 kcal/mol (v2.2)
v
State 2: Latch open (PflDH-binder complex, SmBiT exposed)
|
| Step 2: lucKey binds exposed SmBiT
| dG_lucKey = -2.32 kcal/mol
v
State 3: lucKey bound (NanoLuciferase reconstituted -> LIGHT)

```

The total cascade free energy is the sum of all steps. All must be negative (thermodynamically favorable) for the sensor to produce signal. The latch opening has a cost (+4.09 kcal/mol) because the latch is designed to be closed by default — the binder-PflDH binding energy must overcome this cost.

Net equation for latch opening:

$$dG_{\text{net}} = dG_{\text{latch_cost}} + dG_{\text{binding}}$$

$$dG_{\text{latch_cost}} = -RT * \ln(K_{\text{open}}) = -0.592 * \ln(1e-3) = +4.09 \text{ kcal/mol}$$

$$\text{v1.0: } dG_{\text{net}} = +4.09 + (-12.0) = -7.91 \text{ kcal/mol} \quad \text{--> FAVORABLE}$$

$$\text{v2.2: } dG_{\text{net}} = +4.09 + (-16.6) = -12.51 \text{ kcal/mol} \quad \text{--> FAVORABLE}$$

Full cascade:

$$dG_{\text{lucKey}} = -RT * \ln([lucKey]/K_{\text{CK}}) = -0.592 * \ln(50) = -2.32 \text{ kcal/mol}$$

$$\text{v1.0 total: } -7.91 + (-2.32) = -10.23 \text{ kcal/mol}$$

$$\text{v2.2 total: } -12.51 + (-2.32) = -14.83 \text{ kcal/mol}$$

2. Constants and Parameters

Constant	Symbol	Value	Derivation / Source
Gas constant x temperature	RT	0.592 kcal/mol	$R=8.314\text{e-}3 \text{ kcal/mol/K} \times T=310.15 \text{ K (37 C)}$
Cage opening equilibrium	K_open	1.0e-3	Quijano-Rubio et al. 2021; measured in vitro
$\Delta G(\text{latch opening cost})$	dG_open	+4.09 kcal/mol	$-RT * \ln(K_{\text{open}}) = -0.592 * \ln(1\text{e-}3)$
lucKey-SmBiT binding Kd	K_CK	1.0e-8 M (10 nM)	Quijano-Rubio et al. 2021
lucKey assay concentration	[lucKey]	500 nM = 5e-7 M	Assay design (50x K_CK)
lucKey/K_CK ratio	—	50	$[\text{lucKey}]/K_{\text{CK}} = 500\text{nM}/10\text{nM}$; dominates FC denominator
$\Delta G(\text{lucKey binding})$	dG_lucKey	-2.32 kcal/mol	$-RT * \ln([\text{lucKey}]/K_{\text{CK}}) = -0.592 * \ln(50)$
v1.0 binder-PfLDH Kd	K_LT	~100 pM	ADCP docking -12.0 kcal/mol -> Boltzmann
v2.2 tandem Kd	K_LT_tandem	~42 fM	ADCP docking -16.6 kcal/mol -> Boltzmann
$\Delta\Delta G (\text{v2.2 vs v1.0})$	DDG	-4.6 kcal/mol	$-16.6 - (-12.0) = -4.6 \text{ kcal/mol}$
Kd improvement factor	—	2369x	$\exp(-DDG/RT) = \exp(4.6/0.592)$

Table 1. All thermodynamic constants used in ECLIPSE analysis.

Boltzmann-Kd relationship (used throughout):

$$dG = -RT * \ln(1/Kd) = RT * \ln(Kd)$$

$$\text{Rearranged: } Kd = \exp(dG / RT)$$

Example (v1.0): $Kd = \exp(-12.0 / 0.592) = \exp(-20.27) = 1.57\text{e-}9 \text{ M}$
(anchored to 100 pM from MD work; ADCP gives relative differences)

$$\begin{aligned} \text{Example (v2.2): } Kd_{\text{v22}} &= Kd_{\text{v10}} * \exp(DDG / RT) \\ &= 100\text{e-}12 * \exp(-4.6 / 0.592) \\ &= 100\text{e-}12 * \exp(-7.77) \\ &= 100\text{e-}12 * 4.22\text{e-}4 \\ &= 4.22\text{e-}14 \text{ M} = 42.2 \text{ fM} \end{aligned}$$

3. The K_{CK} Problem — Charge Optimization Thermodynamics

The original RFDiffusion binder LISDAELEAIFAEELDC has 6 charged D/E residues (positions 4, 6, 8, 13, 14, 16) that sit at the lucKey-cage interface. This creates a thermodynamic penalty that collapses K_{CK} and eliminates all signal regardless of how well PFLDH binds.

Calculation:

Each charged residue at the lucKey interface imposes ~0.8 kcal/mol electrostatic penalty

Penalty total = 6 residues × 0.8 kcal/mol = 4.8 kcal/mol

Original K_{CK}: 1e-8 M

dG_{CK} (original) = -RT * ln(1/K_{CK}) = -0.592 * ln(1e8) = -10.9 kcal/mol

After grafting original binder:

dG_{CK} (grafted) = -10.9 + 4.8 = -6.1 kcal/mol

K_{CK} (grafted) = exp(-6.1 / 0.592) = exp(-10.31) = 3.3e-5 M

-> K_{CK} drops from 1e-8 to 3.3e-5 (~3300x weaker)

Effect on signal -- lucKey occupancy of the open SmBiT site:

occupancy = [lucKey] / ([lucKey] + K_{CK})

Healthy (optimized binder, K_{CK}=1e-8):

occupancy = 500e-9 / (500e-9 + 1e-8) = 500/510 = 98.0%

(lucKey nearly saturates SmBiT once latch opens)

Degraded (original binder, K_{CK}=3.3e-5):

occupancy = 500e-9 / (500e-9 + 3.3e-5) = 1.5%

(lucKey almost never bound, even if latch is wide open)

Reduction factor: 98.0% / 1.5% ~ 66x less usable signal

CORE PROBLEM: even with the latch fully open, ~98.5% of cage molecules would have no lucKey bound. The sensor is structurally intact but the light-producing step is crippled.

NOTE: this occupancy calculation is independent of the separate fold-change model in Section 7 (theta/pull), which describes how PFLDH concentration modulates LATCH opening, not lucKey occupancy. The two are sequential steps in the cascade and are treated separately here for clarity -- see Section 1 for how they combine into the full cascade dG.

Solution — D/E to Alanine mutations:

Preserve PFLDH interface residues: positions 1, 2, 11, 12, 15

Mutate remaining D/E to Alanine: positions 4, 6, 8, 13, 14, 16

Mutations: D4A, E6A, E8A, E13A, E14A, D16A

Original: LISDAELEAIFAEELDC (6 D/E at lucKey interface)

Optimized: LISAAALAAIFAAALAC (0 D/E at lucKey interface)

Penalty removed: 6 × 0.8 = 4.8 kcal/mol -> 0

K_{CK} restored: ~1e-8 M

Signal equation comparison:

	Original binder	Optimized binder
ADCP affinity (PfLDH)	-26.3 kcal/mol	-12.0 kcal/mol
K _d (PfLDH)	~0.1 fM (est.)	~100 pM
K _{CK} (lucKey binding)	~3.3e-5 M	1e-8 M
lucKey occupancy at 500nM	~1.5%	~98.0%
Absolute signal	Severely reduced (~66x)	Functional
Sensor works?	No	Yes

Table 2. Original vs optimized binder thermodynamic comparison. Despite stronger PfLDH affinity, the original binder produces zero signal.

4. ADCP Docking Affinities and Kd Conversion

ADCP (AutoDock CrankPep) was used to dock each binder sequence against the PflDH receptor. 20 independent runs x 1,000,000 steps each. Cluster size is as important as affinity — large clusters indicate reproducible binding at a genuine energy minimum.

Script 1 — Original charged binder docking

```
# Receptor prep (30A box)
agfr -r pflhd.pdbqt -o receptor/pflhd \
      -b user 40.1 0.0 45.35 30 30 30

# Dock original binder
adcp -t pflhd.trg \
      -s LISDAELEAIFAEELDC \
      -N 20 -n 1000000 \
      -o original_charged_binder
```

Output:

```
Detected 24 cores, using 24 cores
Docking performed in 47.15 seconds
```

mode	affinity (kcal/mol)	clust. rmsd	clust. size	energy avg.
1	-26.3	0.0	23	-24.2
2	-23.6	2.1	31	-18.5

The original binder docks at -26.3 kcal/mol — the highest raw affinity of all three binders. The D/E residues were actively forming electrostatic contacts with PflDH. Cluster size of 23 is moderate, suggesting multiple non-specific binding modes due to promiscuous electrostatic attraction.

Script 2 — v1.0 optimized binder docking

```
adcp -t pflhd.trg \
      -s LISAAALAAIFAAALAC \
      -N 20 -n 1000000 \
      -o v1_single
```

Output:

```
Docking performed in 23.07 seconds
```

mode	affinity (kcal/mol)	clust. rmsd	clust. size	energy avg.
1	-12.0	0.0	45	-10.0
2	-7.3	2.6	52	-4.3

Affinity -12.0 kcal/mol, cluster 45. Lower affinity than original but larger, tighter cluster — binding is more specific and reproducible. The -12.0 kcal/mol corresponds to Kd ~100 pM (anchored from MD simulations).

Script 3 — v2.2 tandem binder docking

```
# Larger 50A box needed for 35 aa tandem peptide
agfr -r pflhd.pdbqt -o receptor/pflhd_large \
      -b user 40.1 0.0 45.35 50 50 50

adcp -t pflhd_large.trg \
      -s LISAAALAAIFAAALACGLISAAALAAIFAAALAC \
```

```
-N 20 -n 1000000 \
-o v2_tandem
```

Output:

Docking performed in 62.12 seconds

mode	affinity (kcal/mol)	clust. rmsd	clust. size	energy avg.
1	-16.6	0.0	70	-14.0
2	-10.4	2.8	12	-9.7
3	-9.1	3.0	53	-6.8
4	-7.4	21.0	39	-6.1

The tandem binder reaches -16.6 kcal/mol with 70 poses in the top cluster — the most reproducible binding geometry of all three. The avidity effect (two binder copies) recovers binding energy lost during charge optimization and then significantly exceeds it.

Script 4 — DDG and Kd improvement calculation

```
import math

RT = 0.592 # kcal/mol at 37 C

dG_original = -26.3 # kcal/mol
dG_v10      = -12.0
dG_v22      = -16.6

# Kd from Boltzmann (v1.0 anchored to 100 pM from MD)
Kd_v10 = 100e-12
Kd_v22 = Kd_v10 * math.exp((dG_v22 - dG_v10) / RT)

# Kd improvement
ddG_opt_vs_orig = dG_v10 - dG_original # cost of optimization
ddG_v22_vs_v10  = dG_v22 - dG_v10     # gain from avidity

print(f"=== BINDING AFFINITY PROGRESSION ===")
print(f"Original:      {dG_original:.1f} kcal/mol")
print(f"v1.0:            {dG_v10:.1f} kcal/mol")
print(f"v2.2:            {dG_v22:.1f} kcal/mol")
print()
print(f"Optimization cost (orig -> v1.0): DDG = {ddG_opt_vs_orig:.1f} kcal/mol")
print(f"Avidity gain (v1.0 -> v2.2):      DDG = {ddG_v22_vs_v10:.1f} kcal/mol")
print()
print(f"Kd v1.0 = {Kd_v10*1e12:.1f} pM")
print(f"Kd v2.2 = {Kd_v22*1e15:.2f} fM")
print(f"Kd improvement (v2.2 vs v1.0) = {Kd_v10/Kd_v22:.0f}x")
print()
print(f"Tradeoff analysis:")
print(f"  Charge optimization sacrificed: +14.3 kcal/mol PfLDH affinity")
print(f"  But restored K CK (essential for any signal)")
print(f"  Avidity recovered: -4.6 kcal/mol")
print(f"  Net vs original: {dG_v22 - dG_original:.1f} kcal/mol (still weaker)")
```

Output:

```
=== BINDING AFFINITY PROGRESSION ===
Original:  -26.3 kcal/mol
v1.0:      -12.0 kcal/mol
v2.2:      -16.6 kcal/mol

Optimization cost (orig -> v1.0): DDG = +14.3 kcal/mol
Avidity gain (v1.0 -> v2.2):      DDG = -4.6 kcal/mol
```

Kd v1.0 = 100.0 pM
 Kd v2.2 = 42.21 fM
 Kd improvement (v2.2 vs v1.0) = 2369x

Tradeoff analysis:

Charge optimization sacrificed: +14.3 kcal/mol PfLDH affinity

But restored K_{CK} (essential for any signal)

Avidity recovered: -4.6 kcal/mol

Net vs original: -9.7 kcal/mol (still weaker in raw PfLDH affinity)

Binder	Affinity	DDG vs v1.0	Kd	Cluster	K _{CK}	Signal
Original	- 26.3 kcal/mol	+14.3	~0.1 fM*	23	3.3e-5	No
v1.0 optimized	- 12.0 kcal/mol	baseline	~100 pM	45	1e-8	Yes
v2.2 tandem	- 16.6 kcal/mol	-4.6	~42 fM	70	1e-8	Yes

Table 3. Complete docking comparison. *Original Kd estimated; non-functional due to K_{CK} collapse.

5. Avidity Kd Estimate (Pre-Docking Bounds)

Before running ADCP for v2.2, theoretical bounds were calculated to verify the docking result would be physically reasonable.

Script 5 — Simple and cooperative Kd estimates

```
import math

Kd_single = 100e-12 # 100 pM from v1.0

# Lower bound: simple additive (two independent binders)
Kd_simple = Kd_single / 2

# Upper bound: cooperative binding (both engage simultaneously)
# For perfectly cooperative two-site binding: Kd_app = Kd_single^(3/2)
Kd_cooperative = Kd_single ** 1.5

print(f"Single binder Kd:           {Kd_single*1e12:.0f} pM")
print(f"Simple model (2x):          ~{Kd_simple*1e12:.0f} pM")
print(f"Cooperative model (^1.5):    ~{Kd_cooperative*1e15:.2f} fM")
print()
print(f"Expected ADCP result should fall between 50 pM and 1 fM")
```

Output:

```
Single binder Kd:           100 pM
Simple model (2x):          ~50 pM
Cooperative model (^1.5):    ~1.00 fM
```

Expected ADCP result should fall between 50 pM and 1 fM

ADCP returned 42 fM — well within the predicted bounds and in the cooperative regime, confirming the tandem binder achieves genuine avidity rather than just simple additive binding.

6. Free Energy of Latch Opening

The core thermodynamic question: does PflDH binding provide enough energy to open the latch against its natural closed-state stability?

Script 6 — Full latch opening and cascade analysis

```
import math

RT      = 0.592    # kcal/mol at 37 C
K_open  = 1e-3      # latch opening equilibrium
K_CK    = 1e-8      # lucKey-SmBiT Kd
lucKey   = 500e-9   # assay lucKey concentration

# Cost to open the latch (from K_open)
dG_open_cost = -RT * math.log(K_open)
print(f"1. LATCH OPENING COST")
print(f"    K_open = {K_open}")
print(f"    dG = -RT*ln(K_open) = -{RT}*ln({K_open}) = +{dG_open_cost:.2f} kcal/mol")
print()

# Binder-PflDH binding energies from ADCP
dG_v10 = -12.0
dG_v22 = -16.6
print(f"2. BINDER-PFLDH BINDING (from ADCP docking)")
print(f"    v1.0: {dG_v10:.1f} kcal/mol")
print(f"    v2.2: {dG_v22:.1f} kcal/mol")
print()

# Net free energy of latch opening
print(f"3. NET FREE ENERGY OF LATCH OPENING")
for label, dG_b in [("v1.0", dG_v10), ("v2.2", dG_v22)]:
    dG_net = dG_open_cost + dG_b
    K_eq    = math.exp(-dG_net / RT)
    pct     = K_eq / (1 + K_eq) * 100
    verdict = 'FAVORABLE' if dG_net < 0 else 'UNFAVORABLE'
    print(f"    {label}: {dG_open_cost:.2f} + ({dG_b:.1f}) = {dG_net:.2f} kcal/mol")
    print(f"        K_eq = exp(-({dG_net:.2f})/0.592) = {K_eq:,.0f}")
    print(f"        Fraction open = {pct:.1f}% --> {verdict}")
print()

# lucKey binding step
dG_lucKey = -RT * math.log(lucKey / K_CK)
print(f"4. lucKey CASCADE STEP")
print(f"    [lucKey]/K_CK = {lucKey:.0e}/{K_CK:.0e} = {lucKey/K_CK:.0f}")
print(f"    dG = -RT*ln([lucKey]/K_CK) = -{RT}*ln({lucKey/K_CK:.0f}) = {dG_lucKey:.2f} kcal/mol")
print()

# Full cascade
print(f"5. FULL CASCADE dG (latch open + lucKey bind)")
for label, dG_b in [("v1.0", dG_v10), ("v2.2", dG_v22)]:
    dG_net = dG_open_cost + dG_b
    cascade = dG_net + dG_lucKey
    print(f"    {label}: {dG_net:.2f} + ({dG_lucKey:.2f}) = {cascade:.2f} kcal/mol")
```

Output:

1. LATCH OPENING COST
 $K_{\text{open}} = 0.001$
 $dG = -RT \ln(K_{\text{open}}) = -0.592 \ln(0.001) = +4.09 \text{ kcal/mol}$
2. BINDER-PflDH BINDING (from ADCP docking)

v1.0: -12.0 kcal/mol
v2.2: -16.6 kcal/mol

3. NET FREE ENERGY OF LATCH OPENING

v1.0: $+4.09 + (-12.0) = -7.91$ kcal/mol
 $K_{eq} = \exp(-(-7.91)/0.592) = 635,721$
 Fraction open = 100.0% --> FAVORABLE

v2.2: $+4.09 + (-16.6) = -12.51$ kcal/mol
 $K_{eq} = \exp(-(-12.51)/0.592) = 1,506,093,819$
 Fraction open = 100.0% --> FAVORABLE

4. lucKey CASCADE STEP

$[lucKey]/K_{CK} = 5e-7/1e-8 = 50$
 $dG = -RT \cdot \ln([lucKey]/K_{CK}) = -0.592 \cdot \ln(50) = -2.32$ kcal/mol

5. FULL CASCADE dG (latch open + lucKey bind)

v1.0: $-7.91 + (-2.32) = -10.23$ kcal/mol
 v2.2: $-12.51 + (-2.32) = -14.83$ kcal/mol

Quantity	v1.0	v2.2	Meaning
dG(latch cost)	+4.09 kcal/mol	+4.09 kcal/mol	Cost to open latch (same for both)
dG(binder-PfLDH)	-12.0 kcal/mol	-16.6 kcal/mol	Energy released by PfLDH binding
Net dG(latch open)	-7.91 kcal/mol	-12.51 kcal/mol	Net: latch opening is favorable
K_{eq} (latch)	635,721	1,506,093,819	Open state dominates overwhelmingly
Latch open %	100.0%	100.0%	Latch fully open when PfLDH bound
dG(lucKey)	-2.32 kcal/mol	-2.32 kcal/mol	lucKey binding always favorable
Total cascade dG	-10.23 kcal/mol	-14.83 kcal/mol	End-to-end signal chain: favorable

Table 4. Free energy analysis of the full signal cascade.

7. LOCKR Fold-Change Model

The LOCKR system is modeled as a three-state equilibrium. Fold-change is the ratio of signal-state to baseline-state luciferase reconstitution.

Core equations:

$$\theta = [\text{PflDH}] / ([\text{PflDH}] + K_d) \quad (\text{fraction binder occupied})$$

$$K_{\text{open_eff}} = K_{\text{open}} * (1 + \text{pull} * \theta) \quad (\text{effective latch opening w/ PflDH})$$

$$f_{\text{base}} = K_{\text{open}} / (1 + K_{\text{open}} + [\text{lucKey}]/K_{\text{CK}}) \quad (\text{baseline fraction open})$$

$$f_{\text{signal}} = K_{\text{open_eff}} / (1 + K_{\text{open_eff}} + [\text{lucKey}]/K_{\text{CK}}) \quad (\text{signal fraction open})$$

$$FC = f_{\text{signal}} / f_{\text{base}}$$

Substituting numbers:

$$[\text{lucKey}]/K_{\text{CK}} = 500\text{nM}/10\text{nM} = 50$$

$$f_{\text{base}} = 1\text{e-}3 / (1 + 0.001 + 50) = 1\text{e-}3/51.001 = 1.96\text{e-}5$$

At pull=10, theta=1 (saturating PflDH):

$$K_{\text{open_eff}} = 1\text{e-}3 * (1 + 10*1) = 1.1\text{e-}2$$

$$f_{\text{signal}} = 1.1\text{e-}2 / (1 + 0.011 + 50) = 1.1\text{e-}2/51.011 = 2.16\text{e-}4$$

$$FC = 2.16\text{e-}4 / 1.96\text{e-}5 = 11.0\text{x}$$

Script 7 — Thermodynamic model: all variants and pull strengths

```
import numpy as np

K_open = 1e-3
K_CK = 1e-8
lucKey = 500e-9

def fold_change(pfldh, Kd, pull):
    theta = pfldh / (pfldh + Kd)
    Koe = K_open * (1 + pull * theta)
    f_base = K_open / (1 + K_open + lucKey/K_CK)
    f_signal = Koe / (1 + Koe + lucKey/K_CK)
    return f_signal / f_base

def max_fc_ec50(Kd, pull):
    pfldh = np.logspace(-14, -5, 500)
    fcs = [fold_change(c, Kd, pull) for c in pfldh]
    mfc = max(fcs)
    half = (mfc + 1) / 2
    ec50 = pfldh[np.argmin(np.abs(np.array(fcs) - half))]
    return mfc, ec50

Kd_v10 = 100e-12 # 100 pM
Kd_v22 = 42.21e-15 # 42.21 fM

print(f'Variant:<25 {\'pull\':>5} {\'Max FC\':>8} {\'EC50\':>12} {\'LOD\':>12}')
print("-" * 65)
for label, Kd in [("v1.0 (Kd=100pM)", Kd_v10), ("v2.2 (Kd=42fM)", Kd_v22)]:
    for pull in [10, 20, 30]:
        mfc, ec50 = max_fc_ec50(Kd, pull)
        lod = ec50 * 0.1
```

```
print(f"{label:<25} {pull:>5}    {mfc:>6.0f}x    "
      f"{ec50*1e9:>10.4f} nM    ~{lod*1e12:>8.2f} pM")
```

Output:

Variant	pull	Max FC	EC50	LOD
v1.0 (Kd=100pM)	10	11x	0.1008 nM	~10.08 pM
v1.0 (Kd=100pM)	20	21x	0.1008 nM	~10.08 pM
v1.0 (Kd=100pM)	30	31x	0.1008 nM	~10.08 pM
v2.2 (Kd=42fM)	10	11x	~0.0000 nM	~0.00 pM
v2.2 (Kd=42fM)	20	21x	~0.0000 nM	~0.00 pM
v2.2 (Kd=42fM)	30	31x	~0.0000 nM	~0.00 pM

Script 8 — Full model output with interpretation**Output (expanded):**

v1.0 single (pull=10, conservative)

Max fold-change: 11x | EC50: 0.1008 nM | LOD: ~10.08 pM

v1.0 single (pull=20, optimistic)

Max fold-change: 21x | EC50: 0.1008 nM | LOD: ~10.08 pM

v2.2 tandem (pull=10)

Max fold-change: 11x | EC50: ~0 nM | LOD: ~0 pM

v2.2 tandem (pull=20)

Max fold-change: 21x | EC50: ~0 nM | LOD: ~0 pM

v2.2 tandem (pull=30, best case)

Max fold-change: 31x | EC50: ~0 nM | LOD: ~0 pM

Note: v2.2 EC50 ~0 because Kd (42 fM) << any clinical PfLDH concentration.

Sensor is always fully saturated. Performance ceiling = pull strength.

The fold-change ceiling (11/21/31x) is identical for v1.0 and v2.2 because it is set by cage mechanics (K_{open} and K_{CK}), not binder affinity. What changes between variants is the EC50 — v2.2 is saturated at essentially zero PfLDH concentration, meaning it hits maximum fold-change even at trace infection levels where v1.0 would still be on the rising phase of its sigmoidal response curve.

8. lucKey Competition Analysis — Why Latch Mutations Don't Help

When we attempted to create a v1.1 variant with latch mutations I346S + I356S (analogous to the Baker Lab's V223S/I238S), the thermodynamic model predicted no meaningful improvement. This section shows the math.

The denominator problem:

$$f_{\text{base}} = K_{\text{open}} / (1 + K_{\text{open}} + [\text{lucKey}]/K_{\text{CK}})$$

Numerically:

$$K_{\text{open}} = 1\text{e-}3 = 0.001$$

$$[\text{lucKey}]/K_{\text{CK}} = 500\text{nM}/10\text{nM} = 50$$

$$\begin{aligned} f_{\text{base}} &= 0.001 / (1 + 0.001 + 50) \\ &= 0.001 / 51.001 \end{aligned}$$

The 50 term DOMINATES. $K_{\text{open}} = 0.001$ is negligible.

Even with v1.1 moderate estimate ($K_{\text{open}} = 2.93\text{e-}2$, 30x increase):

$$\begin{aligned} f_{\text{base}} &= 0.0293 / (1 + 0.0293 + 50) \\ &= 0.0293 / 51.029 \\ &= 5.74\text{e-}4 \quad (\text{only } 29\text{x larger } f_{\text{base}}) \end{aligned}$$

For FC calculation, f_{base} and f_{signal} both scale similarly
-> ratio (FC) barely changes

Conclusion: $[\text{lucKey}]/K_{\text{CK}} = 50 \gg K_{\text{open}}$ in all scenarios.

The fold-change ceiling is set by lucKey competition, not latch stability.

Script 9 — v1.1 latch mutation thermodynamics

```
import math, numpy as np

RT      = 0.592
K_CK    = 1e-8
lucKey   = 500e-9
Kd       = 100e-12 # same binder, same Kd

dG_latch_v10 = -RT * math.log(1e-3) # +4.09 kcal/mol

# I->S destabilization: 0.5-1.5 kcal/mol per mutation
scenarios = [
    ("Conservative (0.5/mut = 1.0 total)", 1.0),
    ("Moderate (1.0/mut = 2.0 total)", 2.0),
    ("Optimistic (1.5/mut = 3.0 total)", 3.0),
]

def max_fc_ec50(K_o, pull, Kd):
    pflhdh = np.logspace(-14, -5, 500)
    fcs = []
    for c in pflhdh:
        theta = c/(c+Kd)
        Koe = K_o*(1+pull*theta)
        fb = K_o/(1+K_o+lucKey/K_CK)
        fs = Koe/(1+Koe+lucKey/K_CK)
        fcs.append(fs/fb)
    mfc = max(fcs)
```

```

ec50 = pflhd[np.argmin(np.abs(np.array(fcs)-(mfc+1)/2))]
return mfc, ec50

print(f"v1.0 baseline: K_open=1e-3 dG(open)=+{dG_latch_v10:.2f} kcal/mol")
for label, destab in scenarios:
    dG11 = dG_latch_v10 - destab
    Ko11 = math.exp(-dG11/RT)
    print(f"\n{label}")
    print(f" dG(latch) = +{dG11:.2f} K_open = {Ko11:.3e}")
    for pull in [10,20,30]:
        mfc, ec50 = max_fc_ec50(Ko11, pull, Kd)
        print(f" pull={pull}: FC={mfc:.0f}x EC50={ec50*1e9:.4f} nM")

print(f"\n--- v1.0 baseline ---")
for pull in [10,20,30]:
    mfc, ec50 = max_fc_ec50(1e-3, pull, Kd)
    print(f" pull={pull}: FC={mfc:.0f}x EC50={ec50*1e9:.4f} nM")

```

Output:

v1.0 baseline: K_open=1e-3 dG(open)=+4.09 kcal/mol

Conservative (0.5/mut = 1.0 total)
dG(latch) = +3.09 K_open = 5.415e-03
pull=10: FC=11x EC50=0.1009 nM
pull=20: FC=21x EC50=0.1009 nM
pull=30: FC=31x EC50=0.1009 nM

Moderate (1.0/mut = 2.0 total)
dG(latch) = +2.09 K_open = 2.932e-02
pull=10: FC=11x EC50=0.1009 nM
pull=20: FC=21x EC50=0.1009 nM
pull=30: FC=30x EC50=0.0968 nM

Optimistic (1.5/mut = 3.0 total)
dG(latch) = +1.09 K_open = 1.588e-01
pull=10: FC=11x EC50=0.0968 nM
pull=20: FC=20x EC50=0.0929 nM
pull=30: FC=28x EC50=0.0929 nM

--- v1.0 baseline ---
pull=10: FC=11x EC50=0.1009 nM
pull=20: FC=21x EC50=0.1009 nM
pull=30: FC=31x EC50=0.1009 nM

All three v1.1 scenarios produce fold-change and EC50 values essentially identical to v1.0. In the optimistic scenario, fold-change at pull=30 actually decreases from 31x to 28x because the raised K_{open} compresses the ratio by lifting baseline signal. Decision: do not synthesize v1.1. Test v1.0 and v2.2 instead.

When would latch mutations help?

[lucKey]	[lucKey]/K _{CK}	K _{open} matters?	Expected v1.1 benefit
500 nM	50	No (50 >> 0.001)	< 1% — negligible
100 nM	10	Marginally	5-10% at best
50 nM	5	Yes (5 ~ K _{open} range)	Moderate, observable
10 nM	1	Yes (1 ~ K _{open})	Significant

Table 5. lucKey concentration determines whether K_{open} changes affect FC. At 500 nM, the effect is negligible. Titrating [lucKey] in the wet lab assay is the correct test — no extra gene synthesis required.

9. Complete Thermodynamic Results Summary

Quantity	Original binder	v1.0	v2.2
ADCP affinity	-26.3 kcal/mol	-12.0 kcal/mol	-16.6 kcal/mol
DDG vs v1.0	+14.3 kcal/mol	baseline	-4.6 kcal/mol
K _d (PfLDH)	~0.1 fM (non-func.)	~100 pM	~42.21 fM
K _d improvement	—	baseline (1x)	2369x
K _{CK} (lucKey)	~3.3e-5 (FATAL)	1e-8	1e-8
dG(latch cost)	—	+4.09 kcal/mol	+4.09 kcal/mol
dG(binding)	—	-12.0 kcal/mol	-16.6 kcal/mol
dG(net, latch)	—	-7.91 kcal/mol	-12.51 kcal/mol
K _{eq} (latch)	—	635,721	1,506,093,819
Latch opens?	—	100%	100%
dG(lucKey)	—	-2.32 kcal/mol	-2.32 kcal/mol
Full cascade dG	—	-10.23 kcal/mol	-14.83 kcal/mol
Max FC (pull=10)	—	11x	11x
Max FC (pull=20)	—	21x	21x
Max FC (pull=30)	—	31x	31x
EC ₅₀ (pull=10)	—	~0.10 nM	~0 nM
LOD (pull=10)	—	~10 pM	~0 pM
Signal produced?	No	Yes	Yes

Table 6. Complete thermodynamic results for all three binder variants. Original binder is included for comparison but is non-functional.

10. Wet Lab: Measuring Pull Strength from Luminescence Data

Pull strength is the one thermodynamic parameter that cannot be determined computationally — it measures how efficiently the mechanical force of binder-PfLDH binding transfers to latch opening (allosteric coupling). It is extracted by fitting the experimental luminescence curve.

Method: curve fitting to the fold-change model

```
from scipy.optimize import curve_fit
import numpy as np

# Fixed constants (from literature + ADCP)
K_open = 1e-3
K_CK = 1e-8
lucKey = 500e-9
Kd = 100e-12 # v1.0; use 42.21e-15 for v2.2

def fold_change_model(pfldh_conc, pull_strength):
    theta = pfldh_conc / (pfldh_conc + Kd)
    Koe = K_open * (1 + pull_strength * theta)
    f_base = K_open / (1 + K_open + lucKey/K_CK)
    f_signal = Koe / (1 + Koe + lucKey/K_CK)
    return f_signal / f_base

# pfldh_conc = array of PfLDH concentrations tested (in M)
# fc_measured = array of measured fold-changes at each concentration
pfldh_conc = np.array([1e-12, 10e-12, 100e-12, 1e-9, 10e-9, 100e-9, 1e-6])
fc_measured = np.array([...]) # fill in from experiment

popt, pcov = curve_fit(
    fold_change_model,
    pfldh_conc,
    fc_measured,
    p0=[10], # initial guess: pull=10
    bounds=(0, 100) # physically reasonable range
)

pull_strength = popt[0]
pull_std = np.sqrt(pcov[0,0])
print(f"Pull strength = {pull_strength:.1f} +/- {pull_std:.1f}")
print(f"Expected fold-change at pull={pull_strength:.0f}: {1+pull_strength:.0f}x")
```

Interpretation guide:

Pull strength	Expected max FC	Interpretation
< 5	< 6x	Weak allosteric coupling; latch is stiff
5-15	6-16x	Moderate coupling; consistent with literature
15-30	16-31x	Strong coupling; excellent design
> 30	> 31x	Very strong; near-ideal allosteric transfer

Table 7. Pull strength interpretation. Baker Lab LOCKR designs typically range 10-20x.

If fold-change is lower than expected:

Observed FC	Pull strength	Likely issue	Next experiment
< 5x	< 4	Latch not opening fully	Try [lucKey]=50nM; check expression by CD

Observed FC	Pull strength	Likely issue	Next experiment
5-10x	4-9	Moderate coupling	Try I346S+I356S (v1.0 only)
> 10x (v1.0)	9+	Design works	Proceed to clinical testing
> 50x (v2.2)	49+	Avidity fully expressed	Paper-worthy result

Table 8. Decision tree based on experimental fold-change results.

ECLIPSE iGEM 2026 · Denmark High School · Thermodynamics Documentation · June 2026 ·
Computational Lead: Vishnu T.