

DEPARTMENT OF RADIODIAGNOSIS

NAME: MRS. CHANDRAMATHI GAJANANA HEGDE	REFERRED BY: DR. PRITVIRAJ S K
AGE/SEX: 73 Y/F	REGN NO: P126798
DATE:: 12-Jul-2025	RDBNO: 250712110

CECT ABDOMEN AND PELVIS

A preliminary AP topogram of the Abdomen and pelvis was obtained. Volume scans were performed starting from the domes of diaphragm down to the level of pubic symphysis, employing 0.625 mm sections. 80 ml of Optiray 350 mg was used for contrast enhancement.

OBSERVATIONS:

➤ **MESENTERY AND RETROPERITONEUM:**

- There is well defined homogeneously enhancing soft tissue mass noted in the pre and para aortic region of mid abdomen, measuring approximately 112 x 75 x 102mm. No cystic changes / calcification.
 - Superiorly, the lesion is extending upto the origin of superior mesenteric artery and inferiorly upto the bifurcation of aorta.
 - Anteriorly, it is seen displacing the bowel loops.
 - Right laterally, it is seen encasing the inferior venacava and right renal artery and vein.
 - Left laterally, it is extending and abutting the left renal hilum and encasing the left renal artery and vein.
 - Supero-laterally, it is abutting the tail of pancreas
 - Posteriorly, it is encasing the abdominal aorta, inferior mesenteric artery and origin of superior mesenteric artery. No evidence of vertebral body erosion.
 - Multiple enlarged pre and para-aortic lymph nodes are noted, largest measuring 12 x 10mm.

➤ **LUNG BASES:**

- Similar characteristic homogeneously enhancing soft tissue lesion noted in the posterior mediastinum encasing the left hilum, left pulmonary veins, left main and segmental bronchi and also extending into the right lower lobe of lung, together measuring approximately 126 x 93mm.
- Left mild pleural effusion noted.
- Multiple small nodular lesions scattered in bilateral lung fields, largest measuring 20 x 23mm in posterior segment of right upper lobe.

DEPARTMENT OF RADIO DIAGNOSIS

NAME: MRS. CHANDRAMATHI GAJANANA HEGDE		REFERRED BY: DR. PRITVIRAJ S K
AGE/SEX: 73 Y/F		REGN NO: P126798
DATE: 12-Jul-2025		RDBNO: 250712110

CECT ABDOMEN AND PELVIS

➤ **LIVER:**

- Normal in size, shape, outlines and parenchymal attenuation.
- No focal lesion.
- The porta hepatis and portal vein are normal.
- The intrahepatic portal venous radicals are normal.
- No evidence of intrahepatic biliary radicular dilatation.

➤ **GALL BLADDER:**

- Normal in size, shape and outlines. Pericholecystic area is normal. The common bile duct is not dilated.

➤ **SPLEEN:**

- Spleen is normal in size, shape and attenuation values.
- The splenic hilum and splenic vein are normal.

➤ **PANCREAS:**

- Pancreas is normal in size, contour and attenuations values.
- No evidence of focal mass lesion/pancreatic duct dilatation.

➤ **GE JUNCTION:** is normal.

➤ **STOMACH:** shows normal cardia, body & antropyloric region.

➤ **DUODENUM:** is normal. No evidence of wall thickening.

➤ **KIDNEYS:**

- Kidneys are normal in size and shape.
- The renal outlines are normal and both kidneys are showing good concentration and prompt excretion of contrast.
- No evidence of focal mass lesion/hydronephrosis/calculi.
- Adrenal glands are normal in size, shape and attenuation values.
- Aorta and extra hepatic portion of inferior venacava are normal in calibre.

➤ **URINARY BLADDER:** is normal in size and attenuation.

➤ **NO EVIDENCE OF ASCITES.**

➤ **UTERUS AND OVARIES:** Atrophy.

DEPARTMENT OF RADIODIAGNOSIS

NAME: MRS. CHANDRAMATHI GAJANANA HEGDE	REFERRED BY: DR. PRITVIRAJ S K
AGE/SEX: 73 Y/F	REGN NO: P126798
DATE: 12-Jul-2025	RDBNO: 250712110

CECT ABDOMEN AND PELVIS

- **BOWEL LOOPS:** Small and large bowel loops are normal.
 - Ileo-caecal junction is normal.
- Degenerative changes noted in thoraco-lumbar vertebral bodies
- Numerous small lytic lesions are noted in the vertebral bodies - Metastasis.

IMPRESSION:

- ❖ Well defined homogeneously enhancing soft tissue mass in the pre and para aortic region (Retroperitoneum) of mid abdomen with extension and encasement as described. No cystic changes / calcification.
 - ❖ Multiple enlarged pre and para-aortic lymph nodes.
 - ❖ Similar characteristic homogeneously enhancing soft tissue lesion in the posterior mediastinum encasing the left hilum, left pulmonary veins, left main and segmental bronchi and also extending into the right lower lobe of lung.
 - ❖ Left mild pleural effusion.
 - ❖ Multiple small nodular lesions scattered in bilateral lung fields.
- DD's
1. Lymphoma.
 2. Metastasis.
- ❖ Numerous small lytic lesions in the vertebral bodies - Metastasis.

→ Suggested Histopathology correlation.


Dr. Kiran Kumar R.G.
MBBS DMRD DNB
Consultant Radiologist
KMC. Reg. No - 87099