



CENTER FOR ADVANCED NUCLEAR
MEDICINE THERANOSTICS



Name :	Mr. Manjunatha G M	PET No :	KMIO20256701 (202411951)
Age/Sex :	45 Years/ Male	UHID :	20240007016
Ref/Dept :	Medical Oncology	Date :	11.06.2025 (13.12.2024)

¹⁸F FDG PET-CT Report

CLINICAL INDICATION: Case of periampullary carcinoma status post stenting.

TECHNIQUE: Whole body images (vertex to mid thigh) were acquired in 3-D mode 60 min after i.v. injection of 4.3mCi of ¹⁸F-FDG using a uMI 550 80 slice Digital PET CT. SIPM with TOF detector to crystal coupling. FBS before trace injection was 104mg/dl. IV contrast was used. Reconstruction of the acquired data was performed so as to obtain fused PET-CT images in all views.

FINDINGS:

Abdomen and Pelvis:

- Post-surgical changes noted in periampullary region. Biliary stent noted in situ. No evidence of FDG avid lesion at surgical site.
- Interval resolution of FDG uptake in portocaval (previously 1.8 x 1.3 cm SUVmax 4.5), para-aortic and right external iliac lymph nodes.
- Liver appears enlarged (size 19.9 cm). No evidence of FDG avid lesion in liver. Spleen appears normal. Gall bladder appears normal with no evidence of calculus. Stomach and small intestine appear normal. Both kidneys and adrenals appear normal. Prostate gland appears unremarkable. Urinary bladder appears normal. No evidence of free fluid.

Head and neck:

- No evidence of FDG avid cervical lymph nodes. Pharyngeal mucosal space, larynx, thyroid and the cervical trachea appear normal. Parotid glands and



CENTER FOR ADVANCED NUCLEAR
MEDICINE THERANOSTICS



Name :	Mr. Manjunatha G M	PET No :	KMIO20256701 (202411951)
Age/Sex :	45 Years/ Male	UHID :	20240007016
Ref/Dept :	Medical Oncology	Date :	11.06.2025 (13.12.2024)

submandibular glands are unremarkable. No focal mass lesion/mass effect seen within the visualized brain parenchyma.

Chest:

- Both lungs appear normal with no evidence of nodules. No evidence of pleural effusion/ thickening. No evidence of FDG avid mediastinal and axillary lymphadenopathy. Esophagus appears normal. Physiological tracer uptake is noted in myocardium.

Musculoskeletal system:

- No abnormal FDG uptake is seen in bones of axial and appendicular skeleton. No evidence of lytic or sclerotic lesions.

IMPRESSION:

- *Post surgical changes in periampullary region. Biliary stent in situ. No evidence of FDG avid mass lesion at surgical site.*
- *Interval resolution of metabolic activity in portocaval, retroperitoneal and right external iliac lymph node correlation.*
- *No evidence of active disease in the current study.*

Dr. Harish. B DRM, DNB, FEBNM, PGDHA/HM
Professor and Head,
Department of Nuclear Medicine