



VIJAYA DIAGNOSTIC CENTRE®

3-6-16 & 17, Street No. 19, Opp. Lane to Tanishq, Chandra Nagar, Himayatnagar, Hyderabad - 500029

TEST REPORT

Name : **Mr. VEERABHADRA RAO R**
Age/Gender : **62 Years / Male**
Registration ID : **251600016694**
Ref. By : **Dr. SHRUTHI S**
Sample Type : **Whole Blood - EDTA**

Registered on : 18-Aug-2025 07:37
Collected on : 18-Aug-2025 07:39
Released on : 18-Aug-2025 10:10
Printed on : 18-Aug-2025 14:57
Regn Centre : **Yousufguda -160**

GLYCATED HAEMOGLOBIN (HbA1C)

TEST NAME	RESULT	UNIT	BIOLOGICAL REFERENCE INTERVAL
Glycated Haemoglobin	: 10.6	%	Non Diabetic : < 5.7 Pre - Diabetic Range : 5.7 - 6.4 Diabetic Range : >= 6.5

Method: High performance liquid chromatography

Interpretation / Comments:

Reference Range as per American Diabetes Association (ADA) 2025 Guidelines:

- * HbA1c is used to diagnose Diabetes, monitor glycemic control, evaluate the need to change therapy and predict the development of microvascular complications.
- * The Glycemic target for non-pregnant adults with diabetes is <7.0%. Lowering A1c value below or around 7.0% has been shown to reduce microvascular complications and macrovascular disease.
- * Trends in HbA1C values, at interval of 3 months are a better indicator of Glycemic control than a single test.
- * Glycated hemoglobin is proportional to mean plasma glucose level during the previous 6-12 weeks.
- * HbA1c result may not correlate with the blood glucose level in presence of Hemoglobin variants, conditions that affect red cell turnover, Systemic Inflammatory Diseases, Chronic renal failures and liver diseases, recent blood transfusions and hypertriglyceridemia. Clinical Correlation is advised in interpretation of low Values.
- * ADA guidelines 2025 is being followed.



VIJAYA DIAGNOSTIC CENTRE®

3-6-16 & 17, Street No. 19, Opp. Lane to Tanishq, Chandra Nagar, Himayatnagar, Hyderabad - 500029

TEST REPORT

Name : **Mr. VEERABHADRA RAO R**
Age/Gender : **62 Years / Male**
Registration ID : **251600016694**
Ref. By : **Dr. SHRUTHI S**
Sample Type : **Whole Blood - EDTA**

Registered on : 18-Aug-2025 07:37
Collected on : 18-Aug-2025 07:39
Released on : 18-Aug-2025 10:10
Printed on : 18-Aug-2025 14:57
Regn Centre : **Yousufguda -160**

GLYCATED HAEMOGLOBIN (HbA1C)


G11 V03.11 2025-08-18 10:08:53
ID 3251600016694
Sample No. 2025081810040031 SL 0002 - 07
Patient ID
Name
Comment

CALIB (N)			
Y = 1.1201X + 0.6283			
Name	%	Time	Area
FP			
A1A	0.7	0.17	7.06
A1B	0.9	0.24	9.14
F	1.2	0.30	12.16
LA1C+	3.2	0.38	31.04
SA1C	10.6	0.48	86.06
A0	86.2	0.72	832.53
H-VAR			

Total Area 977.99

HbA1c 10.6 %

HbF 1.2 %



18-08-2025 10:08:54 VDC

1 / 1

VDC
HIMAYATH NAGAR



A. Manugna

DR. MANUGNA A

MD BIOCHEMISTRY

Registration No: TSMC/FMR/31118



VIJAYA DIAGNOSTIC CENTRE®

3-6-16 & 17, Street No. 19, Opp. Lane to Tanishq, Chandra Nagar, Himayatnagar, Hyderabad - 500029

TEST REPORT

Name : **Mr. VEERABHADRA RAO R**
Age/Gender : **62 Years / Male**
Registration ID : **251600016694**
Ref. By : **Dr. SHRUTHI S**
Sample Type : **Fluoride Plasma**

Registered on : 18-Aug-2025 07:37
Collected on : 18-Aug-2025 10:08
Released on : 18-Aug-2025 13:12
Printed on : 18-Aug-2025 14:57
Regn Centre : **Yousufguda -160**

POST LUNCH GLUCOSE

TEST NAME	RESULT	UNIT	BIOLOGICAL REFERENCE INTERVAL
Post Lunch Glucose	: 221	mg/dL	Normal : <140 Impaired Glucose Tolerance : 140 - 199 Diabetes : >= 200

Method: Hexokinase

Interpretation / comments:

- Discordant Plasma glucose level (i.e. Post prandial plasma glucose level lower than Fasting plasma glucose level) can be observed, which could be due to the glycemic index and the response to the food consumed, increased insulin secretion, Effects of oral hypoglycemic drugs and Insulin treatment, defective absorption, insufficient dietary intake, endocrine disorders and reactive hypoglycemia etc.
- The diagnosis of Diabetes requires a fasting plasma glucose of ≥ 126 mg/dL and/or a random / 2 hr post glucose value of ≥ 200 mg/dL on at least 2 occasions.
- Very high glucose levels (>450 mg/dL in adults) may result in Diabetic Ketoacidosis & is considered critical.
- ADA (American Diabetes Association) guidelines – 2025 is adopted for the evaluation of diabetes status.



MC-2657

DR. N SUJANA
Consultant Biochemist
Registration No: 50478



VIJAYA DIAGNOSTIC CENTRE®

3-6-16 & 17, Street No. 19, Opp. Lane to Tanishq, Chandra Nagar, Himayatnagar, Hyderabad - 500029

TEST REPORT

Name : **Mr. VEERABHADRA RAO R**
Age/Gender : **62 Years / Male**
Registration ID : **251600016694**
Ref. By : **Dr. SHRUTHI S**
Sample Type : **Fluoride Plasma**

Registered on : 18-Aug-2025 07:37
Collected on : 18-Aug-2025 07:39
Released on : 18-Aug-2025 10:47
Printed on : 18-Aug-2025 14:57
Regn Centre : **Yousufguda -160**

FASTING PLASMA GLUCOSE

TEST NAME	RESULT	UNIT	BIOLOGICAL REFERENCE INTERVAL
Fasting Plasma Glucose <i>Method: Hexokinase</i>	214	mg/dL	Normal : 70 - 100 Impaired Fasting Glucose : 101 - 125 Diabetes : ≥ 126

Interpretation / Comments :

- The diagnosis of Diabetes requires a fasting plasma glucose of ≥ 126 mg/dL and/or a random / 2 hr post glucose value of ≥ 200 mg/dL on at least 2 occasions.
- Very high glucose levels (>450 mg/dL in adults) may result in Diabetic Ketoacidosis & is considered critical.
- ADA (American Diabetes Association) guidelines – 2025 is adopted for the evaluation of diabetes status.



MC-2657

A. Manugna

DR. MANUGNA A
MD BIOCHEMISTRY
Registration No: TSMC/FMR/31118