



ENGINEERING MATHEMATICS I

T. R.Geetha

Department of Science and Humanities

ENGINEERING MATHEMATICS I

HIGHER ORDER DIFFERENTIAL EQUATIONS

T R Geetha

Department of Science and Humanities

ENGINEERING MATHEMATICS I

HIGHER ORDER DIFFERENTIAL EQUATIONS



CLASS CONTENT

- TO SOLVE A NON – HOMOGENEOUS LINEAR DIFFERENTIAL EQUATION OF THE TYPE $f(D)y = X$ WHEN $X =$.

ENGINEERING MATHEMATICS I

NON - HOMOGENEOUS LDE



RULES TO FIND PARTICULAR INTEGRAL

Type (ii); When $X = \sin(ax + b)$ or $\cos(ax + b)$

$$\frac{\sin(ax + b)}{f(D^2)} = \frac{\sin(ax + b)}{f(-a^2)} \quad \text{provided } f(-a^2) \neq 0$$

$$\frac{\cos(ax + b)}{f(D^2)} = \frac{\cos(ax + b)}{f(-a^2)} \quad \text{provided } f(-a^2) \neq 0$$

ENGINEERING MATHEMATICS I

NON - HOMOGENEOUS LDE



Type (ii) continued,

Case of failure; If $f(-a^2) = 0$, then

$$\frac{\sin(ax + b)}{f(D^2)} = x \cdot \frac{\sin(ax + b)}{f'(-a^2)} \quad \text{provided } f'(-a^2) \neq 0$$

$$\frac{\cos(ax + b)}{f(D^2)} = x \cdot \frac{\cos(ax + b)}{f'(-a^2)} \quad \text{provided } f'(-a^2) \neq 0$$

ENGINEERING MATHEMATICS I

NON - HOMOGENEOUS LDE



Type (ii) continued,

If $f'(-a^2) = 0$, then

$$\frac{\sin(ax+b)}{f(D^2)} = x^2 \cdot \frac{\sin(ax+b)}{f''(-a^2)} \quad \text{provided } f''(-a^2) \neq 0$$

$$\frac{\cos(ax+b)}{f(D^2)} = x^2 \cdot \frac{\cos(ax+b)}{f''(-a^2)} \quad \text{provided } f''(-a^2) \neq 0$$

ENGINEERING MATHEMATICS I

NON - HOMOGENEOUS LDE



Type (ii) continued,

The same formula can be used to find the PI for the following functions also.

When $X = \sin ax \cdot \cos bx$ or $\cos ax \cdot \cos bx$
or $\sin ax \cdot \sin bx$

When $X = \sin^2 ax$ or $\cos^2 ax$

When $X = \sin^3 ax$ or $\cos^3 ax$

ENGINEERING MATHEMATICS I

NON - HOMOGENEOUS LDE

Solve; $(D^3 + 1)y = \cos(2x + 3)$

To Find CF

AE is $m^3 + 1 = 0$

$$(m + 1)(m^2 - m + 1) = 0$$

Roots are $m = -1$ and $\frac{1}{2} \pm \frac{\sqrt{3}}{2}i$

$$y_c = c_1 e^{-x} + e^{\frac{x}{2}} \left(c_2 \cos\left(\frac{\sqrt{3}}{2}x\right) + c_3 \sin\left(\frac{\sqrt{3}}{2}x\right) \right)$$

To Find PI

$$\begin{aligned}y_p &= \frac{\cos(2x+3)}{D^3+1} = \frac{\cos(2x+3)}{D.D^2+1} = \frac{\cos(2x+3)}{D.(-2^2)+1} \\&= \frac{\cos(2x+3)}{(1-4D)} = \frac{(1+4D)\cos(2x+3)}{(1+4D)(1-4D)} = \frac{(1+4D)\cos(2x+3)}{(1-16D^2)} \\&= \frac{(1+4D)\cos(2x+3)}{65} = \frac{1}{65}(1+4D)\cos(2x+3) \\&= \frac{1}{65}(\cos(2x+3) - 8\sin(2x+3))\end{aligned}$$

$$\therefore y = y_c + y_p$$

ENGINEERING MATHEMATICS I

NON - HOMOGENEOUS LDE

$$\text{Solve; } (D - 1)^2 (D^2 + 1)y = e^x + \cos^2\left(\frac{x}{2}\right)$$

To Find CF

$$\text{AE is } (m - 1)^2 (m^2 + 1) = 0$$

Roots are $m = 1, 1, \pm i$

$$\text{CF} = (c_1 + c_2 x)e^x + c_3 \cos x + c_4 \sin x$$

ENGINEERING MATHEMATICS I

NON - HOMOGENEOUS LDE



To Find PI

$$PI = \frac{e^x}{(D-1)^2 (D^2+1)} + \frac{\cos^2 \frac{x}{2}}{(D-1)^2 (D^2+1)}$$

$$= PI_1 + PI_2$$

$$\text{where } PI_1 = \frac{e^x}{(D-1)^2 (D^2+1)} \text{ and } PI_2 = \frac{\cos^2 \frac{x}{2}}{(D-1)^2 (D^2+1)}$$

ENGINEERING MATHEMATICS I

NON - HOMOGENEOUS LDE



To Find PI_1

$$PI_1 = \frac{e^x}{(D-1)^2 (D^2+1)} = \frac{1}{(D-1)^2} \left(\frac{e^x}{D^2+1} \right) = \frac{e^x}{2(D-1)^2}$$

$$= x^2 \frac{e^x}{2.2} = \frac{x^2 \cdot e^x}{4}$$

$$\therefore PI_1 = \frac{x^2 \cdot e^x}{4}$$

ENGINEERING MATHEMATICS I

NON - HOMOGENEOUS LDE



To Find PI_2

$$PI_2 = \frac{\cos^2 \frac{x}{2}}{(D-1)^2 (D^2+1)} = \frac{(1 + \cos x) / 2}{(D-1)^2 (D^2+1)}$$

$$= \frac{1}{2(D-1)^2 (D^2+1)} + \frac{\cos x}{2(D-1)^2 (D^2+1)}$$

$$= \frac{1}{4} + \frac{1}{2(D^2+1)} \left(\frac{\cos x}{D^2 - 2D + 1} \right)$$

ENGINEERING MATHEMATICS I

NON - HOMOGENEOUS LDE



To Find PI_2

$$PI_2 = \frac{1}{4} + \frac{1}{2(D^2 + 1)} \left(\frac{\cos x}{-2D} \right) = \frac{1}{4} - \frac{\sin x}{4(D^2 + 1)}$$

$$= \frac{1}{4} - x \frac{\sin x}{4(2D)} = \frac{1}{4} + x \frac{\cos x}{8}$$

$$\therefore PI_2 = \frac{1}{4} + x \frac{\cos x}{8}$$

Thus, $PI = PI_1 + PI_2$ And the GS is $y = CF + PI$



THANK YOU

T R Geetha

Department of Science and Humanities