



# ENGINEERING PHYSICS

---

**Gajanan V Honnavar, Ph.D.**

Department of Science and Humanities

### Class #15

- *Definition of a state vector*
- *Basis and inner products*
- *Orthogonality and Orthonormality of state vectors*
- *Projection operators & their properties*

## State Vectors

- In quantum mechanics, the **state** of a system is described by a **state vector** (also called a *ket*) written as:  $|\psi\rangle$
- In the **wave function picture**, the state is described by  $\psi(x)$ , the probability amplitude in position space.
- In the **vector picture**,  $|\psi\rangle$  is more general — the wave function is just its representation in a **chosen basis**:

$$\psi(x) = \langle x | \psi \rangle$$

Here,  $|x\rangle$  is the position eigenstate.

- The state vector contains all measurable information about the system.

- In quantum mechanics, the state of a system is described by a state vector (ket) in a complex vector space called Hilbert space.
- To describe this state mathematically, we need a set of **basis vectors** — just like in regular geometry we need  $\hat{i}, \hat{j}, \hat{k}$  to describe any 3D vector.
- The inner product of two ket vectors  $|u\rangle$  and  $|v\rangle$  is:

$$\langle u | v \rangle = (\langle u |) | v \rangle = (|u\rangle^*)(|v\rangle) = \sum_i u_i^* v_i$$

It's a complex number that encodes the “overlap” between two states.

- Two vectors  $|u\rangle$  and  $|v\rangle$  are **orthogonal** if their inner product is zero:  $\langle u | v \rangle = 0$
- This means the two states are completely distinct and share no probability amplitude overlap
- The norm (length) of a state vector  $|\psi\rangle$  is:  
$$\|\psi\| = \langle \psi | \psi \rangle$$
- If  $\|\psi\| = 1$ , the vector is normalized.

## Orthonormal Basis

---

A set of vectors  $\{|e_1\rangle, |e_2\rangle, \dots\}$  is an **orthonormal basis** if:

$$\langle e_i | e_j \rangle = \delta_{ij}$$

and the set spans the entire space, so any state can be written as:

$$|\psi\rangle = \sum_i c_i |e_i\rangle$$

With  $c_i = \langle e_i | \psi \rangle$ .

## Example

- If  $|u\rangle$  and  $|v\rangle$  are orthogonal unit vectors, then:  
 $\langle u | v \rangle = 0, \langle u | u \rangle = 1, \langle v | v \rangle = 1$

$$\langle u + v | u - v \rangle = ?$$

Using linearity:

$$\langle u+v|u-v\rangle=\langle u|u\rangle-\langle u|v\rangle+\langle v|u\rangle-\langle v|v\rangle = 1 - 0 + 0 - 1 = 0$$

## Projection Operators

- A **projection operator** projects a state vector onto a specific direction (state) in Hilbert space.
- If  $|v\rangle$  is a normalized vector ( $\langle v | v \rangle = 1$ ), the projection operator onto  $|v\rangle$  is:  $P = |v\rangle\langle v|$
- When  $P$  acts on any state  $|\psi\rangle$ :  $P |\psi\rangle = |v\rangle\langle v | \psi \rangle$
- This gives the component of  $|\psi\rangle$  along  $|v\rangle$ .



- If  $|v\rangle$  is normalized and  $P = |v\rangle\langle v|$

$$P |v\rangle = |v\rangle\langle v | v\rangle = |v\rangle(1) = |v\rangle$$

- So projection operators leave their target state unchanged.

**Q1. Which of the following is the correct representation of an orthonormal basis?**

A.  $\langle e_i | e_j \rangle = 0$  for all  $i, j$

B.  $\langle e_i | e_j \rangle = \delta_{ij}$

C.  $\langle e_i | e_j \rangle = 1$  for all  $i, j$

D.  $\langle e_i | e_j \rangle = e_i + e_j$

Answer: B.  $\langle e_i | e_j \rangle = \delta_{ij}$

**Q2.** A normalized state vector is written as:

$$|\psi\rangle = \frac{1}{\sqrt{2}} (|0\rangle + |1\rangle)$$

If a projection operator  $P = |0\rangle\langle 0|$  acts on  $|\psi\rangle$ , what is the resulting state?

**Answer:**  $\frac{1}{\sqrt{2}} |0\rangle$



# THANK YOU

---

**Gajanan V Honnavar, Ph.D.**

Professor, Department of Science and Humanities

**[Gajanan.honnavar@pes.edu](mailto:Gajanan.honnavar@pes.edu)**