



ENGINEERING MATHEMATICS I

T R Geetha

Department of Science and Humanities

ENGINEERING MATHEMATICS I

HIGHER ORDER DIFFERENTIAL EQUATIONS

T R Geetha

Department of Science and Humanities

ENGINEERING MATHEMATICS I

HIGHER ORDER DIFFERENTIAL EQUATIONS



UNIT CONTENT:

- Linear Differential Equation of higher order
- Homogeneous and Non homogeneous Linear Differential Equation with constant coefficients
- Linear Differential Equation of higher order with variable coefficients
- Method of variation of parameters to solve second order Linear Differential Equation
- Applications to solve problems on Electric Circuits and SHM.

ENGINEERING MATHEMATICS I

HIGHER ORDER DIFFERENTIAL EQUATIONS



CLASS CONTENT:

- Introduction to Linear Differential Equation
- Higher Order Linear Differential Equation
- Different Forms of Linear Differential Equation
- Linear Differential Equation with constant coefficients and variable coefficients
- Complementary Function and Particular Integral

ENGINEERING MATHEMATICS I

HIGHER ORDER DIFFERENTIAL EQUATIONS

Ordinary Differential Equation



```
graph TD; A[Ordinary Differential Equation] --> B[Linear Differential Equation]; A --> C[Non - Linear Differential Equation];
```

Linear Differential Equation

$$\text{Eg. } \frac{dy}{dx} + 2xy = x^2$$

Non - Linear Differential Equation

$$\text{Eg. } y^2 \cdot \frac{dy}{dx} + \sin y = x^2 e^{2y}$$

ENGINEERING MATHEMATICS I

LINEAR DIFFERENTIAL EQUATION



A differential equation in which the dependent variable and its derivative occur in first degree and are not multiplied together is a Linear Differential Equation.

$$\frac{dy}{dx} + P \cdot y = Q \text{ where } P \text{ and } Q \text{ are functions of } x \text{ alone}$$

ENGINEERING MATHEMATICS I

HIGHER ORDER DIFFERENTIAL EQUATION



LINEAR DIFFERENTIAL EQUATIONS OF DIFFERENT ORDER

A linear differential equation of first order is of the form
$$\frac{dy}{dx} + P_1 y = Q$$
 where P and Q are functions of x alone.

A linear differential equation of order two is of the form
$$\frac{d^2y}{dx^2} + P_1 \frac{dy}{dx} + P_2 y = X$$
 where P_1 and P_2 and X are functions of x alone.

A linear differential equation of order 'n' is of the form

$$\frac{d^n y}{dx^n} + P_1 \frac{d^{n-1} y}{dx^{n-1}} + P_2 \frac{d^{n-2} y}{dx^{n-2}} + \dots + P_n y = X$$

where P_1, P_2, \dots, P_n and X are functions of x alone.

ENGINEERING MATHEMATICS I

HIGHER ORDER LINEAR DIFFERENTIAL EQUATION



A general linear differential equation of order n is of the form

$$\frac{d^n y}{dx^n} + a_1 \frac{d^{n-1} y}{dx^{n-1}} + a_2 \frac{d^{n-2} y}{dx^{n-2}} + \dots + a_n y = X$$

where a_1, a_2, \dots, a_n are functions of x or constants
and X is a function of x alone.

Two forms of HigherOrder linear differential equation of order n

1. Linear differential equation with constant coefficients

In this type, the coefficient of derivatives are constants.

Eg. $2 \frac{d^3 y}{dx^3} - 5 \frac{d^2 y}{dx^2} + 7 \frac{dy}{dx} + y = x^2$

ENGINEERING MATHEMATICS I

HIGHER ORDER LINEAR DIFFERENTIAL EQUATION



Two forms of Higher Order linear differential equation of order n

2. Linear differential equation with variable coefficients

$$\frac{d^n y}{dx^n} + a_1 \frac{d^{n-1} y}{dx^{n-1}} + a_2 \frac{d^{n-2} y}{dx^{n-2}} + \dots + a_n y = X$$

In this type, the coefficient of derivatives $a_1, a_2 \dots a_n$ are functions of x .

$$\text{Eg. } 2x^3 \frac{d^3 y}{dx^3} - (5x^2 + 2) \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + y = x^2 \sin x$$

ENGINEERING MATHEMATICS i

HIGHER ORDER LINEAR DIFFERENTIAL EQUATION

Linear differential equation with constant coefficients

$$\frac{d^n y}{dx^n} + a_1 \frac{d^{n-1} y}{dx^{n-1}} + a_2 \frac{d^{n-2} y}{dx^{n-2}} + \dots + a_n y = X$$

HOMOGENEOUS LDE

RHS X is zero

Eg. $2 \frac{d^2 y}{dx^2} - 3 \frac{dy}{dx} + 9 y = 0$

NON HOMOGENEOUS LDE

RHS X is a nonzero function of x

Eg. $2 \frac{d^2 y}{dx^2} - 3 \frac{dy}{dx} + 9 y = x \cdot \sinh x$

ENGINEERING MATHEMATICS I

HIGHER ORDER LINEAR DIFFERENTIAL EQUATION



DIFFERENTIAL OPERATOR

$$D = \frac{d}{dx}, \quad \frac{d^2}{dx^2} = D^2, \quad \frac{d^3}{dx^3} = D^3, \quad \dots \quad \frac{d^n}{dx^n} = D^n$$

Thus, in terms of operator notation

$$\frac{d^n y}{dx^n} + a_1 \frac{d^{n-1} y}{dx^{n-1}} + a_2 \frac{d^{n-2} y}{dx^{n-2}} + \dots + a_n y = X \text{ can be written as}$$

$$D^n y + a_1 D^{n-1} y + a_2 D^{n-2} y + \dots + a_n y = X$$

$$f(D)y = X$$

$$\text{where } f(D) = D^n + a_1 D^{n-1} + a_2 D^{n-2} + \dots + a_n$$

called linear differential operator.

ENGINEERING MATHEMATICS I

LINEAR DIFFERENTIAL EQUATION WITH CONSTANT COEFFICIENTS



Consider a non homogeneous equation of the type

$$\frac{d^n y}{dx^n} + a_1 \frac{d^{n-1} y}{dx^{n-1}} + a_2 \frac{d^{n-2} y}{dx^{n-2}} + \dots + a_n y = X$$

$$f(D)y = X$$

The general solution of the non homogeneous linear differential equation has two components:

Complementary Function and particular Integral.

The complementary function is the complete solution of homogeneous linear differential equation.

Linear Dependence and Linear Independent Functions

A set of functions $f_1(x), f_2(x), \dots, f_n(x)$ are said to be linearly dependent on an interval 'I' if there exists constants c_1, c_2, \dots, c_n not all zero, such that $c_1 f_1 + c_2 f_2 + \dots + c_n f_n = 0$.

If not linearly dependent, the functions are called linearly independent functions.

In other words, if the set is linearly independent, then

$$c_1 f_1 + c_2 f_2 + \dots + c_n f_n = 0 \quad \Rightarrow \quad c_1 = c_2 = c_3 = \dots = c_n = 0$$

ENGINEERING MATHEMATICS I

LINEAR DIFFERENTIAL EQUATION WITH CONSTANT COEFFICIENTS



Linear Dependent and Linear Independent Functions

Two functions y_1 and y_2 are linearly independent when $C_1 y_1 + C_2 y_2 = 0$ for non zero constants C_1 and C_2 .

-

ENGINEERING MATHEMATICS I

LINEAR DIFFERENTIAL EQUATION WITH CONSTANT COEFFICIENTS



SECOND ORDER HOMOGENEOUS LINEAR DIFFERENTIAL EQUATION WITH CONSTANT COEFFICIENTS:

Consider a second order LDE with constant coefficients of the type

$$\frac{d^2y}{dx^2} + k_1 \frac{dy}{dx} + k_2 y = 0$$

If y_1 and y_2 are two linearly independent solutions of a second order homogeneous linear differential equation, then $y_1 + y_2$ is also a solution of the differential equation.

It is the complete solution of the differential equation.

ENGINEERING MATHEMATICS I

LINEAR DIFFERENTIAL EQUATION WITH CONSTANT COEFFICIENTS



EXTENSION OF THE PROPERTY TO A LINEAR DIFFERENTIAL EQUATION OF ORDER 'n' :

If the nth order LDE with constant coefficients

$\frac{d^n y}{dx^n} + a_1 \frac{d^{n-1} y}{dx^{n-1}} + a_2 \frac{d^{n-2} y}{dx^{n-2}} + \dots + a_n y = 0$ has 'n' linearly independent solutions

$y_1, y_2, y_3, \dots, y_n$, then their linear combination

$c_1 y_1 + c_2 y_2 + \dots + c_n y_n$ is also a solution of the differential equation.

$y = c_1 y_1 + c_2 y_2 + \dots + c_n y_n$, the complete solution is called

COMPLEMENTARY FUNCTION of the differential equation.

ENGINEERING MATHEMATICS I

LINEAR DIFFERENTIAL EQUATION WITH CONSTANT COEFFICIENTS



NON HOMOGENEOUS LINEAR DIFFERENTIAL EQUATION WITH
CONSTANT COEFFICIENTS:

Consider a non homogeneous linear differential equation $f(D)y = X$

If $y = y_c$ is the complete solution of $f(D)y = 0$ and $y = y_p$ is a particular solution of the differential equation $f(D)y = X$, then the complete solution is $y = y_c + y_p$

The particular solution ' y_p ' is called PARTICULAR INTEGRAL.



THANK YOU

T R Geetha

Department of Science and Humanities