



**PES**  
UNIVERSITY  
**O N L I N E**

## **ENGINEERING PHYSICS**

---

**Radhakrishnan S, Ph.D.**

Department of Science and Humanities

### Week #1

Cl#1    Review of Electric and magnetic fields

**Cl#2**    *EM Wave equation*

Cl#3    Energy transported by EM Waves

Cl#4    Max Planck's Black Body Radiation equation

### Class #2

- *Maxwell's equation in differential form*
- *Maxwell's Equations in free space*
- *Ideas of Electric and Magnetic waves*
- *Wave equation*

### ➤ *Suggested Reading*

1. *Fundamentals of Physics, Halliday, Resnik, Chapter 34*
2. *NCERT Physics Book I grade 12 – Chapter 8*

### ➤ *Reference Videos*

1. <https://nptel.ac.in/courses/108/106/108106073/>
2. *Engineering Physics week1 class1*

### Electric fields

*the divergence of the electric field is proportional to the charge density*

$$\nabla \cdot \vec{E} = \frac{\rho}{\epsilon_0}$$

### Magnetic Fields

*the divergence of the magnetic flux is uniformly zero*

$$\nabla \cdot \vec{B} = 0$$

### *Faraday's law of electromagnetic induction*

*The curl of the induced electric field in a closed loop is proportional to the rate of change of magnetic flux linked with the loop*

$$\nabla \times \vec{E} = - \frac{\partial \vec{B}}{\partial t}$$

### Ampere - Maxwell circuital law

*The curl of the magnetic field in a closed loop is equal to the sum of the current density and the displacement current due to the time varying electric field*

$$\nabla \times \vec{B} = \mu_0 \vec{J} + \mu_0 \epsilon_0 \frac{\partial \vec{E}}{\partial t}$$

# ENGINEERING PHYSICS

## Maxwell's equations in free space

Free space implies – charges and currents do not exist



$$\vec{\nabla} \cdot \vec{E} = 0$$

1.

$$\vec{\nabla} \cdot \vec{B} = 0$$

2.

$$\vec{\nabla} \times \vec{E} = - \frac{\partial \vec{B}}{\partial t}$$

3

$$\vec{\nabla} \times \vec{B} = +\mu_0 \epsilon_0 \frac{\partial \vec{E}}{\partial t}$$

4.

# ENGINEERING PHYSICS

## Electric waves in free space

*The curl of equation 3.*  $\vec{\nabla} \times (\vec{\nabla} \times \vec{E}) = \vec{\nabla} \times \left( -\frac{\partial \vec{B}}{\partial t} \right)$

$$\vec{\nabla}(\vec{\nabla} \cdot \vec{E}) - \nabla^2 \vec{E} = \left( -\frac{\partial \vec{\nabla} \times \vec{B}}{\partial t} \right)$$

For free space  $\vec{\nabla} \cdot \vec{E} = 0$

$$-\nabla^2 \vec{E} = \left( -\frac{\partial \vec{\nabla} \times \vec{B}}{\partial t} \right)$$

Substituting for curl of B from equation 4.

$$\nabla^2 \vec{E} = \left( \mu_o \epsilon_o \frac{\partial^2 E}{\partial t^2} \right)$$

$$\nabla^2 \vec{E} = \left( \frac{1}{c^2} \frac{\partial^2 \vec{E}}{\partial t^2} \right) \dots\dots$$

With  $\mu_o \epsilon_o = \frac{1}{c^2}$

A wave equation for electric wave propagating in free space !

- *The curl of Maxwell's equation 4.*

$$\vec{\nabla} \times \vec{\nabla} \times \vec{B} = \vec{\nabla} \times (+\mu_0 \epsilon_0 \frac{\partial \vec{E}}{\partial t}) \rightarrow$$
$$\nabla^2 \vec{B} = \left( \frac{1}{c^2} \frac{\partial^2 \vec{B}}{\partial t^2} \right) \rightarrow$$

*a magnetic wave propagating in free space with speed of light*

- *Both electric and magnetic waves propagate with the speed of light*
- *Maxwell's predication of light (radiation) as electromagnetic waves*

# ENGINEERING PHYSICS

## Electromagnetic waves in free space

---



***Light waves are transverse waves ---***


***Electromagnetic waves are transverse waves !***

***The electric and magnetic fields are perpendicular to the direction of propagation of the electromagnetic wave***

# ENGINEERING PHYSICS

## Electromagnetic waves in free space

- Let a plane EM wave be propagating along the z direction ( $E_z = 0$ )
- Instantaneous electric field  $E_x = E_{xo} \sin(\omega t + kz)$



- *Plane wave implies  $E_y = 0$*
- *From the third Maxwell's equation*


$$\bullet \quad \vec{\nabla} \times \vec{E}_x = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ E_x & E_y & E_z \end{vmatrix} = -\hat{j} \frac{\partial E_x}{\partial z} = - \frac{\partial \vec{B}}{\partial t}$$
$$\bullet \quad \frac{\partial \vec{B}}{\partial t} \text{ can be evaluated since } E_x \text{ is known}$$

- Integrate  $\frac{\partial \vec{B}}{\partial t}$  with respect to time to obtain
- $B_y = \hat{j} \frac{E_{xo}}{\omega/k} \sin(\omega t + kz)$
- $\frac{\omega}{k} = \text{the velocity of the wave} = c$
- magnetic field is oriented in the y direction
- perpendicular to the Electric field (x direction)
- E and B are mutually perpendicular to each other

- $E_x = \hat{i}E_{xo} \sin(\omega t + kz)$
- $B_y = \hat{j} \frac{E_{xo} \sin(\omega t + kz)}{\omega/k}$
- ***in phase with the electric field variations***
- **$\omega/k = c$  is the phase velocity of the wave**
- ***Thus  $B_y$  is perpendicular to  $E$  and  $|B_y| = \frac{|E_x|}{c}$***

## Electromagnetic waves in free space

- *Electromagnetic wave*
- *E and B are mutually perpendicular to each other and the direction of propagation*



**The concepts which are not correct are....**

1. Electric waves in free space are longitudinal
2. Magnetic waves in free space are transverse
3. The curl of a magnetic field is uniformly zero
4. The divergence of a magnetic field can be non zero
5. The curl of an electric field is always linked to a time varying magnetic field
6. The divergence of a vector field is also a scalar



**THANK YOU**

---

**Radhakrishnan S, Ph.D.**

Professor, Department of Science and Humanities

**sradhakrishnan@pes.edu**

+91 80 21722683 Extn 759