



ELSEVIER

Available at
WWW.MATHEMATICSWEB.ORG
 POWERED BY SCIENCE @ DIRECT®

Journal of Statistical Planning and
 Inference 113 (2003) 385–402

journal of
 statistical planning
 and inference

www.elsevier.com/locate/jspi

On the dependence of the Berry–Esseen bound on dimension

V. Bentkus*,¹

Vilnius Institute of Mathematics and Informatics, Akademijos 4, 232600 Vilnius, Lithuania

Received 14 July 2001; received in revised form 30 October 2001; accepted 20 November 2001

Abstract

Let X be a random vector with values in \mathbb{R}^d . Assume that X has mean zero and identity covariance. Write $\beta = \mathbf{E}|X|^3$. Let S_n be a normalized sum of n independent copies of X . For $\Delta_n = \sup_{A \in \mathcal{C}} |\mathbb{P}\{S_n \in A\} - \nu(A)|$, where \mathcal{C} is the class of convex subsets of \mathbb{R}^d , and ν is the standard d -dimensional normal distribution, we prove a Berry–Esseen bound $\Delta_n \leq 400d^{1/4}\beta/\sqrt{n}$. Whether one can remove or replace the factor $d^{1/4}$ by a better one (eventually by 1), remains an open question. © 2002 Published by Elsevier Science B.V.

MSC: 60F05

Keywords: Multidimensional; Berry–Esseen bound; Dependence on dimension

1. Introduction and results

We provide a Berry–Esseen type bound with the best known dependence on dimension. Our proof is less technically involved compared to the classical proofs using characteristic functions (see Bhattacharya and Rao, 1986). It seems to be a difficult problem to prove the result using the Fourier method.

Let \mathbb{R}^d be a real Euclidean d -dimensional space of vectors $x = (x_1, \dots, x_d)$ with the norm $|x|^2 = x_1^2 + \dots + x_d^2$ and the scalar product $\langle x, x \rangle = |x|^2$.

Let X, X_1, \dots, X_n be i.i.d. random vectors with common mean $\mathbf{E}X_j = 0$ and the identity covariance, so that $\mathbf{E}\langle X, x \rangle^2 = |x|^2$. Write $\beta = \mathbf{E}|X|^3$. By Y, Y_1, \dots, Y_n we denote i.i.d. standard normal random vectors, that is Gaussian random vectors with mean zero and

* Corresponding author.

¹ Research supported by the Max Planck Institute for Mathematics, Bonn.

identity covariance operator. We assume throughout that X 's, Y 's and other random vectors and variables are independent if the contrary is not clear from the context.

Let \mathcal{A} be a class of (measurable) subsets $A \subset \mathbb{R}^d$ (below we assume that all sets and functions are measurable). We shall obtain upper bounds for the quantity

$$\Delta_n \equiv \Delta_n(\mathcal{A}) = \sup_{A \in \mathcal{A}} |\mathbb{P}\{S_n \in A\} - \mathbb{P}\{Y \in A\}|,$$

where $S_n = (X_1 + \dots + X_n)/\sqrt{n}$. The bounds have the shape

$$\Delta_n \leq c_d(\mathcal{A})\beta/\sqrt{n}, \quad \beta = \mathbf{E}|X|^3 \quad (1.1)$$

with some constant $c_d(\mathcal{A})$ which depends only on the dimension and the class \mathcal{A} . For example, for the class \mathcal{B} of (Euclidean) balls we have $\Delta_n(\mathcal{B}) \leq c\beta/\sqrt{n}$, where c is an absolute constant. For the class \mathcal{C} of convex subsets of \mathbb{R}^d , it holds $\Delta_n(\mathcal{C}) \leq 400d^{1/4}\beta/\sqrt{n}$. Whether the dependence on d in the last bound is optimal, remains an open question. One can show (see Nagaev, 1976) that for the class of convex sets $\Delta_n(\mathcal{C}) \geq c_0\beta/\sqrt{n}$, where $c_0 > 0$ is an absolute positive constant. Hence, the upper and lower bounds for $\Delta_n(\mathcal{C})$ differ by the factor $d^{1/4}$.

We need some definitions and conditions related to \mathcal{A} and the standard normal distribution $\nu(\mathcal{A}) = \mathbb{P}\{Y \in A\}$.

For a class, say \mathcal{A} , of subsets $A \subset \mathbb{R}^d$, consider the following conditions:

- (i) \mathcal{A} is invariant under rescaling, that is, $aA \in \mathcal{A}$, if $A \in \mathcal{A}$ and a number $a \in \mathbb{R}$ is positive;
- (ii) \mathcal{A} is shift invariant, that is, $x + A \in \mathcal{A}$, if $A \in \mathcal{A}$ and $x \in \mathbb{R}^d$;
- (iii) \mathcal{A} is invariant under taking of ε -neighborhoods, for $\varepsilon > 0$. More precisely, $A^\varepsilon, A^{-\varepsilon} \in \mathcal{A}$ if $A \in \mathcal{A}$. Here

$$A^\varepsilon = \{x \in \mathbb{R}^d : \rho_A(x) \leq \varepsilon\} \quad \text{and} \quad A^{-\varepsilon} = \{x \in A : B_\varepsilon(x) \subset A\},$$

where $\rho_A(x) = \inf_{y \in A} |x - y|$ is the distance between $A \subset \mathbb{R}^d$ and $x \in \mathbb{R}^d$, and $B_\varepsilon(x) = \{y \in \mathbb{R}^d : |x - y| \leq \varepsilon\}$.

For example, the class \mathcal{C} of convex subsets of \mathbb{R}^d satisfies the conditions (i)–(iii). Another example provides the class \mathcal{B} of Euclidean balls $B_r(x)$, $r \geq 0$, $x \in \mathbb{R}^d$. The class of rectangles does not satisfy the condition (iii) since an ε -neighborhood of a rectangle is not a rectangle.

We consider the following condition on a class \mathcal{A} of subsets of \mathbb{R}^d and the standard normal distribution $\nu(A) = \mathbb{P}\{Y \in A\}$: there exist constants, say $a_d = a_d(\mathcal{A})$, depending only on d and \mathcal{A} such that

$$\mathbb{P}\{Y \in A^\varepsilon \setminus A\} \leq a_d \varepsilon, \quad \mathbb{P}\{Y \in A \setminus A^{-\varepsilon}\} \leq a_d \varepsilon \quad \text{for all } A \in \mathcal{A} \text{ and } \varepsilon > 0. \quad (1.2)$$

We shall refer to $a_d(\mathcal{A})$ as to the isoperimetric constant of \mathcal{A} . For the class \mathcal{B} of balls we have $\sup_{d \geq 1} a_d(\mathcal{B}) < \infty$. Ball (1993) has proved that

$$e^{-1} \sqrt{\ln d} \leq \sup_{A \in \mathcal{C}} \int_{\partial A} p(x) ds \leq 4d^{1/4}, \quad (1.3)$$

where $p(x)$ is the standard normal d -dimensional density and ds is the surface measure on the boundary ∂A of A . Using (1.3) one can show that for the class of convex sets

$$e^{-1}\sqrt{\ln d} \leq a_d(\mathcal{C}) \leq 4d^{1/4}. \quad (1.4)$$

To find precise dependence of $a_d(\mathcal{C})$ on d as $d \rightarrow \infty$ was a longstanding open problem. Recently Nazarov (2001a, b) showed that up to an absolute factor $a_d(\mathcal{C})$ behaves as $d^{1/4}$, see Lemma 2.6 below, where we provide Nazarov's lower bound. Indirectly this result supports a conjecture that the dependence on d as in Theorem 1.1 below is optimal or near to optimal since in all cases which we can analyze the isoperimetric constant controls the constant in the Berry–Esseen bound up to an absolute factor.

Our result is the following theorem.

Theorem 1.1. *Let a class \mathcal{A} of convex sets satisfy the conditions (i)–(iii). Furthermore, assume that \mathcal{A} and the standard normal distribution satisfy the condition (1.2). Then*

$$\Delta_n \leq 100b_d\beta/\sqrt{n}, \quad b_d = \max\{1, a_d\}.$$

In particular, for the class of convex sets we have $\Delta_n \leq 400d^{1/4}\beta/\sqrt{n}$. In the case of the class of all Euclidean balls the bound improves to $\Delta_n \leq c\beta/\sqrt{n}$, where c is an absolute constant.

The constants in Theorem 1.1 can be considerably improved, especially for larger d . Our aim was just to provide explicit constants keeping the proof and notation as simple as possible. Conditions (i)–(iii) on class \mathcal{A} can be relaxed using a more refined techniques.

The question of the dependence of the Berry–Esseen bound on dimension has drawn attention of many authors, let us mention here only Nagaev (1976), Senatov (1980), Sazonov (1981), Götze (1991), where the bounds depend on the dimension as d . Our approach is very close to that of the author in Bentkus (1986a), where the bound depends on d as \sqrt{d} .

The proof of Theorem 1.1 based on induction in n and Taylor expansions. Usually such proofs are very robust against extensions to general spaces and abstract settings, and can be used as well to prove large deviation theorems (see Bentkus, 1986b). Direct inductual proofs is a commonly used tool to obtain convergence rates in the CLT, see, for example, books Sazonov (1981), Paulauskas and Račkauskas (1989), Senatov (1998). A feature of our proof is that we apply a special smoothing with differentiable functions approximating indicator functions of convex sets. The smoothness properties of these functions rely on good differentiability properties of Euclidean distances. In the context of limit theorems smoothness of Euclidean distances were earlier applied by Senatov (1984) to estimate Prokhorov's metric.

2. Auxiliary lemmas

We provide the proof of Theorem 1.1 a little later since first we would like to formulate and prove several auxiliary lemmas.

Results like the next (very simple technically) Lemma 2.1 usually are called smoothing inequalities or lemmas. Smoothing inequalities allow to replace probabilities like $\mathbb{P}\{S_n \in A\} = \mathbf{E}\mathbb{I}\{S_n \in A\}$, which are expectations of non-differentiable indicator functions $x \mapsto \mathbb{I}\{x \in A\}$, by expectations of smooth functions, say φ . Such a replacement allows to apply the differential calculus. In Lemma 2.1 we use smooth functions $x \mapsto \varphi(x)$ which approximate indicator functions. Another often used way to make smoothing is via convolutions with certain measures and Fourier transforms (or characteristic functions), that is, via the family of exponential functions $\varphi_t(x) = \exp\{i\langle t, x \rangle\}$, $t \in \mathbb{R}^d$, see Bhattacharya and Rao (1986).

Lemma 2.1. *Let $\varepsilon > 0$ be a positive number. Let a set $A \subset \mathbb{R}^d$ and functions $0 \leq \varphi_1, \varphi_2 \leq 1$ satisfy*

$$\varphi_1(x) = 1 \quad \text{for } x \in A, \quad \varphi_1(x) = 0 \quad \text{for } x \notin A^\varepsilon,$$

$$\varphi_2(x) = 1 \quad \text{for } x \in A^{-\varepsilon}, \quad \varphi_2(x) = 0 \quad \text{for } x \notin A.$$

Then, for arbitrary r. v. X and Y , the difference $\gamma(A) = \mathbb{P}\{X \in A\} - \mathbb{P}\{Y \in A\}$ satisfies

$$|\gamma(A)| \leq \max_{k=1,2} |\gamma(\varphi_k)| + \max\{\mathbb{P}\{Y \in A^\varepsilon \setminus A\}, \mathbb{P}\{Y \in A \setminus A^{-\varepsilon}\}\}, \quad (2.1)$$

where $\gamma(\varphi_k) = \mathbf{E}\varphi_k(X) - \mathbf{E}\varphi_k(Y)$.

Let a class \mathcal{A} of sets satisfy condition (iii). Assume that we have a family, say $\{\varphi_{\varepsilon,A}: A \in \mathcal{A}\}$, of functions $\varphi_{\varepsilon,A}: \mathbb{R}^d \rightarrow \mathbb{R}$ such that $0 \leq \varphi_{\varepsilon,A} \leq 1$ and $\varphi_{\varepsilon,A}(x) = 1$ for $x \in A$, $\varphi_{\varepsilon,A}(x) = 0$, for $x \notin A^\varepsilon$. Then

$$\begin{aligned} \sup_{A \in \mathcal{A}} |\gamma(A)| &\leq \sup_{A \in \mathcal{A}} |\gamma(\varphi_{\varepsilon,A})| \\ &\quad + \max \left\{ \sup_{A \in \mathcal{A}} \mathbb{P}\{Y \in A^\varepsilon \setminus A\}, \sup_{A \in \mathcal{A}} \mathbb{P}\{Y \in A \setminus A^{-\varepsilon}\} \right\} \end{aligned} \quad (2.2)$$

with $\gamma(\varphi_{\varepsilon,A}) = \mathbf{E}\varphi_{\varepsilon,A}(X) - \mathbf{E}\varphi_{\varepsilon,A}(Y)$.

Proof. We prove only (2.2) since the proof of (2.1) is simpler. We consider first the case $\gamma(A) \geq 0$. Writing for brevity $\varphi = \varphi_{\varepsilon,A}$, we have

$$\begin{aligned} |\gamma(A)| &= \gamma(A) \leq \mathbf{E}\varphi(X) - \mathbf{E}\varphi(Y) + \mathbf{E}\varphi(Y) - \mathbb{P}\{Y \in A\} \\ &\leq |\mathbf{E}\varphi(X) - \mathbf{E}\varphi(Y)| + \mathbb{P}\{Y \in A^\varepsilon \setminus A\} \end{aligned} \quad (2.3)$$

since the function φ satisfies $0 \leq \varphi \leq 1$ and equals 1 on the set A . Taking $\sup_{A \in \mathcal{A}}$ in (2.3) leads to (2.2).

Now consider the case $\gamma(A) < 0$. We have to estimate $-\gamma(A) = \mathbb{P}\{Y \in A\} - \mathbb{P}\{X \in A\}$. The proof depends on whether $A^{-\varepsilon} = \emptyset$ or $A^{-\varepsilon} \neq \emptyset$.

If $A^{-\varepsilon} = \emptyset$ then $\mathbb{P}\{Y \in A^{-\varepsilon}\} = 0$ and

$$-\gamma(A) \leq \mathbb{P}\{Y \in A\} = \mathbb{P}\{Y \in A\} - \mathbb{P}\{Y \in A^{-\varepsilon}\} \leq \mathbb{P}\{Y \in A \setminus A^{-\varepsilon}\}. \quad (2.4)$$

Taking $\sup_{A \in \mathcal{A}}$ in (2.4) we get (2.2).

If $A^{-\varepsilon} \neq \emptyset$, we can in essence repeat the proof of the case $\gamma(A) \geq 0$. Writing for brevity $\varphi = \varphi_{\varepsilon, A^{-\varepsilon}}$, we have

$$\begin{aligned} -\gamma(A) &\leq \mathbb{P}\{Y \in A\} - \mathbf{E}\varphi(Y) + \mathbf{E}\varphi(Y) - \mathbf{E}\varphi(X) \\ &\leq \mathbb{P}\{Y \in A \setminus A^{-\varepsilon}\} + |\mathbf{E}\varphi(X) - \mathbf{E}\varphi(Y)| \end{aligned} \quad (2.5)$$

since the function φ satisfies $0 \leq \varphi \leq 1$ and equals 1 on the set $A^{-\varepsilon}$. Noting that by the condition (iii) $A^{-\varepsilon} \in \mathcal{A}$ and taking in (2.5) $\sup_{A^{-\varepsilon}} < \sup_A$ we obtain (2.2). \square

Lemma 2.2. *Let $A \subset \mathbb{R}^d$ be a closed convex set. Then the distance function*

$$\rho(x) = \min\{|x - y| : y \in A\}$$

satisfies $|\rho'(x)| \leq 1$, for $x \in A^c = \mathbb{R}^d \setminus A$. Furthermore, its derivative satisfies the following Lipschitz type condition:

$$|\rho'(x) - \rho'(y)| \leq |x - y|/r \quad \text{for all } x, y \in A^c, \quad (2.6)$$

with $r = \min\{\rho(x), \rho(y)\}$.

Proof. Without loss of generality we assume that $r = \rho(X) \geq s = \rho(y)$.

For $x \in A^c$, by x_0 we denote a unique point $x_0 \in A$ closest to x . Consider the unit vector $e_1 = (x - x_0)/|x - x_0|$ and the ray $R(x_0) = \{x_0 + te_1 : t > 0\}$. Then $\rho'(z) = e_1$, for all $z \in R_1$. Let e_2 and $R(y_0)$ be a unit vector and ray corresponding to y_0 . It is clear that $x = x_0 + re_1 \in \mathbb{R}(x_0)$ and $y = y_0 + se_2 \in \mathbb{R}(y_0)$. Introduce the hyper-plane P_1 (resp. P_2) which is a hyper-plane orthogonal to the ray R_1 (resp. R_2) such that $x_0 \in P_1$ (resp. $y_0 \in P_2$). Analytically, P_1 is given by $P_1 = \{x_0 + u \in \mathbb{R}^d : \langle u, e_1 \rangle = 0\}$. It is clear that the set A is a subset of the half-space $P_1^- = \{x_0 + u \in \mathbb{R}^d : \langle u, e_1 \rangle \leq 0\}$.

We consider separately the following two cases:

- (i) $P_1 \cap P_2 \neq \emptyset$;
- (ii) $P_1 \cap P_2 = \emptyset$;

- (i) There exists $z \in P_1 \cap P_2$. Consider the rays

$$R_3 = \{z + te_1 : t > 0\} \quad \text{and} \quad R_4 = \{z + te_2 : t > 0\}$$

and points $\bar{x} = z + re_1 \in R_3$ and $\bar{y} = z + se_2 \in R_4$. It is clear that

$$|\bar{x} - \bar{y}| \leq |x - y|. \quad (2.7)$$

Indeed, $A \subset P_1^- \cap P_2^-$ since A is a convex set. The angle between planes P_1 and P_2 is bounded from above by π since A is convex. Therefore, if we have two rays, say $R(u)$ and $R(v)$, then the distance between points $u + re_1 \in R(u)$ and $v + se_2 \in R(v)$ becomes minimal when $u = v \in P_1 \cap P_2$, which proves (2.7) (this fact is obvious geometrically and it is not difficult to provide an analytical proof of it). Inequality (2.7) and $\rho'(x) = e_1$, $\rho'(y) = e_2$ show that (2.6) is implied by the following inequality:

$$|e_1 - e_2| \leq |\bar{x} - \bar{y}|/r = |e_1 - se_2/r|. \quad (2.8)$$

Inequality (2.8) is just a simple fact from the planar geometry. Nonetheless, let us provide an analytical proof of it. Taking the square, (2.8) reduces to

$$2 - 2\langle e_1, e_2 \rangle \leq 1 + \frac{s^2}{r^2} - 2\frac{s}{r}\langle e_1, e_2 \rangle. \quad (2.9)$$

Note that $0 \leq \langle e_1, e_2 \rangle \leq 1$ since the angle between planes P_1 and P_2 is bounded from above by π . Hence, for $s \geq r$, the right-hand side of (2.9) attains its minimal value $2 - 2\langle e_1, e_2 \rangle$ when $s = r$, which proves (2.9) and (2.6) in the case (i).

(ii) Now the planes P_1 and P_2 are parallel and $e_2 = -e_1$. Therefore $\rho'(x) = e_1$ and $\rho'(y) = -e_1$. The inequality (2.6) takes the form $2 \leq |x_0 - y_0 + (r+s)e_1|/r$. It is clear geometrically that $|x_0 - y_0 + (r+s)e_1|$ assumes its minimal value when $P_1 = P_2$, $x_0 = y_0$ and $s = r$. This value is 2, which proves (ii) and the lemma. \square

Lemma 2.3. *Let a set $A \subset \mathbb{R}^d$ be convex. Then for any $\varepsilon > 0$ there exists a function φ (which depends only on ε and A) such that*

$$\varphi(x) = 1 \quad \text{for } x \in A, \quad \varphi(x) = 0 \quad \text{for } x \in \mathbb{R}^d \setminus A^\varepsilon, \quad 0 \leq \varphi \leq 1, \quad (2.10)$$

and

$$|\varphi'(x)| \leq \frac{2}{\varepsilon}, \quad |\varphi'(x) - \varphi'(y)| \leq \frac{8|x - y|}{\varepsilon^2}. \quad (2.11)$$

Furthermore, we can choose φ to have the form $\varphi(x) = \psi(\rho(x)/\varepsilon)$, where $\rho(x)$ is the distance between x and A , and $\psi: \mathbb{R} \rightarrow \mathbb{R}$ is a continuously differentiable non-negative non-increasing function such that $\int_{\mathbb{R}} |\psi'(t)| dt = 1$.

Proof. Without loss of generality we assume that the set A is closed.

Let $\psi: \mathbb{R} \rightarrow [0, 1]$ be a function such that $\psi''(x) = 0$, for $x \leq 0$ or $x \geq 1$, and $\psi''(x) = -4$, for $0 < x \leq 1/2$, $\psi''(x) = 4$, for $1/2 < x < 1$. We define

$$\psi'(x) = \int_{-\infty}^x \psi''(t) dt \quad \text{and} \quad \psi(x) = 1 + \int_{-\infty}^x \psi'(t) dt.$$

The function ψ is non-increasing and has the following properties: $\psi(x) = 1$, for $x \leq 0$, $\psi(x) = 0$, for $x \geq 1$, $0 \leq \psi \leq 1$,

$$|\psi'(t)| \leq 4|t|, \quad |\psi'(t)| \leq 2, \quad |\psi'(t) - \psi'(s)| \leq 4|t - s|. \quad (2.12)$$

We define $\varphi(x) = \psi(\rho(x)/\varepsilon)$. The function φ clearly satisfies relations (2.10).

Using Lemma 2.2 and the properties of ψ , we have

$$|\varphi'(x)| = \varepsilon^{-1} |\psi'(\rho(x)/\varepsilon) \rho'(x)| \leq 2/\varepsilon,$$

which proves the first inequality in (2.11).

Let us prove the second inequality in (2.11). Let $x, y \in \mathbb{R}^d$.

If both $x, y \in A$, then $\varphi'(x) = \varphi'(y) = 0$, and the second inequality in (2.11) is obviously fulfilled. To check that $\varphi'(x) = 0$, for $x \in A$, we consider two cases: (i) x is in the interior $A \setminus \partial A$ of A ; (ii) $x \in \partial A$. In case (i) $\varphi'(x) = 0$ since φ is a constant

function in the open set $A \setminus \partial A$. In case (ii), using $|\psi'(t)| \leq 4|t|$, we have

$$|\varphi(x+h) - \varphi(x)| \leq \int_0^{\rho(x+h)} |\psi'(t)| dt \leq 4\rho^2(x+h) \leq 4|h|^2$$

which implies $\varphi'(x) = 0$.

Now let both $x, y \in A^c$. We can assume that $r = \rho(x) \leq s = \rho(y)$. Lemma 2.2 and properties of ψ yield

$$\begin{aligned} |\varphi'(x) - \varphi'(y)| &= \varepsilon^{-1} |\psi'(r/\varepsilon)\rho'(x) - \psi'(s/\varepsilon)\rho'(y)| \\ &\leq \varepsilon^{-1} |\psi'(r/\varepsilon)| |\rho'(x) - \rho'(y)| + \varepsilon^{-1} |\psi'(r/\varepsilon) - \psi'(s/\varepsilon)| |\rho'(y)| \\ &\leq \frac{4r}{\varepsilon^2} \frac{|x-y|}{r} + \frac{4}{\varepsilon^2} |r-s| \leq \frac{8}{\varepsilon^2} |x-y| \end{aligned}$$

which proves the second inequality in (2.11).

It remains to consider the cases $x \in A$ and $y \in A^c$. Now $\varphi'(x) = 0$ and the second inequality in (2.11) takes the form $|\varphi'(y)| \leq 8|x-y|/\varepsilon^2$. Write $s = \rho(y)$. Using $|\psi'(t)| \leq 4|t|$, we have

$$|\varphi'(y)| = \varepsilon^{-1} |\psi'(s/\varepsilon)| \leq \frac{4s}{\varepsilon^2} \leq \frac{4|x-y|}{\varepsilon^2}$$

Since $s = \rho(y) \leq |x-y|$. This completes the proof of (2.11) and of the lemma. \square

The next two simple technical lemmas will be often used in the forthcoming proof of Theorem 1.1. To prove the bound $\Delta_n \leq cb_d\beta/\sqrt{n}$ with an inexplicit absolute constant c , all we need is that expectations and integrals in Lemmas 2.4 and 2.5 are bounded from above by $c_0\beta$ (with an absolute constant c_0), which is almost obvious.

Lemma 2.4. Write $\sigma = 2/\sqrt{2\pi}$. We have

$$1 \leq d^{-3/2}\beta, \quad \mathbf{E}|X| \leq \beta, \quad \mathbf{E}|Y| \leq \sigma\beta \leq \beta, \quad \mathbf{E}|Y|^3 \leq 2\sigma\beta \leq 2\beta, \quad (2.13)$$

$$\mathbf{E}|X|^2|\langle X, Y \rangle| \leq \sigma\beta \leq \beta, \quad \mathbf{E}|Y_1|^2|\langle Y_1, Y \rangle| \leq 2\sigma^2\beta \leq 2\beta, \quad (2.14)$$

$$\mathbf{E}|\langle X, Y \rangle| \leq \sigma\beta \leq \beta, \quad \mathbf{E}|\langle Y_1, Y \rangle| \leq \sigma^2\beta \leq \beta,$$

$$\mathbf{E}|\langle X, Y \rangle|^3 \leq 2\sigma\beta \leq 2\beta, \quad \mathbf{E}|\langle Y_1, Y \rangle|^3 \leq 4\sigma^2\beta \leq 3\beta. \quad (2.15)$$

Proof. To prove that $\beta \geq d^3/2$, it suffices to note that $d = \mathbf{E}|X|^2 \leq \beta^{2/3}$. The inequality $\mathbf{E}|X| \leq \beta$ is implied by an application of Hölder inequality and $\beta \geq 1$.

To prove other inequalities, let us consider first the case $d = 1$. Then $\langle X, Y \rangle = XY$, the random variable Y is a standard normal variable, and (2.13)–(2.15) are implied by an application of $\mathbf{E}|Y| = \sigma$ and $\mathbf{E}|Y|^3 = 2\sigma$.

Henceforth we assume that $d \geq 2$. Let η, η_j be independent real standard normal variables. Using that coordinates η_i of $Y = (\eta_1, \dots, \eta_d)$ are independent and identically distributed, $\mathbf{E}\eta^4 = 3$ and $\mathbf{E}\eta^2 = 1$, we have

$$\mathbf{E}|Y|^4 = d\mathbf{E}\eta^4 + (d^2 - d)(\mathbf{E}\eta^2)^2 = d^2 + 2d.$$

Therefore, for $d \geq 2$, we have

$$\mathbf{E}|Y|^3 \leq (\mathbf{E}|Y|^2)^{1/2}(\mathbf{E}|Y|^4)^{1/2} = d^{3/2} \sqrt{1 + 2/d} \leq \sqrt{2}\beta \leq 2\sigma\beta.$$

Using polar coordinates and integrating by parts we have

$$\begin{aligned} \mathbf{E}|Y|^3 &= s_{d-1} \int_0^\infty r^{d+2} \exp\{-r^2/2\} dr = (d+1)s_{d-1} \int_0^\infty r^d \{-r^2/2\} dr \\ &= (d+1)\mathbf{E}|Y|, \end{aligned}$$

where s_{d-1} is the area of the unit sphere $\{x \in \mathbb{R}^d: |x| = 1\}$. Hence, $\mathbf{E}|Y|^3 \geq 2\mathbf{E}|Y|$, for $d \geq 2$. Combining this inequality with the already proved $\mathbf{E}|Y|^3 \leq 2\sigma\beta$, we conclude the proof of (2.13).

Let us prove (2.14) and (2.15). Given X , the random variable $\langle X, Y \rangle$ is a mean zero normal random variable with variance $|X|^2$. Therefore, conditioning on X , we have $\mathbf{E}|X|^2|\langle X, Y \rangle| = \sigma\mathbf{E}|X|^3 = \sigma\beta$. The proof of the second inequality in (2.14) is similar. We have $\mathbf{E}|\langle X, Y \rangle| = \sigma\mathbf{E}|X| \leq \sigma\beta$ and $\mathbf{E}|\langle X, Y \rangle|^3 = 2\sigma\mathbf{E}|X|^3 = 2\sigma\beta$. The proof of the last inequality in (2.15) is similar. \square

Lemma 2.5. *Let*

$$p(y) = (2\pi)^{-d/2} \exp\{-|y|^2/2\}, \quad y \in \mathbb{R}^d, \quad (2.16)$$

be the standard normal d -dimensional density. Then we have

$$\begin{aligned} A &= \mathbf{E} \int_{\mathbb{R}^d} |p'(y)X| dy \leq \sigma\beta \leq \beta, \\ B &= \mathbf{E} \int_{\mathbb{R}^d} |p'(y)Y| dy \leq \sigma^2\beta \leq \beta \end{aligned} \quad (2.17)$$

and

$$\begin{aligned} C &= \mathbf{E} \int_{\mathbb{R}^d} |p'''(y)X^3| dy \leq \zeta\beta \leq 2\beta, \\ D &= \mathbf{E} \int_{\mathbb{R}^d} |p'''(y)Y^3| dy \leq 2\sigma\zeta\beta \leq 3\beta \end{aligned} \quad (2.18)$$

with $\sigma = 2/\sqrt{2\pi}$ and $\zeta = 2(1 + 4e^{-3/2})/\sqrt{2\pi}$.

Proof. We have

$$p'(x)h = -\langle x, h \rangle p(x), \quad p'''(x)h^3 = (3|h|^2\langle x, h \rangle - \langle x, h \rangle^3)p(x). \quad (2.19)$$

Let us prove (2.17). Using (2.19) we can write $A = \mathbf{E}|\langle Y, X \rangle|$. Applying (2.15) we get $A \leq \sigma\beta$. Similarly $B \leq \sigma^2\beta$.

To prove (2.18) we note that

$$\mathbf{E}|3|h|^2\langle Y, h \rangle - \langle Y, h \rangle^3| = |h|^3\mathbf{E}|3\eta - \eta^3| = 2(1 + 4e^{-3/2})/\sqrt{2\pi},$$

where η is a standard real normal random variable. Applying (2.13), we derive (2.18). \square

For a sufficiently smooth function $f: \mathbb{R}^d \rightarrow \mathbb{R}$, we shall apply the Taylor formula

$$f(x+h) = f(x) + f'(x)h + \cdots + \frac{1}{s!} f^{(s)}(x)h^s + \frac{1}{s!} \mathbf{E}(1-\tau)^s f^{(s+1)}(x+\tau h)h^{s+1}, \quad (2.20)$$

where τ is a random variable uniformly distributed on the interval $[0,1]$.

In the next Lemma 2.6, we provide Nazarov's (2001a, b) lower bound for the isoperimetric constant $a_d(\mathcal{C})$. The proof differs in technical details from the original proof of Nazarov. However, the main ingredient, namely, a construction of *random* convex sets instead of *deterministic* ones belongs to Nazarov. Whether it is possible to provide an explicit deterministically given convex set remains an open question.

Lemma 2.6. *Let $a_d = a_d(\mathcal{C})$ be the isoperimetric constants defined by (1.2) for the class \mathcal{C} convex subsets of \mathbb{R}^d . Then we have*

$$\liminf_{d \rightarrow \infty} a_d/d^{1/4} > 0. \quad (2.21)$$

Nazarov (2001a, b) proves more than (2.21). He shows that

$$0.28 < \liminf_{d \rightarrow \infty} a_d/d^{1/4} \quad \text{and} \quad \limsup_{d \rightarrow \infty} a_d/d^{1/4} < 0.64. \quad (2.22)$$

Moreover, if the standard normal distribution ν is replaced by a normal distribution ν_A with the covariance matrix A , then there exist absolute positive constants c_1 and c_2 such that

$$c_1 S \leq a_d \leq c_2 S$$

where

$$S^4 = \sigma_1^{-2} + \cdots + \sigma_d^{-2}$$

and where σ_k^2 are eigenvalues of the covariance matrix A .

Let us describe the sets which realize the lower bound (2.21). Let v_1, \dots, v_{2^d} be vertexes of the unit cube $[-1, 1]^d$. Then one can represent the ball

$$B_1(r) = \{x \in \mathbb{R}^d : |x_1| + \cdots + |x_d| \leq r\} = \bigcap_{k=1}^{2^d} H_k(r) \quad (2.23)$$

of the space \mathbb{R}^d as an intersection of 2^d half-spaces

$$H_k(r) = \{x \in \mathbb{R}^d : \langle x, v_k \rangle \leq r\}.$$

It is well known that the class \mathcal{B}_1 of balls of type (2.23) satisfies $a_d(\mathcal{B}_1) \leq c$ with some absolute constant c , for all d . Nevertheless, it serves as a basis of the construction. Let k_1, k_2, \dots be i.i.d. random variables taking values $1, 2, 3, \dots, 2^d$ with equal probability 2^{-d} . A set, say A , which realizes the lower bound (2.21) is defined as

$$A = H_{k_1}(r) \cap \cdots \cap H_{k_N}(r) \quad \text{with } r = d^{3/4}$$

and an integer N such that $1 \leq Nd^{-1/4} \exp\{-d^{1/2}/2\} \leq 2$. Loosely speaking, A is just “a very much unfinished” ball $B_1(d^{3/4})$ constructed using intersections of a few (that is,

of N , which is much smaller than the needed total amount 2^d for $B_1(d^{3/4})$) randomly chosen half-spaces $H_k(d^{3/4})$. Note, that the largest centered Euclidean ball inscribed in A has radius $d^{1/4}$.

Proof of Lemma 2.6. In the proof we assume that d is sufficiently large such that all quantities below are well defined.

Let Z, Z_1, Z_2, \dots be independent identically distributed random vectors with values in \mathbb{R}^d such that $\sqrt{d}Z = (\varepsilon_1, \dots, \varepsilon_d)$, where ε_k are i.i.d. Rademacher random variables such that they assume values 1 and -1 with equal probability $\frac{1}{2}$. Note that $|Z| = 1$ with probability 1. Write $\rho = d^{1/4}$ and let an integer N satisfy

$$\rho \exp\{\rho^2/2\} \leq N \leq 2\rho \exp\{\rho^2/2\}. \quad (2.24)$$

Introduce the set

$$A(\rho) = \bigcap_{k=1}^N \{x \in \mathbb{R}^d : \langle x, Z_k \rangle \leq \rho\}.$$

In the notation we suppress the dependence of the random set A on Z_1, \dots, Z_N and d . Write

$$b = \mathbb{P}\{\langle x, Z \rangle \leq \rho\}, \quad a = \mathbb{P}\{\langle x, Z \rangle \leq \rho + \delta\}, \quad \delta > 0.$$

Using the Fubini theorem and the i.i.d. assumption imposed on Z, Z_1, \dots, Z_N , we get

$$\begin{aligned} \mathbb{E}v(A(\rho)) &= \mathbb{E} \int_{\mathbb{R}^d} \mathbb{I}\{x \in A(\rho)\} v(dx) = \int_{\mathbb{R}^d} \mathbb{E} \mathbb{I}\{x \in A(\rho)\} v(dx) \\ &= \int_{\mathbb{R}^d} b^N v(dx). \end{aligned} \quad (2.25)$$

Taking the expectation of both sides of the inequality $a_d \delta \geq v(A(\rho + \delta)) - v(A(\rho))$ (the inequality clearly holds since we assume that $|\mathbb{Z}| = 1$), Using the obvious inequality

$$a^N - b^N = (a - b) \sum_{k=1}^N a^{N-k} b^{k-1} \geq N(a - b)b^{N-1} \geq N(a - b)b^N$$

for $1 \geq a \geq b \geq 0$, and applying (2.25), we obtain

$$a_d \delta \geq N \int_{\mathbb{R}^d} b^N \mathbb{P}\{\rho < \langle x, Z \rangle \leq \rho + \delta\} v(dx). \quad (2.26)$$

To estimate b^N we use the following inequality (see Pinelis, 1998):

$$\mathbb{P}\{x_1 \varepsilon_1 + \dots + x_d \varepsilon_d \geq t\} \leq c_1 I(t/|x|), \quad I(t) = 1 - \Phi(t),$$

where Φ is the standard normal distribution function. Here and below by c_k we denote absolute constants. Using in addition $I(t) \leq \varphi(t)/t$, for $t > 0$, where φ is the standard normal density, and $1 - u \geq \exp\{-2u\}$, for $0 \leq u \leq 1/2$, we have

$$\begin{aligned} b^N &= (1 - \mathbb{P}\{\langle x, Z \rangle > \rho\})^N \geq \exp\{-c_2 N I(\rho \sqrt{d}/|x|)\} \geq \exp\{-c_2 N I(\rho)\} \\ &\geq \exp\{-c_3 N \exp\{-\rho^2/2\}/\rho\} \quad \text{for } |x| \leq \sqrt{d}. \end{aligned} \quad (2.27)$$

In view of definition (2.24) of N , bound (2.27) yields $b^N > c_4 > 0$, for $|x| \leq \sqrt{d}$. Hence, (2.26) implies

$$a_d \delta \geq c_4 N J, \quad J \stackrel{\text{def}}{=} \int_{|x| \leq \sqrt{d}} \mathbb{P}\{\rho < \langle x, Z \rangle \leq \rho + \delta\} v(dx). \quad (2.28)$$

Fubini's theorem combined with $|\mathbb{Z}| = 1$ and the rotational symmetry of v yield

$$\begin{aligned} J &= \mathbf{E} \int_{|x| \leq \sqrt{d}} \mathbb{I}\{\rho < \langle x, Z \rangle \leq \rho + \delta\} v(dx) \\ &= \int_{|x| \leq \sqrt{d}} \mathbb{I}\{\rho < x_1 \leq \rho + \delta\} v(dx). \end{aligned} \quad (2.29)$$

By (2.28) we have $a_d \geq c_4 N J / \delta$. Using (2.29), passing to the limit as $\delta \downarrow 0$, and noting that $N \exp\{-\rho^2/2\} \geq \rho = d^{1/4}$ (see (2.24)), we derive

$$\begin{aligned} a_d &\geq c_5 N \exp\{-\rho^2/2\} \int_{\rho^2 + x_2^2 + \dots + x_d^2 \leq d} p(0, x_2, \dots, x_d) dx_2 \dots dx_d \\ &\geq c_5 d^{1/4} \mathbb{P}\{\eta_2^2 + \dots + \eta_d^2 \leq d - \sqrt{d}\}, \end{aligned} \quad (2.30)$$

where η_2, \dots, η_d are independent identically distributed real standard normal random variables and p is the density of v . Write $T_d = (\eta_2^2 - 1 + \dots + \eta_d^2 - 1)/\sqrt{d}$. The Central Limit Theorem applied to the probability in (2.30) yields

$$\liminf_{d \rightarrow \infty} a_d / d^{1/4} \geq c_5 \lim_{d \rightarrow \infty} \mathbb{P}\{T_d \leq d^{-1/2} - 1\} = c_6 > 0,$$

which proves (2.21) and the lemma. \square

3. Proof of Theorem 1.1

The proof is based on induction in n . We shall prove that

$$\Delta_n \leq M b_d \delta, \quad \delta \stackrel{\text{def}}{=} \beta / \sqrt{n} \quad (3.1)$$

holds for all n provided that an absolute constant $M \geq 10$ is sufficiently large (recall that $b_d = \max\{1, a_d\}$).

Since $\beta \geq 1$ and $\Delta_n \leq 1$, for $n \leq M^2 \beta^2$ we have

$$\Delta_n \leq 1 \leq b_d \sqrt{M^2 \beta^2 / n} = M b_d \delta. \quad (3.2)$$

Hence, (3.1) holds for $n \leq 100 \leq M^2 \beta^2$. In particular, (3.1) holds for $n = 1$.

Now assume that (3.1) holds for $1, \dots, n-1$. We shall prove that (3.1) holds for n as well. In the proof we can assume that $n \geq M^2 \beta^2 \geq 100$, cf. (3.2).

Let $\varepsilon > 0$ be a positive number (later we shall choose $\varepsilon = a\delta$ with a sufficiently large absolute constant $a > 0$). Applying the smoothing Lemma 2.1, condition (1.2) and using $a_d \leq b_d$, we have

$$\Delta_n \leq \sup_{A \in \mathcal{A}} |\mathbf{E} \varphi(S_n) - \mathbf{E} \varphi(Y)| + b_d \varepsilon, \quad (3.3)$$

where a function $\varphi = \varphi_{\varepsilon, A}$ satisfies the conditions of Lemma 2.1. Fix $A \in \mathcal{A}$. By Lemma 2.3 we can choose $\varphi(x) = \psi(\rho(x)/\varepsilon)$, where $\rho(x) = \rho_A(x)$ is the distance between x and A , and φ and ψ satisfy the conditions of Lemma 2.3.

To simplify the notation, we write

$$U = X/\sqrt{n}, \quad U_k = X_k/\sqrt{n}, \quad V = Y/\sqrt{n}, \quad V_k = Y_k/\sqrt{n}.$$

Write

$$W_k = V_1 + \cdots + V_{k-1} + U_{k+1} + \cdots + U_n, \quad Z_n = V_1 + \cdots + V_n. \quad (3.4)$$

Due to the i.i.d. assumption, S_n is distributed as $W_1 + U$. Moreover, $Z_n, W_n + V$ and Y are i.i.d. standard normal random vectors. We have

$$\begin{aligned} |\mathbf{E}\varphi(S_n) - \mathbf{E}\varphi(Y)| &= |\mathbf{E}\varphi(W_1 + U) - \mathbf{E}\varphi(W_n + V)| \\ &\leq |\mathbf{E}\varphi(W_1 + U) - \mathbf{E}\varphi(W_1 + V)| + |\mathbf{E}\varphi(W_1 + V) - \mathbf{E}\varphi(W_n + V)| \\ &= |\mathbf{E}\varphi(W_1 + U) - \mathbf{E}\varphi(W_1 + V)| + |\mathbf{E}\varphi(W_2 + U) - \mathbf{E}\varphi(W_n + V)| \end{aligned} \quad (3.5)$$

since by the i.i.d. assumption $\mathbf{E}\varphi(W_1 + V) = \mathbf{E}\varphi(W_2 + U)$. Repetitions of arguments used to obtain (3.5), lead to

$$|\mathbf{E}\varphi(S_n) - \mathbf{E}\varphi(Y)| \leq \gamma_1 + \cdots + \gamma_n, \quad (3.6)$$

where

$$\gamma_k = |\mathbf{E}\varphi(W_k + U) - \mathbf{E}\varphi(W_k + V)|.$$

Introduce

$$\theta^2 = n/(k-1), \quad \alpha^2 = n/(n-k), \quad Z = U_{k+1} + \cdots + U_n. \quad (3.7)$$

Then, for $2 \leq k \leq n$, we can write

$$\gamma_k = |\mathbf{E}\varphi(Y_1/\theta + U + Z) - \mathbf{E}\varphi(Y_1/\theta + V + Z)|. \quad (3.8)$$

Write $n_0 = [n/2]$, where $[z]$ stands for the integer part of z . Below we shall prove that, for any $m = 2, \dots, n_0$,

$$\gamma_1 \leq b_d(c_1\delta + c_2M\delta^2/\varepsilon), \quad (3.9)$$

$$\gamma_2 + \cdots + \gamma_m \leq \frac{\beta b_d \sqrt{m-1}}{\varepsilon^2 n} (c_3 M \delta + c_4 \varepsilon), \quad (3.10)$$

$$\gamma_{m+1} + \cdots + \gamma_{n_0} \leq \frac{c_5 \beta b_d M \delta}{\sqrt{m-1}} + c_6 b_d \delta, \quad (3.11)$$

$$\gamma_{n_0+1} + \cdots + \gamma_n \leq c_7 b_d \delta \quad (3.12)$$

with some absolute constants c_1, \dots, c_7 . These constants can be chosen to be

$$\begin{aligned} c_1 &= (2 + 2\sigma)\sqrt{100/99}, & c_2 &= 2c_1, & c_3 &= 2c_4, & c_4 &= 16\sqrt{2}(\sigma + 2\sigma^2)/3, \\ c_5 &= \zeta(1 + 2\sigma)\sqrt{2/3}, & c_6 &= c_5\sqrt{100/98}, & c_7 &= c_5\sqrt{100/97} \end{aligned} \quad (3.13)$$

with $\sigma = 2/\sqrt{2\pi}$ and $\zeta = 2(1 + 4e^{-3/2})/\sqrt{2\pi}$.

Bounds (3.9)–(3.12) imply the desired bound for Δ_n . Indeed, summing the inequalities (3.9)–(3.12), using (3.3) and setting $\varepsilon = a\delta$ with some $1 \leq a \leq M$ to be chosen

later, we have (below $c_8 = c_8(c_1, \dots, c_7)$ and $c_9 = c_9(c_8)$ are sufficiently large absolute constants)

$$\begin{aligned} \Delta_n &\leq b_d \delta \left(a + c_8 + c_8 \frac{\sqrt{m-1}}{a\beta} + c_8 M \left(\frac{1}{a} + \frac{\sqrt{m-1}}{a^2\beta} + \frac{\beta}{\sqrt{m-1}} \right) \right) \\ &\leq b_d \delta \left(a + c_9 M \left(\frac{1}{a} + \frac{\sqrt{m-1}}{a^2\beta} + \frac{\beta}{\sqrt{m-1}} \right) \right) \end{aligned}$$

using $M/a \geq 1$. Therefore, the inequality $\Delta_n \leq Mb_d \delta$ reduces to checking that

$$a + c_9 M \left(\frac{1}{a} + \frac{\sqrt{m-1}}{a^2\beta} + \frac{\beta}{\sqrt{m-1}} \right) \leq M. \quad (3.14)$$

We choose $m-1$ such that $a^2\beta^2 \leq m-1 \leq 2a^2\beta^2$. Then (3.14) is implied by

$$a + c_{10}M/a \leq M, \quad (3.15)$$

where $c_{10} \geq 1$ is an absolute constant. Let us take $a = 2c_{10}$ and $M = 10c_{10}$. Then (3.15) holds. Note, that we have to check the condition $1 \leq m \leq n/2$. Since $m-1 \leq 2a^2\beta^2$, it suffices to check that $3a^2\beta^2 \leq n/4$ or, equivalently, that $12c_{10}^2\beta^2 \leq n$. This inequality holds since we assume $n \geq M^2\beta^2 = 100c_{10}^2\beta^2$. A little more careful choice of the parameters using numerical values (3.13) of constants shows that it suffices to take $M \leq 100$, which proves theorem.

It remains to prove (3.9)–(3.12). We shall prove (3.9)–(3.12) with

$$c_1 \leq 8, \quad c_2 \leq 16, \quad c_3 \leq 192, \quad c_4 \leq 48, \quad c_5 \leq 20, \quad c_6 \leq 10, \quad c_7 \leq 20.$$

Better constants (3.13) can be obtained using refined versions of inequalities of Lemmas 2.4 and 2.5, applying $n \geq 100$ to estimate better θ and α , and using $E\tau = 1/2$ and $E\tau^2 = 1/3$. In order to keep the notation as simple as possible, we omit related details.

Let us prove (3.9). A little later we prove that

$$\gamma_1 \leq \frac{4\beta}{\varepsilon\sqrt{n}} (2\Delta_{n-1} + \alpha b_d \varepsilon), \quad \alpha^2 = n/(n-1). \quad (3.16)$$

Using the induction assumption $\Delta_{n-1} \leq Mb_d\beta/\sqrt{(n-1)} = Mb_d\delta\alpha$ and estimating $\alpha \leq 2$, for $n \geq 2$, inequality (3.16) yields (3.9).

Let us prove inequality (3.16). Using the Taylor formula (2.20) with $s = 0$, noting that $\varphi'(x) = \varphi'(x)\mathbb{I}\{A^\varepsilon \setminus A\}$ since φ is a constant function outside the set $A^\varepsilon \setminus A^{-\varepsilon}$, and estimating the derivative φ' by (2.11) of Lemma 2.3, we have

$$\gamma_1 = |\mathbf{E}\varphi'(W_1 + \tau U)U - \mathbf{E}\varphi'(W_1 + \tau V)V| \leq 2\mathbf{E}I(U)/\varepsilon + 2\mathbf{E}I(V)/\varepsilon \quad (3.17)$$

with

$$I(U) = |U|\mathbb{I}\{W_1 + \tau U \in A^\varepsilon \setminus A\}, \quad I(V) = |V|\mathbb{I}\{W_1 + \tau V \in A^\varepsilon \setminus A\}.$$

We shall prove that

$$\mathbf{E}I(U) \leq \mathbf{E}|U|(2\Delta_{n-1} + \alpha b_d \varepsilon) \quad \mathbf{E}I(V) \leq \mathbf{E}|V|(2\Delta_{n-1} + \alpha b_d \varepsilon). \quad (3.18)$$

By (2.13) of Lemma 2.4, we have $\mathbf{E}|V|, \mathbf{E}|U| \leq \beta/\sqrt{n}$, and (3.17)–(3.18) imply (3.16).

Let us prove (3.18). We shall prove only the bound for $\mathbf{E}I(U)$ since a proof of the bound for $\mathbf{E}I(V)$ is similar. Conditioning on τ and U , we have

$$\begin{aligned}\mathbf{E}I(U) &= \mathbf{E}|U| \mathbb{P}\{W_1 \in A^\varepsilon \setminus A - \tau U | \tau, U\} \leq \mathbf{E}|U| \sup_{z \in \mathbb{R}^d} \mathbb{P}\{W_1 \in A^\varepsilon \setminus A + z\} \\ &\leq \mathbf{E}|U| \sup_{A \in \mathcal{A}} \mathbb{P}\{W_1 \in A^\varepsilon \setminus A\} = \mathbf{E}|U| \sup_{A \in \mathcal{A}} \mathbb{P}\{S_{n-1}/\alpha \in A^\varepsilon \setminus A\} \\ &\leq \mathbf{E}|U| \sup_{A \in \mathcal{A}} \mathbb{P}\{S_{n-1} \in A^{\alpha\varepsilon} \setminus A\}.\end{aligned}\quad (3.19)$$

We estimate $\mathbb{P}\{S_{n-1} \in A^{\alpha\varepsilon} \setminus A\} = \mathbb{P}\{S_{n-1} \in A^{\alpha\varepsilon}\} - \mathbb{P}\{S_{n-1} \in A\}$ applying the induction assumption. Using in addition condition (1.2) to estimate $\mathbb{P}\{Y \in A^{\alpha\varepsilon} \setminus A\}$, we get

$$\mathbb{P}\{S_{n-1} \in A^{\alpha\varepsilon} \setminus A\} \leq 2\Delta_{n-1} + \mathbb{P}\{Y \in A^{\alpha\varepsilon} \setminus A\} \leq 2\Delta_{n-1} + \alpha b_d \varepsilon. \quad (3.20)$$

Combining (3.19) and (3.20), we obtain (3.18), which concludes the proof of (3.16) and of (3.19).

Let us prove (3.10). We shall show that (now $\alpha^2 = n/(n-k)$)

$$\gamma_k \leq \frac{24\beta}{\varepsilon^2 n \sqrt{k-1}} (2\Delta_{n-k} + b_d \alpha \varepsilon) \quad \text{for } k = 2, \dots, n-1. \quad (3.21)$$

This inequality implies (3.10). Indeed, using the induction assumption we can estimate $\Delta_{n-k} \leq Mb_d \alpha \varepsilon$. Furthermore, $\alpha \leq 2$ since $1 \leq k \leq n_0 = [n/2]$, and

$$\sum_{k=2}^m \frac{1}{\sqrt{k-1}} \leq 1 + \int_1^m \frac{dt}{\sqrt{t}} = 2\sqrt{m} - 1 \leq 2\sqrt{m-1} \quad \text{for } m \geq 2.$$

We note that the Lipschitz condition (2.11) implies

$$|\varphi'(x) - \varphi'(y)| \leq \frac{8|x-y|}{\varepsilon^2} (\mathbb{I}\{x \in A^\varepsilon \setminus A\} + \mathbb{I}\{y \in A^\varepsilon \setminus A\}). \quad (3.22)$$

Indeed, if at least one of x and y belong to $A^\varepsilon \setminus A$, then $\mathbb{I}\{x \in A^\varepsilon \setminus A\} + \mathbb{I}\{y \in A^\varepsilon \setminus A\} \geq 1$, and (3.22) becomes (2.11). If neither x nor y belongs to $A^\varepsilon \setminus A$, then $\varphi'(x) = \varphi'(y) = 0$ since φ is a constant function outside the set $A^\varepsilon \setminus A$, and (3.22) is just the identity $0 = 0$.

Let us return to the proof of (3.21). Using notations (3.7)–(3.8), we can write

$$\gamma_k = |I - J| \quad I \stackrel{\text{def}}{=} \mathbf{E}\varphi(Y_1/\theta + U + Z), \quad J \stackrel{\text{def}}{=} \mathbf{E}\varphi(Y_1/\theta + V + Z).$$

Let us show that

$$I = I_0 + I_1, \quad J = I_0 + J_1 \quad (3.23)$$

with $I_0 = \mathbf{E}\varphi(Y_1/\theta + Z)$ and

$$\begin{aligned}I_1 &= \theta \mathbf{E}\tau \langle Y_1, U \rangle \varphi'(Y_1/\theta + \tau_1 \tau U + Z) U, \\ J_1 &= \theta \mathbf{E}\tau \langle Y_1, V \rangle \varphi'(Y_1/\theta + \tau_1 \tau V + Z) V,\end{aligned}\quad (3.24)$$

where τ and τ_1 are independent random variables uniformly distributed in the interval $[0, 1]$. We have

$$I = \mathbf{E} \int_{\mathbb{R}^d} \varphi(x/\theta + U + Z) p(x) dx, \quad (3.25)$$

where p is standard normal density (2.16). Changing the variable x as $x/\theta + U = y$, we can rewrite (3.25) as

$$I = \mathbf{E} \int_{\mathbb{R}^d} \varphi(y + Z) p(\theta y - \theta U) \theta^d dy. \quad (3.26)$$

Given U , expanding the density

$$\begin{aligned} p(\theta y - \theta U) &= p(\theta y) - \theta \mathbf{E} p'(\theta y - \tau \theta U) U \\ &= p(\theta y) + \theta \mathbf{E} \langle \theta y - \tau \theta U, U \rangle p'(\theta y - \tau \theta U) \end{aligned}$$

(cf. (2.19)) and changing the variable y in integrals as $\theta y = x$ and $\theta y - \tau \theta U = x$, the relation (3.25) yields

$$I = I_0 + I_1, \quad I_1 \stackrel{\text{def}}{=} \theta \mathbf{E} \langle Y_1, U \rangle \varphi(Y_1/\theta + \tau U + Z). \quad (3.27)$$

Expanding the function φ by (2.20) with $s = 0$ and using $\mathbf{E}U = 0$, we have

$$I_1 = \theta \mathbf{E} \tau \langle Y_1, U \rangle \varphi'(Y_1/\theta + \tau_1 \tau U + Z) U. \quad (3.28)$$

Relation (3.25)–(3.28) yield (3.23) for I . The proof for J is similar, we have just to replace everywhere U by V .

Write

$$I_2 = \theta \mathbf{E} \tau \langle Y_1, U \rangle \varphi'(Y_1/\theta + Z) U, \quad J_2 = \theta \mathbf{E} \tau \langle Y_1, V \rangle \varphi'(Y_1/\theta + Z) V$$

and notice that $I_2 = J_2$ since the covariances of U and V are equal. Therefore (3.23) implies

$$\gamma_k = |I - J| = |I_1 - J_1| \leq |I_1 - I_2| + |J_1 - J_2|. \quad (3.29)$$

Applying to the function φ' the Lipschitz condition (3.22), writing for a while

$$\xi = \frac{8\theta}{\varepsilon^2} \tau_1 \tau^2 |U|^2 \langle Y_1, U \rangle$$

and \mathbb{P}_* for the conditional probability given all random variables except Z , we have

$$\begin{aligned} |I_1 - I_2| &\leq \mathbf{E} \xi (\mathbb{P}_* \{Y_1/\theta + \tau_1 \tau U + Z \in A^c \setminus A\} + \mathbb{P}_* \{Y_1/\theta + Z \in A^c \setminus A\}) \\ &\leq 2 \mathbf{E} \xi \sup_{x \in \mathbb{R}^d} \mathbb{P} \{x + Z \in A^c \setminus A\}. \end{aligned} \quad (3.30)$$

Note that Z is a normalized sum of $n - k$ independent copies of X . Therefore, using the induction assumption and arguing similarly to (3.19)–(3.30), we get

$$\mathbb{P} \{x + Z \in A^c \setminus A\} \leq 2\Delta_{n-k} + b_d \alpha \varepsilon. \quad (3.31)$$

In order to estimate $\mathbf{E} \xi$ we notice that $\tau, \tau_1 \leq 1$ and $\mathbf{E} |U|^2 \langle Y_1, U \rangle \leq \beta n^{-3/2}$ (cf. (2.14) of Lemma 2.4). We get

$$\mathbf{E} \xi \leq 8\theta \beta / (\varepsilon^2 n^{3/2}). \quad (3.32)$$

Collecting bounds (3.30)–(3.32), we obtain

$$|I_1 - I_2| \leq 8\theta \beta (2\Delta_{n-k} + b_d \alpha \varepsilon) / (\varepsilon^2 n^{3/2}). \quad (3.33)$$

Similarly to the proof of (3.33) we have

$$|J_1 - J_2| \leq 16\theta \beta (2\Delta_{n-k} + b_d \alpha \varepsilon) / (\varepsilon^2 n^{3/2}). \quad (3.34)$$

Now (3.29) combined with (3.33) and (3.34) yields (3.21) since $\theta = \sqrt{n/(k-1)}$.

Let us prove (3.11) and (3.12). Below we show that

$$\gamma_k \leq 5\beta \Delta_{n-k}(k-1)^{-3/2} + 5\beta(n-k)^{-3/2} \quad \text{for } k = 2, \dots, n. \quad (3.35)$$

$$\gamma_k \leq 5\beta(k-1)^{-3/2} \quad \text{for } k = 2, \dots, n. \quad (3.36)$$

These bounds imply (3.11) and (3.12). To prove (3.11) we use (3.35) and the induction assumption $\Delta_{n-k} \leq 2b_d M\delta$, for $k \leq n_0$. Using

$$\sum_{k=m+1}^{\infty} \frac{1}{(k-1)^{3/2}} \leq \int_{m-1}^{\infty} \frac{dt}{t^{3/2}} = \frac{2}{(m-1)^{1/2}},$$

estimating $n-k \geq n/2$, for $k \leq n_0$, we have

$$\gamma_{m+1} + \dots + \gamma_{n_0} \leq \frac{20b_d \beta M\delta}{\sqrt{m-1}} + \frac{10\beta}{\sqrt{n}}$$

which proves (3.11) since $b_d \geq 1$. To prove (3.12), we use (3.36). Summing the bound (3.36) over $k > n_0 = [n/2]$ and using

$$\sum_{k=n_0+1}^{\infty} \frac{1}{(k-1)^{3/2}} \leq \int_{n_0-1}^{\infty} \frac{dt}{t^{3/2}} = \frac{2}{\sqrt{n_0-1}} \leq \frac{4}{\sqrt{n}},$$

we obtain (3.12).

Let us prove (3.35) and (3.36). We have $\varphi(x) = \psi(\rho_A(x)/\varepsilon)$ and $\int_{\mathbb{R}} |\psi'(t)| dt = 1$ (cf. Lemma 2.3). Integrating by parts and using property (iii) of the class \mathcal{A} , we obtain

$$\begin{aligned} \gamma_k &= \left| \int_{\mathbb{R}} \psi(t/\varepsilon) d(\mathbb{P}\{\rho_A(W_k + U) \leq t\} - \mathbb{P}\{\rho_A(W_k + V) \leq t\}) \right| \\ &\leq \int_{\mathbb{R}} |\psi'(t/\varepsilon)| |\mathbb{P}\{\rho_A(W_k + U) \leq t\} - \mathbb{P}\{\rho_A(W_k + V) \leq t\}| dt/\varepsilon \\ &\leq \sup_{B \in \mathcal{A}} \gamma_k(B) \end{aligned}$$

with

$$\gamma_k(B) = |P\{W_k + U \in B\} - \mathbb{P}\{W_k + V \in B\}|.$$

Hence, it suffices to prove that instead of γ_k the quantity $\gamma_k(B)$ is bounded as γ_k in (3.35) and (3.36). Using notation (3.7)–(3.8), we have

$$\gamma_k(B) = |\mathbb{P}\{Y/\theta + U + Z \in B\} - \mathbb{P}\{Y/\theta + V + Z \in B\}|.$$

Changing the variable y as $y/\theta + U = u$, we get

$$\begin{aligned} \gamma_k(B) &= \left| \mathbf{E} \int_{\mathbb{R}^d} \mathbb{I}\{y/\theta + U + Z \in B\} p(y) dy \right. \\ &\quad \left. - \mathbf{E} \int_{\mathbb{R}^d} \mathbb{I}\{\theta y + V + Z \in B\} p(y) dy \right| = |I_1 - J_1|, \end{aligned}$$

where

$$I_1 = \mathbf{E} \int_{\mathbb{R}^d} \mathbb{I}\{u + Z \in B\} p(\theta u - \theta U) \theta^d du,$$

and J_1 is defined as I_1 just replacing U by V .

Applying the Taylor expansion (2.20) with $s = 2$ to the density p with $h = \theta U$ and $h = \theta V$, using that U and V have the same means and covariances, changing the variables $\theta u - \tau\theta U = y$ and $\theta u - \tau\theta V = y$, we obtain

$$\gamma_k(B) \leq |I_0| + |J_0|$$

with

$$I_0 = \frac{\theta^3}{2} \mathbf{E}(1 - \tau)^2 \int_{\mathbb{R}^d} \mathbb{I}\{y/\theta + \tau U + Z \in B\} p'''(y) U^3 \, dy \quad (3.37)$$

and J_0 defined as I_0 just replacing U by V .

Let us prove (3.36). Using (3.37) and (2.18) of Lemma 2.4, we have

$$|I_0| \leq \theta^3 \mathbf{E} \int_{\mathbb{R}^d} |p'''(y) U^3| \, dy \leq \frac{2\beta\theta^3}{n^{3/2}} = \frac{2\beta}{(k-1)^{3/2}}.$$

Similarly $|J_0| \leq 3\beta/(k-1)^{3/2}$. These bounds and $\gamma_k(B) \leq |I_0| + |J_0|$ prove (3.36).

Let us return to the proof of (3.35). Write $\alpha^2 = n/(n-k)$. The sum Z is a normalized sum of $n-k$ independent copies of X . Conditioning in (3.37) on all random variables except Z and using the induction assumption, we get

$$|I_0| \leq \Delta_{n-k} I_2 + |I_3|, \quad (3.38)$$

where

$$I_2 = \frac{\theta^3}{6} \mathbf{E} \int_{\mathbb{R}^d} |p'''(y) U^3| \, dy, \quad I_3 = \frac{\theta^3}{2} \mathbf{E}(1 - \tau)^2 I_4, \\ I_4 \stackrel{\text{def}}{=} \int_{\mathbb{R}^d} \mathbb{I}\{y/\theta + \tau U + Y_1/\alpha \in B\} p'''(y) U^3 \, dy.$$

We can represent I_4 as

$$I_4 = \theta^3 \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \mathbb{I}\{x + \tau U + z \in B\} p'''(\theta x) U^3 p(\alpha z) \theta^d \, dx \alpha^d \, dz.$$

To estimate I_4 , we integrate by parts. Changing the variable x as $x + z = y$ and writing $\partial_h[z]$ for the partial derivative in direction h with respect to the variable z , we have

$$I_4 = - \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \mathbb{I}\{y + \tau U \in B\} (\partial_U^3[z] p(\theta y - \theta z)) p(\alpha z) \theta^d \, dy \alpha^d \, dz \\ = \alpha^3 \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \mathbb{I}\{y + \tau U \in B\} p(\theta y - \theta z) p'''(\alpha z) U^3 \theta^d \, dy \alpha^d \, dz.$$

Therefore

$$|I_4| \leq \alpha^3 \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} p(\theta y - \theta z) |p'''(\alpha z) U^3| \theta^d \, dy \alpha^d \, dz \\ = \alpha^3 \int_{\mathbb{R}^d} |p'''(z) U^3| \, dz. \quad (3.39)$$

Estimating the last integral in (3.39) by (2.18) of Lemma 2.5 and collecting the bounds in (3.38), we get

$$|I_0| \leq 2\beta(\Delta_{n-k} \theta^3 n^{-3/2} + \alpha^3 n^{-3/2}) = 2\beta(\Delta_{n-k}(k-1)^{-3/2} + (n-k)^{-3/2}). \quad (3.40)$$

A similar proof shows that $|J_0|$ is bounded as I_0 in (3.40) but with the factor 3 instead of 2. Therefore the inequality $\gamma_k(B) \leq |I_0| + |J_0|$ yields (3.35). \square

Acknowledgements

Our attention to the topic, as mentioned in Theorem 1.1 and the text below it, was drawn by R. Bhattacharya. We would like to thank him for the support and numerous helpful discussions.

References

- Ball, K., 1993. The reverse isoperimetric problem for Gaussian measure. *Discrete Comput. Geom.* 10 (4), 411–420.
- Bentkus, V., 1986a. Dependence of the Berry–Esseen bound on the dimension. *Lithuanian Math. J.* 26 (2), 205–210.
- Bentkus, V., 1986b. Large deviations in Banach spaces. *Probab. Theory Appl.* 31 (4), 710–716.
- Bhattacharya, R.N., Rao, R.R., 1986. *Normal Approximation and Asymptotic Expansions*. Krieger, Melbourne.
- Götze, F., 1991. On the rate of convergence in the multivariate CLT. *Ann. Probab.* 19 (2), 724–739.
- Nagaev, S.V., 1976. An estimate of the remainder term in the multidimensional CLT. *Proceeding of the Third Japan–USSR Symposium on Probability Theory, Lecture Notes in Mathematics*, Springer, Berlin, pp. 419–438.
- Nazarov, F., 2001a. On the asymptotic behavior of the maximal perimeter of convex set in \mathbb{R}^n with respect to the Gaussian measure. A manuscript, pp. 1–5.
- Nazarov, F., 2001b. On the maximal perimeter of a convex set in \mathbb{R}^n with respect to the Gaussian measure II (anisotropic case). A manuscript, pp. 1–11.
- Paulauskas, V., Račkauskas, A., 1989. *Approximation Theory in the CLT. Exact Results in Banach Spaces*. Kluwer, Dordrecht.
- Pinelis, I., 1998. Optimal tail comparison based on comparison of moments. *High Dimensional Probability, Progress in Probability*, Vol. 43, Oberwolfach, 1996, Birkhauser, Basel, pp. 297–314.
- Sazonov, V., 1981. *Normal Approximations: Some Recent Advances*. Springer, Berlin, New York.
- Senatov, V., 1980. Several estimates of the rate of convergence in the multidimensional CLT. *Dokl. Akad. Nauk SSSR* 254, 809–812.
- Senatov, V., 1984. An estimate of the Levy–Prokhorov metric. *Theory Probab. Appl.* 29 (1), 109–114.
- Senatov, V., 1998. *Normal Approximations: New Results, Methods and Problems*. VSP, Utrecht.