

## CIE 6020 Assignment 2

WU, Chenhao 117010285

February 18, 2019

1. Let  $X, Y, Z$  be three random variables with a joint probability mass function  $p(x, y, z)$ . The relative entropy between the joint distribution and the product of the marginal is

$$D(p(x, y, z) || p(x)p(y)p(z)) = E\left[\log \frac{p(x, y, z)}{p(x)p(y)p(z)}\right]$$

Expand this in terms of entropies. When is this quantity zero?

**Answer:**

$$\begin{aligned} E\left[\log \frac{p(x, y, z)}{p(x)p(y)p(z)}\right] &= E[\log p(x, y, z) - \log p(x) - \log p(y) - \log p(z)] \\ &= E[\log p(x, y, z)] - E[\log p(x)] - E[\log p(y)] - E[\log p(z)] \\ &= -H(X, Y, Z) + H(X) + H(Y) + H(Z) \end{aligned}$$

in which the quantity is zero iff  $X, Y$  and  $Z$  are mutually independent

i.e.  $p(x, y, z) = p(x)p(y)p(z)$ .

2. Let the random variable  $X$  have three possible outcomes  $a, b, c$ . Consider two distributions on this random variable:

symbol	$p(x)$	$p(y)$
a	$\frac{1}{2}$	$\frac{1}{3}$
b	$\frac{1}{4}$	$\frac{1}{3}$
c	$\frac{1}{4}$	$\frac{1}{3}$

Calculate  $H(p)$ ,  $H(q)$ ,  $D(p||q)$  and  $D(q||p)$ . Verify that in this case,  $D(p||q) \neq D(q||p)$

**Answer:**

$$\begin{aligned}
H(p) &= -\left(\frac{1}{2} \log \frac{1}{2} + \frac{1}{4} \log \frac{1}{4} + \frac{1}{4} \log \frac{1}{4}\right) = \frac{3}{2} \\
H(q) &= -\left(\frac{1}{3} \log \frac{1}{3} + \frac{1}{3} \log \frac{1}{3} + \frac{1}{3} \log \frac{1}{3}\right) = \log 3 \\
D(p||q) &= \frac{1}{2} \log \frac{\frac{1}{2}}{\frac{1}{3}} + \frac{1}{4} \log \frac{\frac{1}{4}}{\frac{1}{3}} + \frac{1}{4} \log \frac{\frac{1}{4}}{\frac{1}{3}} = \log 3 - \frac{3}{2} \\
D(q||p) &= \frac{1}{3} \log \frac{\frac{1}{3}}{\frac{1}{2}} + \frac{1}{3} \log \frac{\frac{1}{3}}{\frac{1}{4}} + \frac{1}{3} \log \frac{\frac{1}{3}}{\frac{1}{4}} = -\log 3 + \frac{5}{3}
\end{aligned}$$

3. Show that  $\ln x \geq 1 - \frac{1}{x}$  for  $x > 0$ , where the equality holds when  $x = 1$ .

**Proof:**

Let  $f(x) = \ln x + \frac{1}{x} - 1$ , and  $f(1) = 0$ .

The derivative of  $f(x)$  is  $\frac{d}{dx}f(x) = \frac{1}{x} - \frac{1}{x^2}$ .

(1). For  $0 < x < 1$ ,  $\frac{d}{dx}f(x) = \frac{x-1}{x^2} < 0$ . Therefore  $f(x)$  is monotonically decreasing over  $(0, 1)$

(2). For  $x > 1$ ,  $\frac{d}{dx}f(x) = \frac{x-1}{x^2} > 0$ . Therefore  $f(x)$  is monotonically increasing over  $(1, +\infty)$

Hence, for  $x > 0$ ,  $\ln x \geq 1 - \frac{1}{x}$ .

4. *Conditioning reduces entropy.* Show that  $H(Y|X) \leq H(Y)$  with equality iff  $X$  and  $Y$  are independent.

**Proof:**

**Claim**  $D(p||q) \geq 0$ , with equality iff  $p(x) = q(x)$  for all  $x$ .

$$\begin{aligned}
 D(p||q) &= \sum_{x \in \mathcal{X}} p(x) \log \frac{p(x)}{q(x)} \\
 &= - \sum_{x \in \mathcal{X}} p(x) \log \frac{q(x)}{p(x)} \\
 &\geq - \log \sum_{x \in \mathcal{X}} p(x) \frac{q(x)}{p(x)} \\
 &= - \log \sum_{x \in \mathcal{X}} q(x) \\
 &= - \log 1 \\
 &= 0
 \end{aligned}$$

Also,  $H(Y) - H(Y|X) = I(X;Y) = D(p(x,y)||p(x)p(y)) \geq 0$  and equality holds iff  $p(x,y) = p(x)p(y)$ , in which  $p(x)$  and  $p(y)$  are independent.

5. Show that  $I(X;Y|Z) \geq 0$  with equality iff  $X \rightarrow Z \rightarrow Y$ .

**Proof:**

Shown in Question 4 that

$$I(X;Y) = D(p(x,y)||p(x)p(y)) \geq 0$$

we can expand the conclusion to conditional mutual information given  $Z$  since that

$$I(X;Y|Z) = \mathbf{E}_{p(x,y,z)} \log \frac{p(X,Y|Z)}{p(X|Z)p(Y|Z)} \geq 0$$

and also equality holds iff  $X$  and  $Y$  are independent conditioning to  $Z$ , which is also a necessary condition of Markov chain  $X \rightarrow Z \rightarrow Y$ .

6. *Data processing.* Let  $X_1 \rightarrow X_2 \rightarrow X_3 \rightarrow \dots \rightarrow X_n$  form a Markov chain, i.e.,

$$p(x_1, x_2, \dots, x_n) = p(x_1)p(x_2|x_1)\dots p(x_n|x_{n-1})$$

Reduce  $I(X_1; X_2, \dots, X_n)$  to its simplest form.

**Answer**

From the chain rule for mutual information we have

$$\begin{aligned} I(X_1; X_2, \dots, X_n) &= I(X_1; X_2) + I(X_1; X_3|X_2) + \dots + I(X_1; X_n|X_2, \dots, X_{n-2}) \\ &= I(X_1; X_2) \end{aligned}$$

7. Let  $X$  and  $Y$  be two random variables and let  $Z$  be independent of  $(X, Y)$ . Show that  $I(X; Y) \geq I(X; g(Y, Z))$  for any function  $g$ .

**Proof:**

$$\begin{aligned} I(X; Y) &= H(X) - H(X|Y) && \text{by definition} \\ &= H(X) - H(X|(Y, Z)) && \text{independence between } \mathbf{X} \text{ and } \mathbf{Z} \\ &\geq H(X) - H(X|g(Y, Z)) && H(X|(Y, Z)) \leq H(X|g(Y, Z)) \\ & && \text{with equality iff } (Y, Z) \text{ is a function of } g(Y, Z) \\ &= I(X; g(Y, Z)) \end{aligned}$$

8. *Bottleneck.* Suppose that a (non-stationary) Markov chain starts in one of  $n$  states, necks down to  $k < n$  states, and then fans back to  $m > k$  states. Thus  $X_1 \rightarrow X_2 \rightarrow X_3$ , that is,  $p(x_1, x_2, x_3) = p(x_1)p(x_2|x_1)p(x_3|x_2)$ , for all  $x_1 \in \{1, 2, \dots, n\}$ ,  $x_2 \in \{1, 2, \dots, k\}$ ,

$x_3 \in \{1, 2, \dots, m\}$ .

(a) Show that the dependence of  $X_1$  and  $X_3$  is limited by the bottleneck by proving that  $I(X_1; X_3) \leq \log k$

(b) Evaluate  $I(X_1; X_3)$  for  $k = 1$ , and conclude that no dependence can survive such a bottleneck.

**Proof:**

(a) By the data processing inequality

$$\begin{aligned} I(X_1; X_3) &\leq I(X_1; X_2) \\ &= H(X_2) - H(X_2|X_1) \\ &\leq H(X_2) \\ &\leq \log k \end{aligned}$$

(b) If  $k = 1$ , then

$$\begin{aligned} I(X_1; X_3) &\leq \log 1 \\ &= 0 \end{aligned}$$

in which  $I(X_1; X_3) \geq 0$ , thus  $I(X_1; X_3) = 0 \rightarrow X_1$  and  $X_3$  are independent.