

# Lecture 24: Symmetric Matrices

## MAT2040 Linear Algebra

## Theorem 24.1

*If  $A$  is a symmetric (real) matrix, then any two eigenvectors of  $A$  from different eigenspaces are orthogonal.*

## Definition 24.2

$A$  is **orthogonally diagonalizable** if  $A = Q\Lambda Q^T$  for some matrix  $Q$  with orthonormal columns (so  $Q^{-1} = Q^T$ ), and  $\Lambda$  a diagonal matrix.

### Example 24.3

Suppose  $A$  is orthogonally diagonalizable. What can we say about  $A^T$ ?

### Example 24.3

Suppose  $A$  is orthogonally diagonalizable. What can we say about about  $A^T$ ?

Converse is true as well!

## Theorem 24.4

*$A$  is orthogonally diagonalizable if and only if  $A$  is a symmetric matrix.*

(No proof.)

## Theorem 24.5 (Spectral Theorem)

*If  $A$  is a symmetric (real) matrix, then  $A$  is diagonalizable, all of  $A$ 's eigenvalues are real, and the eigenvectors of  $A$  are mutually orthogonal.*

## Theorem 24.5 (Spectral Theorem)

*If  $A$  is a symmetric (real) matrix, then  $A$  is diagonalizable, all of  $A$ 's eigenvalues are real, and the eigenvectors of  $A$  are mutually orthogonal.*

*In other words, we can write  $A$  as*

$$A = Q\Lambda Q^T$$

*where  $Q$  is a matrix with orthonormal columns, and  $\Lambda$  is a diagonal matrix.*

(No proof.)



## Corollary 24.6 (Spectral Decomposition)

*If  $A$  is a symmetric (real)  $n \times n$  matrix, then*

$$A = \lambda_1 \mathbf{q}_1 \mathbf{q}_1^T + \lambda_2 \mathbf{q}_2 \mathbf{q}_2^T + \cdots + \lambda_n \mathbf{q}_n \mathbf{q}_n^T$$

*where  $\lambda_1, \lambda_2, \dots, \lambda_n$  are the (real) eigenvalues of  $A$ , and  $\mathbf{q}_1, \mathbf{q}_2, \dots, \mathbf{q}_n$  the corresponding eigenvectors (that form an orthonormal set).*

## Corollary 24.6 (Spectral Decomposition)

*If  $A$  is a symmetric (real)  $n \times n$  matrix, then*

$$A = \lambda_1 \mathbf{q}_1 \mathbf{q}_1^T + \lambda_2 \mathbf{q}_2 \mathbf{q}_2^T + \cdots + \lambda_n \mathbf{q}_n \mathbf{q}_n^T$$

*where  $\lambda_1, \lambda_2, \dots, \lambda_n$  are the (real) eigenvalues of  $A$ , and  $\mathbf{q}_1, \mathbf{q}_2, \dots, \mathbf{q}_n$  the corresponding eigenvectors (that form an orthonormal set).*

Note: Each term is a rank one matrix. In fact,  $\mathbf{q}_i \mathbf{q}_i^T$  is the projection matrix onto  $\text{Span}\{\mathbf{q}_i\}$ .

## Example 24.7

Construct a spectral decomposition of  $A$  where  $A$  is a matrix with the following orthogonal decomposition

$$A = \begin{bmatrix} -6 & 12 \\ 12 & 1 \end{bmatrix} = \begin{bmatrix} \frac{3}{5} & \frac{4}{5} \\ \frac{4}{5} & -\frac{3}{5} \end{bmatrix} \begin{bmatrix} 10 & 0 \\ 0 & -15 \end{bmatrix} \begin{bmatrix} \frac{3}{5} & \frac{4}{5} \\ \frac{4}{5} & -\frac{3}{5} \end{bmatrix}.$$

Spectral decomposition is for symmetric (square) matrices. Can we find a similar decomposition for general (rectangular) matrices?

Spectral decomposition is for symmetric (square) matrices. Can we find a similar decomposition for general (rectangular) matrices?

Yes! For  $m \times n$  matrix  $A$ , the decomposition we will aim for is

$$A = U\Sigma V^T,$$

where  $U$  is an  $m \times m$  matrix with orthonormal columns,  $\Sigma$  is an  $m \times n$  matrix with only nonzero entries in position  $(i, i)$  for  $i = 1, 2, \dots, \min\{m, n\}$ , and  $V$  is an  $n \times n$  matrix with orthonormal columns.

For  $m \times n$  matrix  $A$ , the decomposition we will aim for is

$$A = U\Sigma V^T.$$

Note that  $A^T A$  is a symmetric  $n \times n$  matrix, and  $AA^T$  is a symmetric  $m \times m$  matrix.

For  $m \times n$  matrix  $A$ , the decomposition we will aim for is

$$A = U\Sigma V^T.$$

Note that  $A^T A$  is a symmetric  $n \times n$  matrix, and  $AA^T$  is a symmetric  $m \times m$  matrix.

We will use the eigendecompositions of  $A^T A$  and  $AA^T$  to find the matrices  $U$ ,  $\Sigma$  and  $V$ .

By the spectral decomposition theorem we know we can write

$$A^T A = Q_1 \Lambda_1 Q_1^T$$

and

$$AA^T = Q_2 \Lambda_2 Q_2^T.$$



By the spectral decomposition theorem we know we can write

$$A^T A = Q_1 \Lambda_1 Q_1^T$$

and

$$A A^T = Q_2 \Lambda_2 Q_2^T.$$

Also: if  $A = U \Sigma V^T$  then

$$A^T A = (U \Sigma V^T)^T (U \Sigma V^T) = V \Sigma^T U^T U \Sigma V^T = V \Sigma^T \Sigma V^T.$$

$$\text{and } A A^T = (U \Sigma V^T)(U \Sigma V^T)^T = U \Sigma V^T V \Sigma^T U^T = U \Sigma \Sigma^T U^T.$$

By the spectral decomposition theorem we know we can write

$$A^T A = Q_1 \Lambda_1 Q_1^T$$

and

$$A A^T = Q_2 \Lambda_2 Q_2^T.$$

Also: if  $A = U \Sigma V^T$  then

$$A^T A = (U \Sigma V^T)^T (U \Sigma V^T) = V \Sigma^T U^T U \Sigma V^T = V \Sigma^T \Sigma V^T.$$

$$\text{and } A A^T = (U \Sigma V^T)(U \Sigma V^T)^T = U \Sigma V^T V \Sigma^T U^T = U \Sigma \Sigma^T U^T.$$

So **if** we somehow can find a  $\Sigma$  so that  $\Sigma^T \Sigma = \Lambda_1$  and  $\Sigma \Sigma^T = \Lambda_2$ , then we can set  $V = Q_1$  and  $U = Q_2$  and have the decomposition we want!

By the spectral decomposition theorem we know we can write

$$A^T A = Q_1 \Lambda_1 Q_1^T$$

and

$$A A^T = Q_2 \Lambda_2 Q_2^T.$$

Also: if  $A = U \Sigma V^T$  then

$$A^T A = (U \Sigma V^T)^T (U \Sigma V^T) = V \Sigma^T U^T U \Sigma V^T = V \Sigma^T \Sigma V^T.$$

$$\text{and } A A^T = (U \Sigma V^T)(U \Sigma V^T)^T = U \Sigma V^T V \Sigma^T U^T = U \Sigma \Sigma^T U^T.$$

So **if** we somehow can find a  $\Sigma$  so that  $\Sigma^T \Sigma = \Lambda_1$  and  $\Sigma \Sigma^T = \Lambda_2$ , then we can set  $V = Q_1$  and  $U = Q_2$  and have the decomposition we want! (Almost true — I'll get back to this.)

Let's think about the eigenvalues of  $A^T A$  and  $AA^T$ :

Eigenvalues and eigenvectors of  $A^T A$  are pairs of  $\lambda$  and nonzero  $\mathbf{x}$  so that  $A^T A \mathbf{x} = \lambda \mathbf{x}$ .

Let's think about the eigenvalues of  $A^T A$  and  $AA^T$ :

Eigenvalues and eigenvectors of  $A^T A$  are pairs of  $\lambda$  and nonzero  $\mathbf{x}$  so that  $A^T A \mathbf{x} = \lambda \mathbf{x}$ .

$$\mathbf{x}^T A^T A \mathbf{x} = \mathbf{x}^T \lambda \mathbf{x}$$

Let's think about the eigenvalues of  $A^T A$  and  $AA^T$ :

Eigenvalues and eigenvectors of  $A^T A$  are pairs of  $\lambda$  and nonzero  $\mathbf{x}$  so that  $A^T A \mathbf{x} = \lambda \mathbf{x}$ .

$$\mathbf{x}^T A^T A \mathbf{x} = \mathbf{x}^T \lambda \mathbf{x} = \lambda \mathbf{x}^T \mathbf{x}.$$

Let's think about the eigenvalues of  $A^T A$  and  $AA^T$ :

Eigenvalues and eigenvectors of  $A^T A$  are pairs of  $\lambda$  and nonzero  $\mathbf{x}$  so that  $A^T A \mathbf{x} = \lambda \mathbf{x}$ .

$$\mathbf{x}^T A^T A \mathbf{x} = \mathbf{x}^T \lambda \mathbf{x} = \lambda \mathbf{x}^T \mathbf{x}.$$

$$\text{Also } \mathbf{x}^T A^T A \mathbf{x} = (A \mathbf{x})^T (A \mathbf{x})$$

Let's think about the eigenvalues of  $A^T A$  and  $AA^T$ :

Eigenvalues and eigenvectors of  $A^T A$  are pairs of  $\lambda$  and nonzero  $\mathbf{x}$  so that  $A^T A \mathbf{x} = \lambda \mathbf{x}$ .

$$\mathbf{x}^T A^T A \mathbf{x} = \mathbf{x}^T \lambda \mathbf{x} = \lambda \mathbf{x}^T \mathbf{x}.$$

$$\text{Also } \mathbf{x}^T A^T A \mathbf{x} = (A \mathbf{x})^T (A \mathbf{x}) \geq 0.$$



Let's think about the eigenvalues of  $A^T A$  and  $AA^T$ :

Eigenvalues and eigenvectors of  $A^T A$  are pairs of  $\lambda$  and nonzero  $\mathbf{x}$  so that  $A^T A\mathbf{x} = \lambda\mathbf{x}$ .

$$\mathbf{x}^T A^T A\mathbf{x} = \mathbf{x}^T \lambda\mathbf{x} = \lambda\mathbf{x}^T \mathbf{x}.$$

$$\text{Also } \mathbf{x}^T A^T A\mathbf{x} = (A\mathbf{x})^T (A\mathbf{x}) \geq 0.$$

This means that  $\lambda \geq 0$  (because  $\mathbf{x}^T \mathbf{x} \geq 0$ ).

Let's think about the eigenvalues of  $A^T A$  and  $AA^T$ :

Eigenvalues and eigenvectors of  $A^T A$  are pairs of  $\lambda$  and nonzero  $\mathbf{x}$  so that  $A^T A\mathbf{x} = \lambda\mathbf{x}$ .

$$\mathbf{x}^T A^T A\mathbf{x} = \mathbf{x}^T \lambda\mathbf{x} = \lambda\mathbf{x}^T \mathbf{x}.$$

$$\text{Also } \mathbf{x}^T A^T A\mathbf{x} = (A\mathbf{x})^T (A\mathbf{x}) \geq 0.$$

This means that  $\lambda \geq 0$  (because  $\mathbf{x}^T \mathbf{x} \geq 0$ ).

Same reasoning holds for eigenvalues of  $AA^T$ .

Let's think about the eigenvalues of  $A^T A$  and  $AA^T$ :

Eigenvalues and eigenvectors of  $A^T A$  are pairs of  $\lambda$  and nonzero  $\mathbf{x}$  so that  $A^T A\mathbf{x} = \lambda\mathbf{x}$ .

$$\mathbf{x}^T A^T A\mathbf{x} = \mathbf{x}^T \lambda\mathbf{x} = \lambda\mathbf{x}^T \mathbf{x}.$$

$$\text{Also } \mathbf{x}^T A^T A\mathbf{x} = (A\mathbf{x})^T (A\mathbf{x}) \geq 0.$$

This means that  $\lambda \geq 0$  (because  $\mathbf{x}^T \mathbf{x} \geq 0$ ).

Same reasoning holds for eigenvalues of  $AA^T$ .

**All eigenvalues of  $A^T A$  and  $AA^T$  are nonnegative.**

Let's think about the eigenvalues of  $A^T A$  and  $AA^T$ :

Suppose  $\mathbf{x}$  is an eigenvector of  $A^T A$  with corresponding eigenvalue  $\lambda$ , i.e.,  $A^T A \mathbf{x} = \lambda \mathbf{x}$ .

Let's think about the eigenvalues of  $A^T A$  and  $AA^T$ :

Suppose  $\mathbf{x}$  is an eigenvector of  $A^T A$  with corresponding eigenvalue  $\lambda$ , i.e.,  $A^T A \mathbf{x} = \lambda \mathbf{x}$ .

$$AA^T A \mathbf{x} = A \lambda \mathbf{x}.$$

Let's think about the eigenvalues of  $A^T A$  and  $AA^T$ :

Suppose  $\mathbf{x}$  is an eigenvector of  $A^T A$  with corresponding eigenvalue  $\lambda$ , i.e.,  $A^T A \mathbf{x} = \lambda \mathbf{x}$ .

$$AA^T A \mathbf{x} = A \lambda \mathbf{x}.$$

$$\text{Or: } AA^T (A \mathbf{x}) = \lambda (A \mathbf{x}).$$

Let's think about the eigenvalues of  $A^T A$  and  $AA^T$ :

Suppose  $\mathbf{x}$  is an eigenvector of  $A^T A$  with corresponding eigenvalue  $\lambda$ , i.e.,  $A^T A \mathbf{x} = \lambda \mathbf{x}$ .

$$AA^T A \mathbf{x} = A \lambda \mathbf{x}.$$

$$\text{Or: } AA^T (A \mathbf{x}) = \lambda (A \mathbf{x}).$$

Note that this is true if  $\lambda$  is an eigenvalue of  $AA^T$ , or if  $A \mathbf{x} = \mathbf{0}$ .

Let's think about the eigenvalues of  $A^T A$  and  $AA^T$ :

Suppose  $\mathbf{x}$  is an eigenvector of  $A^T A$  with corresponding eigenvalue  $\lambda$ , i.e.,  $A^T A\mathbf{x} = \lambda\mathbf{x}$ .

$$AA^T A\mathbf{x} = A\lambda\mathbf{x}.$$

$$\text{Or: } AA^T(A\mathbf{x}) = \lambda(A\mathbf{x}).$$

Note that this is true if  $\lambda$  is an eigenvalue of  $AA^T$ , or if  $A\mathbf{x} = \mathbf{0}$ .  
If  $\lambda > 0$  then  $A\mathbf{x} \neq \mathbf{0}$ , because  $A^T A\mathbf{x} = \lambda\mathbf{x}$  and  $\mathbf{x} \neq \mathbf{0}$ .



Let's think about the eigenvalues of  $A^T A$  and  $AA^T$ :

Suppose  $\mathbf{x}$  is an eigenvector of  $A^T A$  with corresponding eigenvalue  $\lambda$ , i.e.,  $A^T A \mathbf{x} = \lambda \mathbf{x}$ .

$$AA^T A \mathbf{x} = A \lambda \mathbf{x}.$$

$$\text{Or: } AA^T (A \mathbf{x}) = \lambda (A \mathbf{x}).$$

Note that this is true if  $\lambda$  is an eigenvalue of  $AA^T$ , or if  $A \mathbf{x} = \mathbf{0}$ .  
If  $\lambda > 0$  then  $A \mathbf{x} \neq \mathbf{0}$ , because  $A^T A \mathbf{x} = \lambda \mathbf{x}$  and  $\mathbf{x} \neq \mathbf{0}$ .

**$AA^T$  and  $A^T A$  have the same positive eigenvalues.**

Let  $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_r > \lambda_{r+1} = 0$  be the positive eigenvalues of  $A^T A$  and  $AA^T$ .

Let  $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_r > \lambda_{r+1} = 0$  be the positive eigenvalues of  $A^T A$  and  $AA^T$ .

Define  $\Sigma =$  
$$\left[ \begin{array}{cccc|c} \sqrt{\lambda_1} & & & & 0 \\ & \sqrt{\lambda_2} & & & \\ & & \ddots & & \\ & & & \sqrt{\lambda_p} & \\ \hline & & & & 0 \end{array} \right]$$

so that  $\Sigma$  is an  $m \times n$  matrix.

Then we have the decompositions

$$A^T A = Q_1 \Lambda_1 Q_1^T$$

and

$$A A^T = Q_2 \Lambda_2 Q_2^T.$$

for  $\Lambda_1 = \Sigma^T \Sigma$  and  $\Lambda_2 = \Sigma \Sigma^T$ !

The only problem is that the orthogonal matrix is not uniquely defined in the eigendecomposition!

The only problem is that the orthogonal matrix is not uniquely defined in the eigendecomposition!

If  $Q = [\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n]$  is a matrix with orthonormal columns, then  $\tilde{Q} = [\pm\mathbf{b}_1, \pm\mathbf{b}_2, \dots, \pm\mathbf{b}_n]$  is also a matrix with orthonormal columns for any choice of pluses and minuses, and the columns are still eigenvectors corresponding to the diagonal elements in  $\Lambda$  in order!

This is easy to fix, though. Note that  $A = U\Sigma V^T$  implies  $AV = U\Sigma$ .

So can choose  $V = Q_1$  for **any**  $Q_1$  so that  $A^T A = Q_1 \Lambda_1 Q_1^T$ , and then find  $U$  from

$$AV = U\Sigma.$$

This is easy to fix, though. Note that  $A = U\Sigma V^T$  implies  $AV = U\Sigma$ .

So can choose  $V = Q_1$  for **any**  $Q_1$  so that  $A^T A = Q_1 \Lambda_1 Q_1^T$ , and then find  $U$  from

$$AV = U\Sigma.$$

We can find  $U$  column by column:  $\sigma_i \mathbf{u}_i = A\mathbf{v}_i$  for  $i = 1, 2, \dots, r$ .



This is easy to fix, though. Note that  $A = U\Sigma V^T$  implies  $AV = U\Sigma$ .

So can choose  $V = Q_1$  for **any**  $Q_1$  so that  $A^T A = Q_1 \Lambda_1 Q_1^T$ , and then find  $U$  from

$$AV = U\Sigma.$$

We can find  $U$  column by column:  $\sigma_i \mathbf{u}_i = A\mathbf{v}_i$  for  $i = 1, 2, \dots, r$ .

(The other columns of  $U$  can be chosen arbitrarily as long as  $U$ 's columns form an orthonormal basis.)

## Theorem 24.8 (Singular Value Decomposition)

*Any  $m \times n$  matrix  $A$  can be written as*

$$A = U\Sigma V^T$$

*where  $U$  is an  $m \times m$  matrix with orthonormal columns,  $\Sigma$  is an  $m \times n$  matrix with only nonzero entries in position  $(i, i)$  for  $i = 1, 2, \dots, \min\{m, n\}$ , and  $V$  is an  $n \times n$  matrix with orthonormal columns.*

## Theorem 24.8 (Singular Value Decomposition)

*Any  $m \times n$  matrix  $A$  can be written as*

$$A = U\Sigma V^T$$

*where  $U$  is an  $m \times m$  matrix with orthonormal columns,  $\Sigma$  is an  $m \times n$  matrix with only nonzero entries in position  $(i, i)$  for  $i = 1, 2, \dots, \min\{m, n\}$ , and  $V$  is an  $n \times n$  matrix with orthonormal columns.*

The  $\min\{m, n\}$  entries in  $\Sigma$  in position  $(i, i)$  (for  $i = 1, 2, \dots, n$ ) are called the **singular values** of  $A$ .

Finding SVD of  $A$ :

- Find eigendecomposition of  $A^T A = Q_1 \Lambda_1 Q_1^T$ .  
Reorder eigenvalues (diagonal elements in  $\Lambda$ ) in nonincreasing order, and order the eigenvectors (columns of  $Q_1$ ) in the same way.

- $V = Q_1$ .

- $$\Sigma = \left[ \begin{array}{ccc|c} \sqrt{\lambda_1} & & & 0 \\ & \sqrt{\lambda_2} & & \\ & & \ddots & \\ & & & \sqrt{\lambda_p} \\ \hline & 0 & & 0 \end{array} \right]$$

- Construct  $U$  as follows:  
 $\mathbf{u}_i = \frac{1}{\sqrt{\lambda_i}} A \mathbf{v}_i$  for  $i = 1, 2, \dots, r$ , where  $\lambda_r$  is the smallest nonzero eigenvalue of  $A^T A$ .  
Finish the orthonormal basis arbitrarily.

Why is this cool?

Why is this cool?

It gives us a handle on Col  $A$  and Row  $A$ :

Why is this cool?

It gives us a handle on Col  $A$  and Row  $A$ :

$A = U\Sigma V^T$  implies  $AV = U\Sigma$ , which means that the first  $r$  columns of  $U$  are in Col  $A$  (where  $r$  is the number of nonzero singular values).

Why is this cool?

It gives us a handle on Col  $A$  and Row  $A$ :

$A = U\Sigma V^T$  implies  $AV = U\Sigma$ , which means that the first  $r$  columns of  $U$  are in Col  $A$  (where  $r$  is the number of nonzero singular values).

$A = U\Sigma V^T$  also implies  $U^T A = \Sigma V^T$ , or  $A^T U = V\Sigma$  which means that the first  $r$  columns of  $V$  are in Row  $A$ .



Even stronger:

$$U\Sigma V^T \mathbf{v}_i =$$

Even stronger:

$$U\Sigma V^T \mathbf{v}_i = U\Sigma \mathbf{e}_i =$$

Even stronger:

$$U\Sigma V^T \mathbf{v}_i = U\Sigma \mathbf{e}_i = \begin{cases} U\sigma_i \mathbf{e}_i = \sigma_i \mathbf{u}_i & \text{for } i \leq r \\ \mathbf{0} & \text{otherwise,} \end{cases}$$


where  $\mathbf{u}_i$  is the  $i$ th column of  $U$ , and  $r$  is the number of nonzero singular values.

Even stronger:

$$U\Sigma V^T \mathbf{v}_i = U\Sigma \mathbf{e}_i = \begin{cases} U\sigma_i \mathbf{e}_i = \sigma_i \mathbf{u}_i & \text{for } i \leq r \\ \mathbf{0} & \text{otherwise,} \end{cases}$$

where  $\mathbf{u}_i$  is the  $i$ th column of  $U$ , and  $r$  is the number of nonzero singular values.

This decomposition is telling is **exactly** where every vector in Row  $A$  is mapped, and how it is scaled!!! (Using orthonormal bases!)



(picture from David Lay, Linear Algebra.)

## Theorem 24.9 (Rank one decomposition)

Any  $m \times n$  matrix  $A$  can be written as  $A = U\Sigma V^T$ , and thus as

$$A = \sigma_1 \mathbf{u}_1 \mathbf{v}_1^T + \sigma_2 \mathbf{u}_2 \mathbf{v}_2^T + \cdots + \sigma_r \mathbf{u}_r \mathbf{v}_r^T$$

where  $r$  is the number of positive singular values of  $A$ .  
(Note that  $r = \text{rank } A$ !)

## Theorem 24.9 (Rank one decomposition)

*Any  $m \times n$  matrix  $A$  can be written as  $A = U\Sigma V^T$ , and thus as*

$$A = \sigma_1 \mathbf{u}_1 \mathbf{v}_1^T + \sigma_2 \mathbf{u}_2 \mathbf{v}_2^T + \cdots + \sigma_r \mathbf{u}_r \mathbf{v}_r^T$$

*where  $r$  is the number of positive singular values of  $A$ .  
(Note that  $r = \text{rank } A$ !)*

Note: Each term is a rank one matrix.