

Lecture 14: More Vector Spaces of Matrices

MAT2040 Linear Algebra

Definition 14.1

The **row space** of a matrix A is the set of all possible linear combinations of the rows of A (written as column vectors).

It is denoted by $\text{Row } A$.

In other words,

$$\text{Row } A = \text{Span}\{\vec{a}_1^T, \vec{a}_2^T, \dots, \vec{a}_m^T\}.$$

Fact 14.2

Let A be an $m \times n$ matrix. $\text{Row } A$ is a subspace of \mathbb{R}^n .

Fact 14.2

Let A be an $m \times n$ matrix. $\text{Row } A$ is a subspace of \mathbb{R}^n .

Fact 14.3

$\text{Row } A = \text{Col } A^T$ for any matrix A .

How can we find a basis for Row A ?

How can we find a basis for Row A ?

We know how to find a basis for $\text{Col } A^T$.

How can we find a basis for Row A ?

We know how to find a basis for $\text{Col } A^T$. Done?

How can we find a basis for Row A ?

We know how to find a basis for $\text{Col } A^T$. Done?

Suppose we want to find a basis for both $\text{Col } A$ and $\text{Row } A$. Do we really need to find a row echelon form for both A and A^T to identify these bases?

How can we find a basis for Row A ?

We know how to find a basis for $\text{Col } A^T$. Done?

Suppose we want to find a basis for both $\text{Col } A$ and $\text{Row } A$. Do we really need to find a row echelon form for both A and A^T to identify these bases? No!

Theorem 14.4

If A and B are row-equivalent matrices then $\text{Row } A = \text{Row } B$.

Theorem 14.4

If A and B are row-equivalent matrices then $\text{Row } A = \text{Row } B$.

Basis for $\text{Row } A$ is easy when A is in row echelon form! (Why?)

Theorem 14.4

If A and B are row-equivalent matrices then $\text{Row } A = \text{Row } B$.

Basis for $\text{Row } A$ is easy when A is in row echelon form! (Why?)

So to find a basis for both $\text{Col } A$ and $\text{Row } A$ we only have to find the row echelon form of A (not of A and A^T both).

Theorem 14.4

If A and B are row-equivalent matrices then $\text{Row } A = \text{Row } B$.

Basis for $\text{Row } A$ is easy when A is in row echelon form! (Why?)

So to find a basis for both $\text{Col } A$ and $\text{Row } A$ we only have to find the row echelon form of A (not of A and A^T both).

Also, we immediately see the following.

Corollary 14.5

$\dim(\text{Col } A) = \dim(\text{Row } A)$ for any matrix A .

Example 14.6

Find Col A and Row A for $A = \begin{bmatrix} 2 & 5 & -2 & -3 & 1 \\ 4 & 10 & -8 & -16 & -6 \\ 6 & 15 & -8 & -8 & 9 \\ 2 & 5 & 0 & -1 & 0 \end{bmatrix}$

Example 14.6

Find Col A and Row A for $A = \begin{bmatrix} 2 & 5 & -2 & -3 & 1 \\ 4 & 10 & -8 & -16 & -6 \\ 6 & 15 & -8 & -8 & 9 \\ 2 & 5 & 0 & -1 & 0 \end{bmatrix}$

This may be helpful: A is row equivalent to $\begin{bmatrix} 2 & 5 & -2 & -3 & 1 \\ 0 & 0 & 2 & 2 & -1 \\ 0 & 0 & 0 & 3 & 5 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$

Definition 14.7

The **rank** of a matrix A is the dimension of $\text{Col } A$. We use $\text{rank}(A)$ to denote this quantity.

Definition 14.7

The **rank** of a matrix A is the dimension of $\text{Col } A$. We use $\text{rank}(A)$ to denote this quantity.

We just saw that $\text{rank}(A)$ therefore also equals the dimension of $\text{Row } A$.

Example 14.8 (continuation of previous example)

$$\text{Let } A = \begin{bmatrix} 2 & 5 & -2 & -3 & 1 \\ 4 & 10 & -8 & -16 & -6 \\ 6 & 15 & -8 & -8 & 9 \\ 2 & 5 & 0 & -1 & 0 \end{bmatrix}.$$

$$A \text{ is row equivalent to } \begin{bmatrix} 2 & 5 & -2 & -3 & 1 \\ 0 & 0 & 2 & 2 & -1 \\ 0 & 0 & 0 & 3 & 5 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

What is $\text{rank}(A)$?

Example 14.8 (continuation of previous example)

$$\text{Let } A = \begin{bmatrix} 2 & 5 & -2 & -3 & 1 \\ 4 & 10 & -8 & -16 & -6 \\ 6 & 15 & -8 & -8 & 9 \\ 2 & 5 & 0 & -1 & 0 \end{bmatrix}.$$

$$A \text{ is row equivalent to } \begin{bmatrix} 2 & 5 & -2 & -3 & 1 \\ 0 & 0 & 2 & 2 & -1 \\ 0 & 0 & 0 & 3 & 5 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

What is $\text{rank}(A)$? Also, what was $\dim(\text{Null } A)$ again?

Example 14.8 (continuation of previous example)

$$\text{Let } A = \begin{bmatrix} 2 & 5 & -2 & -3 & 1 \\ 4 & 10 & -8 & -16 & -6 \\ 6 & 15 & -8 & -8 & 9 \\ 2 & 5 & 0 & -1 & 0 \end{bmatrix}.$$

$$A \text{ is row equivalent to } \begin{bmatrix} 2 & 5 & -2 & -3 & 1 \\ 0 & 0 & 2 & 2 & -1 \\ 0 & 0 & 0 & 3 & 5 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

What is $\text{rank}(A)$? Also, what was $\dim(\text{Null } A)$ again? What about $\text{rank}(A) + \dim(\text{Null } A)$?

Side note: When considering generalizations of matrices (where the entries are not real or complex numbers), it can be the case that the corresponding notion of $\text{Col } A$ and $\text{Row } A$ do **not** have the same dimension.

Side note: When considering generalizations of matrices (where the entries are not real or complex numbers), it can be the case that the corresponding notion of $\text{Col } A$ and $\text{Row } A$ do **not** have the same dimension.

Therefore these dimensions have a separate names — **column rank** and **row rank**. In this course the entries in a matrix are only real or complex numbers, and that means column rank and row rank have the same value. We will thus use “rank” for this quantities.

Summarizing, a basis for $\text{Col } A$ and $\text{Row } A$ can be found from a row echelon form of matrix A (say matrix B):

Summarizing, a basis for $\text{Col } A$ and $\text{Row } A$ can be found from a row echelon form of matrix A (say matrix B):

- ▶ a basis for $\text{Col } A = \{\mathbf{a}_i : \mathbf{a}_i \text{ is a pivot column}\},$
- ▶ a basis for $\text{Row } A = \{\vec{\mathbf{b}}_i^T : \vec{\mathbf{b}}_i \text{ is not an all zero row}\}.$

Summarizing, a basis for $\text{Col } A$ and $\text{Row } A$ can be found from a row echelon form of matrix A (say matrix B):

- ▶ a basis for $\text{Col } A = \{\mathbf{a}_i : \mathbf{a}_i \text{ is a pivot column}\},$
- ▶ a basis for $\text{Row } A = \{\vec{\mathbf{b}}_i^T : \vec{\mathbf{b}}_i \text{ is not an all zero row}\}.$

Note that the first $k = \text{rank}(A)$ rows of A are not necessarily a basis for $\text{Row } A$! (Why?)

Recall:

Corollary 13.13

Let A be an $m \times n$ matrix.

$$\dim(\text{Col } A) + \dim(\text{Null } A) = n.$$

Recall:

Corollary 13.13

Let A be an $m \times n$ matrix.

$$\dim(\text{Col } A) + \dim(\text{Null } A) = n.$$

We can also state this in terms of rank:

$$\text{rank}(A) + \dim(\text{Null } A) = \# \text{ of columns of } A.$$

Recall:

Corollary 13.13

Let A be an $m \times n$ matrix.

$$\dim(\text{Col } A) + \dim(\text{Null } A) = n.$$

We can also state this in terms of rank:

$$\text{rank}(A) + \dim(\text{Null } A) = \# \text{ of columns of } A.$$

This is often referred to as the “Rank Theorem” or “Rank-Nullity Theorem”.

Recall:

Corollary 13.13

Let A be an $m \times n$ matrix.

$$\dim(\text{Col } A) + \dim(\text{Null } A) = n.$$

We can also state this in terms of rank:

$$\text{rank}(A) + \dim(\text{Null } A) = \# \text{ of columns of } A.$$

This is often referred to as the “Rank Theorem” or “Rank-Nullity Theorem”.

($\dim(\text{Null } A)$ is sometimes called “nullity”.)

Special case 1: Matrices with rank 0?

Special case 2: Matrices with rank 1. What can we say about these matrices?

Special case 2: Matrices with rank 1. What can we say about these matrices?

If an $m \times n$ matrix A has rank 1, then $\text{Col } A = \text{Span}\{\mathbf{u}\}$ for some $\mathbf{u} \neq \mathbf{0}$.

Special case 2: Matrices with rank 1. What can we say about these matrices?

If an $m \times n$ matrix A has rank 1, then $\text{Col } A = \text{Span}\{\mathbf{u}\}$ for some $\mathbf{u} \neq \mathbf{0}$.

Note that every column of A is in $\text{Col } A$ (because $\text{Col } A = \text{Span}\{\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n\}$).

Special case 2: Matrices with rank 1. What can we say about these matrices?

If an $m \times n$ matrix A has rank 1, then $\text{Col } A = \text{Span}\{\mathbf{u}\}$ for some $\mathbf{u} \neq \mathbf{0}$.

Note that every column of A is in $\text{Col } A$ (because $\text{Col } A = \text{Span}\{\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n\}$).

And therefore every column of A is a linear combination of the vector \mathbf{u} (because \mathbf{u} spans $\text{Col } A$).

We see that

$$A = [v_1 \mathbf{u}, \quad v_2 \mathbf{u}, \quad \dots, \quad v_n \mathbf{u}]$$

for some $v_1, v_2, \dots, v_n \in \mathbb{R}$.

We see that

$$A = [v_1 \mathbf{u}, \quad v_2 \mathbf{u}, \quad \dots, \quad v_n \mathbf{u}]$$

for some $v_1, v_2, \dots, v_n \in \mathbb{R}$.

We can also interpret this as

$$A = [\mathbf{u}[v_1], \quad \mathbf{u}[v_2], \quad \dots, \quad \mathbf{u}[v_n]]$$

We see that

$$A = [v_1 \mathbf{u}, \quad v_2 \mathbf{u}, \quad \dots, \quad v_n \mathbf{u}]$$

for some $v_1, v_2, \dots, v_n \in \mathbb{R}$.

We can also interpret this as

$$\begin{aligned} A &= [\mathbf{u}[v_1], \quad \mathbf{u}[v_2], \quad \dots, \quad \mathbf{u}[v_n]] \\ &= \mathbf{u} [v_1, \quad v_2, \quad \dots, \quad v_n] \end{aligned}$$

We see that

$$A = [v_1 \mathbf{u}, \quad v_2 \mathbf{u}, \quad \dots, \quad v_n \mathbf{u}]$$

for some $v_1, v_2, \dots, v_n \in \mathbb{R}$.

We can also interpret this as

$$\begin{aligned} A &= [\mathbf{u}[v_1], \quad \mathbf{u}[v_2], \quad \dots, \quad \mathbf{u}[v_n]] \\ &= \mathbf{u} [v_1, \quad v_2, \quad \dots, \quad v_n] \\ &= \mathbf{u} \mathbf{v}^T. \end{aligned}$$

We see that

$$A = [v_1 \mathbf{u}, \quad v_2 \mathbf{u}, \quad \dots, \quad v_n \mathbf{u}]$$

for some $v_1, v_2, \dots, v_n \in \mathbb{R}$.

We can also interpret this as

$$\begin{aligned} A &= [\mathbf{u}[v_1], \quad \mathbf{u}[v_2], \quad \dots, \quad \mathbf{u}[v_n]] \\ &= \mathbf{u} [v_1, \quad v_2, \quad \dots, \quad v_n] \\ &= \mathbf{u} \mathbf{v}^T. \end{aligned}$$

Again, to avoid getting $A = \mathbf{O}$ (the all zero matrix that has rank 0), we do need $\mathbf{v} \neq \mathbf{0}$.

We just proved one implication of the following fact (and the other implication is easy).

Fact 14.9

The $m \times n$ matrix A has rank 1 if and only if there exist nonzero vectors $\mathbf{u} \in \mathbb{R}^m$ and $\mathbf{v} \in \mathbb{R}^n$ so that

$$A = \mathbf{u}\mathbf{v}^T.$$

We just proved one implication of the following fact (and the other implication is easy).

Fact 14.9

The $m \times n$ matrix A has rank 1 if and only if there exist nonzero vectors $\mathbf{u} \in \mathbb{R}^m$ and $\mathbf{v} \in \mathbb{R}^n$ so that

$$A = \mathbf{u}\mathbf{v}^T.$$

($\mathbf{u}\mathbf{v}^T$ is sometimes referred to as the “outer product of \mathbf{u} and \mathbf{v} ”.)

Final special case: matrices for which the rank is “as large as possible” (meaning there do not exist matrices of the same size with higher rank).

Final special case: matrices for which the rank is “as large as possible” (meaning there do not exist matrices of the same size with higher rank).

We will say that these matrices have **full rank**.

Final special case: matrices for which the rank is “as large as possible” (meaning there do not exist matrices of the same size with higher rank).

We will say that these matrices have **full rank**.

Suppose A is an $m \times n$ matrix and it has full rank.
What is $\text{rank}(A)$?

Definition 14.10

The **left null space** of a matrix A is $\text{Null } A^T$.

Example 14.11

Show that the left null space of the $m \times n$ matrix A is the set $\{\mathbf{y} \in \mathbb{R}^m : \mathbf{y}^T A = \mathbf{0}^T\}$.

Example 14.11

Show that the left null space of the $m \times n$ matrix A is the set $\{\mathbf{y} \in \mathbb{R}^m : \mathbf{y}^T A = \mathbf{0}^T\}$.

Note that this shows that the left null space of A is exactly the vectors corresponding to the weights of a linear dependence relations of the rows of A .

Example 14.11

Show that the left null space of the $m \times n$ matrix A is the set $\{\mathbf{y} \in \mathbb{R}^m : \mathbf{y}^T A = \mathbf{0}^T\}$.

Note that this shows that the left null space of A is exactly the vectors corresponding to the weights of a linear dependence relations of the rows of A .

The null space of A can similarly be described as exactly the vectors corresponding to the weights of a linear dependence relations of the columns of A .

Theorem 14.12 (Invertible Matrix Theorem)

Given a $n \times n$ matrix A (square!). The following are equivalent:

1. A is invertible
2. the only solution to $A\mathbf{x} = \mathbf{0}$ is the trivial solution
3. A is row equivalent to I
4. A is a product of elementary matrices
5. the columns of A are linearly independent
6. $A\mathbf{x} = \mathbf{b}$ has a solution for every $\mathbf{b} \in \mathbb{R}^m$
7. the columns of A span \mathbb{R}^n
8. $\text{Null } A = \{\mathbf{0}\}$
9. $\text{Col } A = \mathbb{R}^n$
10. the columns of A form a basis of \mathbb{R}^n
11. $\dim(\text{Col } A) = n$
12. $\dim(\text{Null } A) = 0$
13. $\text{rank}(A) = n$.