

# Lecture 22: Eigenvalues and Eigenvectors

## MAT2040 Linear Algebra

We will again only consider square matrices (for this topic).

Another decomposition!

$$A = PDP^{-1},$$

where  $D$  is a diagonal matrix.

We will again only consider square matrices (for this topic).

Another decomposition!

$$A = PDP^{-1},$$

where  $D$  is a diagonal matrix.

This decomposition does not always exist — we will study when it exists, and how to find it if it does.

We will again only consider square matrices (for this topic).

Another decomposition!

$$A = PDP^{-1},$$

where  $D$  is a diagonal matrix.

This decomposition does not always exist — we will study when it exists, and how to find it if it does.

But first let's recall what we already know about this decomposition and how we can interpret it.

Recall:

### Example 18.2

Consider  $T(\mathbf{x}) = A\mathbf{x}$  where  $A = \begin{bmatrix} 7 & 2 \\ -4 & 1 \end{bmatrix}$ .

What is the  $\mathcal{B}$ -matrix for  $T$  for  $\mathcal{B} = \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ -2 \end{bmatrix} \right\}$ ?

Express  $A$  in terms of the  $\mathcal{B}$ -matrix for  $T$ .

Recall:

### Example 18.2

Consider  $T(\mathbf{x}) = A\mathbf{x}$  where  $A = \begin{bmatrix} 7 & 2 \\ -4 & 1 \end{bmatrix}$ .

What is the  $\mathcal{B}$ -matrix for  $T$  for  $\mathcal{B} = \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ -2 \end{bmatrix} \right\}$ ?

Express  $A$  in terms of the  $\mathcal{B}$ -matrix for  $T$ .

$$A = P \begin{bmatrix} 5 & 0 \\ 0 & 3 \end{bmatrix} P^{-1}.$$

where  $P = \begin{bmatrix} 1 & 1 \\ -1 & -2 \end{bmatrix}$ .

This decomposition is useful if we think about matrix powers:

$$\begin{aligned} A^k &= \overbrace{(PDP^{-1})(PDP^{-1})\cdots(PDP^{-1})}^{k \text{ times}} \\ &= PD(P^{-1}P)D(P^{-1}P)\cdots(P^{-1}P)DP^{-1} \\ &= PD^kP^{-1} \\ &= P \begin{bmatrix} 5^k & 0 \\ 0 & 3^k \end{bmatrix} P^{-1}. \end{aligned}$$

In general, the  $k$ th power of a diagonal matrix  $D$  is the diagonal matrix with entries equal to the  $k$ th power of the entries of  $D$ .

This decomposition is useful if we think about matrix powers:

$$\begin{aligned} A^k &= \overbrace{(PDP^{-1})(PDP^{-1})\cdots(PDP^{-1})}^{k \text{ times}} \\ &= PD(P^{-1}P)D(P^{-1}P)\cdots(P^{-1}P)DP^{-1} \\ &= PD^kP^{-1} \\ &= P \begin{bmatrix} 5^k & 0 \\ 0 & 3^k \end{bmatrix} P^{-1}. \end{aligned}$$

In general, the  $k$ th power of a diagonal matrix  $D$  is the diagonal matrix with entries equal to the  $k$ th power of the entries of  $D$ .

(It also turns out to be very useful for differential equations, where you will often get a *matrix exponential*. The matrix exponential of a diagonal matrix is, again, easy to compute.)



We interpreted this decomposition as performing the same linear transformation in a different **basis**:

We interpreted this decomposition as performing the same linear transformation in a different **basis**:

It's telling us that the transformation  $A\mathbf{x}$  can be interpreted as scaling the vector  $\mathbf{b}_1 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$  by a factor 5, and scaling the vector  $\mathbf{b}_2 = \begin{bmatrix} 1 \\ -2 \end{bmatrix}$  by a factor 3.

We interpreted this decomposition as performing the same linear transformation in a different **basis**:

It's telling us that the transformation  $A\mathbf{x}$  can be interpreted as scaling the vector  $\mathbf{b}_1 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$  by a factor 5, and scaling the vector  $\mathbf{b}_2 = \begin{bmatrix} 1 \\ -2 \end{bmatrix}$  by a factor 3.

(The image of any vector that is not a multiple of  $\mathbf{b}_1$  and  $\mathbf{b}_2$  will **not** just be scaled: if  $\mathbf{v} = t_1\mathbf{b}_1 + t_2\mathbf{b}_2$  ( $t_1 \neq 0$ ,  $t_2 \neq 0$ ) then

$$A\mathbf{v} = t_1A\mathbf{b}_1 + t_2A\mathbf{b}_2 = 5t_1\mathbf{b}_1 + 3t_2\mathbf{b}_2$$

(by linearity of  $A\mathbf{x}$ , or by matrix algebra).)

The interpretation on the previous slide gives us an idea of how to find this decomposition.



Alexander Yakovlev / Fotolia

We want to know what vectors are just **scaled** by a linear transformation  $A\mathbf{x}$  (and the scaling factor).

We want to know what vectors are just **scaled** by a linear transformation  $A\mathbf{x}$  (and the scaling factor).

If these vectors, say  $\mathcal{B}$ , form a basis for  $\mathbb{R}^n$ , we have a **diagonal**  $\mathcal{B}$ -matrix for the transformation  $A\mathbf{x}$ .

(We will prove a precise statement later.)

## Definition 22.1

Let  $A$  be an  $n \times n$  (square!) matrix.

A scalar  $\lambda$  is called an **eigenvalue** of  $A$  if there exists a nonzero vector  $\mathbf{x}$  so that  $A\mathbf{x} = \lambda\mathbf{x}$ . The vector  $\mathbf{x}$  is called an **eigenvector** corresponding to eigenvalue  $\lambda$ .

## Example 22.2

Let  $A = \begin{bmatrix} 1 & 6 \\ 5 & 2 \end{bmatrix}$ .

Is  $\mathbf{u} = \begin{bmatrix} 6 \\ -5 \end{bmatrix}$  an eigenvector of  $A$ ?



## Example 22.2

Let  $A = \begin{bmatrix} 1 & 6 \\ 5 & 2 \end{bmatrix}$ .

Is  $\mathbf{u} = \begin{bmatrix} 6 \\ -5 \end{bmatrix}$  an eigenvector of  $A$ ?

What about  $\mathbf{v} = \begin{bmatrix} 3 \\ -2 \end{bmatrix}$ ?

### Example 22.2 (continued)

Let  $A = \begin{bmatrix} 1 & 6 \\ 5 & 2 \end{bmatrix}$ .

Show that 7 is an eigenvalue of  $A$  and find (all) eigenvectors corresponding to eigenvalue 7.

Given an eigenvalue, we can find corresponding eigenvectors by solving a system of equations.

Given an eigenvalue, we can find corresponding eigenvectors by solving a system of equations.

How can we find eigenvalues?

Given an eigenvalue, we can find corresponding eigenvectors by solving a system of equations.

How can we find eigenvalues?

Want pairs of  $\lambda$  and  $\mathbf{x} \neq \mathbf{0}$  so that

$$A\mathbf{x} = \lambda\mathbf{x}.$$

Given an eigenvalue, we can find corresponding eigenvectors by solving a system of equations.

How can we find eigenvalues?

Want pairs of  $\lambda$  and  $\mathbf{x} \neq \mathbf{0}$  so that

$$A\mathbf{x} = \lambda\mathbf{x}.$$

Equivalently,

$$A\mathbf{x} = \lambda I\mathbf{x}.$$

Given an eigenvalue, we can find corresponding eigenvectors by solving a system of equations.

How can we find eigenvalues?

Want pairs of  $\lambda$  and  $\mathbf{x} \neq \mathbf{0}$  so that

$$A\mathbf{x} = \lambda\mathbf{x}.$$

Equivalently,

$$A\mathbf{x} = \lambda I\mathbf{x}.$$

Rewriting gives

$$(A - \lambda I)\mathbf{x} = \mathbf{0}.$$

We want  $\lambda$  and  $\mathbf{x} \neq \mathbf{0}$  so that

$$(A - \lambda I)\mathbf{x} = \mathbf{0}.$$



We want  $\lambda$  and  $\mathbf{x} \neq \mathbf{0}$  so that

$$(A - \lambda I)\mathbf{x} = \mathbf{0}.$$

Want  $\lambda$  and  $\mathbf{x} \neq \mathbf{0}$  so that  $\mathbf{x} \in \text{Null}(A - \lambda I)$ !

We want  $\lambda$  and  $\mathbf{x} \neq \mathbf{0}$  so that

$$(A - \lambda I)\mathbf{x} = \mathbf{0}.$$

Want  $\lambda$  and  $\mathbf{x} \neq \mathbf{0}$  so that  $\mathbf{x} \in \text{Null}(A - \lambda I)$ !

So in particular, we want  $\lambda$  so that  $\text{Null}(A - \lambda I) \neq \{\mathbf{0}\}$ .

We want  $\lambda$  and  $\mathbf{x} \neq \mathbf{0}$  so that

$$(A - \lambda I)\mathbf{x} = \mathbf{0}.$$

Want  $\lambda$  and  $\mathbf{x} \neq \mathbf{0}$  so that  $\mathbf{x} \in \text{Null}(A - \lambda I)$ !

So in particular, we want  $\lambda$  so that  $\text{Null}(A - \lambda I) \neq \{\mathbf{0}\}$ .

We can solve this without worrying about  $\mathbf{x}$ :

We want  $\lambda$  and  $\mathbf{x} \neq \mathbf{0}$  so that

$$(A - \lambda I)\mathbf{x} = \mathbf{0}.$$

Want  $\lambda$  and  $\mathbf{x} \neq \mathbf{0}$  so that  $\mathbf{x} \in \text{Null}(A - \lambda I)$ !

So in particular, we want  $\lambda$  so that  $\text{Null}(A - \lambda I) \neq \{\mathbf{0}\}$ .

We can solve this without worrying about  $\mathbf{x}$ :

We want  $\lambda$  so that  $A - \lambda I$  is not invertible!  
(Recall that all matrices are square.)

We want  $\lambda$  and  $\mathbf{x} \neq \mathbf{0}$  so that

$$(A - \lambda I)\mathbf{x} = \mathbf{0}.$$

Want  $\lambda$  and  $\mathbf{x} \neq \mathbf{0}$  so that  $\mathbf{x} \in \text{Null}(A - \lambda I)$ !

So in particular, we want  $\lambda$  so that  $\text{Null}(A - \lambda I) \neq \{\mathbf{0}\}$ .

We can solve this without worrying about  $\mathbf{x}$ :

We want  $\lambda$  so that  $A - \lambda I$  is not invertible!  
(Recall that all matrices are square.)

Which, we know, is equivalent to  $\det(A - \lambda I) = 0$ .

We want  $\lambda$  and  $\mathbf{x} \neq \mathbf{0}$  so that

$$(A - \lambda I)\mathbf{x} = \mathbf{0}.$$

Want  $\lambda$  and  $\mathbf{x} \neq \mathbf{0}$  so that  $\mathbf{x} \in \text{Null}(A - \lambda I)$ !

So in particular, we want  $\lambda$  so that  $\text{Null}(A - \lambda I) \neq \{\mathbf{0}\}$ .

We can solve this without worrying about  $\mathbf{x}$ :

We want  $\lambda$  so that  $A - \lambda I$  is not invertible!  
(Recall that all matrices are square.)

Which, we know, is equivalent to  $\det(A - \lambda I) = 0$ .

This is a polynomial equation where  $\lambda$  is the variable!

Conclusion:

The eigenvalues of  $A$  are exactly the solutions to  $\det(A - \lambda I) = 0$  (where  $\lambda$  is the variable).

### Example 22.3

Find the eigenvalues of  $A = \begin{bmatrix} 3 & 2 \\ 3 & -2 \end{bmatrix}$ .



The polynomial

$$p(\lambda) = \det(A - \lambda I)$$

is called the **characteristic polynomial** for matrix  $A$ , and

$$\det(A - \lambda I) = 0$$

is called the **characteristic equation** for matrix  $A$ .

The polynomial

$$p(\lambda) = \det(A - \lambda I)$$

is called the **characteristic polynomial** for matrix  $A$ , and

$$\det(A - \lambda I) = 0$$

is called the **characteristic equation** for matrix  $A$ .

Note that  $\lambda$  is a variable in both the characteristic polynomial and the characteristic equation.

Summarizing:

- ▶ Finding eigenvalues of  $A$  is equivalent to finding roots of the characteristic equation for matrix  $A$ .

Summarizing:

- ▶ Finding eigenvalues of  $A$  is equivalent to finding roots of the characteristic equation for matrix  $A$ .
- ▶ Finding all eigenvectors belonging to a particular eigenvalue  $\bar{\lambda}$  is equivalent to finding  $\text{Null}(A - \bar{\lambda}I)$  (except that  $\mathbf{0}$  is never an eigenvector).

$\text{Null}(A - \bar{\lambda}I)$  is known as the **eigenspace** belonging to eigenvalue  $\bar{\lambda}$ .

### Theorem 22.4

*If  $A$  and  $B$  are  $n \times n$  matrices that are similar, then  $A$  and  $B$  have the same characteristic polynomial.*