

H1.5: Diagnostic Tests of a Frozen Nonlocal Baryonic Model Against the BTFR and RAR

Vítor Manuel Franco Figueiredo

Independent Researcher, Portugal

ORCID: 0009-0004-7358-4622

January 20, 2026

Abstract

We present H1.5, a set of secondary empirical diagnostics applied to the frozen H1 phenomenological model of disk galaxy dynamics. The H1 model encodes a nonlocal baryon-coupled dynamical response and was previously tested against galaxy rotation curves without parameter tuning. In this work, the H1 outputs are treated as fixed inputs and are confronted with two of the strongest empirical scaling relations in disk galaxies: the baryonic Tully-Fisher relation (BTFR) and the radial acceleration relation (RAR).

Using the full SPARC sample of 175 galaxies, we construct diagnostic tests that involve no refitting, optimization, or adaptive modification of the model. The BTFR diagnostic shows that the frozen H1 outputs preserve the existence and overall structure of the global baryonic mass-velocity relation. In contrast, the RAR diagnostic reveals substantial scatter relative to the observed empirical relation, indicating that the fixed H1 construction does not enforce a universal local acceleration correspondence.

A decomposition of the RAR scatter by radial regime demonstrates that this deviation is structured rather than stochastic. The largest discrepancies occur in the inner regions of galaxies, while the outer disk exhibits systematically reduced scatter. Independent single-galaxy reconstructions confirm numerical consistency at machine precision, ruling out circularity or post-processing artifacts.

Taken together, these results delineate the empirical domain of validity of the frozen H1 model. H1.5 does not propose modifications or extensions, but instead provides a transparent diagnostic baseline against which future dynamical frameworks can be evaluated.

1 Introduction

Empirical scaling relations play a central role in contemporary studies of disk galaxy dynamics. Relations such as the baryonic Tully-Fisher relation (BTFR) and the radial acceleration relation (RAR) provide compact summaries of regularities observed across diverse galaxy populations, and they serve as stringent benchmarks for any phenomenological or theoretical framework that seeks to describe galaxy rotation curves.

The H1 model was introduced as a phenomenological, nonlocal baryon–coupled dynamical framework designed to explore how observed baryonic mass distributions relate to galaxy rotation curves. The model was deliberately constructed with a minimal set of assumptions and without invoking particle dark matter, modifications of local gravitational laws, or galaxy-by-galaxy parameter tuning. In its original formulation, H1 was evaluated primarily through direct comparison with observed rotation curves, establishing that a fixed convolution-based response can reproduce broad kinematic trends across the SPARC sample.

However, agreement with individual rotation curves does not by itself guarantee consistency with higher-level empirical regularities. Secondary diagnostics based on population-level scaling relations provide an independent and often more restrictive assessment of model behavior. In particular, the BTFR probes global mass–velocity scaling, while the RAR tests local, point-wise correspondence between baryonic and total accelerations. Together, these relations offer complementary constraints that extend beyond direct curve fitting.

The purpose of the present work, denoted H1.5, is to subject the frozen H1 outputs to such secondary empirical tests. Crucially, H1.5 introduces no new modeling elements. All numerical results analyzed here are derived from precomputed H1 rotation curves treated as fixed inputs. No refitting, optimization, adaptive response, or modification of the underlying model is performed.

It is important to emphasize what H1.5 is not. This work does not propose a new dynamical model, does not attempt to improve agreement with empirical relations through tuning, and does not advance claims regarding gravitational lensing, relativistic effects, or cosmological implications. Instead, H1.5 is strictly diagnostic in scope. Its aim is to delineate the empirical domain of validity of the frozen H1 construction by confronting it with the strongest observational regularities in disk galaxy dynamics.

By explicitly mapping both successes and failures, H1.5 provides a transparent reference point for evaluating the behavior of nonlocal baryon–coupled phenomenological models under global and local empirical constraints.

2 Data and Frozen Pipeline

This work uses the SPARC (Spitzer Photometry and Accurate Rotation Curves) database, which provides high-quality rotation curves and baryonic mass models for disk galaxies across a wide range of masses, morphologies, and surface brightnesses [? ?]. The full SPARC sample employed here consists of 175 galaxies, each with resolved rotation curve measurements and associated stellar and gas mass components.

The diagnostic analyses presented in this paper do not involve any new fitting, tuning, or modification of the underlying dynamical model. Instead, they operate exclusively on a fixed set of precomputed rotation curve outputs generated by the H1 model. These outputs were produced prior to the present study and are treated here as fixed inputs.

For each galaxy, the frozen H1 outputs include the total modeled rotation curve and its baryonic decomposition as a function of radius. All quantities used in the BTFR and RAR diagnostics are derived directly from these stored results. No model parameters were altered, re-optimized, or re-estimated at any stage of the H1.5 analysis.

The H1 numerical pipeline is explicitly frozen: the convolution kernel, grid resolution, normalization procedures, and fitting logic are identical to those used in the archived H1 release. The present work therefore constitutes a purely diagnostic extension, designed to test how the fixed H1 outputs behave when confronted with independent empirical scaling relations.

Reproducibility is a core requirement of this study. The complete frozen H1 outputs, together with the analysis scripts used to construct the BTFR and RAR diagnostics, are publicly available through a versioned GitHub repository (<https://github.com/VitorFigueiredoResearch/H1-5-Secondary-Tests>) and an archived Zenodo record associated with the H1 model. All figures and numerical values reported in this paper can be regenerated directly from these materials without manual intervention.

We emphasize that the results presented below should be interpreted strictly as properties of the frozen H1 model. Any deviations from empirical relations reflect the behavior of that fixed construction and are not the result of post hoc adjustments or adaptive corrections introduced in the present analysis.

3 Baryonic Tully–Fisher Relation Diagnostic

The Baryonic Tully–Fisher Relation (BTFR) provides a global empirical connection between the total baryonic mass of a disk galaxy and its characteristic rotational velocity. As a diagnostic test, the BTFR probes whether a dynamical model preserves large-scale mass–velocity scaling, independent of detailed radial structure.

In this work, the BTFR is evaluated using only frozen outputs from the H1 model. No fitting or optimization is performed. For each galaxy in the SPARC sample, the characteristic velocity V_{flat} is extracted directly from the outer region of the H1-predicted rotation curve, following the same operational definition applied uniformly across the sample. The corresponding baryonic mass M_b is taken from the SPARC mass models without modification.

Figure 1 shows the resulting BTFR diagnostic, plotting $\log_{10}(V_{\text{flat}})$ against $\log_{10}(M_b)$ for all galaxies in the sample. The H1-predicted velocities exhibit a clear monotonic correlation with baryonic mass, recovering the expected global scaling behavior across more than three orders of magnitude in M_b .

No attempt is made here to refit the slope or normalization of the relation. The purpose of this diagnostic is not to optimize agreement with the observed BTFR, but to assess whether the frozen H1 construction preserves the existence and overall structure of the mass–velocity relation. Within this restricted scope, the BTFR is broadly reproduced: galaxies of higher baryonic mass systematically correspond to higher characteristic velocities in the H1 outputs.




Figure 1: Baryonic Tully–Fisher Relation diagnostic using frozen H1 outputs. The characteristic velocity V_{flat} is extracted from the outer regions of the H1-predicted rotation curves and plotted against the SPARC baryonic mass M_b . No fitting or parameter optimization is performed.

Scatter is present, particularly at the low-mass end, and reflects the fixed nature of the model rather than any adaptive response to individual systems. This behavior is expected in a framework that applies a single, frozen dynamical prescription across a heterogeneous galaxy population.

The BTFR diagnostic therefore indicates that the H1 model retains the global baryonic scaling relation at the level of overall structure, even in the absence of tuning or feedback. This result contrasts with the more stringent local constraints imposed by acceleration-based relations, which are examined separately in the following section.

4 Radial Acceleration Relation Diagnostic

The Radial Acceleration Relation (RAR) provides a local, pointwise empirical correlation between the observed total gravitational acceleration g_{obs} and the baryonic acceleration g_{bar} within disk galaxies. Unlike the BTFR, which probes global mass–velocity scaling, the RAR imposes a significantly more stringent constraint on the internal structure of galaxy rotation curves.

In this work, the RAR is evaluated strictly as a diagnostic test applied to the frozen H1 outputs. For each radial sample point in the SPARC rotation curves, the baryonic acceleration g_{bar} is computed directly from the SPARC baryonic mass models, while the observed acceleration g_{obs} is derived from the H1-predicted total rotation velocity at the same radius. No fitting, smoothing, or reweighting is applied.

The full diagnostic sample consists of 5250 individual radial points across the 175 SPARC galaxies. Figure 2 shows the resulting distribution in the $(g_{\text{bar}}, g_{\text{obs}})$ plane, together with the Newtonian baseline and the empirical RAR reference relation derived from SPARC data.

Globally, the frozen H1 outputs do not reproduce a tight RAR. The overall scatter across the full sample is substantial, with an RMS scatter of 0.539 dex and a median absolute deviation of 0.309 dex [?]. This level of scatter exceeds that observed in the empirical RAR and indicates that the fixed H1 construction does not enforce a universal local acceleration law.

To investigate whether this behavior is uniform across galactic radii, the RAR scatter was further decomposed by normalized radial position. Each radial point was assigned a fractional radius r_{frac} , defined relative to the maximum sampled radius of the corresponding galaxy, and grouped into inner ($r_{\text{frac}} < 0.3$), intermediate ($0.3 \leq r_{\text{frac}} < 0.7$), and outer ($r_{\text{frac}} \geq 0.7$) regimes.

The resulting radial decomposition reveals a clear and structured pattern. In the inner regions, the RAR scatter is largest, with an RMS scatter of 0.764 dex and a median deviation of 0.639 dex. The intermediate regime shows reduced but still substantial scatter, with an RMS of 0.435 dex and a median deviation of 0.306 dex. In contrast, the outer regions exhibit the tightest correspondence, with an RMS scatter of 0.364 dex and a median deviation of 0.181 dex [?].

This systematic reduction in scatter with increasing radius indicates that the RAR failure of the frozen H1 model is not uniform. Instead, deviations are most pronounced in the inner regions, where baryonic complexity and local structural features dominate, while the outer disk shows comparatively improved alignment between g_{bar} and g_{obs} .

As a final consistency check, the full RAR assembly was independently verified at the single-galaxy level. For the well-studied galaxy NGC 3198, the values of g_{bar} and g_{obs} extracted from the assembled RAR dataset were recomputed directly from the original rotation curve decomposition. Agreement was found at machine precision, with maximum absolute differences below 10^{-12} in both quantities, confirming the internal consistency of the diagnostic pipeline [?].

Taken together, these results demonstrate that the frozen H1 model does not reproduce the empirical RAR as a universal local relation. However, the structured radial dependence of the




Figure 2: Global RAR diagnostic using frozen H1 outputs. The observed acceleration g_{obs} is derived from the H1-predicted rotation curves, while g_{bar} is computed from the SPARC baryonic mass models. The empirical RAR reference and Newtonian baseline are shown for comparison.

scatter suggests that the deviation is systematic rather than random, with improved correspondence emerging preferentially in the outer, low-acceleration regions of disk galaxies.

5 Sanity and Consistency Checks

Given the diagnostic nature of this study, particular care was taken to verify that the BTFR and RAR results arise from the frozen H1 outputs alone and are not influenced by numerical inconsistencies, circular definitions, or data-handling artifacts. Several independent checks were therefore performed to validate the internal integrity of the analysis pipeline.

5.1 Single-Galaxy Reconstruction

As a direct numerical verification, the RAR assembly was independently reconstructed for a representative galaxy, NGC 3198, which has well-sampled rotation curve data and is frequently used as a benchmark system. For this galaxy, the baryonic acceleration g_{bar} and the total acceleration g_{obs} were recomputed directly from the original SPARC rotation curve decomposition and the corresponding frozen H1 velocity outputs at each sampled radius.

These independently recomputed values were then compared point-by-point with the entries

extracted from the assembled RAR dataset used in Section 4. Agreement was found at machine precision, with maximum absolute differences below 10^{-12} for both g_{bar} and g_{obs} . This confirms that the RAR assembly faithfully reflects the underlying rotation curve data and does not introduce numerical distortion during post-processing.

5.2 Unit Consistency and Numerical Stability

All accelerations are computed and stored internally in physical units of ms^{-2} , with logarithmic transformations applied only at the plotting and diagnostic stages. No rescaling or unit conversion is performed during the construction of the BTFR or RAR datasets. Internal checks confirm consistent unit handling across all analysis scripts.

The diagnostic results were also verified to be stable against numerical resolution and ordering effects. Reassembly of the RAR dataset with shuffled galaxy ordering and independent script execution produced identical statistical outcomes within numerical precision, indicating that the reported scatter values are not sensitive to data traversal or aggregation order.

5.3 Exclusion of Circularity

A potential concern in acceleration-based diagnostics is circularity, whereby the same quantity might implicitly enter both axes of a correlation. This is explicitly avoided here. The baryonic acceleration g_{bar} is computed solely from the SPARC baryonic mass models, while the total acceleration g_{obs} is derived exclusively from the H1-predicted rotation velocity. No observed rotation curve data enter the calculation of g_{obs} , and no H1 model outputs enter the computation of g_{bar} .

As a result, the RAR diagnostic probes a genuine comparison between independent baryonic inputs and the fixed dynamical response encoded in the H1 model.

5.4 Sample Completeness

All galaxies included in the frozen H1 release are retained in the BTFR and RAR diagnostics. No systems are excluded based on morphology, mass, surface brightness, or diagnostic outcome. The full set of 175 SPARC galaxies and 5250 radial sample points therefore contribute to the reported statistics, ensuring that the results reflect the global behavior of the frozen model rather than a selectively curated subset.

Taken together, these checks confirm that the diagnostic results presented in Sections 3 and 4 are numerically robust, internally consistent, and free from post hoc adjustments. Any deviations from empirical scaling relations therefore reflect intrinsic properties of the frozen H1 construction rather than artifacts of the analysis procedure.

6 Interpretation of Diagnostic Outcomes

The BTFR and RAR diagnostics probe fundamentally different aspects of galaxy dynamics. The BTFR constrains a global, integrated relation between total baryonic mass and a characteristic velocity, while the RAR imposes a local, pointwise correspondence between baryonic and total accelerations at each radius. The contrasting outcomes reported in Sections 3 and 4 therefore reflect the differing levels of structural constraint imposed by these relations.

The preservation of a clear BTFR-like trend in the frozen H1 outputs indicates that the model encodes a robust large-scale coupling between baryonic mass distribution and overall rotational support. This behavior is consistent with the nonlocal construction of H1, in which the dynamical response at a given radius reflects an integrated influence of the baryonic mass profile rather than a strictly local dependence. As a result, global mass–velocity scaling emerges naturally even in the absence of tuning or feedback.

By contrast, the failure to reproduce a tight RAR highlights the limitations of applying a single, fixed dynamical prescription to local acceleration constraints. The empirical RAR effectively requires a near-universal mapping between g_{bar} and g_{obs} across a wide range of radii, environments, and internal galaxy structures. Such a requirement is substantially more restrictive than the existence of a global BTFR.

The radial decomposition of the RAR scatter provides further insight into this distinction. The largest deviations occur preferentially in the inner regions of galaxies, where baryonic distributions are complex and local structural features dominate the dynamical landscape. In these regimes, a frozen nonlocal response struggles to accommodate the diversity of central mass concentrations using a single prescription. In contrast, the outer regions exhibit systematically reduced scatter, indicating improved correspondence between baryonic and total accelerations where the mass distribution is smoother and local structural complexity is diminished.

This behavior suggests that the RAR deviation of the frozen H1 model is not stochastic but systematic. The model preserves global coherence while lacking the flexibility required to enforce a universal local acceleration law across all radii. From a diagnostic perspective, this outcome is expected: a framework designed to encode nonlocal baryonic influence without adaptivity or feedback is naturally better suited to reproducing integrated scaling relations than pointwise acceleration constraints.

Importantly, no attempt is made here to remedy these deviations. The purpose of the H1.5 analysis is not to improve agreement with the empirical RAR, but to delineate the domain of validity of the frozen H1 construction. The results therefore identify, rather than resolve, the specific conditions under which the model succeeds and fails.

In this sense, the combined BTFR and RAR diagnostics serve as complementary probes. Together, they demonstrate that global baryonic scaling can be preserved even when local acceleration correspondence breaks down, underscoring the distinction between integrated and pointwise empirical constraints in disk galaxy dynamics.

7 Limits of the Present Framework

The results presented in this work should be interpreted strictly within the limits of the frozen H1 construction. Several important restrictions define the scope of the present analysis.

First, the H1 model is fully fixed. No parameters are tuned, adjusted, or optimized in response to the BTFR or RAR diagnostics. The convolution kernel, normalization, numerical resolution, and fitting logic are identical to those of the archived H1 release. Consequently, the diagnostic outcomes reported here reflect intrinsic properties of the frozen model rather than any adaptive response.

Second, the present framework contains no mechanism for local feedback or environment-dependent modification of the dynamical response. All galaxies are treated uniformly under the same prescription, regardless of mass, morphology, surface brightness, or internal structure. This limitation is particularly relevant for local acceleration relations, which impose stronger pointwise constraints than global scaling laws.

Third, the analysis is purely kinematic. No claims are made regarding gravitational lensing, spacetime geometry, or relativistic effects. The H1.5 diagnostics are restricted to rotation curve-based observables derived from disk galaxy dynamics, and no inference beyond this domain is attempted.

Fourth, the present work does not introduce new physics or propose extensions to the H1 model. While the diagnostic results clearly identify conditions under which the frozen construction succeeds and fails, no corrective mechanisms, additional degrees of freedom, or alternative formulations are explored here.

Finally, all conclusions are contingent on the SPARC dataset and the specific operational definitions employed for characteristic velocities and accelerations. Although these choices are standard and consistently applied, the results should not be extrapolated beyond the tested domain.

These limitations are not shortcomings of the analysis but defining features of its diagnostic purpose. By explicitly constraining the scope of H1.5, the present work aims to provide a clear and reproducible mapping of the frozen model’s empirical behavior without overinterpretation.

8 Outlook

The diagnostic results presented in this work provide a clear and constrained assessment of the empirical behavior of the frozen H1 model. By confronting the model with both the BTFR and the RAR, the present analysis delineates the regimes in which global baryonic scaling is preserved and those in which local acceleration correspondence breaks down.

These findings motivate, but do not themselves constitute, the development of extended frameworks. In particular, any attempt to address the limitations identified here would require additional dynamical freedom beyond the fixed construction examined in H1.5. Such considerations

lie outside the scope of the present work.

Future investigations may explore whether models that allow for adaptive or environment-sensitive responses can reconcile global scaling relations with local acceleration constraints. Likewise, tests involving gravitational lensing or relativistic observables may provide complementary information on the broader consistency of baryon-coupled dynamical frameworks. No such analyses are undertaken here.

The purpose of H1.5 is therefore complete. It establishes a transparent diagnostic baseline against which future developments can be evaluated, and it provides a reproducible reference point for assessing how phenomenological models behave when confronted with the strongest empirical relations in disk galaxy dynamics.