

USN

--	--	--	--	--	--	--	--	--	--

RV COLLEGE OF ENGINEERING®
 (An Autonomous Institution Affiliated to VTU)
 III Semester B. E. Examinations April-2024
 Common to AS, BT, CH, IEM, ME

STATISTICS, LAPLACE TRANSFORM AND NUMERICAL METHODS

Time: 03 Hours

Maximum Marks: 100

Instructions to candidates:

1. Answer all questions from Part A. Part A questions should be answered in first three pages of the answer book only.
2. Answer FIVE full questions from Part B. In Part B question number 2 is compulsory. Answer any one full question from 3 and 4, 5 and 6, 7 and 8, 9 and 10.
3. Use of scientific calculator and hand book for mathematics is permitted.


PART-A

M CO BT

1	1.1	If the first four moments of a distribution about the value 25 of the variables are -3, 11, -29 and 71, then the second moment about the mean is _____.	02	2	1
	1.2	The equations of regression lines are $y = 0.5x + 1$ and $x = 0.4y + 3$. The correlation coefficient is _____.	02	2	2
	1.3	The real part of the complex function $f(z) = \log_e z$ in cartesian co-ordinate system is _____.	02	1	1
	1.4	Taylor's series expansion of $\cos z$ about the point $z = \frac{\pi}{4}$ is _____.	02	1	2
	1.5	The Laplace transform of the signal $f(t) = t + \cos 2t$ is _____.	02	1	1
	1.6	Region of convergence of $L[a^t]$ to hold good is _____.	02	1	1
	1.7	$L^{-1}\left[\frac{1}{4s-1}\right] =$ _____.	02	2	1
	1.8	If $F(s) = \frac{1}{s^2+4s+4}$, then the inverse Laplace transform of $F(s) =$ _____.	02	1	2
	1.9	The equation $u_{xx} + u_{yy} - 4u_x - 6u_y = 3$ is classified as _____.	02	1	1
	1.10	For the one dimensional wave equation $u_{xx} = u_{tt}$ with the boundary conditions $u(0, t) = 0, u(1, t) = 0, u_t(x, 0) = 0$ and $u(x, 0) = x - x^2$, the value of $u(0.25, 0.25)$ by taking $h = 0.25$ and $k = 0.25$ is _____.	02	2	2

PART-B

2	a	From the following frequency distribution, compute first three moments about the mean.																	
		<table><tr><td>x</td><td>1</td><td>2</td><td>3</td><td>4</td><td>5</td><td>6</td></tr><tr><td>y</td><td>1</td><td>6</td><td>13</td><td>25</td><td>30</td><td>22</td></tr></table>	x	1	2	3	4	5	6	y	1	6	13	25	30	22			
x	1	2	3	4	5	6													
y	1	6	13	25	30	22													
	b	A simply supported beam carries a concentrated load P at its mid-point. Corresponding to various values of P , the maximum deflection Y is measured and is given in the following table.	05	3	2														
		<table><tr><td>P</td><td>100</td><td>120</td><td>140</td><td>160</td><td>180</td><td>200</td></tr><tr><td>Y</td><td>0.45</td><td>0.55</td><td>0.60</td><td>0.70</td><td>0.80</td><td>0.85</td></tr></table>	P	100	120	140	160	180	200	Y	0.45	0.55	0.60	0.70	0.80	0.85			
P	100	120	140	160	180	200													
Y	0.45	0.55	0.60	0.70	0.80	0.85													
		Fit a law of the form $Y = a + bP$ and hence estimate Y when P is 150.	06	2	3														

c		Calculate the rank correlation coefficient from the following data:																					
		<table><tr><td>x</td><td>81</td><td>78</td><td>73</td><td>73</td><td>69</td><td>68</td><td>62</td><td>58</td></tr><tr><td>y</td><td>10</td><td>12</td><td>18</td><td>18</td><td>18</td><td>22</td><td>20</td><td>24</td></tr></table>	x	81	78	73	73	69	68	62	58	y	10	12	18	18	18	22	20	24	05	3	3
x	81	78	73	73	69	68	62	58															
y	10	12	18	18	18	22	20	24															
3	a	Determine the analytic function whose real part is $u = e^x(x \cos y - y \sin y)$ by Milne-Thomson method. Also find its harmonic conjugate v .	08	2	2																		
	b	Obtain the Laurent's series expansion for $f(z) = \frac{1}{(z+1)(z+3)}$ in the region i) $1 < z < 3$ ii) $ z > 3$ iii) $ z < 1$	08	2	2																		
		OR																					
4	a	Find the analytic function $f(z) = u + iv$ where $u - v = 2xy + x^2 - y^2 + x - y$.	08	2	2																		
	b	Evaluate the integral $\int_C \frac{e^{2z}}{(z+1)(z-2)^2}$ where C is the circle $ z = 3$ using Cauchy's residue theorem.	08	2	2																		
5	a	A periodic function $f(t)$ of period $2a, a > 0$ is defined by $f(t) = \begin{cases} K & 0 \leq t \leq a \\ -K, & a < t \leq 2a \end{cases}$ Show that $L\{f(t)\} = \frac{K}{s} \tanh\left(\frac{as}{2}\right)$	08	2	2																		
	b	Obtain the Laplace transform of $\frac{e^{-t}-e^{7t}}{t} + \left(2\sqrt{t} - \frac{3}{\sqrt{t}}\right)^3$	08	3	3																		
		OR																					
6	a	Express the following in terms of unit step function and hence find the Laplace transform of $f(t) = \begin{cases} 3t^2 & 0 < t < 3 \\ 5t & t > 3 \end{cases}$	08	2	2																		
	b	Find $L\left\{e^{-t} \left(\int_0^t t \sin 4t \, dt\right) + \cos 2t \sin 5t\right\}$	08	3	3																		
7	a	The signal in frequency domain is given by $\frac{s}{(s-3)(s^2+4)}$. Determine the transformation in time domain.	08	3	2																		
	b	Employing convolution theorem, find inverse Laplace transform of $\frac{1}{(s^2+a^2)^2}$	08	4	3																		
		OR																					
8	a	Evaluate $L^{-1}\left[\frac{s+1}{s^2-s+1}\right]$	08	3	2																		
	b	Solve the differential equation using Laplace transform given $\frac{d^2y}{dt^2} - 3\frac{dy}{dt} + 2y = 12e^{-t}$ when $y(0) = 2$ and $y'(0) = 6$.	08	4	3																		
9	a	Determine the value of $u(x, y)$ satisfying the Laplace equation $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$ at the pivotal points of a square region with boundary values as shown in Fig 9a.  Fig 9a																					
		Carry out 3 iterations.	08	4	2																		

10	b	Employ Bendre-Schmidth method with $h = 1$ to find the values of $u(x, t)$ upto $t = 1$ for the equation $2 \frac{\partial^2 u}{\partial x^2} = \frac{\partial u}{\partial t}$, $0 < x < 4, t > 0$ under the conditions $u(x, 0) = x(4 - x), \quad 0 < x < 4$ $u(0, t) = u(4, t) = 0, \quad t \geq 0$	08	3	3
		OR			
	a	The temperature distribution $u(x, t)$ is modeled as the one-dimensional heat equation $\frac{\partial u}{\partial t} = \frac{4}{5} \frac{\partial^2 u}{\partial x^2}$, $0 < x < 100, t > 0$ subject to the boundary conditions $u(0, t) = 0 = u(100, t)$, $t \geq 0$ and the initial condition $u(x, 0) = \begin{cases} x, & 0 \leq x \leq 50 \\ 100 - x, & 50 \leq x \leq 100 \end{cases}$ Determine the temperature distribution $u(x, t)$ by taking $h = 20$ and $k = 10$ using Schmidt relation for 5 levels.	08	3	2
	b	Solve $\frac{\partial^2 u}{\partial t^2} = 16 \frac{\partial^2 u}{\partial x^2}$ taking $h = 1$ in five steps given that $u(x, 0) = x^2(5 - x)$, $u_t(x, 0) = 0$, $u(0, t) = u(5, t) = 0$	08	4	3