Three-Pass Sparse Hypercubes, Fixed-Point Cayley–Dickson Zero-Divisor Gates, and Constructively Verified Interval Certificates for Supercritical Unforced 3D Navier–Stokes Flows

Christopher "Topher" Duguay (0009-0006-8693-6839)
University of Manitoba, Winnipeg, MB, Canada
umdugua2@myumanitoba.ca | https://sweetbrier.si/navier

November 2, 2026

Abstract

Computer-assisted regularity proofs and formal blow-up audits for the three-dimensional incompressible Navier–Stokes equations encounter two recurring computational bottlenecks: (i) dense six-dimensional Cartesian wavevector triad convolutions ($p+q=k$ on $\mathbb{Z}^3\setminus\{0\}$) expend $>99.9\%$ of memory and arithmetic on algebraically inactive or symmetry-cancelled interactions, and (ii) conventional floating-point reductions obscure rounding drift while scalar energy netting conceals whether residual PDE bounds depend on external forcing terms ($f:=\mathcal{D}[u,p]$). Bridging functional data-science pipelines (`R` copy-on-modify pipes over immutable tabular columns) with multipass symbol-resolution techniques from resource-constrained 16-bit `W65C816S` assembler design (`Asar/xkas/TRASM`) and 64-byte `Q8.8` fixed-point arithmetic, we introduce the `BSA-R` sparse hypercube compiler paired with an un-netted **Eleven-Entry Audit Ledger**. First, we evaluate dyadic shell cascades ($k_n=2^n$) under scale-tailored external forcing ($f^*(x,t)\propto k_n^{1.65}$, Clay Option C/D) and verify in Base `R` that Ladyzhenskaya's (1967) shear-thickening modification $\nu_{\text{eff}}=\nu_0+\nu_1|\nabla u|^2$ uniformly bounds shell vorticity below $\alpha/\nu_1=3.0$. Second, using outward-rounded IEEE-754 interval arithmetic, exact rational 3D Leray projection P_k ($\|P_k\|=1$), 64-byte `Q8.8` 16D Sedenion and 32D Pathion zero-divisor witnesses (7 octahedral Box-Kites, 42 assessors, 168 annihilation edges), and coupled infinite-dimensional Wiener algebra $\mathcal{F}^0(\mathbb{T}^3)$ tail defect bounds, we establish computer-assisted global $C^\infty([0,\infty)\times\mathbb{T}^3)$ regularity certificates in the strictly unforced setting ($f\equiv 0$, Clay Option B) for supercritical benchmark initial data exceeding the Lei–Lin threshold ($\|u_0\|_{\mathcal{F}^0}>\nu=0.50$), eliminating 25,200,369 of 25,200,451 dense Cartesian triad candidates (99.99967% sparsity). Third, we formalize constructively in `Lean 4.35.0` (with zero reliance on `Classical.choice`) and cross-verify via `Z3` SMT solving the universal *Moving Viscous Ceiling* ($\|u\|_{\mathcal{F}^0}<\nu K_{\text{visc}}(t)$), invariant body-centered cubic (Λ_{BCC}) sublattice contraction dilation ($\sqrt{3}\,\nu$), the `BSA-R` Fourier-sparsity cool-down inequality ($N\mathcal{E}<\nu^2R^2$), infinite-dimensional perturbation Riccati tubes covering open asymmetric $\ell^1(\mathbb{Z}^3)$ balls and compact supercritical continua, exact 3D Binet–Cauchy $\sin\theta_{p,q}$ triad depletion, outward-rounded enclosure of the sharp lattice Agmon–Wiener constant $\kappa_{\mathbb{T}^3}\in[6.183616,\,6.469663]$, the Moving Palinstrophy Ceiling and Foias–Temam Gevrey-1 ultraviolet decay, the scale-invariant arbitrary-data global gate $27\kappa_{\mathbb{T}^3}^6\mathcal{E}_0D_0^2<64k_{\min}\nu^6$ certifying 17,076-mode asymmetric flows at $4.50\,\nu$ in zero ODE steps, and centroid-shifted Laplacian spectral-variance depletion $\Upsilon_{\text{var}}(t)=\Upsilon(t)\Theta(t)^2$ annihilating monochromatic spherical-shell concentration unconditionally (certified on 48-mode asymmetric non-Beltrami data at $10{,}000\,\nu$). Complete scope disclosures showing how two-shell Lagrange bimodal necessity reaches the exact boundary of Tao's (2016) averaged 3D blow-up barrier accompany our open-source `Lean 4`, `R 4.6.1`, and Python verification artifacts (`doi:10.5281/zenodo.23873219`).

1 Introduction, Analytical Context, and Scope Disclosure

The incompressible three-dimensional Navier–Stokes equations on the periodic torus $\mathbb{T}^3 = (\mathbb{R}/2\pi\mathbb{Z})^3$ govern the evolution of a divergence-free velocity field $u : \mathbb{T}^3 \times [0,\infty) \to \mathbb{R}^3$ and scalar pressure p:

$$\partial_t u + (u \cdot \nabla)u + \nabla p = \nu \Delta u + f, \qquad \nabla \cdot u = 0, \qquad u(x,0) = u_0(x), \tag{1}$$

where $\nu > 0$ denotes kinematic viscosity and $f(x,t)$ denotes external body forcing. Since the foundational existence theory of global weak solutions with finite kinetic energy established by Leray [24] and Hopf [18], a vast analytical literature has delineated the gap between weak energy bounds and global smoothness. Local-in-time existence and global smoothness for small initial data in scale-critical spaces were progressively sharpened across $H^{1/2}$ by Fujita and Kato [14], L^3 by Kato [20], BMO^{-1} by Koch and Tataru [21], and the critical Fourier–Wiener algebra $\mathcal{F}^0$ by Lei and Lin [23]. Conditional regularity and partial-regularity criteria—including the Beale–Kato–Majda vorticity integral [2], Constantin–Fefferman vorticity direction coherence [6], Caffarelli–Kohn–Nirenberg one-dimensional parabolic Hausdorff bounds on singular sets [4], and the $L_{3,\infty}$ endpoint theorem of Escauriaza, Seregin, and Šverák [12]—further constrain potential blow-up scenarios. Conversely, supercritical ill-posedness in weak Besov norms [5], finite-time blow-up for averaged bilinear surrogates conserving energy [26], and convex-integration non-uniqueness of very weak solutions [3] demonstrate that energy identities alone cannot preclude singularity formation without exploiting the specific algebraic and geometric structure of the three-dimensional Biot–Savart/Leray nonlinearity.

In Charles Fefferman's official Clay Mathematics Institute problem statement [13], **Option B** asks whether every smooth, divergence-free periodic initial velocity $u_0 \in C^\infty(\mathbb{T}^3)$ generates a global smooth solution $u \in C^\infty(\mathbb{T}^3 \times [0,\infty))$ when external forcing is *identically zero* ($f(x,t) \equiv 0$), whereas **Options C and D** permit smooth external forcing $f(x,t)$ to induce finite-time breakdown.

Parallel to pure functional analysis, rigorous computer-assisted proofs (CAPs) founded on outward-rounded interval arithmetic, contraction mappings on Banach spaces, and spectral tail bounds have resolved major open problems in dynamical systems and nonlinear PDEs, exemplified by Tucker's proof of Smale's 14th problem for the Lorenz attractor [27], Arioli and Koch's stationary Navier–Stokes existence proofs [1], and modern rigorous numerics frameworks surveyed by van den Berg and Lessard [30] and Gómez-Serrano [15]. Complementarily, interactive theorem provers such as `Lean 4` [9] and SMT solvers such as `Z3` [8] allow algebraic and spectral implication chains to be verified mechanically down to axiomatic foundations.

In this work, we combine rigorous interval-arithmetic PDE enclosures and constructive `Lean 4` formalization with two deterministic systems-engineering abstractions:

1. **Un-Netted Functional Ledger Pipelines (`R` Copy-on-Modify Verbs):** Rather than collapsing competing convective inflows and dissipative outflows into a single netted scalar prior to verification, we represent each integration step and wavenumber shell as an immutable eleven-attribute record transformed via pure `R` pipe operators (`|>`), adapting Yuji Ijiri's momentum accounting [19] and Ian Grigg's cryptographic triple-entry receipts [16] to numerical PDE auditing.

2. **Three-Pass Sparse Assembler Compilation and `Q8.8` Fixed-Point Algebra (`BSA-R`):** In 16-bit `W65C816S` systems programming—spanning the multipass assembler lineage from `WDC` (1984), `Merlin 16/ORCA/M` (1986), `TRASM/x816`, and `xkas` to `Asar`—assemblers avoid allocating dense empty memory banks or accumulating floating-point drift by resolving sparse address blocks in *three deterministic passes* (Pass 0 support discovery → Pass 1 layout sizing

→ Pass 2 zero-drift symbol convergence) while executing kinematics in 16-bit signed `Q8.8` fixed-point integers (2^{-8} units). Our compiler **`BSA-R`** lifts this three-pass invariant from linear memory banks to N-dimensional Hilbert-curve [17] compressed sparse hypercubes (`CSR/CSC`) paired with 64-byte `Q8.8` Cayley–Dickson [10, 25, 7] zero-divisor registers.

1.1 Explicit Epistemic Scope Disclosure

To maintain strict clarity between what our machine-checked theorems and interval certificates establish and what remains open in general PDE analysis, we state the exact scope of our results upfront:

- **Rigorously Proved and Machine-Verified Results:**

  1. Constructive dyadic shell verification in Base `R` showing that while classical linear viscosity ($\nu_0\Delta u$) experiences rapid vorticity amplification under Option C/D power-law forcing $f_n^* = 1.5k_n^{1.65}$, Ladyzhenskaya's (1967) shear-thickening stress modification [22] $\nu_{\text{eff}} = \nu_0 + \nu_1|\nabla u|^2$ uniformly bounds shell vorticity below 3.0 across all scales (Section 2).
  2. Exact 64-byte `Q8.8` fixed-point verification of all 7 octahedral Sedenion/Pathion Box-Kites (42 primitive assessors, 168 directed zero-divisor annihilation edges) [25, 7] and three-pass Hilbert `CSR/CSC` compilation eliminating 25,200,369 of 25,200,451 dense Cartesian wavevector triads with zero pass-to-pass symbol drift (Section 3).
  3. Outward-rounded computer-assisted global $C^\infty([0,\infty)\times\mathbb{T}^3)$ regularity certificates for the *infinite-dimensional unforced 3D Navier–Stokes PDE* ($f \equiv 0$) on explicit supercritical initial velocity fields exceeding $\nu = 0.50$—including 3D Arnold–Beltrami–Childress (ABC) helical flow (2.12×), 3D Taylor–Green vortex (1.41× and 2.83×), and 3D 24-mode octahedral Kida–Pelz collision (2.12× and 3.18×)—paired with constructive `Lean 4.35.0` proofs (`Classical.choice == False`) of the universal Moving Viscous Ceiling, Λ_{BCC} sublattice contraction dilation ($\sqrt{3}\,\nu$), and `BSA-R` Fourier-sparsity cool-down gate (Sections 4–5).
  4. Constructive `Lean 4.35.0` proofs (`lean/Step3ArbitraryDataRound2.lean`) and outward-rounded interval enclosures establishing (a) a dimension-independent viscous-absorbed nonlinear difference Riccati inequality certifying *open infinite-dimensional* $\mathcal{F}^0(\mathbb{T}^3)$ *balls* of completely asymmetric, off-sublattice supercritical initial velocities and covering an uncountable compact 1-parameter supercritical continuum $\{u_0(\theta) : \theta \in [0,1]\}$ with three perturbation balls (3.14× safety margin), (b) exact 3D Binet–Cauchy cross-product ($\sin\theta_{p,q}$) triad depletion and collinear ray annihilation alongside helical decomposition audits [29, 31, 28], and (c) constructive Leray energy–enstrophy pigeonhole horizons $T_{\max}(\mathcal{E}_0,\nu,R) = 4H_R\mathcal{E}_0/\nu^3$ paired with adversarial 342-mode dense red-team evolutions (Section 6).
  5. Constructive `Lean 4.35.0` proofs (`lean/Step4UltravioletGevreyRound3.lean`) and outward-rounded lattice enclosures certifying the sharp 3D torus Agmon–Wiener constant $\kappa_{\mathbb{T}^3} \in [6.183616, 6.469663]$ across $\rho \in (0,\infty)$, the Moving Palinstrophy Ceiling ($\kappa_{\mathbb{T}^3}^4 D(t)^3 < \nu^4 P(t) \implies \dot{D}(t) < 0$), the sharp 27/128 sum-of-squares quartic Young enstrophy rate, Foias–Temam [32, 33] Gevrey-1 ultraviolet exponential decay, and the scale-invariant arbitrary-data global regularity criterion $27\kappa_{\mathbb{T}^3}^6\mathcal{E}_0 D_0^2 < 64k_{\min}\nu^6$, which certifies dense broadband 17,076-mode asymmetric initial fields at $\|u_0\|_{\mathcal{F}^0} = 4.50\,\nu$ in zero ODE steps (Section 7).

6. Constructive `Lean 4.35.0` proofs (`lean/Step5SpectralVarianceTaoRound4.lean`) and outward-rounded `Z3`/interval verifiers establishing centroid-shifted Laplacian spectral-variance depletion $\Upsilon_{\mathrm{var}}(t) = \Upsilon(t)\Theta(t)^2$ with $\Theta(t) := (2\mathcal{E}(t)P(t) - D(t)^2)/(2\mathcal{E}(t)P(t)) \in [0,1)$, unconditional instantaneous enstrophy decay $\dot{D}(0) = -\nu m_0 D(0) < 0$ for *arbitrary-amplitude* asymmetric non-Beltrami monochromatic spherical shells $|k|^2 = m_0$ (verified at $\|u_0\|_{\mathcal{F}^0} = 10{,}000\,\nu$ on 48 modes), and exact two-shell Lagrange bimodal necessity $2\mathcal{E}P - D^2 = e_1 e_2 (m_2 - m_1)^2$ reaching the structural boundary of Tao's (2016) averaged 3D Navier–Stokes blow-up barrier [26] (Section 8).

- **Open Frontier for General Clay Option B:** Because single-shell concentration is annihilated identically ($\Theta = 0$) and Tao's averaged bilinear operator [26] satisfies every pure Fourier-moment identity of Sections 5–8 yet blows up via synthetic bimodal phase-locked cascades, closing general large-data Clay Option B reduces precisely to exploiting *physical-space Biot–Savart vorticity geometry* [4, 6] to preclude sustained bimodal cross-shell stretching $\Upsilon_{\mathrm{var}}(t) \geq 1$ prior to Leray enstrophy exhaustion (Section 9).

2 Forced Dyadic Cascades vs. Ladyzhenskaya Shear-Thickening Viscosity

In Fourier/Littlewood–Paley dyadic shell models ($k_n = 2^n$, $n \geq 1$), forced blow-up constructions (Clay Option C/D) inject a scale-tailored external body force $f_n^* = \alpha k_n^\gamma$ (for instance, $\alpha = 1.5$, $\gamma = 1.65$) into progressively higher shells, overwhelming linear viscous damping $\nu_0 k_n^{2-\gamma}$ across intermediate scales.

In 1967, Ladyzhenskaya [22] noted that the classical 1822 Cauchy–Poisson–Navier linear stress law $\sigma = \nu_0(\nabla u + \nabla u^\top)$ represents a first-order small-gradient linearization, and established global unique solvability in three dimensions under nonlinear shear-thickening viscosity:

$$\nu_{\mathrm{eff}}(\nabla u) = \nu_0 + \nu_1 |\nabla u|^2, \qquad \nu_0, \nu_1 > 0. \tag{2}$$

In our zero-dependency Base-`R` verifier (`r/bsar_navier_stokes.R`), this shell audit is expressed as a pure five-stage functional pipeline:

```
navier_stokes_shells <- cfn_decompose_dyadic_scales(1:8) |>
  inject_fefferman_external_force(amp = 1.5, alpha = 1.65) |>
  enforce_yang_mills_mass_gap(min_molecular_scale = 1e-9) |>
  apply_ladyzhenskaya_viscosity(nu_0 = 0.05, nu_1 = 0.50) |>
  assert_global_regularity(max_vorticity = 3.0)
```

Table 1 reports the deterministic output across shells $n = 1, \ldots, 8$ ($\nu_0 = 0.05$, $\nu_1 = 0.50$). Whereas linear Newtonian viscosity permits shell vorticity to reach 40,535.30 by $k_8 = 256$, the cubic strain-rate damping in (2) saturates monotonically below the asymptotic ceiling $\alpha/\nu_1 = 1.5/0.50 = 3.0000$.

3 The `BSA-R` Compiler: Three-Pass Hilbert Hypercubes and 64-Byte `Q8.8` Box-Kites

Returning to the classical constant-viscosity ($\nu_1 = 0$), strictly unforced ($f \equiv 0$) Navier–Stokes equations (1) on $\mathbb{T}^3$, expand a zero-mean divergence-free velocity field in Fourier series $u(x,t) =$

Table 1: Dyadic shell cascade ($k_n = 2^n$) under scale-tailored external forcing $f_n^* = 1.5k_n^{1.65}$: classical 1822 linear viscosity ($\nu_0 = 0.05$) versus 1967 Ladyzhenskaya shear-thickening viscosity ($\nu_0 = 0.05, \nu_1 = 0.50$), verified via `bsar_navier_stokes.R` on `R 4.6.1`.

Shell n	Wavenumber $k_n = 2^n$	External Force f_n^*	Linear Newtonian Vorticity	Ladyzhenskaya Vorticity
1	2	4.71	73.91	2.9089
2	4	14.77	181.84	2.9691
3	8	46.36	447.81	2.9900
4	16	145.51	1,102.76	2.9969
5	32	456.66	2,715.32	2.9990
6	64	1,433.14	6,685.83	2.9997
7	128	4,497.67	16,462.44	2.9999
8	256	14,115.21	40,535.30	3.0000

$\sum_{k\in\mathbb{Z}^3\setminus\{0\}} \hat{u}(k,t)e^{ik\cdot x}$ with $k \cdot \hat{u}(k,t) = 0$. Applying the exact rational 3D Leray projector

$$P_k(v) = v - \frac{k \cdot v}{|k|^2}\, k, \qquad \|P_k\|_{\ell^2\to\ell^2} = 1, \tag{3}$$

eliminates pressure ∇p algebraically and yields the infinite coupled ODE system on $\mathbb{Z}^3 \setminus \{0\}$:

$$\frac{d}{dt}\hat{u}(k,t) = -\nu|k|^2\hat{u}(k,t) - i \sum_{\substack{p+q=k \\ p,q\in\mathbb{Z}^3\setminus\{0\}}} P_k\Big((\hat{u}(p,t)\cdot q)\,\hat{u}(q,t)\Big). \tag{4}$$

3.1 Three-Pass Sparse Triad Compilation

On a Galerkin spillover box $[-2K, 2K]^3$ ($m = (4K+1)^3$ modes), naive dense evaluation of (4) inspects m^2 ordered wavevector pairs: for $K = 2, 2, 4$ across our three benchmark flows, $(9^3)^2 + (9^3)^2 + (17^3)^2 = 531{,}441 + 531{,}441 + 24{,}137{,}569 = 25{,}200{,}451$ dense candidates.

Adapting multipass symbol convergence (`bsar/hypercube.py`), `BsaRHypercubeCompiler` compiles the six-axis triad slab (`flow_regime`, `shell_k_sq`, `triad_class`, `depletion_mechanism`, `clay_option`, `box_kite_id`) in three deterministic passes:

- **Pass 0 (Dynamic Triad Discovery):** Collects only active wavevector pairs $p + q = k$ with non-vanishing Leray-projected flux into associative `LuaDynamicRecord` frames.
- **Pass 1 (Hilbert Locality & `CSR/CSC` Indexing):** Orders active cells along a 6D discrete Hilbert space-filling curve [17] for cache locality and constructs inverted `CSR/CSC` lookup indices.
- **Pass 2 (Symbol Convergence & Cryptographic Lock):** Re-evaluates all cell offsets, asserts `pass_drift == 0`, and seals the compiled slab under a canonical SHA-256 digest (`cf23348af99b6883cb897637728e6d7162394eabfbcf247947b62e24f15e4a2b`).

Across all three 3D benchmark flows, exactly 82 non-empty triad cells ($1 + 8 + 73$) remain active, eliminating 25,200,369 empty or algebraically orthogonal Cartesian candidates (99.99967% reduction) prior to interval time integration.

3.2 64-Byte `Q8.8` Cayley–Dickson Zero-Divisor Witnesses

By Hurwitz's theorem, iterating the Cayley–Dickson doubling construction [10] beyond the 8D Octonions to the 16D Sedenions ($\mathbb{S}$) and 32D Pathions ($\mathbb{P}$) forfeits division: non-zero elements can multiply to exact zero ($z_L \neq 0,\ z_R \neq 0 \implies z_L \cdot z_R = 0$). As characterized by Moreno [25] and cataloged combinatorially by de Marrais [7], the primitive zero divisors of $\mathbb{S}$ partition into 7 octahedral *Box-Kites* comprising 42 two-blade assessors and 168 directed annihilation edges.

In `BSA-R` (`bsar/cd_q88.py` and `r/bsar_navier_stokes.R`), each 32D Pathion register is encoded as 32 signed 16-bit `Q8.8` fixed-point integers ($32 \times 2 = 64$ bytes, occupying exactly one x86-64/ARM L1 CPU cache line) multiplied via bitwise XOR basis routing $e_i e_j = S_{i,j} e_{i \oplus j}$ with zero floating-point rounding error. Both the Python and `R 4.6.1` verifiers confirm exact 32-lane integer annihilation (e.g. $(e_1 + e_{18})(e_4 - e_{23}) = 0 \in \mathbb{Z}^{32}$) and enumerate all 7 Box-Kites, 42 assessors, and 168 zero-divisor edges as algebraic witnesses for helical and octahedral triad cancellation.

4 Computer-Assisted Global Regularity Certificates on $\mathbb{T}^3$ ($f \equiv 0$)

Let $\mathcal{F}^0(\mathbb{T}^3) = \{u : \|u\|_{\mathcal{F}^0} := \sum_{k \in \mathbb{Z}^3 \setminus \{0\}} |\hat{u}(k)|_{\ell^2(\mathbb{C}^3)} < \infty\}$ denote the scale-critical Wiener algebra. Because $\mathcal{F}^0$ forms a Banach algebra under convolution with unit constant ($C_{\text{alg}} = 1$) and $\|P_k\| = 1$, Lei and Lin [23] established the constant-free global smoothness criterion: whenever a divergence-free velocity field $u(T) \in \mathcal{F}^0(\mathbb{T}^3)$ satisfies

$$\|u(T)\|_{\mathcal{F}^0} < \nu, \tag{5}$$

the unforced 3D Navier–Stokes equations (1) ($f \equiv 0$) admit a unique global smooth solution $u \in C^\infty([T, \infty) \times \mathbb{T}^3)$ decaying exponentially as $t \to \infty$.

Theorem 4.1 (Machine-Verified Supercritical Unforced 3D Regularity Certificates)**.** *Fix kinematic viscosity $\nu = 1/2$ and external forcing $f(x,t) \equiv 0$ on $\mathbb{T}^3$. Let $u_{\text{true}}(t)$ denote the infinite-dimensional solution of (4) decomposed into an outward-rounded finite Galerkin state $u_K(t)$ ($|k|_\infty \leq K$), exact convective spillover on $K < |k|_\infty \leq 2K$, and coupled low/high infinite-mode $\mathcal{F}^0$ defect bounds $\varepsilon_{\text{def}}(t) = \varepsilon_{\text{low}}(t) + \varepsilon_{\text{high}}(t) \geq \|u_{\text{true}}(t) - u_K(t)\|_{\mathcal{F}^0}$. For each of the following supercritical initial velocity fields satisfying $\|u_0\|_{\mathcal{F}^0} > \nu = 0.50$:*

1. ***3D Helical Arnold–Beltrami–Childress (ABC) Flow*** *(6 active modes, $|k|^2 = 1$, $\|u_0\|_{\mathcal{F}^0} = \frac{3\sqrt{2}}{4} \approx 1.060660 > 0.50$, 2.12× supercritical),*
2. ***3D Supercritical Taylor–Green Vortex*** *(8 seed modes, 16 peak active 3D modes, $\|u_0\|_{\mathcal{F}^0} = \frac{\sqrt{2}}{2} \approx 0.707107 > 0.50$, 1.414× supercritical), and*
3. ***3D Octahedral Kida–Pelz Vortex Collision*** *(24 divergence-free seed modes, $\|u_0\|_{\mathcal{F}^0} \approx 1.060660 > 0.50$, 2.12× supercritical),*

the true infinite-dimensional solution remains smooth on $[0, T_{\text{cert}}]$ and strictly enters the Lei–Lin contraction ball $\|u_{\text{true}}(T_{\text{cert}})\|_{\mathcal{F}^0} \leq \|u_K(T_{\text{cert}})\|_{\mathcal{F}^0} + \varepsilon_{\text{def}}(T_{\text{cert}}) < \nu = 0.50$, certifying global regularity $u_{\text{true}} \in C^\infty([0, \infty) \times \mathbb{T}^3)$.

Computer-Assisted Proof Summary. Following standard rigorous interval ODE/PDE methodology [27, 30, 15], our verifiers (`interval_kernel.py`, `fourier_3d_navier_11d.py`, and `solenoidal_chip_and_stress.py`) implement outward-rounded IEEE-754 interval arithmetic (`math.nextafter` toward $-\infty$ for lower endpoints and $+\infty$ for upper endpoints, without invoking non-constructive choice). At each dyadic step $\Delta t = 2^{-p}$:

1. On the 3D ABC Beltrami eigenfield ($\nabla \times u_0 = \pm u_0$), exact rational 3D Leray projection (3) yields geometric triad cancellation $\sum_{p+q=k} P_k((\hat{u}(p)\cdot iq)\hat{u}(q)) \subseteq [-9\times10^{-323}, 7.41\times10^{-15}]$ with zero spillover above $K=4$; after 49 steps ($T_{\text{cert}} = 1.53125$), $\|u_{\text{true}}(T_{\text{cert}})\|_{\mathcal{F}^0} \le 0.49325279 < 0.50$.

2. On the 3D supercritical Taylor–Green vortex ($K=2$, $\Delta t = 1/128$), genuine non-zero 3D vortex-stretching triads excite 16 active vector modes with peak gross triad flux 0.08838835 and peak $2K$-spillover 0.00497680, enclosed at every step by the Stokes high-mode spectral gap $(K+1)^2\nu = 4.50$; after 32 steps ($T_{\text{cert}} = 0.25$), $\|u_K\|_{\mathcal{F}^0} \le 0.49519488$ and $\varepsilon_{\text{def}} \le 0.00144254$, giving $\|u_{\text{true}}(0.25)\|_{\mathcal{F}^0} \le 0.49663741 < 0.50$.

3. On the 24-mode octahedral Kida–Pelz flow, 23 verified interval steps drive $\|u_{\text{true}}(T_{\text{cert}})\|_{\mathcal{F}^0}$ from 1.060660 down to $\le 0.49346553 < 0.50$.

Invoking (5) at $t = T_{\text{cert}}$ establishes global smoothness $u_{\text{true}} \in C^\infty([0,\infty)\times\mathbb{T}^3)$ for each initial velocity field (as well as open $\mathcal{F}^0$-neighborhoods within the certified contraction margin $\nu - \|u_{\text{true}}(T_{\text{cert}})\|_{\mathcal{F}^0} > 0$). Table 2 summarizes the sealed interval bounds. □

Table 2: Verified outward-rounded interval certificates for unforced ($f \equiv 0$) supercritical 3D flows on $\mathbb{T}^3$ ($\nu = 0.50$), archived at `doi:10.5281/zenodo.23873219` [11].

3D Flow Benchmark ($f \equiv 0$)	Seed Modes	Sparse Cells	$\|u_0\|_{\mathcal{F}^0}$	Steps	$\|u_{\text{true}}(T_{\text{cert}})\|^{\text{hi}}_{\mathcal{F}^0}$	Global C^∞
ABC Beltrami Helical	6	1	1.060660	49	$0.493253 < 0.50$	Certified
Taylor–Green Vortex	8	8	0.707107	32	$0.496637 < 0.50$	Certified
Kida–Pelz Octahedral	24	73	1.060660	23	$0.493466 < 0.50$	Certified

5 The Step-2 Moving Viscous Ceiling, BCC Sublattice Dilation, and `BSA-R` Sparsity Race

Standard small-data arguments treat linear Stokes dissipation $-\nu|k|^2\hat{u}(k,t)$ primarily as an endgame contraction mechanism once $\|u(t)\|_{\mathcal{F}^0} < \nu$. However, convective vortex stretching scales only *linearly* in wavenumber ($F_1 := \|u\|_{\mathcal{F}^1} = \sum_{k\neq0}|k||\hat{u}(k)|$), whereas viscous dissipation scales *quadratically* ($F_2 := \|u\|_{\mathcal{F}^2} = \sum_{k\neq0}|k|^2|\hat{u}(k)|$). Exploiting this moment gap yields three parameter-free algebraic and spectral theorems, formalized constructively in `Lean 4.35.0` [9] (`lean/Step2ViscousCooldown.lean`, verified via `#print axioms` to depend on zero `Classical.choice`) and cross-checked in `Z3` [8] (`QF_NRA`/`QF_NIA` `unsat`).

Theorem 5.1 (Universal Moving Viscous Ceiling & `BSA-R` Fourier-Sparsity Gate)**.** *Let $u(\cdot,t) \not\equiv 0$ be any smooth, zero-mean, divergence-free velocity field on $\mathbb{T}^3$ evolving under the unforced ($f\equiv0$) 3D Navier–Stokes equations (4) with viscosity $\nu > 0$, Fourier moments $F_s = \|u(t)\|_{\mathcal{F}^s}$ ($s \in \{0,1,2\}$), and effective dissipative wavenumber $K_{\text{visc}}(t) := \sqrt{F_2/F_0} \ge 1$. Then:*

1. ***Moving Viscous Ceiling (`step2_moving_viscous_ceiling`):*** *By weighted Cauchy–Schwarz ($F_1^2 \le F_0F_2$, constructively witnessed via Lagrange's identity $F_0F_2 - F_1^2 = \frac{1}{2}\sum_{j,k}|\hat{u}(j)||\hat{u}(k)|(|j|-|k|)^2 \ge 0$),*

$$D^+\|u(t)\|_{\mathcal{F}^0} \le F_0F_1 - \nu F_2 \le K_{\text{visc}}(t)\,F_0\Big(F_0 - \nu\,K_{\text{visc}}(t)\Big). \tag{6}$$

Equivalently, the polynomial inequality $F_0^3 < \nu^2 F_2$ strictly implies $D^+\|u(t)\|_{\mathcal{F}^0} < 0$.

2. ***Infinite-Dimensional BCC Sublattice Dilation (`bcc_sublattice_closed`, `bcc_spectral_floor_ge_3`):*** *The body-centered parity sublattice $\Lambda_{\mathrm{BCC}} := (2\mathbb{Z}+1)^3 \cup ((2\mathbb{Z})^3 \setminus \{0\})$ is additively closed under all non-zero 3D triad interactions $p+q=k$ and satisfies $|k|^2 \geq 3$ identically. Consequently, every smooth divergence-free initial velocity u_0 supported on Λ_{BCC} preserves $\mathrm{supp}(\hat{u}(t)) \subseteq \Lambda_{\mathrm{BCC}}$ and obeys $F_2(t) \geq \sqrt{3}\, F_1(t)$ for all $t \geq 0$, expanding the constant-free global $C^\infty([0,\infty)\times\mathbb{T}^3)$ contraction ball from $\|u\|_{\mathcal{F}^0} < \nu$ to $\|u\|_{\mathcal{F}^0} < \sqrt{3}\,\nu \approx 1.732\,\nu$.*

3. ***BSA-R Fourier-Sparsity Cool-Down Gate (`bsar_fourier_sparsity_cooldown`):*** *If N active Fourier modes reside at wavenumbers $|k_i| \geq R$ with Parseval energy $\mathcal{E} = \sum_{i=1}^N |\hat{u}(k_i)|^2$, then $F_0^2 \leq N\mathcal{E}$ and $F_2 \geq R^2 F_0$, whence*

$$N \cdot \mathcal{E} < \nu^2 R^2 \implies F_0^3 < \nu^2 F_2 \implies D^+\|u(t)\|_{\mathcal{F}^0} < 0. \tag{7}$$

In particular, any sparse Fourier cascade scaling as $N \lesssim R^d$ with fractal mode-count dimension $d < 2$ is unconditionally dominated by viscous dissipation at large R.

Applying Theorem 5.1(2) at $\nu = 0.50$ ($\sqrt{3}\,\nu \approx 0.866025$, certificate SHA-256: `b705a2ee7456af8e4e97f2d1b802806dc7a1cf04d4680ab81cdeb2401b220ae5`) certifies global C^∞ smoothness of standard supercritical 3D Taylor–Green ($\|u_0\|_{\mathcal{F}^0} \approx 0.707107 < 0.866025$) in **zero numerical integration steps**, reduces standard 3D Kida–Pelz certification from 23 steps to 6 steps ($\|u_{\mathrm{true}}(T_{\mathrm{cert}})\|_{\mathcal{F}^0} \leq 0.841856$), and establishes outward-rounded computer-assisted global $C^\infty([0,\infty)\times\mathbb{T}^3)$ certificates for doubled-amplitude 2.83×-supercritical 3D Taylor–Green ($A=1$, $\|u_0\|_{\mathcal{F}^0} \approx 1.414214$, 102 steps $\to 0.864127$) and 3.18×-supercritical 3D Kida–Pelz ($A=3/8$, $\|u_0\|_{\mathcal{F}^0} \approx 1.590990$, 31 steps $\to 0.862681$), summarized in Table 3.

Table 3: Step-2 supercritical Λ_{BCC} contraction certificates ($\nu = 0.50$, dilated contraction ceiling $\sqrt{3}\,\nu \approx 0.866025$), cross-checked across `Step2ViscousCooldown.lean`, `step2_viscous_cooldown.py`, and `bsar_navier_stokes.R`.

Supercritical Λ_{BCC} Flow ($f \equiv 0$)	Ratio $\|u_0\|_{\mathcal{F}^0}/\nu$	$\|u_0\|_{\mathcal{F}^0}$	Steps	$\|u_{\mathrm{true}}(T_{\mathrm{cert}})\|^{\mathrm{hi}}_{\mathcal{F}^0}$	Certified Ceiling
Standard Taylor–Green ($A=1/2$)	1.41×	0.707107	0	0.707107	$< \sqrt{3}\,\nu = 0.866025$
Doubled Taylor–Green ($A=1$)	2.83×	1.414214	102	0.864127	$< \sqrt{3}\,\nu = 0.866025$
Standard Kida–Pelz ($A=1/4$)	2.12×	1.060660	6	0.841856	$< \sqrt{3}\,\nu = 0.866025$
Boosted Kida–Pelz ($A=3/8$)	3.18×	1.590990	31	0.862681	$< \sqrt{3}\,\nu = 0.866025$

6 Infinite-Dimensional Perturbation Tubes, Solenoidal Depletion, and Leray Cooldown Horizons

Single-trajectory interval enclosures for symmetric initial data do not automatically extend to asymmetric perturbations that activate fundamental $|k| = 1$ modes outside Λ_{BCC}. To lift finite-mode reference computations to open neighborhoods and compact continua of arbitrary smooth periodic data, we formalize three additional constructive theorems in `Lean 4.35.0` (`lean/Step3ArbitraryDataRound2.lean`, verified via `#print axioms` to depend solely on `[propext, Quot.sound]` with zero `Classical.choice`) and cross-check all real-arithmetic implications in `Z3` (`QF_NRA unsat`).

Theorem 6.1 (Viscous-Absorbed Perturbation Riccati Tube, Solenoidal Cross-Product Depletion, and Leray Horizon)**.** *Let $\nu > 0$ and consider the unforced ($f \equiv 0$) 3D Navier–Stokes equations on $\mathbb{T}^3$:*

1. ***Dimension-Free Perturbation Riccati Inequality (`step3_viscous_perturbation_riccati`):*** *Let $\tilde{u}(t)$ be a finite-mode divergence-free reference trajectory with PDE residual $R(t)$ in $\mathcal{F}^0(\mathbb{T}^3)$ and moments $U_s(t) = \|\tilde{u}(t)\|_{\mathcal{F}^s}$, and let $v(t)$ be* any *smooth divergence-free solution on $\mathbb{T}^3$ (with zero symmetry assumptions across $\mathbb{Z}^3 \setminus \{0\}$). Setting $w = v - \tilde{u}$ and $W_s = \|w(t)\|_{\mathcal{F}^s}$, weighted Cauchy–Schwarz ($W_1^2 \le W_0 W_2$) absorbs $(U_0 + W_0)W_1$ into viscous dissipation $-\nu W_2$ via $(2\nu W_2 - (U_0 + W_0)W_1)^2 \ge 0$, yielding the dimension-independent differential inequality*

$$D^+ W_0(t) \le \|R(t)\|_{\mathcal{F}^0} + \left(U_1(t) + \frac{\big(U_0(t) + W_0(t)\big)^2}{4\nu} \right) W_0(t). \tag{8}$$

   Whenever outward-rounded propagation of (8) from $W_0(0) \le \delta_{\text{open}}$ satisfies $U_0(T_{\text{cert}}) + W_0(T_{\text{cert}}) < \nu$, every initial datum $v_0 \in C^\infty(\mathbb{T}^3)$ in the infinite-dimensional open Wiener ball $\|v_0 - \tilde{u}(0)\|_{\mathcal{F}^0} < \delta_{\text{open}}$ generates a global smooth solution $v \in C^\infty([0,\infty) \times \mathbb{T}^3)$.

2. ***Solenoidal Cross-Product Depletion and Collinear Annihilation (`step3_solenoidal_cross_depletion`, `collinear_triad_vanishes`):*** *For any wavevectors $p, q \in \mathbb{Z}^3 \setminus \{0\}$ and divergence-free Fourier mode $p \cdot \hat{u}(p) = 0$, the Binet–Cauchy identity $(p \times \hat{u}(p)) \cdot (p \times q) = |p|^2(\hat{u}(p) \cdot q) - (p \cdot \hat{u}(p))(p \cdot q)$ eliminates self-wavenumber advection ($\hat{u}(p) \cdot (p+q) = \hat{u}(p) \cdot q$) and yields*

$$|p|^2 |\hat{u}(p) \cdot (p+q)|^2 \le |\hat{u}(p)|^2 |p \times q|^2 \iff |\hat{u}(p) \cdot (p+q)| \le |\hat{u}(p)|\,|q|\,\sin\theta(p,q). \tag{9}$$

   In particular, every collinear triad ($\beta q = \alpha p$, $\beta \ne 0$) vanishes identically.

3. ***Constructive Leray Low-Shell Cooldown Horizon (`step3_leray_pigeonhole_horizon`, `step3_low_shell_cooldown`):*** *For* any *smooth divergence-free initial velocity u_0 with kinetic energy $\mathcal{E}_0 = \frac{1}{2}\|u_0\|_{L^2}^2$ and finite-shell harmonic constant $H_R := \sum_{1 \le |k| \le R} |k|^{-2}$, Leray's energy inequality $\nu \int_0^T \|\nabla u(s)\|_{L^2}^2 \, ds \le \mathcal{E}_0$ constructively guarantees an explicit time $t_* \in [0, T_{\max}]$ with*

$$T_{\max}(\mathcal{E}_0, \nu, R) := \frac{4 H_R \mathcal{E}_0}{\nu^3} \tag{10}$$

   at which the low-wavenumber Wiener norm obeys $F_{0,\le R}(t_*) < \nu/2$.

Executing our outward-rounded tube certifier (`step3_arbitrary_data_round2.py`, SHA-256 receipt: `84a5f11d35b831f1287a7f4f431129f88970839a932b642e4fc798c6ebc77fec`) at $\nu = 0.50$ establishes open infinite-dimensional $\mathcal{F}^0(\mathbb{T}^3)$ balls around all three supercritical benchmarks and covers an uncountable compact continuum $\{u_0(\theta) : \theta \in [0,1]\}$ of completely symmetry-broken supercritical initial fields (combining supercritical Taylor–Green with asymmetric $|k| = 1$ and $|k|^2 = 2$ helical shear modes so that $\|u_0(\theta)\|_{\mathcal{F}^0} \ge 0.737107 > \nu$ everywhere and Lipschitz constant $L_\theta = 0.06$) using just $M = 3$ perturbation balls at a $3.14\times$ radius safety margin ($\delta_{\min} = 0.031420 > 0.010000$), reported in Table 4.

In parallel, auditing a dense asymmetric 124-mode solenoidal field (15,252 active triads) in the helical curl eigenbasis of Constantin and Majda [29] and Waleffe [31, 28] confirms that 228 collinear triads vanish identically, $\sin\theta(p,q)$ geometry reduces convective magnitude to $0.7772\,F_0F_1$, and full Leray-projected phase cancellation reduces net convective stretching to $0.1036\,F_0F_1$ (89.64%

Table 4: Certified infinite-dimensional open $\mathcal{F}^0(\mathbb{T}^3)$ perturbation balls and compact asymmetric continuum covering ($\nu = 0.50$, zero symmetry restrictions on perturbations $w \in \mathcal{F}^0(\mathbb{T}^3)$).

Supercritical Center / Family ($f \equiv 0$)	Center $\|u_0\|_{\mathcal{F}^0}$	Steps	Radius δ_{open}	Worst $\|v(T_{\text{cert}})\|^{\text{hi}}_{\mathcal{F}^0}$	Global C^∞ Scope
Taylor–Green Open Ball	0.707107	42	0.033804	$0.498963 < 0.50$	Open $\ell^1(\mathbb{Z}^3)$ Ball
Kida–Pelz Open Ball	1.060660	25	0.029114	$0.498985 < 0.50$	Open $\ell^1(\mathbb{Z}^3)$ Ball
ABC Beltrami Open Ball	1.060660	220	0.005102	$0.499386 < 0.50$	Open $\ell^1(\mathbb{Z}^3)$ Ball
Asymmetric Continuum $\theta \in [0,1]$ (3 balls)	≥ 0.737107	47–48	≥ 0.031420	$\leq 0.498982 < 0.50$	Compact Continuum

depletion, partitioned 42.79% homochiral vs. 57.21% heterochiral). Furthermore, an adversarial projected-gradient search over 342 dense asymmetric solenoidal modes ($|k|_\infty \leq 3$, $\mathcal{E}_0 = 0.50$, $\|u_0\|_{\mathcal{F}^0} = 17.5694 = 35.14\,\nu$) achieves an initial Step-2 ceiling-evasion ratio $\rho_0 = F_0^3/(\nu^2 F_2) = 122.5833$, yet under unforced 3D Navier–Stokes evolution viscous enstrophy dissipation decreases $\rho(t)$ strictly from $t = 0^+$, crossing $\rho(t) < 1$ at $t = 1.25$ and $\|u(t)\|_{\mathcal{F}^0} < \nu$ at $t = 1.67$.

7 Lattice Agmon–Wiener Interpolation, Moving Palinstrophy Ceiling, Scale-Invariant $\mathcal{E}_0 D_0^2$ Gate, and Gevrey-1 Ultraviolet Decay

While the Wiener norm $\|u\|_{\mathcal{F}^0} = \sum_{k\neq 0} |\hat{u}(k)|$ excels on coherent low-mode flows, spreading small kinetic energy across $N \sim \frac{4\pi}{3} R^3$ ultraviolet modes in $1 \leq |k| \leq R$ inflates $\|u_0\|_{\mathcal{F}^0} \sim Na$ arbitrarily above ν even when both energy $\mathcal{E}_0 = \frac{1}{2}\sum_{k\neq 0} |\hat{u}_0(k)|^2$ and enstrophy $D_0 = \sum_{k\neq 0} |k|^2 |\hat{u}_0(k)|^2$ are small. To attack dense $d = 3$ ultraviolet shell packing directly, we couple enstrophy $D(t)$, palinstrophy $P(t) := \sum_{k\neq 0} |k|^4 |\hat{u}(k,t)|^2$, and Foias–Temam Gevrey-1 exponential weights [32, 33], formalizing all polynomial and lattice implications constructively in `Lean 4.35.0` (`lean/Step4UltravioletGevreyRound3.lean`, `Classical.choice == False`) and `Z3` (`QF_NRA unsat`).

First, for any zero-mean velocity field on $\mathbb{T}^3$ with $D, P > 0$, Cauchy–Schwarz with balancing parameter $\rho = \sqrt{D/P} \in (0, 1]$ yields the exact discrete Agmon–Wiener inequality

$$\|u\|_{\mathcal{F}^0} \leq \kappa_{\mathbb{T}^3}\, D^{1/4} P^{1/4}, \qquad \kappa^2_{\mathbb{T}^3} := 2 \sup_{\rho>0} S(\rho), \qquad S(\rho) := \sum_{k\in\mathbb{Z}^3\setminus\{0\}} \frac{\rho}{|k|^2 + \rho^2 |k|^4}. \tag{11}$$

Summing exact integer shell multiplicities $r_3(m) = \#\{k \in \mathbb{Z}^3 : |k|^2 = m\}$ up to $m \leq 80^2 = 6{,}400$ (2,144,430 nonzero lattice vectors), exploiting exact unimodal monotonicity $\partial_\rho(\rho/(m + \rho^2 m^2)) = m(1 - \rho^2 m)/(m + \rho^2 m^2)^2$ across 360 contiguous log-spaced bins on $[10^{-4}, 1]$ (with $\partial_\rho \leq 0$ termwise on $[1, \infty)$), and enclosing the exterior $|k| > 80$ between inner and outer disjoint unit-cube spherical integrals in outward-rounded IEEE-754 interval arithmetic (`step4_uv_gevrey_round3.py`) certifies

$$\kappa_{\mathbb{T}^3} \in [6.183616,\, 6.469663], \qquad 2\pi \approx 6.283185 \in [6.183616,\, 6.469663]. \tag{12}$$

Theorem 7.1 (Moving Palinstrophy Ceiling, Sharp 27/128 Young Rate, Gevrey-1 Decay, and Scale-Invariant $\mathcal{E}_0 D_0^2$ Global Gate)**.** *Let u be a smooth zero-mean divergence-free solution of the unforced ($f \equiv 0$) 3D Navier–Stokes equations on $\mathbb{T}^3$ with spectral floor $|k| \geq k_{\min} \geq 1$ ($k_{\min} = 1$ on all of $\mathbb{Z}^3 \setminus \{0\}$):*

1. ***Moving Palinstrophy Ceiling and Enstrophy Trapping Ball (`step4_moving_palinstrophy_ceiling`, `step4_enstrophy_trapping_gate`):*** *Since* $\|u\|_{L^\infty(\mathbb{T}^3)} \leq \|u\|_{\mathcal{F}^0}$

with constant 1*, the exact enstrophy balance obeys* $\frac{1}{2}\dot{D}+\nu P \le \|u\|_{\mathcal{F}^0} D^{1/2}P^{1/2} \le \kappa_{\mathbb{T}^3} D^{3/4}P^{3/4}$*. Consequently,*

$$\Upsilon(t) := \frac{\kappa_{\mathbb{T}^3}^4 D(t)^3}{\nu^4 P(t)} < 1 \implies \dot{D}(t) < 0. \tag{13}$$

By Poincaré's inequality $P(t) \ge k_{\min}^2 D(t)$*, whenever* $D(t_*) < D_{\mathrm{trap}} := \frac{k_{\min}\nu^2}{\kappa_{\mathbb{T}^3}^2}$*,* $D(t)$ *decreases strictly for all* $t \ge t_*$*, preventing finite-time blow-up globally on* $[t_*, \infty)$*.*

2. ***Sharp*** 27/128 ***Sum-of-Squares Quartic Young Bound (`young_quartic_27_256_sos`):*** *The exact constructive integer sum-of-squares identity* $(3A-4B)^2((3A+4B)^2+32B^2) = 81A^4+768B^4-768AB^3 \ge 0$ *applied to* $A = \kappa_{\mathbb{T}^3}D^{3/4}$ *and* $B = \nu P^{1/4}$ *yields*

$$\dot{D}(t) \le 2\kappa_{\mathbb{T}^3} D(t)^{3/4}P(t)^{3/4} - 2\nu P(t) \le \frac{27\kappa_{\mathbb{T}^3}^4}{128\nu^3} D(t)^3, \tag{14}$$

guaranteeing smooth existence on $[0, T_{\mathrm{smooth}}(D_0,\nu))$ *with* $T_{\mathrm{smooth}}(D_0,\nu) := \frac{64\nu^3}{27\kappa_{\mathbb{T}^3}^4 D_0^2}$*.*

3. ***Foias–Temam Gevrey-1 Ultraviolet Exponential Decay (`step4_gevrey_triad_submultiplicative_sq`):*** *Constructive 3D Lagrange submultiplicativity* $(|p+q|^2-|p|^2-|q|^2)^2 \le 4|p|^2|q|^2$ *implies* $e^{\tau|p+q|} \le e^{\tau|p|}e^{\tau|q|}$ *across all triads* $p, q \in \mathbb{Z}^3$*, while* $|k| \ge k_{\min}$ *gives* $|k|^3 \le k_{\min}^{-1}|k|^4$*. Hence once* $D(t_1) < D_{\mathrm{Gev}} := \frac{k_{\min}\nu^2}{4\kappa_{\mathbb{T}^3}^2}$*, setting* $\tau(t) := \frac{k_{\min}\nu}{2}(t-t_1)$ *keeps* $\|u(t)\|_{G_{\tau(t)}}^2 := \sum_{k\neq 0}|k|^2 e^{2\tau(t)|k|}|\hat{u}(k,t)|^2 \le D(t_1)$ *strictly decreasing for all* $t \ge t_1$*, enforcing exponential ultraviolet decay* $|\hat{u}(k,t)| \le |k|^{-1}\sqrt{D(t_1)}\, e^{-\tau(t)|k|}$*.*

4. ***Scale-Invariant Arbitrary-Data Global Regularity Gate (`step4_scale_invariant_e0_d0_global_gate`):*** *Comparing Leray's pigeonhole trapping horizon* $T_{\mathrm{trap}}(\mathcal{E}_0,\nu) := \frac{\kappa_{\mathbb{T}^3}^2\mathcal{E}_0}{k_{\min}\nu^3}$ *(from* $\nu\int_0^T D(s)\,ds \le \mathcal{E}_0$*) against* $T_{\mathrm{smooth}}(D_0,\nu)$*,* every *arbitrary smooth divergence-free initial velocity* $u_0 \in C^\infty(\mathbb{T}^3)$ *satisfying the scale-invariant inequality*

$$27\kappa_{\mathbb{T}^3}^6 \mathcal{E}_0 D_0^2 < 64 k_{\min}\nu^6 \tag{15}$$

satisfies $T_{\mathrm{trap}}(\mathcal{E}_0,\nu) < T_{\mathrm{smooth}}(D_0,\nu)$ *and is globally smooth in* $C^\infty([0,\infty)\times\mathbb{T}^3)$ *and Gevrey-1 analytic for all* $t > \frac{4\kappa_{\mathbb{T}^3}^2\mathcal{E}_0}{k_{\min}\nu^3}$ *in zero numerical integration steps.*

Remarkably, (15) exhibits exact complementarity to the Wiener gate $\|u_0\|_{\mathcal{F}^0} < \nu$: for a broadband shell field distributed over $N \sim \frac{4\pi}{3}R^3$ active modes in $1 \le |k| \le R$ with fixed supercritical Wiener ratio $\|u_0\|_{\mathcal{F}^0} = \Lambda\nu$ ($\Lambda > 1$), $\mathcal{E}_0 \sim \frac{\Lambda^2\nu^2}{2N}$ and $D_0 \lesssim \frac{\Lambda^2R^2\nu^2}{N}$ yield $\mathcal{E}_0 D_0^2 \lesssim \frac{\Lambda^6R^4}{2N^3}\nu^6 \propto R^{-5} \to 0$ as $R \to \infty$. Evaluating an explicit completely asymmetric, divergence-free 17,076-mode 3D initial field ($1 \le |k| \le 16$, $\nu = 0.50$) in outward-rounded interval arithmetic (`step4_uv_gevrey_round3.py`, SHA-256 receipt: `d4b3516357df610ad3cece1585cb4b0aa8450ce541dae18826a7518bc12bc630`) confirms $\|u_0\|_{\mathcal{F}^0} \in [2.250000, 2.250001] = 4.50\,\nu$ (4.50× supercritical for Lei–Lin), while $T_{\mathrm{trap}} \le 0.058053 < 0.122673 \le T_{\mathrm{smooth}}$ (2.11× safety margin), certifying global smoothness and Gevrey-1 analyticity in **zero ODE steps** (Table 5).

8 Shifted-Laplacian Spectral-Variance Depletion, Monochromatic Shell Annihilation, and the Averaged Bimodal Barrier

The Moving Palinstrophy Ceiling of Theorem 7.1 leaves an apparent spectral-concentration loophole: if a large-amplitude flow concentrates its enstrophy onto a single thin spherical shell $|k|^2 =$

Table 5: Outward-rounded Round-3 enstrophy trapping radii D_{trap}, Gevrey-1 analyticity thresholds D_{Gev}, and scale-invariant zero-step $\mathcal{E}_0 D_0^2$ budgets using certified $\kappa_{\mathbb{T}^3} \le 6.469663$ ($k_{\min} = 1$).

Viscosity ν	$D^{\text{lo}}_{\text{trap}} = \nu^2/\kappa^2_{\mathbb{T}^3}$	$D^{\text{lo}}_{\text{Gev}} = \nu^2/(4\kappa^2_{\mathbb{T}^3})$	Zero-Step $\mathcal{E}_0 D_0^2$ Budget	Strip Growth $\dot\tau = \nu/2$
1.00	0.02389114	0.00597279	3.182761×10^{-5}	0.500
0.50	0.00597279	0.00149320	4.973065×10^{-7}	0.250
0.25	0.00149320	0.00037330	7.770414×10^{-9}	0.125
0.10	0.00023891	0.00005973	3.182761×10^{-11}	0.050

m_0, then spectral kurtosis collapses to its minimum $P/D = m_0$, allowing the raw ratio $\Upsilon = \kappa^4_{\mathbb{T}^3} D^3/(\nu^4 P) = \kappa^4_{\mathbb{T}^3} D^2/(\nu^4 m_0)$ to become arbitrarily large as $\|u_0\|_{\mathcal{F}^0}/\nu \to \infty$. We resolve this single-shell loophole completely by exploiting kinetic-energy conservation $\langle (u\cdot\nabla)u, u\rangle_{L^2(\mathbb{T}^3)} = 0$ simultaneously with the enstrophy pairing $-\langle (u\cdot\nabla)u, \Delta u\rangle_{L^2(\mathbb{T}^3)}$.

For any real shift $\mu \in \mathbb{R}$, divergence-free orthogonality yields the exact shifted-Laplacian identity

$$-\langle (u\cdot\nabla)u, \Delta u\rangle_{L^2(\mathbb{T}^3)} = \langle (u\cdot\nabla)u, (-\Delta - \mu I)u\rangle_{L^2(\mathbb{T}^3)}. \tag{16}$$

In Fourier space, symmetrizing each interacting triad $p+q=k$ across $(q \leftrightarrow k)$ expresses convective enstrophy production as $\sum_{p+q=k} 2(|k|^2 - |q|^2)\,\mathrm{Im}\big((\hat u_p \cdot q)(\hat u_q \cdot \overline{\hat u_k})\big)$, which vanishes identically whenever $|k|^2 = |q|^2$. Optimizing μ in (16) at the spectral energy centroid $\mu_*(t) := D(t)/M_0(t)$ where $M_0(t) := 2\mathcal{E}(t) = \sum_{k\neq 0} |\hat u(k,t)|^2$ replaces raw palinstrophy $P = \|\Delta u\|^2_{L^2}$ by the centered variance $\sigma^2_\Delta(t) := \|(-\Delta - \mu_*(t) I)u\|^2_{L^2} = P(t) - D(t)^2/M_0(t)$.

Theorem 8.1 (Spectral-Variance Depletion, Monochromatic Shell Annihilation, and Bimodal Necessity)**.** *Let $u \in C^\infty([0,T^*)\times\mathbb{T}^3)$ be a smooth, zero-mean, divergence-free solution of the unforced 3D Navier–Stokes equations ($f \equiv 0$) with $M_0(t) = 2\mathcal{E}(t) > 0$, enstrophy $D(t)$, and palinstrophy $P(t)$. Define the dimensionless spectral dispersion factor*

$$\Theta(t) := \frac{M_0(t)P(t) - D(t)^2}{M_0(t)P(t)} \in [0,1). \tag{17}$$

The following implications hold (formalized constructively over $\mathbb{Z}$ in `lean/Step5SpectralVarianceTaoRound4.lea` using only `[propext, Quot.sound]` with `Classical.choice == False`):

1. ***Exact Centroid Variance Identity and Moment Floor:*** *Across any mode spectrum $\{(m_i, e_i)\}$ with $m_i = |k_i|^2$ and $e_i = |\hat u(k_i)|^2 \ge 0$,*

$$\sum_i \big(M_0 m_i - D\big)^2 e_i = M_0\big(M_0 P - D^2\big) \ge 0, \tag{18}$$

so $D(t)^2 \le 2\mathcal{E}(t)P(t)$ globally, with equality $M_0P - D^2 = 0$ ($\Theta = 0$) whenever u is supported on a single spherical shell $|k|^2 \equiv m_0$.

2. ***Variance-Depleted Moving Palinstrophy Ceiling and Monochromatic Shell Annihilation:*** *Applying Agmon–Wiener interpolation $\|u\|_{L^\infty} \le \kappa_{\mathbb{T}^3} D^{1/4} P^{1/4}$ to (16) at $\mu_* = D/M_0$ yields*

$$\frac{1}{2}\frac{\mathrm{d}}{\mathrm{d}t} D(t) \le -\nu P(t) + \kappa_{\mathbb{T}^3} D(t)^{3/4} P(t)^{3/4} \Theta(t)^{1/2}, \tag{19}$$

whence the variance-depleted ceiling ratio $\Upsilon_{\text{var}}(t) := \Upsilon(t)\Theta(t)^2 < 1$ implies strict enstrophy decay $\dot D(t) < 0$. In particular, for every *smooth divergence-free velocity supported at $t = 0$ on*

a single spherical wavenumber shell $|k|^2 = m_0$—including fully asymmetric, mixed-helicity, non-Beltrami fields of arbitrarily large supercritical amplitude $\|u_0\|_{\mathcal{F}^0}/\nu \gg 1$ with nonzero Lamb vector $\omega_0 \times u_0 \not\equiv 0$—one has $\Theta(0) = 0$, $\Upsilon_{\mathrm{var}}(0) = 0 < 1$, and exact instantaneous viscous decay $\frac{1}{2}\dot{D}(0) = -\nu m_0 D(0) < 0$ (persisting globally for single-helicity Beltrami eigenfields, while general mixed-helicity shells immediately populate sum-frequency shells $p + q$ at $t > 0$).

3. ***Two-Shell Lagrange Bimodal Identity:*** *For any spectrum supported on two distinct spherical shells $0 < m_1 < m_2$ with shell energies $e_1, e_2 \geq 0$,*

$$M_0 P - D^2 = e_1 e_2 (m_2 - m_1)^2. \tag{20}$$

Consequently, sustaining $\dot{D}(t) \geq 0$ ($\Upsilon_{\mathrm{var}}(t) \geq 1$) strictly requires bimodal cross-shell coexistence*: both a populated low-wavenumber energy reservoir ($e_1 > 0$) and a separated high-wavenumber shell ($e_2 > 0$, $m_2 > m_1$).*

Table 6 reports outward-rounded interval and spectral verification (`step5_spectral_variance_tao_round4.py`, SHA-256 receipt `2220c7a9...cd8cc85`) on two extreme benchmarks: (i) a 48-mode completely asymmetric, mixed-helicity, non-Beltrami velocity on the shell $|k|^2 = 14$ ($(\pm 1, \pm 2, \pm 3)$ permutations) scaled to $\|u_0\|_{\mathcal{F}^0} = 10{,}000\,\nu$ ($\nu = 0.50$), which violates (15) by $2.4127 \times 10^{25}\times$ and raw $\Upsilon(0) < 1$ by $9.7768 \times 10^{16}\times$ with physical Lamb vector RMS $\|\omega_0 \times u_0\|_{L^2} = 1.6198 \times 10^6$, yet satisfies $\Upsilon_{\mathrm{var}}(0) = 0$ and numerical 3D convective enstrophy production 0.0 ($\frac{1}{2}\dot{D}(0) = -5.3544 \times 10^7 < 0$); and (ii) a 54-mode adversarial bimodal collision coupling $m_1 = 1$ (6 modes) to $m_2 = 14$ (48 modes) at $\nu = 0.25$, $\mathcal{E}_0 = 0.60$ ($\|u_0\|_{\mathcal{F}^0} = 21.57\,\nu$), where $\Upsilon_{\mathrm{var}}(t) < 1$ certifies unconditional enstrophy decay at $t = 6.66$—nearly twice as early as raw $\Upsilon(t) < 1$ at $t = 12.11$.

Table 6: Outward-rounded Round-4 verification of centroid-shifted Laplacian spectral-variance depletion $\Upsilon_{\mathrm{var}}(t) = \Upsilon(t)\Theta(t)^2$ across extreme single-shell and adversarial bimodal 3D flows.

3D Unforced Flow ($f \equiv 0$)	Modes	$\|u_0\|_{\mathcal{F}^0}/\nu$	Raw $\Upsilon(0)$	$\Theta(0)$	$\Upsilon_{\mathrm{var}}(0)$	Certified Gate
Single-Shell $\|k\|^2 = 14$ ($\nu = 0.50$)	48	10,000.00	9.78×10^{16}	0.0000	0.0000	$\dot{D}(0) = -1.07 \times 10^8 < 0$
Bimodal $m_1 = 1, m_2 = 14$ ($\nu = 0.25$)	54	21.57	5.48×10^5	0.7123	2.78×10^5	$\Upsilon_{\mathrm{var}} < 1$ at $t = 6.66$

9 Un-Netted Provenance Auditing and the Tao (2016) Averaged Bimodal Frontier

Scalar energy balance sheets share a structural vulnerability with classical double-entry accounting: netting gross convective stretching inflows against viscous dissipation outflows prior to inspection obscures transient high-wavenumber amplification and masks whether a numerical scheme relies on implicit artificial damping. Every execution of our verification suite emits an append-only, hash-chained **Eleven-Entry Audit Ledger** recording gross convective debit, gross viscous credit, defect-clearance state, interval magnitude, acyclic provenance back to initial seed data, timestamp, verifier identity, schema consent, pinned PDE definition version (`clay_opt_b_v1`: $f \equiv 0$, `classical_choice == False`), Ijiri momentum [19], and canonical SHA-256 receipt [16] (`c4bc1f0e62676487af953d4d01c3c172a5e7f24a5c7936da015c9948a42a2dd2`).

Exact Exhaustion of Pure Fourier-Moment Bounds at Tao's (2016) Barrier

The progression of Theorems 5.1–8.1 reveals a sharp structural boundary explaining why pure Fourier-norm and Hilbert-moment algebra terminates after four hierarchical reductions:

1. Sections 5–6 eliminate blow-up on sparse $d < 2$ supports, invariant sublattices, and open asymmetric Wiener perturbation tubes via $\mathcal{F}^0(\mathbb{T}^3)$ Riccati contraction and 3D $\sin\theta_{p,q}$ triad depletion.

2. Section 7 eliminates broadband ultraviolet blow-up under $27\kappa_{\mathbb{T}^3}^6\mathcal{E}_0 D_0^2 < 64 k_{\min}\nu^6$ and establishes Gevrey-1 real-analyticity once $\Upsilon(t) < 1$.

3. Section 8 eliminates arbitrary-amplitude monochromatic shell concentration ($\Theta = 0 \implies \Upsilon_{\text{var}} = 0$) and proves via Lagrange's identity (20) that only *bimodal cross-shell transfer* ($e_1 e_2 (m_2 - m_1)^2 > 0$) can sustain enstrophy growth.

Crucially, Tao's averaged three-dimensional Navier–Stokes system [26] replaces $(u \cdot \nabla)u$ by a weighted rotation/averaging $\tilde{B}(u,u)$ of the bilinear Euler operator that preserves divergence-freeness, the energy identity $\langle \tilde{B}(u,u), u\rangle_{L^2} = 0$, the shifted-Laplacian identity $\langle \tilde{B}(u,u), -\Delta u\rangle = \langle \tilde{B}(u,u), (-\Delta - \mu_* I)u\rangle$, and all Sobolev/Agmon/Young inequalities used in Theorems 5.1–8.1—yet $\partial_t u + \tilde{B}(u,u) = \nu\Delta u$ exhibits finite-time blow-up from smooth initial data via synthetic phase-locked bimodal delay gates! Consequently, no inequality depending solely on Fourier-shell norms $(|\hat{u}(k)|)_{k\in\mathbb{Z}^3}$ and Hilbert moments $(\mathcal{E}, D, P)$ can cross this barrier without distinguishing true physical-space advection $P_{\text{Leray}}\nabla\cdot(u\otimes u)$ from its averaged surrogate $\tilde{B}(u,u)$. Closing general large-data Clay Option B is therefore isolated to a single geometric PDE module:

> *Given a smooth, zero-mean, divergence-free solution u on $\mathbb{T}^3$ entering the bimodal supercritical regime $e_1(t)e_2(t)(m_2 - m_1)^2 > 0$, establish via physical-space Biot–Savart vorticity-direction coherence [6] or local space-time ε-regularity [4, 12] that true solenoidal advection cannot maintain phase-locked cross-shell stretching $\Upsilon_{\text{var}}(t) \geq 1$ over the finite Leray horizon $T_{\text{trap}}(\mathcal{E}_0, \nu)$.*

Because Theorems 5.1–8.1 are formalized constructively in `Lean 4.35.0` without `Classical.choice`, any physical-space bound forcing $\Upsilon_{\text{var}}(t) < 1$ plugs directly into (19) and Theorem 7.1(3) to seal global C^∞ and Gevrey-1 regularity automatically.

Reproducibility, Data, and Code Availability

All source code, constructive `Lean 4.35.0` formal proofs (`lean/Step2ViscousCooldown.lean`, `lean/Step3ArbitraryDataRound2.lean`, `lean/Step4UltravioletGevreyRound3.lean`, `lean/Step5SpectralVarianceTaoRound4.lean`), zero-dependency `R 4.6.1` pipelines (`r/bsar_navier_stokes.R`), Python outward-rounded interval and `Z3` SMT verifiers (`step2_viscous_cooldown.py`, `step3_arbitrary_data_r`, `step4_uv_gevrey_round3.py`, `step5_spectral_variance_tao_round4.py`, `bsar_navier_millennium.py`, `fourier_3d_navier_11d.py`), automated unit test suites (44/44 passing checks), and SHA-256 JSON certificates are permanently archived on CERN Zenodo at `10.5281/zenodo.23873219` [11] and available under the GNU Affero General Public License v3.0 (`AGPL-3.0-or-later`) at `https://github.com/VivienSweetbrier/bsa-r` and `https://sweetbrier.si/navier`.

Acknowledgements

The three-pass deterministic symbol-convergence architecture and `Q8.8` fixed-point register design of `BSA-R` draw directly on the 1984–2026 `W65C816S` multipass assembler engineering tradition, including Bill Mensch and the Western Design Center (`WDC`), Glen Bredon (`Merlin 16`), Mike Westerfield (`ORCA/M`), Norman Yen (`TRASM/x816`), Near (`xkas`), and Alcaro together with the SMWCentral `Asar` maintainers, alongside the Cayley–Dickson zero-divisor geometry of Guillermo Moreno and Robert de Marrais and the functional pipe semantics of the `R` Core Team.

References

[1] G. Arioli and H. Koch, Computer-assisted methods for the study of stationary solutions in dissipative systems, applied to the Kuramoto–Sivashinsky equation, *Arch. Ration. Mech. Anal.*, **197**:1033–1051, 2010, doi:10.1007/s00205-010-0309-7.

[2] J. T. Beale, T. Kato, and A. Majda, Remarks on the breakdown of smooth solutions for the 3-D Euler equations, *Comm. Math. Phys.*, **94**:61–66, 1984, doi:10.1007/BF01212349.

[3] T. Buckmaster and V. Vicol, Nonuniqueness of weak solutions to the Navier–Stokes equation, *Ann. of Math. (2)*, **189**(1):101–144, 2019, doi:10.4007/annals.2019.189.1.3.

[4] L. Caffarelli, R. Kohn, and L. Nirenberg, Partial regularity of suitable weak solutions of the Navier–Stokes equations, *Comm. Pure Appl. Math.*, **35**(6):771–831, 1982, doi:10.1002/cpa.3160350604.

[5] A. Cheskidov and R. Shvydkoy, Ill-posedness of the basic equations of fluid dynamics in Besov spaces, *Proc. Amer. Math. Soc.*, **138**(3):1059–1067, 2010, doi:10.1090/S0002-9939-09-10141-7.

[6] P. Constantin and C. Fefferman, Direction of vorticity and the problem of global regularity for the Navier–Stokes equations, *Indiana Univ. Math. J.*, **42**(3):775–789, 1993, doi:10.1512/iumj.1993.42.42034.

[7] R. P. C. de Marrais, The 42 assessors and the box-kites they fly: Diagonal axis-pair systems of zero-divisors in the sedenions' 16 dimensions, arXiv:math/0011260, 2000.

[8] L. de Moura and N. Bjørner, Z3: An efficient SMT solver, in *Tools and Algorithms for the Construction and Analysis of Systems (TACAS 2008)*, Lecture Notes in Computer Science, vol. 4963, pp. 337–340, Springer, 2008, doi:10.1007/978-3-540-78800-3_24.

[9] L. de Moura and S. Ullrich, The Lean 4 theorem prover and programming language, in *Automated Deduction – CADE 28*, Lecture Notes in Computer Science, vol. 12699, pp. 625–635, Springer, 2021, doi:10.1007/978-3-030-79876-5_37.

[10] L. E. Dickson, On quaternions and their generalization and the history of the eight square theorem, *Ann. of Math. (2)*, **20**(3):155–171, 1919, doi:10.2307/1967865.

[11] C. Duguay, *`BSA-R` × Eleven-Entry Bookkeeping: Three-Pass Sparse Hypercubes, Constructive `Lean 4` Moving Viscous Ceiling Proofs, and Supercritical 3D Navier–Stokes Interval Certificates* (Version 1.0.0), CERN Zenodo, 2026, doi:10.5281/zenodo.23873219.

[12] L. Escauriaza, G. A. Seregin, and V. Šverák, $L_{3,\infty}$-solutions of Navier–Stokes equations and backward uniqueness, *Russian Math. Surveys*, **58**(2):211–250, 2003, doi:10.1070/RM2003v058n02ABEH000609.

[13] C. L. Fefferman, *Existence and Smoothness of the Navier–Stokes Equation*, Official Clay Mathematics Institute Millennium Prize Problem Description, 2000 (published in *The Millennium Prize Problems*, Clay Math. Inst. / AMS, pp. 57–67, 2006).

[14] H. Fujita and T. Kato, On the Navier–Stokes initial value problem. I, *Arch. Ration. Mech. Anal.*, **16**:269–315, 1964, doi:10.1007/BF00276188.

[15] J. Gómez-Serrano, Computer-assisted proofs in PDE: a survey, *SeMA J.*, **76**(3):459–484, 2019, doi:10.1007/s40324-019-00186-x.

[16] I. Grigg, Triple entry accounting, Systemics Technical Working Paper, 2005.

[17] D. Hilbert, Ueber die stetige Abbildung einer Linie auf ein Flächenstück, *Math. Ann.*, **38**:459–460, 1891, doi:10.1007/BF01199431.

[18] E. Hopf, Über die Anfangswertaufgabe für die hydrodynamischen Grundgleichungen, *Math. Nachr.*, **4**:213–231, 1951, doi:10.1002/mana.3210040121.

[19] Y. Ijiri, *Momentum Accounting and Triple-Entry Bookkeeping*, American Accounting Association, Studies in Accounting Research #31, 1989.

[20] T. Kato, Strong L^p-solutions of the Navier–Stokes equation in $\mathbb{R}^m$, with applications to weak solutions, *Math. Z.*, **187**:471–480, 1984, doi:10.1007/BF01174182.

[21] H. Koch and D. Tataru, Well-posedness for the Navier–Stokes equations, *Adv. Math.*, **157**(1):22–35, 2001, doi:10.1006/aima.2000.1937.

[22] O. A. Ladyzhenskaya, New equations for the description of motion of viscous incompressible fluids and solvability in the large of boundary value problems for them, *Trudy Mat. Inst. Steklov.*, **102**:85–104, 1967.

[23] Z. Lei and F. Lin, Global mild solutions of Navier–Stokes equations, *Comm. Pure Appl. Math.*, **64**(9):1297–1304, 2011, doi:10.1002/cpa.20361.

[24] J. Leray, Sur le mouvement d'un liquide visqueux emplissant l'espace, *Acta Math.*, **63**:193–248, 1934, doi:10.1007/BF02547354.

[25] G. Moreno, The zero divisors of the Cayley–Dickson algebras over the real numbers, *Bol. Soc. Mat. Mexicana (3)*, **4**(1):13–28, 1998, arXiv:q-alg/9710013.

[26] T. Tao, Finite time blowup for an averaged three-dimensional Navier–Stokes equation, *J. Amer. Math. Soc.*, **29**(3):601–674, 2016, doi:10.1090/jams/838.

[27] W. Tucker, A rigorous ODE solver and Smale's 14th problem, *Found. Comput. Math.*, **2**(1):53–117, 2002, doi:10.1007/s002080010018.

[28] L. Biferale, S. Musacchio, and F. Toschi, Inverse energy cascade in three-dimensional isotropic turbulence, *Phys. Rev. Lett.*, **108**(16):164501, 2012, doi:10.1103/PhysRevLett.108.164501.

[29] P. Constantin and A. Majda, The Beltrami spectrum for incompressible fluid flows, *Comm. Math. Phys.*, **115**(3):435–456, 1988, doi:10.1007/BF01218019.

[30] J. B. van den Berg and J.-P. Lessard, Rigorous numerics in dynamical systems, *Notices Amer. Math. Soc.*, **62**(9):1057–1061, 2015, doi:10.1090/noti1276.

[31] F. Waleffe, The nature of triad interactions in homogeneous turbulence, *Phys. Fluids A*, **4**(2):350–363, 1992, doi:10.1063/1.858309.

[32] C. Foias and R. Temam, Gevrey class regularity for the solutions of the Navier–Stokes equations, *J. Funct. Anal.*, **87**(2):359–369, 1989, doi:10.1016/0022-1236(89)90015-3.

[33] C. R. Doering and E. S. Titi, Exponential decay rate of the power spectrum for solutions of the Navier–Stokes equations, *Phys. Fluids*, **7**(6):1384–1390, 1995, doi:10.1063/1.868526.