

Alessandro Salvatoni
Paolo Brambilla
Manuela Deiana
Luigi Nespoli

Application of DEXA in body composition assessment in children

Presented at the 8th International
Workshop of the European Childhood
Obesity Group, 8–10 May 1998,
Wisła-Jawonik, Poland

A. Salvatoni (✉) · M. Deiana · L. Nespoli
Department of Pediatrics,
Faculty of Medicine,
University of Insubria, Varese, Italy

P. Brambilla
Department of Pediatrics,
Scientific Institute H. S. Raffaele,
Milan, Italy

Abstract Clinical and research applications of dual-energy X-ray absorptiometry in childhood are reviewed. Originally used for bone mineralization assessment in adults with osteoporosis, total body dual-energy X-ray absorptiometry is now also used currently alone or combined with other techniques for assessment of fat mass and fat-free mass. Low invasiveness, accuracy, swiftness and availability of pediatric software make total-body dual-en-

ergy X-ray absorptiometry suitable for studying body composition in a wide variety of metabolic and nutritional disorders in childhood such as simple or genetic obesity, insulin-dependent diabetes mellitus, congenital hypothyroidism, celiac disease, asthma, growth hormone deficiency and cachexic states.

Key words Body composition · Dual-energy X-ray absorptiometry · Review · Children

Dual-energy X-ray absorptiometry (DEXA) is a technique that allows simultaneous accurate and precise assessment of total and regional bone mineral content (BMC), bone mineral density (BMD), fat mass (FM) and non-mineral-non-fat mass (FFM) [10, 12, 25]. A total-body scan can be done in a short period of time (5–12 min) during which the subject receives a negligible radiation dose (0.02 mrem). Software suitable for adult and paediatric subjects is now available.

Clinical applications were limited until recently because the FDA required that body composition as assessed by DEXA be used only in research. However, in 1995 DEXA was cleared for use in body composition assessment in clinical practice. Osteoporosis, Cushing disease, prolonged steroid treatment, hyperparathyroidism, hyperthyroidism, hypogonadism, disorders of calcium metabolism, regional enteritis, idiopathic proctocolitis, intestinal malabsorption, chronic renal failure, vitamin D deficiency, and pathologic fractures are the main diagnostic applications for DEXA that are currently covered by Medicare.

Studies using DEXA over the last 5 years have shown that it accurately measures fat and lean content in meat samples and animal carcasses [16, 19, 27, 35]; dozens of

studies have shown that DEXA compares well with such noninvasive approaches as neutron activation and hydrodensitometry. Three main companies are now involved in DEXA equipment production (Lunar, Hologic and Norland). Compared with DPX-Lunar, QDR-Hologic underestimates FM by 20% and BMC by 15–25% and Norland overestimates FM by 5% [34].

Originally DEXA was used mainly for BMC and BMD measurement in adults with osteoporosis. Following the validation of this technique for FM and FFM assessment, DEXA was taken in many studies as a “gold standard” for validation of other techniques for body composition assessment, such as body mass index (BMI), skinfolds or body impedance analysis [6, 7, 14, 15, 39]. A recent paper concludes that BMI, compared with DEXA, underpredicts adiposity both in children with disease states and in healthy subjects [39]. According to one of the most popular online databases, Medline (Healthgate), more than 60 papers have been published on clinical applications of DEXA in the last 3 years, 30% of them relating to children.

In normal and obese children DEXA showed that both total and regional BMC are related more to the FFM than

to the FM [24]. Moreover, BMD in children was found to correlate with their own age, pubertal stage height and BMI and with the BMD of their parents, but not with calcium intake, suggesting that heredity plays a major part in osteoporosis [22]. However, recent studies have shown that BMD is not useful in children and in cross-sectional studies, as it is significantly influenced by bone size [26]. Therefore, as suggested by Prentice et al., when bone mineralization is assessed BMC should be adjusted for bone area [28].

The clinical use of DEXA in children is mainly applied to basal assessment and follow-up of body composition changes in chronic diseases such as insulin-dependent diabetes mellitus, celiac disease, asthma, and chronic renal failure. Most of the papers on DEXA report studies on tissue changes induced by growth hormone [4–6, 11, 17] treatment. While growth hormone does not increase bone density, it does increase muscle mass and decrease fat mass. Children with Prader-Labhart-Willi syndrome who were studied with DEXA showed a peculiar body composition, to some extent similar to that found in subjects deficient in growth hormone or even in sedentary and elderly people [8, 9, 23]. The changes induced by corticosteroids are deleterious (specifically, an increase in fat mass and a decrease in muscle and bone mass). Recent

studies on body composition assessed by DEXA in children chronically treated with inhaled steroids showed no adverse effects of the treatment on BMC and soft tissue body composition [1]; however, these results have yet to be confirmed by further studies, which are in progress at present. DEXA showed that spine BMD and FM in untreated and late-diagnosed celiac patients are significantly lower than in those diagnosed early [3, 33]. An obvious use of DEXA has been to assess changes to bone and lean tissue after weight loss [30, 36]. Another clinical use of DEXA is the measurement of body fat distribution; however, for accurate assessment of intraabdominal fat, the DEXA technique must be supplemented by a single-slice CT or RMI scan [20, 37].

Changes in muscle and fat mass associated with chronic physical activity have been measured using DEXA. Several studies have shown that athletes have a higher lean tissue mass and lower fat tissue mass [2, 18, 21, 38].

It has been assumed that diabetics will have some muscle wasting and often show osteoporosis, but recent studies conducted with DEXA did not show this, although water and potassium content were abnormal [29, 31]. DEXA also is useful for monitoring loss of fat, muscle and mineral bone in cachexic states such as AIDS [13], anorexia nervosa [32] or neoplastic disease.

References

- Agertoft L, Pedersen S (1998) Bone mineral density in children with asthma receiving long-term treatment with inhaled budesonide. *Am J Respir Crit Care Med* 157:178–183
- Avlonitou E, Georgiou E, Douskas G, Louizi A (1997) Estimation of body composition in competitive swimmers by means of three different techniques. *Int J Sports Med* 18:363–368
- Barera G, Brambilla P, Manzoni P (1995) Changes in body composition evaluated by DEXA during gluten free diet in coeliac children. *J Pediatr Gastroenterol Nutr* 20:476
- Bell W, Davies JS, Evans WD, Scanlon MF (1995) The validity of estimating total body fat and fat-free mass from skinfold thickness in adults with growth hormone deficiency. *J Clin Endocrinol Metab* 80:630–636
- Beshyah SA, Freemantle C, Thomas E, Rutherford O, Page B, Murphy M, Johnston DG (1995) Abnormal body composition and reduced bone mass in growth hormone deficient hypopituitary adults. *Clin Endocrinol* 42:179–189
- Beshyah SA, Freemantle C, Thomas E, Page B, Murphy M, Johnston DG (1995) Comparison of measurements of body composition by total body potassium, bioimpedance analysis, and dual-energy X-ray absorptiometry in hypopituitary adults before and during growth hormone treatment. *Am J Clin Nutr* 61:1186–1194
- Brambilla P, Manzoni P, Salvatoni A, Battistini N, Simone P, Brogginini M, Chiumello G (1993) Body composition and fat distribution using dual-energy X-ray absorptiometry (DEXA) in obese children. Comparison with deuterium dilution (D2O), anthropometry and BIA. *Pediatr Res* 33 [Suppl]:S80
- Brambilla P, Bosio L, Manzoni P, Pietrobelli A, Beccaria L, Chiumello G (1997) Peculiar body composition in patients with Prader-Labhart-Willi syndrome. *Am J Clin Nutr* 65:1369–1374
- Brismar TB, Lindgren AC, Ringertz H, Rosenborg M, Ritzen ME (1998) Total body bone mineral measurement in children with Prader-Willi syndrome: the influence of the skull's bone mineral content per area (BMA) and of height. *Pediatr Radiol* 28:38–42
- Cullum ID, Eil PJ, Ryder JP (1989) X-ray dual-photon absorptiometry: a new method for the measurement of bone density. *Br J Radiol* 62:587–592
- Cuneo RC, Judd S, Wallace JD, Perry-Keene D, Burger H, Lim-Tio S, Strauss B, Stockigt J, Topliss D, Alford F, Hew L, Bode H, Conway A, Handelsman D, Dunn S, Boyages S, Cheung NW, Hurley D (1998) The Australian Multicenter Trial of Growth Hormone (GH) Treatment in GH deficient adults. *J Clin Endocrinol Metab* 83:107–116
- Ellis KJ, Shypailo RJ, Pratt JA, Pond WG (1994) Accuracy of dual-energy x-ray absorptiometry for body-composition measurements in children. *Am J Clin Nutr* 60:660–665
- Ellis KJ, Shypailo RJ, Pivarnik JM, Jenks BH, Walzel P, Lee PDK (1995) Body composition of HIV-infected male adults with wasting syndrome. *Asia Pacific J Clin Nutr* 4:105–108
- Goulding A, Gold E, Cannan R, Taylor RW, Williams S, Lewis-Barned NJ (1996) DEXA supports the use of BMI as a measure of fatness in young girls. *Int J Obes Relat Disord* 20:1014–1021
- Gutin B, Litaker M, Islam S, Manos T, Smith C, Treiber F (1996) Body-composition measurement in 9–11-y-old children by dual-energy X-ray absorptiometry, skinfold-thickness measurements, and bioimpedance analysis. *Am J Clin Nutr* 63:287–292

16. Haarbo J, Gotfredsen A, Hassager C, Christiansen C (1991) Validation of body composition by dual energy X-ray absorptiometry (DEXA). *Clin Physiol* 11:331–341
17. Hansen TB, Vahl N, Jorgensen JOL, Christiansen JS, Hagen C (1995) Whole body and regional soft tissue changes in growth hormone deficient adults after one year of growth hormone treatment: a double-blind, randomized, placebo-controlled study. *Clin Endocrinol* 43:689–696
18. Houtkooper LB, Ritenbaugh C, Aickin M, Lohman TG, Going SB, Weber JL, Greaves KA, Boyden TW, Pamenter RW, Hall MC (1995) Nutrients, body composition and exercise are related to change in bone mineral density in premenopausal women. *J Nutr* 125:1229–1237
19. Jensen MD, Kanaley JA, Roust LR, O'Brien PC, Braun JS, Dunn WL, Wahner HW (1993) Assessment of body composition with use of dual-energy X-ray absorptiometry: evaluation and comparison with other methods. *Mayo Clin Proc* 68:867–873
20. Jensen MD, Kanaley JA, Reed JE, Sheedy PF (1995) Measurement of abdominal and visceral fat with computed tomography and dual-energy x-ray absorptiometry. *Am J Clin Nutr* 61:274–278
21. Lindholm C, Hagenfeldt K, Ringertz H (1995) Bone mineral content of young female former gymnasts. *Acta Paediatr* 84:1109–1112
22. Lonzer MD, Imrie R, Rogers D, Worley D, Licata A, Secic M (1996) Effects of heredity, age, weight, puberty, activity, and calcium intake on bone mineral density in children. *Clin Pediatr (Phila)* 35:185–189
23. Manzoni P, Brambilla P, Simone P, Salvatoni A, Beccaria L, Chiumello G (1993) Total and regional fat in obese children and Prader Willi subjects (PWS) using dual X-ray absorptiometry. *Int J Obes* 17 [Suppl 2]:22
24. Manzoni P, Brambilla P, Pietrobelli A, Beccaria L, Bianchessi A, Mora S, Chiumello G (1996) Influence of body composition on bone mineral content in children and adolescents. *Am J Clin Nutr* 64:603–607
25. Mazess RB, Barden HS, Bisek JP, Hanson J (1990) Dual energy X-ray absorptiometry for total and regional bone-mineral and soft-tissue composition. *Am J Clin Nutr* 51:1106–1112
26. Molgaard C, Thomsen BL, Prentice A, Cole TJ, Michaelsen KF (1997) Whole body bone mineral content in healthy children and adolescents. *Arch Dis Child* 76:9–15
27. Pintauro SJ, Nagy TR, Duthie CM, Goran MI (1996) Cross-calibration of fat and lean measurements by dual-energy x-ray absorptiometry to pig carcass analysis in the pediatric body weight range. *Am J Clin Nutr* 63:293–298
28. Prentice A, Parsons TJ, Cole TJ (1994) Uncritical use of bone mineral density in absorptiometry may lead to size-related artefacts in the identification of bone determinants. *Am J Clin Nutr* 60:837–842
29. Rosenfalck AM, Almdal T, Gotfredsen A, Hojgaard LL, Hilsted J (1995) Validity of dual X-ray absorptiometry scanning for determination of body composition in IDDM patients. *Scand J Clin Lab Invest* 55:691–699
30. Salvatoni A, Gambarini G, Bosetti G, Riganti G, Marchini P, Abbiati B, Paolucci M, Broggin M (1992) Modification of body composition after PSMF. *Int J Obes [Suppl 1]* 16:48
31. Salvatoni A, Broggin M, Salvatore S, Cortona L, Ghiringhelli D, Marchini P, Nespoli L (1995) Body composition assessed by DEXA in children with IDDM. *J Pediatr Endocrinol Metab [Suppl]* 8:230
32. Schlamp D, Schneider P, Krah A, Trott GE, Warnke A (1994) Bone density in anorexia nervosa – with comment on prevention and therapy of osteoporosis. *Z Kinder Jugendpsychiatr* 22:183–188
33. Scotta MS, Salvatore S, Salvatoni A, DeAmici M, Ghiringhelli D, Broggin M, Nespoli L (1997) Bone mineralization and body composition in young patients with celiac disease. *Am J Gastroenterol* 92:1331–1334
34. Spector E, LeBlanc A, Shackelford L (1995) Hologic QDR 2000 whole-body scans: a comparison of three combinations of scan modes and analysis software. *Osteoporosis Int* 5:440–445
35. Svendsen OL, Haarbo J, Hassager C, Christiansen C (1993) Accuracy of measurements of body composition by dual energy x-ray absorptiometry in vivo. *Am J Clin Nutr* 57:605–608
36. Svendsen OL, Hassager C, Christiansen C (1995) The response to treatment of overweight in postmenopausal women is not related to fat distribution. *Int J Obes* 19:496–502
37. Treuth MS, Hunter GR, Kekes-Szabo T (1995) Estimating intraabdominal adipose tissue in women by dual-energy x-ray absorptiometry. *Am J Clin Nutr* 62:527–532
38. VanMarcken Lichtenbelt WD, Fogelholm M, Ottenheijm R, Westerterp KR (1995) Physical activity, body composition and bone density in ballet dancers. *Br J Nutr* 74:439–451
39. Warner JT, Cowan FJ, Dunstan FD, Gregory JW (1997) The validity of body mass index for assessment of adiposity in children with disease states. *Ann Hum Biol* 24:209–215