

AURELIUS / CXO — Human-Readable Protocol (All Phases)

Version source: AURELIUS_CXO_v1.29.xlsx • Generated: 2026-01-06

This document is a self-contained operational guide. It lists what to do/take by Phase and Time Block, including key constraints, maintenance notes, and major operator practices (mobility, light, breathwork). It is not medical advice.

Governance & Rules of Use

- Follow phases sequentially unless you have an explicit, logged reason to deviate.
- Change one variable at a time when assessing response (5–7 days between meaningful changes).
- Binder window is a 'clean interval'—avoid stacking supplements inside it unless explicitly listed there.
- If a STOP/ADJUST trigger is hit (sleep collapse, major GI worsening, pressure headaches, agitation), revert to the last stable baseline.

Phase 1 quick anchor (terrain restoration)

EXPECTED PROGRESSION

Week 1: Digestion improves

Week 2: SD reduces, energy stabilizes

Week 3: Smell improves, libido flicker, better sleep

PHASE 1 EXIT CRITERIA

- Stable digestion
- No persistent nausea
- Good sleep
- No bloating
- 1–2 days/week improved well-being

DO NOT USE (PHASE 1 RESTRICTIONS)

- Tongkat, Cistanche, dopamine agents
- PDE5 inhibitors
- Angion protocol
- High-dose Vit D
- DIM / I3C
- Choline supplements
- Alcohol
- Fasting >12h

Phase 1.0

Intent: reduce endotoxin/aldehyde load, stabilize digestion, bile rhythm, sleep, and baseline energy.

On waking

Do / Take

- Hydration Baseline [COP-23]

Source IDs: COP-23

Do / Take

- Black Coffee — 1–3 cups/day [CAS-FOOD-03]

Optional

- Optional

Source IDs: CAS-FOOD-03

Binder window (fasted, mid-morning)

Do / Take

- Psyllium — 5-10g [CAS-GUT-01]

Continuation / maintenance: Carry: Yes | Maint: 2–4×/week or PRN | Stop/Taper: Taper to PRN once stools stable 10–14d; reduce first if constipation.

Source IDs: CAS-GUT-01

Binder window (after psyllium)

Do / Take

- Carrot salad [CAS-FOOD-01]

Continuation / maintenance: Carry: Yes | Maint: 3–5×/week or PRN | Stop/Taper: Maintenance once GI/skin stable 10–14d.

Source IDs: CAS-FOOD-01

Post-binder meal (first meal)

Do / Take

- Taurine — 1g [CAS-LIV-01]

Continuation / maintenance: Carry: Yes | Maint: Daily or 3–5×/week | Stop/Taper: Reduce frequency once bile/GI stable 10–14d.

Source IDs: CAS-LIV-01

Do / Take

- MSM — 750mg [CAS-LIV-02]

Continuation / maintenance: Carry: Yes | Maint: 3–5×/week → discontinue | Stop/Taper: Taper off once skin/gut stable 14d; stop if sulfur intolerance/wired.

Source IDs: CAS-LIV-02

Do / Take

- Molybdenum — 100-200mcg [CAS-LIV-03]

Continuation / maintenance: Carry: Yes | Maint: Daily with MSM; optional otherwise | Stop/Taper: Keep while MSM used; optional after MSM stopped.

Source IDs: CAS-LIV-03

Do / Take

- Beet extract — per label [CAS-VAS-08]

Continuation / maintenance: Carry: PRN | Maint: PRN (training/sex days) | Stop/Taper: Use PRN; stop if low BP/dizziness.

Source IDs: CAS-VAS-08

Do / Take

- Vitamin C — 1g [CAS-ANTIOX-01]

Continuation / maintenance: Carry: Yes | Maint: Daily or PRN with vascular block | Stop/Taper: Optional long-term; reduce if GI irritation.

Source IDs: CAS-ANTIOX-01

Do / Take

- Zinc — 20-30mg [CAS-MICRO-01]

Continuation / maintenance: Carry: Course | Maint: 2–4×/week maintenance or stop | Stop/Taper: Reassess after 6–8 weeks; avoid long-term high-dose without monitoring.

Source IDs: CAS-MICRO-01

Do / Take

- Selenium — 100-200mcg [CAS-MICRO-02]

Continuation / maintenance: Carry: Course | Maint: 2–4×/week maintenance or stop | Stop/Taper: Reassess after 8–12 weeks.

Source IDs: CAS-MICRO-02

With meals (split across meals)

Do / Take

- Undecylenic Acid — titrate 1→5 caps/day [CAS-FUNG-01]

Continuation / maintenance: Carry: Course | Maint: Course; PRN pulses | Stop/Taper: Finite course: taper off after fungal/SD stable 14d; PRN pulses for relapse.

Source IDs: CAS-FUNG-01

Lunch (with food)

Do / Take

- Creatine — 5g [CAS-MITO-01]

Continuation / maintenance: Carry: Yes | Maint: Daily | Stop/Taper: Long-term OK.

Source IDs: CAS-MITO-01

Dinner (with food)

Do / Take

- TUDCA — 250mg — Daily (from Week 2) [CAS-LIV-04]

Continuation / maintenance: Carry: Yes | Maint: 2–4×/week maintenance | Stop/Taper: Taper once motility + bile symptoms stable 10–14d; hold if bile reflux/RUQ pain.

Source IDs: CAS-LIV-04

Bedtime (PM)

Do / Take

- Magnesium glycinate — 200-300mg [CAS-MICRO-03]

Continuation / maintenance: Carry: Yes | Maint: Daily | Stop/Taper: Long-term OK.

Source IDs: CAS-MICRO-03

Do / Take

- Glycine — 2-3g [CAS-AMINO-01]

Continuation / maintenance: Carry: Yes | Maint: Daily or PRN | Stop/Taper: Optional long-term; reduce if daytime sedation.

Source IDs: CAS-AMINO-01

Practices (today)

Do / Take

- Reverse Kegels — Daily — Any, especially pre/ post-sex [COP-01]

Source IDs: COP-01

Do / Take

- Psoas Release — 3–7x/week — Evening or off days [COP-02]

Source IDs: COP-02

Do / Take

- Adductor & Hamstring Lengthening — 3–5x/week — Post-walk or training [COP-03]

Source IDs: COP-03

Do / Take

- Pelvic Mobility Drills — Daily — AM or PM [COP-04]

Source IDs: COP-04

Do / Take

- Nasal PBM — 3–7x/week — Evening or morning [COP-06]

Source IDs: COP-06

Do / Take

- Diaphragmatic Breathwork — Daily — Any, especially pre-sleep [COP-12]

Source IDs: COP-12

Do / Take

- Glute Activation Sequences — 3–5x/week — Pre-walk or training [COP-13]

Source IDs: COP-13

Do / Take

- Sleep Optimization Practices — Daily — Evening [COP-14]

Source IDs: COP-14

Do / Take

- Liver Pumping / Thoracic Mobilization — 3–7x/week — Evening [COP-15]

Source IDs: COP-15

Do / Take

- Vagal Reset Maneuvers — PRN + daily micro-practices — Any [COP-16]

Optional

- Optional

Source IDs: COP-16

Do / Take

- Hydration Baseline — Daily — On waking + throughout day [COP-23]

Source IDs: COP-23

Watch-for / STOP-ADJUST

Do / Take

- Avoid choline, lecithin, ox bile, DIM, I3C, high-dose vitamin D, aggressive training.

Do / Take

- Avoid starting at very high dose

Do / Take

- caution if severe herx or constipation.

Do / Take

- Avoid if severe diarrhea, strong bile reflux sensations, or new right-upper-quadrant pain.

Advance when

Do / Take

- Stools more regular, nausea absent, energy modestly improved, SD beginning to ease, no worsening of symptoms.

Phase 1.5

Intent: continue terrain stabilization while introducing gentle anti-inflammatory/lipid and support rails.

On waking

Do / Take

- Hydration Baseline [COP-23]

Source IDs: COP-23

Do / Take

- Black Coffee — 1–3 cups/day [CAS-FOOD-03]

Optional

- Optional

Source IDs: CAS-FOOD-03

Binder window (fasted, mid-morning)

Do / Take

- Psyllium — 5-10g [CAS-GUT-01]

Continuation / maintenance: Carry: Yes | Maint: 2–4×/week or PRN | Stop/Taper: Taper to PRN once stools stable 10–14d; reduce first if constipation.

Source IDs: CAS-GUT-01

Binder window (after psyllium)

Do / Take

- Carrot salad [CAS-FOOD-01]

Continuation / maintenance: Carry: Yes | Maint: 3–5×/week or PRN | Stop/Taper: Maintenance once GI/skin stable 10–14d.

Source IDs: CAS-FOOD-01

Post-binder meal (first meal)

Do / Take

- Taurine — 1g [CAS-LIV-01]

Continuation / maintenance: Carry: Yes | Maint: Daily or 3–5×/week | Stop/Taper: Reduce frequency once bile/GI stable 10–14d.

Source IDs: CAS-LIV-01

Do / Take

- MSM — 750mg [CAS-LIV-02]

Continuation / maintenance: Carry: Yes | Maint: 3–5×/week → discontinue | Stop/Taper: Taper off once skin/gut stable 14d; stop if sulfur intolerance/wired.

Source IDs: CAS-LIV-02

Do / Take

- Molybdenum — 100-200mcg [CAS-LIV-03]

Continuation / maintenance: Carry: Yes | Maint: Daily with MSM; optional otherwise | Stop/Taper: Keep while MSM used; optional after MSM stopped.

Source IDs: CAS-LIV-03

Do / Take

- Beet extract — per label [CAS-VAS-08]

Continuation / maintenance: Carry: PRN | Maint: PRN (training/sex days) | Stop/Taper: Use PRN; stop if low BP/dizziness.

Source IDs: CAS-VAS-08

Do / Take

- Vitamin C — 1g [CAS-ANTIOX-01]

Continuation / maintenance: Carry: Yes | Maint: Daily or PRN with vascular block | Stop/Taper: Optional long-term; reduce if GI irritation.

Source IDs: CAS-ANTIOX-01

Do / Take

- Zinc — 20-30mg [CAS-MICRO-01]

Continuation / maintenance: Carry: Course | Maint: 2–4×/week maintenance or stop | Stop/Taper: Reassess after 6–8 weeks; avoid long-term high-dose without monitoring.

Source IDs: CAS-MICRO-01

Do / Take

- Selenium — 100-200mcg [CAS-MICRO-02]

Continuation / maintenance: Carry: Course | Maint: 2–4×/week maintenance or stop | Stop/Taper: Reassess after 8–12 weeks.

Source IDs: CAS-MICRO-02

Do / Take

- Benfotiamine — 150-300mg [CAS-MITO-05]

Continuation / maintenance: Carry: Yes | Maint: 3–5×/week maintenance | Stop/Taper: Taper once energy stable 14d.

Source IDs: CAS-MITO-05

Do / Take

- ALCAR — 500mg [CAS-MITO-06]

Continuation / maintenance: Carry: PRN | Maint: PRN (training/drive days) | Stop/Taper: Use PRN; stop if insomnia/wired.

Source IDs: CAS-MITO-06

Do / Take

- Calcium D-Glucarate — 500mg [CAS-GUT-02]

Continuation / maintenance: Carry: PRN | Maint: 2–4×/week or PRN | Stop/Taper: Use during mobilization; taper when stable 14d.

Source IDs: CAS-GUT-02

Do / Take

- PHGG — 1-2g [CAS-GUT-03]

Continuation / maintenance: Carry: Yes | Maint: 3–5×/week maintenance | Stop/Taper: Taper once stools stable 10–14d.

Source IDs: CAS-GUT-03

Do / Take

- MegaSporeBiotic — 0.5→1 cap [CAS-GUT-04]

Continuation / maintenance: Carry: Course | Maint: Stop after course | Stop/Taper: Course 8–12 weeks then stop and reassess.

Source IDs: CAS-GUT-04

Do / Take

- Glutathione (oral, generic) — 250–500mg — Daily (optional) [CAS-DET-01]

Optional

- Optional

Continuation / maintenance: Carry: PRN | Maint: 2–4×/week or short blocks | Stop/Taper: Optional: use in short blocks if tolerated.

Source IDs: CAS-DET-01

Do / Take

- NAC — 600mg [CAS-BIO-02]

Optional

- Optional

Continuation / maintenance: Carry: PRN | Maint: 2–4×/week or short blocks | Stop/Taper: Optional: use in short blocks; stop if wired/insomnia.

Source IDs: CAS-BIO-02

Do / Take

- Sulforaphane (Broccoli Sprout Extract) — per label (e.g. 20–40mg) — 3–5×/week [CAS-INF-02]

Continuation / maintenance: Carry: PRN | Maint: 3–5×/week cycles | Stop/Taper: Optional/cycled; reduce frequency if headache/insomnia.

Source IDs: CAS-INF-02

Do / Take

- Methylene Blue (low-dose) — 0.5–1mg/kg — 3–5×/week [CAS-MITO-15]

Continuation / maintenance: Carry: PRN | Maint: 3–5×/week cycles (never continuous) | Stop/Taper: Cycled only; stop if insomnia/wired.

Source IDs: CAS-MITO-15

With meals (split across meals)

Do / Take

- Undecylenic Acid — titrate 1→5 caps/day [CAS-FUNG-01]

Continuation / maintenance: Carry: Course | Maint: Course; PRN pulses | Stop/Taper: Finite course: taper off after fungal/SD stable 14d; PRN pulses for relapse.

Source IDs: CAS-FUNG-01

Do / Take

- Niacinamide — 250–500mg [CAS-B-01]

Continuation / maintenance: Carry: Yes | Maint: Daily with food or 3–5×/week | Stop/Taper: Reduce if over-sedation or GI effects.

Source IDs: CAS-B-01

Lunch (with food)

Do / Take

- Creatine — 5g [CAS-MITO-01]

Continuation / maintenance: Carry: Yes | Maint: Daily | Stop/Taper: Long-term OK.

Source IDs: CAS-MITO-01

Do / Take

- Aspirin — 81mg — EOD [CAS-INF-01]

Continuation / maintenance: Carry: Course | Maint: EOD course then reassess | Stop/Taper: Course; taper off once inflammation stable; stop if bruising/GI issues.

Source IDs: CAS-INF-01

Do / Take

- Vitamin K2 — MK7 100-200mcg [CAS-MICRO-04]

Continuation / maintenance: Carry: Yes | Maint: Daily with food | Stop/Taper: Long-term OK.

Source IDs: CAS-MICRO-04

Do / Take

- Omega-3 — 500mg [CAS-LIP-01]

Continuation / maintenance: Carry: Yes | Maint: Daily with food | Stop/Taper: Long-term OK.

Source IDs: CAS-LIP-01

Dinner (with food)

Do / Take

- TUDCA — 250mg — Daily (from Week 2) [CAS-LIV-04]

Continuation / maintenance: Carry: Yes | Maint: 2–4×/week maintenance | Stop/Taper: Taper once motility + bile symptoms stable 10–14d; hold if bile reflux/RUQ pain.

Source IDs: CAS-LIV-04

Bedtime (PM)

Do / Take

- Magnesium glycinate — 200-300mg [CAS-MICRO-03]

Continuation / maintenance: Carry: Yes | Maint: Daily | Stop/Taper: Long-term OK.

Source IDs: CAS-MICRO-03

Do / Take

- Glycine — 2-3g [CAS-AMINO-01]

Continuation / maintenance: Carry: Yes | Maint: Daily or PRN | Stop/Taper: Optional long-term; reduce if daytime sedation.

Source IDs: CAS-AMINO-01

Do / Take

- ALA — 100-200mg [CAS-MITO-07]

Continuation / maintenance: Carry: PRN | Maint: 2–4×/week or cycles | Stop/Taper: Optional/cycled; stop if insomnia/wired.

Source IDs: CAS-MITO-07

Do / Take

- DHM — 100-300mg [CAS-NEURO-10]

Continuation / maintenance: Carry: PRN | Maint: PRN (evening) | Stop/Taper: PRN for aldehyde load; stop if daytime sedation.

Source IDs: CAS-NEURO-10

Practices (today)

Do / Take

- Reverse Kegels — Daily — Any, especially pre/ post-sex [COP-01]

Source IDs: COP-01

Do / Take

- Psoas Release — 3–7x/week — Evening or off days [COP-02]

Source IDs: COP-02

Do / Take

- Adductor & Hamstring Lengthening — 3–5x/week — Post-walk or training [COP-03]

Source IDs: COP-03

Do / Take

- Pelvic Mobility Drills — Daily — AM or PM [COP-04]

Source IDs: COP-04

Do / Take

- Nasal PBM — 3–7x/week — Evening or morning [COP-06]

Source IDs: COP-06

Do / Take

- Fascia & Pelvic Floor Neuro-Mechanics — 3–5x/week — Any [COP-07]

Source IDs: COP-07

Do / Take

- Sauna / Cold Contrast — 1–4x/week — Evening [COP-11]

Source IDs: COP-11

Do / Take

- Diaphragmatic Breathwork — Daily — Any, especially pre-sleep [COP-12]

Source IDs: COP-12

Do / Take

- Glute Activation Sequences — 3–5x/week — Pre-walk or training [COP-13]

Source IDs: COP-13

Do / Take

- Sleep Optimization Practices — Daily — Evening [COP-14]

Source IDs: COP-14

Do / Take

- Liver Pumping / Thoracic Mobilization — 3–7x/week — Evening [COP-15]

Source IDs: COP-15

Do / Take

- Vagal Reset Maneuvers — PRN + daily micro-practices — Any [COP-16]

Optional

- Optional

Source IDs: COP-16

Do / Take

- Foot–Core Activation Chain — 3–5x/week — Pre-walk/training [COP-17]

Source IDs: COP-17

Do / Take

- Olfactory Training Protocol — Daily — Any [COP-19]

Source IDs: COP-19

Do / Take

- CO₂ Tolerance Breathwork — 3–5x/week — Any [COP-20]

Source IDs: COP-20

Do / Take

- Hydration Baseline — Daily — On waking + throughout day [COP-23]

Source IDs: COP-23

Watch-for / STOP-ADJUST

Do / Take

- Avoid choline, lecithin, ox bile, DIM, I3C, high-dose vitamin D, aggressive training.

Do / Take

- Avoid starting at very high dose

Do / Take

- caution if severe herx or constipation.

Do / Take

- Avoid if severe diarrhea, strong bile reflux sensations, or new right-upper-quadrant pain.

Do / Take

- Avoid stacking new stimulants

Do / Take

- continue to avoid choline, ox bile and aggressive DIM/I3C usage.

Do / Take

- Caution in severe liver disease

Do / Take

- watch for unusual fatigue or paradoxical agitation.

Do / Take

- May provoke mild detox/headache in sensitive individuals

Do / Take

- start at lower frequency if needed.

Do / Take

- Avoid with serotonergic drugs (SSRIs, MAOIs)

Do / Take

- caution in G6PD deficiency

Do / Take

- may discolor urine/teeth.

Advance when

Do / Take

- Bloating and gut reactivity reduced, sleep stable, baseline libido flickers returning, stress tolerance better.

Phase 2.0

Intent: cautiously add neuroendocrine / vascular support while keeping terrain stability.

On waking

Do / Take

- Hydration Baseline [COP-23]

Source IDs: COP-23

Do / Take

- Black Coffee — 1–3 cups/day [CAS-FOOD-03]

Optional

- Optional

Source IDs: CAS-FOOD-03

Binder window (fasted, mid-morning)

Do / Take

- Psyllium — 5-10g [CAS-GUT-01]

Continuation / maintenance: Carry: Yes | Maint: 2–4×/week or PRN | Stop/Taper: Taper to PRN once stools stable 10–14d; reduce first if constipation.

Source IDs: CAS-GUT-01

Binder window (after psyllium)

Do / Take

- Carrot salad [CAS-FOOD-01]

Continuation / maintenance: Carry: Yes | Maint: 3–5×/week or PRN | Stop/Taper: Maintenance once GI/skin stable 10–14d.

Source IDs: CAS-FOOD-01

Post-binder meal (first meal)

Do / Take

- Taurine — 1g [CAS-LIV-01]

Continuation / maintenance: Carry: Yes | Maint: Daily or 3–5×/week | Stop/Taper: Reduce frequency once bile/GI stable 10–14d.

Source IDs: CAS-LIV-01

Do / Take

- MSM — 750mg [CAS-LIV-02]

Continuation / maintenance: Carry: Yes | Maint: 3–5×/week → discontinue | Stop/Taper: Taper off once skin/gut stable 14d; stop if sulfur intolerance/wired.

Source IDs: CAS-LIV-02

Do / Take

- Molybdenum — 100-200mcg [CAS-LIV-03]

Continuation / maintenance: Carry: Yes | Maint: Daily with MSM; optional otherwise | Stop/Taper: Keep while MSM used; optional after MSM stopped.

Source IDs: CAS-LIV-03

Do / Take

- Zinc — 20-30mg [CAS-MICRO-01]

Continuation / maintenance: Carry: Course | Maint: 2–4×/week maintenance or stop | Stop/Taper: Reassess after 6–8 weeks; avoid long-term high-dose without monitoring.

Source IDs: CAS-MICRO-01

Do / Take

- Selenium — 100-200mcg [CAS-MICRO-02]

Continuation / maintenance: Carry: Course | Maint: 2–4×/week maintenance or stop | Stop/Taper: Reassess after 8–12 weeks.

Source IDs: CAS-MICRO-02

Do / Take

- Benfotiamine — 150-300mg [CAS-MITO-05]

Continuation / maintenance: Carry: Yes | Maint: 3–5×/week maintenance | Stop/Taper: Taper once energy stable 14d.

Source IDs: CAS-MITO-05

Do / Take

- Calcium D-Glucarate — 500mg [CAS-GUT-02]

Continuation / maintenance: Carry: PRN | Maint: 2–4×/week or PRN | Stop/Taper: Use during mobilization; taper when stable 14d.

Source IDs: CAS-GUT-02

Do / Take

- PHGG — 1-2g [CAS-GUT-03]

Continuation / maintenance: Carry: Yes | Maint: 3–5×/week maintenance | Stop/Taper: Taper once stools stable 10–14d.

Source IDs: CAS-GUT-03

Do / Take

- MegaSporeBiotic — 0.5→1 cap [CAS-GUT-04]

Continuation / maintenance: Carry: Course | Maint: Stop after course | Stop/Taper: Course 8–12 weeks then stop and reassess.

Source IDs: CAS-GUT-04

Do / Take

- Glutathione (oral, generic) — 250–500mg — Daily (optional) [CAS-DET-01]

Optional

- Optional

Continuation / maintenance: Carry: PRN | Maint: 2–4×/week or short blocks | Stop/Taper: Optional: use in short blocks if tolerated.

Source IDs: CAS-DET-01

Do / Take

- NAC — 600mg [CAS-BIO-02]

Optional

- Optional

Continuation / maintenance: Carry: PRN | Maint: 2–4×/week or short blocks | Stop/Taper: Optional: use in short blocks; stop if wired/insomnia.

Source IDs: CAS-BIO-02

Do / Take

- Sulforaphane (Broccoli Sprout Extract) — per label (e.g. 20–40mg) — 3–5×/week [CAS-INF-02]

Continuation / maintenance: Carry: PRN | Maint: 3–5×/week cycles | Stop/Taper: Optional/cycled; reduce frequency if headache/insomnia.

Source IDs: CAS-INF-02

Do / Take

- Methylene Blue (low-dose) — 0.5–1mg/kg — 3–5×/week [CAS-MITO-15]

Continuation / maintenance: Carry: PRN | Maint: 3–5×/week cycles (never continuous) | Stop/Taper: Cycled only; stop if insomnia/wired.

Source IDs: CAS-MITO-15

Do / Take

- Tongkat Ali — 200-400mg [CAS-ANDRO-01]

Continuation / maintenance: Carry: Yes | Maint: 5 on / 2 off or blocks | Stop/Taper: Cycle to avoid tolerance; stop if irritability/insomnia.

Source IDs: CAS-ANDRO-01

Do / Take

- Cistanche — 500-1500mg [CAS-ANDRO-02]

Continuation / maintenance: Carry: Yes | Maint: 5 on / 2 off or blocks | Stop/Taper: Cycle to avoid tolerance; stop if irritability/insomnia.

Source IDs: CAS-ANDRO-02

Do / Take

- Pregnenolone — 5-10mg [CAS-STERO-01]

Continuation / maintenance: Carry: Yes | Maint: Daily or 5 on / 2 off | Stop/Taper: Reassess with symptoms/labs; stop if insomnia/anxiety.

Source IDs: CAS-STERO-01

Do / Take

- DHEA — 5mg [CAS-STERO-02]

Continuation / maintenance: Carry: Yes | Maint: Daily or 5 on / 2 off | Stop/Taper: Reassess with symptoms/labs; stop if acne/irritability/insomnia.

Source IDs: CAS-STERO-02

Do / Take

- SAM-e — 200mg [CAS-MET-01]

Continuation / maintenance: Carry: PRN | Maint: 3–5×/week or PRN | Stop/Taper: Optional; stop if agitation/insomnia.

Source IDs: CAS-MET-01

Do / Take

- 7,8-Dihydroxyflavone (7,8-DHF) — 10–40mg — Daily (10 days on / 10 days off) – optional [CAS-NEURO-05]

Optional

- Optional

Continuation / maintenance: Carry: PRN | Maint: 10 on / 10 off | Stop/Taper: Optional/cycled; stop if insomnia/anxiety.

Source IDs: CAS-NEURO-05

Do / Take

- TTFD (Thiamine Tetrahydrofurfuryl Disulfide) — 50–200mg [CAS-MITO-06]

Continuation / maintenance: Carry: PRN | Maint: PRN (training/drive days) | Stop/Taper: Use PRN; stop if insomnia/wired.

Source IDs: CAS-MITO-06

Do / Take

- Boron — 6–9mg [CAS-MIN-01]

Continuation / maintenance: Carry: Yes | Maint: 2 weeks on / 1 off | Stop/Taper: As written: cycle.

Source IDs: CAS-MIN-01

Do / Take

- Uridine Monophosphate — 50–150mg [CAS-NEURO-08]

Continuation / maintenance: Carry: PRN | Maint: Cycles only | Stop/Taper: Optional/cycled.

Source IDs: CAS-NEURO-08

Do / Take

- Piracetam — 1200–2400mg [CAS-NOOT-01]

Continuation / maintenance: Carry: PRN | Maint: Cycles only | Stop/Taper: Optional/cycled.

Source IDs: CAS-NOOT-01

Do / Take

- Sulbuthiamine — 200–400mg — 5 days on / 2 off [CAS-NEURO-09]

Continuation / maintenance: Carry: PRN | Maint: 5 on / 2 off | Stop/Taper: Cycled only (as written).

Source IDs: CAS-NEURO-09

Do / Take

- Methylated B-Complex — Label-equivalent [CAS-VIT-05]

Continuation / maintenance: Carry: Yes | Maint: 3–5×/week or daily | Stop/Taper: Optional long-term; reduce if overstimulation.

Source IDs: CAS-VIT-05

Do / Take

- Semax — 0.1–0.3mg IN — 5 days on / 2 off [CAS-PEPT-02]

Continuation / maintenance: Carry: PRN | Maint: 5 on / 2 off | Stop/Taper: Cycled only; stop if agitation/insomnia.

Source IDs: CAS-PEPT-02

Do / Take

- Mucuna Pruriens — 100–200mg — 10 on / 20 off [CAS-NEURO-10]

Continuation / maintenance: Carry: PRN | Maint: PRN (evening) | Stop/Taper: PRN for aldehyde load; stop if daytime sedation.

Source IDs: CAS-NEURO-10

Do / Take

- HMB (β -Hydroxy β -Methylbutyrate) — 1.5 g/day [CAS-NEURO-14]

Continuation / maintenance: Carry: PRN | Maint: Training/sex days or blocks | Stop/Taper: Use with food; choose one HMB lane; stop if GI upset.

Source IDs: CAS-NEURO-14

With meals (split across meals)

Do / Take

- Undecylenic Acid — titrate 1→5 caps/day [CAS-FUNG-01]

Continuation / maintenance: Carry: Course | Maint: Course; PRN pulses | Stop/Taper: Finite course: taper off after fungal/SD stable 14d; PRN pulses for relapse.

Source IDs: CAS-FUNG-01

Do / Take

- Niacinamide — 250–500mg [CAS-B-01]

Continuation / maintenance: Carry: Yes | Maint: Daily with food or 3–5×/week | Stop/Taper: Reduce if over-sedation or GI effects.

Source IDs: CAS-B-01

Do / Take

- Endotelon (grape seed OPC) — 150–300mg/day (often 150mg BID) [CAS-VAS-15]

Continuation / maintenance: Carry: Course | Maint: Block 8–12 w | Stop/Taper: Block 8–12 weeks then reassess; avoid stacking with aspirin/polyphenols if bruising.

Source IDs: CAS-VAS-15

Do / Take

- Tributyrin — 300–600mg [CAS-GUT-07]

Continuation / maintenance: Carry: Yes | Maint: 3–5×/week | Stop/Taper: Taper once stools stable 14d.

Source IDs: CAS-GUT-07

Do / Take

- Lactobacillus reuteri — 1–2B CFU [CAS-PRO-02]

Continuation / maintenance: Carry: Course | Maint: Stop after course or cycle | Stop/Taper: Course 8–12 weeks then reassess.

Source IDs: CAS-PRO-02

Lunch (with food)

Do / Take

- Creatine — 5g [CAS-MITO-01]

Continuation / maintenance: Carry: Yes | Maint: Daily | Stop/Taper: Long-term OK.

Source IDs: CAS-MITO-01

Do / Take

- Aspirin — 81mg — EOD [CAS-INF-01]

Continuation / maintenance: Carry: Course | Maint: EOD course then reassess | Stop/Taper: Course; taper off once inflammation stable; stop if bruising/GI issues.

Source IDs: CAS-INF-01

Do / Take

- Vitamin K2 — MK7 100-200mcg [CAS-MICRO-04]

Continuation / maintenance: Carry: Yes | Maint: Daily with food | Stop/Taper: Long-term OK.

Source IDs: CAS-MICRO-04

Do / Take

- Omega-3 — 500mg [CAS-LIP-01]

Continuation / maintenance: Carry: Yes | Maint: Daily with food | Stop/Taper: Long-term OK.

Source IDs: CAS-LIP-01

Do / Take

- LCLT — 1-2g [CAS-MITO-02]

Continuation / maintenance: Carry: Yes | Maint: Daily with food | Stop/Taper: Long-term OK; stop if fishy odor or GI intolerance.

Source IDs: CAS-MITO-02

Do / Take

- Reduced Glutathione (liposomal) — 250–500mg — Daily (optional) [CAS-DET-02]

Optional

- Optional

Continuation / maintenance: Carry: PRN | Maint: 2–4×/week or blocks | Stop/Taper: Optional mitochondrial/antioxidant support; use in short blocks if tolerated.

Source IDs: CAS-DET-02

Dinner (with food)

Do / Take

- TUDCA — 250mg — Daily (from Week 2) [CAS-LIV-04]

Continuation / maintenance: Carry: Yes | Maint: 2–4×/week maintenance | Stop/Taper: Taper once motility + bile symptoms stable 10–14d; hold if bile reflux/RUQ pain.

Source IDs: CAS-LIV-04

Pre-workout / pre-sex (acute signaling)

Do / Take

- Citrulline — 6g [CAS-VAS-01]

Continuation / maintenance: Carry: PRN | Maint: Training/sex days only | Stop/Taper: Acute tool for training/sex days.

Source IDs: CAS-VAS-01

Do / Take

- Icariin — 150-200mg [CAS-VAS-02]

Continuation / maintenance: Carry: PRN | Maint: Training/sex days only | Stop/Taper: Acute tool for training/sex days.

Source IDs: CAS-VAS-02

Do / Take

- Histidine — 500-1000mg [CAS-VAS-03]

Continuation / maintenance: Carry: PRN | Maint: PRN | Stop/Taper: Optional acute adjunct; stop if histamine symptoms.

Source IDs: CAS-VAS-03

Do / Take

- Hesperidin — 250mg [CAS-VAS-04]

Continuation / maintenance: Carry: PRN | Maint: PRN | Stop/Taper: Optional adjunct; low priority if low BP.

Source IDs: CAS-VAS-04

Do / Take

- Pycnogenol — 50-100mg [CAS-VAS-05]

Continuation / maintenance: Carry: PRN | Maint: PRN or blocks | Stop/Taper: Prefer as block (8–12 w) if used chronically; watch bleeding with aspirin.

Source IDs: CAS-VAS-05

Do / Take

- Vitamin C — 500-1000mg [CAS-ANTIOX-01]

Continuation / maintenance: Carry: Yes | Maint: Daily or PRN with vascular block | Stop/Taper: Optional long-term; reduce if GI irritation.

Source IDs: CAS-ANTIOX-01

Do / Take

- Beet nitrate — 300-600mg [CAS-VAS-08]

Continuation / maintenance: Carry: PRN | Maint: PRN (training/sex days) | Stop/Taper: Use PRN; stop if low BP/dizziness.

Source IDs: CAS-VAS-08

Bedtime (PM)

Do / Take

- Magnesium glycinate — 200-300mg [CAS-MICRO-03]

Continuation / maintenance: Carry: Yes | Maint: Daily | Stop/Taper: Long-term OK.

Source IDs: CAS-MICRO-03

Do / Take

- Glycine — 2-3g [CAS-AMINO-01]

Continuation / maintenance: Carry: Yes | Maint: Daily or PRN | Stop/Taper: Optional long-term; reduce if daytime sedation.

Source IDs: CAS-AMINO-01

Do / Take

- ALA — 100-200mg [CAS-MITO-07]

Continuation / maintenance: Carry: PRN | Maint: 2–4×/week or cycles | Stop/Taper: Optional/cycled; stop if insomnia/wired.

Source IDs: CAS-MITO-07

Do / Take

- Magnesium β -Hydroxybutyrate (MHB) — 500–1500mg — 3x/week [CAS-NEURO-13]

Optional

- Optional

Continuation / maintenance: Carry: PRN | Maint: 2–4×/week cycles | Stop/Taper: Cycled only; stop if oversedation/dizziness.

Source IDs: CAS-NEURO-13

Practices (today)

Do / Take

- Reverse Kegels — Daily — Any, especially pre/ post-sex [COP-01]

Source IDs: COP-01

Do / Take

- Psoas Release — 3–7x/week — Evening or off days [COP-02]

Source IDs: COP-02

Do / Take

- Adductor & Hamstring Lengthening — 3–5x/week — Post-walk or training [COP-03]

Source IDs: COP-03

Do / Take

- Pelvic Mobility Drills — Daily — AM or PM [COP-04]

Source IDs: COP-04

Do / Take

- Perineal PBM (red/IR light) — 3–7x/week — Evening [COP-05]

Source IDs: COP-05

Do / Take

- Nasal PBM — 3–7x/week — Evening or morning [COP-06]

Source IDs: COP-06

Do / Take

- Fascia & Pelvic Floor Neuro-Mechanics — 3–5x/week — Any [COP-07]

Source IDs: COP-07

Do / Take

- Sensate Focus Mapping — 2–4x/week — Private time [COP-08]

Source IDs: COP-08

Do / Take

- Feather-Touch Neuroerotic Training — 2–4x/week — Private time [COP-09]

Source IDs: COP-09

Do / Take

- Angion v2 Microvascular Training — 2–3x/week — Pre- or post-vascular stack [COP-10]

Source IDs: COP-10

Do / Take

- Sauna / Cold Contrast — 1–4x/week — Evening [COP-11]

Source IDs: COP-11

Do / Take

- Diaphragmatic Breathwork — Daily — Any, especially pre-sleep [COP-12]

Source IDs: COP-12

Do / Take

- Glute Activation Sequences — 3–5x/week — Pre-walk or training [COP-13]

Source IDs: COP-13

Do / Take

- Sleep Optimization Practices — Daily — Evening [COP-14]

Source IDs: COP-14

Do / Take

- Liver Pumping / Thoracic Mobilization — 3–7x/week — Evening [COP-15]

Source IDs: COP-15

Do / Take

- Vagal Reset Maneuvers — PRN + daily micro-practices — Any [COP-16]

Optional

- Optional

Source IDs: COP-16

Do / Take

- Foot–Core Activation Chain — 3–5x/week — Pre-walk/training [COP-17]

Source IDs: COP-17

Do / Take

- Infrared Sauna Protocol — 1–3x/week — Evening [COP-18]

Source IDs: COP-18

Do / Take

- Olfactory Training Protocol — Daily — Any [COP-19]

Source IDs: COP-19

Do / Take

- CO₂ Tolerance Breathwork — 3–5x/week — Any [COP-20]

Source IDs: COP-20

Do / Take

- Micro-Arousal Sensory Conditioning — 2–3x/week — Private time [COP-21]

Source IDs: COP-21

Do / Take

- Glans Sensitivity Reversal Protocol — 2–4x/week — Private time [COP-22]

Source IDs: COP-22

Do / Take

- Hydration Baseline — Daily — On waking + throughout day [COP-23]

Source IDs: COP-23

Watch-for / STOP-ADJUST

Do / Take

- Avoid choline, lecithin, ox bile, DIM, I3C, high-dose vitamin D, aggressive training.

Do / Take

- No daily Angion

Do / Take

- avoid overtraining

Do / Take

- cycle dopaminergics

Do / Take

- monitor for anxiety, insomnia or excessive sympathetic tone.

Do / Take

- Avoid starting at very high dose

Do / Take

- caution if severe herx or constipation.

Do / Take

- Avoid if severe diarrhea, strong bile reflux sensations, or new right-upper-quadrant pain.

Do / Take

- Avoid stacking new stimulants

Do / Take

- continue to avoid choline, ox bile and aggressive DIM/I3C usage.

Do / Take

- Caution in severe liver disease

Do / Take

- watch for unusual fatigue or paradoxical agitation.

Do / Take

- May provoke mild detox/headache in sensitive individuals

Do / Take

- start at lower frequency if needed.

Do / Take

- Avoid with serotonergic drugs (SSRIs, MAOIs)

Do / Take

- caution in G6PD deficiency

Do / Take

- may discolor urine/teeth.

Do / Take

- Avoid with sedatives, alcohol, or severe hypotension

Do / Take

- discontinue if dizziness or excessive somnolence.

Do / Take

- None significant

Do / Take

- avoid taking fasted.

Advance when

Do / Take

- Morning erections more consistent, libido clearly improved, workout recovery good, mood and drive significantly higher.

Phase 3.0

Intent: advanced optimization; stronger levers remain gated; keep RESET track separate unless deliberately entered.

On waking

Do / Take

- Hydration Baseline [COP-23]

Source IDs: COP-23

Do / Take

- Black Coffee — 1–3 cups/day [CAS-FOOD-03]

Optional

- Optional

Source IDs: CAS-FOOD-03

Binder window (fasted, mid-morning)

Do / Take

- Psyllium — 5-10g [CAS-GUT-01]

Continuation / maintenance: Carry: Yes | Maint: 2–4×/week or PRN | Stop/Taper: Taper to PRN once stools stable 10–14d; reduce first if constipation.

Source IDs: CAS-GUT-01

Binder window (after psyllium)

Do / Take

- Carrot salad [CAS-FOOD-01]

Continuation / maintenance: Carry: Yes | Maint: 3–5×/week or PRN | Stop/Taper: Maintenance once GI/skin stable 10–14d.

Source IDs: CAS-FOOD-01

Post-binder meal (first meal)

Do / Take

- Taurine — 1g [CAS-LIV-01]

Continuation / maintenance: Carry: Yes | Maint: Daily or 3–5×/week | Stop/Taper: Reduce frequency once bile/GI stable 10–14d.

Source IDs: CAS-LIV-01

Do / Take

- MSM — 750mg [CAS-LIV-02]

Continuation / maintenance: Carry: Yes | Maint: 3–5×/week → discontinue | Stop/Taper: Taper off once skin/gut stable 14d; stop if sulfur intolerance/wired.

Source IDs: CAS-LIV-02

Do / Take

- Molybdenum — 100-200mcg [CAS-LIV-03]

Continuation / maintenance: Carry: Yes | Maint: Daily with MSM; optional otherwise | Stop/Taper: Keep while MSM used; optional after MSM stopped.

Source IDs: CAS-LIV-03

Do / Take

- Zinc — 20-30mg [CAS-MICRO-01]

Continuation / maintenance: Carry: Course | Maint: 2–4×/week maintenance or stop | Stop/Taper: Reassess after 6–8 weeks; avoid long-term high-dose without monitoring.

Source IDs: CAS-MICRO-01

Do / Take

- Selenium — 100-200mcg [CAS-MICRO-02]

Continuation / maintenance: Carry: Course | Maint: 2–4×/week maintenance or stop | Stop/Taper: Reassess after 8–12 weeks.

Source IDs: CAS-MICRO-02

Do / Take

- Benfotiamine — 150-300mg [CAS-MITO-05]

Continuation / maintenance: Carry: Yes | Maint: 3–5×/week maintenance | Stop/Taper: Taper once energy stable 14d.

Source IDs: CAS-MITO-05

Do / Take

- Calcium D-Glucarate — 500mg [CAS-GUT-02]

Continuation / maintenance: Carry: PRN | Maint: 2–4×/week or PRN | Stop/Taper: Use during mobilization; taper when stable 14d.

Source IDs: CAS-GUT-02

Do / Take

- PHGG — 1-2g [CAS-GUT-03]

Continuation / maintenance: Carry: Yes | Maint: 3–5×/week maintenance | Stop/Taper: Taper once stools stable 10–14d.

Source IDs: CAS-GUT-03

Do / Take

- MegaSporeBiotic — 0.5→1 cap [CAS-GUT-04]

Continuation / maintenance: Carry: Course | Maint: Stop after course | Stop/Taper: Course 8–12 weeks then stop and reassess.

Source IDs: CAS-GUT-04

Do / Take

- Glutathione (oral, generic) — 250–500mg — Daily (optional) [CAS-DET-01]

Optional

- Optional

Continuation / maintenance: Carry: PRN | Maint: 2–4×/week or short blocks | Stop/Taper: Optional: use in short blocks if tolerated.

Source IDs: CAS-DET-01

Do / Take

- NAC — 600mg [CAS-BIO-02]

Optional

- Optional

Continuation / maintenance: Carry: PRN | Maint: 2–4×/week or short blocks | Stop/Taper: Optional: use in short blocks; stop if wired/insomnia.

Source IDs: CAS-BIO-02

Do / Take

- Sulforaphane (Broccoli Sprout Extract) — per label (e.g. 20–40mg) — 3–5×/week [CAS-INF-02]

Continuation / maintenance: Carry: PRN | Maint: 3–5×/week cycles | Stop/Taper: Optional/cycled; reduce frequency if headache/insomnia.

Source IDs: CAS-INF-02

Do / Take

- Methylene Blue (low-dose) — 0.5–1mg/kg — 3–5×/week [CAS-MITO-15]

Continuation / maintenance: Carry: PRN | Maint: 3–5×/week cycles (never continuous) | Stop/Taper: Cycled only; stop if insomnia/wired.

Source IDs: CAS-MITO-15

Do / Take

- SAM-e — 200mg [CAS-MET-01]

Continuation / maintenance: Carry: PRN | Maint: 3–5×/week or PRN | Stop/Taper: Optional; stop if agitation/insomnia.

Source IDs: CAS-MET-01

Do / Take

- 7,8-Dihydroxyflavone (7,8-DHF) — 10–40mg — Daily (10 days on / 10 days off) – optional [CAS-NEURO-05]

Optional

- Optional

Continuation / maintenance: Carry: PRN | Maint: 10 on / 10 off | Stop/Taper: Optional/cycled; stop if insomnia/anxiety.

Source IDs: CAS-NEURO-05

Do / Take

- TTFD (Thiamine Tetrahydrofurfuryl Disulfide) — 50–200mg [CAS-MITO-06]

Continuation / maintenance: Carry: PRN | Maint: PRN (training/drive days) | Stop/Taper: Use PRN; stop if insomnia/wired.

Source IDs: CAS-MITO-06

Do / Take

- Boron — 6–9mg [CAS-MIN-01]

Continuation / maintenance: Carry: Yes | Maint: 2 weeks on / 1 off | Stop/Taper: As written: cycle.

Source IDs: CAS-MIN-01

Do / Take

- Uridine Monophosphate — 50–150mg [CAS-NEURO-08]

Continuation / maintenance: Carry: PRN | Maint: Cycles only | Stop/Taper: Optional/cycled.

Source IDs: CAS-NEURO-08

Do / Take

- Piracetam — 1200–2400mg [CAS-NOOT-01]

Continuation / maintenance: Carry: PRN | Maint: Cycles only | Stop/Taper: Optional/cycled.

Source IDs: CAS-NOOT-01

Do / Take

- Sulbutiamine — 200–400mg — 5 days on / 2 off [CAS-NEURO-09]

Continuation / maintenance: Carry: PRN | Maint: 5 on / 2 off | Stop/Taper: Cycled only (as written).

Source IDs: CAS-NEURO-09

Do / Take

- Methylated B-Complex — Label-equivalent [CAS-VIT-05]

Continuation / maintenance: Carry: Yes | Maint: 3–5×/week or daily | Stop/Taper: Optional long-term; reduce if overstimulation.

Source IDs: CAS-VIT-05

Do / Take

- Semax — 0.1–0.3mg IN — 5 days on / 2 off [CAS-PEPT-02]

Continuation / maintenance: Carry: PRN | Maint: 5 on / 2 off | Stop/Taper: Cycled only; stop if agitation/insomnia.

Source IDs: CAS-PEPT-02

Do / Take

- Mucuna Pruriens — 100–200mg — 10 on / 20 off [CAS-NEURO-10]

Continuation / maintenance: Carry: PRN | Maint: PRN (evening) | Stop/Taper: PRN for aldehyde load; stop if daytime sedation.

Source IDs: CAS-NEURO-10

Do / Take

- Tongkat Ali — 400-600mg [CAS-ANDRO-01]

Continuation / maintenance: Carry: Yes | Maint: 5 on / 2 off or blocks | Stop/Taper: Cycle to avoid tolerance; stop if irritability/insomnia.

Source IDs: CAS-ANDRO-01

Do / Take

- Cistanche — 1000-2000mg [CAS-ANDRO-02]

Continuation / maintenance: Carry: Yes | Maint: 5 on / 2 off or blocks | Stop/Taper: Cycle to avoid tolerance; stop if irritability/insomnia.

Source IDs: CAS-ANDRO-02

Do / Take

- Proviron — 12.5mg [CAS-ANDRO-10]

Continuation / maintenance: Carry: No | Maint: No (trial only) | Stop/Taper: Late-game trial only; requires monitoring; stop if adverse androgenic/prostate effects.

Source IDs: CAS-ANDRO-10

Do / Take

- Pregnenolone — 10-20mg [CAS-STERO-01]

Continuation / maintenance: Carry: Yes | Maint: Daily or 5 on / 2 off | Stop/Taper: Reassess with symptoms/labs; stop if insomnia/anxiety.

Source IDs: CAS-STERO-01

Do / Take

- DHEA — 5-15mg [CAS-STERO-02]

Continuation / maintenance: Carry: Yes | Maint: Daily or 5 on / 2 off | Stop/Taper: Reassess with symptoms/labs; stop if acne/irritability/insomnia.

Source IDs: CAS-STERO-02

Do / Take

- HMB — 1.5-3g [CAS-MITO-03]

Continuation / maintenance: Carry: PRN | Maint: Training/sex days or blocks | Stop/Taper: Choose one HMB lane; use with food; stop if GI upset.

Source IDs: CAS-MITO-03

Do / Take

- BPC-157 — 250-500mcg — Cycle [CAS-PEPT-01]

Continuation / maintenance: Carry: PRN | Maint: Short cycles | Stop/Taper: Cycled only (injury/recovery blocks).

Source IDs: CAS-PEPT-01

Do / Take

- Niacin (flush) — 50-100mg [CAS-MET-07]

Continuation / maintenance: Carry: PRN | Maint: PRN or low-frequency | Stop/Taper: Flush niacin: optional; stop if hypotension/flush intolerance.

Source IDs: CAS-MET-07

Do / Take

- 4'-DMA-7,8-DHF — Experimental microdose — Short microcycle (5–10 days max, optional) [CAS-NEURO-06]

Optional

- Optional

Continuation / maintenance: Carry: No | Maint: No (microcycle) | Stop/Taper: Experimental microcycle only (max 5–10 days).

Source IDs: CAS-NEURO-06

Do / Take

- Phenylpiracetam — 50–100mg — 2–4×/week [CAS-NOOT-02]

Continuation / maintenance: Carry: PRN | Maint: 2–4×/week cycles | Stop/Taper: 2–4×/week only; stop if insomnia/anxiety.

Source IDs: CAS-NOOT-02

Do / Take

- HMB (β -Hydroxy β -Methylbutyrate) — 2–3 g/day — Daily or pre-performance [CAS-NEURO-14]

Continuation / maintenance: Carry: PRN | Maint: Training/sex days or blocks | Stop/Taper: Use with food; choose one HMB lane; stop if GI upset.

Source IDs: CAS-NEURO-14

With meals (split across meals)

Do / Take

- Undecylenic Acid — titrate 1→5 caps/day [CAS-FUNG-01]

Continuation / maintenance: Carry: Course | Maint: Course; PRN pulses | Stop/Taper: Finite course: taper off after fungal/SD stable 14d; PRN pulses for relapse.

Source IDs: CAS-FUNG-01

Do / Take

- Niacinamide — 250–500mg [CAS-B-01]

Continuation / maintenance: Carry: Yes | Maint: Daily with food or 3–5×/week | Stop/Taper: Reduce if over-sedation or GI effects.

Source IDs: CAS-B-01

Do / Take

- Endotelon (grape seed OPC) — 150–300mg/day (often 150mg BID) [CAS-VAS-15]

Continuation / maintenance: Carry: Course | Maint: Block 8–12 w | Stop/Taper: Block 8–12 weeks then reassess; avoid stacking with aspirin/polyphenols if bruising.

Source IDs: CAS-VAS-15

Do / Take

- Tributyrin — 300–600mg [CAS-GUT-07]

Continuation / maintenance: Carry: Yes | Maint: 3–5×/week | Stop/Taper: Taper once stools stable 14d.

Source IDs: CAS-GUT-07

Do / Take

- *Lactobacillus reuteri* — 1–2B CFU [CAS-PRO-02]

Continuation / maintenance: Carry: Course | Maint: Stop after course or cycle | Stop/Taper: Course 8–12 weeks then reassess.

Source IDs: CAS-PRO-02

Lunch (with food)

Do / Take

- Creatine — 5g [CAS-MITO-01]

Continuation / maintenance: Carry: Yes | Maint: Daily | Stop/Taper: Long-term OK.

Source IDs: CAS-MITO-01

Do / Take

- Aspirin — 81mg — EOD [CAS-INF-01]

Continuation / maintenance: Carry: Course | Maint: EOD course then reassess | Stop/Taper: Course; taper off once inflammation stable; stop if bruising/GI issues.

Source IDs: CAS-INF-01

Do / Take

- Vitamin K2 — MK7 100-200mcg [CAS-MICRO-04]

Continuation / maintenance: Carry: Yes | Maint: Daily with food | Stop/Taper: Long-term OK.

Source IDs: CAS-MICRO-04

Do / Take

- Omega-3 — 500mg [CAS-LIP-01]

Continuation / maintenance: Carry: Yes | Maint: Daily with food | Stop/Taper: Long-term OK.

Source IDs: CAS-LIP-01

Do / Take

- Reduced Glutathione (liposomal) — 250–500mg — Daily (optional) [CAS-DET-02]

Optional

- Optional

Continuation / maintenance: Carry: PRN | Maint: 2–4×/week or blocks | Stop/Taper: Optional mitochondrial/antioxidant support; use in short blocks if tolerated.

Source IDs: CAS-DET-02

Do / Take

- LCLT — 2g [CAS-MITO-02]

Continuation / maintenance: Carry: Yes | Maint: Daily with food | Stop/Taper: Long-term OK; stop if fishy odor or GI intolerance.

Source IDs: CAS-MITO-02

Dinner (with food)

Do / Take

- TUDCA — 250mg — Daily (from Week 2) [CAS-LIV-04]

Continuation / maintenance: Carry: Yes | Maint: 2–4×/week maintenance | Stop/Taper: Taper once motility + bile symptoms stable 10–14d; hold if bile reflux/RUQ pain.

Source IDs: CAS-LIV-04

Pre-workout / pre-sex (acute signaling)

Do / Take

- Citrulline — 6g [CAS-VAS-01]

Continuation / maintenance: Carry: PRN | Maint: Training/sex days only | Stop/Taper: Acute tool for training/sex days.

Source IDs: CAS-VAS-01

Do / Take

- Icaritin — 150-200mg [CAS-VAS-02]

Continuation / maintenance: Carry: PRN | Maint: Training/sex days only | Stop/Taper: Acute tool for training/sex days.

Source IDs: CAS-VAS-02

Do / Take

- Histidine — 500-1000mg [CAS-VAS-03]

Continuation / maintenance: Carry: PRN | Maint: PRN | Stop/Taper: Optional acute adjunct; stop if histamine symptoms.

Source IDs: CAS-VAS-03

Do / Take

- Hesperidin — 250mg [CAS-VAS-04]

Continuation / maintenance: Carry: PRN | Maint: PRN | Stop/Taper: Optional adjunct; low priority if low BP.

Source IDs: CAS-VAS-04

Do / Take

- Pycnogenol — 50-100mg [CAS-VAS-05]

Continuation / maintenance: Carry: PRN | Maint: PRN or blocks | Stop/Taper: Prefer as block (8–12 w) if used chronically; watch bleeding with aspirin.

Source IDs: CAS-VAS-05

Do / Take

- Vitamin C — 500-1000mg [CAS-ANTIOX-01]

Continuation / maintenance: Carry: Yes | Maint: Daily or PRN with vascular block | Stop/Taper: Optional long-term; reduce if GI irritation.

Source IDs: CAS-ANTIOX-01

Do / Take

- Beet nitrate — 300-600mg [CAS-VAS-08]

Continuation / maintenance: Carry: PRN | Maint: PRN (training/sex days) | Stop/Taper: Use PRN; stop if low BP/dizziness.

Source IDs: CAS-VAS-08

Do / Take

- Agmatine — 500-1000mg [CAS-VAS-07]

Continuation / maintenance: Carry: PRN | Maint: Training/sex days only | Stop/Taper: Acute adjunct; stop if low BP.

Source IDs: CAS-VAS-07

Do / Take

- Berberine — 150-300mg [CAS-MET-03]

Continuation / maintenance: Carry: PRN | Maint: PRN or short blocks | Stop/Taper: Use with meals or training days; stop if GI upset/hypoglycemia.

Source IDs: CAS-MET-03

Do / Take

- L-Arginine — 1-3g [CAS-VAS-12]

Optional

- Optional

Continuation / maintenance: Carry: PRN | Maint: Training/sex days only | Stop/Taper: Acute tool; reduce first if low BP/headache.

Source IDs: CAS-VAS-12

Bedtime (PM)

Do / Take

- Magnesium glycinate — 200-300mg [CAS-MICRO-03]

Continuation / maintenance: Carry: Yes | Maint: Daily | Stop/Taper: Long-term OK.

Source IDs: CAS-MICRO-03

Do / Take

- Glycine — 2-3g [CAS-AMINO-01]

Continuation / maintenance: Carry: Yes | Maint: Daily or PRN | Stop/Taper: Optional long-term; reduce if daytime sedation.

Source IDs: CAS-AMINO-01

Do / Take

- ALA — 100-200mg [CAS-MITO-07]

Continuation / maintenance: Carry: PRN | Maint: 2–4×/week or cycles | Stop/Taper: Optional/cycled; stop if insomnia/wired.

Source IDs: CAS-MITO-07

Do / Take

- Nattokinase — 100-200mg [CAS-VAS-11]

Continuation / maintenance: Carry: No | Maint: No (trial only) | Stop/Taper: Avoid stacking with aspirin; trial only; stop if bruising/bleeding.

Source IDs: CAS-VAS-11

Do / Take

- Retinol — 5000IU — 2–3×/week [CAS-VIT-01]

Continuation / maintenance: Carry: PRN | Maint: 2–3×/week | Stop/Taper: 2–3×/week only; stop if headaches/dryness.

Source IDs: CAS-VIT-01

Do / Take

- Magnesium β-Hydroxybutyrate (MHB) — 500–1500mg — 2–4x/week [CAS-NEURO-13]

Continuation / maintenance: Carry: PRN | Maint: 2–4×/week cycles | Stop/Taper: Cycled only; stop if oversedation/dizziness.

Source IDs: CAS-NEURO-13

Practices (today)

Do / Take

- Reverse Kegels — Daily — Any, especially pre/ post-sex [COP-01]

Source IDs: COP-01

Do / Take

- Psoas Release — 3–7x/week — Evening or off days [COP-02]

Source IDs: COP-02

Do / Take

- Adductor & Hamstring Lengthening — 3–5x/week — Post-walk or training [COP-03]

Source IDs: COP-03

Do / Take

- Pelvic Mobility Drills — Daily — AM or PM [COP-04]

Source IDs: COP-04

Do / Take

- Perineal PBM (red/IR light) — 3–7x/week — Evening [COP-05]

Source IDs: COP-05

Do / Take

- Nasal PBM — 3–7x/week — Evening or morning [COP-06]

Source IDs: COP-06

Do / Take

- Fascia & Pelvic Floor Neuro-Mechanics — 3–5x/week — Any [COP-07]

Source IDs: COP-07

Do / Take

- Sensate Focus Mapping — 2–4x/week — Private time [COP-08]

Source IDs: COP-08

Do / Take

- Feather-Touch Neuroerotic Training — 2–4x/week — Private time [COP-09]

Source IDs: COP-09

Do / Take

- Angion v2 Microvascular Training — 2–3x/week — Pre- or post-vascular stack [COP-10]

Source IDs: COP-10

Do / Take

- Sauna / Cold Contrast — 1–4x/week — Evening [COP-11]

Source IDs: COP-11

Do / Take

- Diaphragmatic Breathwork — Daily — Any, especially pre-sleep [COP-12]

Source IDs: COP-12

Do / Take

- Glute Activation Sequences — 3–5x/week — Pre-walk or training [COP-13]

Source IDs: COP-13

Do / Take

- Sleep Optimization Practices — Daily — Evening [COP-14]

Source IDs: COP-14

Do / Take

- Liver Pumping / Thoracic Mobilization — 3–7x/week — Evening [COP-15]

Source IDs: COP-15

Do / Take

- Vagal Reset Maneuvers — PRN + daily micro-practices — Any [COP-16]

Optional

- Optional

Source IDs: COP-16

Do / Take

- Foot–Core Activation Chain — 3–5x/week — Pre-walk/training [COP-17]

Source IDs: COP-17

Do / Take

- Infrared Sauna Protocol — 1–3x/week — Evening [COP-18]

Source IDs: COP-18

Do / Take

- Olfactory Training Protocol — Daily — Any [COP-19]

Source IDs: COP-19

Do / Take

- CO₂ Tolerance Breathwork — 3–5x/week — Any [COP-20]

Source IDs: COP-20

Do / Take

- Micro-Arousal Sensory Conditioning — 2–3x/week — Private time [COP-21]

Source IDs: COP-21

Do / Take

- Glans Sensitivity Reversal Protocol — 2–4x/week — Private time [COP-22]

Source IDs: COP-22

Do / Take

- Hydration Baseline — Daily — On waking + throughout day [COP-23]

Source IDs: COP-23

Watch-for / STOP-ADJUST

Do / Take

- Avoid choline, lecithin, ox bile, DIM, I3C, high-dose vitamin D, aggressive training.

Do / Take

- No daily Angion

Do / Take

- avoid overtraining

Do / Take

- cycle dopaminergics

Do / Take

- monitor for anxiety, insomnia or excessive sympathetic tone.

Do / Take

- Avoid starting at very high dose

Do / Take

- caution if severe herx or constipation.

Do / Take

- Avoid if severe diarrhea, strong bile reflux sensations, or new right-upper-quadrant pain.

Do / Take

- Avoid stacking new stimulants

Do / Take

- continue to avoid choline, ox bile and aggressive DIM/I3C usage.

Do / Take

- Caution in severe liver disease

Do / Take

- watch for unusual fatigue or paradoxical agitation.

Do / Take

- May provoke mild detox/headache in sensitive individuals

Do / Take

- start at lower frequency if needed.

Do / Take

- Avoid with serotonergic drugs (SSRIs, MAOIs)

Do / Take

- caution in G6PD deficiency

Do / Take

- may discolor urine/teeth.

Do / Take

- Avoid stacking too many stimulants or vasodilators

Do / Take

- cap Angion at 3×/week

Do / Take

- ensure deload weeks.

Do / Take

- Avoid with sedatives, alcohol, or severe hypotension

Do / Take

- discontinue if dizziness or excessive somnolence.

Do / Take

- None significant

Do / Take

- avoid taking fasted.

Advance when

Do / Take

- Sustained high libido and EQ, robust energy, low crash frequency, stable mood and sleep even under stress.

RESET TRACK (Optional) — Separate from Mainline

Higher-volatility option. Do not blend into mainline. Maintain Aurelius stabilizers and use single-variable changes.

Track Component	Related CAS ID(s)	Suggested Range (Reset)	Purpose	Prerequisites / Gates
High-dose NMN (NAD+ push)	CAS-MITO-08	≥1000 mg/day (experimental reset)	Epi-genetic / NAD+ drive; may increase Phase 1 tolerance	NOT Phase 1 use only (after stable baseline)
TMG (methyl-buffer)	CAS-MET-08	500–2000 mg/day with meals	Buffers methyl demand; supports histone acetylation; daily NMN push, tolerable	Start once baseline acylation; daily NMN push, tolerable
Hops extract (8■PN) — RESET ONLY	CAS-RESET-01	Standardized 8■PN; clinician-grade	Reduce estrogenic lever in reset	Avoid if prior estrogenization via high prep
Keep Aurelius stabilizers (do not change mainline CAS list)	As per Phase + baseline	As per Phase + baseline	Maintain terrain stability while testing	Do not reset multiple new agents at once; one c
Governance: RESET track is OPTIONAL and higher-volatility. Keep it separate from Phase 1 / mainline. Prefer: baseline labs → single-variable change → track → decision				