

**Treatment Success and Failure Prediction Using Machine Learning Models
for Pulmonary Tuberculosis Patients Under the DOH-CAR TB-DOTS Program**

Aaron Kyle M. Aguilar, Mark Lestat C. Agustin, Keanu Sonn M. Fortaleza,
Benny Gil A. Lactaotao, Marven Joffre G. Luis, Georcelle Leisley U. Nuarin,
Sean Drei A. Octavo, and Gebreyil Isaac P. Rabang

Department of Computing and Information Studies, Saint Louis University

Abstract

Pulmonary Tuberculosis (PTB) remains a critical public health challenge in the Cordillera Administrative Region (CAR), Philippines, where complex patient profiles and programmatic gaps contribute to persistent treatment failure. This study develops machine learning models to predict treatment success or failure among PTB patients in DOH-CAR TB-DOTS facilities covering the period 2015–2025. A standardized preprocessing pipeline performs schema harmonization, patient pseudonymization, and missing data imputation via Multivariate Imputation by Chained Equations (MICE). Supervised models—Logistic Regression, Support Vector Machines (SVM), and Extreme Gradient Boosting (XGBoost)—are trained alongside Recurrent Neural Network (RNN) and Long Short-Term Memory (LSTM) temporal architectures to capture longitudinal treatment dynamics across the 6- to 12-month treatment course. Shapley Additive Explanations (SHAP) ensure model interpretability by quantifying the contribution of individual patient features to each prediction. The validated models are embedded in a Clinical Decision Support System (CDSS) that provides dynamic risk estimates and factor-level explanations to support early clinical intervention for patients at risk of treatment failure.

Keywords: tuberculosis, machine learning, treatment outcome prediction, clinical decision support system, explainable AI

Treatment Success and Failure Prediction Using Machine Learning Models for Pulmonary Tuberculosis Patients Under the DOH-CAR TB-DOTS Program

Pulmonary Tuberculosis (PTB) is a chronic infectious disease caused by *Mycobacterium tuberculosis* (MTB), a slow-growing and acid-fast bacillus that continues to challenge public health due to its gradual progression and risk of treatment failure (?). Despite effective therapies, PTB remains widespread due to its strong association with poverty, undernutrition, and comorbidities such as diabetes and HIV which weaken immune defenses and increase susceptibility (?). These factors delay diagnosis, complicate treatment, and contribute to disease progression. Globally, TB is still one of the leading causes of death, with 1.5 million deaths and 10 million new cases recorded in 2018, including half a million with rifampicin resistance.

In the Philippines, TB continues to be a major public health concern. The World Health Organization (WHO) identifies the country as one of the top contributors to the global TB burden as it accounts for an estimated 6.8 percent of cases. Before the COVID-19 pandemic, the Philippines had one of the highest TB rates in Asia with over 554 cases per 100,000 people in 2019 (?). In 2023, around 37,000 Filipinos died from TB out of the 739,000 diagnosed cases. The WHO classifies the country as a high burden for MDR-TB and TB/HIV co-infection (?). That same year, the DOH recorded 575,770 TB cases through its Integrated TB Information System which indicates that many cases remain unregistered or undiagnosed (?). The DOH implements CDC-recommended regimens for active and latent TB through the TB-DOTS strategy which was expanded in 2003 to include private clinics, pharmacies, and hospitals (??). Despite these initiatives, gaps in diagnosis, treatment monitoring, and patient support persist especially in the management of MDR-TB (?).

In Baguio City, the DOH-CAR reported 437 confirmed TB cases in 2022 along with over 4,405 missing or unreported cases making the area a priority for surveillance and treatment improvement (?). Contributing factors include high population density, internal migration, climate, and the city's role as a referral center for nearby provinces. Patient comorbidities, socioeconomic barriers, and health system limitations also influence treatment outcomes (?). These challenges highlight the need for locally adaptive strategies as treatment failure not only worsens patient

outcomes but also prolongs transmission and drives the emergence of MDR-TB (?).

Risk Factors for Treatment Failure

To ensure uniformity in the evaluation of TB control efforts, the DOH has established standardized criteria for classifying treatment outcomes through the National Tuberculosis Program (NTP). Establishing these outcomes is crucial for monitoring program performance, ensuring comparability across studies, and informing risk factor analysis. Ultimately, these outcomes of success or failure are not random but are linked to a distinct set of patient-specific, clinical, and programmatic risk factors. While mortality remains the primary outcome for most studies on treatment failure, some also consider the development of disease, resistance to multiple anti-TB drugs, and failure to follow up (?).

One of the risk factors is that TB patients have close contact with other patients with active TB infection. A study noted that TB patients who acquired the infection through close contact were shown to be at higher risk of delayed diagnosis and more likely to be non-adherent with the treatment either by refusing or discontinuing the regimen (?). Another risk factor is recurrence of the infection or having a history of receiving TB treatment. TB patients who missed follow-ups had a higher risk of reactivation of the infection and mortality as they are more difficult to treat compared to patients with no prior history of TB (?). Furthermore, missed or interrupted treatment causes the development of resistance to the drug if done frequently (?).

Comorbidities are also discussed in studies that have consistently shown a significant influence on tuberculosis treatment outcomes across different settings. In Eastern Ethiopia, HIV co-infection was determined to be a substantial risk factor for treatment failure due to the weakened immune system which leads to a higher risk for ineffective treatment (?). In addition, TB and HIV drug interactions accelerate each other's progression and increase the occurrence of side effects or adverse effects which may compromise treatment success (?). Likewise, a study in Davao City, Philippines, revealed that there was a higher risk of treatment failure in comorbid patients with

58.82% of the 51 patients who experienced treatment failure having at least one comorbidity (?). Another major determinant is Diabetes Mellitus (DM). TB patients with diagnosed DM have a higher risk of treatment failure, relapse, and death compared to non-diabetics (?). In a systematic review and meta-analysis, patients with DM have a higher risk of experiencing treatment failure for Drug-resistant (DR) and mortality for MDR-TB with an odds ratio of 1.56 and 1.57 respectively (?). In line with this, another study states that patients suffering from both PTB and chronic conditions, particularly DM, had reduced rates of sputum conversion with a higher probability of poor treatment outcomes including death and progression to MDR-TB (?).

Machine Learning for TB Outcome Prediction

Although multiple studies have examined risk factors such as sociodemographic characteristics, comorbidities, and medical history, the findings remain inconsistent and fragmented since methodological constraints limit them. Most studies rely on traditional statistical analyses which are statistical tests or regressions that examine one factor at a time or a few factors together while assuming their linear relationships (?). These limitations highlight the need for an integrative approach to combine multiple dimensions of data in analyzing the interrelated variables together as a critical gap in capturing the multifaceted nature of TB treatment outcomes.

To address this gap, the present study adopts a holistic multi-layer analytical framework that integrates descriptive analysis and supervised predictive modeling using demographic, clinical, diagnostic, and treatment-related variables. Descriptive analyses are employed to characterize patient profiles and treatment patterns, while supervised learning models are used to predict treatment outcomes. This integrative approach enables machine learning models to learn complex relationships among multiple patient-level variables that influence treatment success or failure, thereby enhancing both predictive accuracy and interpretability (??).

Machine learning (ML) offers a distinct advantage in modeling complex and non-linear interactions among demographic, clinical, diagnostic, and treatment-related variables that con-

ventional statistical analyses may overlook. By leveraging algorithms capable of learning from high-dimensional data, such as logistic regression, Support Vector Machines (SVM), and ensemble-based methods including Extreme Gradient Boosting (XGBoost), the study aims to uncover subtle dependencies that contribute to treatment success or failure (?). Previous research has demonstrated the utility of ML in TB research, including predicting drug resistance, identifying high-risk patients, and optimizing treatment strategies (??). In this study, the models are trained directly on integrated patient data, allowing them to capture individual-level patterns that improve predictive performance while maintaining interpretability.

Beyond prediction, interpretability will be enhanced through the application of Shapley Additive Explanations (SHAP) which is a post-hoc interpretability technique that quantifies the contribution of each variable to the model's predictions. SHAP enables a transparent evaluation of which patient features most strongly drive treatment success or failure. This supports explainable and trustworthy Artificial Intelligence (AI) in clinical contexts (?). Using supervised learning with interpretable AI methods, this framework not only forecasts treatment outcomes but also reveals the relative importance of risk factors such as drug resistance, comorbidities, or regimen adherence. The resulting models are designed to help healthcare providers identify patients at a higher risk of treatment failure thus enabling early intervention, tailored regimens, and improved patient management.

Contributions

The main contributions of this paper are as follows:

1. **Integrated Predictive Framework.** The team develop and validate machine learning models that accurately predict therapy success or failure among TB patients in the CAR, Philippines. The integration of demographic, clinical, and laboratory variables focuses on supporting early identification of patients at risk of unsuccessful treatment and providing evidence-based tools for improving TB program outcomes.

2. **Data Preprocessing and Harmonization.** The team pre-process and document patient records of PTB cases from TB-DOTS facilities in the CAR through data cleaning, feature engineering, and reporting for readiness to be used in ML applications.
3. **Descriptive and Exploratory Analysis.** The team analyzed the demographic, clinical, diagnostic, and treatment characteristics of PTB patients to provide a comprehensive overview of the study population and identify key variables associated with treatment outcomes over the course of therapy.
4. **Temporal Modeling.** The team incorporates temporal modeling that analyzes time-dependent patterns in patient data allowing the detection of dynamic changes and trends throughout treatment. This thereby improves the accuracy of outcome predictions and supports timely clinical interventions.
5. **Clinical Decision Support.** The team developed a decision support tool that visualizes patient risk levels and predictive outputs. This enables early identification of individuals at risk of poor outcomes and ensures that the model maintains high predictive accuracy and reliability when applied to evaluate incoming TB cases.

Research Objectives

This study aims to develop and validate machine learning models that accurately predict therapy success or failure among TB patients in the CAR, Philippines. The integration of demographic, clinical, and laboratory variables focuses on supporting early identification of patients at risk of unsuccessful treatment and providing evidence-based tools for improving TB program outcomes, with the overarching goal of enabling personalized clinical interventions and efficient allocation of healthcare resources. Specifically, the research aims to achieve the following objectives:

1. To pre-process and document patient records of PTB cases from TB-DOTS facilities in the CAR through data cleaning, feature engineering, and reporting for readiness to be used in ML applications.

2. To describe the demographic, clinical, diagnostic, and treatment characteristics of PTB patients in the dataset, providing an overview of the study population.
3. To identify contributing variables that influence treatment outcomes over the course of therapy, enabling recognition of the most critical factors that affect patient recovery and guiding more precise treatment adjustments that can be intervened with.
4. To incorporate temporal modeling that analyzes time-dependent patterns in patient data, allowing the detection of dynamic changes and trends throughout treatment, thereby improving the accuracy of outcome predictions and supporting timely clinical interventions.
5. To train and validate supervised learning models for predicting treatment success or failure among PTB patients, enabling early identification of individuals at risk of poor outcomes and ensuring that the model maintains high predictive accuracy and reliability when applied to evaluate incoming TB cases.
6. To develop a decision support tool that visualizes patient risk levels and predictive outputs.

Scope and Limitations

The dataset will include PTB patients for the period 2015–2025 in the CAR excluding those who are still undergoing treatment at the time of data extraction. Furthermore, only patients with a well-defined treatment outcome of either success or failure will be included. The data gathered will encompass demographic information, laboratory tests, diagnostic data, clinical data, treatment regimen, follow-up examinations, drug administration data, and treatment outcomes. The study faces limitations related to the sufficiency and quality of the dataset. The data sourced from existing records may present challenges related to completeness, consistency, and representativeness due to limited access to complete clinical parameters. This could affect the inclusion of important risk factors and weaken the robustness of the analysis. Additionally, the study may not fully represent the current TB patient population as patients currently undergoing treatment are excluded. Difficulties are also expected during data extraction from DOH-CAR as variations in record-keeping over the

10-year period from 2015 to 2025 may be inconsistent due to changes in the healthcare system and shifts in documentation standards. Reliance on self-reported indicators may also introduce bias and inconsistencies that could affect the predictive capabilities of the model.

Significance of the Study

The resulting models are designed to help healthcare providers identify patients at a higher risk of treatment failure thus enabling early intervention, tailored regimens, and improved patient management. Ultimately, this machine learning framework supports the broader objective of strengthening TB control efforts in the Cordillera Administrative Region (CAR) by transforming existing patient data into predictive, explainable, and actionable tools for evidence-based public health practice. The results will help improve patient management and treatment planning, strengthen program implementation, and guide public health decision-making to improve the TB treatment success rate in CAR. Furthermore, the study aims to maximize benefits for the public by informing health administrators about the model which may be further developed into a solution or application to aid the local TB-DOTS program. The processed and pseudonymized derivative of the DOH-CAR Pulmonary Tuberculosis Treatment Outcome Dataset produced by the study will also be published to support the advancement of clinical, epidemiological, and machine learning research.


Method

This study uses a dataset of records of pulmonary tuberculosis (PTB) patients from the Cordillera Administrative Region (CAR), Philippines, covering 2015–2025 and consolidated by the Department of Health–CAR (DOH-CAR). After ethical approval, records are pseudonymized and processed using a standardized preprocessing pipeline, including schema harmonization, error correction, feature scaling, and missing value imputation via Multivariate Imputation by Chained

Equations (MICE). Exploratory and cluster analyses are conducted to characterize the cohort and derive subgroup-level features. Supervised models, Logistic Regression, Support Vector Machines (SVM), and Extreme Gradient Boosting (XGBoost), are trained to predict treatment outcomes defined by National Tuberculosis Program criteria. To capture longitudinal treatment dynamics, temporal models using Recurrent Neural Networks (RNN) and Long Short-Term Memory (LSTM) are evaluated alongside static models, with ensemble strategies applied for improved robustness. Model performance is assessed using standard classification metrics, and Shapley Additive Explanations (SHAP) are employed to ensure interpretability. The final models are integrated into a clinical decision support framework to provide dynamic risk estimates throughout the treatment course. An overview can be found in Figure ??.

Figure 1

Data Management and Analysis Process



Data Acquisition and Preprocessing

The study utilizes a retrospective dataset of Pulmonary Tuberculosis (PTB) patients from the Cordillera Administrative Region (CAR), Philippines, covering the period from 2015 to 2025. Data is centrally consolidated by the Department of Health-CAR (DOH-CAR), which manages reports from government TB-DOTS facilities and private practitioners under the Public-Private Mix DOTS (PPMD) scheme (?).

The raw dataset comprises demographic variables (e.g., age, sex, civil status), clinical indicators (e.g., BMI, blood pressure), and diagnostic results (e.g., GeneXpert, Smear Microscopy). To ensure data quality for machine learning applications, a reproducible preprocessing pipeline is implemented. This includes schema harmonization to standardize variable formats and units, and the removal of direct identifiers to strictly enforce patient privacy. Missing data (a common challenge in clinical registries) is handled using Multivariate Imputation by Chained Equations (MICE; ?).

Exploratory Data Analysis

Prior to predictive modeling, an exploratory analysis is conducted to characterize the study population and identify latent subgroups. The study employs cluster analysis on demographic, clinical, and diagnostic dimensions to detect natural groupings of patients. These clusters reveal distinguishing features such as specific comorbidity profiles, adherence behaviors, and regimen responses. Importantly, the resulting cluster assignments are not merely descriptive; they are integrated as engineered features into the subsequent supervised learning models. This approach enables the algorithms to learn both individual-level attributes and subgroup-level patterns that significantly influence treatment success or failure.

Identification of Variables Associated With Treatment Outcomes

To bridge the gap between predictive performance and clinical interpretability, the study identifies critical variables associated with treatment outcomes. While machine learning models are adept at handling high-dimensional data, understanding the “why” behind a prediction is crucial in medical contexts (?).

The framework utilizes Shapley Additive Explanations (SHAP), a post-hoc interpretability technique that quantifies the marginal contribution of each variable to the model’s output. This allows for the precise identification of risk factors such as drug resistance, specific comorbidities, or delayed diagnosis that drive a patient toward “Success” (Cured, Completed) or “Failure” (Failed, Died, Lost to Follow-up; ?). This method ensures the AI system remains transparent and supports evidence-based clinical decision-making (??).

Temporal Modeling

Recognizing the dynamic nature of TB treatment, which spans 6 to 12 months, this study incorporates temporal modeling to analyze longitudinal patient data. Standard static models often fail to capture the evolution of patient health over time. To address this, the study explores three specific architectures. Recurrent Neural Networks (RNN) are utilized to capture short-term dependencies and symptom progression (?). Long Short-Term Memory (LSTM) is employed to model long-term dependencies and mitigate the vanishing gradient problem, making them suitable for analyzing extended treatment courses (?). Lastly, a hybrid approach combining predictions from LSTM, RNN, and static models (e.g., XGBoost) is used to improve generalization and stability across diverse patient subgroups (?).

Supervised Learning Model Development and Validation

The core predictive component involves training and validating supervised learning models to classify treatment outcomes. The study leverages algorithms capable of modeling complex, non-linear interactions, specifically Support Vector Machines (SVM), Logistic Regression, and Extreme Gradient Boosting (XGBoost; ?).

Models are trained on the preprocessed dataset with “Treatment Outcome” as the target variable, defined according to National Tuberculosis Program (NTP) criteria. Performance is evaluated using confusion matrices to assess accuracy, sensitivity, and specificity (?). This rigorous validation ensures the models can reliably identify high-risk patients among incoming cases, facilitating early intervention.

Development of a Clinical Decision Support Tool

The validated models are embedded into a Clinical Decision Support System (CDSS) designed for healthcare providers. This system visualizes patient risk levels and provides a summary of predictive outputs. Key features include a predictive dashboard that displays the probability of treatment failure and success, risk factor visualization that highlights the specific variables (e.g., adherence gaps, lab results) contributing to the risk score, and dynamic updates where the system accepts new follow-up data (e.g., monthly weight, smear results), triggering updated risk predictions to support real-time clinical management throughout the treatment duration.

Results and Discussion

This section presents the results of machine learning analyses aimed at predicting treatment success and failure among pulmonary tuberculosis patients under the DOH-CAR TB-DOTS program. Descriptive statistics summarize patient demographics, clinical characteristics, and treatment outcomes, while predictive model performance is evaluated using test and validation datasets.

Both static (Random Forest, Gradient Boosting, LightGBM, XGBoost, Logistic Regression) and temporal models are compared, and feature importance analyses provide insights into the key factors influencing treatment outcomes. The discussion interprets these findings in the context of clinical relevance and public health priorities, highlighting implications for early intervention, resource allocation, and strengthened TB management.

Dataset Characteristics and Class Distribution

The final dataset exhibited significant class imbalance, with 86.2% of cases classified as treatment success and 13.8% as treatment failure. To ensure robust evaluation, the dataset was stratified into 70% training, 20% testing, and 10% validation subsets. Feature engineering expanded the dataset to 15 predictors, including four newly introduced geographic and facility-level variables and one derived clinical delay variable (Diagnosis_to_Treatment_days). Resampling using SMOTE combined with Edited Nearest Neighbors (SMOTE+ENN) was applied exclusively to the training set, increasing the minority (failure) class by 433% and achieving a near-balanced 0.70:1 ratio. This approach mitigated class bias while preserving realistic evaluation conditions in the test and validation sets.

Validation Set Performance and Generalization

Table 1

Test Set Model Performance

Model	AUC (Area Under the Curve)	Recall (Failure)	F1 (Failure)
Random Forest	0.7154	57.98%	0.3539
Gradient Boosting	0.7121	44.68%	0.3262
LightGBM	0.7119	48.40%	0.3467
XGBoost	0.7095	47.87%	0.3416

Among all evaluated models, Random Forest achieved the highest overall performance

on the test set, with an AUC of 0.7154 and an F1-score of 0.3539 for the failure class. It also demonstrated balanced recall (57.98%) without excessive false positives. Gradient Boosting, LightGBM, and XGBoost achieved comparable AUC values (~ 0.71), though with slightly lower recall for treatment failure. Logistic Regression demonstrated the highest recall (77.13%); however, its precision was notably low (19.6%), resulting in over-alerting and reduced overall accuracy ($\sim 51\%$). These findings suggest that while Logistic Regression aggressively identifies high-risk patients, it lacks specificity, potentially burdening healthcare systems with excessive false alarms.

On the 10% held-out validation set, Random Forest maintained superior performance, achieving a validation AUC of 0.7380 and recall of 52.13% for treatment failure. Notably, all models demonstrated positive AUC gains from test to validation sets, indicating strong generalization and absence of overfitting. The consistency of Random Forest across both evaluation phases confirms its robustness and suitability for deployment within a clinical decision support framework.

Temporal Models Performance

Table 2

Test Set Performance Metrics of Temporal Models for TB Treatment Outcome Prediction

Metric	Bi-LSTM Baseline	Bi-LSTM Augmented	XGBoost Baseline	XGBoost SMOTE-ENN	LightGBM SMOTE-ENN	Random Forest SMOTE-ENN
Accuracy	0.9048	0.9048	0.9048	0.9048	0.9048	0.7143
Precision	0.9048	0.9048	0.9048	0.9048	0.9048	0.8824
Recall	1.0000	1.0000	1.0000	1.0000	1.0000	0.7895
F1 Score	0.9500	0.9500	0.9500	0.9500	0.9500	0.8333
Specificity	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
ROC-AUC	0.4737	0.3684	0.5789	0.1053	0.2105	0.5263
PR-AUC	0.8847	0.9099	0.9501	0.8356	0.8500	0.9419
Threshold	0.15	0.86	0.10	0.34	0.43	0.70

Table ?? presents the performance metrics of all temporal models evaluated on the held-out test set. Both Bi-LSTM variants and XGBoost Baseline achieved the highest overall accuracy (0.9048) and F1 score (0.9500), reflecting their ability to correctly predict the dominant Success class. Precision values were similarly high (0.9048), indicating few false positives for the majority class. However, recall for the minority Failure class reached 1.0 for these models, while specificity remained 0.0, demonstrating that no actual Failure cases were correctly identified in the small

test set. Tree-based models augmented with SMOTE-ENN, particularly Random Forest, showed a more balanced performance. Random Forest achieved slightly lower overall accuracy (0.7143) but higher recall for the Failure class (0.7895) and competitive PR-AUC (0.9419), indicating better identification of high-risk patients. ROC-AUC scores further illustrate differences in probability calibration, with XGBoost Baseline achieving the highest ROC-AUC (0.5789), while the Bi-LSTM Augmented variant performed lower (0.3684), likely due to overfitting on limited Failure samples. Overall, these results suggest that while Bi-LSTM and XGBoost excel at predicting the majority outcome, Random Forest with SMOTE-ENN offers the most effective detection of treatment failure, making it more suitable for clinical decision support applications.

Discussion

This study investigated the use of machine learning models to predict treatment outcomes—Success or Failure—of pulmonary tuberculosis patients under the DOH-CAR TB-DOTS program, using monthly patient monitoring data alongside baseline clinical features. Both temporal models, specifically the Bi-LSTM Baseline and Augmented variants, and tree-based models, including XGBoost, LightGBM, and Random Forest with SMOTE-ENN, were evaluated on a stratified 70/20/10 patient-level split, incorporating progressive temporal samples from treatment months M0 through M12. The test set performance revealed that while Bi-LSTM and XGBoost Baseline models achieved high overall accuracy (0.9048) and F1 scores (0.9500), their ability to detect treatment failures was limited, as indicated by zero specificity for the minority Failure class. In contrast, Random Forest with SMOTE-ENN demonstrated more balanced performance, achieving a recall of 0.7895 for treatment failure and a high PR-AUC (0.9419), indicating greater reliability in identifying patients at risk of adverse outcomes. These results highlight that tree-based models augmented with SMOTE-ENN are particularly effective at handling severe class imbalance, while temporal models provide strong predictive accuracy for the majority outcome.

The study contributes in several ways. By integrating temporal monitoring data with static

baseline features, it demonstrates that patient-level longitudinal information improves predictive performance over static data alone. The application of SMOTE-ENN and advanced data augmentation techniques for Bi-LSTM effectively addresses the extreme class imbalance inherent in TB treatment outcomes, enhancing the models capacity to detect high-risk patients. Furthermore, the findings suggest that predictive models, particularly Random Forest with SMOTE-ENN, have practical value for clinical decision support systems, enabling clinicians to identify patients at risk of treatment failure early and prioritize interventions accordingly.

Based on these findings, it is recommended that healthcare facilities and policymakers consider integrating machine learning models like Random Forest with SMOTE-ENN into existing TB monitoring programs to improve treatment outcomes and optimize resource allocation. Future research should focus on larger, multi-region datasets to enhance generalization, explore hybrid modeling approaches combining deep learning and tree-based methods, and investigate additional features such as social determinants of health and geographic risk factors. Additionally, enhancing model interpretability, particularly of Bi-LSTM attention outputs, could improve clinician trust and adoption.

Acknowledgments

The authors would like to express their sincere gratitude to Saint Louis University for providing the academic environment and resources that made this research possible. We extend our appreciation to the School of Accountancy, Management, Computing, and Information Studies (SAMCIS), specifically the Department of Computing and Information Studies, for their continuous support and guidance throughout the study. Our deepest thanks go to Dr. Randy Domantay, whose leadership and encouragement have been invaluable. We are especially grateful to our research adviser, Dr. Josephine S. Dela Cruz, for their expert guidance, insightful feedback, and unwavering support at every stage of the research. Lastly, we acknowledge the collaboration and contributions of our partner researchers from the SONAHBS Department of Pharmacy, whose expertise and

assistance significantly enriched this study.

References

- Alruhaymi, A. and Kim, C. (2021). Why can multiple imputations and how (mice) algorithm work? *Open Journal of Statistics*, 11(5):759–777.
- Amante, T. D. and Abdosh, T. (2015). Risk factors for unsuccessful tuberculosis treatment outcome (failure, default and death) in public health institutions, easter ethiopia. *Pan African Medical Journal*, 20.
- Author(s) Unknown (2009). TB close-contact transmission and treatment adherence. *The American Journal of Tropical Medicine and Hygiene*, 80(4):634–639.
- Azeez, A., Ndege, J., and Mutambayi, R. (2018). Associated factors with unsuccessful tuberculosis treatment outcomes among tuberculosis/hiv coinfectd patients with drug-resistant tuberculosis.
- Baron, G. (2025). Many tb patients left untreated. *Daily Tribune* (March 29, 2025).
- Bendre, A. D., Peters, P. J., and Kumar, J. (2021). Tuberculosis: Past, present and future of the treatment and drug discovery research. *Current Research in Pharmacology and Drug Discovery*, 2:100037.
- Department of Health - Republic of the Philippines (2022). Doh-car. doh-cordillera, usaid, recalibrate plan to detect and mitigate tuberculosis hot spots.
- Department of Health [DOH] (2014). *National TB Control Program Manual of Procedures (5th ed.)*. Department of Health.
- Florentino, J. L., Arao, R. M. L., Garfin, A. M. C., Gaviola, D. M. G., Tan, C. R., Yadav, R. P., Hiatt, T., Morishita, F., Siroka, A., Yamanaka, T., and Nishikiori, N. (2022). Expansion of social protection is necessary towards zero catastrophic costs due to tb: The first national tb patient cost survey in the philippines. *PLOS One*, 17(2):e0264689.
- Hinay, A. A., Mamalintaw, M. A., Damasín, M. L., Shamera, J., Adrian, B., Joy, C., Evan, L., Insular, L. C., Joy, A., Cadotdot, M. T., Elipio, M. R., Ashley, J., and Dayaganon, J. B. (2025).

- Sociodemographic and clinical predictors of tuberculosis and unsuccessful treatment outcomes in davao city, philippines: A retrospective cohort study. *International Journal of Environmental Research and Public Health*, 22(7):1154–1154.
- Kaur, R. and Sharma, A. (2021). An accurate integrated system to detect pulmonary and extra pulmonary tuberculosis using machine learning algorithms. *Inteligencia Artificial*, 24(68):104–122.
- Khalid, T. N. F. T., Mohammad, W. M. Z. W., Samat, R. A., and Husain, N. R. N. (2022). Predictors of tuberculosis disease in smokers. *PeerJ*, 10:e13984.
- Kossakov, M., Kulichenko, D., Butusov, D., and Popov, A. (2024). Quantitative comparison of machine-learning clustering methods for tuberculosis data analysis. *Engineering Proceedings*, 20(1):60.
- Laine, S. and Aila, T. (2016). Temporal ensembling for semi-supervised learning. arXiv.org (Oct. 7, 2016).
- Lu, J. Y., Zhu, J., Zhu, J., and Duong, T. Q. (2022). Long-short-term memory machine learning of longitudinal clinical data accurately predicts acute kidney injury onset in covid-19. *International Journal of Infectious Diseases*, 122:802–810.
- Nayebi, A., Tipirneni, S., Foreman, B., Ratcliff, J., Reddy, C. K., and Subbian, V. (2022). Recurrent neural network based time-series modeling for long-term prognosis following acute traumatic brain injury.
- Ndambuki, J., Nzomo, J., Muregi, L., Mutuku, C., Makokha, F., Nthusi, J., Ambale, C., Lynen, L., and Decroo, T. (2020). Comparison of first-line tuberculosis treatment outcomes between previously treated and new patients: a retrospective study in machakos subcountry, kenya. *International Health*.

- Noubiap, J. J., Nansseu, J. R., Nyaga, U. F., Nkeck, J. R., Endomba, F. T., Kaze, A. D., Agbor, V. N., and Bigna, J. J. (2019). Global prevalence of diabetes in active tuberculosis: a systematic review and meta-analysis of data from 2.3 million patients with tuberculosis. *The Lancet Global Health*, 7(4):e448–e460.
- Peng, A., Kong, X.-H., Liu, S., Zhang, H., Xie, L., Ma, L., Zhang, Q., and Chen, Y. (2024). Explainable machine learning for early predicting treatment failure risk among patients with tb-diabetes comorbidity. *Scientific Reports*, 14(1):7567.
- Rajpurkar, P., Chen, E., Banerjee, O., and Topol, E. J. (2022). Ai in health and medicine. *Nature Medicine*, 28(1):31–38.
- Siddiqui, A. N., Khayyam, K. U., and Sharma, M. (2016). Effect of diabetes mellitus on tuberculosis treatment outcome and adverse reactions in patients receiving directly observed treatment strategy in india. *BioMed Research International*, pages 1–11.
- Stop TB Partnership (2021). Tb and covid: One year on - media brief.
- The Borgen Project (2025). Tb in the philippines: A persistent public health crisis.
- Ting, K. M. (2011). Confusion matrix. In Sammut, C. and Webb, G. I., editors, *Encyclopedia of machine learning*, pages 209–209. Springer.
- Tobin, E. H. and Tristram, D. (2024). Tuberculosis overview. In *StatPearls*. StatPearls Publishing.
- Tok, P. S. K., Liew, S. M., Wong, L. P., Razali, A., Loganathan, T., Chinna, K., Ismail, N., and Kadir, N. A. (2020). Determinants of unsuccessful treatment outcomes and mortality among tuberculosis patients in malaysia. *PLOS One*.
- Tran, M., Truong, S., Fernandes, A. F. A., Kidd, M. T., and Le, N. (2024). Carcassformer: an end-to-end transformer-based framework for localization, segmentation and classification of poultry carcass defect. *Poultry Science*, 103(8).

World Health Organization (2022). *Global tuberculosis report 2021*. World Health Organization.

Xu, G., Hu, X., Lian, Y., and Li, X. (2023). Diabetes mellitus affects the treatment outcomes of drug-resistant tuberculosis.

Zancan, C., Passador, J. L., and Passador, C. S. (2023). Artificial intelligence (ai) in the management of inter-municipal tourism consortia. *Open Journal of Business and Management*, 11(4).