

Let's calculate smth with a given function: $f(x, y) = \sin x \cdot y^{2.000}$
Firstly, let's insert all constants and simplify this expression: $f(x, y) = \sin x \cdot y^{2.000}$
BRITISH SCIENTISTS WERE SHOCKED, WHEN THEY COUNT IT!!!
In the point ($x = 3.000, y = 2.000$) it's value = 0.564
Personally, I've always thought about first derivation of something like that function... Haven't you?
But now, by using informatics and math skills I feel that I'm prepared enough to calculate it!
1 step. finding a derivation of:

While preparing for exams, I learned a lot of new things, for example: $(y)' = 1.000$
2 step. finding a derivation of:

It's really easy to find: $(y^{2.000})' = 2.000 \cdot y$
3 step. finding a derivation of:

My roommate mumbled it in his sleep all night: $(x)' = 1.000$
4 step. finding a derivation of:

Sounds logical that it is the same as: $(\sin x)' = \cos x$
5 step. finding a derivation of:

For centuries, people have hunted for the secret knowledge that: $(\sin x \cdot y^{2.000})' = \cos x \cdot y^{2.000} + 2.000 \cdot y \cdot \sin x$
Congratulations! The first derivation of the expression is:

In the point ($x = 3.000, y = 2.000$) it's value = -3.395
Let's calculate the 4 derivation of the expression:
Calculating the 1 derivation of the expression:
1 step. finding a derivation of:

Sounds logical that it is the same as: $(y)' = 1.000$
2 step. finding a derivation of:

It's really easy to find: $(y^{2.000})' = 2.000 \cdot y$
3 step. finding a derivation of:

My roommate mumbled it in his sleep all night: $(x)' = 1.000$
4 step. finding a derivation of:

What if it equals: $(\sin x)' = \cos x$
5 step. finding a derivation of:

It's really easy to find: $(\sin x \cdot y^{2.000})' = \cos x \cdot y^{2.000} + 2.000 \cdot y \cdot \sin x$
Calculating the 2 derivation of the expression:
1 step. finding a derivation of:

Even my two-aged sister knows that it equals: $(x)' = 1.000$
2 step. finding a derivation of:

When I was child, my father always told me: "Remember, son: $(\sin x)' = \cos x$
3 step. finding a derivation of:

I spend the hole of my life to find the answer and finally it's: $(y)' = 1.000$
4 step. finding a derivation of:

Man... Just look: $(2.000)' = 0.000$
5 step. finding a derivation of:

For centuries, people have hunted for the secret knowledge that: $(2.000 \cdot y)' = 2.000$
6 step. finding a derivation of:

It's really easy to find: $(2.000 \cdot y \cdot \sin x)' = 2.000 \cdot \sin x + \cos x \cdot 2.000 \cdot y$
7 step. finding a derivation of:

It's simple as fuck: $(y)' = 1.000$
8 step. finding a derivation of:

thanks to the results of my colleagues' scientific work, I know that it equals: $(y^{2.000})' = 2.000 \cdot y$
9 step. finding a derivation of:

When I was child, my father always told me: "Remember, son: $(x)' = 1.000$
10 step. finding a derivation of:

It's really easy to find: $(\cos x)' = (-1.000) \cdot \sin x$

CrInGeCrInGeProduction.Supereringeintroductionhere :

$$y$$

$$y^{2.000}$$

$$x$$

$$\sin x$$

$$\sin x \cdot y^{2.000}$$

$$\cos x \cdot y^{2.000} + 2.000 \cdot y \cdot \sin x$$

$$y$$

$$y^{2.000}$$

$$x$$

$$\sin x$$

$$\sin x \cdot y^{2.000}$$

$$x$$

$$\sin x$$

$$y$$

$$2.000$$

$$2.000 \cdot y$$

$$2.000 \cdot y \cdot \sin x$$

$$y$$

$$y^{2.000}$$

$$x$$

$$\cos x$$

$$1$$

11 step. finding a derivation of:

I was asked not to tell anyone that: $(\cos x \cdot y^{2.000})' = (-1.000) \cdot \sin x \cdot y^{2.000} + 2.000 \cdot y \cdot \cos x$
12 step. finding a derivation of:

For centuries, people have hunted for the secret knowledge that: $(\cos x \cdot y^{2.000} + 2.000 \cdot y \cdot \sin x)' = (-1.000) \cdot \sin x \cdot y^{2.000} + 2.000 \cdot y \cdot \cos x + 2.000 \cdot \sin x + \cos x \cdot 2.000 \cdot y$
Calculating the 3 derivation of the expression:
1 step. finding a derivation of:

My roommate mumbled it in his sleep all night: $(y)' = 1.000$
2 step. finding a derivation of:

What if it equals: $(2.000)' = 0.000$
3 step. finding a derivation of:

Even my two-aged sister knows that it equals: $(2.000 \cdot y)' = 2.000$
4 step. finding a derivation of:

I spend the hole of my life to find the answer and finally it's: $(x)' = 1.000$
5 step. finding a derivation of:

Even my two-aged sister knows that it equals: $(\cos x)' = (-1.000) \cdot \sin x$
6 step. finding a derivation of:

While preparing for exams, I learned a lot of new things, for example: $(\cos x \cdot 2.000 \cdot y)' = (-1.000) \cdot \sin x \cdot 2.000 \cdot y + 2.000 \cdot \cos x$
7 step. finding a derivation of:

When I was child, my father always told me: "Remember, son: $(x)' = 1.000$
8 step. finding a derivation of:

Sounds logical that it is the same as: $(\sin x)' = \cos x$
9 step. finding a derivation of:

A true prince must know that it equals: $(2.000)' = 0.000$
10 step. finding a derivation of:

My roommate mumbled it in his sleep all night: $(2.000 \cdot \sin x)' = 2.000 \cdot \cos x$
11 step. finding a derivation of:

My roommate mumbled it in his sleep all night: $(2.000 \cdot \sin x + \cos x \cdot 2.000 \cdot y)' = 2.000 \cdot \cos x + (-1.000) \cdot \sin x \cdot 2.000 \cdot y + 2.000 \cdot \cos x$
12 step. finding a derivation of:

If someone asked me that in the middle of the night, I wouldn't hesitate to say: $(x)' = 1.000$
13 step. finding a derivation of:

A true prince must know that it equals: $(\cos x)' = (-1.000) \cdot \sin x$
14 step. finding a derivation of:

My roommate mumbled it in his sleep all night: $(y)' = 1.000$
15 step. finding a derivation of:

While preparing for exams, I learned a lot of new things, for example: $(2.000)' = 0.000$
16 step. finding a derivation of:

It's really easy to find: $(2.000 \cdot y)' = 2.000$
17 step. finding a derivation of:

It's really easy to find: $(2.000 \cdot y \cdot \cos x)' = 2.000 \cdot \cos x + (-1.000) \cdot \sin x \cdot 2.000 \cdot y$
18 step. finding a derivation of:

When I was child, my father always told me: "Remember, son: $(y)' = 1.000$
19 step. finding a derivation of:

What if it equals: $(y^{2.000})' = 2.000 \cdot y$
20 step. finding a derivation of:

If someone asked me that in the middle of the night, I wouldn't hesitate to say: $(x)' = 1.000$
21 step. finding a derivation of:

thanks to the results of my colleagues' scientific work, I know that it equals: $(\sin x)' = \cos x$
22 step. finding a derivation of:

A true prince must know that it equals: $((-1.000))' = 0.000$

$$\cos x \cdot y^{2.000}$$

$$\cos x \cdot y^{2.000} + 2.000 \cdot y \cdot \sin x$$

$$y$$

$$2.000$$

$$2.000 \cdot y$$

$$x$$

$$\cos x$$

$$\cos x \cdot 2.000 \cdot y$$

$$x$$

$$\sin x$$

$$2.000$$

$$2.000 \cdot \sin x$$

$$2.000 \cdot \sin x + \cos x \cdot 2.000 \cdot y$$

$$x$$

$$\cos x$$

$$y$$

$$2.000$$

$$2.000 \cdot y$$

$$2.000 \cdot y \cdot \cos x$$

$$y$$

$$y^{2.000}$$

$$x$$

$$\sin x$$

$$(-1.000)$$

$$2$$

23 step. finding a derivation of:

A true prince must know that it equals: $((-1.000) \cdot \sin x)' = (-1.000) \cdot \cos x$

24 step. finding a derivation of:

When I was child, my father always told me: "Remember, son: $((-1.000) \cdot \sin x \cdot y^{2.000})' = (-1.000) \cdot \cos x \cdot y^{2.000} + 2.000 \cdot y \cdot (-1.000) \cdot \sin x$

25 step. finding a derivation of:

For centuries, people have hunted for the secret knowledge that: $((-1.000) \cdot \sin x \cdot y^{2.000} + 2.000 \cdot y \cdot \cos x)' = (-1.000) \cdot \cos x \cdot y^{2.000} + 2.000 \cdot y \cdot (-1.000) \cdot \sin x + 2.000 \cdot \cos x + (-1.000) \cdot \sin x \cdot 2.000 \cdot y$

26 step. finding a derivation of:

A true prince must know that it equals: $((-1.000) \cdot \sin x \cdot y^{2.000} + 2.000 \cdot y \cdot \cos x + 2.000 \cdot \sin x + \cos x \cdot 2.000 \cdot y)' = (-1.000) \cdot \cos x \cdot y^{2.000} + 2.000 \cdot y \cdot (-1.000) \cdot \sin x + 2.000 \cdot \cos x + (-1.000) \cdot \sin x \cdot 2.000 \cdot y + 2.000 \cdot \cos x + (-1.000) \cdot \sin x \cdot 2.000 \cdot y + 2.000 \cdot \cos x$

Calculating the 4 derivation of the expression:

1 step. finding a derivation of:

I spend the hole of my life to find the answer and finally it's: $(x)' = 1.000$

2 step. finding a derivation of:

It's simple as fuck: $(\cos x)' = (-1.000) \cdot \sin x$

3 step. finding a derivation of:

For centuries, people have hunted for the secret knowledge that: $(2.000)' = 0.000$

4 step. finding a derivation of:

It's really easy to find: $(2.000 \cdot \cos x)' = 2.000 \cdot (-1.000) \cdot \sin x$

5 step. finding a derivation of:

It's really easy to find: $(y)' = 1.000$

6 step. finding a derivation of:

I spend the hole of my life to find the answer and finally it's: $(2.000)' = 0.000$

7 step. finding a derivation of:

I was asked not to tell anyone that: $(2.000 \cdot y)' = 2.000$

8 step. finding a derivation of:

If someone asked me that in the middle of the night, I wouldn't hesitate to say: $(x)' = 1.000$

9 step. finding a derivation of:

Even my two-aged sister knows that it equals: $(\sin x)' = \cos x$

10 step. finding a derivation of:

I spend the hole of my life to find the answer and finally it's: $((-1.000))' = 0.000$

11 step. finding a derivation of:

It's really easy to find: $((-1.000) \cdot \sin x)' = (-1.000) \cdot \cos x$

12 step. finding a derivation of:

It's really easy to find: $((-1.000) \cdot \sin x \cdot 2.000 \cdot y)' = (-1.000) \cdot \cos x \cdot 2.000 \cdot y + 2.000 \cdot (-1.000) \cdot \sin x$

13 step. finding a derivation of:

It's simple as fuck: $((-1.000) \cdot \sin x \cdot 2.000 \cdot y + 2.000 \cdot \cos x)' = (-1.000) \cdot \cos x \cdot 2.000 \cdot y + 2.000 \cdot (-1.000) \cdot \sin x + 2.000 \cdot (-1.000) \cdot \sin x$

14 step. finding a derivation of:

It's simple as fuck: $(x)' = 1.000$

15 step. finding a derivation of:

A true prince must know that it equals: $(\cos x)' = (-1.000) \cdot \sin x$

16 step. finding a derivation of:

My roommate mumbled it in his sleep all night: $(2.000)' = 0.000$

17 step. finding a derivation of:

I was asked not to tell anyone that: $(2.000 \cdot \cos x)' = 2.000 \cdot (-1.000) \cdot \sin x$

18 step. finding a derivation of:

I spend the hole of my life to find the answer and finally it's: $(2.000 \cdot \cos x + (-1.000) \cdot \sin x \cdot 2.000 \cdot y + 2.000 \cdot \cos x)' = 2.000 \cdot (-1.000) \cdot \sin x + (-1.000) \cdot \cos x \cdot 2.000 \cdot y + 2.000 \cdot (-1.000) \cdot \sin x + 2.000 \cdot (-1.000) \cdot \sin x$

19 step. finding a derivation of:

It's really easy to find: $(y)' = 1.000$

20 step. finding a derivation of:

What if it equals: $(2.000)' = 0.000$

$$(-1.000) \cdot \sin x$$

$$(-1.000) \cdot \sin x \cdot y^{2.000}$$

$$(-1.000) \cdot \sin x \cdot y^{2.000} + 2.000 \cdot y \cdot \cos x$$

$$(-1.000) \cdot \sin x \cdot y^{2.000} + 2.000 \cdot y \cdot \cos x + 2.000 \cdot \sin x + \cos x \cdot 2.000 \cdot y$$

$$x$$

$$\cos x$$

$$2.000$$

$$2.000 \cdot \cos x$$

$$y$$

$$2.000$$

$$2.000 \cdot y$$

$$x$$

$$\sin x$$

$$(-1.000)$$

$$(-1.000) \cdot \sin x$$

$$(-1.000) \cdot \sin x \cdot 2.000 \cdot y$$

$$(-1.000) \cdot \sin x \cdot 2.000 \cdot y + 2.000 \cdot \cos x$$

$$x$$

$$\cos x$$

$$2.000$$

$$2.000 \cdot \cos x$$

$$2.000 \cdot \cos x + (-1.000) \cdot \sin x \cdot 2.000 \cdot y + 2.000 \cdot \cos x$$

$$y$$

$$2.000$$

$$3$$

21 step. finding a derivation of:

While preparing for exams, I learned a lot of new things, for example: $(2.000 \cdot y)' = 2.000$

22 step. finding a derivation of:

Even my two-aged sister knows that it equals: $(x)' = 1.000$

23 step. finding a derivation of:

What if it equals: $(\sin x)' = \cos x$

24 step. finding a derivation of:

It's simple as fuck: $((-1.000))' = 0.000$

25 step. finding a derivation of:

My roommate mumbled it in his sleep all night: $((-1.000) \cdot \sin x)' = (-1.000) \cdot \cos x$

26 step. finding a derivation of:

It's simple as fuck: $((-1.000) \cdot \sin x \cdot 2.000 \cdot y)' = (-1.000) \cdot \cos x \cdot 2.000 \cdot y + 2.000 \cdot (-1.000) \cdot \sin x$

27 step. finding a derivation of:

A true prince must know that it equals: $(x)' = 1.000$

28 step. finding a derivation of:

My roommate mumbled it in his sleep all night: $(\cos x)' = (-1.000) \cdot \sin x$

29 step. finding a derivation of:

A true prince must know that it equals: $(2.000)' = 0.000$

30 step. finding a derivation of:

A true prince must know that it equals: $(2.000 \cdot \cos x)' = 2.000 \cdot (-1.000) \cdot \sin x$

31 step. finding a derivation of:

If someone asked me that in the middle of the night, I wouldn't hesitate to say: $(2.000 \cdot \cos x + (-1.000) \cdot \sin x \cdot 2.000 \cdot y)' = 2.000 \cdot (-1.000) \cdot \sin x + (-1.000) \cdot \cos x \cdot 2.000 \cdot y + 2.000 \cdot (-1.000) \cdot \sin x$

32 step. finding a derivation of:

I spend the hole of my life to find the answer and finally it's: $(x)' = 1.000$

33 step. finding a derivation of:

When I was child, my father always told me: "Remember, son: $(\sin x)' = \cos x$

34 step. finding a derivation of:

If someone asked me that in the middle of the night, I wouldn't hesitate to say: $((-1.000))' = 0.000$

35 step. finding a derivation of:

thanks to the results of my colleagues' scientific work, I know that it equals: $((-1.000) \cdot \sin x)' = (-1.000) \cdot \cos x$

36 step. finding a derivation of:

For centuries, people have hunted for the secret knowledge that: $(y)' = 1.000$

37 step. finding a derivation of:

For centuries, people have hunted for the secret knowledge that: $(2.000)' = 0.000$

38 step. finding a derivation of:

What if it equals: $(2.000 \cdot y)' = 2.000$

39 step. finding a derivation of:

If someone asked me that in the middle of the night, I wouldn't hesitate to say: $(2.000 \cdot y \cdot (-1.000) \cdot \sin x)' = 2.000 \cdot (-1.000) \cdot \sin x + (-1.000) \cdot \cos x \cdot 2.000 \cdot y$

40 step. finding a derivation of:

I was asked not to tell anyone that: $(y)' = 1.000$

41 step. finding a derivation of:

A true prince must know that it equals: $(y^{2.000})' = 2.000 \cdot y$

42 step. finding a derivation of:

Man... Just look: $(x)' = 1.000$

43 step. finding a derivation of:

My roommate mumbled it in his sleep all night: $(\cos x)' = (-1.000) \cdot \sin x$

44 step. finding a derivation of:

It's simple as fuck: $((-1.000))' = 0.000$

45 step. finding a derivation of:

$$2.000 \cdot y$$

$$x$$

$$\sin x$$

$$(-1.000)$$

$$(-1.000) \cdot \sin x$$

$$(-1.000) \cdot \sin x \cdot 2.000 \cdot y$$

$$x$$

$$\cos x$$

$$2.000$$

$$2.000 \cdot \cos x$$

$$2.000 \cdot \cos x + (-1.000) \cdot \sin x \cdot 2.000 \cdot y$$

$$x$$

$$\sin x$$

$$(-1.000)$$

$$(-1.000) \cdot \sin x$$

$$y$$

$$2.000$$

$$2.000 \cdot y$$

$$2.000 \cdot y \cdot (-1.000) \cdot \sin x$$

$$y$$

$$y^{2.000}$$

$$x$$

$$\cos x$$

$$(-1.000)$$

$$(-1.000) \cdot \cos x$$

I was asked not to tell anyone that: $((-1.000) \cdot \cos x)' = (-1.000) \cdot (-1.000) \cdot \sin x$
46 step. finding a derivation of:

If someone asked me that in the middle of the night, I wouldn't hesitate to say: $((-1.000) \cdot \cos x \cdot y^{2.000})' = (-1.000) \cdot (-1.000) \cdot \sin x \cdot y^{2.000} + 2.000 \cdot y \cdot (-1.000) \cdot \cos x$
47 step. finding a derivation of:

Man... Just look: $((-1.000) \cdot \cos x \cdot y^{2.000} + 2.000 \cdot y \cdot (-1.000) \cdot \sin x)' = (-1.000) \cdot (-1.000) \cdot \sin x \cdot y^{2.000} + 2.000 \cdot y \cdot (-1.000) \cdot \cos x + 2.000 \cdot (-1.000) \cdot \sin x + (-1.000) \cdot \cos x \cdot 2.000 \cdot y$
48 step. finding a derivation of:

When I was child, my father always told me: "Remember, son: $((-1.000) \cdot \cos x \cdot y^{2.000} + 2.000 \cdot y \cdot (-1.000) \cdot \sin x + 2.000 \cdot \cos x + (-1.000) \cdot \sin x \cdot 2.000 \cdot y)' = (-1.000) \cdot (-1.000) \cdot \sin x \cdot y^{2.000} + 2.000 \cdot y \cdot (-1.000) \cdot \cos x + 2.000 \cdot (-1.000) \cdot \sin x + (-1.000) \cdot \cos x \cdot 2.000 \cdot y + 2.000 \cdot (-1.000) \cdot \sin x$
49 step. finding a derivation of:

It's really easy to find: $((-1.000) \cdot \cos x \cdot y^{2.000} + 2.000 \cdot y \cdot (-1.000) \cdot \sin x + 2.000 \cdot \cos x + (-1.000) \cdot \sin x \cdot 2.000 \cdot y + 2.000 \cdot \cos x + (-1.000) \cdot \sin x \cdot 2.000 \cdot y + 2.000 \cdot \cos x)' = (-1.000) \cdot (-1.000) \cdot \sin x \cdot y^{2.000} + 2.000 \cdot y \cdot (-1.000) \cdot \cos x + 2.000 \cdot (-1.000) \cdot \sin x + (-1.000) \cdot \cos x \cdot 2.000 \cdot y + 2.000 \cdot (-1.000) \cdot \sin x + (-1.000) \cdot \cos x \cdot 2.000 \cdot y + 2.000 \cdot (-1.000) \cdot \sin x + 2.000 \cdot (-1.000) \cdot \sin x + (-1.000) \cdot \cos x \cdot 2.000 \cdot y + 2.000 \cdot (-1.000) \cdot \sin x + 2.000 \cdot (-1.000) \cdot \sin x$
Finally... The 4 derivation of the expression:

BRITISH SCIENTISTS WERE SHOCKED AGAIN, WHEN THEY COUNT THE 4 DERIVATION OF THIS EXPRESSION!!!

In the point (x = 3.000, y = 2.000) it's value = 14.711
Partial derivation of the expression on the variable x:
 $\frac{\partial f}{\partial x} = 4.000 \cdot \cos x$
In the point (x = 3.000, y = 2.000) it's value = -3.959970 !!!
Partial derivation of the expression on the variable y:
 $\frac{\partial f}{\partial y} = 0.141 \cdot 2.000 \cdot y$
In the point (x = 3.000, y = 2.000) it's value = 0.564480 !!!
Full derivation:

In the point (x = 3.000, y = 2.000) it's value = 4.000 !!!

Let's consider the expression as a function of x variable: $f(x) = 4.000 \cdot \sin x$

Maklorens formula for $x \rightarrow 3.000$: $f(x) = 0.564 + (-3.960) \cdot (x - 3.000) + (-0.282) \cdot (x - 3.000)^{2.000} + 0.660 \cdot (x - 3.000)^{3.000} + 0.024 \cdot (x - 3.000)^{4.000} + (-0.033) \cdot (x - 3.000)^{5.000} + o((x - 3.000)^{5.000})$

Graph f(x):

$$(-1.000) \cdot \cos x \cdot y^{2.000}$$

$$(-1.000) \cdot \cos x \cdot y^{2.000} + 2.000 \cdot y \cdot (-1.000) \cdot \sin x$$

$$(-1.000) \cdot \cos x \cdot y^{2.000} + 2.000 \cdot y \cdot (-1.000) \cdot \sin x + 2.000 \cdot \cos x + (-1.000) \cdot \sin x \cdot 2.000 \cdot y$$

$$(-1.000) \cdot \cos x \cdot y^{2.000} + 2.000 \cdot y \cdot (-1.000) \cdot \sin x + 2.000 \cdot \cos x + (-1.000) \cdot \sin x \cdot 2.000 \cdot y + 2.000 \cdot \cos x + (-1.000) \cdot \sin x \cdot 2.000 \cdot y + 2.000 \cdot \cos x$$

$$\sqrt{(4.000 \cdot \cos x)^{2.000} + (0.141 \cdot 2.000 \cdot y)^{2.000}}$$

Tangent equation in the point $x = 0.000$: $f(x) = 4.000 \cdot x$
Normal equation in the point $x = 0.000$: $f(x) = (-0.250) \cdot (x - 0.000) + 0.000$