

1 Introduction

CrInGeCrInGeProduction.Supercringeintroductionhere :

2 Some basic knowledge about researching problem...

Parameters and constants we use in this work:

Constants (3):  
e = 2.718282  
pi = 3.141593  
AbObA = 1337.228690

Variables (3):  
a = 3.141500  
kek = 13.000000  
x = 1.000000

Parameters of exploration :  
Number of differentiates = 2  
Macloren's accuracy = 3  
Tanget point = 0.200000  
Delta coverage of tangent point = 2.500000  
Graph diapasone = [-1 : 15]

So let's calculate smth with a given function:

f(a, kek, x) = cos(a + kek / 1.000AbObA) + ln(1.000 + x · kek · (1.000ln e - 0.000))

Firstly, let's insert all constants:

f(a, kek, x) = cos(a + kek / 1.0001337.229) + ln(1.000 + x · kek · (1.000ln 2.718) - 0.000))

And simplify this expression (if possible):

f(a, kek, x) = cos(a + kek) + ln(1.000 + x · kek)

3 Exploration the expression as a function of multiple variables

- Calculation a value of function in the point

BRITISH SCIENTISTS WERE SHOCKED, WHEN THEY COUNT IT!!!  
In the point M0(a0, kek0, x0) = (3.142, 13.000, 1.000) expression's value = 1.73157

- Finding the first derivation of function

Personally, I've always thought about first derivation of something like that function... Haven't you?  
But now, by using informatics and math skills I feel that I'm prepared enough to calculate it!

1 step: Finding a derivation of kek

While preparing for exams, I learned a lot of new things, for example:  
(kek)' =  
= 1.000

2 step: Finding a derivation of x

Only after two cups of beer you might understand it:  
(x)' =  
= 1.000

3 step: Finding a derivation of x · kek

Never say it to girls:  
(x · kek)' =  
= kek + x

4 step: Finding a derivation of 1.000

Only by using special skills we might know::  
(1.000)' = ... = [top secret] = ... =  
= 0.000

5 step: Finding a derivation of 1.000 + x · kek

What if:  
(1.000 + x · kek)' =  
= kek + x

6 step: Finding a derivation of ln(1.000 + x · kek)

Even my two-aged sister knows that:  
(ln(1.000 + x · kek))' =  
= 1.000 / (1.000 + x · kek) · (kek + x)

7 step: Finding a derivation of kek

The first task in MIPT was to calculate:  
(kek)' =

$$= 1.000$$

8 step: Finding a derivation of  $a$

Never say it to girls:

$$\begin{aligned}(a)' &= \\ &= 1.000\end{aligned}$$

9 step: Finding a derivation of  $a + kek$

It's simple as fuck:

$$\begin{aligned}(a + kek)' &= \dots = \text{[top secret]} = \dots = \\ &= 2.000\end{aligned}$$

10 step: Finding a derivation of  $\cos(a + kek)$

As we know:

$$\begin{aligned}(\cos(a + kek))' &= \\ &= 2.000 \cdot (-1.000) \cdot \sin(a + kek)\end{aligned}$$

11 step: Finding a derivation of  $\cos(a + kek) + \ln(1.000 + x \cdot kek)$

I was asked not to tell anyone that:

$$\begin{aligned}(\cos(a + kek) + \ln(1.000 + x \cdot kek))' &= \\ &= 2.000 \cdot (-1.000) \cdot \sin(a + kek) + \frac{1.000}{1.000 + x \cdot kek} \cdot (kek + x)\end{aligned}$$

Congratulations! **The first derivation of the expression** is:

$$\begin{aligned}f'(a, kek, x) &= 2.000 \cdot (-1.000) \cdot \sin(a + kek) + \frac{1.000}{1.000 + x \cdot kek} \cdot (kek + x) \\ \text{In the point } M_0(a_0, kek_0, x_0) &= (3.142, 13.000, 1.000) \text{ it's value} = 1.84017\end{aligned}$$

**Finding the 2 derivation** Let's find **the 1 derivation** of the expression:

1 step: Finding a derivation of  $kek$

Only after two cups of beer you might understand it:

$$\begin{aligned}(kek)' &= \dots = \text{[top secret]} = \dots = \\ &= 1.000\end{aligned}$$

2 step: Finding a derivation of  $x$

Even my two-aged sister knows that:

$$\begin{aligned}(x)' &= \\ &= 1.000\end{aligned}$$

3 step: Finding a derivation of  $x \cdot kek$

Even my two-aged sister knows that:

$$\begin{aligned}(x \cdot kek)' &= \\ &= kek + x\end{aligned}$$

4 step: Finding a derivation of 1.000

When I was a child, my father always told me: "Remember, son:

$$\begin{aligned}(1.000)' &= \\ &= 0.000\end{aligned}$$

5 step: Finding a derivation of  $1.000 + x \cdot kek$

I have no words to describe this fact:

$$\begin{aligned}(1.000 + x \cdot kek)' &= \dots = \text{[top secret]} = \dots = \\ &= kek + x\end{aligned}$$

6 step: Finding a derivation of  $\ln(1.000 + x \cdot kek)$

My roommate mumbled it in his sleep all night:

$$\begin{aligned}(\ln(1.000 + x \cdot kek))' &= \dots = \text{[top secret]} = \dots = \\ &= \frac{1.000}{1.000 + x \cdot kek} \cdot (kek + x)\end{aligned}$$

7 step: Finding a derivation of  $kek$

I have no words to describe this fact:

$$\begin{aligned}(kek)' &= \dots = \text{[top secret]} = \dots = \\ &= 1.000\end{aligned}$$

8 step: Finding a derivation of  $a$

While preparing for exams, I learned a lot of new things, for example:

$$\begin{aligned}(a)' &= \\ &= 1.000\end{aligned}$$

9 step: Finding a derivation of  $a + kek$

It's really easy to find:

$$\begin{aligned}(a + kek)' &= \\ &= 2.000\end{aligned}$$

10 step: Finding a derivation of  $\cos(a + kek)$

What if:

$$(\cos(a + kek))' = \dots = \text{[top secret]} = \dots =$$

$$= 2.000 \cdot (-1.000) \cdot \sin(a + kek)$$

$$\text{11 step: Finding a derivation of } \cos(a + kek) + \ln(1.000 + x \cdot kek)$$

You should be aware of the fact that:

$$\begin{aligned} (\cos(a + kek) + \ln(1.000 + x \cdot kek))' &= \\ &= 2.000 \cdot (-1.000) \cdot \sin(a + kek) + \frac{1.000}{1.000 + x \cdot kek} \cdot (kek + x) \end{aligned}$$

So the 1 derivation of the expression is:

$$2.000 \cdot (-1.000) \cdot \sin(a + kek) + \frac{1.000}{1.000 + x \cdot kek} \cdot (kek + x)$$

Let’s find **the 2 derivation** of the expression:

$$\text{1 step: Finding a derivation of } x$$

A true prince must know that:

$$\begin{aligned} (x)' &= \\ &= 1.000 \end{aligned}$$

$$\text{2 step: Finding a derivation of } kek$$

For centuries, people have hunted for the secret knowledge that:

$$\begin{aligned} (kek)' &= \\ &= 1.000 \end{aligned}$$

$$\text{3 step: Finding a derivation of } kek + x$$

I spend the hole of my life to find the answer and finally it’s:

$$\begin{aligned} (kek + x)' &= \dots = \text{[top secret]} = \dots = \\ &= 2.000 \end{aligned}$$

$$\text{4 step: Finding a derivation of } kek$$

Never say it to girls:

$$\begin{aligned} (kek)' &= \\ &= 1.000 \end{aligned}$$

$$\text{5 step: Finding a derivation of } x$$

It’s really easy to find:

$$\begin{aligned} (x)' &= \\ &= 1.000 \end{aligned}$$

$$\text{6 step: Finding a derivation of } x \cdot kek$$

Sometimes I hear the same voice in my head, it always says:

$$\begin{aligned} (x \cdot kek)' &= \dots = \text{[top secret]} = \dots = \\ &= kek + x \end{aligned}$$

$$\text{7 step: Finding a derivation of } 1.000$$

Even my two-aged sister knows that:

$$\begin{aligned} (1.000)' &= \\ &= 0.000 \end{aligned}$$

$$\text{8 step: Finding a derivation of } 1.000 + x \cdot kek$$

Only by using special skills we might know::

$$\begin{aligned} (1.000 + x \cdot kek)' &= \\ &= kek + x \end{aligned}$$

$$\text{9 step: Finding a derivation of } 1.000$$

My friends always beat me, because I didn’t know that:

$$\begin{aligned} (1.000)' &= \dots = \text{[top secret]} = \dots = \\ &= 0.000 \end{aligned}$$

$$\text{10 step: Finding a derivation of } \frac{1.000}{1.000 + x \cdot kek}$$

A true prince must know that:

$$\begin{aligned} \left(\frac{1.000}{1.000 + x \cdot kek}\right)' &= \dots = \text{[top secret]} = \dots = \\ &= \frac{(-1.000) \cdot (kek + x)}{(1.000 + x \cdot kek)^{2.000}} \end{aligned}$$

$$\text{11 step: Finding a derivation of } \frac{1.000}{1.000 + x \cdot kek} \cdot (kek + x)$$

Sometimes I hear the same voice in my head, it always says:

$$\begin{aligned} \left(\frac{1.000}{1.000 + x \cdot kek} \cdot (kek + x)\right)' &= \\ &= \frac{(-1.000) \cdot (kek + x)}{(1.000 + x \cdot kek)^{2.000}} \cdot (kek + x) + 2.000 \cdot \frac{1.000}{1.000 + x \cdot kek} \end{aligned}$$

$$\text{12 step: Finding a derivation of } kek$$

Only by using special skills we might know::

$$\begin{aligned} (kek)' &= \dots = \text{[top secret]} = \dots = \\ &= 1.000 \end{aligned}$$

$$\text{13 step: Finding a derivation of } a$$

While preparing for exams, I learned a lot of new things, for example:

$$(a)' =$$

$$= 1.000$$

14 step: Finding a derivation of  $a + kek$

She: please, never speak with my dad about math... Me: ok) Also me after homework of matan:  
 $(a + kek)' = \dots = \text{[top secret]} = \dots =$   
 $= 2.000$

15 step: Finding a derivation of  $\sin(a + kek)$

My roommate mumbled it in his sleep all night:  
 $(\sin(a + kek))' = \dots = \text{[top secret]} = \dots =$   
 $= 2.000 \cdot \cos(a + kek)$

16 step: Finding a derivation of  $-1.000$

A true prince must know that:  
 $(-1.000)' = \dots = \text{[top secret]} = \dots =$   
 $= 0.000$

17 step: Finding a derivation of  $(-1.000) \cdot \sin(a + kek)$

A true prince must know that:  
 $((-1.000) \cdot \sin(a + kek))' =$   
 $= (-1.000) \cdot 2.000 \cdot \cos(a + kek)$

18 step: Finding a derivation of  $2.000$

If someone asked me that in the middle of the night, I wouldn't hesitate to say:  
 $(2.000)' =$   
 $= 0.000$

19 step: Finding a derivation of  $2.000 \cdot (-1.000) \cdot \sin(a + kek)$

When I was a child, my father always told me: "Remember, son:  
 $(2.000 \cdot (-1.000) \cdot \sin(a + kek))' = \dots = \text{[top secret]} = \dots =$   
 $= 2.000 \cdot (-1.000) \cdot 2.000 \cdot \cos(a + kek)$

20 step: Finding a derivation of  $2.000 \cdot (-1.000) \cdot \sin(a + kek) + \frac{1.000}{1.000+x \cdot kek} \cdot (kek + x)$

thanks to the results of my colleagues' scientific work, I know that:  
 $(2.000 \cdot (-1.000) \cdot \sin(a + kek) + \frac{1.000}{1.000+x \cdot kek} \cdot (kek + x))' =$   
 $= 2.000 \cdot (-1.000) \cdot 2.000 \cdot \cos(a + kek) + \frac{(-1.000) \cdot (kek+x)}{(1.000+x \cdot kek)^{2.000}} \cdot (kek + x) + 2.000 \cdot \frac{1.000}{1.000+x \cdot kek}$

So the 2 derivation of the expression is:  
 $2.000 \cdot (-1.000) \cdot 2.000 \cdot \cos(a + kek) + \frac{(-1.000) \cdot (kek+x)}{(1.000+x \cdot kek)^{2.000}} \cdot (kek + x) + 2.000 \cdot \frac{1.000}{1.000+x \cdot kek}$

**Finally... The 2 derivation of the expression:**  
 $f^{(2)}(a, kek, x) = 2.000 \cdot (-1.000) \cdot 2.000 \cdot \cos(a + kek) + \frac{(-1.000) \cdot (kek+x)}{(1.000+x \cdot kek)^{2.000}} \cdot (kek + x) + 2.000 \cdot \frac{1.000}{1.000+x \cdot kek}$   
 BRITISH SCIENTISTS WERE SHOCKED AGAIN, WHEN THEY COUNT THE 2 DERIVATION OF THIS EXPRESSION!!!  
 In the point  $M_0(a_0, kek_0, x_0) = (3.142, 13.000, 1.000)$  it's value = 2.77280

**Finding partical derivations** Partial derivation of the expression on the variable a:

$\frac{\partial f}{\partial a} = (-1.000) \cdot \sin(a + 13.000)$   
 In the point  $M_0(a_0, kek_0, x_0) = (3.142, 13.000, 1.000)$  it's value = 0.42008 !!!  
 Partial derivation of the expression on the variable kek:  
 $\frac{\partial f}{\partial kek} = (-1.000) \cdot \sin(3.142 + kek) + \frac{1.000}{1.000+kek}$   
 In the point  $M_0(a_0, kek_0, x_0) = (3.142, 13.000, 1.000)$  it's value = 0.49151 !!!  
 Partial derivation of the expression on the variable x:  
 $\frac{\partial f}{\partial x} = 13.000 \cdot \frac{1.000}{1.000+13.000 \cdot x}$   
 In the point  $M_0(a_0, kek_0, x_0) = (3.142, 13.000, 1.000)$  it's value = 0.92857 !!!

**Finding full derivation** Full derivation:


$\sqrt{((-1.000) \cdot \sin(a + 13.000))^{2.000} + ((-1.000) \cdot \sin(3.142 + kek) + \frac{1.000}{1.000+kek})^{2.000} + (13.000 \cdot \frac{1.000}{1.000+13.000 \cdot x})^{2.000}}$   
 In the point  $M_0(a_0, kek_0, x_0) = (3.142, 13.000, 1.000)$  it's value = 1.13150 !!!

## 4 Exploration the expression as a function of the first variable

Now let's consider the expression as a function of the first variable a:  $f(a) = \cos(a + 13.000) + 2.639$

**Decomposing on Macloren's formula** Maklorems formula for  $a \rightarrow a_0 = 3.142$ :  
 $f(a) = 1.732 + 0.420 \cdot (a - 3.142) + 0.454 \cdot (a - 3.142)^{2.000} + (-0.070) \cdot (a - 3.142)^{3.000} + o((a - 3.142)^{3.000})$

**Graphics** Graph  $f(a) = \cos(a + 13.000) + 2.639$  on the diapasone  $a \in [-1 : 15]$  :




**Equations in the point**    **Tangent equation** in the point  $a_0 = 0.200$ :

$$f(a) = (-0.592) \cdot (a - 0.200) + 3.445$$

**Normal equation** in the point  $a_0 = 0.200$ :

$$f(a) = 1.689 \cdot (a - 0.200) + 3.445$$

Their graphs in  $\delta = 2.50000$  coverage of the point  $a_0 = 0.200000$



## 5 Conclusion

*Ultrarcringeconclusionhere :*