

1 Introduction

CrInGeCrInGeProduction.Supercringeintroductionhere :

2 Some basic knowledge about researching problem...

Parameters and constants we use in this work:

Constants (3):
e = 2.718282
pi = 3.141593
AbObA = 1337.228690

Variables (3):
a = 3.141500
kek = 13.000000
x = 1.000000

Parameters of exploration :
Number of differentiates = 2
Macloren's accuracy = 3
Tanget point = 0.200000
Delta coverage of tangent point = 2.500000
Graph diapasone = [-1 : 15]

So let's calculate smth with a given function:

f(a, kek, x) = cos(a + kek / 1.000AbObA) + ln(1.000 + x · kek · (1.000^(ln e) - 0.000))

Firstly, let's insert all constants:

f(a, kek, x) = cos(a + kek / 1.0001337.229) + ln(1.000 + x · kek · (1.000^(ln 2.718) - 0.000))

And simplify this expression (if possible):

f(a, kek, x) = cos(a + kek) + ln(1.000 + x · kek)

3 Exploration the expression as a function of multiple variables

- Calculation a value of function in the point

BRITISH SCIENTISTS WERE SHOCKED, WHEN THEY COUNT IT!!!
In the point M0(a0, kek0, x0) = (3.142, 13.000, 1.000) expression's value = 1.73157

- Finding the first derivation of function

Personally, I've always thought about first derivation of something like that function... Haven't you?
But now, by using informatics and math skills I feel that I'm prepared enough to calculate it!

1 step: Finding a derivation of kekWhile preparing for exams, I learned a lot of new things, for example: (kek)' =

= 1.000

2 step: Finding a derivation of xOnly after two cups of beer you might understand it: (x)' =

= 1.000

3 step: Finding a derivation of x · kekNever say it to girls: (x · kek)' =

= kek + x

4 step: Finding a derivation of 1.000Only by using special skills we might know:: (1.000)' = ... = [top secret] = ... =

= 0.000

5 step: Finding a derivation of 1.000 + x · kekWhat if: (1.000 + x · kek)' =

= kek + x

6 step: Finding a derivation of ln(1.000 + x · kek)Even my two-aged sister knows that: (ln(1.000 + x · kek))' =

= 1.000 / (1.000 + x · kek) · (kek + x)

7 step: Finding a derivation of kekThe first task in MIPT was to calculate: (kek)' =

= 1.000

8 step: Finding a derivation of a Never say it to girls: $(a)' =$

$= 1.000$

9 step: Finding a derivation of $a + kek$ It's simple as fuck: $(a + kek)' = \dots = \text{[top secret]} = \dots =$

$= 2.000$

10 step: Finding a derivation of $\cos(a + kek)$ As we know: $(\cos(a + kek))' =$

$= 2.000 \cdot (-1.000) \cdot \sin(a + kek)$

11 step: Finding a derivation of $\cos(a + kek) + \ln(1.000 + x \cdot kek)$ I was asked not to tell anyone that: $(\cos(a + kek) + \ln(1.000 + x \cdot kek))' =$

$= 2.000 \cdot (-1.000) \cdot \sin(a + kek) + \frac{1.000}{1.000 + x \cdot kek} \cdot (kek + x)$

Congratulations! **The first derivation of the expression** is:
 $f'(a, kek, x) = 2.000 \cdot (-1.000) \cdot \sin(a + kek) + \frac{1.000}{1.000 + x \cdot kek} \cdot (kek + x)$
In the point $M_0(a_0, kek_0, x_0) = (3.142, 13.000, 1.000)$ it's value = 1.84017

Finding the 2 derivation Let's find **the 1 derivation** of the expression:

1 step: Finding a derivation of kek Only after two cups of beer you might understand it: $(kek)' = \dots = \text{[top secret]} = \dots =$

$= 1.000$

2 step: Finding a derivation of x Even my two-aged sister knows that: $(x)' =$

$= 1.000$

3 step: Finding a derivation of $x \cdot kek$ Even my two-aged sister knows that: $(x \cdot kek)' =$

$= kek + x$

4 step: Finding a derivation of 1.000 When I was a child, my father always told me: "Remember, son: $(1.000)' =$

$= 0.000$

5 step: Finding a derivation of $1.000 + x \cdot kek$ I have no words to describe this fact: $(1.000 + x \cdot kek)' = \dots = \text{[top secret]} = \dots =$

$= kek + x$

6 step: Finding a derivation of $\ln(1.000 + x \cdot kek)$ My roommate mumbled it in his sleep all night: $(\ln(1.000 + x \cdot kek))' = \dots = \text{[top$

$\text{secret]} = \dots =$
 $= \frac{1.000}{1.000 + x \cdot kek} \cdot (kek + x)$

7 step: Finding a derivation of kek I have no words to describe this fact: $(kek)' = \dots = \text{[top secret]} = \dots =$

$= 1.000$

8 step: Finding a derivation of a While preparing for exams, I learned a lot of new things, for example: $(a)' =$

$= 1.000$

9 step: Finding a derivation of $a + kek$ It's really easy to find: $(a + kek)' =$

$= 2.000$

10 step: Finding a derivation of $\cos(a + kek)$ What if: $(\cos(a + kek))' = \dots = \text{[top secret]} = \dots =$

$$= 2.000 \cdot (-1.000) \cdot \sin(a + kek)$$

$$\text{11 step: Finding a derivation of } \cos(a + kek) + \ln(1.000 + x \cdot kek) \text{You should be aware of the fact that: } (\cos(a + kek) + \ln(1.000 + x \cdot kek))'$$

$$=$$

$$= 2.000 \cdot (-1.000) \cdot \sin(a + kek) + \frac{1.000}{1.000+x \cdot kek} \cdot (kek + x)$$

$$\text{So the 1 derivation of the expression is:} \\ 2.000 \cdot (-1.000) \cdot \sin(a + kek) + \frac{1.000}{1.000+x \cdot kek} \cdot (kek + x)$$

$$\text{Let's find \textbf{the 2 derivation} of the expression:}$$

$$\text{1 step: Finding a derivation of } x \text{A true prince must know that: } (x)' =$$

$$= 1.000$$

$$\text{2 step: Finding a derivation of } kek \text{For centuries, people have hunted for the secret knowledge that: } (kek)' =$$

$$= 1.000$$

$$\text{3 step: Finding a derivation of } kek + x \text{I spend the hole of my life to find the answer and finally it's: } (kek + x)' = \dots = \text{[top secret]}$$

$$= \dots =$$

$$= 2.000$$

$$\text{4 step: Finding a derivation of } kek \text{Never say it to girls: } (kek)' =$$

$$= 1.000$$

$$\text{5 step: Finding a derivation of } x \text{It's really easy to find: } (x)' =$$

$$= 1.000$$

$$\text{6 step: Finding a derivation of } x \cdot kek \text{Sometimes I hear the same voice in my head, it always says: } (x \cdot kek)' = \dots = \text{[top secret]} =$$

$$\dots =$$

$$= kek + x$$

$$\text{7 step: Finding a derivation of } 1.000 \text{Even my two-aged sister knows that: } (1.000)' =$$

$$= 0.000$$

$$\text{8 step: Finding a derivation of } 1.000 + x \cdot kek \text{Only by using special skills we might know:: } (1.000 + x \cdot kek)' =$$

$$= kek + x$$

$$\text{9 step: Finding a derivation of } 1.000 \text{My friends always beat me, because I didn't know that: } (1.000)' = \dots = \text{[top secret]} = \dots =$$

$$= 0.000$$

$$\text{10 step: Finding a derivation of } \frac{1.000}{1.000+x \cdot kek} \text{A true prince must know that: } (\frac{1.000}{1.000+x \cdot kek})' = \dots = \text{[top secret]} = \dots =$$

$$= \frac{(-1.000) \cdot (kek+x)}{(1.000+x \cdot kek)^{2.000}}$$

$$\text{11 step: Finding a derivation of } \frac{1.000}{1.000+x \cdot kek} \cdot (kek + x) \text{Sometimes I hear the same voice in my head, it always says: } (\frac{1.000}{1.000+x \cdot kek} \cdot (kek + x))'$$

$$=$$

$$= \frac{(-1.000) \cdot (kek+x)}{(1.000+x \cdot kek)^{2.000}} \cdot (kek + x) + 2.000 \cdot \frac{1.000}{1.000+x \cdot kek}$$

$$\text{12 step: Finding a derivation of } kek \text{Only by using special skills we might know:: } (kek)' = \dots = \text{[top secret]} = \dots =$$

$$= 1.000$$

$$\text{13 step: Finding a derivation of } a \text{While preparing for exams, I learned a lot of new things, for example: } (a)' =$$

$$= 1.000$$

14 step: Finding a derivation of $a + kek$ She: please, never speak with my dad about math... Me: ok) Also me after homework of matan:

$$(a + kek)' = \dots = \text{[top secret]} = \dots = \\ = 2.000$$

$$\textbf{15 step:}$$
 Finding a derivation of $\sin(a + kek)$ My roommate mumbled it in his sleep all night: $(\sin(a + kek))' = \dots = \text{[top secret]} = \dots = \\ \dots = \\ = 2.000 \cdot \cos(a + kek)$

$$\textbf{16 step:}$$
 Finding a derivation of -1.000 A true prince must know that: $(-1.000)' = \dots = \text{[top secret]} = \dots = \\ = 0.000$

$$\textbf{17 step:}$$
 Finding a derivation of $(-1.000) \cdot \sin(a + kek)$ A true prince must know that: $((-1.000) \cdot \sin(a + kek))' = \\ = (-1.000) \cdot 2.000 \cdot \cos(a + kek)$

$$\textbf{18 step:}$$
 Finding a derivation of 2.000 If someone asked me that in the middle of the night, I wouldn't hesitate to say: $(2.000)' = \\ = 0.000$

$$\textbf{19 step:}$$
 Finding a derivation of $2.000 \cdot (-1.000) \cdot \sin(a + kek)$ When I was a child, my father always told me: "Remember, son: $(2.000 \cdot (-1.000) \cdot \sin(a + kek))' = \dots = \text{[top secret]} = \dots = \\ = 2.000 \cdot (-1.000) \cdot 2.000 \cdot \cos(a + kek)$

$$\textbf{20 step:}$$
 Finding a derivation of $2.000 \cdot (-1.000) \cdot \sin(a + kek) + \frac{1.000}{1.000 + x \cdot kek} \cdot (kek + x)$ thanks to the results of my colleagues' scientific work, I know that: $(2.000 \cdot (-1.000) \cdot \sin(a + kek) + \frac{1.000}{1.000 + x \cdot kek} \cdot (kek + x))' =$

$$= 2.000 \cdot (-1.000) \cdot 2.000 \cdot \cos(a + kek) + \frac{(-1.000) \cdot (kek + x)}{(1.000 + x \cdot kek)^{2.000}} \cdot (kek + x) + 2.000 \cdot \frac{1.000}{1.000 + x \cdot kek}$$

So the 2 derivation of the expression is:

$$2.000 \cdot (-1.000) \cdot 2.000 \cdot \cos(a + kek) + \frac{(-1.000) \cdot (kek + x)}{(1.000 + x \cdot kek)^{2.000}} \cdot (kek + x) + 2.000 \cdot \frac{1.000}{1.000 + x \cdot kek}$$

Finally... The 2 derivation of the expression:

$$f^{(2)}(a, kek, x) = 2.000 \cdot (-1.000) \cdot 2.000 \cdot \cos(a + kek) + \frac{(-1.000) \cdot (kek + x)}{(1.000 + x \cdot kek)^{2.000}} \cdot (kek + x) + 2.000 \cdot \frac{1.000}{1.000 + x \cdot kek}$$

BRITISH SCIENTISTS WERE SHOCKED AGAIN, WHEN THEY COUNT THE 2 DERIVATION OF THIS EXPRESSION!!!

In the point $M_0(a_0, kek_0, x_0) = (3.142, 13.000, 1.000)$ it's value = 2.77280

Finding partial derivations Partial derivation of the expression on the variable a:

$$\frac{\partial f}{\partial a} = (-1.000) \cdot \sin(a + 13.000)$$

In the point $M_0(a_0, kek_0, x_0) = (3.142, 13.000, 1.000)$ it's value = 0.42008 !!!

Partial derivation of the expression on the variable kek:

$$\frac{\partial f}{\partial kek} = (-1.000) \cdot \sin(3.142 + kek) + \frac{1.000}{1.000 + kek}$$

In the point $M_0(a_0, kek_0, x_0) = (3.142, 13.000, 1.000)$ it's value = 0.49151 !!!

Partial derivation of the expression on the variable x:

$$\frac{\partial f}{\partial x} = 13.000 \cdot \frac{1.000}{1.000 + 13.000 \cdot x}$$

In the point $M_0(a_0, kek_0, x_0) = (3.142, 13.000, 1.000)$ it's value = 0.92857 !!!

Finding full derivation Full derivation:

$$\sqrt{((-1.000) \cdot \sin(a + 13.000))^{2.000} + ((-1.000) \cdot \sin(3.142 + kek) + \frac{1.000}{1.000 + kek})^{2.000} + (13.000 \cdot \frac{1.000}{1.000 + 13.000 \cdot x})^{2.000}}$$

In the point $M_0(a_0, kek_0, x_0) = (3.142, 13.000, 1.000)$ it's value = 1.13150 !!!


4 Exploration the expression as a function of the first variable

Now let's consider the expression as a function of the first variable a: $f(a) = \cos(a + 13.000) + 2.639$

Decomposing on Macloren's formula Maklorems formula for $a \rightarrow a_0 = 3.142$:

$$f(a) = 1.732 + 0.420 \cdot (a - 3.142) + 0.454 \cdot (a - 3.142)^{2.000} + (-0.070) \cdot (a - 3.142)^{3.000} + o((a - 3.142)^{3.000})$$

Graphics Graph $f(a) = \cos(a + 13.000) + 2.639$ on the diapasone $a \in [-1 : 15]$:




Equations in the point **Tangent equation** in the point $a_0 = 0.200$:

$$f(a) = (-0.592) \cdot (a - 0.200) + 3.445$$

Normal equation in the point $a_0 = 0.200$:

$$f(a) = 1.689 \cdot (a - 0.200) + 3.445$$

Their graphs in $\delta = 2.50000$ coverage of the point $a_0 = 0.200000$



5 Conclusion

Ultrarcringeconclusionhere :