

1 Introduction

CrInGeCrInGeProduction.Supercringeintroductionhere :

2 Some basic knowledge about researching problem...

Parameters and constants we use in this work:

Constants (3):  
e = 2.718282  
pi = 3.141593  
AbObA = 1337.228690

Variables (3):  
a = 3.141500  
kek = 13.000000  
x = 1.000000

Parameters of exploration :  
Number of differentiates = 2  
Macloren's accuracy = 3  
Tanget point = 0.200000  
Delta coverage of tangent point = 2.500000  
Graph diapasone = [-1 : 15]

So let's calculate smth with a given function:

f(a, kek, x) = cos(a + kek / 1.000AbObA) + ln(1.000 + x · kek · (1.000ln e - 0.000))

Firstly, let's insert all constants:

f(a, kek, x) = cos(a + kek / 1.0001337.229) + ln(1.000 + x · kek · (1.000ln 2.718) - 0.000))

And simplify this expression (if possible):

f(a, kek, x) = cos(a + kek) + ln(1.000 + x · kek)

3 Exploration the expression as a function of multiple variables

- Calculation a value of function in the point

BRITISH SCIENTISTS WERE SHOCKED, WHEN THEY COUNT IT!!!  
In the point M0(a0, kek0, x0) = (3.142, 13.000, 1.000) expression's value = 1.73157

- Finding the first derivation of function

Personally, I've always thought about first derivation of something like that function... Haven't you?  
But now, by using informatics and math skills I feel that I'm prepared enough to calculate it!

1 step: Finding a derivation of kek  
While preparing for exams, I learned a lot of new things, for example:

(kek)' =

= 1.000

2 step: Finding a derivation of x  
Only after two cups of beer you might understand it:

(x)' =

= 1.000

3 step: Finding a derivation of x · kek  
Never say it to girls:

(x · kek)' =

= kek + x

4 step: Finding a derivation of 1.000  
Only by using special skills we might know::

(1.000)' = ... = [top secret] = ... =

$$= 0.000$$

**5 step:** Finding a derivation of  $1.000 + x \cdot kek$   
 What if:

$$(1.000 + x \cdot kek)' =$$

$$= kek + x$$

**6 step:** Finding a derivation of  $\ln(1.000 + x \cdot kek)$   
 Even my two-aged sister knows that:

$$(\ln(1.000 + x \cdot kek))' =$$

$$= \frac{1.000}{1.000+x \cdot kek} \cdot (kek + x)$$

**7 step:** Finding a derivation of  $kek$   
 The first task in MIPT was to calculate:

$$(kek)' =$$

$$= 1.000$$

**8 step:** Finding a derivation of  $a$   
 Never say it to girls:

$$(a)' =$$

$$= 1.000$$

**9 step:** Finding a derivation of  $a + kek$   
 It's simple as fuck:

$$(a + kek)' = \dots = \text{[top secret]} = \dots =$$

$$= 2.000$$

**10 step:** Finding a derivation of  $\cos(a + kek)$   
 As we know:

$$(\cos(a + kek))' =$$

$$= 2.000 \cdot (-1.000) \cdot \sin(a + kek)$$

**11 step:** Finding a derivation of  $\cos(a + kek) + \ln(1.000 + x \cdot kek)$   
 I was asked not to tell anyone that:

$$(\cos(a + kek) + \ln(1.000 + x \cdot kek))' =$$

$$= 2.000 \cdot (-1.000) \cdot \sin(a + kek) + \frac{1.000}{1.000+x \cdot kek} \cdot (kek + x)$$

Congratulations! **The first derivation of the expression** is:  
 $f'(a, kek, x) = 2.000 \cdot (-1.000) \cdot \sin(a + kek) + \frac{1.000}{1.000+x \cdot kek} \cdot (kek + x)$   
 In the point  $M_0(a_0, kek_0, x_0) = (3.142, 13.000, 1.000)$  it's value = 1.84017

**Finding the 2 derivation** Let's find **the 1 derivation** of the expression:

**1 step:** Finding a derivation of  $kek$   
 Only after two cups of beer you might understand it:

$$(kek)' = \dots = \text{[top secret]} = \dots =$$

$$= 1.000$$

**2 step:** Finding a derivation of  $x$   
 Even my two-aged sister knows that:

$$(x)' =$$

$$= 1.000$$

**3 step:** Finding a derivation of  $x \cdot kek$   
 Even my two-aged sister knows that:

$$(x \cdot kek)' =$$

$$= kek + x$$

**4 step:** Finding a derivation of 1.000  
 When I was a child, my father always told me: "Remember, son:

$$(1.000)' =$$

$$= 0.000$$

**5 step:** Finding a derivation of  $1.000 + x \cdot kek$   
 I have no words to describe this fact:

$$(1.000 + x \cdot kek)' = \dots = \text{[top secret]} = \dots =$$

$$= kek + x$$

**6 step:** Finding a derivation of  $\ln(1.000 + x \cdot kek)$   
 My roommate mumbled it in his sleep all night:

$$(\ln(1.000 + x \cdot kek))' = \dots = \text{[top secret]} = \dots =$$

$$= \frac{1.000}{1.000+x \cdot kek} \cdot (kek + x)$$

**7 step:** Finding a derivation of  $kek$   
 I have no words to describe this fact:

$$(kek)' = \dots = \text{[top secret]} = \dots =$$

$$= 1.000$$

**8 step:** Finding a derivation of  $a$   
 While preparing for exams, I learned a lot of new things, for example:

$$(a)' =$$

$$= 1.000$$

**9 step:** Finding a derivation of  $a + kek$   
 It's really easy to find:

$$(a + kek)' =$$

$$= 2.000$$

**10 step:** Finding a derivation of  $\cos(a + kek)$   
 What if:

$$(\cos(a+kek))' = \dots = \text{[top secret]} = \dots =$$

$$= 2.000 \cdot (-1.000) \cdot \sin(a+kek)$$

**11 step:** Finding a derivation of  $\cos(a+kek) + \ln(1.000+x \cdot kek)$   
 You should be aware of the fact that:

$$(\cos(a+kek) + \ln(1.000+x \cdot kek))' =$$

$$= 2.000 \cdot (-1.000) \cdot \sin(a+kek) + \frac{1.000}{1.000+x \cdot kek} \cdot (kek+x)$$

So the 1 derivation of the expression is:  
 $2.000 \cdot (-1.000) \cdot \sin(a+kek) + \frac{1.000}{1.000+x \cdot kek} \cdot (kek+x)$

Let’s find **the 2 derivation** of the expression:

**1 step:** Finding a derivation of  $x$   
 A true prince must know that:

$$(x)' =$$

$$= 1.000$$

**2 step:** Finding a derivation of  $kek$   
 For centuries, people have hunted for the secret knowledge that:

$$(kek)' =$$

$$= 1.000$$

**3 step:** Finding a derivation of  $kek+x$   
 I spend the hole of my life to find the answer and finally it’s:

$$(kek+x)' = \dots = \text{[top secret]} = \dots =$$

$$= 2.000$$

**4 step:** Finding a derivation of  $kek$   
 Never say it to girls:

$$(kek)' =$$

$$= 1.000$$

**5 step:** Finding a derivation of  $x$   
 It’s really easy to find:

$$(x)' =$$

$$= 1.000$$

**6 step:** Finding a derivation of  $x \cdot kek$   
 Sometimes I hear the same voice in my head, it always says:

$$(x \cdot kek)' = \dots = \text{[top secret]} = \dots =$$

$$= kek+x$$

**7 step:** Finding a derivation of 1.000

Even my two-aged sister knows that:

$(1.000)' =$

$= 0.000$

**8 step:** Finding a derivation of  $1.000 + x \cdot kek$   
Only by using special skills we might know::

$(1.000 + x \cdot kek)' =$

$= kek + x$

**9 step:** Finding a derivation of 1.000  
My friends always beat me, because I didn't know that:

$(1.000)' = \dots = \text{[top secret]} = \dots =$

$= 0.000$

**10 step:** Finding a derivation of  $\frac{1.000}{1.000+x \cdot kek}$   
A true prince must know that:

$(\frac{1.000}{1.000+x \cdot kek})' = \dots = \text{[top secret]} = \dots =$

$= \frac{(-1.000) \cdot (kek+x)}{(1.000+x \cdot kek)^{2.000}}$

**11 step:** Finding a derivation of  $\frac{1.000}{1.000+x \cdot kek} \cdot (kek + x)$   
Sometimes I hear the same voice in my head, it always says:

$(\frac{1.000}{1.000+x \cdot kek} \cdot (kek + x))' =$

$= \frac{(-1.000) \cdot (kek+x)}{(1.000+x \cdot kek)^{2.000}} \cdot (kek + x) + 2.000 \cdot \frac{1.000}{1.000+x \cdot kek}$

**12 step:** Finding a derivation of  $kek$   
Only by using special skills we might know::

$(kek)' = \dots = \text{[top secret]} = \dots =$

$= 1.000$

**13 step:** Finding a derivation of  $a$   
While preparing for exams, I learned a lot of new things, for example:

$(a)' =$

$= 1.000$

**14 step:** Finding a derivation of  $a + kek$   
She: please, never speak with my dad about math... Me: ok) Also me after homework of matan:

$(a + kek)' = \dots = \text{[top secret]} = \dots =$

$= 2.000$

**15 step:** Finding a derivation of  $\sin(a + kek)$   
My roommate mumbled it in his sleep all night:

$(\sin(a + kek))' = \dots = \text{[top secret]} = \dots =$

$$= 2.000 \cdot \cos(a + kek)$$

16 **step:** Finding a derivation of  $-1.000$   
A true prince must know that:

$$(-1.000)' = \textcolor{red}{...} = \textcolor{red}{[\text{top secret}]} = \textcolor{red}{...} =$$

$$= 0.000$$

17 **step:** Finding a derivation of  $(-1.000) \cdot \sin(a + kek)$   
A true prince must know that:

$$((-1.000) \cdot \sin(a + kek))' =$$

$$= (-1.000)' \cdot 2.000 \cdot \cos(a + kek)$$

18 **step:** Finding a derivation of  $2.000$   
If someone asked me that in the middle of the night, I wouldn't hesitate to say:

$$(2.000)' =$$

$$= 0.000$$

19 **step:** Finding a derivation of  $2.000 \cdot (-1.000) \cdot \sin(a + kek)$   
When I was a child, my father always told me: "Remember, son:

$$(2.000 \cdot (-1.000) \cdot \sin(a + kek))' = \textcolor{red}{...} = \textcolor{red}{[\text{top secret}]} = \textcolor{red}{...} =$$

$$= 2.000 \cdot (-1.000)' \cdot 2.000 \cdot \cos(a + kek)$$

20 **step:** Finding a derivation of  $2.000 \cdot (-1.000) \cdot \sin(a + kek) + \frac{1.000}{1.000+x \cdot kek} \cdot (kek + x)$   
thanks to the results of my colleagues' scientific work, I know that:

$$(2.000 \cdot (-1.000) \cdot \sin(a + kek) + \frac{1.000}{1.000+x \cdot kek} \cdot (kek + x))' =$$

$$= 2.000 \cdot (-1.000)' \cdot 2.000 \cdot \cos(a + kek) + \frac{(-1.000) \cdot (kek+x)}{(1.000+x \cdot kek)^{2.000}} \cdot (kek + x) + 2.000 \cdot \frac{1.000}{1.000+x \cdot kek}$$

So the 2 derivation of the expression is:  
 $2.000 \cdot (-1.000)' \cdot 2.000 \cdot \cos(a + kek) + \frac{(-1.000) \cdot (kek+x)}{(1.000+x \cdot kek)^{2.000}} \cdot (kek + x) + 2.000 \cdot \frac{1.000}{1.000+x \cdot kek}$

**Finally... The 2 derivation of the expression:**  
 $f^{(2)}(a, kek, x) = 2.000 \cdot (-1.000)' \cdot 2.000 \cdot \cos(a + kek) + \frac{(-1.000) \cdot (kek+x)}{(1.000+x \cdot kek)^{2.000}} \cdot (kek + x) + 2.000 \cdot \frac{1.000}{1.000+x \cdot kek}$   
BRITISH SCIENTISTS WERE SHOCKED AGAIN, WHEN THEY COUNT THE 2 DERIVATION OF THIS EXPRESSION!!!  
In the point  $M_0(a_0, kek_0, x_0) = (3.142, 13.000, 1.000)$  it's value = 2.77280

**Finding partial derivations**    Partial derivation of the expression on the variable a:  
 $\frac{\partial f}{\partial a} = (-1.000)' \cdot \sin(a + 13.000)$   
In the point  $M_0(a_0, kek_0, x_0) = (3.142, 13.000, 1.000)$  it's value = 0.42008 !!!  
Partial derivation of the expression on the variable kek:  
 $\frac{\partial f}{\partial kek} = (-1.000)' \cdot \sin(3.142 + kek) + \frac{1.000}{1.000+kek}$   
In the point  $M_0(a_0, kek_0, x_0) = (3.142, 13.000, 1.000)$  it's value = 0.49151 !!!  
Partial derivation of the expression on the variable x:  
 $\frac{\partial f}{\partial x} = 13.000 \cdot \frac{1.000}{1.000+13.000 \cdot x}$   
In the point  $M_0(a_0, kek_0, x_0) = (3.142, 13.000, 1.000)$  it's value = 0.92857 !!!


**Finding full derivation**    Full derivation:  
 $\sqrt{((-1.000)' \cdot \sin(a + 13.000))^{2.000} + ((-1.000)' \cdot \sin(3.142 + kek) + \frac{1.000}{1.000+kek})^{2.000} + (13.000 \cdot \frac{1.000}{1.000+13.000 \cdot x})^{2.000}}$   
In the point  $M_0(a_0, kek_0, x_0) = (3.142, 13.000, 1.000)$  it's value = 1.13150 !!!

## 4    Exploration the expression as a function of the first variable

Now let's consider the expression as a function of the first variable a:  $f(a) = \cos(a + 13.000) + 2.639$

**Decomposing on Macloren's formula**    **Maklorems formula for  $a \rightarrow a_0 = 3.142$ :**  
 $f(a) = 1.732 + 0.420 \cdot (a - 3.142) + 0.454 \cdot (a - 3.142)^{2.000} + (-0.070) \cdot (a - 3.142)^{3.000} + o((a - 3.142)^{3.000})$

**Graphics**    **Graph**  $f(a) = \cos(a + 13.000) + 2.639$  on the diapasone  $a \in [-1 : 15]$  :




**Equations in the point**    **Tangent equation** in the point  $a_0 = 0.200$ :

$f(a) = (-0.592) \cdot (a - 0.200) + 3.445$

**Normal equation** in the point  $a_0 = 0.200$ :

$f(a) = 1.689 \cdot (a - 0.200) + 3.445$

Their graphs in  $\delta = 2.50000$  coverage of the point  $a_0 = 0.200000$



5    **Conclusion**

*Ultrarcringeconclusionhere :*