

Let’s calculate smth with expression given:  $f(x, y) =$

$$x \cdot y^{2.000}$$

Firstly, let’s insert all constants and simplify it:

$$x \cdot y^{2.000}$$

BRITISH SCIENTISTS WERE SHOCKED, WHEN THEY COUNT IT!!! IN THE POINT ( $x = 3.000$ ,  $y = 2.000$ )IT’S VALUE = 12.000 !!!

1 step: finding a derivation of function:

$$y$$

here it is:

$$1.000$$

2 step: finding a derivation of function:

$$y^{2.000}$$

here it is:

$$2.000 \cdot y$$

3 step: finding a derivation of function:

$$x$$

here it is:

$$1.000$$

4 step: finding a derivation of function:

$$x \cdot y^{2.000}$$

here it is:

$$y^{2.000} + 2.000 \cdot y \cdot x$$

Congratulations! The first derivation of the expression is:

$$y^{2.000} + 2.000 \cdot y \cdot x$$

IN THE POINT ( $x = 3.000$ ,  $y = 2.000$ )IT’S VALUE = 16.000 !!!

Let’s calculate the 3 derivation of the expression:

Calculating the 1 derivation of the expression:

1 step: finding a derivation of function:

$$y$$

here it is:

$$1.000$$

2 step: finding a derivation of function:

$$y^{2.000}$$

here it is:

$$2.000 \cdot y$$

3 step: finding a derivation of function:

$$x$$

here it is:

$$1.000$$

4 step: finding a derivation of function:

$$x \cdot y^{2.000}$$

here it is:

$$y^{2.000} + 2.000 \cdot y \cdot x$$

Calculating the 2 derivation of the expression:

1 step: finding a derivation of function:

$$x$$

here it is:

$$1.000$$

2 step: finding a derivation of function:

here it is:

3 step: finding a derivation of function:

here it is:

4 step: finding a derivation of function:

here it is:

5 step: finding a derivation of function:

here it is:

6 step: finding a derivation of function:

here it is:

7 step: finding a derivation of function:

here it is:

8 step: finding a derivation of function:

here it is:

Calculating the 3 derivation of the expression:

1 step: finding a derivation of function:

here it is:

2 step: finding a derivation of function:

here it is:

3 step: finding a derivation of function:

here it is:

4 step: finding a derivation of function:

here it is:

5 step: finding a derivation of function:

here it is:

$$y$$

$$1.000$$

$$2.000$$

$$0.000$$

$$2.000 \cdot y$$

$$2.000$$

$$2.000 \cdot y \cdot x$$

$$2.000 \cdot x + 2.000 \cdot y$$

$$y$$

$$1.000$$

$$y^{2.000}$$

$$2.000 \cdot y$$

$$y^{2.000} + 2.000 \cdot y \cdot x$$

$$2.000 \cdot y + 2.000 \cdot x + 2.000 \cdot y$$

$$y$$

$$1.000$$

$$2.000$$

$$0.000$$

$$2.000 \cdot y$$

$$2.000$$

$$x$$

$$1.000$$

$$2.000$$

$$0.000$$

6 step: finding a derivation of function:

$$2.000 \cdot x$$

here it is:

$$2.000$$

7 step: finding a derivation of function:

$$2.000 \cdot x + 2.000 \cdot y$$

here it is:

$$4.000$$

8 step: finding a derivation of function:

$$y$$

here it is:

$$1.000$$

9 step: finding a derivation of function:

$$2.000$$

here it is:

$$0.000$$

10 step: finding a derivation of function:

$$2.000 \cdot y$$

here it is:

$$2.000$$

11 step: finding a derivation of function:

$$2.000 \cdot y + 2.000 \cdot x + 2.000 \cdot y$$

here it is:

$$6.000$$

Finally... The 3 derivation of the expression:

$$6.000$$

BRITISH SCIENTISTS WERE SHOCKED, WHEN THEY COUNT THE 3 DERIVATION OF THIS EXPRESSION!!! IN THE POINT (x = 3.000, y = 2.000)IT'S VALUE = 6.000 !!!

Partial derivation of the expression on the variable 'x':

$$4.000$$

IN THE POINT (x = 3.000, y = 2.000) IT'S VALUE = 4.000000 !!!

Partial derivation of the expression on the variable 'y':

$$3.000 \cdot 2.000 \cdot y$$

IN THE POINT (x = 3.000, y = 2.000) IT'S VALUE = 12.000000 !!!

Full derivation:

$$\sqrt{16.000 + (3.000 \cdot 2.000 \cdot y)^{2.000}}$$

IN THE POINT (x = 3.000, y = 2.000)IT'S VALUE = 12.649 !!!

Let's consider the expression as a function of x variable: f(x) =

$$4.000 \cdot x$$

Maklorens formula for x near to 3.000000:

$$12.000 + 4.000 \cdot (x - 3.000)$$

And remainig member is o maloe from:

$$(x - 3.000)^{4.000}$$

Graph f(x):

Tangent equation in point -2.000: f(x) =

$$4.000 \cdot (x - (-2.000)) + (-8.000)$$

Normal equation in point -2.000: f(x) =

$$(-0.250) \cdot (x - (-2.000)) + (-8.000)$$