

Let's calculate smth with a given function: $f(x, y) = \sin x \cdot y^{2.000000000}$

Firstly, let's insert all constants and simplify this expression: $f(x, y) = \sin x \cdot y^{2.000000000}$

BRITISH SCIENTISTS WERE SHOCKED, WHEN THEY COUNT IT!!!

In the point $M_0(x_0, y_0) = (3.000000000, 2.000000000)$ **it's value** = 0.564480032

Personally, I've always thought about first derivation of something like that function... Haven't you?

But now, by using informatics and math skills I feel that I'm prepared enough to calculate it!

1 step. finding a derivation of:

y

While preparing for exams, I learned a lot of new things, for example:

$(y)' = \dots = [\text{top secret}] = \dots =$

= 1.000000000

2 step. finding a derivation of:

$y^{2.000000000}$

It's really easy to find:

$(y^{2.000000000})' = \dots = [\text{top secret}] = \dots =$

= 2.000000000 · y

3 step. finding a derivation of:

x

My roommate mumbled it in his sleep all night:

$(x)' = \dots = [\text{top secret}] = \dots =$

= 1.000000000

4 step. finding a derivation of:

$\sin x$

Sounds logical that it is the same as:

$(\sin x)' = \dots = [\text{top secret}] = \dots =$

= $\cos x$

5 step. finding a derivation of:

$\sin x \cdot y^{2.000000000}$

For centuries, people have hunted for the secret knowledge that:

$(\sin x \cdot y^{2.000000000})' = \dots = [\text{top secret}] = \dots =$

= $\cos x \cdot y^{2.000000000} + 2.000000000 \cdot y \cdot \sin x$

Congratulations! **The first derivation of the expression is:**

$\cos x \cdot y^{2.000000000} + 2.000000000 \cdot y \cdot \sin x$ In the point $M_0(x_0, y_0) = (3.000000000, 2.000000000)$ it's value = -3.395489954

Let's calculate the 0 derivation of the expression:

Finally... The 0 derivation of the expression:

$\sin x \cdot y^{2.000000000}$

BRITISH SCIENTISTS WERE SHOCKED AGAIN, WHEN THEY COUNT THE 0 DERIVATION OF THIS EXPRESSION!!!

In the point $M_0(x_0, y_0) = (3.000000000, 2.000000000)$ it's value = 0.564480032

Partial derivation of the expression on the variable x:

$\frac{\partial f}{\partial x} = 4.000000000 \cdot \cos x$

In the point $M_0(x_0, y_0) = (3.000000000, 2.000000000)$ it's value = -3.959970 !!!

Partial derivation of the expression on the variable y:

$\frac{\partial f}{\partial y} = 0.141120008 \cdot 2.000000000 \cdot y$

In the point $M_0(x_0, y_0) = (3.000000000, 2.000000000)$ it's value = 0.564480 !!!

Full derivation:

$\sqrt{(4.000000000 \cdot \cos x)^{2.000000000} + (0.141120008 \cdot 2.000000000 \cdot y)^{2.000000000}}$

In the point $M_0(x_0, y_0) = (3.000000000, 2.000000000)$ it's value = 4.000000000 !!!

Now let's consider the expression as a function of x variable: $f(x) = 4.000000000 \cdot \sin x$

Maklorens formula for $x \rightarrow x_0 = 3.000000000$:

$f(x) = 0.564480032 + (-3.959969986) \cdot (x - 3.000000000) + (-0.282240016) \cdot (x - 3.000000000)^{2.000000000} + 0.659994998 \cdot (x - 3.000000000)^{3.000000000}$

Graph $f(x) = 4.000000000 \cdot \sin x$ on the diapason $x \in [-10 : 10]$:

Tangent equation in the point $x_0 = 0.000000000$: $f(x) = 4.000000000 \cdot x$

Normal equation in the point $x_0 = 0.000000000$: $f(x) = (-0.250000000) \cdot (x - 0.000000000) + 0.000000000$