CrInGeCrInGeProduction. Supercringeint roduction here:

Let's calculate smth with expression given: f(x, y) = $x \cdot y^{2.000}$

Firstly, let's insert all constants and simplify it: $f(x, y) = x \cdot y^{2.000}$

BRITISH SCIENTISTS WERE SHOCKED, WHEN THEY COUNT IT!!!

In the point (x = 3.000, y = 2.000)it's value = 12.000

1 step: finding a derivation: y

If someone asked me that in the middle of the night, I wouldn't hesitate to say:

1.000

2 step: finding a derivation: $y^{2.000}$

It's really easy to find:

 $2.000 \cdot y$

3 step: finding a derivation: x

Even my two-aged sister knows that it equals:

1.000

4 step: finding a derivation: $x \cdot y^{2.000}$

It's simple as fuck:

 $y^{2.000} + 2.000\cdot y\cdot x$

Congratulations! The first derivation of the expression is:

 $y^{2.000} + 2.000 \cdot y \cdot x$

In the point (x = 3.000, y = 2.000) it's value = 16.000

Let's calculate the 3 derivation of the expression:

Calculating the 1 derivation of the expression:

1 step: finding a derivation: y

If someone asked me that in the middle of the night, I wouldn't hesitate to say:

1.000

2 step: finding a derivation: $y^{2.000}$

It's simple as fuck:

 $2.000 \cdot y$

3 step: finding a derivation: x

It's really easy to find:

1.000

4 step: finding a derivation: $x \cdot y^{2.000}$

Even my two-aged sister knows that it equals:

 $y^{2.000} + 2.000\cdot y\cdot x$

Calculating the 2 derivation of the expression:

1 step: finding a derivation: x

Man... Just look:

1.000

2 step: finding a derivation: y

It's really easy to find:

1.000

3 step: finding a derivation: 2.000

Even my two-aged sister knows that it equals:

0.000

4 step: finding a derivation: $2.000 \cdot y$

Even my two-aged sister knows that it equals:

2.000

5 step: finding a derivation: $2.000 \cdot y \cdot x$

It's simple as fuck:

 $2.000 \cdot x + 2.000 \cdot y$

6 step: finding a derivation: y

Man... Just look:

1.000

7 step: finding a derivation: $y^{2.000}$

If someone asked me that in the middle of the night, I wouldn't hesitate to say:

 $2.000 \cdot y$

8 step: finding a derivation: $y^{2.000} + 2.000 \cdot y \cdot x$

It's really easy to find:

 $2.000 \cdot y + 2.000 \cdot x + 2.000 \cdot y$

Calculating the 3 derivation of the expression:

1 step: finding a derivation: y

It's simple as fuck:

1.000

2 step: finding a derivation: 2.000

It's really easy to find:

0.000

3 step: finding a derivation: $2.000 \cdot y$

Even my two-aged sister knows that it equals:

2.000

4 step: finding a derivation: x

It's really easy to find:

1.000

 $5~\mathrm{step} \colon \mathrm{finding}$ a derivation: 2.000

It's really easy to find:

0.000

6 step: finding a derivation: $2.000 \cdot x$

If someone asked me that in the middle of the night, I wouldn't hesitate to say:

2.000

7 step: finding a derivation: $2.000 \cdot x + 2.000 \cdot y$

Even my two-aged sister knows that it equals:

4.000

8 step: finding a derivation: y

Man... Just look:

1.000

9 step: finding a derivation: 2.000

Even my two-aged sister knows that it equals:

0.000

10 step: finding a derivation: $2.000 \cdot y$

It's simple as fuck:

2.000

11 step: finding a derivation: $2.000 \cdot y + 2.000 \cdot x + 2.000 \cdot y$

Even my two-aged sister knows that it equals:

6.000

Finally... The 3 derivation of the expression:

6.000

BRITISH SCIENTISTS WERE SHOCKED AGAIN, WHEN THEY COUNT THE 3 DERIVATION OF THIS EXPRESSION!!!

In the point (x = 3.000, y = 2.000)it's value = 6.000

Partial derivation of the expression on the variable 'x':

 $\frac{\partial f}{\partial x} = 4.000$

In the point (x = 3.000, y = 2.000) it's value = 4.000000!!!

Partial derivation of the expression on the variable 'y':

 $\frac{\partial f}{\partial y} = 3.000 \cdot 2.000 \cdot y$

In the point (x = 3.000, y = 2.000) it's value = 12.000000!!!

Full derivation:

$$\sqrt{16.000 + (3.000 \cdot 2.000 \cdot y)^{2.000}}$$

In the point (x = 3.000, y = 2.000)it's value = 12.649 !!!

Let's consider the expression as a function of x variable: f(x) = $4.000 \cdot x$

Maklorens formula for $x \to 3.000$: $f(x) = 12.000 + 4.000 \cdot (x - 3.000) + o((x - 3.000)^{4.000})$

Graph f(x):

Tangent equation in point -2.000:

 $f(x) = 4.000 \cdot (x - (-2.000)) + (-8.000)$

Normal equation in point -2.000: $f(x) = (-0.250) \cdot (x - (-2.000)) + (-8.000)$