

**Exercise 1.** Show that  $(S^m, *) \wedge (S^n, *) \cong (S^{m+n}, *)$ .

*Solution.* Using exercises 3 and 6 from the previous tutorial, we have

$$\begin{aligned} (S^m, *) \wedge (S^n, *) &\cong (D^m/S^{m-1}) \wedge (D^n/S^{n-1}) \cong \\ &\cong (D^m \times D^n)/(S^{m-1} \times D^m \cup D^n \times S^{n-1}) \cong \\ &\cong (I^m \times I^n)/(\partial I^m \times I^n \cup \partial I^n \times I^m) = \\ &= I^{m+n}/(\partial(I^{m+n})) \cong D^{m+n}/\partial D^{m+n} \cong S^{m+n}. \end{aligned}$$

□

**Exercise 2.** Show that  $\mathbb{C}P^n$  is a CW-complex.

*Solution.* Clearly  $\mathbb{C}P^0$  is a point. Next, we have

$$\begin{aligned} \mathbb{C}P^n &= \mathbb{C}^{n+1} \setminus \{0\} / \{v \sim \lambda v, \lambda \in \mathbb{C} \setminus \{0\}\} \cong S^{2n+1} \setminus \{0\} / \{v \sim \lambda v, |\lambda| = 1\} \cong \\ &\cong \{(w, \sqrt{1-|w|^2}), w \in D^{2n}\} / \{w \sim \lambda w \text{ for } |w| = 1\} \cong \\ &\cong (D^{2n} \cup S^{2n-1}) / \{w \sim \lambda w \text{ for } w \in S^{2n-1}\} = D^{2n} \cup_f \mathbb{C}. \end{aligned}$$

Taking the canonical projection  $S^{2n-1} \rightarrow \mathbb{C}P^{n-1} \cong S^{2n-1}/\sim$  as the attaching map now yields a CW-complex with one cell in every even dimension and none in the odd ones. □

**Exercise 3.** From the lecture we know that  $A := \{\frac{1}{n}, n \in \mathbb{N}\} \cup \{0\}$  with the subset topology from  $\mathbb{R}$  is not a CW-complex. Show that  $X := I \times \{0\} \cup A \times I$  is not a CW-complex.

*Solution.* Suppose that  $X$  is a CW-complex. Then it cannot contain cells of dimension  $\geq 2$ , because it becomes disconnected after removing any point. In fact, the space obtained after removing any point  $(a, 0)$  with  $a \in A$  has more than two connected components (three, to be exact), so these points cannot lie inside a 1-cell. Therefore these points must form 0-cells, but we already know that  $A$  does not have discrete topology, a contradiction. □

**Exercise 4.** Show that the Hawaiian earring given by

$$X = \{(x, y) \in \mathbb{R}^2, (x - \frac{1}{n})^2 + y^2 = \frac{1}{n^2} \text{ for some } n\}$$

is not a CW-complex.

*Solution.* Suppose that  $X$  is a CW-complex. Using similar arguments as in the previous exercise, we can see that  $(0, 0)$  must be a 0-cell and that  $X$  must have either infinitely many 0-cells, or infinitely many 1-cells. But since  $X$  is compact, exercise 5 implies that  $X$  can have only finitely many cells, a contradiction. □

**Exercise 5.** Prove that every compact set  $A$  in a CW-complex  $X$  can have a nonempty intersection with only finitely many cells.

*Solution.*  $X$  is comprised of cells that are indexed by elements of some set  $J$ . Let  $B$  be a set containing exactly one point from each intersection  $A \cap e^\beta$ ,  $\beta \in J$ . We need to show that  $B$  is closed and discrete, which will imply that  $B$  is compact (since  $B \subseteq A$ ) and discrete, hence finite. We know that a set  $C \subseteq X^n$  is closed iff both  $C \cap X^{n-1}$  and  $C \cap e_\alpha^n$  for each  $\alpha \in J$  are closed, because  $D^n \cup_f X^{n-1}$  is a pushout. Using induction, this implies that  $C \subseteq X$  is closed iff  $C \cap e_\alpha$  is closed for each  $\alpha \in J$ . Since  $B \cap e_\alpha$  contains at most one point for any  $\alpha \in J$  and  $X$  is  $T_1$  (even Hausdorff), this shows that  $B$  is closed. Using the same argument,  $B$  with any one point removed is closed. Therefore  $B$  is also discrete and we are done.  $\square$

**Exercise 6.** Show that for a short exact sequence  $0 \rightarrow A \xrightarrow{f} B \xrightarrow{g} C \rightarrow 0$  of abelian groups (or more generally modules over a commutative ring) the following are equivalent:

- (1) There exists  $p : B \rightarrow A$  such that  $pf = \text{id}_A$ .
- (2) There exists  $q : C \rightarrow B$  such that  $gq = \text{id}_C$ .
- (3) There exist  $p : B \rightarrow A$  and  $q : C \rightarrow B$  such that  $fp + qg = \text{id}_B$ .

(Another equivalent condition is  $B \cong A \oplus C$ , with  $(p, g)$  and  $f + q$  being the respective inverse isomorphisms.)

*Solution.*

(1)  $\implies$  (2) and (3):

Since  $g$  is surjective, for any  $c \in C$  there is some  $b \in B$  such that  $g(b) = c$ . Moreover, for any other  $b' \in B$  such that also  $g(b') = c$ , we have  $b - fp(b) = b' - fp(b')$ , since  $b - b' \in \ker g = \text{im} f$ , so that  $b - b' = f(a)$  and

$$fp(b - b') = fpf(a) = f(a) = b - b'.$$

This shows that we can correctly define  $q(c) := b - fp(b)$  for any such  $b$ . Then we have

$$gq(c) = g(b) - gfp(b) = g(b) = c$$

(since  $gf = 0$ ), which shows that  $gq = \text{id}_C$ , and also  $qg(b) = b - fp(b)$ , hence  $fp + qg = \text{id}_B$ .

(3)  $\implies$  (1) and (2):

Applying  $f$  from the right to the equation  $fp + qg = \text{id}_B$  yields  $fpf = f$  (since  $gf = 0$ ), which together with the fact that  $f$  is injective implies  $pf = \text{id}_A$ . Similarly, applying  $g$  from the left yields  $gqg = g$ , which together with the fact that  $g$  is surjective implies  $gq = \text{id}_C$ .  $\square$

**Exercise 7.** Let  $0 \rightarrow A_* \xrightarrow{f} B_* \xrightarrow{g} C_* \rightarrow 0$  be a short exact sequence of chain modules. We have defined the connecting homomorphism  $\partial_* : H_n(C) \rightarrow H_{n-1}(A)$  by the formula  $\partial_*[c] = [a]$ , where  $\partial c = 0$ ,  $f(a) = \partial b$  and  $g(b) = c$ . Show that this definition does not depend on  $a$  nor  $b$ .

*Solution.* We have  $\partial a = 0$  iff  $f(\partial a) = 0$  (using injectivity of  $f$ ) iff  $0 = \partial f(a) = \partial \partial b$ , and the last condition is true.

Now let  $b, b' \in B$  be such that  $g(b) = g(b') = c$  with  $a, a' \in A$  such that  $f(a) = b, f(a') = b'$ . Then  $b - b' \in \ker g = \operatorname{im} f$ , so  $b - b' = f(\bar{a})$  for some  $\bar{a} \in A$ . Therefore  $f(\partial \bar{a}) = \partial b - \partial b' = f(a - a')$  and the injectivity of  $f$  implies  $\partial \bar{a} = a - a'$ , hence  $[0] = [\partial a'] = [a] - [a']$  and we are done.  $\square$