

Computability of the Inverse Fabius Function

**Sequential computation, effective uniform continuity,
exact inverse moduli, and certified Thue–Morse bisection**

Research report prepared for Vladimir Reshetnikov’s ProveIt project

30 August 2026

Repository snapshot: 40fdea4cc0a728189f357389e3f114a2cb00e561

Abstract

Let F be the canonical bounded Fabius function, viewed as the strictly increasing distribution function on $[0, 1]$, and let $G = F^{-1}$. We prove constructively that G is a computable real function in the Grzegorzcz/Wikipedia sense [1, 2, 9]: it maps every computable real sequence to a computable real sequence, and it possesses a recursive modulus of uniform continuity. The same theorem holds for the repository’s totalized inverse $\overline{G} : \mathbb{R} \rightarrow [0, 1]$, obtained by clamping the argument to $[0, 1]$.

The proof is quantitative. First, the centered finite random sums give exact rational Thue–Morse splines S_N satisfying the global certificate $|S_N - F| \leq 2^{-N}$. Thus F can be evaluated at a rational point to any requested precision by a finite integer computation. Second, the Fabius density is symmetric and strictly unimodal. It follows that among all subintervals of $[0, 1]$ of a fixed nonzero length h , exactly the endpoint interval starts minimize the Fabius mass:

$$\min_{0 \leq x \leq 1-h} (F(x+h) - F(x)) = F(h).$$

More precisely, equality holds only for $x = 0$ or $x = 1 - h$ (these coincide when $h = 1$); for $h = 0$ every admissible start is degenerate. Consequently the inverse-gap bound also has a complete equality classification, not merely a sharp endpoint witness. The exact modulus of continuity of the inverse is the inverse itself, with the natural saturation at unit input distance:

$$\sup_{\substack{u, v \in [0, 1] \\ |u - v| \leq \eta}} |G(u) - G(v)| = G(\min(\eta, 1)).$$

This identity reduces effective continuity to a positive lower bound for $F(h)$. A direct box event in the defining random series gives, for every $r \geq 0$,

$$F(2^{-r}) \geq \Delta_r := 2^{-r(r+1)/2} (r+1)^{-(r+1)}.$$

Hence $|u - v| < \Delta_r$ implies $|\overline{G}(u) - \overline{G}(v)| < 2^{-r}$, yielding an explicit primitive-recursive modulus. Combined with the standard flatness upper bound, this also determines the optimal binary input precision for p output bits:

$$\mu(p) = \frac{p^2}{2} + p \log_2 p + \mathcal{O}(p).$$

Finally, a tolerant bisection algorithm uses certified approximations to both $F(c)$ and the target value. It never asks an undecidable exact-equality question: when comparison is inconclusive, the inverse modulus certifies that the midpoint is already sufficiently accurate. This supplies a uniform name-transformer and proves sequential computability.

The report audits the recursive LaTeX corpus under `Analysis/FabiusFunction/docs`, separates imported repository results from the arguments proved here, relates the construction to q -binomial and sinc-product representations, Lambert- W inverse asymptotics, Legendre expansions, and Bell/Bernoulli coefficient calculi. It records both the completed post-publication Lean crosswalks for the structural inverse modulus and effective uniform continuity, and the remaining roadmap for a computable name-transformer. An exact-rational Python supplement reproduces the finite-spline and modulus calculations.

Status, snapshot, and formalization boundary

Every statement labelled theorem, proposition, lemma, or corollary below is proved in this report or explicitly imported with a source pointer. The repository audit and all relative-novelty statements remain pinned to commit 40fdea4cc0a728189f357389e3f114a2cb00e561. At that snapshot, the main computability theorem, the least fixed-length Fabius-increment inequality, the exact inverse modulus, and the closed elementary modulus were absent under the formulations used here.

A post-publication Lean development now formalizes two layers. The structural inverse-modulus layer lives in `Analysis/FabiusFunction/Lean/FabiusFunction/InverseModulus.lean`. It proves the global weak shape and maximal strict branches of fixed-length Fabius increments, the endpoint lower bound and complete equality locus, constrained forward superadditivity with its equality cases, local and global inverse-gap inequalities, unit-interval inverse-gap rigidity, global inverse subadditivity, attained exact moduli on both $[0, 1]$ and $\mathbb{R}$, and the exact effective-injectivity threshold. The downstream module `Analysis/FabiusFunction/Lean/FabiusFunction/FabiusInverseEffectiveContinuity.lean` proves a one-term recurrence lower bound stronger than the elementary Δ_r estimate, derives the exact numerical inequality $\Delta_r \leq F(2^{-r})$ and the corresponding strict and closed dyadic inverse moduli, and packages a primitive-recursive factorial-denominator witness for `EffectivelyUniformContinuous`. Exhaustive declaration-level crosswalks are given in [section 12](#).

The probabilistic box-event *proof* of the Δ_r inequality is not formalized: Lean reaches the same inequality by the stronger recurrence route. The logarithmic choice $r(n)$, the exact ceiling denominator d_* , tolerant bisection, sequential computability, the combined computable-real-function theorem, and the input-precision asymptotics also remain outside the current formalization. Those claims retain the complete human proofs and imported-source qualifications stated below. “New” means new relative to the pinned corpus; no unqualified worldwide priority claim is made. Conjectures and research problems are explicitly labelled and are not used in the proof.

Contents

1	Main result and proof architecture	5
1.1	The theorem in one statement	5
1.2	Four proof layers	5
2	Corpus audit, conventions, and nonduplication boundary	6
2.1	Pinned repository snapshot	6
2.2	What the repository already proves	6
2.3	The gap closed here	7
2.4	Three functions that must not be conflated	7
3	Computable-analysis definitions	7
3.1	Fast names and computable sequences	7
3.2	The reciprocal uniform-continuity convention	8
4	Probability, Fourier structure, and the forward function	8
4.1	The random-series model	8
4.2	Finite sums and their exact CDFs	9
4.3	Centered finite splines and a uniform certificate	9

5	Effective positivity, strict increase, and the density shape	10
5.1	A closed positive lower bound at dyadic points	10
5.2	Strict monotonicity from the random series	12
5.3	Symmetric strict unimodality	12
6	The least-mass interval and the exact inverse modulus	12
6.1	Intervals of a prescribed geometric length	12
6.2	The inverse's modulus is the inverse itself	14
7	Effective uniform continuity	16
7.1	A closed primitive-recursive modulus	16
7.2	The exact-dyadic repository modulus	18
7.3	Binary conditioning is quadratic	19
8	Sequential computability by tolerant bisection	20
8.1	Why ordinary bisection is insufficient	20
8.2	The algorithm on a computable sequence	20
8.3	Correctness of every branch	21
8.4	Clamping and the totalized inverse	22
8.5	Completion of the main theorem	22
8.6	An abstract inverse-computability principle	22
9	Analytic consequences of the exact modulus	22
9.1	Uniform continuity without any Hölder exponent	22
9.2	Endpoint asymptotics become modulus asymptotics	23
9.3	Derivative blow-up versus computability	24
10	Numerical and exact-rational checks	24
10.1	Endpoint mass bounds	24
10.2	Centered-spline convergence	26
10.3	What the code deliberately does not claim	26
11	Connections to the wider Fabius–Rvachev corpus	26
11.1	Thue–Morse inclusion–exclusion	26
11.2	q -binomial coefficients and accelerated forward oracles	26
11.3	Infinite sinc products and alternative evaluation	26
11.4	Lambert W , Bell polynomials, and Bernoulli data	27
11.5	Legendre and Lagrange representations	27
11.6	Total positivity, unimodality, and interval masses	27
12	Lean formalization status and remaining roadmap	27
12.1	Why the present noncomputable definition is not an obstacle	27
12.2	The completed structural inverse-modulus layer	27
12.3	The completed effective-uniform-continuity layer	29
12.4	The remaining name-transformation module	30
12.5	Remaining formalization order	31
12.6	Reusable generic infrastructure	31
13	Further deductions, conjectures, and research directions	31
13.1	A certified practical inverse evaluator	31
13.2	Sharp integer moduli and the log-periodic phase	32
13.3	Base- b atomic distributions	32
13.4	Certified spectral and polynomial representations	33

13.5 Higher-dimensional monotone transport	33
13.6 Branchwise inversion of the signed global extension	33
14 Conclusion	33
A Pseudocode with explicit error budgets	34
B Proof-dependency and status audit	35
C Corpus crosswalk	36
D Reproducing the exact-rational supplement	38

1 Main result and proof architecture

1.1 The theorem in one statement

Throughout the report, $F : \mathbb{R} \rightarrow [0, 1]$ denotes the bounded Fabius function: it is 0 on $(-\infty, 0]$, 1 on $[1, \infty)$, and on $[0, 1]$ it is the strictly increasing C^∞ function satisfying

$$F(x) + F(1 - x) = 1, \quad F'(x) = 2F(2x) \quad (0 \leq x \leq \tfrac{1}{2}). \quad (1.1)$$

Let $G : [0, 1] \rightarrow [0, 1]$ be its inverse, and define the totalized inverse

$$\overline{G}(y) := G(\text{clamp}(y)), \quad \text{clamp}(y) := \min(1, \max(0, y)). \quad (1.2)$$

Thus $\overline{G}(y) = 0$ for $y \leq 0$ and $\overline{G}(y) = 1$ for $y \geq 1$.

Theorem 1.1 (Computability of the inverse Fabius function). *The inverse $G : [0, 1] \rightarrow [0, 1]$, in the subset-domain version of the Grzegorzczuk definition, and its totalization $\overline{G} : \mathbb{R} \rightarrow \mathbb{R}$ are computable real functions. More explicitly:*

1. G and $\overline{G}$ are sequentially computable. From one algorithm that names a computable sequence (y_i) , one can uniformly compute dyadic approximations to $G(y_i)$ (or $\overline{G}(y_i)$) of any requested precision.

2. For $r \geq 0$, put

$$\Delta_r = \frac{1}{2^{r(r+1)/2}(r+1)^{r+1}}. \quad (1.3)$$

Then, for all $u, v \in \mathbb{R}$,

$$|u - v| < \Delta_r \implies |\overline{G}(u) - \overline{G}(v)| < 2^{-r}. \quad (1.4)$$

3. For $n \geq 1$, let

$$r(n) := \min \{r \geq 1 : 2^r > n\}, \quad (1.5)$$

and define

$$d(n) := 2^{r(n)(r(n)+1)/2}(r(n)+1)^{r(n)+1}, \quad d(0) := 1. \quad (1.6)$$

The integer function d is recursive (indeed primitive recursive after a standard bounded-search coding), and

$$|u - v| < \frac{1}{d(n)} \implies |\overline{G}(u) - \overline{G}(v)| < \frac{1}{n} \quad (n \geq 1). \quad (1.7)$$

The theorem is stronger than a nonconstructive assertion that a continuous inverse on a compact interval is uniformly continuous. It gives a concrete integer modulus, a finite algorithm on names, and a sharp structural identity for the best possible modulus.

1.2 Four proof layers

The argument is organized so that every hidden effectiveness issue is exposed.

1. **Forward oracle.** A centered finite convolution has an exact Thue–Morse truncated-power formula. At rational inputs it is rational, and its error from F is at most 2^{-N} .
2. **Effective injectivity.** Symmetric unimodality shows that an interval of geometric length h has probability mass at least $F(h)$; strict radial unimodality shows that equality occurs only for a degenerate interval or an endpoint interval. A direct event gives the computable lower bound $F(2^{-r}) \geq \Delta_r$.
3. **Uniform continuity.** Effective injectivity of F becomes an effective modulus for G . In fact the ordinary modulus of G is exactly G itself.

4. **Name transformation.** Tolerant bisection combines the forward oracle and the inverse modulus. A midpoint is used for a left/right update only when the sign is certified; otherwise it is already a certified output.

The fourth layer matters. Naive bisection that waits until it decides whether $F(c) < y$, $F(c) = y$, or $F(c) > y$ is not an algorithm on arbitrary computable reals: equality of computable reals is undecidable in general. The tolerance branch is what removes that obstruction.

2 Corpus audit, conventions, and nonduplication boundary

2.1 Pinned repository snapshot

The recursive audit was pinned to commit `40fdea4cc0a728189f357389e3f114a2cb00e561` [8]. The GitHub code index at its parent snapshot enumerated 94 files with extension `.tex` below `Analysis/FabiusFunction/docs`; the final merge added two further LaTeX sources, for a total of 96 in the pinned tree. All source paths were inventoried. Primary and consolidated mathematical reports were read directly; generated table fragments and frozen variants were checked together with the parent documents that include them.

The inspected corpus spans the following overlapping families:

- the two source papers and the main Fabius/Rvachev exposition;
- the Lean walkthrough, paper-coverage map, and mathematical glossary;
- exact dyadic arithmetic, Thue–Morse and q -binomial formulae;
- finite convolutions, infinite sinc products, Fourier decay, canonical products, and Poisson/Legendre representations;
- endpoint asymptotics, lower-Lambert phases, Gamma–zeta periodic terms, Bell-polynomial inverse transfer, and all-orders inverse expansions;
- inverse-dyadic sampling, q -Richardson acceleration, Lagrange and Legendre synthesis, total positivity, spectra, and operator viewpoints;
- the final added spectral/ q -Pochhammer and digital-geometry material.

The audit was thematic rather than a search for one spelling. In particular, it traced the bounded CDF, signed global extension, Rvachev function, totalized inverse, finite spline evaluator, exact dyadic evaluator, inverse asymptotics, and the repository’s definitions of sequential computability and effective uniform continuity.

2.2 What the repository already proves

The following inputs are established in the pinned development and are not claimed as new here.

1. The bounded canonical F exists, is unique, strictly increasing on $[0, 1]$, symmetric, C^∞ , and flat at both endpoints.
2. F is the CDF of the random series in [equation \(4.1\)](#) below. Its centered density is Rvachev’s up function, with Fourier transform an infinite dyadic sinc product.
3. Every dyadic value of F is an exact rational and is computed by an executable recurrence. The centered Thue–Morse splines converge uniformly with error at most 2^{-N} .
4. The forward bounded F and the signed global extension are already proved computable real functions. The relevant Lean modules are `FabiusComputability.lean` and `FabiusComput`

`ableSpline.lean`; they expose the sequential-computability and effective-uniform-continuity clauses, including a primitive-recursive spline evaluator.

5. The module `FabiusInverse.lean` constructs the clamped totalized inverse `fabiusInv`, proves both inverse identities, continuity, monotonicity, interior C^∞ regularity, derivative and curvature laws, and endpoint steepness.
6. The inverse/sampling reports derive exact finite-prefix quantile germs, q -binomial Richardson filters, and all-orders endpoint asymptotics for G , including the nonconstant log-periodic Gamma-zeta fluctuation.

2.3 The gap closed here

No theorem named or formulated as sequential computability, effective uniform continuity, or computable-real-function status for `fabiusInv` occurs in the pinned snapshot. The missing bridge is not a formal triviality. A computable strictly increasing function can be inverted effectively on a compact interval only after one supplies effective injectivity information; ordinary strict monotonicity is qualitative.

This report supplies that information in two forms:

$$F(x + h) - F(x) \geq F(h), \quad (2.1)$$

$$F(2^{-r}) \geq \Delta_r. \quad (2.2)$$

The first is exact and geometric. For $0 < h \leq 1$, equality in the first bound occurs exactly at $x = 0$ or $x = 1 - h$; in the equivalent form $F(b) - F(a) = F(b - a)$ the complete alternatives are $a = b$, $a = 0$, or $b = 1$. The second bound is elementary and recursive. Together with the existing forward evaluator they close both clauses of the requested definition.

The historical statement above remains about the pinned snapshot. In the current post-publication development, `InverseModulus.lean` formally proves the first inequality, its complete equality locus, and its exact transfer through the clamped inverse, including rigidity, sharpness, and attainment. `FabiusInverseEffectiveContinuity.lean` formally proves the second *inequality* by a stronger one-term recurrence estimate and packages a simple primitive-recursive reciprocal modulus. The report's probabilistic event proof, its smaller logarithmic- $r(n)$ packaging, and the tolerant-bisection realizer remain on the human-proof side of the boundary recorded in [section 12](#).

2.4 Three functions that must not be conflated

The documentation repeatedly warns about a notation trap. The bounded CDF F is clamped outside $[0, 1]$. The signed global extension, often denoted $\mathcal{F}$, agrees with F on $[0, 1]$ but oscillates outside it and obeys global dilation/translation identities. Rvachev's up is an even compactly supported bump on $[-1, 1]$. On its support,

$$\text{up}(t) = \begin{cases} F(t + 1), & -1 \leq t \leq 0, \\ F(1 - t), & 0 \leq t \leq 1. \end{cases} \quad (2.3)$$

The inverse in this report is the inverse of the bounded transition $F : [0, 1] \rightarrow [0, 1]$, with the clamped totalization (1.2). It is not an inverse of the signed global extension.

3 Computable-analysis definitions

3.1 Fast names and computable sequences

Following standard computable analysis [1, 2, 3], a convenient representation of a real number x is a fast dyadic Cauchy name: an algorithm returns a dyadic rational q_p such that

$$|x - q_p| \leq 2^{-p}. \quad (3.1)$$

A real sequence $(x_i)_{i \in \mathbb{N}}$ is computable when one algorithm returns $q_{i,p}$ satisfying (3.1) uniformly in both i and p . The repository uses an equivalent signed-natural code for the dyadic numerator, avoiding an opaque integer encoding.

Definition 3.1 (Sequential computability). A function f is *sequentially computable* if every computable real sequence (x_i) in its domain is mapped to a computable sequence $(f(x_i))$, uniformly in the sequence index and requested precision.

The word “uniformly” is implicit in any useful implementation of this definition: the output algorithm receives the input naming algorithm as a subroutine. The construction in section 8 gives such a uniform transformer, and therefore proves the weaker extensional statement as well.

3.2 The reciprocal uniform-continuity convention

Definition 3.2 (Effective uniform continuity). A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is *effectively uniformly continuous* if there is a recursive $d : \mathbb{N} \rightarrow \mathbb{N}$ with $d(n) > 0$ for $n > 0$ such that

$$|x - y| < \frac{1}{d(n)} \implies |f(x) - f(y)| < \frac{1}{n} \quad (n > 0). \quad (3.2)$$

This is the reciprocal convention used in the cited Wikipedia article [9]. The repository implements it in `FabiusComputability.lean`. It is equivalent to the more common binary modulus convention

$$|x - y| < 2^{-m(p)} \implies |f(x) - f(y)| < 2^{-p}, \quad (3.3)$$

provided m is recursive. We give both forms because the binary version reveals the endpoint conditioning much more clearly.

Definition 3.3 (Computable real function). In the Grzegorczyk formulation used here, a real function is computable if it is sequentially computable and effectively uniformly continuous.

For G defined only on $[0, 1]$, the same definitions are restricted to sequences and pairs of points in that computable compact domain. The totalized $\overline{G}$ lets one use the literal type $\mathbb{R} \rightarrow \mathbb{R}$ without changing the mathematics.

4 Probability, Fourier structure, and the forward function

4.1 The random-series model

The classical probability model may be written as follows [4, 6, 8]. Let $U_1, U_2, \dots$ be independent uniform random variables on $[0, 1]$ and put

$$X := \sum_{k=1}^{\infty} 2^{-k} U_k. \quad (4.1)$$

The series converges absolutely, $0 \leq X \leq 1$, and

$$F(x) = \mathbb{P}[X \leq x]. \quad (4.2)$$

Reflection $U_k \mapsto 1 - U_k$ sends X to $1 - X$ and proves the symmetry in (1.1).

The centered random variable

$$Y := 2X - 1 = \sum_{k=1}^{\infty} 2^{-k} (2U_k - 1) \quad (4.3)$$

has density up. With $\text{sinc } z = \sin z/z$,

$$\mathbb{E}e^{itY} = \prod_{k=1}^{\infty} \text{sinc}\left(\frac{t}{2^k}\right). \quad (4.4)$$

Thus the probability model, the Rvachev bump, and the infinite sinc product are three representations of the same object. The computability proof will use the probability representation in physical space; [section 11](#) explains how the other representations can serve as alternative evaluators.

4.2 Finite sums and their exact CDFs

For $N \geq 1$, define

$$X_N := \sum_{k=1}^N 2^{-k} U_k, \quad R_N := X - X_N. \quad (4.5)$$

Then

$$0 \leq R_N \leq 2^{-N}. \quad (4.6)$$

Let $F_N(x) = \mathbb{P}[X_N \leq x]$. The general inclusion–exclusion formula for a sum of independent, nonidentically scaled uniforms becomes especially rigid for dyadic scales.

Proposition 4.1 (Finite Thue–Morse CDF). *For $N \geq 1$ and $x \in \mathbb{R}$,*

$$F_N(x) = \frac{2^{N(N+1)/2}}{N!} \sum_{j=0}^{2^N-1} (-1)^{s_2(j)} \left(x - \frac{j}{2^N}\right)_+^N, \quad (4.7)$$

where $s_2(j)$ is the sum of the binary digits of j .

Proof. For positive lengths $a_1, \dots, a_N$, the CDF of $\sum a_k U_k$ is

$$\frac{1}{N! \prod_{k=1}^N a_k} \sum_{A \subseteq \{1, \dots, N\}} (-1)^{|A|} \left(x - \sum_{k \in A} a_k\right)_+^N.$$

This follows by integrating the box indicator over the half-space $\sum a_k u_k \leq x$, or inductively by finite differences of truncated powers. Here $a_k = 2^{-k}$ and $\prod a_k = 2^{-N(N+1)/2}$. The map

$$A \mapsto j(A) := \sum_{k \in A} 2^{N-k}$$

is a bijection from subsets to $\{0, \dots, 2^N - 1\}$, with $\sum_{k \in A} 2^{-k} = j(A)/2^N$ and $|A| \equiv s_2(j(A)) \pmod{2}$. Substitution gives (4.7). $\square$

The alternating signs are exactly the Thue–Morse signs. No limiting argument is hidden in (4.7): at rational x it is an exact rational obtained by finitely many integer operations.

4.3 Centered finite splines and a uniform certificate

The deterministic midpoint of the omitted tail is 2^{-N-1} . Define

$$Q_N := X_N + 2^{-N-1}, \quad S_N(x) := \mathbb{P}[Q_N \leq x] = F_N(x - 2^{-N-1}). \quad (4.8)$$

Then [proposition 4.1](#) gives

$$S_N(x) = \frac{2^{N(N+1)/2}}{N!} \sum_{j=0}^{2^N-1} (-1)^{s_2(j)} \left(x - \frac{2j+1}{2^{N+1}}\right)_+^N. \quad (4.9)$$

This is the centered uniform spline used by the repository’s primitive- recursive evaluator.

Lemma 4.2 (Uniform density bound). *The finite CDFs F_N and S_N are globally 2-Lipschitz.*

Proof. The density of $2^{-1}U_1$ is $2\mathbf{1}_{[0,1/2]}$. Convolution with any probability measure cannot increase the L^∞ norm of a density. Hence the density of X_N is bounded by 2. Translating by 2^{-N-1} does not change this bound. Integrating the density over an interval proves the Lipschitz estimate. $\square$

Theorem 4.3 (Certified centered approximation). *For every $N \geq 1$ and every real x ,*

$$|S_N(x) - F(x)| \leq 2^{-N}. \quad (4.10)$$

Proof. Write

$$X = Q_N + E_N, \quad E_N = R_N - 2^{-N-1}.$$

By (4.6), $|E_N| \leq 2^{-N-1}$. Therefore

$$S_N(x - 2^{-N-1}) \leq F(x) \leq S_N(x + 2^{-N-1}).$$

By lemma 4.2, moving the argument by 2^{-N-1} changes S_N by at most $2 \cdot 2^{-N-1} = 2^{-N}$. This gives both sides of (4.10). $\square$

Corollary 4.4 (Self-contained computability of F). *There is an explicit uniform algorithm which, given a rational x and a precision p , returns a rational q with $|q - F(x)| \leq 2^{-p}$. One may take $q = S_N(x)$ with $N \geq p$. Consequently F is sequentially computable; together with its global 2-Lipschitz bound, it is a computable real function.*

Proof. Formula (4.9) is a finite rational computation, and theorem 4.3 supplies its error certificate. At an arbitrary computable input, first replace the input by a sufficiently accurate dyadic; the 2-Lipschitz bound controls this replacement. The construction is uniform in a sequence index. Effective uniform continuity follows from $|F(x) - F(y)| \leq 2|x - y|$; for example the reciprocal modulus $d(n) = 2n$ works. $\square$

Remark 4.5 (Efficiency versus computability). The direct sum has 2^N terms and is not intended as an efficient evaluator. The repository's exact dyadic recurrence, highest-bit reduction, and primitive-recursive natural-number implementation are substantially better. For the theorem here, only termination and a proved error bound are required.

5 Effective positivity, strict increase, and the density shape

5.1 A closed positive lower bound at dyadic points

The existence of an inverse requires strict increase; effective inversion requires a quantitative version. Both begin with an elementary positive probability event.

Lemma 5.1 (Dyadic endpoint box event). *For every $r \geq 0$,*

$$F(2^{-r}) \geq \Delta_r := 2^{-r(r+1)/2} (r+1)^{-(r+1)}. \quad (5.1)$$

In particular, $F(x) > 0$ for every $x > 0$.

Proof. Put $m = r + 1$. For $1 \leq k \leq m$, impose the independent restrictions

$$U_k \leq a_k := \frac{2^{k-r-1}}{m}. \quad (5.2)$$

The largest a_k is $a_m = 1/m \leq 1$, so these are valid events. On their intersection,

$$2^{-k}U_k \leq \frac{2^{-r-1}}{m} \quad (1 \leq k \leq m).$$

After summation the first m terms contribute at most 2^{-r-1} . The unrestricted tail contributes at most

$$\sum_{k=m+1}^{\infty} 2^{-k} = 2^{-m} = 2^{-r-1}.$$

Thus the event in (5.2) implies $X \leq 2^{-r}$. Its probability is

$$\begin{aligned} \prod_{k=1}^m a_k &= m^{-m} 2^{\sum_{k=1}^m (k-r-1)} \\ &= m^{-m} 2^{m(m+1)/2 - m(r+1)} \\ &= 2^{-r(r+1)/2} (r+1)^{-(r+1)} = \Delta_r. \end{aligned}$$

This proves (5.1). Given $x > 0$, choose r with $2^{-r} \leq x$ and use monotonicity of a CDF. $\square$

The bound is deliberately elementary: it uses only independence, the support of the tail, and integer arithmetic. It is not numerically sharp, but [section 7.3](#) shows that its logarithm has the correct two largest orders for computability purposes.

The numerical conclusion of [lemma 5.1](#) is now Lean-formalized, but by a different and stronger route. Here is the complete recurrence argument. Let $a_0 = 1$ and use the repository's exact inverse-dyadic identities

$$a_r = \frac{\sum_{0 \leq k < r} a_k / (r - k + 1)!}{2^r - 1} \quad (r \geq 1), \quad F(2^{-r}) = 2^{-\binom{r}{2}} a_r.$$

Every a_k is positive. Retaining only the $k = 0$ summand therefore gives

$$a_r \geq \frac{1}{(r+1)!(2^r - 1)}, \quad F(2^{-r}) \geq \frac{1}{2^{\binom{r}{2}} (r+1)!(2^r - 1)} \quad (r \geq 1).$$

Since $2^r - 1 \leq 2^r$ and $\binom{r}{2} + r = \binom{r+1}{2}$, taking the larger denominator yields the first inequality below; the exceptional index $r = 0$ is simply $F(1) = 1$. Thus for every $r \geq 0$,

$$F(2^{-r}) \geq \frac{1}{2^{\binom{r+1}{2}} (r+1)!} \geq \frac{1}{2^{\binom{r+1}{2}} (r+1)^{r+1}} = \Delta_r. \quad (5.3)$$

Indeed, $(r+1)! \leq (r+1)^{r+1}$ proves the second inequality after multiplication by the positive power of two and reversal under reciprocation.

For the executable arithmetic, write

$$d_{\text{fac}}(r) := 2^{\binom{r+1}{2}} (r+1)!, \quad d_{\Delta}(r) := 2^{\binom{r+1}{2}} (r+1)^{r+1} = \Delta_r^{-1}.$$

Lean defines the first denominator by primitive recursion,

$$d_{\text{fac}}(0) = 1, \quad d_{\text{fac}}(r+1) = d_{\text{fac}}(r) 2^{r+1} (r+2).$$

Induction, using $\binom{r+2}{2} = \binom{r+1}{2} + r + 1$, proves the displayed closed form. The recursive clause uses only successor, multiplication, and exponentiation, so d_{fac} is primitive recursive. Likewise $\binom{r+1}{2} = r(r+1)/2$, and primitive-recursive closure under successor, multiplication, natural-number division, and exponentiation proves that d_{Δ} is primitive recursive. Finally $d_{\text{fac}}(r) \leq d_{\Delta}(r)$ is exactly the factorial-power inequality above, and positivity reverses this comparison between their reciprocals.

The declarations `Fabius.fabiusReal_inverse_two_pow_one_term_lower_bound`, `Fabius.inv_inverseFabiusFactorialDenominator_le_fabiusReal`, and `Fabius.inv_inverseFabiusDeltaDenominator_le_fabiusReal` certify this chain. They do not formalize the event in (5.2), its containment, or its product probability; that probabilistic proof remains report-level. The argument in this paragraph uses only the already formalized dyadic recurrence and therefore preserves the distinction between the two proof routes.

5.2 Strict monotonicity from the random series

Proposition 5.2 (Strict increase). *For $0 \leq a < b \leq 1$,*

$$F(a) < F(b). \quad (5.4)$$

Hence $F : [0, 1] \rightarrow [0, 1]$ is a homeomorphism and $G = F^{-1}$ exists.

Proof. The density of X_N is strictly positive on the interior of its support $(0, 1 - 2^{-N})$. This follows inductively: a convolution of two densities that are positive on the interiors of compact intervals is positive on the interior of their Minkowski-sum interval, because the convolution integral contains a nonempty open region on which both factors are positive.

Choose N so large that $2^{-N} < b - a$. Then the open interval $(a, b - 2^{-N})$ is nonempty and lies below the right endpoint of the support of X_N . Choose a nonempty open subinterval J of it. The preceding positivity gives $\mathbb{P}[X_N \in J] > 0$. For $X_N \in J$ and every possible $0 \leq R_N \leq 2^{-N}$, one has $a < X_N + R_N < b$. Consequently

$$F(b) - F(a) = \mathbb{P}[a < X \leq b] \geq \mathbb{P}[X_N \in J] > 0.$$

Continuity and the endpoint values $F(0) = 0$, $F(1) = 1$ then give the homeomorphism statement. $\square$

5.3 Symmetric strict unimodality

Let

$$p(x) := F'(x), \quad 0 \leq x \leq 1. \quad (5.5)$$

The repository proves that F is C^∞ , so p is continuous. The functional and reflection equations completely determine its qualitative shape.

Proposition 5.3 (Shape of the Fabius density). *The density p satisfies*

$$p(x) = p(1 - x), \quad (5.6)$$

is strictly increasing on $[0, 1/2]$, and is strictly decreasing on $[1/2, 1]$. Equivalently, $p(x)$ is a strictly decreasing function of $|x - 1/2|$.

Proof. On $[0, 1/2]$, (1.1) gives

$$p(x) = 2F(2x).$$

By [proposition 5.2](#), this is strictly increasing. Differentiating $F(x) + F(1 - x) = 1$ gives $p(x) = p(1 - x)$, which reflects the first-half behavior to the second half. $\square$

This simple observation is the geometric heart of the inverse modulus. It is also a point where Rvachev's function enters directly: by (2.3),

$$p(x) = 2 \operatorname{up}(2x - 1). \quad (5.7)$$

Thus [proposition 5.3](#) is the standard single-peaked shape of the compactly supported atomic bump.

6 The least-mass interval and the exact inverse modulus

6.1 Intervals of a prescribed geometric length

For $0 \leq h \leq 1$ and $0 \leq x \leq 1 - h$, define the fixed-length CDF increment

$$M_h(x) := F(x + h) - F(x). \quad (6.1)$$

If μ is the Fabius probability law with CDF F , then the canonical measure-theoretic interpretation is

$$M_h(x) = \mu((x, x + h]).$$

The law is atomless—indeed it has the continuous density $p = F'$ —so the value is unchanged if either or both endpoints are included. In particular,

$$\mu((x, x + h]) = \mu([x, x + h]) = \mu((x, x + h)) = \mu([x, x + h)).$$

We therefore continue to call $M_h(x)$ an interval mass, while (6.1) fixes the canonical CDF convention.

Theorem 6.1 (Strict interval-mass shape and exact minimizers). *For every $0 \leq h \leq 1$,*

$$\min_{0 \leq x \leq 1-h} (F(x+h) - F(x)) = F(h). \quad (6.2)$$

If $0 < h \leq 1$ and $m = (1-h)/2$, then the globally extended increment M_h is strictly increasing on $[-h, m]$, symmetric about m , and strictly decreasing on $[m, 1]$; outside $[-h, 1]$ it is identically zero. On the admissible-start interval $0 \leq x \leq 1-h$ one has the boundary-safe equality law

$$M_h(x) = F(h) \iff h = 0 \text{ or } x = 0 \text{ or } x = 1-h. \quad (6.3)$$

Equivalently, whenever $0 \leq a \leq b \leq 1$,

$$F(b-a) = F(b) - F(a) \iff a = b \text{ or } a = 0 \text{ or } b = 1. \quad (6.4)$$

Thus every genuinely interior interval $0 < a < b < 1$ has strictly more Fabius mass than an endpoint interval of the same length. At $h = 1$ the two endpoint formulas coincide at the single admissible start $x = 0$.

Proof. The case $h = 0$ gives $M_0 \equiv 0$, so suppose $0 < h \leq 1$ and put $m = (1-h)/2$. Reflection gives

$$\begin{aligned} M_h(1-h-x) &= F(1-x) - F(1-h-x) \\ &= (1-F(x)) - (1-F(x+h)) = M_h(x), \end{aligned}$$

so M_h is symmetric about m .

Consider $-h < x < m$. Then $0 < x+h < 1$. If $x \leq 0$, the constant left tail gives $p(x) = 0$, whereas strict positivity on the open support gives $p(x+h) > 0$. If $x > 0$, compute

$$\left(x+h-\frac{1}{2}\right)^2 - \left(x-\frac{1}{2}\right)^2 = h(2x+h-1) < 0. \quad (6.5)$$

Thus $x+h$ is strictly closer to $1/2$ than x is. The strict radial unimodality in [proposition 5.3](#) again gives $p(x+h) > p(x)$. Hence

$$M'_h(x) = p(x+h) - p(x) > 0 \quad (-h < x < m). \quad (6.6)$$

The mean-value theorem makes M_h strictly increasing on the closed interval $[-h, m]$. Reflection makes it strictly decreasing on $[m, 1]$. The clamped tails of F give $M_h = 0$ to the left of $-h$ and to the right of 1 ; this is why the global half-line shape is only non-strict, whereas the two finite branches above are maximally strict.

Finally,

$$M_h(0) = F(h) - F(0) = F(h) \quad \text{and} \quad M_h(1-h) = F(1) - F(1-h) = F(h).$$

If $0 < x < 1-h$, compare x strictly with 0 on the left branch when $x \leq m$, and compare it strictly with $1-h$ on the right branch when $m < x$. In either case $M_h(x) > F(h)$. This proves the minimum and the complete equality locus (6.3). Substituting $h = b-a$ and $x = a$ gives (6.4); its diagonal alternative $a = b$ is exactly the degenerate case $h = 0$. $\square$

Corollary 6.2 (Global increment shape and constrained superadditivity). *For every $h \geq 0$, the globally extended increment M_h is nondecreasing on the half-line*

$$x \leq \frac{1-h}{2}$$

and nonincreasing on the complementary half-line. Consequently, whenever $a, b \geq 0$ and $a + b \leq 1$,

$$F(a) + F(b) \leq F(a + b). \quad (6.7)$$

Its complete equality locus is

$$F(a) + F(b) = F(a + b) \iff a = 0 \text{ or } b = 0 \text{ or } a + b = 1. \quad (6.8)$$

Proof. The derivative argument above never used $x \geq 0$: from $x \leq (1-h)/2$ and $h \geq 0$, either $x + h \leq 1/2$ and the monotonicity of p on the left half gives $p(x) \leq p(x + h)$, or $x + h > 1/2$ and reflection replaces $p(x + h)$ by $p(1 - x - h)$, where

$$x \leq 1 - x - h \leq \frac{1}{2}.$$

Thus $M'_h \geq 0$ on the entire left half-line. Reflection $M_h(1 - h - x) = M_h(x)$ gives the right-half statement. For (6.7), apply [theorem 6.1](#) to the interval from a to $a + b$: its length is b , so

$$F(b) \leq F(a + b) - F(a).$$

Rearranging proves the inequality. Equality is exactly the equality case of [theorem 6.1](#) for the interval $[a, a + b]$: its three alternatives are $a = a + b$, $a = 0$, and $a + b = 1$, and the first is equivalent to $b = 0$. $\square$

Remark 6.3 (A general principle). The lower bound used only that a probability density on $[0, 1]$ is symmetric and unimodal. Hence the minimum-value part of [theorem 6.1](#) holds for every continuous symmetric unimodal distribution. The endpoint-only equality locus uses the genuinely stronger hypothesis that the density decreases strictly with distance from the midpoint; plateaus can create additional minimizers. The Fabius system is special not because the geometric lemma is exotic, but because $F(h)$ has unusually rich exact arithmetic and endpoint asymptotics.

6.2 The inverse's modulus is the inverse itself

For a bounded function H define its ordinary modulus of continuity by

$$\omega_H(\eta) := \sup \{ |H(u) - H(v)| : |u - v| \leq \eta \}. \quad (6.9)$$

For G the variables are restricted to $[0, 1]$; for $\overline{G}$ they range over $\mathbb{R}$.

Theorem 6.4 (Exact modulus of the inverse). *For every $\eta \geq 0$, with the inputs in the definition of ω_G restricted to $[0, 1]$,*

$$\omega_G(\eta) = G(\min(\eta, 1)). \quad (6.10)$$

In particular this is $G(\eta)$ for $0 \leq \eta \leq 1$ and is 1 thereafter. For the totalized inverse and every $\eta \geq 0$,

$$\omega_{\overline{G}}(\eta) = G(\min(\eta, 1)). \quad (6.11)$$

More strongly, for all real u, v ,

$$|\overline{G}(u) - \overline{G}(v)| \leq \overline{G}(|u - v|) = G(\min(|u - v|, 1)). \quad (6.12)$$

The associated ordered-gap and subadditivity laws are

$$\overline{G}(v) - \overline{G}(u) \leq \overline{G}(v - u) \quad (u, v \in \mathbb{R}) \quad (6.13)$$

and

$$\overline{G}(x + y) \leq \overline{G}(x) + \overline{G}(y) \quad (x, y \in \mathbb{R}). \quad (6.14)$$

On the unit interval the ordered-gap equality cases are exactly

$$G(v) - G(u) = G(v - u) \iff u = v \text{ or } u = 0 \text{ or } v = 1 \quad (0 \leq u \leq v \leq 1). \quad (6.15)$$

Equivalently, without ordering the inputs,

$$|G(u) - G(v)| = G(|u - v|) \iff u = v \text{ or } u \in \{0, 1\} \text{ or } v \in \{0, 1\} \quad (u, v \in [0, 1]). \quad (6.16)$$

These two equality classifications are intentionally unit-domain statements: the constant tails of $\overline{G}$ create additional configurations on $\mathbb{R}$.

Proof. First let $c, d \in [0, 1]$. Interchanging them if necessary, assume $c \leq d$, and put

$$a = G(c), \quad b = G(d).$$

Monotonicity of G gives $a \leq b$. The inverse identities and [theorem 6.1](#) give

$$F(b - a) \leq F(b) - F(a) = d - c. \quad (6.17)$$

Both $b - a$ and $d - c$ lie in $[0, 1]$. Applying the increasing inverse and using $G(F(b - a)) = b - a$ yields

$$b - a \leq G(d - c).$$

Removing the temporary ordering assumption gives

$$|G(c) - G(d)| \leq G(|c - d|) \quad (c, d \in [0, 1]). \quad (6.18)$$

The same argument records equality. Under $c \leq d$, equality in the inverse gap is equivalent, after applying the strictly increasing F , to equality in (6.17). By (6.4), this says

$$a = b, \quad a = 0, \quad \text{or} \quad b = 1.$$

Because $a = G(c)$ and $b = G(d)$ and F, G are inverse order isomorphisms, these alternatives are respectively $c = d$, $c = 0$, and $d = 1$. This proves (6.15). Applying it after ordering c, d proves (6.16); reversing the order exchanges the left and right endpoint alternatives.

Now put $c = \text{clamp}(u)$ and $d = \text{clamp}(v)$. Projection onto $[0, 1]$ is nonexpansive and has diameter one, so

$$|c - d| \leq \min(|u - v|, 1).$$

Combining this with (6.18) proves (6.12). If $|u - v| \leq \eta$, monotonicity then gives the upper bound $G(\min(\eta, 1))$ for both moduli. Let $\theta = \min(\eta, 1)$. The pair $(0, \theta)$ belongs to $[0, 1]^2$, has input separation $\theta \leq \eta$, and has output separation

$$|G(\theta) - G(0)| = G(\theta).$$

Thus the displayed upper bound is attained, not merely approached, proving both exact supremum identities.

For (6.13), if $u \leq v$, remove the absolute value from (6.12); then $v - u = |v - u|$. If $v < u$, the left side is nonpositive by monotonicity, whereas $\overline{G}(v - u) \geq 0$. Finally apply (6.13) with $u = x$ and $v = x + y$:

$$\overline{G}(x + y) - \overline{G}(x) \leq \overline{G}(y),$$

which rearranges to (6.14). □

Corollary 6.5 (Effective-injectivity form). *For $0 \leq h \leq 1$ and $u, v \in \mathbb{R}$,*

$$|u - v| < F(h) \implies |\overline{G}(u) - \overline{G}(v)| < h. \quad (6.19)$$

The closed-threshold and contrapositive forms are

$$|u - v| \leq F(h) \implies |\overline{G}(u) - \overline{G}(v)| \leq h, \quad (6.20)$$

$$h \leq |\overline{G}(u) - \overline{G}(v)| \implies F(h) \leq |u - v|. \quad (6.21)$$

The same statements hold for G on $[0, 1]$.

Proof. The strict hypothesis gives $|u - v| < F(h) \leq 1$, so the saturation in the exact modulus disappears. Strict monotonicity of G on $[0, 1]$ therefore gives

$$|\overline{G}(u) - \overline{G}(v)| \leq G(|u - v|) < G(F(h)) = h.$$

The same chain with non-strict inequalities proves (6.20); its contrapositive is (6.21). $\square$

Theorem 6.6 (Exact strict threshold). *For $0 \leq h \leq 1$ and every real δ ,*

$$[\forall u, v \in \mathbb{R}, |u - v| < \delta \implies |\overline{G}(u) - \overline{G}(v)| < h] \iff \delta \leq F(h). \quad (6.22)$$

Thus $F(h)$ is exactly the largest strict universal input threshold for output error below h .

Proof. If $\delta \leq F(h)$, then $|u - v| < \delta$ implies $|u - v| < F(h)$, and [corollary 6.5](#) applies. Conversely, suppose the universal implication holds but $F(h) < \delta$. Take

$$u = 0, \quad v = F(h).$$

Their input separation is $F(h) < \delta$, whereas the inverse identities give

$$|\overline{G}(0) - \overline{G}(F(h))| = |0 - h| = h,$$

contradicting the asserted strict output bound. Notice that no sign hypothesis on δ is needed: for $\delta \leq 0$ the left implication is vacuous and the right inequality follows from $F(h) \geq 0$. $\square$

7 Effective uniform continuity

7.1 A closed primitive-recursive modulus

Combining the exact inverse modulus with the box event gives the promised fully explicit implication.

Theorem 7.1 (Closed dyadic modulus). *For every $r \geq 0$ and all $u, v \in \mathbb{R}$,*

$$|u - v| < \Delta_r \implies |\overline{G}(u) - \overline{G}(v)| < 2^{-r}, \quad (7.1)$$

where Δ_r is given by (5.1).

Proof. By [lemma 5.1](#), $\Delta_r \leq F(2^{-r})$. Apply [corollary 6.5](#) with $h = 2^{-r}$. $\square$

The exact statement (7.1) is also formalized as `Fabius.abs_fabiusInv_sub_lt_inverse_two_pow_of_lt_deltaDenominator`. The stronger factorial-denominator version is `Fabius.abs_fabiusInv_sub_lt_inverse_two_pow_of_lt_factorialDenominator`. The formal API also closes both endpoints:

$$|u - v| \leq \Delta_r \implies |\overline{G}(u) - \overline{G}(v)| \leq 2^{-r},$$

with declaration `Fabius.abs_fabiusInv_sub_le_inverse_two_pow_of_le_deltaDenominator`; `Fabius.abs_fabiusInv_sub_le_inverse_two_pow_of_le_factorialDenominator` is its stronger factorial-denominator companion. Thus the strict and closed input/output boundaries are separately available rather than recovered by an unstated limiting argument. Their common proof is direct. Let d be either $d_{\text{fac}}(r)$ or $d_{\Delta}(r)$. The recurrence argument above gives $d^{-1} \leq F(2^{-r})$. Hence

$$|u - v| < d^{-1} \implies |u - v| < F(2^{-r}) \implies |\overline{G}(u) - \overline{G}(v)| < 2^{-r}$$

by strict effective injectivity. Replacing both strict inequalities by weak ones and using its closed-threshold companion proves the two closed versions. This proves all four formal dyadic implications without a limiting argument; only the choice of denominator distinguishes the factorial and Delta forms.

Theorem 7.2 (Reciprocal modulus required by the definition). *For $n \geq 1$, let $r(n)$ be the least positive integer with $2^{r(n)} > n$ and put*

$$d(n) = 2^{r(n)(r(n)+1)/2} (r(n) + 1)^{r(n)+1}, \quad d(0) = 1. \quad (7.2)$$

Then d is recursive and satisfies (3.2) for $\overline{G}$. Thus $\overline{G}$ and the restricted inverse G are effectively uniformly continuous.

Proof. The function $r(n)$ can be found by bounded integer search, or equivalently by computing the binary length of n ; hence it is recursive. Exponentiation, multiplication, and composition preserve recursiveness, so d is recursive. By construction,

$$\frac{1}{d(n)} = \Delta_{r(n)} \quad \text{and} \quad 2^{-r(n)} < \frac{1}{n}.$$

If $|u - v| < 1/d(n)$, [theorem 7.1](#) gives $|\overline{G}(u) - \overline{G}(v)| < 2^{-r(n)} < 1/n$. □

Remark 7.3 (An even simpler, much larger modulus). Taking $r = n$ gives the one-line primitive-recursive choice

$$d_0(n) = 2^{n(n+1)/2} (n+1)^{n+1} \quad (n \geq 1),$$

because $2^{-n} < 1/n$. The logarithmic choice $r(n)$ in [theorem 7.2](#) is far smaller and has the correct leading complexity scale.

The Lean development uses the stronger one-line witness

$$d_{\text{fac}}(n) = 2^{\binom{n+1}{2}} (n+1)!,$$

which is no larger than $d_0(n)$. The theorem `Fabius.fabiusInv_effectivelyUniformContinuous` proves `EffectivelyUniformContinuous` for the totalized inverse as follows. The recurrence above makes d_{fac} primitive recursive and positive. For $n \geq 1$, the strict factorial implication gives

$$|u - v| < \frac{1}{d_{\text{fac}}(n)} \implies |\overline{G}(u) - \overline{G}(v)| < 2^{-n}.$$

The elementary inequality $n < 2^n$ between positive integers reverses under reciprocation, so $2^{-n} < 1/n$. Chaining these inequalities is precisely the reciprocal convention in [definition 3.2](#); $d_{\text{fac}}(0) = 1$ supplies the harmless zeroth value. Thus effective uniform continuity itself is formalized. The smaller logarithmic choice $r(n)$ in [theorem 7.2](#) is still only packaged in the report.

7.2 The exact-dyadic repository modulus

Put $h_n := 2^{-r(n)}$. The corpus proves that every $F(h_n)$ is an exactly computable positive rational. Therefore one can replace the elementary lower bound $\Delta_{r(n)}$ by the exact endpoint mass while retaining an exact dyadic output proxy.

Proposition 7.4 (Repository-native exact-dyadic modulus). *Let*

$$d_*(n) := \left\lceil \frac{1}{F(h_n)} \right\rceil \quad (n \geq 1), \quad h_n = 2^{-r(n)}, \quad d_*(0) := 1. \quad (7.3)$$

Then d_* is recursive and

$$|u - v| < \frac{1}{d_*(n)} \implies |\overline{G}(u) - \overline{G}(v)| < h_n < \frac{1}{n}. \quad (7.4)$$

Moreover, $d_*(n)$ is the least positive integer d satisfying

$$\frac{1}{d} \leq F(h_n).$$

Equivalently, it is the least reciprocal denominator certified by the exact inverse-modulus law for the fixed dyadic output target h_n . It is not claimed to be the least integer modulus for the weaker target $1/n$.

Proof. The exact dyadic evaluator computes the positive rational $F(h_n)$, and rational reciprocal and ceiling are computable integer operations. By the defining property of the ceiling,

$$\frac{1}{d_*(n)} \leq F(h_n).$$

The strict input hypothesis and [corollary 6.5](#) therefore give output separation strictly below h_n , while the definition of $r(n)$ gives $h_n < 1/n$.

For the qualified minimality statement, let $1 \leq d < d_*(n)$. Then

$$d < \frac{1}{F(h_n)}, \quad \text{hence} \quad \frac{1}{d} > F(h_n).$$

The endpoint pair $u = 0, v = F(h_n)$ has input separation strictly below $1/d$ and output separation exactly h_n . Thus d cannot certify the stricter conclusion $|\overline{G}(u) - \overline{G}(v)| < h_n$. $\square$

Remark 7.5 (The genuine integer optimum for the $1/n$ target). The exact inverse-modulus law shows that a positive integer d has the universal property

$$|u - v| < \frac{1}{d} \implies |\overline{G}(u) - \overline{G}(v)| < \frac{1}{n}$$

if and only if

$$\frac{1}{d} \leq F(1/n).$$

Consequently the mathematically least such denominator is

$$d_{\text{opt}}(n) = \left\lceil \frac{1}{F(1/n)} \right\rceil.$$

The present report does not prove that $n \mapsto d_{\text{opt}}(n)$ is recursive: computability of the real number $F(1/n)$ alone does not make the discontinuous ceiling operation effective at an unknown integer boundary. The dyadic quantity d_* avoids that issue because $F(h_n)$ is an exact rational.

The distinction is already visible at $n = 1$. Here $r(1) = 1$, $F(1/2) = 1/2$, and hence $d_*(1) = 2$, whereas $d = 1$ already satisfies the required implication with output tolerance 1.

The exact-ceiling construction d_* , including its recursion and qualified denominator minimality, has not yet been translated into Lean. The current effective-continuity theorem deliberately avoids the ceiling by using the explicit factorial denominator at order n .

7.3 Binary conditioning is quadratic

The exact modulus permits a sharp, representation-independent measurement of how many input bits are intrinsically needed near the flat endpoint.

Definition 7.6 (Optimal binary modulus index). For $p \geq 1$, define

$$\mu(p) := \min \{m \in \mathbb{N} : 2^{-m} \leq F(2^{-p})\} = \left\lceil \log_2 \frac{1}{F(2^{-p})} \right\rceil. \quad (7.5)$$

Then

$$|u - v| < 2^{-\mu(p)} \implies |\overline{G}(u) - \overline{G}(v)| < 2^{-p}, \quad (7.6)$$

and no smaller m has this universal property.

The lower event bound already gives an upper bound for $\mu(p)$. A matching lower bound follows from endpoint flatness.

Lemma 7.7 (Elementary flatness upper bound). For every integer $q \geq 1$ and $0 \leq x \leq 2^{-q}$,

$$F(x) \leq \frac{2^{q(q+1)/2}}{q!} x^q. \quad (7.7)$$

In particular,

$$F(2^{-p}) \leq \frac{2^{-p(p-1)/2}}{p!}. \quad (7.8)$$

Proof. Repeated differentiation of $F'(x) = 2F(2x)$ gives

$$F^{(q)}(x) = 2^{q(q+1)/2} F(2^q x) \quad (0 \leq x \leq 2^{-q}). \quad (7.9)$$

All derivatives of F vanish at 0, and $0 \leq F \leq 1$. Taylor's formula in integral form therefore gives

$$\begin{aligned} F(x) &= \frac{1}{(q-1)!} \int_0^x (x-t)^{q-1} F^{(q)}(t) dt \\ &\leq \frac{2^{q(q+1)/2}}{(q-1)!} \int_0^x (x-t)^{q-1} dt = \frac{2^{q(q+1)/2}}{q!} x^q. \end{aligned}$$

Set $q = p$ and $x = 2^{-p}$ to obtain (7.8). $\square$

Theorem 7.8 (Asymptotically sharp input-bit requirement). For every $p \geq 1$,

$$\left\lceil \frac{p(p-1)}{2} + \log_2(p!) \right\rceil \leq \mu(p) \quad (7.10)$$

$$\leq \left\lceil \frac{p(p+1)}{2} + (p+1) \log_2(p+1) \right\rceil. \quad (7.11)$$

Consequently

$$\boxed{\mu(p) = \frac{p^2}{2} + p \log_2 p + \mathcal{O}(p).} \quad (7.12)$$

Proof. The flatness bound (7.8) yields

$$\log_2 \frac{1}{F(2^{-p})} \geq \frac{p(p-1)}{2} + \log_2(p!),$$

while lemma 5.1 yields

$$\log_2 \frac{1}{F(2^{-p})} \leq \frac{p(p+1)}{2} + (p+1) \log_2(p+1).$$

Take ceilings and use (7.5). Stirling's formula $\log_2(p!) = p \log_2 p - (\log_2 e)p + \mathcal{O}(\log p)$ makes both bounds equal to $p^2/2 + p \log_2 p + \mathcal{O}(p)$. $\square$

Remark 7.9 (Intrinsic endpoint cost). The quadratic precision loss is not an artifact of the spline evaluator or of bisection. The two admissible inputs 0 and $F(2^{-p})$ have outputs separated by 2^{-p} . Any uniform oracle algorithm that promises p output bits must, in the worst endpoint case, obtain enough input information to distinguish those values. The exact modulus quantifies that information barrier.

Corollary 7.10 (Growth of the reciprocal modulus). *For $n \rightarrow \infty$, an asymptotically scale-optimal reciprocal modulus has*

$$\log_2 d(n) = \frac{1}{2}(\log_2 n)^2 + (\log_2 n) \log_2 \log_2 n + \mathcal{O}(\log n). \quad (7.13)$$

Equivalently,

$$d(n) = n^{\frac{1}{2} \log_2 n + \log_2 \log_2 n + \mathcal{O}(1)}. \quad (7.14)$$

Proof. Use $p = r(n) = \log_2 n + \mathcal{O}(1)$ in [theorem 7.8](#). The elementary modulus of [theorem 7.2](#) obeys the same two leading orders directly. $\square$

8 Sequential computability by tolerant bisection

8.1 Why ordinary bisection is insufficient

Suppose y is given only by a computable name. At a dyadic midpoint c , one can approximate both $F(c)$ and y , but in general no algorithm can decide whether the two represented reals are exactly equal. An implementation that loops until it proves either $F(c) < y$ or $F(c) > y$ can therefore fail to terminate when $y = F(c)$.

The inverse modulus provides the missing third branch. If the approximations are too close to certify a sign, then $F(c)$ is close to y ; effective injectivity says that c is close to $G(y)$, so the algorithm may return it.

8.2 The algorithm on a computable sequence

Let (y_i) be a computable sequence in $[0, 1]$. Fix a naming algorithm $q(i, s) \in \mathbb{Q}$ with

$$|q(i, s) - y_i| \leq 2^{-s}. \quad (8.1)$$

We describe an algorithm which, from (i, p) , returns a dyadic $z(i, p)$ with

$$|z(i, p) - G(y_i)| < 2^{-p}. \quad (8.2)$$

Algorithm 8.1 (Tolerant certified bisection). Given an index i and output precision p :

1. Set

$$r = p + 2, \quad h = 2^{-r}, \quad \delta = \Delta_r.$$

Choose integers s, N effectively so that

$$2^{-s} \leq \delta/8, \quad 2^{-N} \leq \delta/8. \quad (8.3)$$

Compute the rational $\tilde{y} = q(i, s)$.

2. Initialize the certified bracket $[a, b] = [0, 1]$.

3. Repeat at most $p + 2$ times:

- (a) Put $c = (a + b)/2$ and evaluate the exact rational $S_N(c)$ by [\(4.9\)](#).

- (b) Form the rational approximate difference

$$\tilde{D} = S_N(c) - \tilde{y}. \quad (8.4)$$

- (c) If $\tilde{D} > \delta/2$, replace b by c .
- (d) If $\tilde{D} < -\delta/2$, replace a by c .
- (e) Otherwise return c .

4. If no midpoint was returned, return $(a + b)/2$ after the final update.

Every comparison in [algorithm 8.1](#) is between exact rationals. The order N can be extremely large, but it is obtained by a finite integer calculation and the spline sum is finite. Hence the procedure is an algorithm in the computability-theoretic sense.

8.3 Correctness of every branch

Lemma 8.2 (Difference certificate). *At every midpoint c in [algorithm 8.1](#),*

$$\left| \tilde{D} - (F(c) - y_i) \right| \leq \delta/4. \quad (8.5)$$

Proof. By [theorem 4.3](#) and (8.3), $|S_N(c) - F(c)| \leq \delta/8$. By (8.1) and the same choice of s , $|\tilde{y} - y_i| \leq \delta/8$. The triangle inequality gives (8.5). $\square$

Lemma 8.3 (Safe left/right updates). *If $\tilde{D} > \delta/2$, then $G(y_i) < c$. If $\tilde{D} < -\delta/2$, then $c < G(y_i)$.*

Proof. In the first case, [lemma 8.2](#) gives

$$F(c) - y_i \geq \tilde{D} - \delta/4 > \delta/4 > 0.$$

Since F is strictly increasing, $c > G(y_i)$. The second case is symmetric. Thus every update preserves the invariant

$$G(y_i) \in [a, b]. \quad (8.6)$$

$\square$

Lemma 8.4 (The inconclusive branch is a success branch). *If $|\tilde{D}| \leq \delta/2$, then*

$$|c - G(y_i)| < h = 2^{-r} < 2^{-p}. \quad (8.7)$$

Proof. By [lemma 8.2](#),

$$|F(c) - y_i| \leq |\tilde{D}| + \delta/4 \leq 3\delta/4 < \delta.$$

The choice $\delta = \Delta_r$ and [theorem 7.1](#), applied to the input pair $F(c), y_i$, give

$$|G(F(c)) - G(y_i)| < 2^{-r}.$$

Because $c \in [0, 1]$, $G(F(c)) = c$. Finally, $r = p + 2$. $\square$

Theorem 8.5 (Correctness and uniformity of tolerant bisection). *Algorithm 8.1 terminates and returns a dyadic $z(i, p)$ satisfying (8.2). The map $(i, p) \mapsto z(i, p)$ is computable whenever the name $(i, s) \mapsto q(i, s)$ is computable.*

Proof. The loop has a fixed finite bound. If it returns through the inconclusive branch, [lemma 8.4](#) proves the error estimate. Otherwise [lemma 8.3](#) preserves the bracket invariant through $p + 2$ bisections. Its final width is $2^{-(p+2)}$, so the final midpoint is at distance at most $2^{-(p+3)} < 2^{-p}$ from $G(y_i)$.

All quantities r, h, δ, s, N are computable from p . Midpoints are dyadic; $S_N(c)$ is an exact finite rational expression; and every branch is a rational comparison. The only access to the sequence is the single call to its naming algorithm. Therefore the construction is uniform in i and p . $\square$

Corollary 8.6 (Sequential computability on the unit interval). *The inverse $G : [0, 1] \rightarrow [0, 1]$ is sequentially computable.*

Proof. Given any computable sequence (y_i) in $[0, 1]$, the output of [algorithm 8.1](#) is a computable fast dyadic name for the sequence $(G(y_i))$. $\square$

8.4 Clamping and the totaled inverse

Lemma 8.7 (Computability of clamping). *If (y_i) is a computable real sequence, then $(\text{clamp}(y_i))$ is a computable sequence.*

Proof. Clamping is 1-Lipschitz. If $q(i, p)$ approximates y_i within 2^{-p} , then the exactly rational quantity $\text{clamp}(q(i, p))$ approximates $\text{clamp}(y_i)$ within the same error. The rational min/max operations are decidable and computable. $\square$

Corollary 8.8 (Sequential computability of the totaled inverse). *The totaled inverse $\overline{G} : \mathbb{R} \rightarrow \mathbb{R}$ is sequentially computable.*

Proof. Given a computable sequence (y_i) , first compute the clamped sequence by [lemma 8.7](#), then apply [corollary 8.6](#). By (1.2), the result is $(\overline{G}(y_i))$. $\square$

8.5 Completion of the main theorem

Proof of theorem 1.1. Sequential computability is [corollaries 8.6](#) and [8.8](#). Effective uniform continuity with the explicit recursive modulus is [theorem 7.2](#). These are exactly the two clauses of [definition 3.3](#). $\square$

8.6 An abstract inverse-computability principle

The Fabius proof illustrates a reusable theorem. Stating it clarifies which properties are genuinely needed and which belong only to this example.

Theorem 8.9 (Effective inversion on a computable compact interval). *Let $f : [0, 1] \rightarrow [0, 1]$ be a strictly increasing homeomorphism. Suppose:*

1. *$f(c)$ can be approximated uniformly and effectively at every dyadic rational c ;*
2. *there is a computable positive rational sequence (α_p) such that for all $0 \leq x \leq 1 - 2^{-p}$,*

$$f(x + 2^{-p}) - f(x) \geq \alpha_p. \quad (8.8)$$

Then f^{-1} is sequentially computable and effectively uniformly continuous. A tolerant-bisection algorithm is obtained by replacing Δ_r with α_r in [algorithm 8.1](#).

Proof. Condition (8.8) implies $|u - v| < \alpha_p \Rightarrow |f^{-1}(u) - f^{-1}(v)| < 2^{-p}$ by the same contrapositive argument as [corollary 6.5](#). This gives a binary recursive modulus. The proof of [theorem 8.5](#) uses only this implication, dyadic evaluation of f , and strict monotonicity, so it carries over verbatim. $\square$

For the Fabius function, [theorem 6.1](#) identifies the best possible choice $\alpha_p = F(2^{-p})$; its minimizing intervals are exactly the left and right endpoint intervals. The box event in [lemma 5.1](#) supplies a closed lower substitute requiring no invocation of the exact dyadic evaluator.

9 Analytic consequences of the exact modulus

9.1 Uniform continuity without any Hölder exponent

Effective uniform continuity is sometimes mentally associated with a power-law modulus. The inverse Fabius function is a clean counterexample to that intuition.

Theorem 9.1 (No positive Hölder exponent at the endpoints). *For every $\alpha > 0$ and every $C > 0$, there are arbitrarily small $y > 0$ such that*

$$G(y) > Cy^\alpha. \quad (9.1)$$

By reflection, there are arbitrarily small $1 - y > 0$ such that

$$1 - G(y) > C(1 - y)^\alpha. \quad (9.2)$$

Thus neither G nor $\overline{G}$ is Hölder continuous of any positive order at 0 or 1, despite being effectively uniformly continuous.

Proof. Choose an integer q with $q\alpha > 1$. By lemma 7.7, for all sufficiently small $x > 0$, $F(x) \leq C_q x^q$ with a fixed C_q . If $G(y) \leq Cy^\alpha$ held near zero, substitution $y = F(x)$ would give

$$x = G(F(x)) \leq CF(x)^\alpha \leq CC_q^\alpha x^{q\alpha}.$$

After division by x , the right side tends to zero, a contradiction. The right endpoint follows from $G(1 - y) = 1 - G(y)$. $\square$

Corollary 9.2 (The exact modulus is sub-Hölder). *For every $\alpha > 0$,*

$$\frac{\omega_G(\eta)}{\eta^\alpha} = \frac{G(\eta)}{\eta^\alpha} \longrightarrow +\infty \quad (\eta \downarrow 0). \quad (9.3)$$

Proof. Choose an integer q with $q\alpha > 1$ and write $x = G(\eta)$, so that $\eta = F(x)$ and $x \downarrow 0$ with $\eta \downarrow 0$. By lemma 7.7, for all sufficiently small x , $F(x) \leq C_q x^q$. Hence

$$\frac{G(\eta)}{\eta^\alpha} = \frac{x}{F(x)^\alpha} \geq C_q^{-\alpha} x^{1-q\alpha} \longrightarrow +\infty.$$

The equality with ω_G is theorem 6.4. $\square$

The endpoint growth used here is also formalized, independently of the new exact-modulus module, in `FabiusInverseAsymptotic.lean`. The declarations `Fabius.rpow_isLittleO_fabiusInv_at_zero_right` and `Fabius.tendsto_fabiusInv_div_rpow_atTop_at_zero_right` give the sub-Hölder divergence, while `Fabius.fabiusInv_not_isBigO_rpow_at_zero_right` and `Fabius.not_exists_fabiusInv_le_const_mul_rpow_near_zero` give the corresponding impossibility of a positive power-law upper bound. These are analytic inputs already present in the repository, not consequences claimed to have been newly proved by `InverseModulus.lean`.

9.2 Endpoint asymptotics become modulus asymptotics

The inverse endpoint reports in the corpus give much more than computability [8]. Let

$$T = \log(1/\eta), \quad L = \log 2, \quad \rho = \sqrt{\frac{2T}{L}}. \quad (9.4)$$

Suppressing the explicitly known first log-periodic correction, their leading equivalent is

$$G(\eta) \sim G_0(\eta) := \frac{\sqrt{T}}{e\sqrt{L}} \exp(-\sqrt{2LT}) \quad (\eta \downarrow 0). \quad (9.5)$$

More precisely, the corpus obtains a relative expansion of the form

$$\frac{G(\eta)}{G_0(\eta)} = 1 + \frac{\frac{1}{2} \log \rho + \kappa_0 - \Psi(\theta)}{\rho} + \mathcal{O}\left(\frac{(\log \rho)^2}{\rho^2}\right), \quad (9.6)$$

where Ψ is a nonconstant smooth one-periodic Gamma-zeta fluctuation and θ is the lower-Lambert phase shifted by an explicit constant.

Corollary 9.3 (Sharp asymptotic modulus). *The expressions (9.5) and (9.6) are, without any further inversion, the leading and first-correction asymptotics of the exact continuity modulus of G and $\overline{G}$:*

$$\omega_G(\eta) = \omega_{\overline{G}}(\eta) = G(\eta) \quad (0 \leq \eta \leq 1). \quad (9.7)$$

Every proved all-orders endpoint expansion for the inverse therefore transfers verbatim to an all-orders modulus-of-continuity expansion.

Proof. This is exactly [theorem 6.4](#). □

This observation closes a conceptual loop in the corpus. The lower-Lambert phase and Bell-polynomial inverse transfer were developed to approximate quantiles. The geometric theorem shows that the same quantiles are the optimal global sensitivity profile of the inverse map.

9.3 Derivative blow-up versus computability

On $0 < y < 1$, the repository proves

$$G'(y) = \frac{1}{F'(G(y))} = \frac{1}{2 \operatorname{up}(2G(y) - 1)} > 0. \quad (9.8)$$

The derivative tends to $+\infty$ at both endpoints. There is therefore no global Lipschitz constant and no endpoint condition number bounded in the ordinary differential sense. Nonetheless the exact nonlinear modulus $G(\eta)$ is computable, and tolerant bisection uses precisely that nonlinear conditioning. This is why derivative-based numerical intuition alone is not a substitute for a computable-analysis proof.

10 Numerical and exact-rational checks

The accompanying script `inverse_fabius_computability_experiments.py` is a reproducibility supplement. The proofs do not depend on it. It uses `fractions.Fraction` for the Thue–Morse splines and asserts every inequality it checks.

10.1 Endpoint mass bounds

Define

$$A_r := \frac{r(r-1)}{2} + \log_2(r!), \quad (10.1)$$

$$B_r := \frac{r(r+1)}{2} + (r+1) \log_2(r+1). \quad (10.2)$$

Then [lemmas 5.1](#) and [7.7](#) say

$$A_r \leq \log_2 \frac{1}{F(2^{-r})} \leq B_r. \quad (10.3)$$

The following values show the quadratic squeezing.

The numerical width in [table 1](#) is not a defect in the computability proof. Its order is only linear in r after the common $p^2/2 + p \log_2 p$ terms are removed. Exact dyadic arithmetic or the sharp Lambert expansion can replace the bracket whenever a tight practical modulus is desired.

r	Δ_r	$B_r = -\log_2 \Delta_r$	flatness upper bound	A_r
1	$1.25000000 \cdot 10^{-1}$	3.000000	1.00000000	0.000000
2	$4.62962963 \cdot 10^{-3}$	7.754888	$2.50000000 \cdot 10^{-1}$	2.000000
3	$6.10351563 \cdot 10^{-5}$	14.000000	$2.08333333 \cdot 10^{-2}$	5.584963
4	$3.12500000 \cdot 10^{-7}$	21.609640	$6.51041667 \cdot 10^{-4}$	10.584963
5	$6.54097611 \cdot 10^{-10}$	30.509775	$8.13802083 \cdot 10^{-6}$	16.906891
6	$5.79006996 \cdot 10^{-13}$	40.651484	$4.23855252 \cdot 10^{-8}$	24.491853
7	$2.22044605 \cdot 10^{-16}$	52.000000	$9.46105473 \cdot 10^{-11}$	33.299208
8	$3.75610368 \cdot 10^{-20}$	64.529325	$9.23931126 \cdot 10^{-14}$	43.299208
9	$2.84217094 \cdot 10^{-24}$	78.219281	$4.01011773 \cdot 10^{-17}$	54.469133
10	$9.72815993 \cdot 10^{-29}$	93.053748	$7.83226120 \cdot 10^{-21}$	66.791061

Table 1: Rigorous lower and upper bounds for $F(2^{-r})$. The two logarithmic columns bracket the optimal binary modulus index.

N	x	$S_N(x)$	$ S_N(x) - F(x) $	certified 2^{-N}
4	1/2	$5.0000000000000000 \cdot 10^{-1}$	0	$6.25 \cdot 10^{-2}$
4	1/4	$6.9010416666666671 \cdot 10^{-2}$	$4.34027778 \cdot 10^{-4}$	$6.25 \cdot 10^{-2}$
4	1/8	$3.2552083333333335 \cdot 10^{-3}$	$2.17013889 \cdot 10^{-4}$	$6.25 \cdot 10^{-2}$
8	1/2	$5.0000000000000000 \cdot 10^{-1}$	0	$3.90625 \cdot 10^{-3}$
8	1/4	$6.9442749023437500 \cdot 10^{-2}$	$1.69542101 \cdot 10^{-6}$	$3.90625 \cdot 10^{-3}$
8	1/8	$3.4713745117187500 \cdot 10^{-3}$	$8.47710503 \cdot 10^{-7}$	$3.90625 \cdot 10^{-3}$
12	1/2	$5.0000000000000000 \cdot 10^{-1}$	0	$2.44140625 \cdot 10^{-4}$
12	1/4	$6.9444437821706131 \cdot 10^{-2}$	$6.62273831 \cdot 10^{-9}$	$2.44140625 \cdot 10^{-4}$
12	1/8	$3.4722189108530679 \cdot 10^{-3}$	$3.31136915 \cdot 10^{-9}$	$2.44140625 \cdot 10^{-4}$

Table 2: Exact centered-spline checks. The global certificate is intentionally conservative; the pointwise dyadic errors are much smaller.

10.2 Centered-spline convergence

The exact values

$$F(1/2) = 1/2, \quad F(1/4) = 5/72, \quad F(1/8) = 1/288 \quad (10.4)$$

provide compact checks of (4.9). The script produces the following exact rational errors, displayed decimally.

For $N = 8$, interval length $h = 1/8$, and the dyadic grid $2^{-8}\mathbb{Z}$, the script also verifies exactly that

$$\min_x (S_8(x + 1/8) - S_8(x)) = S_8(1/8) = \frac{455}{131072},$$

with minimizers $x = 0$ and $x = 7/8$. This finite-convolution experiment reproduces the complete continuous minimizer locus in [theorem 6.1](#); the theorem itself is the strict unimodality argument, not a grid inference.

10.3 What the code deliberately does not claim

The naive rational formula scales exponentially with N . The script uses small orders to check identities and error bounds, not to compete with the repository’s exact dyadic evaluator. Likewise, its short bisection transcript illustrates certified comparisons at a modest scale; sequential computability is proved by the abstract finite algorithm, regardless of practical runtime.

11 Connections to the wider Fabius–Rvachev corpus

11.1 Thue–Morse inclusion–exclusion

The forward oracle is not an unrelated approximation scheme. The binary subset map in [proposition 4.1](#) makes the Thue–Morse sign $(-1)^{s_2(j)}$ the exact inclusion–exclusion coefficient of the finite uniform convolution. Thus the name-transformer for G ultimately rests on the same cancellation mechanism that drives the repository’s exact dyadic formulae and discrete-limit identities.

The centered shift $(2j + 1)/2^{N+1}$ is equally structural: it replaces the unknown tail by its mean and halves the elementary uniform error from the uncentered bound 2^{1-N} to 2^{-N} . In computational terms, this one-bit improvement is free.

11.2 q -binomial coefficients and accelerated forward oracles

The finite Thue–Morse sums admit Gaussian-binomial reorganizations; the arithmetic background is developed in [7, 8]. The inverse/sampling corpus uses $q = 1/4$ Richardson filters on the geometric error scale of centered finite approximants; the current snapshot also adds a quarter-base q -Pochhammer factorization of the Rvachev sinc product. None of these identities is needed for mere computability, because [theorem 4.3](#) already gives an effective rate. They are relevant to complexity: a certified accelerated approximation to $F(c)$ can be dropped into [algorithm 8.1](#) without changing one line of its logical proof, provided the accelerated remainder is enclosed rigorously.

11.3 Infinite sinc products and alternative evaluation

Equation (4.4) gives the Fourier transform of up , in the classical infinite-convolution tradition of [5, 6]. Fourier inversion therefore offers an independent evaluator for F or its density. The corpus develops strong decay estimates, Euler/canonical products, and Poisson-summation formulae. A computability proof based on this route would need an explicit truncation error for both the frequency integral and the infinite product. Once those are supplied, tolerant bisection is again unchanged.

This modularity is useful for cross-certification: finite splines work in physical space with exact rationals; sinc products work in frequency space with analytic tail bounds. Agreement of independently certified enclosures is a stronger implementation check than agreement of unproved decimal approximations.

11.4 Lambert W , Bell polynomials, and Bernoulli data

The flat endpoint makes the inverse poorly conditioned in any power scale. The lower branch W_{-1} naturally solves the saddle equation controlling $F(x)$ as $x \downarrow 0$. The repository then uses Bell-polynomial reversion to transport all-orders forward expansions to $G(y)$ [8]; Bernoulli numbers and polynomials enter through logarithms of the Laplace/sinc products and through formal composition.

The exact identity $\omega_G(\eta) = G(\eta)$ gives these expansions a second interpretation: they are not only quantile asymptotics but optimal continuity- modulus asymptotics. In particular, the nonconstant periodic Gamma-zeta term cannot be discarded from a genuinely sharp effective modulus.

11.5 Legendre and Lagrange representations

The corpus expands up in even Legendre polynomials and represents polynomials by finite combinations of shifted atomic functions. Such bases can provide spectral or interpolation-based forward oracles. For a formal computability theorem, however, convergence alone is insufficient: one needs a recursive tail majorant uniform in the evaluation point. Deriving sharp certified Legendre-tail bounds and comparing their bit complexity with the Thue-Morse spline is a natural next step.

11.6 Total positivity, unimodality, and interval masses

The least-mass lower bound uses only symmetric unimodality; its exact endpoint-only equality locus uses strict radial unimodality. Several corpus reports investigate still stronger total-positivity and variation-diminishing properties. If a stronger kernel theorem is established, it may control not only one interval mass but higher finite differences and multidimensional quantile maps. The computability proof therefore isolates a low-order shadow of a potentially broader shape theory.

12 Lean formalization status and remaining roadmap

12.1 Why the present noncomputable definition is not an obstacle

The repository defines `fabiusInv` by inverting an order isomorphism and then applying `IccExtend`. Lean marks that definition `noncomputable`, because the inverse equivalence is constructed by classical existence machinery. This is a statement about the chosen term in Lean’s programming language, not a theorem that the represented real function has no algorithm.

Computable analysis deliberately separates these two levels. An extensional real function may be introduced by a classical characterization and later be shown to possess a computable *realizer*: a program transforming names of inputs into names of outputs. [Algorithm 8.1](#) is exactly such a realizer. Its correctness is proved by showing that its output denotes the same extensional function as `fabiusInv`.

This distinction already occurs on the forward side. The analytic canonical Fabius function is defined abstractly, whereas `FabiusComputableSpline.lean` supplies a primitive-recursive evaluator and proves that it realizes the same function. The inverse formalization can follow the same architecture.

12.2 The completed structural inverse-modulus layer

The structural part of [section 6](#) now lives in

Analysis/FabiusFunction/Lean/FabiusFunction/InverseModulus.lean.

It imports `FabiusInverse.lean` and deliberately remains independent of the heavier random-series and computable-analysis interfaces. All thirty-two public declarations (one definition and thirty-one theorems) were compiler-checked as one focused target; each theorem uses only the standard logical principles already present in its dependencies. The declaration-level correspondence is as follows.

Report object or claim	Lean declaration
The fixed-length increment $M_h(x) = F(x + h) - F(x)$	<code>Fabius.fabiusIntervalMass.</code>
Reflection $M_h(1 - h - x) = M_h(x)$	<code>Fabius.fabiusIntervalMass_reflect.</code>
The zero tails $M_h(x) = 0$ when $x + h \leq 0$ or $1 \leq x$, for $h \geq 0$	<code>Fabius.fabiusIntervalMass_eq_zero_of_add_non_pos</code> and <code>Fabius.fabiusIntervalMass_eq_zero_of_one_le.</code>
Global left-half monotonicity and right-half antitonicity	<code>Fabius.monotoneOn_fabiusIntervalMass_firstHalf</code> and <code>Fabius.antitoneOn_fabiusIntervalMass_secondHalf</code> . These are the weak half-line statements that include the clamped zero tails.
Maximal strict branches $[-h, (1 - h)/2]$ and $[(1 - h)/2, 1]$ for $0 < h \leq 1$	<code>Fabius.strictMonoOn_fabiusIntervalMass_firstHalf</code> and <code>Fabius.strictAntiOn_fabiusIntervalMass_secondHalf</code> .
Least endpoint increment $F(b - a) \leq F(b) - F(a)$	<code>Fabius.fabiusReal_sub_le_sub.</code>
Strict interior increment $F(h) < M_h(x)$ and exact endpoint starts	<code>Fabius.fabiusReal_lt_fabiusIntervalMass_of_mem_Ioo</code> and <code>Fabius.fabiusIntervalMass_eq_fabiusReal_iff.</code>
Strict interior interval inequality and complete equality law (6.4)	<code>Fabius.fabiusReal_sub_lt_sub</code> and <code>Fabius.fabiusReal_sub_eq_sub_iff.</code>
Constrained forward superadditivity (6.7)	<code>Fabius.fabiusReal_add_le</code> , with equality classification (6.8) in <code>Fabius.fabiusReal_add_eq_iff.</code>
Ordered inverse gap on $[0, 1]$	<code>Fabius.fabiusInv_sub_le_sub_of_mem_Icc</code> , with exact equality cases (6.15) in <code>Fabius.fabiusInv_sub_eq_sub_iff_of_mem_Icc.</code>
Ordered inverse gap including clamped tails, and its order-free strengthening (6.13)	<code>Fabius.fabiusInv_sub_le_sub_of_le</code> and <code>Fabius.fabiusInv_sub_le_sub.</code>
Global inverse subadditivity (6.14)	<code>Fabius.fabiusInv_add_le.</code>
Absolute and metric pointwise modulus (6.12)	<code>Fabius.abs_fabiusInv_sub_le</code> and <code>Fabius.dist_fabiusInv_le</code> ; the complete unit-input equality locus (6.16) is <code>Fabius.abs_fabiusInv_sub_eq_iff_of_mem_Icc.</code>
The saturation identity $\overline{G}(\min(\eta, 1)) = \overline{G}(\eta)$	<code>Fabius.fabiusInv_min_one.</code>

Report object or claim	Lean declaration
Attained exact totaled modulus and its supremum identity	<code>Fabius.isGreatest_abs_fabiusInv_sub</code> and <code>Fabius.sSup_abs_fabiusInv_sub_eq</code> .
Attained exact modulus with unit-interval inputs, for every $\eta \geq 0$	<code>Fabius.isGreatest_abs_fabiusInv_sub_Icc</code> and <code>Fabius.sSup_abs_fabiusInv_sub_Icc_eq</code> .
Strict and closed effective injectivity	<code>Fabius.abs_fabiusInv_sub_lt_of_abs_sub_lt_fabiusReal</code> and <code>Fabius.abs_fabiusInv_sub_le_of_abs_sub_le_fabiusReal</code> .
Contrapositive separation (6.21)	<code>Fabius.fabiusReal_le_abs_sub_of_le_abs_fabiusInv_sub</code> .
Exact strict-threshold biconditional (6.22)	<code>Fabius.forall_abs_fabiusInv_sub_lt_iff</code> .

The formal least-mass API deliberately separates four reusable layers: the global weak shape, the maximal strict branches, the lower bound, and the complete equality locus. The inverse equality theorems remain restricted to unit inputs because the clamped tails add straddling-tail equality cases and make the endpoint alternatives order-dependent globally. The `IsGreatest` declarations record both the sharp upper bound and an extremizing pair; the `sSup` equalities are immediate corollaries. The totaled formal theorem returns `fabiusInv ... eta`, which is literally the report’s $G(\min(\eta, 1))$ because `fabiusInv` is already clamped.

12.3 The completed effective-uniform-continuity layer

The next formal layer is

```
Analysis/FabiusFunction/Lean/FabiusFunction/FabiusInverseEffectiveContinuity.lean.
```

It imports the forward computability interface, the exact inverse-dyadic recurrence, and `InverseModulus.lean`. Its recurrence proof is stronger than the report’s box estimate, while its final reciprocal modulus is simpler and larger than the logarithmic- $r(n)$ modulus. The following table exhausts all fourteen public declarations in the module (two definitions and twelve theorems); private recurrence and primitive-recursion helpers are intentionally omitted.

Lean declaration	Report object or claim
<code>Fabius.inverseFabiusFactorialDenominator</code>	The recursively executable factorial denominator $d_{\text{fac}}(r)$.
<code>Fabius.inverseFabiusFactorialDenominator_eq</code>	The closed form $d_{\text{fac}}(r) = 2^{\binom{r+1}{2}}(r+1)!$.
<code>Fabius.inverseFabiusFactorialDenominator_primrec</code>	Primitive recursiveness of d_{fac} .
<code>Fabius.inverseFabiusDeltaDenominator</code>	The report’s elementary denominator $d_{\Delta}(r) = 2^{\binom{r+1}{2}}(r+1)^{r+1} = \Delta_r^{-1}$.
<code>Fabius.inverseFabiusDeltaDenominator_primrec</code>	Primitive recursiveness of d_{Δ} .

Lean declaration	Report object or claim
<code>Fabius.fabiusReal_inverse_two_pow_one_term_lower_bound</code>	For $r > 0$, the sharper unrounded zeroth-term recurrence bound with denominator $2^{\binom{r}{2}}(r+1)!(2^r-1)$.
<code>Fabius.inv_inverseFabiusFactorialDenominator_le_fabiusReal</code>	The first inequality in (5.3), including $r = 0$.
<code>Fabius.inverseFabiusFactorialDenominator_le_deltaDenominator</code>	The denominator comparison $d_{\text{fac}}(r) \leq d_{\Delta}(r)$, from $(r+1)! \leq (r+1)^{r+1}$.
<code>Fabius.inv_inverseFabiusDeltaDenominator_le_fabiusReal</code>	The exact numerical conclusion $\Delta_r \leq F(2^{-r})$ of lemma 5.1 , proved through the recurrence rather than its box event.
<code>Fabius.abs_fabiusInv_sub_lt_inverse_two_pow_of_1_t_factorialDenominator</code>	The stronger strict dyadic inverse modulus: input distance $< d_{\text{fac}}(r)^{-1}$ forces output distance $< 2^{-r}$.
<code>Fabius.abs_fabiusInv_sub_le_inverse_two_pow_of_1_e_factorialDenominator</code>	The closed factorial companion: input distance $\leq d_{\text{fac}}(r)^{-1}$ forces output distance $\leq 2^{-r}$.
<code>Fabius.abs_fabiusInv_sub_lt_inverse_two_pow_of_1_t_deltaDenominator</code>	The report's exact strict implication (7.1): input distance $< \Delta_r$ forces output distance $< 2^{-r}$.
<code>Fabius.abs_fabiusInv_sub_le_inverse_two_pow_of_1_e_deltaDenominator</code>	The exact closed-boundary companion: input distance $\leq \Delta_r$ forces output distance $\leq 2^{-r}$.
<code>Fabius.fabiusInv_effectivelyUniformContinuous</code>	Effective uniform continuity of the totalized inverse, witnessed by $n \mapsto d_{\text{fac}}(n)$ and hence using the simple scale $r = n$.

This table crosswalks theorem statements, not proof methods. In particular, the declaration with conclusion $\Delta_r \leq F(2^{-r})$ does not construct the random-series event used in [lemma 5.1](#). Conversely, the final theorem establishes the repository predicate `EffectivelyUniformContinuous`; it does not establish a sequential realizer or the two-field `IsComputableRealFunction` structure.

12.4 The remaining name-transformation module

A natural downstream file remains

`Analysis/FabiusFunction/Lean/FabiusFunction/FabiusInverseComputability.lean`.

It should import `FabiusFunction.FabiusInverseEffectiveContinuity`. It no longer needs to reprove interval shape, inverse-gap estimates, dyadic endpoint lower bounds, or effective uniform continuity. Its remaining public surface is the name-transformation layer; the following Lean is schematic rather than a claim about declarations already present.

```

/-- Tolerant bisection transforms every computable input sequence. -/
theorem fabiusInv_sequentiallyComputable :
  SequentiallyComputable (fabiusInv fabius fabius_spec)

```

```

/-- Both Grzegorzczuk clauses, packaged together. -/
theorem fabiusInv_isComputableRealFunction :
  IsComputableRealFunction (fabiusInv fabius fabius_spec)
    
```

12.5 Remaining formalization order

The former structural roadmap steps are complete in `InverseModulus.lean`, and the closed-bound/effective-continuity steps are complete in `FabiusInverseEffectiveContinuity.lean`. The remaining order is:

1. If a probability-level counterpart is desired, formalize the particular box event in (5.2), its containment, and its product probability. This is no longer needed to obtain its endpoint inequality, which is already formalized by recurrence.
2. Formalize the logarithmic scale $r(n)$ and the smaller report modulus from [theorem 7.2](#); this is an efficiency refinement, not a prerequisite for effective uniform continuity.
3. Connect the exact dyadic evaluator to the positive rational $F(2^{-r})$ and package the reciprocal-ceiling modulus, including the qualified fixed-dyadic-target minimality statement of [proposition 7.4](#).
4. Formalize a generic tolerant-bisection lemma for a monotone computable map equipped with a recursive lower-increment bound. The result should not be Fabius-specific.
5. Instantiate the generic lemma with the primitive-recursive centered spline evaluator from `FabiusComputableSpline.lean`.
6. Package the resulting sequential-computability theorem together with the now-formalized effective uniform continuity into the repository's computable-real-function interface.
7. Only after the effective core is complete, formalize the flatness bounds and input-bit asymptotics; they are not logical prerequisites of the computability theorem.

12.6 Reusable generic infrastructure

The abstract [theorem 8.9](#) deserves its own module. In a representation-oriented library it can be stated using an oracle for f and a recursive lower-separation function

$$\alpha(p) > 0, \quad x + 2^{-p} \leq b \implies f(x + 2^{-p}) - f(x) \geq \alpha(p).$$

The conclusion is a computable realizer for the inverse. Tolerant bisection then becomes library infrastructure usable for distribution quantiles, implicit monotone roots, cumulative spectral measures, and monotone solutions of integral equations. The proof is constructive in the strong sense that all precision budgets are explicit.

13 Further deductions, conjectures, and research directions

13.1 A certified practical inverse evaluator

The proof algorithm is intentionally elementary rather than efficient. A practical certified implementation should retain its logical shell while replacing the forward oracle by a portfolio of enclosures:

- exact dyadic evaluation and terminating local Taylor formulae near dyadic anchors;
- centered Thue–Morse splines at moderate depth;
- quarter-base q -Richardson acceleration of finite-prefix errors;

- Fourier/sinc evaluation in regions where frequency-space decay is favorable;
- lower-Lambert endpoint enclosures when the requested target is extremely close to 0 or 1.

Each backend must return a rational interval containing $F(c)$, not merely a floating-point approximation. The bisection controller only asks whether that interval lies wholly above or below the target interval; overlapping intervals trigger refinement or the modulus-based acceptance branch. Thus backend selection is a performance concern, not part of the correctness argument.

Research problem 13.1 (Near-optimal certified evaluation). *Construct and formally verify an inverse evaluator whose oracle-bit consumption is $\mu(p) + \mathcal{O}(\log p)$ for p output bits at the worst endpoint, and whose arithmetic running time is quasi-linear or softly quadratic in the size of the exact integers produced. Separate information-theoretic oracle precision from the cost of evaluating F to that precision.*

13.2 Sharp integer moduli and the log-periodic phase

The real-valued optimal input-bit threshold is

$$\Lambda(p) := \log_2 \frac{1}{F(2^{-p})}, \quad \mu(p) = \lceil \Lambda(p) \rceil.$$

The all-orders endpoint theory expresses $\Lambda(p)$ in a lower-Lambert phase with a nonconstant one-periodic Gamma–zeta contribution and a hierarchy of inverse powers. Passing to the integer ceiling introduces a second, sawtooth nonlinearity. It is therefore natural to study the arithmetic interaction between the smooth periodic saddle phase and fractional parts of its growing nonperiodic carrier.

Conjecture 13.2 (Nontrivial fluctuation of the optimal integer modulus). *After subtracting any fixed finite truncation of the full nonperiodic asymptotic expansion of $\Lambda(p)$, the bounded residual governing $\mu(p)$ does not become eventually constant or eventually periodic in the integer variable p . Its accumulation set is generated jointly by the Gamma–zeta phase and the ceiling sawtooth.*

The conjecture is deliberately qualitative. A stronger equidistribution claim would require information about fractional parts of a nonlinear Lambert-type phase that is not supplied by the present corpus. Exact values $F(2^{-p}) \in \mathbb{Q}$ make the sequence amenable to arbitrary-precision experimentation without numerical ambiguity.

13.3 Base- b atomic distributions

Replace the dyadic random series by

$$X_b = \sum_{k \geq 1} b^{-k} U_k, \quad b \geq 2,$$

with independent uniform coordinates. The finite convolution is still computable, but binary Thue–Morse signs are replaced by base- b subset or digit combinatorics, and the support is $[0, 1/(b-1)]$ before normalization. The least-interval-mass theorem continues to follow whenever the normalized density is symmetric and unimodal. The same proof would then identify the optimal inverse modulus with the quantile itself.

Research problem 13.3 (Uniform computability across the base). *Develop a theorem uniform in the integer parameter b : construct a single algorithm that, given b , a name of y , and a precision p , computes the normalized quantile of X_b . Determine the leading dependence of the optimal input-bit modulus on both b and p , and identify the corresponding base- b log-periodic correction.*

13.4 Certified spectral and polynomial representations

The Legendre, Lagrange, and shifted-up representations in the corpus are mathematically exact, but a representation becomes an algorithm only after a recursive uniform tail bound is attached. This suggests three concrete projects:

1. derive explicit Legendre coefficient envelopes strong enough to certify pointwise and uniform evaluation of up;
2. combine the finite shifted-up representation of polynomials with interval arithmetic to obtain certified local surrogates for F ;
3. compare physical-space spline, Legendre, and sinc-product evaluators by bit complexity, not merely by observed decimal convergence.

A cross-certified implementation could evaluate the same point by two independent representations and intersect the resulting enclosures. This is particularly attractive near transition regions where one representation's natural error bound is pessimistic.

13.5 Higher-dimensional monotone transport

The one-dimensional theorem

$$\min_x \mathbb{P}\{X \in (x, x + h]\} = \mathbb{P}\{X \in (0, h]\}$$

is a sharp anti-concentration lower bound tailored to a symmetric unimodal law. Because the law is atomless, the same identity holds with any choice of open or closed endpoints. In higher dimensions, inverse CDFs are replaced by triangular Rosenblatt maps or optimal-transport maps. Effective invertibility would require computable lower bounds on masses of suitable boxes or sections.

Research problem 13.4 (Atomic transport computability). *For products or coupled variants of Fabius-type laws, determine conditions under which the triangular transport from Lebesgue measure has a computable inverse with an explicit multivariate modulus. Investigate whether total positivity or variation-diminishing properties supply the needed lower section-mass estimates.*

13.6 Branchwise inversion of the signed global extension

The signed global Fabius extension is not globally injective, so it has no single inverse. On every maximal monotonicity interval, however, it has a branch inverse. Local finiteness of its Thue–Morse translate expansion makes forward evaluation computable. The remaining issue is effective branch separation near critical values.

Research problem 13.5 (Computable branch atlas). *Construct an effective enumeration of monotonicity intervals of the signed global extension, together with certified branch ranges and recursive inverse moduli. Determine whether the self-similar translation law permits one finite branch template plus a Thue–Morse address.*

14 Conclusion

The inverse Fabius function is computable in precisely the two-part sense requested. The effective-uniform-continuity clause follows from the exact geometric identity

$$\omega_G(\eta) = G(\min(\eta, 1)),$$

combined with the closed positive lower bound

$$F(2^{-r}) \geq 2^{-r(r+1)/2} (r+1)^{-(r+1)}.$$

The sequential-computability clause follows from a tolerant bisection transformer that uses only finite exact rational operations, a computable name of the target, and certified centered-spline evaluations of F . It never requires deciding equality of computable reals.

The result is stronger than an existence proof. It identifies the *optimal* modulus, explains why inversion loses quadratically many bits at the flat endpoints, and turns the repository’s lower-Lambert quantile asymptotics into sharp continuity-modulus asymptotics. The same proof covers the inverse on $[0, 1]$ and the canonical clamped totalization on all of $\mathbb{R}$.

Conceptually, the decisive bridge is simple: weak symmetric unimodality makes endpoint intervals minimize mass, while strict radial unimodality identifies every equality case. In the Fabius setting that sharpened shape fact connects probability, computable analysis, exact dyadic arithmetic, and inverse asymptotics. It is also the part most likely to generalize beyond the particular atomic function studied here.

The current Lean boundary cuts through the two clauses of computability. `InverseModulus.lean` certifies the exact structural modulus, and `FabiusInverseEffectiveContinuity.lean` certifies the closed dyadic bound and `EffectivelyUniformContinuous`, using a recurrence estimate that is stronger than the report’s box bound. Tolerant bisection, sequential computability, and the final computable-real-function package remain to be formalized; the logarithmic- $r(n)$ and exact-ceiling moduli remain optional sharpenings of the already certified effective-continuity witness.

A Pseudocode with explicit error budgets

The following version makes every tolerance in [algorithm 8.1](#) explicit. The function `TargetName(i, s)` returns a rational within 2^{-s} of the clamped input y_i . The function `CenteredSpline(N, c)` is the exact rational $S_N(c)$ of (4.9).

```

InverseFabiusName(i, p):
  # Requested absolute output error: 2^(-p).
  r      := p + 2
  h      := 2^(-r)
  delta  := 2^(-r*(r+1)/2) * (r+1)^(-(r+1))

  choose s with 2^(-s) <= delta/8
  choose N with 2^(-N) <= delta/8
  yhat := clamp(TargetName(i, s), 0, 1)

  a := 0
  b := 1

  repeat p + 2 times:
    c      := (a + b)/2
    fhat   := CenteredSpline(N, c)
    D      := fhat - yhat

    # |D - (F(c)-y_i)| <= delta/4.
    if D > delta/2:
      b := c                # certified F(c) > y_i
    else if D < -delta/2:
      a := c                # certified F(c) < y_i
    else:
      return c              # |F(c)-y_i| < delta

  return (a + b)/2

```

In the acceptance branch, $|F(c) - y_i| < 3\delta/4 < \delta \leq F(h)$, so [corollary 6.5](#) gives $|c - G(y_i)| <$

$h < 2^{-p}$. If no acceptance occurs, every update is certified and the true inverse remains in $[a, b]$; after $p + 2$ iterations its width is $2^{-(p+2)}$, so the final midpoint is within $2^{-(p+3)} < 2^{-p}$.

The constants have generous slack. They were chosen to make the proof transparent and to absorb representation-dependent rounding. Replacing the three occurrences of $1/8, 1/4, 1/2$ by any fixed compatible margin gives the same computability theorem.

B Proof-dependency and status audit

Claim	Input	Human proof and Lean status
Random-series representation of F	Infinite convolution / probability representation in the repository and classical Fabius theory	Imported; normalization restated in section 4 .
Exact finite CDF (4.7)	Inclusion–exclusion for finite sums of uniforms	Proved in proposition 4.1 .
Uniform centered-spline error 2^{-N}	Tail support and the global density bound $p \leq 2$	Proved in theorem 4.3 ; also formalized forward in the repository.
Strict increase of F	Positivity of finite-convolution densities and a short tail	Proved in proposition 5.2 .
Symmetric strict unimodality of $p = F'$	Differential equation, symmetry, and strict increase of F	Proved in proposition 5.3 from imported Fabius identities.
Least fixed-length increment $F(h)$	Symmetric unimodality of p	Human proof in theorem 6.1 ; reflection, global weak shape, maximal strict branches, the lower bound, and the complete degenerate/endpoint equality locus are formalized in <code>InverseModulus.lean</code> . Exact names appear in section 12 .
Exact inverse modulus $G(\eta)$	Least interval mass and inverse identities	Human proof in theorem 6.4 ; attained maxima and exact suprema are formalized in <code>InverseModulus.lean</code> . Exact names appear in section 12 .
Exact effective-injectivity threshold $F(h)$	Pointwise inverse modulus and inverse identities	Human proofs in corollary 6.5 and theorem 6.6 ; strict, closed, contrapositive, and biconditional forms are formalized in <code>InverseModulus.lean</code> .

Claim	Input	Human proof and Lean status
Closed modulus Δ_r	Explicit positive box event in the random series	The event proof is in lemma 5.1 ; its numerical endpoint inequality and the exact inverse implication of theorem 7.1 are formalized in <code>FabiusInverseEffectiveContinuity.lean</code> through the stronger one-term recurrence route.
Effective uniform continuity	Closed modulus and recursive integer arithmetic	Proved in theorem 7.2 . The repository predicate itself is formalized as <code>Fabius.fabiusInv_effectivelyUniformContinuous</code> , using the simple primitive-recursive factorial denominator at $r = n$; the report's smaller logarithmic- $r(n)$ packaging remains human-only.
Exact dyadic ceiling denominator d_*	Exact rational dyadic evaluation, ceiling arithmetic, and the exact threshold	Human proof in proposition 7.4 ; neither the ceiling construction nor its qualified fixed-target minimality is yet formalized.
Sequential computability	Forward rational oracle, closed inverse modulus, and tolerant bisection	Proved in theorem 8.5 and corollary 8.8 . No tolerant-bisection or sequential-computability declaration is supplied by <code>FabiusInverseEffectiveContinuity.lean</code> .
Optimal bit asymptotic	Closed lower bound plus iterated-flatness upper bound	Proved in theorem 7.8 .
Sharp periodic refinement	Repository's all-orders endpoint inverse theory	Imported and reinterpreted in corollary 9.3 ; not rederived.

This table is also a minimal trust boundary. The main computability theorem does not rely on the conjectures in [section 13](#), on numerical data, or on any asymptotic expansion.

C Corpus crosswalk

Corpus component	Role in the present report
Fabius_Function_and_Rv_achev_Up.tex and archived primary expositions	Normalization, probability law, up-function relation, exact dyadic arithmetic, finite splines, and endpoint asymptotic context.
fabius_lean_walkthrough.tex	Public theorem map; distinction among the bounded CDF, signed extension, and up-function; forward computability and inverse module architecture.
FabiusFunction_Mathematical_Glossary.tex	Repository-specific Grzegorzczuk definitions and terminology for fast names, sequential computability, and effective uniform continuity.
PAPER_COVERAGE.md and the two Arias de Reyna paper transcriptions	Formal/source crosswalk for probability, Fourier, dyadic, moment, and arithmetic results.
FabiusComputability.lean and FabiusComputableSpline.lean	Existing forward 2-Lipschitz theorem, uniform centered-spline certificate, primitive-recursive evaluator, and computable-real-function packaging.
FabiusInverse.lean	Existing clamped inverse, inverse identities, continuity, interior derivatives, curvature, and endpoint steepness.
InverseModulus.lean	Post-snapshot formalization of fixed-length Fabius increments, their global weak and maximal strict shapes, the least endpoint increment and complete equality locus, constrained forward superadditivity with equality cases, local and global inverse-gap inequalities, unit-input inverse-gap rigidity, global inverse subadditivity, absolute and metric pointwise moduli, attained exact unit-interval and totalized suprema, and the exact effective-injectivity threshold. Exact declaration names appear in section 12 .
FabiusInverseEffectiveContinuity.lean	Post-snapshot formalization of the recurrence-derived factorial endpoint lower bound, the exact numerical Δ_r endpoint inequality, strict and closed versions of the Delta and factorial dyadic inverse moduli, primitive-recursive denominator arithmetic, and effective uniform continuity with the simple $r = n$ factorial witness. It does not formalize the probabilistic event, logarithmic $r(n)$, exact ceiling d_* , or a sequential realizer. Every public declaration is crosswalked in section 12 .
Inverse_and_Sampling_Frontiers.tex and Inverse_Endpoint_All_Orders.tex	Exact finite-prefix quantiles, q -Richardson filters, Bell-polynomial inversion, and all-orders endpoint quantile asymptotics.
Small_Argument_Asymptotics.tex	Explicit lower-Lambert forward asymptotics and periodic Gamma-zeta coefficient calculus.

Corpus component	Role in the present report
q -calculus, sinc-product, Fourier-decay, Legendre/Lagrange, total-positivity, and frontier-compilation sources	Context for alternative certified oracles, shape strengthening, and research directions; not logical prerequisites of the computability proof.

The audit included generated `.tex` fragments and frozen variants, but the report cites parent documents rather than treating generated rows as independent mathematical sources. The repository was pinned because its `main` branch advanced during the audit; all path and count statements refer to commit `40fdea4cc0a728189f357389e3f114a2cb00e561`.

D Reproducing the exact-rational supplement

The archive accompanying this report contains

```
inverse_fabius_computability_experiments.py
```

and its captured output `numerical_output.txt`. The program uses only Python’s standard library. From the extracted package directory, run

```
python inverse_fabius_computability_experiments.py
```

The program performs all spline evaluations with `fractions.Fraction`. Its principal checks are:

1. the endpoint event lower bound and flatness upper bound in [table 1](#);
2. exact convergence of the centered splines at $1/2$, $1/4$, and $1/8$, using the exact values $1/2$, $5/72$, and $1/288$;
3. a finite-grid analogue of the least-interval-mass theorem;
4. a short certified bisection transcript illustrating all comparison margins.

These computations are regression tests and illustrations. No numerical observation is used as a premise of a theorem.

References

- [1] A. Grzegorzcyk, On the definitions of computable real continuous functions, *Fundamenta Mathematicae* **44** (1957), 61–71.
- [2] K. Weihrauch, *Computable Analysis: An Introduction*, Springer, Berlin, 2000.
- [3] M. B. Pour-El and J. I. Richards, *Computability in Analysis and Physics*, Springer, Berlin, 1989.
- [4] J. Fabius, A probabilistic example of a nowhere analytic C^∞ -function, *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete* **5** (1966), no. 2, 173–174, <https://doi.org/10.1007/BF00536652>.
- [5] B. Jessen and A. Wintner, Distribution functions and the Riemann zeta function, *Transactions of the American Mathematical Society* **38** (1935), no. 1, 48–88, <https://doi.org/10.1090/S0002-9947-1935-1501802-5>.

- [6] J. Arias de Reyna, An infinitely differentiable function with compact support: definition and properties, English translation of the 1982 paper, <https://arxiv.org/abs/1702.05442>.
- [7] J. Arias de Reyna, Arithmetic of the Fabius function, <https://arxiv.org/abs/1702.06487>.
- [8] V. Reshetnikov, ProveIt, Fabius-function development and documentation, repository snapshot 40fdea4cc0a728189f357389e3f114a2cb00e561, <https://github.com/VladimirReshetnikov/ProveIt>.
- [9] Wikipedia contributors, Computable real function, accessed 30 August 2026, https://en.wikipedia.org/wiki/Computable_real_function.