

Critical Transseries at a Moving Fold

A uniform two-sheet atlas,
large-sector crossover, and resonant action splitting

Research continuation for the ProveIt project

29 September 2026

Abstract

The exponential sectors in the inverse of a logarithmic–exponential map have poles of order $2n - 1$ at a critical core value. Their deepest-pole sum is a square root, but that formal sum does not locate the moving branch point and is not a uniform approximation at the transition. We resolve this explicitly posed project-local problem for

$$\delta + \log(1 + \delta/w) + wze^{-\delta} = 0.$$

Writing $s = w + 1$ and $z = s^2\zeta$, we construct an exact holomorphic two-sheet atlas on an explicit complex domain. Its discriminant gives $z_c(s) = s^2/2 + s^3/3 + 5s^4/8 + \dots$ and an all-order polynomial profile expansion with a geometric error bound valid at the branch point itself. A uniform coefficient theorem then proves the double-scaling law $a_n(\tau/n)/a_n(0) \rightarrow e^{-2\tau/3}$ and its first correction, where $a_n(s) = s^{2n-1}b_n(-1+s)$ normalizes the n th exponential sector. Finally, for general analytic folds driven by finitely many exponential actions, we prove an absolutely convergent, resonance-aware transseries representation and give a constructive finite positive generating set for its support. Explicit cancellations can change the first fractional action or remove the splitting altogether. The arguments use classical local analytic and coefficient-extraction methods; the contribution is the quantitative specialization, crossover calculation, and support construction, not a new general theory of analytic folds.

Scope and verification. This manuscript answers Research question 11.8 of the predecessor *Support-Controlled Reversion and the Exact One-Exponential Substitution Group* in its independent-parameter local setting. The new results are not Lean-formalized. Exact symbolic checks and 100-digit numerical diagnostics are supplied, but are not substitutes for the proofs or for independent review. Worldwide priority is not claimed.

Contents

1	Research target, contribution, and boundaries	2
1.1	The gap between a formal pole sum and a uniform inverse	2
1.2	What is established here	2
1.3	Classical input and the meaning of novelty	3
1.4	An essential parameter qualification	3
2	From the sector equation to a regular critical chart	3
2.1	The known sector functions	3
2.2	Blowing up the coalescing point	4
3	An explicit holomorphic two-sheet atlas	5
3.1	Quantitative perturbation estimates	5
3.2	The critical value and a useful enclosure	5
3.3	Exact preparation by contour moments	6
4	The moving discriminant and the nearby branch points	7
4.1	Critical-point expansions	7
4.2	Branch points when the perturbation parameter is fixed	8
5	Polynomial profiles with an error bound at the fold	8
5.1	The regular sheet coordinate	8
6	Failure of the unshifted model and fractional actions	10
6.1	A sharp diagnostic path	10
6.2	An explicit convergent critical transseries	10
6.3	How much discriminant accuracy is needed?	11
7	Uniform high-sector asymptotics and a crossover law	11
7.1	The singular amplitude	11
7.2	A uniform coefficient lemma	12
7.3	Application to the normalized sector functions	13
7.4	Why the exponential crossover is structurally stable	14
8	Analytic folds with finitely many exponential actions	14
8.1	Exact quadratic preparation without an unproved normal form	14
8.2	Grouping equal actions before taking a square root	15
8.3	A constructive finite positive support grid	16
8.4	A cancellation that changes the first fractional action	17
8.5	Relation to the broader transseries questions	17
9	Residual certificates that remain valid at the fold	17
10	Computation, reproducibility, and proof status	18
10.1	What the symbolic checks verify	18
10.2	Computing high sector indices without symbolic expression swell	19
10.3	Numerical checks at and close to the branch point	19
10.4	Numerical checks of the joint limit	20
10.5	Formalization boundaries and a feasible first tranche	20
11	Further research questions and concrete next steps	20
12	Conclusion	22
A	Dependency map and source audit	22
B	Reproduction instructions	23

1 Research target, contribution, and boundaries

1.1 The gap between a formal pole sum and a uniform inverse

The ProveIt transseries program separates several logically different levels: finite algebraic identities, formal support-controlled operations, analytic realization, and quantitative error bounds. This separation is particularly important near a zero of the derivative of the function being inverted. A coefficientwise expansion can be perfectly correct while its truncations fail precisely where infinitely many of its singular coefficients compete.

The inspected repository module `QuadraticCoreCatalan.lean` formalizes the finite Catalan algebra underlying the quadratic equation

$$A\varepsilon + c\varepsilon^2 + \sigma q = 0.$$

The package README and the module itself distinguish that algebra from the assembled power-series existence statement and from the identification of deepest poles in special-function inversions [1, 2]. We maintain that distinction here.

The predecessor manuscript [3] applies exact-core inversion to the logarithmic–exponential example. It proves an exact rational formula for the sector functions $b_n(w)$ and identifies their leading poles at $w = -1$. Its Research question 11.8 asks whether the resulting formal square-root expression can be promoted to a uniform complex transition approximation, with error bounds and the nearby branch points actually located. This is the central target of the present article.

The answer is affirmative, but the branch point must move. Keeping the quadratic model’s branch point fixed gives an approximation whose error is of order $s^{3/2}$ on a natural test path, rather than the s^2 order one might infer from a regular correction expansion. The corrected coordinate is

$$q^2 = 1 - \frac{z}{z_c(s)}, \quad s = w + 1, \quad (1)$$

not $1 - 2z/s^2$.

1.2 What is established here

The main results have three complementary forms.

First, an explicit two-sheet analytic representation is proved for $|s| \leq 10^{-2}$ and $|z/s^2| < 3/4$. It yields a convergent expansion

$$\delta = s \left(\sum_{j \geq 0} s^j V_j(q) - 1 \right), \quad V_j \in \mathbb{Q}[q], \quad \deg V_j \leq j + 1,$$

with a bound uniform in the truncation order and valid for $|q| \leq 1/2$, including $q = 0$. The first five profiles are given explicitly.

Second, the same atlas controls the limit of large sector index. If $a_n(s) = s^{2n-1}b_n(-1+s)$, then uniformly for bounded complex τ ,

$$\frac{a_n(\tau/n)}{a_n(0)} = e^{-2\tau/3} \left[1 + \frac{\tau/3 - 37\tau^2/36}{n} + \mathcal{O}(n^{-2}) \right]. \quad (2)$$

Thus the first drift of the branch point has an order-one effect on high sectors in the window $s \asymp n^{-1}$.

Third, an analytic fold perturbed by finitely many exponentials admits an absolutely convergent transseries whose leading splitting action is half the first *noncancelled* action in its discriminant.

Equal actions must be combined before selecting this first term. We give a finite, explicitly enumerable positive generating set for all subsequent action gaps. This answers a substantial analytic subclass of the predecessor's multi-action questions, not its unrestricted formal or resurgent versions.

1.3 Classical input and the meaning of novelty

Square-root branching at a simple critical point is classical. The Lambert W branch expansion is an especially relevant example [4]. Extraction of coefficient asymptotics from such a singularity is also classical [5]. General transseries background, including the need for appropriate support conditions, can be found in [6].

We do not claim to replace these theories. Instead, we prove the precise uniform analytic statement that the earlier formal calculation left open, compute its moving discriminant and polynomial profiles, and derive a specific joint critical/large-sector limit. The proofs below are self-contained at the level of elementary complex analysis, including Cauchy's formula, Rouché's theorem, the holomorphic implicit-function theorem, and the gamma-function coefficient formula. A constructive finite-grid refinement is then added for analytic finite-action folds.

1.4 An essential parameter qualification

The variable z is an *independent analytic perturbation parameter*. For the original fixed map $y + \log y + e^{-y}$, putting $x = w + \log w$ gives the specialization $z = e^{-x} = e^{-w}/w$. As $w \rightarrow -1$ on a local logarithm branch, that value tends to $-e$, not to zero. Therefore our small- (s, z) critical chart is not, by itself, an analysis of the fixed-coupling map at $w = -1$.

More generally the equation studied here is the local form of

$$y + \log y + \lambda e^{-y} = w + \log w, \quad \lambda = we^w z.$$

It is a small-coupling, independent-parameter analytic family. Its sector functions are exactly those of the predecessor, so its singularity analysis directly controls their critical behavior. Global continuation to the physical specialization $\lambda = 1$ is a different question and is not claimed in this article.

2 From the sector equation to a regular critical chart

2.1 The known sector functions

For $w \neq 0, -1$, let $\delta(w, z)$ denote the solution near $z = 0$ with $\delta(w, 0) = 0$ of

$$\delta + \log \left(1 + \frac{\delta}{w} \right) + wz e^{-\delta} = 0. \quad (3)$$

The logarithm in this equation is the local branch equal to zero when $\delta = 0$. Write

$$\delta(w, z) = \sum_{n \geq 1} b_n(w) z^n. \quad (4)$$

The predecessor gives

$$b_n(w) = \frac{(-1)^n}{n!} (T - n)^{n-1} \frac{w^{n+1}}{w+1}, \quad T = \frac{w}{w+1} \frac{d}{dw}. \quad (5)$$

For reference,

$$\begin{aligned} b_1(w) &= -\frac{w^2}{w+1}, \\ b_2(w) &= -\frac{w^3(2w^2+2w-1)}{2(w+1)^3}, \\ b_3(w) &= -\frac{w^4(9w^4+18w^3-11w+1)}{6(w+1)^5}. \end{aligned} \quad (6)$$

Its deepest-pole identity is

$$\lim_{s \rightarrow 0} s^{2n-1} b_n(-1+s) = -\frac{(2n-3)!!}{n!}, \quad (-1)!! = 1. \quad (7)$$

Consequently the most singular part of every sector has the formal sum

$$-\sum_{n \geq 1} \frac{(2n-3)!!}{n!} \frac{z^n}{s^{2n-1}} = s \left(\sqrt{1 - \frac{2z}{s^2}} - 1 \right). \quad (8)$$

Equations (5)–(8) are predecessor results, not new assertions of priority here.

2.2 Blowing up the coalescing point

Set

$$s = w + 1, \quad t = y + 1 = s + \delta, \quad h(t) = t + \log(1-t) = -\sum_{j \geq 2} \frac{t^j}{j}. \quad (9)$$

For $|s|, |t| < 1$, direct substitution in (3) gives

$$z = Z(s, t) := \frac{e^{t-s}}{1-s} (h(t) - h(s)). \quad (10)$$

Indeed $1 + \delta/w = (1-t)/(1-s)$, so the logarithmic difference in (3) is exactly the local difference used in (10).

The variables t and s vanish at the same scale. Put $t = sv$ and $z = s^2 \zeta$. Define

$$\Phi(s, v) = \frac{e^{s(v-1)}}{1-s} \frac{h(sv) - h(s)}{s^2}. \quad (11)$$

The apparent singularity at $s = 0$ is removable because

$$\frac{h(sv) - h(s)}{s^2} = -\sum_{j \geq 2} \frac{v^j - 1}{j} s^{j-2}. \quad (12)$$

Thus

$$\Phi(0, v) = P(v) := \frac{1-v^2}{2}, \quad \Phi(s, 1) = 0, \quad \Phi_v(s, 1) = -\frac{1}{(1-s)^2}. \quad (13)$$

There is a unique inverse germ $v(s, \zeta)$ near $\zeta = 0$ satisfying $v(s, 0) = 1$. Write

$$v(s, \zeta) = 1 + \sum_{n \geq 1} a_n(s) \zeta^n. \quad (14)$$

The analytic dependence on s removes the original coefficient poles:

$$b_n(-1+s) = s^{1-2n} a_n(s) \quad (s \neq 0), \quad a_n(0) = -\frac{(2n-3)!!}{n!}. \quad (15)$$

The latter identity follows from $v(0, \zeta) = \sqrt{1-2\zeta}$, with the square root equal to one at zero.

3 An explicit holomorphic two-sheet atlas

3.1 Quantitative perturbation estimates

Lemma 3.1. *For $|s| \leq r := 1/100$ and $|v| \leq 2$,*

$$|\Phi(s, v) - P(v)| < \frac{7}{50}, \quad |\Phi_v(s, v) + v| < \frac{4}{25}. \quad (16)$$

Proof. Let $H = (h(sv) - h(s))/s^2$ and $E = e^{s(v-1)}/(1-s)$, so $\Phi = EH$. The series (12) gives

$$|H - P| \leq R_h := \frac{r}{3} \left(\frac{8}{1-2r} + \frac{1}{1-r} \right).$$

Since $|s(v-1)| \leq 3r$, the elementary bound $e^u \leq (1-u)^{-1}$ for $0 \leq u < 1$ yields

$$|E| \leq E_0 := \frac{1}{(1-3r)(1-r)}, \quad |E - 1| \leq E_0 - 1.$$

Also $|P| \leq 5/2$, hence

$$|\Phi - P| \leq \frac{5}{2}(E_0 - 1) + E_0 R_h = \frac{37787705}{279504918} < \frac{7}{50}.$$

Differentiating the convergent series for H gives $|H_v + v| \leq 4r/(1-2r)$. Since $E_v = sE$,

$$\begin{aligned} |\Phi_v + v| &\leq 2(E_0 - 1) + E_0 \left[\frac{4r}{1-2r} + r \left(\frac{5}{2} + R_h \right) \right] \\ &= \frac{21177832}{139752459} < \frac{4}{25}. \end{aligned}$$

All estimates are absolute complex estimates; no real positivity of s is assumed. $\square$

Lemma 3.2. *For $|s| \leq 1/100$ and $|\zeta| < 3/4$, the equation $\Phi(s, v) = \zeta$ has exactly two roots in $|v| < 2$, counted with multiplicity. There is exactly one critical point $v_c(s)$ in $|v| < 2$, and it is simple as a zero of Φ_v . Its critical value is denoted by $\rho(s) = \Phi(s, v_c(s))$.*

Proof. On $|v| = 2$,

$$|P(v) - \zeta| = \frac{1}{2}|v^2 - (1 - 2\zeta)| \geq \frac{3}{4} \quad \text{when } |\zeta| \leq \frac{3}{4}.$$

The first estimate in Lemma 3.1 and Rouché's theorem compare $\Phi - \zeta$ with $P - \zeta$. Both roots of the latter lie strictly within $|v| = 2$. The second estimate compares Φ_v with $-v$ on the same circle, proving that the total number of critical points, with multiplicity, is one. The unique zero must consequently be simple. Holomorphic dependence of v_c follows from the implicit-function theorem. $\square$

3.2 The critical value and a useful enclosure

Differentiate (10). The critical-point equation is

$$k(t_c(s)) = h(s), \quad k(t) := h(t) + h'(t) = t + \log(1-t) - \frac{t}{1-t}. \quad (17)$$

Here $k(0) = 0$ and $k'(0) = -1$, so t_c is holomorphic near zero. Moreover,

$$v_c(s) = \frac{t_c(s)}{s}, \quad z_c(s) = s^2 \rho(s) = \frac{e^{t_c(s)-s}}{1-s} \frac{t_c(s)}{1-t_c(s)}. \quad (18)$$

The quotients have removable singularities at $s = 0$.

Lemma 3.3. *For $0 < |s| \leq 1/100$, the critical point satisfies $|t_c(s)| < |s|^2$. Its normalized critical value obeys*

$$\left| \rho(s) - \frac{1}{2} \right| < \frac{1}{250}. \quad (19)$$

In particular, ρ is nonzero throughout this disk and $|\rho| < 0.504$.

Proof. The exact series

$$k(t) = -t - \sum_{j \geq 2} \frac{j+1}{j} t^j$$

implies, on $|t| = |s|^2$,

$$|k(t) + t| \leq \frac{3|s|^4}{2(1 - |s|^2)}, \quad |h(s)| \leq \frac{|s|^2}{2(1 - |s|)}.$$

The sum of these bounds is strictly smaller than $|s|^2$ for $|s| \leq 1/100$. Rouché's theorem applied to $k(t) - h(s)$ and $-t$ therefore proves the asserted location. This solution is the critical point of Lemma 3.2 because $|t_c/s| < |s| < 2$.

Dividing (17) by s^2 and separating the first term of $h(s)$ also gives

$$\left| \frac{t_c(s)}{s^2} - \frac{1}{2} \right| \leq \frac{|s|}{3(1 - r)} + \frac{3|s|^2}{2(1 - r^2)}. \quad (20)$$

In the exponential factor of (18), write

$$\frac{e^{t_c - s}}{1 - s} = \exp(t_c - s - \log(1 - s)).$$

The exponent has absolute value at most $L = r^2(1 + 1/[2(1 - r)])$. Put $E_c = (1 - L)^{-1}$. The same exponential estimate gives a bound E_c for this factor and $E_c - 1$ for its difference from one. Substitution of (20) into (18) now bounds $|\rho - 1/2|$ by

$$\left[\frac{r}{3(1 - r)} + \frac{3r^2}{2(1 - r^2)} \right] \frac{E_c}{1 - r^2} + \frac{1}{2} \left(\frac{E_c}{1 - r^2} - 1 \right) = \frac{21851394853}{5997897210294} < \frac{1}{250}.$$

The estimate extends to $s = 0$ by continuity. □

3.3 Exact preparation by contour moments

Theorem 3.4 (Uniform two-sheet atlas). *There are holomorphic functions $A(s, \zeta)$ and $B(s, \zeta)$ on*

$$|s| < \frac{1}{100}, \quad |\zeta| < \frac{3}{4},$$

with B nowhere zero, such that the two roots in $|v| < 2$ are exactly

$$v_{\pm}(s, \zeta) = A(s, \zeta) \pm B(s, \zeta) \sqrt{1 - \frac{\zeta}{\rho(s)}}. \quad (21)$$

The normalization is $A(0, \zeta) = 0$, $B(0, \zeta) = 1$. The plus branch, with the square root equal to one at $\zeta = 0$, is the inverse germ $v(s, 0) = 1$. Its only branch value within this disk is $\rho(s)$; that branching is genuinely square-root. Its Taylor series about $\zeta = 0$ has radius of convergence exactly $|\rho(s)|$.

Proof. For $j = 1, 2$ define

$$S_j(s, \zeta) = \frac{1}{2\pi i} \int_{|v|=2} v^j \frac{\Phi_v(s, v)}{\Phi(s, v) - \zeta} dv. \quad (22)$$

The denominator never vanishes on the contour in the parameter domain, so these are holomorphic functions. By the argument principle they equal $v_1^j + v_2^j$, with multiplicities, even though the individual roots need not be single-valued. Set

$$A = \frac{S_1}{2}, \quad \Delta = \frac{S_2}{2} - A^2.$$

For distinct roots $\Delta = (v_1 - v_2)^2/4$, and this identity extends through a double root. A zero of Δ occurs precisely when both roots coalesce, hence precisely at $\zeta = \rho(s)$.

For completeness, simplicity of this discriminant zero follows directly from a local change of variable. Write $\Phi(s, v) - \rho(s) = (v - v_c(s))^2 G(s, v)$, where $G(s, v_c(s)) = \Phi_{vv}(s, v_c(s))/2 \neq 0$. The coordinate $\eta = (v - v_c)\sqrt{G(s, v)}$ has nonzero derivative at the critical point and therefore an analytic local inverse. The two roots are obtained by setting $\eta = \pm\sqrt{\zeta - \rho}$. Squaring their half-difference shows $\Delta = 2(\zeta - \rho)/\Phi_{vv}(s, v_c) + \mathcal{O}((\zeta - \rho)^2)$. Thus the zero is simple. It follows that

$$C(s, \zeta) := \frac{\Delta(s, \zeta)}{1 - \zeta/\rho(s)}$$

extends holomorphically and never vanishes. The extension is joint in (s, ζ) : a holomorphic function vanishing on $\zeta = \rho(s)$ is divisible by $\zeta - \rho(s)$, for example by integrating its ζ derivative along the segment from $\rho(s)$ to ζ in a local neighborhood. Nonvanishing at the zero follows from simplicity.

The parameter polydisk is simply connected, so the nonvanishing function C has a holomorphic square root B . At $s = 0$ the roots are $\pm\sqrt{1 - 2\zeta}$, hence $C(0, \zeta) = 1$; choose $B = 1$ there. This proves (21). At $\zeta = 0$ the plus branch agrees with $v = 1$ at $s = 0$, and remains that simple root by holomorphic continuation.

On $|\zeta| < |\rho(s)|$ the square root normalized at zero is single-valued and holomorphic, proving the lower bound on the Taylor radius. At $\zeta = \rho(s)$ its coefficient $B(s, \rho(s))$ is nonzero, so this inverse has a nonremovable square-root singularity there. The opposite inequality follows. The contour argument also excludes any additional local root collision or singularity of this two-root inverse inside $|\zeta| < 3/4$. $\square$

Corollary 3.5. *For $0 < |s| < 1/100$, the series (4) at $w = -1 + s$ has radius of convergence exactly $|z_c(s)|$. This is a statement about the independent variable z , with w fixed.*

4 The moving discriminant and the nearby branch points

4.1 Critical-point expansions

Equation (17) is both an exact definition and a recursive method. Because the linear coefficient of k is -1 , the coefficient of each successive power of s in t_c is determined by the already known ones. The result begins

$$t_c(s) = \frac{s^2}{2} + \frac{s^3}{3} - \frac{s^4}{8} - \frac{3s^5}{10} + \frac{s^6}{48} + \frac{323s^7}{840} + \frac{1123s^8}{5760} + \mathcal{O}(s^9), \quad (23)$$

$$\rho(s) = \frac{1}{2} + \frac{s}{3} + \frac{5s^2}{8} + \frac{7s^3}{10} + \frac{19s^4}{24} + \frac{361s^5}{420} + \mathcal{O}(s^6). \quad (24)$$

The second series follows by substituting the first in (18). In particular,

$$z_c(s) = \frac{s^2}{2} + \frac{s^3}{3} + \frac{5s^4}{8} + \frac{7s^5}{10} + \mathcal{O}(s^6). \quad (25)$$

The first omitted correction in the deepest-pole model is therefore already cubic in s . Although smaller than the quadratic term as a function value, it is not uniformly negligible after taking the square root near its zero.

The logarithm of the normalized discriminant is also useful:

$$\log(2\rho(s)) = \frac{2s}{3} + \frac{37s^2}{36} + \frac{539s^3}{810} + \mathcal{O}(s^4). \quad (26)$$

All displayed coefficients in (23)–(26) are checked independently by exact rational symbolic substitution in the accompanying script.

4.2 Branch points when the perturbation parameter is fixed

The equation for branch values in the (s, z) plane is $z = z_c(s) = s^2\rho(s)$. It can be inverted without an ambiguous formal choice of a square root. Define

$$U(s) = s\sqrt{2\rho(s)}, \quad U'(0) = 1,$$

using the holomorphic square root equal to one at $s = 0$. Let J be its local holomorphic inverse.

Proposition 4.1. *For every sufficiently small nonzero z , the two nearby branch points in the w plane of the independent-parameter equation are*

$$w_{\pm}(z) = -1 + J(\pm\sqrt{2z}), \quad (27)$$

where

$$J(u) = u - \frac{u^2}{3} - \frac{25u^3}{72} + \frac{137u^4}{540} + \frac{6529u^5}{17280} + \mathcal{O}(u^6). \quad (28)$$

They are simple square-root branch points and coalesce at $w = -1$ as $z \rightarrow 0$. There are exactly two such points in a sufficiently small fixed neighborhood of $w = -1$.

Proof. The equation $z = s^2\rho(s)$ is exactly $U(s)^2 = 2z$. Since U is biholomorphic near zero, it has precisely the two solutions in (27). Series substitution verifies (28). For nonzero small s , $z'_c(s) = s + \mathcal{O}(s^2) \neq 0$, so the discriminant zero is transverse to the fixed- z slice. The factor B in Theorem 3.4 is nonzero. Multiplication by s in $\delta = s(v - 1)$ does not remove the square-root singularity because $s \neq 0$ at these points. $\square$

This proposition locates actual local branch points of the two-variable family, not merely zeros of a truncated quadratic surrogate. The qualification in Section 1.4 remains in force: it is not a global classification for the map with coupling fixed at one.

5 Polynomial profiles with an error bound at the fold

5.1 The regular sheet coordinate

Substitute $\zeta = \rho(s)(1 - q^2)$ into the atlas and define

$$V(s, q) = A(s, \rho(s)(1 - q^2)) + qB(s, \rho(s)(1 - q^2)). \quad (29)$$

Changing q to $-q$ gives the other sheet. Since $|\rho(s)(1 - q^2)| < 0.504 \cdot 1.25 < 3/4$ for $|q| \leq 1/2$, this expression is holomorphic for $|s| < 1/100$, $|q| \leq 1/2$, and

$$|V(s, q)| < 2, \quad V(0, q) = q. \quad (30)$$

At $q = 0$ it equals the double root $t_c(s)/s$. The coalescence has become an ordinary point of the (s, q) parametrization.

Theorem 5.1 (Polynomial profiles and uniform remainder). *There are polynomials $V_j(q) \in \mathbb{Q}[q]$ such that*

$$V(s, q) = \sum_{j \geq 0} s^j V_j(q), \quad V_0(q) = q, \quad \deg V_j \leq j + 1. \quad (31)$$

For $j \geq 1$, $V_j(1) = 0$ as a polynomial identity. The first five are

$$\begin{aligned} V_0(q) &= q, \\ V_1(q) &= \frac{1}{2} + \frac{q}{3} - \frac{5q^2}{6}, \\ V_2(q) &= \frac{1}{3} - \frac{49q}{72} - \frac{5q^2}{9} + \frac{65q^3}{72}, \\ V_3(q) &= -\frac{1}{8} - \frac{799q}{1080} + \frac{9q^2}{8} + \frac{65q^3}{72} - \frac{157q^4}{135}, \\ V_4(q) &= -\frac{3}{10} + \frac{22739q}{51840} + \frac{31q^2}{18} - \frac{3373q^3}{1728} - \frac{628q^4}{405} + \frac{28369q^5}{17280}. \end{aligned} \quad (32)$$

For $N \geq 0$, put

$$\delta_N(s, q) = s \left(\sum_{j=0}^N s^j V_j(q) - 1 \right). \quad (33)$$

Then for $|s| < 1/100$ and $|q| \leq 1/2$,

$$|s(V(s, q) - 1) - \delta_N(s, q)| \leq \frac{2|s|(100|s|)^{N+1}}{1 - 100|s|}. \quad (34)$$

The estimate is valid at the branch point $q = 0$ and is independent of the sheet once its sign of q has been chosen.

Proof. Holomorphy gives a convergent s expansion with coefficients initially holomorphic in q . We prove their stronger polynomial property from

$$\Phi(s, V(s, q)) = \rho(s)(1 - q^2).$$

Write $\Phi(s, v) = \sum_{i \geq 0} s^i \Phi_i(v)$. Equation (12) and the exponential factor show that $\Phi_i \in \mathbb{Q}[v]$ and $\deg \Phi_i \leq i + 2$, with $\Phi_0(v) = (1 - v^2)/2$.

Suppose $V_0, \dots, V_{j-1}$ are known. At order s^j the only occurrence of V_j is $\Phi'_0(q)V_j = -qV_j$. Let $R_j(q)$ be the coefficient of s^j in

$$\Phi \left(s, \sum_{\ell=0}^{j-1} s^\ell V_\ell(q) \right) - \rho(s)(1 - q^2). \quad (35)$$

Thus $qV_j = R_j$. By induction R_j is a rational polynomial. For a term of degree $d \leq i + 2$ in Φ_i , the coefficient at total order j after substitution has degree at most $d + (j - i) \leq j + 2$. The subtraction on the right also obeys this bound. Hence $\deg R_j \leq j + 2$.

The analytically constructed coefficient V_j is regular at $q = 0$. Consequently $R_j(0) = 0$ and the polynomial R_j is divisible by q . It follows that $V_j \in \mathbb{Q}[q]$ and $\deg V_j \leq j + 1$. This argument is important: formal division by q alone would not justify regularity at the fold. The exact atlas supplies that missing fact.

Near $q = 1$ the same expression (29) is defined for small s , because its argument ζ is near zero. At $q = 1$ it equals the root $v = 1$, proving $V_j(1) = 0$. This is a statement about the coefficient polynomials; the quantitative error domain asserted here remains $|q| \leq 1/2$. Direct execution of the recurrence gives (32).

Finally, (30) and Cauchy's estimate on circles $|s| = r' < r$ give $|V_j(q)| \leq 2(r')^{-j}$ uniformly for $|q| \leq 1/2$. Let $r' \uparrow r$, sum the geometric tail, and multiply by $|s|$. This proves (34) for every N . $\square$

Remark 5.2 (What the error bound does and does not require). The estimate concerns an exact input value of q , equivalently an exact moving critical value $z_c(s)$. It does not silently include errors in computing z_c , nor does it include a separate asymptotic truncation of an exact logarithmic core at infinity. Errors in those quantities need additional propagation bounds, especially when q is small. A residual-based bound valid at a fold is given in Section 9.

6 Failure of the unshifted model and fractional actions

6.1 A sharp diagnostic path

Proposition 6.1. *Let $s \rightarrow 0^+$ and set $z = s^2/2$. On the sheet with positive q ,*

$$\delta = -s + \sqrt{\frac{2}{3}} s^{3/2} + \frac{s^2}{2} + \mathcal{O}(s^{5/2}). \quad (36)$$

The deepest-pole sum (8) equals $-s$ on this path. Its error is therefore asymptotic to $\sqrt{2/3} s^{3/2}$ and is not $\mathcal{O}(s^2)$.

Proof. Equation (24) gives

$$q^2 = 1 - \frac{1}{2\rho(s)} = \frac{2}{3}s + \frac{29}{36}s^2 + \mathcal{O}(s^3), \quad q = \sqrt{\frac{2}{3}} s^{1/2} (1 + \mathcal{O}(s)).$$

The polynomial profiles give $\delta = s(q - 1) + s^2(1/2 + q/3 - 5q^2/6) + \mathcal{O}(s^3)$ uniformly as $q \rightarrow 0$. Substitution proves (36). The leading nonzero error divided by s^2 diverges, proving the last assertion. $\square$

This does not contradict the predecessor's coefficientwise pole theorem. At each fixed n , all less singular Laurent terms disappear under the normalization s^{2n-1} . On a path where z/s^2 approaches $1/2$, however, infinitely many sectors are relevant and those discarded corrections move the singularity itself. The interchange of these two operations is the invalid step.

6.2 An explicit convergent critical transseries

Choose a real $a > 0$ and specialize

$$s = e^{-aX}, \quad z = \frac{1}{2} e^{-2aX}, \quad X \rightarrow +\infty.$$

The plus branch has the convergent exponential transseries

$$\delta(X) = -e^{-aX} + \sqrt{\frac{2}{3}} e^{-3aX/2} + \frac{1}{2} e^{-2aX} + \mathcal{O}(e^{-5aX/2}). \quad (37)$$

Convergence, not just formal asymptotic validity, follows because q is $s^{1/2}$ times a holomorphic unit and V is holomorphic in (s, q) . After setting $r = s^{1/2}$, the whole expression is an ordinary convergent power series in r near zero.

On the exact critical curve $z = z_c(s)$, by contrast, $q = 0$ and

$$\delta = t_c(s) - s = -s + \frac{1}{2}s^2 + \frac{1}{3}s^3 - \frac{1}{8}s^4 + \dots$$

There are only integer multiples of the action a on this curve. The fractional action in (37) is caused by detuning from the moving critical curve, not by a failure of the analytic parametrization.

6.3 How much discriminant accuracy is needed?

Let $z_c^{[M]}(s)$ be the Taylor polynomial of $z_c(s)$ through degree $M \geq 2$, and suppose

$$z_c(s) - z_c^{[M]}(s) = c_{M+1}s^{M+1} + \mathcal{O}(s^{M+2}), \quad c_{M+1} \neq 0.$$

On the path $z = z_c^{[M]}(s)$ one has

$$q^2 = 2c_{M+1}s^{M-1}(1 + \mathcal{O}(s)).$$

Since the half-difference between the two roots $\delta_{\pm}$ is $sqB(s, \zeta)$ and $B \rightarrow 1$, it follows that

$$\frac{\delta_+ - \delta_-}{2} \sim \sqrt{2c_{M+1}} s^{(M+1)/2}, \quad (38)$$

with the chosen square-root convention. This statement concerns splitting about the analytic center, so center shifts do not contaminate it.

For $M = 2$, the coefficient is $c_3 = 1/3$ and the exponent is $3/2$. For $M = 3$, $c_4 = 5/8$ and the splitting is asymptotic to $(\sqrt{5}/2)s^2$. This gives a simple accuracy rule: losing one order in the critical value can lose half that order in the inverse near the fold.

7 Uniform high-sector asymptotics and a crossover law

7.1 The singular amplitude

Let

$$\beta(s) = B(s, \rho(s)) = \partial_q V(s, 0), \quad \gamma(s) = -\rho(s)B_{\zeta}(s, \rho(s)) = [q^3]V(s, q). \quad (39)$$

The second identity for γ follows by substituting $\zeta = \rho(s)(1 - q^2)$ into (29); the contribution from A is even in q .

Lemma 7.1. *The amplitude β is holomorphic and nonzero for $|s| < 1/100$, with $\beta(0) = 1$ and*

$$\beta(s)^2 = \frac{2t_c(s)(1 - t_c(s))}{s^2(1 + t_c(s) - t_c(s)^2)}. \quad (40)$$

In particular,

$$\beta(s) = 1 + \frac{s}{3} - \frac{49s^2}{72} - \frac{799s^3}{1080} + \frac{22739s^4}{51840} + \mathcal{O}(s^5), \quad \gamma(s) = \mathcal{O}(s^2). \quad (41)$$

Proof. Expand Φ in v at the critical point. If $v = v_c + \beta q + \mathcal{O}(q^2)$ and $\zeta = \rho(1 - q^2)$, comparison of quadratic terms gives

$$\beta^2 = -\frac{2\rho}{\Phi_{vv}(s, v_c)}.$$

Writing $E = e^{t_c - s}/(1 - s)$, differentiation of (11) and use of (17) yield

$$\Phi_{vv}(s, v_c) = Ek'(t_c), \quad k'(t) = -\frac{1 + t - t^2}{(1 - t)^2}.$$

Combine this with (18) to obtain (40). The analytic choice of its square root is fixed by $\beta(0) = 1$. Equation (41) also follows by taking the coefficients of q and q^3 in (32). $\square$

7.2 A uniform coefficient lemma

We record the elementary transfer argument needed below. It is a specialized version of the coefficient-transfer principle in [5], with the uniformity made explicit.

Lemma 7.2. *Suppose $\mathcal{A}_s(Z)$ and $\mathcal{B}_s(Z)$ are holomorphic in $|Z| < R$ for some $R > 1$, uniformly bounded on smaller closed disks, with s in a compact parameter set. Suppose*

$$f_s(Z) = \mathcal{A}_s(Z) + \mathcal{B}_s(Z)(1 - Z)^{1/2}$$

as germs at zero. Set $\beta_s = \mathcal{B}_s(1)$ and $\gamma_s = -\mathcal{B}'_s(1)$. Then, uniformly in s ,

$$[Z^n]f_s(Z) = -\frac{\beta_s}{2\sqrt{\pi}}n^{-3/2} + \left(-\frac{3\beta_s}{16\sqrt{\pi}} + \frac{3\gamma_s}{4\sqrt{\pi}}\right)n^{-5/2} + \mathcal{O}(n^{-7/2}). \quad (42)$$

An expansion to every algebraic order in $1/n$ follows from higher Taylor coefficients of $\mathcal{B}_s$ at one.

Proof. The binomial coefficient identity, for the half-integers used here, is

$$[Z^n](1 - Z)^\alpha = \frac{\Gamma(n - \alpha)}{\Gamma(-\alpha)\Gamma(n + 1)}. \quad (43)$$

It follows either by multiplying the factors of the generalized binomial coefficient or by induction in n . The gamma-ratio expansion gives

$$\begin{aligned} [Z^n](1 - Z)^{1/2} &= -\frac{1}{2\sqrt{\pi}}n^{-3/2} \left(1 + \frac{3}{8n} + \mathcal{O}(n^{-2})\right), \\ [Z^n](1 - Z)^{3/2} &= \frac{3}{4\sqrt{\pi}}n^{-5/2}(1 + \mathcal{O}(n^{-1})). \end{aligned}$$

The needed constants are also obtained directly from the first two terms of Stirling's formula; no parameter occurs in these scalar estimates.

Taylor division at $Z = 1$ gives

$$\mathcal{B}_s(Z) = \beta_s + \gamma_s(1 - Z) + (1 - Z)^2\mathcal{C}_s(Z),$$

where $\mathcal{C}_s$ is holomorphic on the same disk and uniformly bounded on every smaller disk. To justify the coefficient remainder, fix $1 < R_1 < R$. Its coefficients satisfy $|[Z^j]\mathcal{C}_s| \leq MR_1^{-j}$ uniformly in s . The coefficients d_k of $(1 - Z)^{5/2}$ satisfy $|d_k| \leq M'(k + 1)^{-7/2}$, after enlarging M' for finitely many small indices. The convolution is bounded by

$$MM' \sum_{j=0}^n R_1^{-j} (n - j + 1)^{-7/2}.$$

For $j \leq n/2$ this is $\mathcal{O}(n^{-7/2})$ by a geometric sum. For $j > n/2$ it is exponentially small, hence has the same bound. The analytic term $\mathcal{A}_s$ likewise has exponentially small coefficients. This proves (42) with a uniform remainder. Repeating Taylor division to any fixed order gives the final assertion by exactly the same convolution estimate. $\square$

7.3 Application to the normalized sector functions

Theorem 7.3 (Uniform large-sector expansion). *Uniformly for $|s| \leq 1/200$, as $n \rightarrow \infty$,*

$$a_n(s) = -\frac{\beta(s)}{2\sqrt{\pi}}\rho(s)^{-n}n^{-3/2} \left[1 + \frac{1}{n} \left(\frac{3}{8} - \frac{3\gamma(s)}{2\beta(s)} \right) + \mathcal{O}(n^{-2}) \right]. \quad (44)$$

The expansion extends to every fixed algebraic order, with coefficients holomorphic in s and a remainder uniform on this closed disk.

Proof. Put $Z = \zeta/\rho(s)$. Theorem 3.4 gives

$$v(s, \rho(s)Z) = \mathcal{A}_s(Z) + \mathcal{B}_s(Z)(1 - Z)^{1/2},$$

where $\mathcal{A}_s(Z) = A(s, \rho(s)Z)$ and $\mathcal{B}_s(Z) = B(s, \rho(s)Z)$. By Lemma 3.3, these functions are holomorphic in $|Z| < 7/5$ since $0.504 \cdot 7/5 < 3/4$. Restricting $|s| \leq 1/200$ leaves a compact subset of the s domain, so the uniform boundedness hypotheses of Lemma 7.2 hold on smaller Z disks.

The coefficient of Z^n is $\rho(s)^n a_n(s)$ for $n \geq 1$. Apply the lemma and factor out the leading term. The division is uniform because β is nonzero and therefore bounded away from zero on the compact s disk. The all-order statement follows from the last assertion of the lemma and holomorphic dependence of every local Taylor coefficient. $\square$

The theorem includes $s = 0$, where the coefficients are exactly

$$a_n(0) = -\frac{2^{n-1}\Gamma(n-1/2)}{\sqrt{\pi}\Gamma(n+1)}. \quad (45)$$

For nonzero s , multiplication by s^{1-2n} gives the corresponding uniformly normalized asymptotic statement for the original rational functions $b_n(-1+s)$.

Theorem 7.4 (Critical/large-sector crossover). *For every $T > 0$, uniformly for complex $|\tau| \leq T$, one has*

$$\frac{a_n(\tau/n)}{a_n(0)} = e^{-2\tau/3} \left[1 + \frac{\tau/3 - 37\tau^2/36}{n} + \mathcal{O}_T(n^{-2}) \right] \quad (n \rightarrow \infty). \quad (46)$$

In particular, the limit is $e^{-2\tau/3}$, not one unless this exponential happens to equal one.

Proof. For n sufficiently large in terms of T , $s = \tau/n$ lies in the disk of Theorem 7.3. Dividing that theorem by its value at $s = 0$ and using (41) gives

$$\frac{a_n(s)}{a_n(0)} = \beta(s)(2\rho(s))^{-n}(1 + \mathcal{O}_T(n^{-2})).$$

Indeed the coefficient of $1/n$ in (44) is $3/8 + \mathcal{O}(s^2)$, so its change contributes no larger error; the uniform remainders suffice. The logarithm in (26) is the analytic one vanishing at zero. Hence

$$\begin{aligned} \beta(\tau/n) &= 1 + \frac{\tau}{3n} + \mathcal{O}_T(n^{-2}), \\ (2\rho(\tau/n))^{-n} &= \exp \left(-\frac{2\tau}{3} - \frac{37\tau^2}{36n} + \mathcal{O}_T(n^{-2}) \right). \end{aligned}$$

Multiplication proves (46). $\square$

7.4 Why the exponential crossover is structurally stable

The phenomenon is not tied to the particular coefficients above. Suppose an analytic family has an atlas of the same form, with a unique dominant simple square-root singularity at $\rho(s)$, nonzero amplitude $\beta(s)$, and a common continuation disk after normalizing the singularity to one. The proof of Theorem 7.3 then gives

$$\frac{a_n(s)}{a_n(0)} \sim \frac{\beta(s)}{\beta(0)} \left(\frac{\rho(s)}{\rho(0)} \right)^{-n}$$

uniformly for small s as $n \rightarrow \infty$. If $\rho(s)/\rho(0) = 1 + \kappa s^k + \mathcal{O}(s^{k+1})$, with $\kappa \neq 0$ and $k \geq 1$, then any regime with $s \rightarrow 0$ and $ns^k \rightarrow \tau$ yields the limit $e^{-\kappa\tau}$. This follows by taking an analytic logarithm; $ns^{k+1} \rightarrow 0$ in such a regime. It is a conditional universality statement with the analytic continuation and nonvanishing hypotheses stated explicitly, not a claim about arbitrary formal transseries.

8 Analytic folds with finitely many exponential actions

The previous example produces fractional exponential actions along a critical path. We now isolate the general analytic mechanism and address resonances without pretending that independent formal multi-indices remain distinct after exponential specialization.

8.1 Exact quadratic preparation without an unproved normal form

Lemma 8.1 (Local analytic fold preparation). *Let $F(u, \mathbf{q})$ be holomorphic near $(0, \mathbf{0}) \in \mathbb{C} \times \mathbb{C}^d$ and suppose*

$$F(0, \mathbf{0}) = F_u(0, \mathbf{0}) = 0, \quad F_{uu}(0, \mathbf{0}) \neq 0. \quad (47)$$

After shrinking the neighborhood, there are holomorphic functions $A(\mathbf{q})$, $D(\mathbf{q})$ and a nonvanishing holomorphic function $\mathcal{U}(u, \mathbf{q})$ such that

$$F(u, \mathbf{q}) = \mathcal{U}(u, \mathbf{q})((u - A(\mathbf{q}))^2 - D(\mathbf{q})), \quad A(\mathbf{0}) = D(\mathbf{0}) = 0. \quad (48)$$

The two nearby roots, with multiplicity, are $u_{\pm} = A(\mathbf{q}) \pm \sqrt{D(\mathbf{q})}$.

Proof. Choose a sufficiently small circle $|u| = r$ on which $F(u, \mathbf{0})$ is nonzero and encloses no zero other than its double zero at zero. For sufficiently small $\mathbf{q}$, Rouché's theorem preserves the count of two zeros. Define the first two root moments by the integrals of $u^j F_u/F$ over this circle, exactly as in (22). Their symmetric combinations define A and D , and the monic quadratic $P(u, \mathbf{q}) = (u - A)^2 - D$ has exactly these two zeros with their multiplicities.

For each fixed $\mathbf{q}$, F/P is holomorphic within the circle after removing its apparent zero-over-zero singularities. Joint holomorphy can be verified without choosing the individual roots: for $|u| < r$ set

$$\mathcal{U}(u, \mathbf{q}) = \frac{1}{2\pi i} \int_{|\xi|=r} \frac{F(\xi, \mathbf{q})}{P(\xi, \mathbf{q})(\xi - u)} d\xi.$$

The denominator on the contour is nonzero, so this is jointly holomorphic. For fixed $\mathbf{q}$, Cauchy's formula identifies it with F/P where the quotient is already defined, and the equality extends everywhere inside. At the origin its value is $F_{uu}(0, \mathbf{0})/2 \neq 0$; shrink the neighborhood once more to make it nonvanishing. This proves the factorization and root formula. $\square$

The lemma is the classical quadratic preparation mechanism, included with a proof to make the later support construction independent of any unstated normal-form theorem. No novelty is claimed for analytic quadratic preparation itself.

8.2 Grouping equal actions before taking a square root

Fix real $\mu_1, \dots, \mu_d > 0$ and specialize

$$q_i(X) = e^{-\mu_i X}, \quad X \rightarrow +\infty. \quad (49)$$

Let

$$S = \{\mathbf{n} \cdot \boldsymbol{\mu} : \mathbf{n} \in \mathbb{N}^d\}. \quad (50)$$

This monoid is locally finite: for $\mathbf{n} \cdot \boldsymbol{\mu} \leq M$, each coordinate satisfies $n_i \leq M/\mu_i$, so there are only finitely many possible multi-indices. In particular, each action has only finitely many representations by such a multi-index.

Expand the discriminant of Lemma 8.1 as

$$D(\mathbf{q}) = \sum_{\mathbf{n} \in \mathbb{N}^d} d_{\mathbf{n}} \mathbf{q}^{\mathbf{n}}.$$

The expansion is absolutely convergent in a small polydisk. Under (49), combine terms with the same action:

$$D(X) = \sum_{\nu \in S} d_{\nu} e^{-\nu X}, \quad d_{\nu} = \sum_{\mathbf{n} \cdot \boldsymbol{\mu} = \nu} d_{\mathbf{n}}. \quad (51)$$

These finite sums are the relevant coefficients. They are independent of how the original monomials representing a given action are listed. Because $D(\mathbf{0}) = 0$, the coefficient d_0 is zero.

Theorem 8.2 (Resonance-aware action splitting). *Under (47) and (49), exactly one of the following alternatives holds.*

If all grouped coefficients d_{ν} vanish, then $D(X) = 0$ for all sufficiently large X , and the two roots coincide there.

Otherwise there is a least $\lambda > 0$ with $d_{\lambda} \neq 0$. Then the roots have absolutely convergent exponential transseries

$$u_{\pm}(X) = A(X) \pm \sqrt{d_{\lambda}} e^{-\lambda X/2} \sum_{k \geq 0} \binom{1/2}{k} R(X)^k, \quad R(X) = \sum_{\nu > \lambda} \frac{d_{\nu}}{d_{\lambda}} e^{-(\nu-\lambda)X}. \quad (52)$$

For either consistent choice of $\sqrt{d_{\lambda}}$,

$$u_+(X) - u_-(X) \sim 2\sqrt{d_{\lambda}} e^{-\lambda X/2}. \quad (53)$$

The expansions in (52), including expansion of each power R^k and grouping equal output actions, are absolutely convergent for all sufficiently large X .

Proof. Absolute convergence of the original multivariable Taylor expansion for large X justifies grouping by action. Local finiteness of S implies that every nonempty subset of S has a least member: choose one member, then minimize over the finite set of smaller members. Thus the second alternative has a well-defined first action λ .

Factor $d_{\lambda} e^{-\lambda X}$ from (51). All exponents remaining in R are positive. More is needed than pointwise smallness: we establish smallness of the absolute coefficient sum. Choose X_0 large enough that the original expansion converges absolutely. If the tail is nonempty, it has a least positive gap $\eta = \min\{\nu - \lambda : \nu > \lambda, d_{\nu} \neq 0\}$. Then

$$\begin{aligned} \sum_{\nu > \lambda} \left| \frac{d_{\nu}}{d_{\lambda}} \right| e^{-(\nu-\lambda)X} &\leq e^{-\eta(X-X_0)} \sum_{\nu > \lambda} \left| \frac{d_{\nu}}{d_{\lambda}} \right| e^{-(\nu-\lambda)X_0} \\ &\longrightarrow 0. \end{aligned}$$

When the tail is empty the conclusion is immediate. For large X this absolute sum is less than one. Therefore the binomial series in (52) converges absolutely even after all its products are expanded: it is majorized by $\sum_k |(\frac{1}{2})^k| \|R\|_X^k$, with $\|R\|_X < 1$. It is exactly a square root of $1 + R(X)$ tending to one. Substitution in Lemma 8.1 proves the root formula and leading splitting. If every grouped coefficient vanishes, the absolutely convergent sum for $D(X)$ is identically zero, proving the first alternative. $\square$

Editorial note (ProveIt, 2026-09-29). The later package *Beyond Finite-Action Folds*, filed in batch 47 as `Critical_Hahn_Transseries_Beyond_Finite_Action_Folds/` in the series directory `Analysis/Transseries/docs/series-and-transseries/`, continues Theorem 8.2 from finitely many to countably many actions, for the feedback kernel $U = \beta \sum_j a_j q^j e^{jU}$ with linear slopes and power-law amplitudes $a_j = a j^{-1-\alpha}$, $1 < \alpha < 2$. At the critical coupling the square-root fold becomes a nonanalytic boundary with exponent $1/\alpha$ and a convergent Hahn chart (its `thm:hahn`); its truncations to finitely many actions are ordinary folds of the kind treated here, whose critical-value and curvature drift it computes (its `thm:fold-drift`, `thm:hard`). It is a model-specific extension, not a generalization of Theorem 8.2, and it does not settle the higher-multiplicity question of Section 11.

8.3 A constructive finite positive support grid

There is a useful stronger conclusion than mere local finiteness of the resulting support. Although the differences $\nu - \lambda$ need not belong to the original positive monoid S , they lie in a finitely generated positive extension that can be constructed explicitly.

Theorem 8.3 (Finite-grid support after a fold). *In the nonzero-discriminant alternative of Theorem 8.2, put $\mu_{\max} = \max_i \mu_i$ and define*

$$E_\lambda = \{\mathbf{n} \in \mathbb{N}^d : \mathbf{n} \cdot \boldsymbol{\mu} > \lambda\}.$$

Let M_λ be its componentwise minimal elements. Then M_λ is finite and is contained in the explicit box

$$0 \leq m_i \leq \left\lfloor \frac{\lambda + \mu_{\max}}{\mu_i} \right\rfloor \quad (1 \leq i \leq d). \quad (54)$$

Define the positive numbers $\beta_{\mathbf{m}} = \mathbf{m} \cdot \boldsymbol{\mu} - \lambda$ for $\mathbf{m} \in M_\lambda$, and let H be the additive monoid generated by

$$\mu_1, \dots, \mu_d, \quad \{\beta_{\mathbf{m}} : \mathbf{m} \in M_\lambda\}. \quad (55)$$

Then

$$\text{supp } A(X) \subseteq H, \quad \text{supp } \sqrt{D(X)} \subseteq \lambda/2 + H, \quad \text{supp } u_\pm \subseteq H \cup (\lambda/2 + H). \quad (56)$$

In particular, adjoining $\lambda/2$ to the finite list (55) gives a single finite positive generating set containing both root supports. If all μ_i lie in $(a/N)\mathbb{N}$ for some $a > 0$ and positive integer N , both supports lie in $(a/(2N))\mathbb{N}$.

Proof. The set E_λ is upward closed under the componentwise order. If $\mathbf{m}$ is minimal and $m_i > 0$, then $(\mathbf{m} - \mathbf{e}_i) \cdot \boldsymbol{\mu} \leq \lambda$, for otherwise minimality would fail. Hence $\mathbf{m} \cdot \boldsymbol{\mu} \leq \lambda + \mu_i \leq \lambda + \mu_{\max}$. Every coordinate obeys (54), so the set of minimal indices is finite. This also gives a direct finite search for it, once λ and exact comparisons among the actions are available.

For each $\mathbf{n} \in E_\lambda$, decrease coordinates while remaining in E_λ until no further decrease is possible. The process terminates because the sum of coordinates decreases. It produces $\mathbf{m} \in M_\lambda$ with $\mathbf{m} \leq \mathbf{n}$. Therefore

$$\mathbf{n} \cdot \boldsymbol{\mu} - \lambda = \beta_{\mathbf{m}} + (\mathbf{n} - \mathbf{m}) \cdot \boldsymbol{\mu} \in H.$$

Every positive gap appearing in R has this form. Thus $\text{supp } R \subseteq H$, and every expanded power R^k has support in H . The absolute convergence proved in Theorem 8.2 justifies collecting these powers; local finiteness also follows because H has finitely many strictly positive generators. Multiplication by $e^{-\lambda X/2}$ gives the second inclusion in (56). The Taylor expansion of A has support in $S \subseteq H$, proving the other inclusions. If the original actions lie in $(a/N)\mathbb{N}$, then so do λ and all the positive differences β_m ; halving λ only requires the stated doubled denominator. $\square$

The finite box is a sufficient construction, not an assertion that its resulting generating list is minimal. A smaller list may be obtained from the actual support of the discriminant. Also, identifying the first nonzero grouped coefficient is not claimed to be an automatically decidable operation for an arbitrary analytic input. When the nonzero case is known and exact coefficient and action comparisons are available, a search in increasing action terminates; the identically zero case may require a separate identity proof.

8.4 A cancellation that changes the first fractional action

Example 8.4. Consider

$$F(u, q_1, q_2) = u^2 - (q_1^2 - q_2 + q_1^3), \quad (\mu_1, \mu_2) = (1, 2).$$

Before specialization, the terms q_1^2 and $-q_2$ are distinct monomials. After specialization they have the same action and cancel exactly:

$$D(X) = e^{-2X} - e^{-2X} + e^{-3X} = e^{-3X}.$$

Consequently $u_{\pm} = \pm e^{-3X/2}$. A rule that selected one monomial of smallest action before combining resonances would incorrectly predict splitting at action 1 instead of $3/2$. If the term q_1^3 is removed, the discriminant vanishes identically on the specialized curve and the two roots coincide there, even though the multivariable discriminant $q_1^2 - q_2$ is not the zero function.

This example distinguishes an analytic identity in all perturbation variables from an identity after restriction to an exponential curve. The action theorem needs only the latter, and its grouping prescription makes that distinction explicit.

8.5 Relation to the broader transseries questions

Theorems 8.2 and 8.3 concern analytic germs in a *finite* collection of exponentially small parameters. Their convergence follows from ordinary Taylor convergence and a binomial majorant. They do not assert convergence for arbitrary Hahn series, realization of unrestricted formal supports, Borel summability of Gevrey series, or preservation of resurgent Stokes data. Those are separate problems. The result is nevertheless more than a formal square-root manipulation: it proves representation-independent handling of resonances, absolute summability after all regroupings, and a finite positive support grid with a concrete search bound.

9 Residual certificates that remain valid at the fold

A derivative-based inverse error bound usually divides by a quantity that tends to zero at a critical point. Quadratic preparation gives a replacement that is uniform across the collision.

Proposition 9.1 (Root-set residual bound). *Assume a factorization (48) is known on a domain containing a candidate $\tilde{u}$ and both local roots. Suppose $|\mathcal{U}(\tilde{u}, \mathbf{q})| \geq m > 0$, and put*

$$r = F(\tilde{u}, \mathbf{q}), \quad R = |r|/m, \quad d = \min\{|\tilde{u} - u_+|, |\tilde{u} - u_-|\}.$$

Then

$$d \leq \sqrt{R}. \quad (57)$$

If $D(\mathbf{q}) \neq 0$, the stronger combined estimate is

$$d \leq \min \left\{ \sqrt{R}, \frac{R}{|\sqrt{D(\mathbf{q})}|} \right\}. \quad (58)$$

The square-root exponent in (57) is optimal at a double root in general.

Proof. The preparation gives

$$|(\tilde{u} - u_+)(\tilde{u} - u_-)| \leq R.$$

The product is at least d^2 , proving (57). If the roots are distinct, their separation is $2|\sqrt{D}|$. By the triangle inequality at least one of the two candidate-to-root distances is at least $|\sqrt{D}|$. Multiplying the smaller distance by this lower bound proves $d \leq R/|\sqrt{D}|$. For $F(u) = u^2$, the exact distance to the root is $\sqrt{|F(u)|}$, so no uniform replacement by a higher power of the residual is possible. $\square$

The estimate certifies proximity to the *root set*, not a preferred sheet. A sheet-specific certificate requires continuation from a known base point, an isolating region, or other branch information. This qualification is necessary near a collision: the residual alone cannot distinguish two very close roots.

A practical verification pipeline can therefore combine a polynomial profile to obtain a candidate with a rigorous enclosure of the residual and the unit factor. Away from the fold, the second term in (58) recovers linear error transport. At the fold, the first term remains finite. The accompanying script does not implement interval enclosures of the unit factor; the proposition gives the mathematical certificate that such an implementation should check.

Editorial note (ProveIt, 2026-09-29). In the real, strictly monotone case the linear error transport recovered by the second term of (58) is the residual bracket machine-checked in the repository as the Lean theorem `Fabius.exists_eq_in_residual_interval` (`Analysis/FabiusFunction/Lean/FabiusFunction/MeanValueBracket.lean`): if $f' \geq m > 0$ on a convex set containing $[x - |f(x) - z|/m, x + |f(x) - z|/m]$, then f takes the value z in that interval. Proposition 9.1 is the complex fold counterpart and is not covered by that lemma. This article uses convergent profiles and contains no least-term truncation.

10 Computation, reproducibility, and proof status

10.1 What the symbolic checks verify

The package contains `verify.py`, using SymPy and mpmath. The exact part checks the critical-point equation through degree eight, ρ through degree five, the first five profile polynomials, the amplitude formula through degree four, the logarithm (26), and the branch-point reversion (28). The rational inequalities used for the explicit domain are checked as exact rational comparisons, not by a floating-point threshold test.

These tests can detect incorrect coefficients or arithmetic. They do not prove the existence of the holomorphic atlas, all-order polynomiality, the uniform coefficient theorem, or the finite-grid result. Those claims are established by the arguments in the preceding sections. The script also independently evaluates the first three known sector formulas (6) as a consistency check on the normalization.

10.2 Computing high sector indices without symbolic expression swell

A scalar Lagrange formula provides an $\mathcal{O}(n^2)$ arithmetic algorithm for $a_n(s)$ at a fixed numerical value of s . Put $u = v - 1$. Direct expansion of (11) gives

$$\Phi(s, 1+u) = -\frac{e^{su}u}{(1-s)^2}J_s(u), \quad J_s(u) = 1 + \sum_{j \geq 1} \frac{(s/(1-s))^{j-1}}{(1-s)(j+1)}u^j. \quad (59)$$

For $s = 0$, the removable interpretation is $J_0(u) = 1 + u/2$. To verify the formula directly, use

$$h(s(1+u)) - h(s) = su + \log \left(1 - \frac{s}{1-s}u \right)$$

and separate its linear term in u .

Since $\Phi_v(s, 1) \neq 0$, ordinary scalar Lagrange inversion gives

$$a_n(s) = \frac{(-1)^n(1-s)^{2n}}{n} [u^{n-1}] (e^{-nsu} J_s(u)^{-n}). \quad (60)$$

This formula is also a purely formal identity in u over the rational function field in s ; it does not require a convergence assumption to extract a fixed coefficient.

Write $J_s(u) = \sum_{j \geq 0} J_j u^j$, $J_0 = 1$, and $J_s(u)^{-n} = \sum_{k \geq 0} C_k u^k$. Differentiating $C = J_s^{-n}$ gives $JC' = -nJ'C$. Comparing coefficients yields

$$C_0 = 1, \quad C_k = \frac{1}{k} \sum_{j=1}^k ((1-n)j - k) J_j C_{k-j}. \quad (61)$$

Computing $C_0, \dots, C_{n-1}$ takes $\mathcal{O}(n^2)$ arithmetic operations; a final convolution with e^{-nsu} evaluates (60). This is an operation count, not a claim of quadratic bit complexity or numerical stability at arbitrary precision. The supplied numerical calculations use 100 decimal digits.

10.3 Numerical checks at and close to the branch point

Table 1 compares the order- $N = 4$ profile truncation with a high-precision root. At $q = 0$ the reference root is the independently computed $t_c(s) - s$. For $q \neq 0$, the local root is computed directly from (10), initialized on the sheet selected by the profile.

s	q	Observed absolute error	Bound (34)
0.001	0	2.122×10^{-20}	2.223×10^{-8}
0.001	0.25	2.233×10^{-19}	2.223×10^{-8}
0.004	0.1	5.465×10^{-16}	1.366×10^{-4}
-0.004	0.25	9.103×10^{-16}	1.366×10^{-4}
0.001	0.000001	2.122×10^{-20}	2.223×10^{-8}

Table 1: Order-four profile checks, including the exact branch point and an extremely close point. The bound column is rounded upward at the shown precision. The observations are floating-point diagnostics, not interval-arithmetic certificates.

The proved bound is intentionally conservative: its coefficient majorant comes from a common complex domain and the simple root bound $|v| < 2$, not from optimized estimates for these particular real inputs. Its purpose is to prove uniform validity through coalescence, rather than to predict the actual error sharply.

10.4 Numerical checks of the joint limit

Table 2 evaluates $a_n(\tau/n)/a_n(0)$ by (60) and compares it with both the limiting exponential and the first corrected expression in Theorem 7.4. The decreasing corrected errors are consistent with the proved $\mathcal{O}_T(n^{-2})$ remainder.

τ	n	$a_n(\tau/n)/a_n(0)$	$e^{-2\tau/3}$	Corrected error
-0.5	100	1.389698524	1.395612425	1.93×10^{-6}
-0.5	200	1.392656005	1.395612425	4.35×10^{-7}
-0.5	400	1.394134330	1.395612425	1.03×10^{-7}
0.5	100	0.715865297	0.716531311	1.91×10^{-5}
0.5	200	0.716203127	0.716531311	4.75×10^{-6}
0.5	400	0.716368411	0.716531311	1.18×10^{-6}
1.0	200	0.511619442	0.513417119	1.50×10^{-5}
1.0	400	0.512522036	0.513417119	3.73×10^{-6}

Table 2: Large-sector crossover diagnostics. “Corrected error” is the absolute difference from the full displayed right-hand side of (46) with its $\mathcal{O}_T(n^{-2})$ term omitted.

The JSON output records the decimal precision, package versions, exact expressions, and all test rows. The code writes its output relative to its own directory, so running it from another working directory does not silently overwrite unrelated data files.

10.5 Formalization boundaries and a feasible first tranche

No new Lean file is included or claimed verified. The repository’s existing Catalan module supplies useful finite algebra, but it does not prove the complex analytic statements of this manuscript. The following separation would keep a future formalization honest and manageable.

Layer	Mathematical content	Present status
Finite algebra	Displayed rational coefficients, polynomial recurrences, Catalan normalization	Exact symbolic checks; related Catalan core already has a repository Lean module.
Local analysis	Root count, holomorphic moments, moving discriminant, uniform profile remainder	Proofs here; no new machine-checked proof.
Coefficient asymptotics	Common continuation disk, half-integer transfer, joint scaling limit	Proofs here and numerical diagnostics; not formalized.
Action support	Resonance grouping, finite positive grid, absolute summability	Proofs here; only the toy resonance is numerically checked.

A useful first formalization target is the finite polynomial recurrence, together with a theorem that an independently supplied analytic solution forces the divisibility by q . A second target is the purely ordered finite-box argument in Theorem 8.3. These avoid claiming that formal coefficient identities already imply analytic convergence. The contour atlas and uniform transfer theorem are substantial later targets.

11 Further research questions and concrete next steps

The following questions distinguish genuine continuations from results already proved above. They range from implementable certification tasks to broader analytic and transseries problems.

Research question 11.1 (Sharp domains and practical certified evaluation). How far can the explicit atlas domain be enlarged while retaining a single simple critical value and a common contour enclosing precisely two roots? Replace the conservative constants by interval-certified bounds, and implement evaluation of t_c , ρ , the profiles, and the residual certificate without losing reliability when q is close to zero. A useful deliverable is a branch-aware evaluator with an absolute error guarantee, not merely a higher-order symbolic expansion.

Research question 11.2 (The fixed-coupling global continuation problem). Determine the singularity geometry of the inverse of $y + \log y + e^{-y}$ under its actual fixed coupling. Which critical points and asymptotic values obstruct continuation of the positive real inverse? How does the small-coupling chart here continue as λ approaches one? The relation $\lambda = we^w z$ must be enforced rather than treating z as independent at the end of a local argument.

Research question 11.3 (Two-stage truncation at infinity). For the real positive inverse, replace the exact logarithmic core w by a truncated power–logarithmic expansion while also truncating the exponential sectors. Prove a bound uniform in both truncation orders and specify how they may grow with the large argument. The present theorem is uniform near a critical auxiliary chart and does not settle the predecessor’s distinct two-stage uniformity question at infinity.

Research question 11.4 (Higher critical multiplicity). For analytic equations with $F_u = \dots = F_{u^{m-1}} = 0$ and $F_{u^m} \neq 0$, construct quantitative m -sheet profiles analogous to Theorem 5.1. Identify precisely which coefficients remain polynomial after a moving-discriminant normalization and when genuinely multivariate Newton polygons are unavoidable. Determine sharp support grid enlargements beyond the simple denominator doubling of a fold.

Editorial note (ProveIt, 2026-09-29). Still open. The countable-action package `Critical_Hahn_Transseries_Beyond_Finite_Action_Folds/` of the same series states explicitly that it does not settle this question: its critical boundary is a nonanalytic point with exponent $1/\alpha$, not an analytic critical point of higher multiplicity. The later exponent-two endpoint packages of the series (logarithmic, marginal, confluent and stable–Gaussian) study the logarithmic boundary singularity of the same kernel at $\alpha = 2$ and do not address it either; the finite-action folds of the logarithmic and marginal packages are ordinary square-root folds.

Research question 11.5 (Competing singularities and new coefficient windows). Extend the uniform large-sector theorem to a family in which two or more critical values have comparable modulus, or themselves collide. Derive joint limits that include interference among their coefficient contributions. The common single-singularity continuation disk used here will generally fail, so simply adding two fixed-parameter asymptotic formulas is not sufficient.

Research question 11.6 (Resurgence beyond analytic perturbation parameters). Replace convergent Taylor coefficients in the perturbation variables by Gevrey or resurgent coefficient series. Under what explicit hypotheses does moving-core inversion preserve Borel summability, and how are singular locations and Stokes constants transformed? The absolute convergence proof in Section 8 supplies none of these conclusions for a divergent input and should not be used as a substitute for them.

Research question 11.7 (Effective cancellation and minimal action grids). For an effectively represented analytic discriminant, design certificates for the first nonzero grouped action and for exact vanishing on a given exponential curve. In the rationally dependent case, identify useful algebraic classes where the resonance identities reduce to a finite ideal-membership problem. In the incommensurate case, specify exact comparison assumptions. Minimize the sufficient grid (55) without erasing cancellations or introducing negative generators.

Research question 11.8 (Actions depending on the large parameter). What remains true when $\mu_i = \mu_i(X)$ or when an external parameter makes distinct actions approach one another? The positive gap in the proof of Theorem 8.2 may cease to be uniform. Develop a joint scaling theory that distinguishes a genuine resonance from a near-resonance and identifies when a fixed finite support grid must be replaced by a parameter-dependent description.

Editorial note (ProveIt, 2026-09-29). Still open. The countable-action package `Critical_Hahn_Transseries_Beyond_Finite_Action_Folds/` keeps its actions fixed; its own research question 6 (incommensurate actions and shrinking gaps) proposes a different part of this question as further work.

Research question 11.9 (Proof-producing critical inversion). Build an implementation that emits a finite certificate for the root count, critical-value enclosure, polynomial-profile residual, and root-set error bound. A small trusted checker should reject unsupported sheet choices and unproved nonvanishing assumptions. Begin with the rational majorants and finite support-grid construction; then connect these certificates to a Lean formalization of the analytic hypotheses.

Research question 11.10 (Gamma and Barnes cores). Apply the method to analytic families genuinely underlying the repository’s Gamma and Barnes deepest-pole calculations. Determine the moving critical values and corresponding joint large-sector limits. Matching a quadratic coefficient recurrence is not enough: the analytic family, its continuation domain, and the nonvanishing singular amplitude must each be supplied and verified for those special functions.

12 Conclusion

The coefficientwise pole hierarchy and its formal square-root sum do not by themselves provide a uniform inverse near a critical point. For the mixed logarithmic–exponential sector equation, the missing object is now explicit: a moving discriminant, an exact two-sheet holomorphic atlas, and a convergent polynomial profile expansion with an error bound valid at the collision itself. This resolves the predecessor’s stated local critical-transition question within its independent-parameter family.

The same atlas controls a different limit that the fixed-sector pole calculation does not see. High sector index amplifies the cubic drift of the critical value, producing the nontrivial crossover $e^{-2\tau/3}$ when $s = \tau/n$. The proof identifies both the leading limit and its first finite- n correction.

For analytic folds with several exponential inputs, the decisive datum is the first nonzero *grouped* discriminant action. Its half controls branch splitting, while a constructive finite positive grid controls all higher terms. Resonances can delay the first splitting action or remove it altogether. The resulting series are absolutely convergent for the analytic finite-action class considered here.

These conclusions do not solve general transseries realization, unrestricted resurgence, or the global singularity geometry of the fixed-coupling original map. They do provide a proved, quantitative bridge from formal deepest poles to local analytic inversion, and a concrete next set of questions with their remaining hypotheses visible.

A Dependency map and source audit

The principal proof dependencies are

$$\text{explicit bounds} \implies \text{two-root contour atlas} \implies \begin{cases} \text{uniform polynomial profiles,} \\ \text{uniform high-sector transfer,} \end{cases}$$

followed by the explicit critical-value expansion for the crossover. The general action theorem uses a separate local fold-preparation lemma, absolute Taylor convergence, and the finite positive-gap construction. No general theorem of formal transseries composition or inversion is assumed in either chain.

The repository was inspected at commit

20301601562cf84173f2b503b0dc9c3fbf21cd0c.

The package README and `QuadraticCoreCatalan.lean` were read at that pinned revision. The large canonical transseries source could not be retrieved in full through the available repository fetch operation; therefore this manuscript does not claim a complete audit of that volume or the whole repository. The directly inspected status materials are sufficient for the limited formalization distinctions made here.

The predecessor source `reversion_and_one_exponential.tex` was retrieved independently from the user's saved document library. Its mixed-inverse equation, exact sector formula, leading-pole statement, uniform real-positive sector bound, and critical-transition research question were read directly. It is cited as a supplied research manuscript, not as an externally peer-reviewed result. The main local critical analysis in this article is proved from equation (3), so its validity does not require accepting any broader claims of that predecessor.

B Reproduction instructions

The package is self-contained for compilation: all tables used by the article are included. From the package directory, run

```
python -m pip install -r requirements.txt
python verify.py
latexmk -pdf -interaction=nonstopmode -halt-on-error article.tex
```

The verification script has no network dependency after installation. Its default run includes sector indices up to 400; `-quick` omits those final large-index diagnostics. The symbolic checks are identical in both modes. Re-running the script intentionally replaces the package's own `data/verification.json` and table fragments; it does not modify any repository or Library source. Editorial note (ProveIt, 2026-09-29): in the filed package a `-quick` run writes into `data/quick/` and leaves the full-run files that this article inputs unchanged, and every output is written with LF line endings.

The supplied PDF was compiled from this source and visually checked. The package records a mathematical proof manuscript and reproducible checks, not a claim of independent peer review, a formal verification certificate, or established worldwide priority.

References

- [1] Vladimir Reshetnikov, *ProveIt: Transseries_And_Inversion package README*, inspected at commit 20301601562cf84173f2b503b0dc9c3fbf21cd0c, 29 September 2026. [Pinned repository source](#). The cited status concerns the distinctions between finite algebra, assembled formal-series theorems, and analytic identification. Editorial note (ProveIt, 2026-09-29): the pinned link still resolves at the cited commit, but since the repository split the README is at `Analysis/Transseries/docs/series-and-transseries/Transseries_And_Inversion/README.md`, beside the canonical volume `transseries_and_inversion.tex`, which carries the three labels cited in the next entry.
- [2] Vladimir Reshetnikov, ProveIt module *QuadraticCoreCatalan.lean*, same pinned revision. [Pinned Lean source](#). Relevant volume labels are `p6:prop:quadratic-core-catalan`, `p6:lem:quadratic-core`,

and `p6:thm:deepest-pole`; the module states its formalization boundaries. Editorial note (ProveIt, 2026-09-29): the module stayed in the Fabius Lean library; the path above is current and the file is unchanged since the pinned commit.

- [3] *Support-Controlled Reversion and the Exact One-Exponential Substitution Group*, supplied research manuscript, 29 September 2026. Source file `reversion_and_one_exponential.tex`; retrieved from the user’s saved documents. In particular, the mixed inverse example, its deepest-pole resummation, and Research question 11.8. This is project-local predecessor material, not a claim of peer-reviewed publication. Editorial note (ProveIt, 2026-09-29): the predecessor is now filed in the same series as `Analysis/Transseries/docs/series-and-transseries/Support_Controlled_Reversion_One_Exponential/reversion_and_one_exponential.tex`.
- [4] R. M. Corless, G. H. Gonnet, D. E. G. Hare, D. J. Jeffrey, and D. E. Knuth, *On the Lambert W function*, *Advances in Computational Mathematics* **5** (1996), 329–359. [doi:10.1007/BF02124750](https://doi.org/10.1007/BF02124750). The branch-point expansion is discussed on pp. 350–351.
- [5] P. Flajolet and A. Odlyzko, *Singularity analysis of generating functions*, *SIAM Journal on Discrete Mathematics* **3**, no. 2 (1990), 216–240. [Author-hosted article](#). The half-integer coefficient identity and its asymptotic expansion are special cases of the coefficient formulas on p. 219.
- [6] G. A. Edgar, *Transseries for beginners*, *Real Analysis Exchange* **35** (2010), 253–310. [arXiv:0801.4877v5](https://arxiv.org/abs/0801.4877v5), revised 15 July 2009.