

Planar Eulerian orientation asymptotics

Exact Lambert sectors, logarithmic corrections and inverse indices

Report 171

3 October 2026

Abstract

We give explicit, reproducible refinements of the known asymptotics of the planar Eulerian orientation sequences [A324312](#), [A277493](#), and the quartic specialization [A324314](#). The leading equivalents and global analytic continuation are inputs from Bousquet-Mélou and Elvey Price. We derive six universal logarithmic correction polynomials, an arbitrary fixed-order decomposition into exact analytic Lambert models, and smooth inverse charts with rigorous asymptotic ceiling bounds. Keeping the entire first Lambert model reveals a relative correction asymptotic to $2/(m \log m)$, where $m = n + 2$; no fixed logarithmic truncation resolves this smaller scale. A bounded executable companion checks integer counts, independent Lagrange inversion, symbolic identities and numerical models. Floating-point diagnostics and asymptotic existence bounds are explicitly distinguished from certified finite-index inequalities.

[write] **Status and trust boundary, added 6 October 2026, batch 109.** This is bundle Report 171 of an external AI-assisted research session, filed in ProveIt as the single-source research report `a324312-planar-eulerian-orientations` (provenance, the sources as the write read them, what was checked, relation to the repository, the collected non-claims and the reading conventions are in [Section 1.1](#)). It is unrefereed and not formalized. Its author line and its PDF author field are empty, and the title block reads “Report 171”: it names no person or tool. *What rests on what.* Every statement about the counts is conditional on one published input that is not re-proved ([Lemma 6.1](#), about the model coefficients h_m alone, is not): the global Δ -continuation of the inverse series R with its leading singular equivalent, Bousquet-Mélou and Elvey Price’s Propositions 8.2 (quartic) and 8.3 (general). For the general family, which carries [A324312](#) and [A277493](#), its authors write that they “will not give any details” ([Remark 3.1](#)). Granted that input, [Theorems 4.1](#), [5.1](#), [6.2](#) and [7.1](#), like [Lemma 6.1](#), have conventional proofs (the hypergeometric continuation formula of the DLMF, a contraction argument for the Lambert branch, analytic implicit functions, and absolute-value contour bounds); no proof uses a computation. The explicit U_1, U_2, U_3 and $Q_0, \dots, Q_5$ are outputs of finite generators, computed by the shipped programs; the write rederived all of them with its own code. Every decimal is a diagnostic. The leading equivalents are Bousquet-Mélou and Elvey Price’s; the inverse-log and reciprocal-Gamma tools are Elvey Price and Guttman’s. The write adds [Remarks 1.1](#), [3.1](#) and [7.2](#) (the OEIS entries quoted; the analytic input as published, with a citation note; the inverses against the transseries volume), dated notes, and one question in [Section 9](#). No statement, proof or number of the source was changed.

Contents

1	Scope and principal results	2
1.1	Provenance, sources and reading conventions	4

2	Exact objects and normalizations	6
3	Analytic input and local singular equation	7
4	Exact analytic Lambert models	9
4.1	A sufficient continuation domain	9
4.2	The first two sectors	9
5	Any fixed number of exact sectors	10
5.1	Joint analytic reversion and branch identification	10
5.2	Triangular sector generator	10
5.3	The full coefficient remainder argument	11
6	Universal logarithmic correction polynomials	12
6.1	Hankel transfer with a finite derivative expansion	12
6.2	Finite generator and explicit formulas	13
7	Smooth inverse charts and integer thresholds	14
8	Reproducible computations and numerical use	17
8.1	Independent exact checks	17
8.2	Numerical coefficient extraction of the exact models	17
8.3	Reproducibility and limits	18
9	Further research directions	18
A	The fourth local sector	20
B	Source map and theorem dependencies	20

1 Scope and principal results

A planar map is a connected graph embedded in the oriented sphere, considered up to orientation-preserving homeomorphism. Loops and multiple edges are allowed. An Eulerian orientation directs every edge so that the indegree and outdegree agree at every vertex. The general family is counted by edges; the quartic family, in which every vertex has degree four, is counted by vertices.

The essential exponential rates are 4π and $4\sqrt{3}\pi$, with a factor $n^{-2}(\log n)^{-2}$. These leading results are already theorems of Bousquet-Mélou and Elvey Price [1], not new conclusions about previously unknown growth. Our purpose is to make the slower logarithmic corrections, smaller algebraic sectors and inverse problem explicit and reproducible.

For either family, write $m = n + 2$, $L = \log m$, and $T = \log L + C + \gamma$. The constants κ_0, μ, C

are specified in Section 2. The three useful levels of approximation are

$$a_n = \frac{\kappa_0 \mu^m}{m^2 L^2} \left\{ \sum_{j=0}^J \frac{Q_j(T)}{L^j} + \mathcal{O}_J \left(\frac{(\log L)^{J+1}}{L^{J+1}} \right) \right\}, \quad (1.1)$$

$$a_n = \kappa_0 \mu^m \left\{ h_m - \frac{1}{2} k_m + \mathcal{O}(m^{-4} L^{-3}) \right\}, \quad (1.2)$$

$$\frac{a_n}{\kappa_0 \mu^m h_m} = 1 + \frac{2}{mL} (1 + o(1)). \quad (1.3)$$

Here J is any fixed nonnegative integer, $Q_0 = 1$, $Q_1 = 3 - 2T$, and h_m, k_m are coefficients of actual analytic functions, not of a truncated logarithmic series. The exact definitions and all fixed-order error arguments appear below. Equation (1.3) is relative to the exact coefficient model h_m ; it must not be read as the next correction after a finite sum in (1.1).

Attribution and boundary. The enumeration, leading equivalents, analytic logarithmic inversion and Hankel–Gamma transfer have substantial antecedents [1, 2, 3, 5]. The contribution here is a worked refinement: the displayed Q_0 – Q_5 , a constructive exact-sector organization, and rounding-aware inverse charts with an executable companion. The particular explicit formulas were not located in the inspected primary sources; this bounded inspection is not a claim of universal priority. The refined enumeration preprint [4], inspected as version 3 of 2 June 2026, discusses a predicted general-weight singularity picture in Section 6.4. We prove no general-weight phase diagram. The retrieved OEIS versions on 3 October 2026 supplied exact descriptions but no asymptotic formulas; this dated observation does not certify their latest uncached revisions.

Remark 1.1 ([write] The OEIS entries, quoted, 6 October 2026). The entries were read again on 6 October 2026 in the OEIS internal format; the quotations are verbatim.

- (a) A324312 (revision #18, 2 October 2025), offset 1: “Expansion of the generating function of rooted planar Eulerian orientations, counted by edges.”, data 1, 5, 33, 252, 2108, ... (22 terms), formulas “G.f.: $(1/(4t^2)) * (t - 2t^2 - R(t))$ where $R(t)$ is A324311.” and “ $a(n) = A277493(n)/2$. – _Andrew Elvey Price_, Dec 12 2024”.
- (b) A277493 (revision #53, 21 December 2024), offset 0: “Number of planar rooted Eulerian orientations (i.e., planar directed Eulerian maps with in-degree and out-degree equal for each vertex) with n edges.”, data 1, 2, 10, 66, 504, ... (23 terms; “ $a(16)$ onwards added by _Andrew Howroyd_, Dec 12 2024”), formulas “G.f.: $(1/(2t^2)) * (t - R(t))$ where $R(t)$ is the g.f. of A324311.” and “ $a(n) = 2 * A324312(n)$ for $n > 0$.” (Elvey Price, 12 December 2024).
- (c) A324314 (revision #7, 22 February 2019), offset 1: “Expansion of the generating function of quartic rooted planar Eulerian orientations, counted by vertices.”, data 4, 35, 402, 5334, ... (20 terms), formula “G.f.: $(1/(3t^2)) * (t - 3t^2 - R(t))$ where $R(t)$ is A324313.”.

These are the three generating functions of (2.2). No entry states an asymptotic formula, as the source found on 3 October 2026. The write computed r_m , $m \leq 40$, by its own exact Lagrange inversion of (2.1) (rational arithmetic), found the residual of (8.1) zero through t^{38} for both families, and compared $-r_{n+2}/d_0$ with every printed term (22, 22 positive-index and 20): all agree, and

A277493 is twice A324312 on every printed positive index. The shipped fixture transcribes 22, 5 and 20 of these terms; all agree with the live data. Nothing was submitted to the OEIS.

1.1 Provenance, sources and reading conventions

[write] **Added 6 October 2026, batch 109.** *Provenance.* This report is Report 171 of the session bundle that ProveIt’s docs/incoming intake processed as batch 109: the archive `Planar_Eulerian_Orientations_Lambert_Expansions_and_Inverses_Source.zip` (3,455,433 bytes, SHA-256 `f0b45e63f38b...482a01cf2c7d1`, 25 files, no wrapper directory), arrival commit `60f54ea06`, placed in `f7e9e5c2f`. The text is the delivered `Report171.tex` (564 lines, dated 3 October 2026), shipped as `article.tex`. The programs are shipped under `code/` (the root `build.py` as `code/build.py`), the fixtures and recorded outputs as `data/fixtures-*` and `data/outputs-*`, and `BUILD-INFO.json` and `requirements.txt` under `data/` (the report README maps every name). Not shipped, and retrievable from the arrival commit: the delivered 14-page PDF; the checksum manifest `SHA256SUMS.json` (24 entries, verified at placement); the delivered README, replaced by the report README; and `outputs/exact.json` (3,024,729 bytes), which no shipped program reads and which the companion regenerates byte for byte (the README gives the command; at the write it took 18 s). The fixture-hash file `fixtures/SHA256.json` is shipped as `data/fixtures-SHA256.json` because the companion reads it. The package records no ProveIt commit and names no repository path; it carries no “prepared for private review” line and no personal data. It is a single-source report, so no merge choice was made. The write prefixed the 58 delivered labels with `peo:` and updated the 45 references to them; it added labels to Sections 1 and 9, to Appendix B and to Sections 5.1 and 6.2, and added this subsection, the notes marked [write], Remarks 1.1, 3.1 and 7.2, and one question in Section 9. The added statements are the last of their sections, this subsection follows the last delivered text of Section 1, and no display, table or statement was added before a delivered one, so every theorem, section, equation and table of the source keeps its delivered number.

The sources, as the write read them (6 October 2026).

- OEIS A324312, A277493 and A324314, in the internal format (Remark 1.1).
- Bousquet-Mélou and Elvey Price [1], arXiv:1803.08265v2 (the version the source names as inspected): Theorems 1.1 and 1.2, the root convention of Section 2, Section 7, and Section 8 (the differential equations of Section 8.1 and Propositions 8.2 and 8.3 of Section 8.2, with the sentences around them); Remark 3.1.
- The refined enumeration [4]: the abstract page (versions 1, 19 March 2025; 2, 1 June 2026; 3, 2 June 2026) and, in version 3, the title and opening of Section 6.4, “Predicted singular behaviour”.
- The transseries volume `Analysis/Transseries/docs/series-and-transseries/Transseries_And_Inversion/transseries_and_inversion.tex`: `p0:thm:lambert-core`, `p0:thm:core-reversion`, `p0:def:three-inverses`, `p0:thm:staircase`, `plt:def:lw-monomial-log-datum` and `plt:thm:lw-template`, read for Remark 7.2.
- Not read by the write: the published version of [1], Bousquet-Mélou and Courtiel [2], Elvey Price and Guttmann [3], and the DLMF pages [5]; the source’s account of them is reported, not confirmed (the constants H , C , δ_j that the source derives from the DLMF formula were recomputed from the digamma definition (3.4)).

What was checked at intake. The batch-109 dossier read the manuscript in full and found no error; it checked the relative correction $2/(mL)$ of (1.3) from the leading terms of (4.6) and the contour argument of Theorem 5.1, and it reran the companion on copies (`exact` in about 50 s, byte-identical to the delivered `outputs/exact.json`; `symbolic`, `diagnostics` and the three `inverse` runs identical; the `test_companion.py` receipt equal as JSON; `test_build.py` fails on Windows, which writes its synthetic files with CRLF). *What the write checked.* Every proof line by line, with no mathematical error found: Table 1 (r_0 from the hypergeometric form of Ω' , $\rho = \Omega(r_0)$ numerically, A , κ_0 and the amplitudes π^2 , $2\pi^2$, $8\pi^2/3$); H , C and $\delta_1, \dots, \delta_4$ from the digamma definition (3.4) (both families, matching Appendix A); the contraction argument

of Section 4 ($\operatorname{Re} \Lambda > C - \log S_0 = 1$ on D_s); the joint equation (5.1), q_1 and (5.4); the contour estimates of the proof of Theorem 5.1; the reciprocal-Gamma identities ($1/\Gamma(-1-w) = w(1+w)/\Gamma(1-w)$), and the coefficient -2 of w in $1/\Gamma(-2-w)$ and the Hankel integral of Lemma 6.1; $-f' = 1/(v(v-1))$ and the two equivalents of (4.6); E_1 , E_2 ; the chart (7.5); and (8.4), (8.5). With its own code (SymPy 1.14.0, mpmath 1.3.0) it rederived $Q_0, \dots, Q_5$ from the generator (6.7) and the coefficients c_j , and U_1, U_2, U_3 of both families from (5.6): all as printed; it checked (8.4) and (8.6) against direct Taylor coefficients of F_0 (agreement to 10^{-52}), and recomputed Table 2, the “about 165% and 185%” and the three values of $x_J(y) - 1000$ (note at the end of Section 8). It reran **exact**, **symbolic**, **diagnostics**, the three **inverse** runs and **test_companion.py** from the shipped files (README, Route B): all outputs byte-identical, the test receipt equal as JSON. Finite and floating computations are observations, not certificates.

Relation to the repository. Before batch 109 no file of the repository named A324312, A277493 or A324314, and no report counted planar Eulerian orientations. The reports **a110058-square-contingency-tables** and **a380592-tied-football-seasons**, in the same directory **SetTheory/Cardinals/docs/reports/generating-functions-and-asymptotics/oeis-sequence-asymptotics/**, cite a cumulant expansion for counting Eulerian orientations of a fixed graph, a different object; no result is shared. No statement of this report is formalized in Lean or Rocq, and its place in the collection confers no formal status. The inverses of Sections 4–7 are compared with the transseries volume in Remark 7.2.

[write] **What is not claimed, collected from the source.** The leading equivalents and the global analytic continuation are Bousquet-Mélou and Elvey Price’s and are not re-proved; for the general family their published proposition omits the proof details. The enumeration, the analytic logarithmic inversion and the Hankel–Gamma transfer have antecedents, and the analytic tools of Elvey Price and Guttmann are not claimed as new; the explicit formulas “were not located in the inspected primary sources; this bounded inspection is not a claim of universal priority”. No general-weight phase diagram is proved; the refined preprint’s prediction is motivation only. Theorem 5.1 is a fixed- K statement: no convergence of the sector sum as $K \rightarrow \infty$, no uniformity in growing K , no optimal truncation and no complete transseries with exponentially small terms. No effective constants or starting indices are supplied; no uniform $o(1)$ approximation of $N(y)$ and no formula $N(y) = \lceil x_J(y) \rceil$ is claimed, and exact finite-index identification needs exact counts. Universality of Q_j is only across the two inverse hypergeometric families. The diagnostics use no interval arithmetic; finite checks do not certify the analytic remainders. The OEIS observations are dated. The write adds: its own checks are floating or finite; Remark 3.1 corrects one citation and no mathematics; Remark 7.2 claims no novelty for any inversion.

Reading conventions (letters the source reuses, with the tempting false readings; no symbol of the source was renamed; symbols of the transseries volume carry the subscript “vol” in Remark 7.2).

Symbol	Meaning here (where)	Not to be read as
a, a_n, a_j	the family parameter $\frac{1}{2}, \frac{1}{3}$; the counts; the ${}_2F_1$ coefficients (3.4)	each other
h_m, h_j, h	$[z^m]F_0$ (4.4); digamma combinations (3.4); the model F_0 in (8.4)	each other; the volume’s h_{vol}
$q, q_\ell, q(\eta, w)$	Y/b (Section 7); the reversion unknown and its coefficients (Section 5); also u/η	each other
C, c_j, c	$1 + \log 4$ or $1 + \log 6$; reciprocal-Gamma coefficients (6.1); $\text{Re } \Lambda$, and the ODE constant $c \in \{16, 27\}$	each other
δ, δ_j	δ_2 ; the local constants (3.5)	each other
A, α, κ_0	r_0/α ; $a(1-a)$; A/d_0	sequence names; the volume’s κ_{vol}
T, t, T_q	$\log L + C + \gamma$; the series variable, also $\log L + C$ in (6.7) and $z = 1 - t/m$ in Lemma 6.1; T at L_q	each other
$L, \mathcal{L}_J, L_q$	$\log m$; the comparison (7.3); $\log q$	the volume’s L_{vol}
w, W_{-1}	$1/v$ (Section 5); in (6.7) $v - L - t$; in (6.1) a formal variable; Lambert’s function	each other
D, D_s	$-x^2\partial_x + x\partial_t$; the slit disk (4.1)	each other
M_j, M_J	sector models (Theorem 5.1); the comparison root (Section 7)	each other
E_K, E_j	the coefficient remainder; inverse-log coefficients	each other
K, K_J, k, k_m	sector order; the constant of Theorem 7.1; $\sin(\pi a)/\pi$; $[z^m]F_1$	each other
ρ, μ, r, r_m	the radius, ρ^{-1} ; the Δ -radius and the variable of Ω ; $[t^m]R$	the volume’s $\rho_{\text{vol}}, \mu_{\text{vol}}$

2 Exact objects and normalizations

The root convention matters. In the paper [1], the orientation of the root edge must agree with the direction used to root the map. This is the convention for A324312 and A324314. A277493 allows both root-edge orientations and has twice the positive-index counts of A324312. The size-zero value of A277493 is 1 and is treated separately.

Let $R(t) = \sum_{m \geq 1} r_m t^m$ be the compositional inverse of Ω , with $R(0) = 0$. The defining series are

$$\Omega_g(r) = \sum_{j \geq 0} \binom{2j}{j}^2 \frac{r^{j+1}}{j+1}, \quad \Omega_q(r) = \sum_{j \geq 0} \binom{2j}{j} \binom{3j}{j} \frac{r^{j+1}}{j+1}. \quad (2.1)$$

The generating functions from [1, Theorems 1.1–1.2] and the OEIS entries are

$$\begin{aligned}\sum_{n \geq 1} \text{A324312}(n)t^n &= \frac{t - 2t^2 - R_g(t)}{4t^2}, \\ \sum_{n \geq 0} \text{A277493}(n)t^n &= \frac{t - R_g(t)}{2t^2}, \\ \sum_{n \geq 1} \text{A324314}(n)t^n &= \frac{t - 3t^2 - R_q(t)}{3t^2}.\end{aligned}\tag{2.2}$$

Consequently, for $n \geq 1$ and always with $m = n + 2$,

$$a_n = -r_m/d_0, \quad d_0 = 4, 2, 3 \quad \text{for A324312, A277493, A324314 respectively.} \tag{2.3}$$

Their first positive-index terms are 1, 5, 33, 252; 2, 10, 66, 504; and 4, 35, 402, 5334. The index origins are 1, 0 and 1 respectively. The second component of an OEIS offset is metadata, not an additional index shift.

Parameter	General	Quartic
a	$1/2$	$1/3$
$\alpha = a(1 - a)$	$1/4$	$2/9$
r_0	$1/16$	$1/27$
ρ	$1/(4\pi)$	$1/(4\sqrt{3}\pi)$
$\mu = \rho^{-1}$	4π	$4\sqrt{3}\pi$
$A = r_0/\alpha$	$1/4$	$1/6$
C	$1 + \log 4$	$1 + \log 6$
δ	$-5/2$	-3

Table 1: Family parameters. r_0 is the limiting value of R at $t = \rho$, not a coefficient r_m .

The amplitude $\kappa_0 = A/d_0$ is $1/16, 1/8, 1/18$ in the order of (2.3). Thus

$$a_n \sim \frac{\kappa_0 \mu^{n+2}}{(n+2)^2 \log^2(n+2)}.\tag{2.4}$$

If only a leading equivalent is wanted, replacing $n+2$ by n gives amplitudes $\pi^2, 2\pi^2, 8\pi^2/3$ multiplying $\mu^n/(n^2 \log^2 n)$. Such replacement is unsuitable when resolving algebraic-order corrections.

3 Analytic input and local singular equation

We use the following published global input. By [1, Propositions 8.2 and 8.3], the quartic and general inverse series respectively continue analytically into a domain

$$|t| < r, \quad t \neq \rho, \quad |\text{Arg}(t - \rho)| > \phi, \quad r > \rho, \quad 0 < \phi < \pi/2,\tag{3.1}$$

and have the stated leading singular equivalents there. In particular, the domain includes every other point of $|t| = \rho$. This is a global continuation assertion, stronger than a local sector expansion. The quartic result uses the earlier spanning-forest analysis [2, Proposition 8.4]; for the general proposition, the authors of [1] omit proof details and refer to the preceding argument. We take their published proposition as input, and do not independently re-prove global continuation in this report.

Set

$$z = t/\rho, \quad s = 1 - z, \quad u = 1 - R(t)/r_0, \quad k = \frac{\sin(\pi a)}{\pi}. \quad (3.2)$$

All local logarithms are continued from positive s, u near zero. Taylor coefficient matching in (2.1) gives

$$\Omega'(r) = {}_2F_1(a, 1 - a; 1; r/r_0). \quad (3.3)$$

The zero-balanced hypergeometric continuation formula [5, 15.8.10, $m = 0$] gives

$$\Omega'(r_0(1-u)) = k \sum_{j \geq 0} a_j u^j (-\log u + h_j), \quad a_j = \frac{(a)_j (1-a)_j}{(j!)^2}, \quad h_j = 2\psi(j+1) - \psi(j+a) - \psi(j+1-a). \quad (3.4)$$

It converges for $|u| < 1$, $|\arg u| < \pi$, uniformly on compact subsectors. The regularized hypergeometric notation in the cited formula causes no extra factor here, because the lower parameter is 1. Integrating to the singular point yields

$$s = \alpha \sum_{j \geq 1} b_j u^j (-\log u + H + \delta_j), \quad b_j = \frac{a_{j-1}}{j}, \quad H = h_0 + 1, \quad \delta_j = h_{j-1} - h_0 + \frac{1}{j} - 1. \quad (3.5)$$

Indeed $r_0 k/\rho = \alpha$ and $\int_0^u x^{j-1} (-\log x + h) dx = u^j (-\log u + h + 1/j)/j$. The endpoint logarithm is integrable. Here $b_1 = 1$, $b_2 = \alpha/2$, $\delta_1 = 0$, and

$$\delta_2 = \delta = \frac{3}{2} - \frac{1}{\alpha}, \quad H = \begin{cases} 1 + 4 \log 2 & \text{general,} \\ 1 + 3 \log 3 & \text{quartic,} \end{cases} \quad C = H + \log \alpha. \quad (3.6)$$

These identities follow from digamma reflection and multiplication; for example, $h_1 - h_0 = 2 - 1/a - 1/(1-a) = 2 - 1/\alpha$.

Remark 3.1 ([write] The analytic input as published, 6 October 2026). In arXiv:1803.08265v2 of [1], Section 8.2 defines a Δ -domain of radius ρ exactly as (3.1) and states: Proposition 8.2, for the quartic series R of its Theorem 1.1, radius $\rho = \sqrt{3}/(12\pi)$, analyticity in a Δ -domain, and $R(t) - \frac{1}{27} \sim \frac{1}{6}(1 - t/\rho)/\log(1 - t/\rho)$ there, introduced by “the first part of the following result is the case $u = -1$ of [17, Prop. 8.4]” ([17] is [2]); and Proposition 8.3, for the general series of its Theorem 1.2, radius $\rho = 1/(4\pi)$ and $R(t) - \frac{1}{16} \sim \frac{1}{4}(1 - t/\rho)/\log(1 - t/\rho)$, followed by “The proof closely follows the proof of Proposition 8.2 given in [17, Sec. 8.3], and we will not give any details.” So the source’s account in this section is accurate, and its trust boundary is real: every statement here about A324312 and A277493 rests on a proposition whose proof is only indicated (question 6). The two leading forms agree with this report’s: $F_0 = s/v$ and $v \sim -\log s$ give $R(\rho z) - r_0 \sim As/\log s$, with $A = \frac{1}{4}, \frac{1}{6}$ as in Table 1, and $\sqrt{3}/(12\pi) = 1/(4\sqrt{3}\pi)$. *A citation note.* The source attributes the differential equations (8.1) to “[1, Section 7]” (Section 8 and Appendix B). In arXiv:1803.08265v2 they are in Section 8.1, “Nature of the series” (as $R(27R - 1)R'' = 6tR'^3$ and $R(16R - 1)R'' = 4tR'^3$); Section 7 solves the functional equations of general orientations and contains no such equation. The published version was not read. The equations themselves are right: the write checked them to order t^{38} (Remark 1.1), and the companion to t^{1001} . (Added after the independent check of 7 October 2026: the equations are in Section 8.1 in every version available, arXiv:1803.08265v1 (22 March 2018) as well as v2, and the HAL deposit `hal-01738160v2` (5 November 2019), which HAL cites as the JCTA paper and whose Section 8 is word for word that of v2; the journal’s typeset version was still not read. Proposition 8.4 of [2] is in its Section 8.3, “When $u < 0$ ”, and covers $u \in [-1, 0)$, as BMEP’s two references to it require.)

4 Exact analytic Lambert models

4.1 A sufficient continuation domain

Put $S_0 = e^{C-1}$, equal to 4 or 6, and define

$$D_s = \{s : 0 < |s| < S_0, |\operatorname{Arg} s| < \pi\}. \quad (4.1)$$

For $s \in D_s$, set $\Lambda = C - \operatorname{Log} s$ and let v be the solution

$$v - \operatorname{Log} v = \Lambda. \quad (4.2)$$

This defines the continuation of the real solution $v > 1$; on the positive real axis it is conventionally written $-W_{-1}(-e^{-C}s)$. Away from that axis the implicit continuation is our definition, avoiding software conventions for the two lips of the Lambert branch [5, 4.13].

For completeness, let $c = \operatorname{Re} \Lambda > 1$. On the closed complete half-plane $\operatorname{Re} v \geq c$, the map $v \mapsto \Lambda + \operatorname{Log} v$ takes values in that half-plane because $\operatorname{Re}(\Lambda + \operatorname{Log} v) \geq c + \log c > c$. Integration of $1/v$ along a straight segment shows its Lipschitz constant is at most $1/c < 1$. The contraction theorem gives a unique fixed point there. The derivative $1 - 1/v$ is nonzero, so the analytic implicit-function theorem gives local analytic dependence; uniqueness glues these solutions on D_s . In particular $\operatorname{Re} v > 1$.

Choose any $1 < r_* < S_0 - 1$. Then $|z| < r_*$, $z \notin [1, \infty)$ implies $1 - z \in D_s$. All rational functions of v with possible finite poles only at 0 and 1 therefore give analytic models on that slit disk. Intersect it with (3.1), after scaling by ρ . This common domain is all that the coefficient proofs require; no larger slit-plane continuation is asserted.

In a smaller closed Δ -sector at $s = 0$, $\Im \Lambda$ is bounded, as is $\Im v = \Im \Lambda + \arg v$. The fixed-point equation gives

$$v = \Lambda + \mathcal{O}(\log \Lambda), \quad |v| \asymp \log(1/|s|), \quad (4.3)$$

uniformly there.

4.2 The first two sectors

Define the analytic functions

$$F_0(z) = \frac{1-z}{v(z)}, \quad F_1(z) = \frac{(1-z)^2(v(z) + \delta)}{v(z)^2(v(z) - 1)}, \quad h_m = [z^m]F_0, \quad k_m = [z^m]F_1. \quad (4.4)$$

They are analytic at $z = 0$, with the specific branch just defined. “Exact model” means these function and coefficient definitions are exact. Finite-precision extraction of their coefficients is a numerical approximation to that definition.

Theorem 4.1. *For either family, as $m \rightarrow \infty$,*

$$r_m \rho^m = -A h_m + \frac{A}{2} k_m + \mathcal{O}(m^{-4}(\log m)^{-3}). \quad (4.5)$$

Thus (1.2) holds for each of the three sequences with the constants of Section 2. Moreover,

$$h_m \sim \frac{1}{m^2 \log^2 m}, \quad k_m \sim -\frac{4}{m^3 \log^3 m}, \quad (4.6)$$

and (1.3) follows.

The local inversion and coefficient remainder are proved in the next section. The leading coefficient estimates in (4.6) are proved in Section 6. Notice that the F_1 contribution is positive in the count, although its coefficients are asymptotically negative.

5 Any fixed number of exact sectors

5.1 Joint analytic reversion and branch identification

Set $\eta = s/(\alpha v)$, $w = 1/v$, and $u = \eta q$. Equation (4.2) implies $-\log \eta + H = v$. Dividing (3.5) by $s = \alpha \eta v$ gives

$$1 = q(1 - w \log q) + \sum_{j \geq 2} b_j \eta^{j-1} q^j \{1 + w(\delta_j - \log q)\}. \quad (5.1)$$

This is a genuinely analytic implicit equation, not just a formal one. Gamma and digamma asymptotics give $a_j = \mathcal{O}((j+1)^{-1})$, $b_j = \mathcal{O}(j^{-2})$ and $\delta_j = \mathcal{O}(1)$. With q in a small disk about 1 and its analytic logarithm, the sum and all its derivatives converge normally on smaller product neighborhoods satisfying $|\eta q| < r < 1$. It is jointly analytic near $(\eta, w, q) = (0, 0, 1)$.

The derivative of the right side in q is 1 at that base point, and at $(0, w, 1)$ it is $1 - w$. Hence there is a unique analytic solution $q(\eta, w)$ near $(0, 0)$, with $q(0, w) = 1$. For each fixed $K \geq 1$ it has the uniform expansion

$$q(\eta, w) = 1 + \sum_{\ell=1}^{K-1} q_\ell(w) \eta^\ell + \mathcal{O}_K(\eta^K). \quad (5.2)$$

The leading singular equivalent furnished by the published analytic input gives $u/\eta \rightarrow 1$ uniformly in the smaller Δ -sector. The branch of the actual inverse R therefore lies in this uniqueness neighborhood and is the analytic solution just constructed. This identifies the implicit expansion with the desired generating function.

At first order,

$$q_1(w) = -\frac{\alpha}{2} \frac{1 + \delta w}{1 - w}, \quad u = \eta - \frac{\alpha}{2} \eta^2 \frac{v + \delta}{v - 1} + \mathcal{O}(\eta^3). \quad (5.3)$$

Consequently

$$R(\rho z) = r_0 - A F_0(z) + \frac{A}{2} F_1(z) + \mathcal{O}(s^3/v^3). \quad (5.4)$$

5.2 Triangular sector generator

Let $\xi = s/\alpha$. There are rational functions $U_\ell(v)$ such that, for each fixed K ,

$$u = \sum_{j=1}^K \xi^j U_{j-1}(v) + \mathcal{O}_K((s/v)^{K+1}), \quad U_0(v) = 1/v. \quad (5.5)$$

A finite generator substitutes $p(\xi) = \sum_{\ell \geq 0} \xi^\ell U_\ell(v)$ into

$$1 = \sum_{j \geq 1} b_j \xi^{j-1} p(\xi)^j \{v - \log(vp(\xi)) + \delta_j\}. \quad (5.6)$$

At coefficient ξ^ℓ the new U_ℓ has multiplier $v - 1$. This makes the recurrence triangular. Equivalently, $U_{j-1}(v) = v^{-j} q_{j-1}(1/v)$ with $q_0 = 1$. The induction in (5.1) divides only by $1 - w$, so each q_ℓ is

rational with possible finite pole only at $w = 1$. Accordingly U_ℓ has possible finite poles only at $v = 0, 1$.

For reference, the first nontrivial functions are

$$\begin{aligned} U_1^g(v) &= -\frac{2v-5}{16v^2(v-1)}, & U_1^q(v) &= -\frac{v-3}{9v^2(v-1)}, \\ U_2^g(v) &= -\frac{8v^3-20v^2-20v+23}{512v^3(v-1)^3}, & U_2^q(v) &= -\frac{12v^3-35v^2-38v+43}{729v^3(v-1)^3}. \end{aligned} \quad (5.7)$$

The next functions U_3 are given in Appendix A and checked independently by substitution in the companion.

Theorem 5.1 (Fixed-sector coefficient expansion). *For each fixed integer $K \geq 1$ define actual analytic functions $M_j(z) = (1-z)^j U_{j-1}(v(z))$, $1 \leq j \leq K$. Then*

$$r_m \rho^m = -r_0 \sum_{j=1}^K \alpha^{-j} [z^m] M_j(z) + \mathcal{O}_K \left(m^{-K-2} (\log m)^{-K-1} \right). \quad (5.8)$$

All implied constants may depend on the family and the fixed order K .

This is a fixed- K coefficient theorem. The joint analytic reversion gives a locally convergent series near $s = 0$, but does not justify exchanging its infinite sum with coefficient extraction at $z = 0$. In particular, the theorem does not establish convergence of the coefficient-sector sum as $K \rightarrow \infty$, uniformity when K grows with m , optimal truncation, or a complete transseries including exponentially small terms. It supplies a rigorous algebraic-sector hierarchy in exact Lambert models.

5.3 The full coefficient remainder argument

Proof of Theorem 5.1. The analytic reversion above gives (5.5). Subtract the corresponding analytic models on the common global Δ -domain and write

$$E_K(z) = R(\rho z) - r_0 + r_0 \sum_{j=1}^K \alpha^{-j} M_j(z). \quad (5.9)$$

For small $s = 1 - z$ in that domain,

$$|E_K(z)| \leq B_K |s|^{K+1} (\log(1/|s|))^{-K-1}. \quad (5.10)$$

Choose $\pi/2 < \theta < \pi - \phi$ in a common smaller domain. Deform the coefficient contour into an arc $|s| = 1/m$, rays $s = re^{\pm i\theta}$ up to a fixed sufficiently small radius ε , and closing curves away from $z = 1$ with $|z| > 1 + \varepsilon_1$.

On the small arc the Cauchy kernel $|z|^{-m-1}$ is bounded, the arc length is $\mathcal{O}(m^{-1})$, and (5.10) is $\mathcal{O}_K(m^{-K-1}(\log m)^{-K-1})$. Its contribution is of the asserted order. On the rays put $r = t/m$. For a small enough terminal radius, $|1 - s|^{-m-1} \leq Be^{-ct}$ with fixed $c > 0$. For $1 \leq t \leq \sqrt{m}$, $\log(1/|s|) \geq (\log m)/2$, so the absolute integral is bounded by a constant times

$$m^{-K-2} (\log m)^{-K-1} \int_1^\infty e^{-ct} t^{K+1} dt. \quad (5.11)$$

The remaining ray portions $\sqrt{m} \leq t \leq \varepsilon m$ are exponentially small: the remainder is bounded on the fixed local contour, and the kernel decays exponentially. The closing curves are also exponentially small by analyticity and $|z| > 1 + \varepsilon_1$.

Thus $[z^m]E_K = \mathcal{O}_K(m^{-K-2}(\log m)^{-K-1})$. The constant r_0 has zero coefficient for $m \geq 1$, proving (5.8). Taking $K = 2$ and using (5.4) proves (4.5). $\square$

The proof uses absolute-value bounds. It remains valid when the singular power $K + 1$ is an integer, and does not assume an extra cancellation at a reciprocal-Gamma zero for an unspecified remainder. No effective constants or starting indices are supplied.

6 Universal logarithmic correction polynomials

6.1 Hankel transfer with a finite derivative expansion

Let $f(\ell) = 1/v$, where $v - \log v = \ell + C$ in the large-real branch, and put

$$c_j = [w^j] \frac{1}{\Gamma(-1-w)}, \quad \frac{1}{\Gamma(-1-w)} = w(1+w) \exp\left(-\gamma w - \sum_{k \geq 2} \frac{\zeta(k)}{k} w^k\right). \quad (6.1)$$

In particular $c_0 = 0$, $c_1 = 1$, $c_2 = 1 - \gamma$.

Lemma 6.1. *For every fixed integer $P \geq 1$, with $L = \log m$,*

$$h_m = \frac{1}{m^2} \sum_{j=1}^P c_j (-1)^j f^{(j)}(L) + \mathcal{O}_P(m^{-2}L^{-P-2}) + \mathcal{O}_P(m^{-3}L^{-2}). \quad (6.2)$$

Proof. Use the same indented contour as above and put $z = 1 - t/m$. Use the Hankel contour $\mathcal{H}$ that runs inward on the lower ray of argument $-\theta$, counterclockwise around the circle of radius 1, and outward on the upper ray of argument θ . With this orientation the leading integral has amplitude $m^{-2}e^t t f(L - \log t) dt$. The operator identity

$$\frac{d}{d\ell} = \frac{v}{v-1} \frac{d}{dv} \quad (6.3)$$

shows $f^{(j)}(\ell) = \mathcal{O}_j((\operatorname{Re} \ell)^{-j-1})$ uniformly on the complex neighborhoods used here. On $1 \leq |t| \leq \sqrt{m}$, the straight interpolation between L and $L - \log t$ has real part at least $L/2$ and bounded imaginary part. Taylor expansion to order P therefore has remainder $\mathcal{O}_P(L^{-P-2}(1 + |\log t|)^{P+1})$. The same bound holds on the small arc. Integration against the exponentially decreasing ray kernel bounds its contribution by $\mathcal{O}_P(m^{-2}L^{-P-2})$.

The discretization error needs one extra step. Subtract the polynomial $(1 - z)f(L)$ before replacing the Cauchy kernel; its coefficient is exactly zero for $m > 1$. The remaining amplitude satisfies

$$f(L - \log t) - f(L) = \mathcal{O}\left(L^{-2}(1 + |\log t|)\right) \quad (6.4)$$

on the central portion. There,

$$(1 - t/m)^{-m-1} = e^t \left\{ 1 + \mathcal{O}\left(\frac{|t| + |t|^2}{m}\right) \right\}, \quad (6.5)$$

with a uniform exponentially decaying majorant on the rays, after reducing the terminal radius if necessary. Multiplying by the subtracted amplitude and integrating gives $\mathcal{O}(m^{-3}L^{-2})$. Without polynomial subtraction this argument would yield only the weaker inverse-log bound.

The portions beyond $|t| = \sqrt{m}$ and the outer closing contour are exponentially small. One may therefore complete each central logarithmic moment to the full Hankel contour. The reciprocal-Gamma integral gives

$$\frac{1}{2\pi i} \int_{\mathcal{H}} e^t t (\log t)^j dt = j! c_j. \quad (6.6)$$

Taylor's factor $(-1)^j/j!$ produces $c_j(-1)^j f^{(j)}(L)$. The constant term vanishes since $c_0 = 0$, completing the proof. $\square$

This lemma is a finite-order derivative formula. It does not assert convergence of the infinite derivative series. Since $-f'(L) = 1/(v(v-1)) \sim L^{-2}$, it proves the first equivalent in (4.6).

For $F_1 = s^2 g(\ell)$, where $g = (v + \delta)/(v^2(v-1)) = v^{-2} + \mathcal{O}(v^{-3})$, the identical proof uses $1/\Gamma(-2-w)$. Its coefficient of w is -2 , and $g'(L) \sim -2L^{-3}$. Subtracting $s^2 g(L)$, again a polynomial, gives $k_m \sim 2m^{-3}g'(L) \sim -4m^{-3}L^{-3}$. Together with Theorem 4.1, this proves (1.3); the safe remainder divided by h_m is $\mathcal{O}(m^{-2}L^{-1})$, smaller than $1/(mL)$.

6.2 Finite generator and explicit formulas

Put $x = 1/L$ and $t = \log L + C$. Formally write $v = L + t + w$. The Lambert equation becomes

$$w = \log(1 + x(t + w)), \quad f = \frac{x}{1 + x(t + w)}. \quad (6.7)$$

For any requested finite order this determines polynomial coefficients in t recursively; the analytic implicit-function theorem in (x, xt) near $(0, 0)$ justifies the finite expansion and its polynomial-log remainder. Differentiation with respect to L acts as

$$D = -x^2 \partial_x + x \partial_t. \quad (6.8)$$

Expand $\sum_j c_j (-D)^j f$ only to the required finite order, substitute $t = T - \gamma$, and take the coefficient of x^{j+2} to obtain $Q_j(T)$. For Q_0 through Q_J , derivative orders through $J+1$ suffice, with all intermediate series truncated at degree $J+2$.

Theorem 6.2. *For every fixed integer $J \geq 0$, all three sequences satisfy (1.1). The universal*

polynomials through degree five are

$$\begin{aligned} Q_0 &= 1, \\ Q_1 &= 3 - 2T, \\ Q_2 &= 3T^2 - 11T + 6 - \frac{\pi^2}{2}, \\ Q_3 &= -4T^3 + 25T^2 + (2\pi^2 - 35)T + 10 - \frac{25\pi^2}{6} - 8\zeta(3), \end{aligned} \tag{6.9}$$

$$\begin{aligned} Q_4 &= 5T^4 - \frac{137}{3}T^3 + \left(\frac{225}{2} - 5\pi^2\right)T^2 \\ &\quad + \left(-85 + 40\zeta(3) + \frac{137\pi^2}{6}\right)T + 15 - \frac{75\pi^2}{4} - \frac{274\zeta(3)}{3} + \frac{\pi^4}{12}, \end{aligned} \tag{6.10}$$

$$\begin{aligned} Q_5 &= -6T^5 + \frac{147}{2}T^4 + \left(10\pi^2 - \frac{812}{3}\right)T^3 \\ &\quad + \left(\frac{735}{2} - \frac{147\pi^2}{2} - 120\zeta(3)\right)T^2 \\ &\quad + \left(-175 - \frac{\pi^4}{2} + 588\zeta(3) + \frac{406\pi^2}{3}\right)T \\ &\quad + 21 - \frac{245\pi^2}{4} - \frac{1624\zeta(3)}{3} - 144\zeta(5) + \frac{49\pi^4}{40} + 20\pi^2\zeta(3). \end{aligned} \tag{6.11}$$

Proof. The finite expansion (6.7), substituted into Lemma 6.1 at sufficiently high fixed derivative order, gives the stated terms and a relative remainder $\mathcal{O}_J((\log L)^{J+1}/L^{J+1})$. The coefficient error in Theorem 5.1 with $K = 1$ is $\mathcal{O}(m^{-3}L^{-2})$ and is absorbed in any such fixed logarithmic remainder. Finite algebra gives (6.9)–(6.11). The companion independently verifies the generated identities using a second reversion equation for $f = xF$, $1/F + x \log F = 1 + xt$. $\square$

The leading polynomial term is $(-1)^j(j+1)T^j$. Universality here is only across the two specified inverse hypergeometric families: their first-sector Lambert equations differ through C . Later exact sectors depend on additional local constants such as δ . At accessible indices a higher logarithmic truncation need not improve an approximation monotonically.

[write] Rederived, 6 October 2026. With its own code the write solved $w = \log(1 + x(t + w))$ to order x^8 , applied $\sum_j c_j(-D)^j f$ with c_j from (6.1), and substituted $t = T - \gamma$: the coefficients of x^0 and x^1 vanish and those of $x^2, \dots, x^7$ are exactly $Q_0, \dots, Q_5$ of (6.9)–(6.11) (SymPy); the leading coefficients are 1, $-2, 3, -4, 5, -6$, as stated. This checks the finite algebra with the same generator, not the analytic remainder, which rests on Lemma 6.1 and Theorem 5.1.

7 Smooth inverse charts and integer thresholds

Let $b = \log \mu$, and for large positive real y put

$$Y = \log(y/\kappa_0), \quad q = Y/b, \quad L_q = \log q, \quad T_q = \log L_q + C + \gamma. \tag{7.1}$$

Let $E_j(T)$ be the coefficient of x^j in the formal logarithm $\log(\sum_{k \geq 0} Q_k(T)x^k)$. Only finitely many Q_k are needed for each E_j ; in particular

$$E_1 = 3 - 2T, \quad E_2 = T^2 - 5T + \frac{3}{2} - \frac{\pi^2}{2}. \tag{7.2}$$

For fixed $J \geq 0$ define the smooth comparison

$$\mathcal{L}_J(m) = bm - 2 \log m - 2 \log \log m + \log \kappa_0 + \sum_{j=1}^J \frac{E_j(\log \log m + C + \gamma)}{(\log m)^j}. \quad (7.3)$$

Its derivative tends to $b > 0$. For large y it has a unique large real solution $M_J(y)$ of $\mathcal{L}_J(M_J) = \log y$; set $\xi_J = M_J - 2$. This is a smooth comparison root, not an integer-valued inverse.

The explicit chart is

$$x_J(y) = q - 2 + \frac{2}{b}(\log q + \log \log q) - \frac{1}{b} \sum_{j=1}^J \frac{E_j(T_q)}{L_q^j}. \quad (7.4)$$

In particular,

$$\begin{aligned} x_2(y) = q - 2 + \frac{2}{b}(\log q + \log \log q) + \frac{2T_q - 3}{bL_q} \\ + \frac{-T_q^2 + 5T_q - 3/2 + \pi^2/2}{bL_q^2}. \end{aligned} \quad (7.5)$$

Theorem 7.1. *For each fixed J , as $y \rightarrow \infty$,*

$$x_J(y) - \xi_J(y) = \mathcal{O}_J(\log q/q). \quad (7.6)$$

Define $N(y) = \min\{n : a_n \geq y\}$ with the sequence's own index origin. There are constants $K_J > 0$, y_J such that for $y \geq y_J$, setting

$$\varepsilon_J(y) = K_J \left\{ \frac{(\log \log q)^{J+1}}{(\log q)^{J+1}} + \frac{\log q}{q} \right\}, \quad (7.7)$$

one has

$$\lceil x_J(y) - \varepsilon_J(y) \rceil \leq N(y) \leq \lceil x_J(y) + \varepsilon_J(y) \rceil. \quad (7.8)$$

Proof. Taking logarithms in Theorem 6.2 gives

$$\log a_n - \mathcal{L}_J(n+2) = \mathcal{O}_J \left(\frac{(\log \log(n+2))^{J+1}}{(\log(n+2))^{J+1}} \right). \quad (7.9)$$

Substitute $m = x_J + 2$ into (7.3). Because $m = q + \mathcal{O}(\log q)$, replacing m by q in its slowly varying terms costs $\mathcal{O}_J(\log q/q)$; the terms displayed in (7.4) cancel all the retained corrections at q . The derivative $b + o(1)$ and the mean-value theorem imply (7.6).

The asymptotic formula also implies $a_{n+1}/a_n \rightarrow \mu > 1$, hence eventual strict increase. Enlarge y_J beyond all values in any finite initial exceptional portion. Uniformly for integers near ξ_J , the discrepancy in (7.9) transports through the increasing smooth comparison with derivative bounded below by $b/2$. Choosing K_J large enough and including (7.6) shows every integer below $x_J - \varepsilon_J$ has $a_n < y$, and the integer $\lceil x_J + \varepsilon_J \rceil$ has $a_n \geq y$. This proves (7.8). $\square$

The constants and starting point in this theorem are not explicitly evaluated or certified here. The two ceiling values may differ near an integer boundary, even though $\varepsilon_J(y) \rightarrow 0$. No uniform $o(1)$ approximation to the step function $N(y)$, and no unconditional formula $N(y) = \lceil x_J(y) \rceil$, is claimed. Exact finite-index identification requires exact count comparisons. The forward Lambert model uses W_{-1} ; the explicit inverse chart itself uses only ordinary real logarithms.

Remark 7.2 ([write] The inversions and the transseries volume, 6 October 2026). Compared, statement by statement, with `Analysis/Transseries/docs/series-and-transseries/Transseries_And_Inversion/transseries_and_inversion.tex`; symbols of the volume carry the subscript “vol”.

- (a) *Instance.* The threshold $N(y)$ is the integer staircase N_* of `p0:def:three-inverses` with $A_n = a_n$ and n_1 the onset of strict increase (eventual, as the proof above shows; the first 1000 terms of each sequence are strictly increasing, by the exact counts), for y above the finitely many earlier values; `p0:thm:staircase(1)` gives $N(y) = \lceil \nu_{\mathcal{A}}(y) \rceil$ for every admissible interpolation. The source does not use this.
- (b) *Theorem 7.1: an analogue as stated and proved; also a consequence of `p0:thm:staircase(1)` once an interpolation is supplied.* The source brackets $N(y)$ directly about the explicit chart x_J , which is not the inverse of an interpolation of the counts; the conclusion has the form of `p0:thm:staircase(2)`. A second route (added by the write): let $E(n) = \log a_n - \mathcal{L}_J(n+2)$, which is $\mathcal{O}_J((\log \log n)^{J+1}/(\log n)^{J+1})$ by (7.9), let $\tilde{E}$ be its piecewise-linear interpolation, and put $\log \mathcal{A}(x) = \mathcal{L}_J(x+2) + \tilde{E}(x)$. Since $\mathcal{L}'_J \rightarrow b > 0$ and $|\tilde{E}'| \leq 2 \max |E| \rightarrow 0$ on the unit intervals, $\mathcal{A}$ is eventually continuous, strictly increasing and equal to a_n at the integers. For $\nu = \nu_{\mathcal{A}}(y)$ one has $\mathcal{L}_J(\nu+2) - \mathcal{L}_J(M_J) = -\tilde{E}(\nu)$, so, once $\mathcal{L}'_J \geq b/2$ on the segment, $|\nu - \xi_J| \leq (2/b)|\tilde{E}(\nu)| \leq (2/b) \max(|E(\lfloor \nu \rfloor)|, |E(\lceil \nu \rceil)|)$, which is $\mathcal{O}_J((\log \log q)^{J+1}/(\log q)^{J+1})$ because $\nu \sim q$. With (7.6), $|\nu - x_J| \leq \varepsilon_J$ for a suitable K_J , and `p0:thm:staircase(1)` with the monotonicity of the ceiling gives (7.8).
- (c) *Instance on the real axis.* For real $s \in (0, S_0)$ the Lambert equation (4.2), $v - \log v = \Lambda$, is `p0:thm:lambert-core` with $a_{\text{vol}} = 1$, $b_{\text{vol}} = -1$: its branch rule gives two positive solutions exactly when $\Lambda > L_c = |b_{\text{vol}}|(1 - \log(|b_{\text{vol}}|/a_{\text{vol}})) = 1$, the large one $-W_{-1}(-e^{-\Lambda}) = -W_{-1}(-e^{-C}s)$, the source’s branch; the threshold 1 is the source’s condition $\text{Re } \Lambda > 1$ on D_s . The complex continuation by the contraction argument is the source’s own.
- (d) *Formal instance.* The large- Λ expansion of v behind the generator (6.7) is `plt:thm:lw-template` with $\xi_{\text{vol}} = \Lambda = L + C$, $Z_{\text{vol}} = v$ and data $(\rho_{\text{vol}}, \mu_{\text{vol}}, \beta_{\text{vol}}, B_{\text{vol}}) = (1, -1, 0, 0)$, the pure Lambert case $q_N = a_{\text{vol}}^{N+1} \text{LamP}_N$ with $a_{\text{vol}} = -1$; the source re-expands it in $x = 1/L$ and $t = \log L + C$ instead of in $1/(L + C)$.
- (e) *Formal instance after a change of variable.* With $q = 1 + E$, the joint equation (5.1) reads $\Lambda_{\text{vol}}E + h_{\text{vol}}(E) + f_{\text{vol}}(\eta, E) = 0$ with $\Lambda_{\text{vol}} = 1 - w$, $h_{\text{vol}}(u) = -w((1+u)\log(1+u) - u)$ and $f_{\text{vol}}(\eta, u) = \sum_{j \geq 2} b_j \eta^{j-1} (1+u)^j (1+w\delta_j - w\log(1+u))$, which is `p0:thm:core-reversion` over the ring $\mathbb{Q}[w, (1-w)^{-1}]$; its triangular recurrence gives each q_ℓ rational in w with poles only at $w = 1$, as the source says. The convergence, by the analytic implicit-function theorem, and the branch identification are the source’s own.
- (f) *Not shown to be an instance of `plt:thm:lw-template`.* The comparison equation $\mathcal{L}_J(m) = \log y$ behind the chart (7.4) has the core $bm - 2 \log m$, which is `p0:thm:lambert-core` with $a_{\text{vol}} = b$, $b_{\text{vol}} = -2$; the source inverts it by a direct logarithmic expansion about q instead. In the chart $Z_{\text{vol}} = m$, $\xi_{\text{vol}} = \log y$ the terms $-2 \log \log m$ and $E_j(\log \log m + C + \gamma)/(\log m)^j$ have t -order 0 ($t = 1/m$) and are not polynomial in $\log Z$, so hypothesis (H3) fails. The write does not claim that no other chart makes the full equation an instance; (7.6) is the source’s own analytic estimate.

No novelty is claimed for any of these inversions.

8 Reproducible computations and numerical use

8.1 Independent exact checks

The companion supplies bounded exact integer calculations and deliberately separate floating-point diagnostics. The equations from [1, Section 7] are

$$R(cR - 1)R'' = dt(R')^3, \quad R(0) = 0, \quad R'(0) = 1, \quad (c, d) = (16, 4) \text{ or } (27, 6). \quad (8.1)$$

For $m \geq 2$, let $R_{<m} = \sum_{j < m} r_j t^j$. The coefficient of the new r_m in the left-minus-right residual is $-m(m-1)r_m$, so

$$r_m = \frac{[t^{m-1}]\{R_{<m}(cR_{<m} - 1)R''_{<m} - dt(R'_{<m})^3\}}{m(m-1)}. \quad (8.2)$$

Integer divisibility is checked before division. A second exact route, independent of this differential recurrence, is Lagrange inversion:

$$r_m = \frac{1}{m} \sum_{k=0}^{m-1} (-1)^k \binom{m+k-1}{k} [r^{m-1}] \left(\frac{\Omega(r)}{r} - 1 \right)^k. \quad (8.3)$$

The package checks a bounded prefix against this formula and against frozen OEIS terms, tests the full generated differential residual, and checks rooting divisibility. It also verifies the Q identities and the sector residuals independently rather than merely printing the stored expressions. The command reference, exact output sizes and resource limits are in the included README.

8.2 Numerical coefficient extraction of the exact models

For $h(z) = F_0(z)$, differentiation of (4.2) yields

$$hh'' = -(h')^3, \quad h(0) = 1/v_0, \quad h'(0) = -1/(v_0 - 1), \quad v_0 - \log v_0 = C, \quad v_0 > 1. \quad (8.4)$$

It also gives

$$F_1 = h^2 - (\delta + 1)h^2 h' = h^2 - \frac{\delta + 1}{3}(h^3)'. \quad (8.5)$$

Coefficient comparison in (8.4) is triangular; maintaining the needed convolutions yields a quadratic-arithmetic-operation recurrence. An independent small-prefix check sets $h = h_0(1 - q)$ and inverts

$$z = h_0 \left\{ (v_0 - 1)q + \sum_{j \geq 2} \frac{q^j}{j(j-1)} \right\} \quad (8.6)$$

by Lagrange's formula. Agreement between these routes is a valuable implementation diagnostic; without interval arithmetic it is not a proof of their rounding error.

Family at $n = 1000$	F_0 model	$F_0 - F_1/2$ model
A324312	$-1.46974833746 \times 10^{-4}$	$+3.81810914855 \times 10^{-8}$
A324314	$-1.31150652412 \times 10^{-4}$	$+4.44844452278 \times 10^{-8}$

Table 2: Relative error (approximation/ a_n) $- 1$ from high-precision numerical extraction of the exact analytic models, using $m = n + 2$. These are diagnostics, not interval-certified error bounds.

The values in Table 2 were reproduced at 110 decimal digits, and the first 60 h_m were independently compared with (8.6). The independent route agreed to better than 10^{-90} relative discrepancy on that checked prefix. This is empirical agreement, not a certified accuracy statement. A277493 has the same relative errors as A324312. In contrast, the shifted leading equivalent alone at $n = 1000$ overestimates by about 165% and 185% in the two families. This slow logarithmic convergence explains the practical value of keeping the first analytic model whole.

The inverse chart also need not improve monotonically with truncation. At $y = A324312(1000)$, the numerical values of $x_J(y) - 1000$ for $J = 0, 1, 2$ are approximately -0.391352 , -0.00264436 , $+0.0300876$. These numbers test the explicit chart at a known level; they do not turn the asymptotic ceiling theorem into a finite certified bound.

8.3 Reproducibility and limits

The distribution contains this source and PDF, executable Python, fixed formula and OEIS fixtures with dated provenance, exact generated outputs, tests, build instructions and checksums. Validation uses explicit exceptions and remains active under Python’s optimized mode. Deliberate failures include invalid bounds, malformed or inconsistent fixtures, failed divisibility and attempted output overwrite. A clean build uses fixed PDF metadata and deterministic archive metadata, refuses to clobber output, and includes the final PDF and source in the complete ZIP. No source PDFs from the cited papers are redistributed.

All exact identities and integer checks can be repeated without network access. Numerical diagnostics require the listed arbitrary-precision package; symbolic verification requires the listed symbolic package. The README distinguishes bounded exact verification commands from the slower floating-point diagnostics; the full archive build regenerates both. The asymptotic statements rest on the proofs and cited analytic input, not on the finite checks.

[write] **Rechecked, 6 October 2026.** From the regenerated exact counts and its own model coefficients (the recurrence (8.4) with incremental convolutions at 60 digits, and k_m from (8.5)), the write reproduced all four entries of Table 2 to the printed digits, the leading overestimates $+1.6475$ and $+1.8500$ (the “about 165% and 185%”), and $x_J(y) - 1000 = -0.39135215$, -0.0026443594 , $+0.030087643$ ($J = 0, 1, 2$) at $y = A324312(1000)$. The $(1 + o(1))$ of (1.3) is slow: at $n = 1000$, $mL(a_n/(\kappa_0 \mu^m h_m) - 1) = 1.0177$ (general) and 0.9082 (quartic), not near 2, while $mL(-k_m/(2h_m)) = 1.0180$ and 0.9085 ; so the exact first sector accounts for almost all of the discrepancy, and its slow approach to 2 is a property of the exact models. Neither number tests the theorem, which is asymptotic. The exact file `outputs/exact.json` is not shipped in this repository; the README gives the command that regenerates it and its SHA-256.

9 Further research directions

1. **Effective error constants.** Quantify the common Δ -domain, implicit-function bounds and contour constants to obtain explicit finite-index forward inequalities. Interval-certified Lambert coefficient extraction would then support rigorous threshold localization.
2. **Large sector order.** Determine the growth of U_j and the optimal truncation scale. The fixed- K theorem alone does not answer whether the infinite coefficient-sector representation converges, or how additional distant singularities enter exponentially small corrections.
3. **Higher coefficient accuracy.** Expand the Cauchy kernel beyond its leading Hankel approximation while retaining exact sector separation. This can organize discretization corrections that

a finite inverse-log truncation otherwise obscures.

4. **Parameter-dependent models.** Where a rigorous global continuation theorem is available for weighted refinements, derive the corresponding local regular constants and test whether the same first-sector normalization persists. The predicted phase diagram in [4] is a motivation, not an assumption already established here.
5. **Certified inversion algorithms.** Combine the smooth chart with interval model bounds and a short exact recurrence window. Such a method could turn the asymptotic ceiling sandwich into a practical, certified integer inverse at specified large levels.

[write] **Added 6 October 2026, under Vladimir’s standing rule of 4 October 2026.** The source’s five directions stand. One question is added: the input on which every count statement for the general family rests is published with its proof only indicated (Remark 3.1).

6. **Continuation for the general family.** Write out the proof of Proposition 8.3 of [1]: radius $1/(4\pi)$, analyticity of the general inverse series R in a Δ -domain, and $R(t) - \frac{1}{16} \sim \frac{1}{4}(1-t/\rho)/\log(1-t/\rho)$ there. Its authors say that the proof “closely follows the proof of Proposition 8.2 given in [17, Sec. 8.3]” and give no details, and the source takes the proposition as input (Section 3). Sketch: the quartic argument of [2], Section 8.3, with Ω_g for Ω_q ; BMEP name nonnegativity of the coefficients of $t - R(t)$ as one key ingredient. Missing: the written argument, in particular that ρ is the only singularity on $|t| = \rho$ and the continuation beyond it. Until then, Theorems 4.1, 5.1, 6.2 and 7.1 are conditional for A324312 and A277493. (Added after the independent check of 7 October 2026: the same authors regard the proposition as proved. Version 3 of [4], Section 6.4 (p. 38), says of the logarithmic singular behaviour of its equation (85) that it “has been proven to be correct in [14] for $\omega = 0$ and $\omega = 1$ ”, where [14] is [1] and $\omega = 0$ is the general family ($Q(t, 0, 1) = 2G(t)$); the other weights are called predicted there. The check found no written proof elsewhere, so “conditional” here means: resting on a refereed proposition whose proof is only indicated, not on a conjecture.)

[write] **Independent check of the write, 7 October 2026.** An adversarial check made by the intake after the write (commit 05dcc17a8) re-derived the statements marked [write] with its own code, after fetching again A324312 (revision #18), A277493 (#53) and A324314 (#7), arXiv:1803.08265v1 and v2, the HAL deposit of [1], version 3 of [4] and [2]. *Counts.* Its own fixed-point reversion of Ω (exact rationals, $m \leq 42$) reproduces every printed term of the three entries; r_m to $m = 1002$ from the recurrence (8.2) agree with it for $m \leq 42$ and satisfy $\Omega(R) \equiv t$ modulo t^{1003} and three primes near 10^6 (a test independent of (8.1); a single perturbed coefficient fails it); the first 1000 terms of each sequence are strictly increasing, as Remark 7.2(a) says. *Remark 3.1.* Every quotation of Section 8 of [1], the Δ -domain, the radii, the two leading forms (and $\rho\alpha = \sqrt{3}/(54\pi)$, $1/(16\pi)$), BMEP’s coefficients of $\varepsilon \log \varepsilon$ and the citation note confirmed; the note is extended there, and question 6 gains the authors’ own later statement. *Remark 7.2.* Each instance re-derived against the volume’s statements: the staircase and its admissible interpolation (b), the branch rule and $L_c = 1$ (c), the data $(1, -1, 0, 0)$ and $q_N = (-1)^{N+1} \text{LamP}_N$ (d), $(1-w)E + h_{\text{vol}}(E) + f_{\text{vol}}(\eta, E) = 0$ with h_{vol} of order 2 in u , f_{vol} divisible by η , over $\mathbb{Q}[w, (1-w)^{-1}]$ (e), and the failure of (H3) in the chart $Z_{\text{vol}} = m$ (f). *Notes.* $Q_0, \dots, Q_5$ and E_1, E_2 recomputed by a different organization of the same transfer (the moments $\tau^j \mapsto j! c_j$ substituted in $f(L - \tau)$, no derivative operator); U_1, U_2, U_3 and $\delta_1, \dots, \delta_4$ of both families by substitution in the integrated equation; Table 1, H and C . The model coefficients h_{1002} , k_{1002} were computed as integrals over the two lips of the cut (no recurrence; validated at $m = 30$ against Taylor coefficients and a circle integral): all four entries of Table 2, 1.6474952 and 1.8500125, $mL(a_n/(\kappa_0 \mu^m h_m) - 1) = 1.0177405$ and 0.9081502, $mL(-k_m/(2h_m)) = 1.0180049$ and 0.9084582, and the three values of $x_J(y) - 1000$ as stated (the write’s four-digit values are rounded). *Provenance.* Archive facts, the 58 delivered label numbers (aux of a build of the delivered text) and the README’s file sizes and counts confirmed. No defect was found in the write.

A The fourth local sector

The finite generator (5.6) gives

$$U_3^g(v) = -\frac{40v^5 - 158v^4 + 112v^3 - 82v^2 + 107v - 46}{8192v^4(v-1)^5}, \quad (\text{A.1})$$

$$U_3^q(v) = -\frac{105v^5 - 434v^4 + 176v^3 - 156v^2 + 383v - 182}{19683v^4(v-1)^5}. \quad (\text{A.2})$$

The local constants needed through this order are

Family	δ_1	δ_2	δ_3	δ_4
General	0	$-5/2$	-3	$-193/60$
Quartic	0	-3	$-211/60$	$-3139/840$

These give $M_4 = s^4 U_3(v)$ in Theorem 5.1, whose remainder for $K = 4$ is $\mathcal{O}(m^{-6}(\log m)^{-5})$. The companion checks vanishing residual coefficients after substituting U_0, U_1, U_2, U_3 in the local equation.

B Source map and theorem dependencies

The exact inverse descriptions and root convention come from [1, Theorems 1.1–1.2 and Section 2]; its Section 7 supplies (8.1). The global continuation and leading singular equivalents used to identify the actual branch are [1, Propositions 8.2–8.3]. For the quartic antecedent, [2, Theorem 7.1, equation (53), Proposition 8.4] gives analytic logarithmic inversion, the local regular constant and global continuation. The general proposition’s omitted proof details are acknowledged in Section 3.

The hypergeometric and Lambert facts are the specific DLMF sections cited in the body. Elvey Price and Guttmann [3] discuss logarithmic test singularities, inverse-log corrections and reciprocal-Gamma derivative transfer, including the limiting integer-power issue. We therefore do not claim those analytic tools as new. The 2026 version of [4], Section 6.4, explicitly distinguishes predicted general-weight behavior from the specializations proved in the earlier paper. The frozen OEIS fixtures reproduce the displayed terms and their source URLs as retrieved on 3 October 2026; any future proposed OEIS update should recheck the then-current entry.

References

- [1] M. Bousquet-Mélou and A. Elvey Price, *The generating function of planar Eulerian orientations*, Journal of Combinatorial Theory, Series A **172** (2020), 105183. doi:10.1016/j.jcta.2019.105183. Inspected preprint: [arXiv:1803.08265v2](#), 10 October 2019.
- [2] M. Bousquet-Mélou and J. Courtiel, *Spanning forests in regular planar maps*, Journal of Combinatorial Theory, Series A **135** (2015), 1–59. Inspected preprint: [arXiv:1306.4536v1](#), 19 June 2013.
- [3] A. Elvey Price and A. J. Guttmann, *Counting planar Eulerian orientations*. Inspected preprint: [arXiv:1707.09120v3](#), 10 January 2018.

- [4] M. Bousquet-Mélou and A. Elvey Price, *Refined enumeration of planar Eulerian orientations*. [arXiv:2503.15046v3](#), 2 June 2026.
- [5] NIST Digital Library of Mathematical Functions, [equation 15.8.10](#) and [Section 4.13](#).
- [6] OEIS Foundation Inc., *The On-Line Encyclopedia of Integer Sequences*, [A277493](#), [A324312](#), [A324314](#). Retrieved versions inspected 3 October 2026.