

Rank Ultra-Log-Concavity of Preorder Gamma and Directed Support Polynomials at the Actual Degree

Degree four, independent role activities, small covers
and universal-sink clouds

Research manuscripts prepared for Vladimir Reshetnikov
(PDF metadata of sources 11, 12, 15, 16, 20, 22 and 25: “prepared with OpenAI”)
merged in ProveIt’s batch-73 intake

1 October 2026

Abstract

For a finite loopless directed relation D , for instance the off-diagonal relation of a finite preorder, let γ_k count the ordered pairs (S, T) of disjoint k -subsets of the ground set that admit a bijection $S \rightarrow T$ along arcs, each pair counted once however many matchings witness it, and possibly weighted by independent tail and head activities. For a preorder, $\Gamma_D = \sum_k \gamma_k t^k$ is the gamma polynomial of *preorder-root-polytopes*. That report asks (its Research question 96, `mr:q:gamma`) whether Γ_D is ultra-log-concave after normalization by its *actual* degree, and names degree three as the first open target. This report merges eleven manuscripts of ProveIt’s batch 73 that answer the question in three directions. Part I (sources 25, 16 and 23) proves it for every finite preorder of actual degree at most four with unit activities: degree three by a Gallai–Edmonds reduction and exact integer-population certificates, with the minimum order fifteen of a nonreal-rooted cubic; degree four by bounded cores, 76 four-attachment templates and 50 global binomial-square identities, with a smaller second route for the largest certificate. Part II (sources 20, 12, 11, 15 and 22) treats independent tail and head activities: real stability of the signed monomer polynomial under every two by two role cover, coefficientwise Rayleigh inequalities for a three-vertex shore, rank-ULC under every physical vertex cover of size three and on bipartite physical graphs, a pendant-head stability criterion, and rank-ULC of every independently weighted preorder of actual degree at most three. Part III (sources 14, 10, 34 and 06) treats an arbitrary directed core with a cloud of universal sinks: a sharp last-coefficient bound for every core size, rank-ULC at every actual degree for four-vertex cores, and for five-vertex cores at actual degrees five and four. The degree-three and degree-four theorems are computer-assisted; the structural theorems have ordinary proofs. Nothing here is formalized in Lean or Rocq, and nothing is refereed.

Contents

1	Guide to this report	4
1.1	Sources, status and plan	4
1.2	What the neighbouring reports already prove	4
1.3	The common object	5
1.4	Notions and notation	6
 I	 Unit activities: finite preorders through actual degree four	 8

2	Degree at most three (source 25)	9
2.1	Statement and mathematical setting	9
2.2	The universal first inequality	10
2.3	A canonical bounded core from Gallai and Edmonds	10
2.4	Finite enumeration and exact positivity methods	12
2.5	No attachments and one attachment	14
2.6	Two attachments	15
2.7	Three attachments and seventeen templates	16
2.8	Completion of the main theorem	18
2.9	Real roots and an explicit obstruction	18
2.10	An infinite family with three negative roots	19
2.11	Reproducibility and scope	20
2.12	Explicit discriminant certificates for the ordinal family	21
3	Degree four (source 16, the base)	22
3.1	The support polynomial and the questions	23
3.2	An exterior-only Lorentzian construction	24
3.3	The two by two role cover	24
3.4	The unbalanced one-source three-sink family	25
3.5	A finite normal form for bounded matching rank	26
3.6	Exact support kernels and coverage	27
3.7	The bounded branches and the degree four theorem	27
3.8	Integer population certificates	29
3.9	Verification boundary and remaining research	31
4	A second route for the last gap of template 26 (source 23)	32
4.1	Supports and the precise family	32
4.2	The support decomposition	33
4.3	Reducing deletion mixtures to pairs	34
4.4	The ten exact quartic identities	34
4.5	The top ratio with the internal edge present	35
4.6	Compatibility and the final square	36
II	Independent role activities and small covers	37
5	Two by two role covers (source 20)	38
5.1	Support polynomials and normalization	38
5.2	Real-rootedness for a complete balanced core	39
5.3	Proof and verification boundary	45
5.4	Appendix of source 20: Rayleigh identities for two disjoint core edges	46
6	Coefficientwise Rayleigh inequalities for a three-vertex shore (source 12)	47
6.1	Supports and the main inequality	47

6.2	Complete coefficient localization	48
6.3	Full Rayleigh differences and negative correlation	50
6.4	A scalar cubic consequence	51
6.5	An exact obstruction at four cores	51
6.6	Proof status and reproducibility	52
7	A physical vertex cover of size three (source 11)	52
7.1	Statement and normalization	53
7.2	The exterior boundary input	54
7.3	Restoring every internal core arc	54
7.4	Bipartite only strictness results	55
7.5	Proof status and independent replay	57
8	Bipartite physical graphs and pendant heads (source 15)	57
8.1	Supports and the normalization issue	57
8.2	A polynomial for bipartite physical graphs	58
8.3	Pendant heads attached to a stable transversal seed	60
8.4	An explicit stable seed in arbitrary rank	61
8.5	Reproducibility and the remaining question	62
9	Independent role activities through actual degree three (source 22)	63
9.1	Definitions and the main theorem	63
9.2	Degree, zero activities, and the universal first gap	64
9.3	An ordinary all-rank articulation argument	65
9.4	Three active tails: a rank-three Rayleigh proof	66
9.5	Role covers and a structural corollary	68
9.6	The finite theorem through seven vertices	69
9.7	Half-plane-property side matroids and the final cases	72
9.8	Two attachments and arbitrary exterior populations	75
9.9	The Gallai–Edmonds reduction and the global proof	77
9.10	A cubic with nonreal zeros	79
9.11	Verification and dependencies	79
9.12	Further questions and mathematical boundaries	80
	Appendix	81
A	Provenance and merge record	81
A.1	The archives	81
A.2	What each source contributed	82
A.3	Where the merge had to choose	83
A.4	Shipped files	84

1 Guide to this report

1.1 Sources, status and plan

This report merges research manuscripts delivered to ProveIt’s incoming-report intake on 1 October 2026 (batch 73, arrival commit `f8c3a392a`, placement commit `8f5c53106`). Sources are numbered by their batch-73 manuscript numbers; a package delivered in several archives takes the lowest number: source 16 is the four archives 16–19 (`preorder-degree-four`, parts 4 to 1), and source 06 is the two archives 06 and 07 (`universal-sink-five-core-boundary`, parts 2 and 1; its manuscript is in archive 07). Eighteen archives are behind the eleven printed sources. Two further archives are earlier versions of printed sources and are not sources: 31 (vertex activities, superseded by source 22) and 13 (a weaker four-core inequality, superseded by source 10). Appendix A records every archive.

The report has three Parts.

- Part I, unit activities on finite preorders through actual degree four: source 25 (degree three), source 16 (degree four, the base of the merge) and source 23 (a second route for the largest degree-four certificate).
- Part II, independent role activities and small covers: sources 20, 12, 11, 15 and 22.
- Part III, universal-sink clouds over an arbitrary directed core: sources 14, 10, 34 and 06.

Every source is printed with its statements, proofs, examples, remarks, limitations, verification notes and research questions; each source’s section opens with its abstract as delivered. A result proved by several sources is printed once, and the other proofs are kept as marked second routes or, where the proof is the same, replaced by a pointer recorded at the place. A result that *preorder-root-polytopes* or *matching-rank-normalization* already proves is cited by its label there and is not presented as new. Editorial additions are marked “[*Merge note*: ...]”.

Status. The sources are AI-assisted research manuscripts and notes: the title pages of sources 11, 12, 15, 16, 20, 22 and 25 read “Research manuscript prepared for Vladimir Reshetnikov” and their PDF metadata “Research manuscript prepared with OpenAI”; sources 06, 10, 14 and 34 read “Research note”; source 23 has no author line. Several main theorems are computer-assisted: their proofs include finite exact certificate computations shipped with this report (Section A.4). The report is unrefereed, and nothing in it is formalized in a proof assistant. No source claims literature-wide priority.

1.2 What the neighbouring reports already prove

Two reports of the collection precede this one. Their results are cited by label and are not reprinted. Line numbers are not used: the labels are stable.

- *preorder-root-polytopes* (in `enumerative-combinatorics/`). Theorem `gam:thm:main` expresses the h -polynomial of a finite preorder through the support numbers of this report, and Theorem `gam:thm:deficiency` says $\deg \Gamma_\tau = \nu(\text{Comp}(\tau))$. Theorem `lor:thm:weightedULC` proves two-shore ultra-log-concavity normalized by the smaller shore, and Corollary `lor:cor:gammaULC` all height-two gamma sequences. Theorems `ade:mat:thm:stable-bridge` and `ade:mat:thm:vertex-stable` treat stability of signed matching supports of bipartite graphs. In Part XII, Theorem `mr:thm:first` (with Lemma `mr:lem:MS`) is the universal first inequality for directed relations with non-negative activities, with a merge note crediting its vanishing-activity form; Theorem `mr:thm:rankthree` and Corollary `lc:cor:rankthree` settle bipartite matching rank three; Corollary `mr:cor:preorderfirst` gives the first inequality and hence every degree at most

two for preorder gamma polynomials; and Questions `mr:q:gamma` (Research question 96) and `mr:q:certificates` are the questions this report answers or bears on.

- *matching-rank-normalization* (beside this report). Corollary `mrn:w:rf:cor:rankfive`: every weighted bipartite graph of matching number at most five is ultra-log-concave of order its matching number. Theorem `mrn:w:sb:thm:main`: a bipartite graph with a vertex cover of two vertices on each shore has a real-stable support polynomial. Equation `mrn:s:tv:eq:rayleigh`: the scalar rank-three Rayleigh inequality (Wagner). Its Part V prints batch-73 manuscript 26 (source 77 there): Theorem `mrn:v:rf4:thm:main`, normalization by the actual degree through rank four, and Theorem `mrn:v:rf4:thm:mixed`, real roots under a two by two cover.

Sources 16 and 25 cite *preorder-root-polytopes* at its commit `d5e863bba`; the labels they cite are unchanged at the placement commit.

1.3 The common object

The following definitions are common to all eleven sources; each source restates them in its own words, and those restatements are kept in their sections because they fix each source's precise scope. Let D be a finite loopless directed relation on a set V . For a preorder, D is its off-diagonal relation: an arc $i \rightarrow j$ means $i \neq j$ and i precedes j , and equivalent elements give opposite arcs.

Definition 1.1 (Feasible supports). A feasible support of size k is an ordered pair (S, T) of disjoint subsets of V , each of cardinality k , for which there exists a bijection $\pi : S \rightarrow T$ with $s \rightarrow \pi(s)$ for every $s \in S$. For nonnegative tail activities u_i and head activities v_i put

$$\gamma_k(D) = \sum_{(S,T) \text{ feasible, } |S|=k} u^S v^T, \quad u^S v^T = \prod_{i \in S} u_i \prod_{j \in T} v_j, \quad \Gamma_D(t) = \sum_{k \geq 0} \gamma_k(D) t^k.$$

The empty support contributes $\gamma_0 = 1$. A support contributes once, regardless of the number of witnessing bijections. With unit activities γ_k is the number of feasible supports of size k .

The same-ground-set disjointness is essential. Passing to independent tail and head copies without subsequently excluding their simultaneous use changes this polynomial. Counting individual matchings also changes it. A vertex may have different activities in its two roles; with zero activities the polynomial degree can decrease.

Definition 1.2 (Rank-ULC). A polynomial $g(t) = \sum_{k=0}^d g_k t^k$ is *rank-ultra-log-concave*, or rank-ULC, if its coefficients are nonnegative with no internal zeros and

$$\left(\frac{g_k}{\binom{d}{k}} \right)^2 \geq \frac{g_{k-1}}{\binom{d}{k-1}} \frac{g_{k+1}}{\binom{d}{k+1}} \quad (1 \leq k < d),$$

where d is the actual degree.

Normalization by the ground-set size, by a cover size, or by a presumed degree after some activities vanish is a different assertion, and Section 1.4 lists where a source uses one.

Lemma 1.3 (Degree; sources 16, 20, 25). *For unit activities, the actual degree of Γ_D equals the matching number of the underlying undirected graph. Its coefficients have no internal zeros.*

Proof. A witnessing bijection yields an undirected matching. Conversely orient each edge of an undirected matching in a direction present in D . Its disjoint endpoint sets form a feasible support. Submatchings establish positivity at every smaller degree. $\square$

[Merge note: for preorders this is *preorder-root-polytopes*, Theorem `gam:thm:deficiency`. Sources 16, 20 and 25 state and prove it as their Lemma “degree” with the proof above (source 25 for preorders, in its own words); source 22 adds a filtering argument for zero role activities, printed in Part II (Lemma 9.4). The statement is not new.]

Lemma 1.4 (Motzkin–Straus bound; source 25). *If a simple graph H has clique number $d \geq 1$ and nonnegative vertex weights w_v , then*

$$\sum_{uv \in E(H)} w_u w_v \leq \frac{d-1}{2d} \left(\sum_v w_v \right)^2.$$

Proof. Normalize the total weight to one, the zero-total case being immediate. If nonadjacent vertices u, v both have positive weight, keep their sum fixed and move all of it to the one with the larger weighted neighbor sum. The objective does not decrease, since it is affine in this division, and its positive support shrinks. Repetition leaves a clique of size $q \leq d$. On that clique the objective is $(1 - \sum_v w_v^2)/2 \leq (1 - 1/q)/2 \leq (d-1)/(2d)$. This is the classical compression argument of Motzkin and Straus [11]. $\square$

Theorem 1.5 (Universal first inequality). *Let D be a finite loopless directed relation with nonnegative independent tail and head activities, and let $d \geq 2$ be the actual degree of Γ_D . Then*

$$\gamma_1^2 \geq \frac{2d}{d-1} \gamma_2. \quad (1.1)$$

Transitivity is not needed.

Proof. Discard every arc of zero weight $u_i v_j$; this changes no coefficient. Make a graph whose vertices are the remaining directed arcs and whose edges join arcs with four distinct physical endpoints. A clique is a positive-weight matching, and every positive-weight support has such a witness, so the clique number is exactly d . Give arc $i \rightarrow j$ weight $u_i v_j$; the total weight is γ_1 . Every feasible support of size two is witnessed by at least one edge of this graph, and every witness has the prescribed support weight; a fixed pair of arcs determines only one ordered support. The weighted edge sum is consequently at least γ_2 , and Lemma 1.4 bounds it by $(d-1)\gamma_1^2/(2d)$. $\square$

[Merge note: this is *preorder-root-polytopes*, Theorem `mr:thm:first`, with the same proof; the form at the surviving degree for vanishing activities is credited in the merge note after that theorem. It is re-proved, by the same route and without novelty claim, as source 16’s Theorem “Universal first inequality”, source 25’s Proposition “first” (unit activities, with the lemma above), source 22’s Proposition “first”, source 11’s Lemma “first” and the first comparisons of sources 10, 34 and 06; sources 10, 11, 34 and 06 prove Lemma 1.4 by the same mass-shifting argument, and source 12 gives its triangle-free case for actual degree two. Those texts are replaced by pointers to this theorem. Source 25 adds, and we keep, that counting matchings instead of supports would change the second cubic gap, and that the comparison in this proof cannot be used backwards to prove it.]

In degrees two, three and four the inequality reads $\gamma_1^2 \geq 4\gamma_2$, $\gamma_1^2 \geq 3\gamma_2$ and $3\gamma_1^2 \geq 8\gamma_2$. In particular every Γ_D of actual degree at most two is rank-ULC and real-rooted; for preorders this is *preorder-root-polytopes*, Corollary `mr:cor:preorderfirst`.

1.4 Notions and notation

The eleven sources share one object but not one vocabulary. The following conventions hold throughout; every source section uses its own letters inside its own scope, and the table below says where a letter means different things.

Notion	Meaning in this report	Not to be confused with
Γ_D, γ_k	ordered <i>disjoint</i> endpoint supports on the <i>same</i> ground set, each counted once (Definition 1.1)	the matching polynomial of D , and the support polynomial of the two-copy bipartite enlargement; the latter is the h -type setting of <i>preorder-root-polytopes</i> Parts XI–XIII and of <i>matching-rank-normalization</i>
Preorder gamma	for a preorder τ , $\Gamma_\tau = \Gamma_D$ with D its off-diagonal relation (<i>preorder-root-polytopes</i> , <code>gam:thm:main</code>)	the h -polynomial $h_\tau = \sum_k \gamma_k t^k (1+t)^{n-2k}$
Unit activities	all $u_i = v_i = 1$ (Part I)	“unweighted” in source 16 means unit activities; its “weighted quotient” means positive integer class sizes
Vertex activities	one activity $x_i = u_i = v_i$ per vertex (archive 31, recorded as a corollary in Part II; source 23)	independent role activities
Independent role activities	u_i (tail) and v_i (head), independent and nonnegative (sources 06, 10, 11, 12, 14, 15, 20, 22, 34)	a zero tail activity does not delete a head role (sources 06 and 22)
Actual degree	$\deg \Gamma_D$ after zero activities are imposed	the ground-set size, a cover size, or a degree presumed before activities vanish
Order- r ULC	log-concavity of $\gamma_k / \binom{r}{k}$ for a fixed $r \geq d$ (source 16’s exterior theorem, source 15’s shore order)	rank-ULC; the two agree only when $r = d$
Role cover	sets P (tails), Q (heads) with every arc $i \rightarrow j$ having $i \in P$ or $j \in Q$; P and Q may overlap (sources 11, 15, 20, 22)	a physical vertex cover of the underlying graph (source 11), and the bipartite cover of <i>matching-rank-normalization</i> with two vertices on each shore (the disjoint bipartite case)
Bipartite physical graph	the underlying undirected graph is bipartite; arcs may run both ways across (source 15)	a bipartite <i>role</i> graph (tail and head copies), which every relation has
Template, attachment	Gallai–Edmonds normal form: $a = A $ attachments, exterior types with integer populations (sources 16, 23, 25)	cores with exterior types in general; a universal-sink cloud is the case of one exterior type
Rayleigh	coefficientwise (source 12, in formal activities); scalar on the nonnegative orthant (Wagner’s rank-three theorem, sources 16, 22, 23); all-real (the stability criterion, sources 20 and 22)	these are three different statements and are never interchanged
Real stable	nonvanishing when all variables lie in $\mathbb{H} = \{\operatorname{Im} z > 0\}$	the half-plane property (positive real parts), equivalent for homogeneous polynomials up to rotation (source 22)

Letters with several meanings.

- P, Q : tail and head sets of a role cover in sources 15, 16, 20 and 22, and in source 11’s Corollary; Q is also a polynomial in sources 16, 22 and 23 ($Q(t) = 1 + Lt + Bt^2 + Tt^3$, $Q = \Gamma_{H-h}$), and P the polynomial $\Gamma_H - Q$ in source 22. The report *preorder-root-polytopes* is never called P in the text.

- A, B : in Part III (sources 06, 10, 34) the moment parameters $A = \sqrt{p_2}$, $B = p_1 - A$ of the sink activities; in source 22 A_i , B_{ij} , C_0 are three-core head sums; in sources 16 and 23 A, B, C, D and T are coefficients of Q and R ; in source 16's role-cover material B_P, L_P are basis and element sums.
- K : the core $V \setminus I$ of the Gallai–Edmonds normal form in sources 16, 22 and 25; a number of two-element supports in source 23; $K(r, m)$, the sharp last-gap constant of source 14; the kernel $p_0q_0 + p_1q_1$ in source 20's appendix.
- D : the directed relation, and also the second cubic gap $\gamma_2^2 - 3\gamma_1\gamma_3$ in source 25, the set of deficient vertices in the Gallai–Edmonds decomposition (sources 16, 22, 25), and a coefficient of R in source 16.
- e_j and E_j (Part III): elementary symmetric functions of the core tail activities and of the sink head activities. e_j is the elementary symmetric polynomial itself.
- ρ : the actual degree in the notation of sources 06, 10 and 34; d in the others.

No symbol of a source has been renamed. Where a source's normalization differs from the actual degree (source 16's Theorem on the exterior construction and source 15's shore order), the source says so at the place, and that sentence is kept.

Labels and file names. Labels of this report carry the prefix `pgr:`, with `pgr:u:`, `pgr:r:` and `pgr:s:` for Parts I, II and III and a two-letter infix per source (README). Each source refers to files of its delivered package by their delivered paths; the report README maps them to the shipped files in `code/` and `data/`.

Part I

Unit activities: finite preorders through actual degree four

This Part answers Research question `mr:q:gamma` of *preorder-root-polytopes* for unit activities through actual degree four. Section 2 (source 25) proves rank-ULC for every finite preorder of actual degree at most three, determines the least order of a preorder with a nonreal-rooted cubic gamma polynomial, and gives a general bounded-core normal form. Section 3 (source 16, the base of this report) proves rank-ULC for every finite preorder of actual degree four. Its two by two role-cover material is source 20's article and is printed once, in Part II. Section 4 (source 23) replaces the largest certificate of source 16 by a smaller certificate-assisted argument. Together:

Theorem 1.6 (Sources 25 and 16). *Every finite preorder whose unweighted gamma polynomial has actual degree at most four is rank-ULC at that degree.*

The proof is Theorem 2.3 for degrees at most three and Theorem 3.9 for degree four (which quotes the former). Both are computer-assisted. Neither extends to activities other than one: Part II proves the independently weighted statement through degree three, and the weighted statement in degree four, and every statement in degree five or more, are open. Real-rootedness fails already in degree three (Section 2.9).

2 Degree at most three (source 25)

Source 25 is *Rank ultra log concavity for preorder gamma polynomials of degree at most three* (14 pages as delivered; archive `preorder-gamma-degree-three-package`). Its code, certificates and receipts are shipped with the prefix `25-degree-three-`.

Abstract of source 25, as delivered. We prove that the gamma polynomial of every finite preorder of actual degree at most three is ultra-log-concave after normalization by the binomial coefficients of that degree. The essential new case is the second cubic Newton inequality. A Gallai–Edmonds decomposition reduces arbitrary ground-set size to four cases: components of order at most seven; cores of order at most seven with one pendant population; cores of order at most five with two attachment vertices; and seventeen three-vertex-cover templates. Exact finite enumeration and rational polynomial certificates settle these cases for all nonnegative integer populations. The certificate package includes independent coverage and support-count checks, rather than only numerical tests of small instances. The distinction from real-rootedness is essential: an explicit fifteen-element height-two poset has gamma polynomial $1+24z+162z^2+208z^3$ with negative discriminant. A separate exhaustive use of the same structural reduction proves that fifteen is the minimum order of such a nonreal-rooted cubic: every degree-three preorder on at most fourteen vertices is real-rooted, and at order fifteen the obstruction is unique up to isomorphism and order reversal. We also record a real-rooted infinite ordinal-sum family and describe the scope and reproducibility of the computation.

2.1 Statement and mathematical setting

A finite preorder τ is a reflexive transitive relation on a finite set V . Its off-diagonal directed relation has an arc $u \rightarrow v$ precisely when $u \neq v$ and $u \tau v$. Antisymmetry is not assumed: distinct equivalent vertices give arcs in both directions. Write $G = \text{Comp}(\tau)$ for the simple undirected graph with edge uv whenever at least one of $u \rightarrow v$ and $v \rightarrow u$ occurs.

Definition 2.1 (Disjoint directed supports). A support of size k is an ordered pair (S, T) of disjoint subsets of V , with $|S| = |T| = k$, for which a bijection $\pi : S \rightarrow T$ satisfies $s \rightarrow \pi(s)$ for every $s \in S$. Let $\gamma_k(\tau)$ be the number of these ordered pairs and put

$$\Gamma_\tau(z) = \sum_{k \geq 0} \gamma_k(\tau) z^k, \quad \gamma_0(\tau) = 1.$$

A support is counted once even when several bijections realize it.

This is the gamma polynomial identified in Part IV of the supplied root-polytope report [1], in particular its theorem labeled `gam:thm:main`. The present paper proves its statements directly for this explicitly defined polynomial. Thus the combinatorial result is independent of how that identification is implemented in the larger report. The convention differs both from counting ordinary matchings and from counting arbitrary row–column supports in two copies of the ground set: here the two endpoint sets must be disjoint in the original set.

[Merge note: source 25’s Lemma “degree” (for a preorder τ : $d = \deg \Gamma_\tau = \nu(G)$, and all γ_k with $0 \leq k \leq d$ are positive) is Lemma 1.3, printed once in Section 1.3 with the same proof; for preorders it is *preorder-root-polytopes*, Theorem `gam:thm:deficiency`.]

Definition 2.2 (Normalization by actual degree). A polynomial $g(z) = \sum_{k=0}^d g_k z^k$ with nonnegative coefficients and no internal zeros is *rank-ultra-log-concave*, or rank-ULC, if

$$\left(\frac{g_k}{\binom{d}{k}} \right)^2 \geq \frac{g_{k-1}}{\binom{d}{k-1}} \frac{g_{k+1}}{\binom{d}{k+1}} \quad (1 \leq k < d). \quad (2.1)$$

The parameter is the actual polynomial degree, not $|V|$ or $\lfloor |V|/2 \rfloor$.

Theorem 2.3 (Main theorem). *Every finite preorder τ with $\deg \Gamma_\tau \leq 3$ has a rank-ULC gamma polynomial. In actual degree three this says*

$$\gamma_1^2 \geq 3\gamma_2, \quad \gamma_2^2 \geq 3\gamma_1\gamma_3. \quad (2.2)$$

In actual degree two the required inequality is the stronger $\gamma_1^2 \geq 4\gamma_2$.

The proof has a classical structural input, elementary support arguments, and finite exact certificates. It is a computational proof with independently checked generators and identities; it is not a proof-assistant formalization. In particular, the conclusion quantifies over arbitrarily large finite preorders. No upper bound on the population sizes is used to prove Theorem 2.3.

The original report [1, Part XII, question `mr:q:gamma`] asks whether every preorder gamma polynomial is log-concave or rank-ULC and identifies degree three beyond the bipartite case as the first unresolved target. The theorem closes that degree-three target. It does not settle degree four or higher, a weighted version, or real-rootedness. A global literature-priority claim is not made.

[Merge note: this target is now closed further. Degree four with unit activities is source 16’s Theorem 3.9. The weighted version in degree three, with independent tail and head activities, is source 22’s Theorem 9.3; its case $u_i = v_i = 1$ is Theorem 2.3, and the proof printed below, by integer-population certificates, and source 22’s proof, by a ledger of role certificates, are independent routes. Real-rootedness fails (Section 2.9).]

2.2 The universal first inequality

The first inequality is already established in the report [1, `mr:thm:first`]. Source 25 includes its short proof to make the normalization and logical dependencies explicit; that proof is printed once, in Section 1.3.

[Merge note: source 25’s Lemma “ms” (the Motzkin–Straus bound, with its compression proof) and Proposition “first” (for any finite off-diagonal directed relation with $d = \nu(G) \geq 2$, $\gamma_1^2 \geq \frac{2d}{d-1}\gamma_2$; transitivity is not needed) are Lemma 1.4 and the unit-weight case of Theorem 1.5, printed once in Section 1.3. Source 25’s proof of the proposition is the proof given there: different supports have disjoint sets of realizing pairs of arcs, so γ_2 is at most the number of edges of the arc-disjointness graph.]

It follows immediately that all degree-two cases are rank-ULC and real-rooted, and that the first degree-three inequality always holds. Degrees zero and one have no internal inequalities. The remaining problem is precisely

$$D = \gamma_2^2 - 3\gamma_1\gamma_3 \geq 0 \quad \text{when } d = 3. \quad (2.3)$$

Counting matchings instead of supports would change D , and the comparison in the proof of Proposition 1.5 cannot be used backwards to prove this second inequality.

2.3 A canonical bounded core from Gallai and Edmonds

For any finite simple graph G , let D_G be the vertices omitted by at least one maximum matching, let $A = N(D_G) \setminus D_G$, and let $C = V(G) \setminus (A \cup D_G)$. The Gallai–Edmonds theorem states that the components D_i of $G[D_G]$ are factor-critical, $G[C]$ has a perfect matching, and every maximum matching matches the vertices of A into distinct D_i , with internal near-perfect matchings in all D_i and a perfect matching in C . We use the theorem in the form proved by West [17, Theorem 5]. In particular,

$$\nu(G) = |A| + \frac{|C|}{2} + \sum_i \frac{|D_i| - 1}{2}. \quad (2.4)$$

Assume from now on that $\nu(G) = 3$. Let I be the union of the singleton components of $G[D_G]$ and put $K = V(G) \setminus I$. We call K the core. The set I is independent and $N(I) \subseteq A$. Write $a = |A|$, $|C| = 2c$, and $|D_i| = 2d_i + 1$ for each nonsingleton component, where $d_i \geq 1$. Equation (2.4) gives

$$a + c + \sum_i d_i = 3, \quad |K| = a + 2c + \sum_i (2d_i + 1) \leq 9 - 2a. \quad (2.5)$$

Here $2d_i + 1 \leq 3d_i$ and $2c \leq 3c$. Hence $a \in \{0, 1, 2, 3\}$.

2.3.1 Why the preorder boundary has no equivalence ambiguity

An arbitrary vertex cover can cut through a preorder equivalence class. That would permit bidirectional exterior incidences and invalidate a fixed-sign exterior classification. The canonical choice above avoids this issue.

Lemma 2.4. *Every vertex of I is a singleton preorder equivalence class. At each attachment vertex $v \in A$, all arcs incident with nonisolated vertices of I have the same direction.*

Proof. Equivalent vertices u, v have identical relations to every third vertex by transitivity. In G they are adjacent true twins, so their transposition is an automorphism. The sets D_G, A, C are automorphism-invariant by their definitions. If an exterior vertex $u \in I$ were equivalent to $v \neq u$, then v would also lie in D_G and be adjacent to u , contradicting that u is a singleton component of $G[D_G]$.

There are therefore no bidirectional arcs crossing from I to K . If distinct $x, y \in I$ satisfied $x \rightarrow v \rightarrow y$, transitivity would force $x \rightarrow y$, contradicting independence. This excludes opposite directions at each fixed attachment. $\square$

After discarding isolated exterior vertices, each exterior vertex is determined by a nonempty neighborhood $T \subseteq A$, together with the direction fixed at each member of T . There are at most $2^a - 1$ possible types. Let $n_T \in \mathbb{Z}_{\geq 0}$ be their populations.

Lemma 2.5 (One-vertex compatibility is sufficient). *Fix a core preorder and a direction for each attachment. Keep precisely the nonempty neighborhood types for which adjoining one vertex gives a preorder. Then arbitrary nonnegative populations of all these types give a preorder.*

Proof. A transitivity failure has a witness $u \rightarrow v \rightarrow w$ with $u \not\rightarrow w$. A witness involving at most one exterior vertex is already excluded by the core and single-type checks. A witness involving two distinct exterior vertices would have an attachment as its middle vertex and require both exterior directions there, which is forbidden. There are no exterior-to-exterior arcs, so these are all possibilities. $\square$

Thus enlarging the family by allowing all nonnegative integer populations is safe. The canonical decomposition can impose additional neighborhood-surplus conditions, but there is no need to impose them for a positivity proof on the larger family.

a	Structural consequence	Verification needed
0	Every nontrivial component has at most 7 vertices	Finite real-rootedness lemma
1	$ K \leq 7$, one pendant-twin population	2050 pairs of core polynomials
2	$ K \leq 5$, at most three exterior types	1084 templates, 448 polynomial families
3	$K = A$, $ K = 3$, at most seven types	17 templates

2.3.2 A reusable normal form at every fixed degree

Theorem 2.6 (General bounded-core normal form). *Let τ be a finite preorder with comparability matching number r . The same canonical construction gives an attachment set A , $a = |A| \leq r$, and an independent exterior I such that*

$$|K| \leq 3r - 2a, \quad \nu(G[K \setminus A]) = r - a.$$

Every exterior point is a singleton equivalence class, every attachment has one fixed exterior direction, and there are at most $2^a - 1$ nonisolated exterior types. The gamma coefficients have the finite expansion (2.7), of total population degree at most a .

Conversely, fix a core of size at most $3r - 2a$, a marked set A of size a , fixed directions, and the compatible exterior types. If $\nu(G[K \setminus A]) \leq r - a$, every population blow-up has matching number at most r . Thus all degree- r preorders are represented by finitely many bounded-core polynomial families with unbounded nonnegative integer populations.

Proof. Replace 3 by r in (2.4). Writing $r = a + c + \sum d_i$ gives the same bound $|K| \leq a + 3(c + \sum d_i) = 3r - 2a$. The graph on $K \setminus A$ is the disjoint union of $G[C]$ and the nonsingleton factor-critical components, with matching number $c + \sum d_i = r - a$. Lemmas 2.4 and 2.5 do not depend on r . In any matching, each selected exterior vertex requires its own attachment, proving the quota bound and (2.7). For the converse, at most a matching edges touch A and every other edge lies in $K \setminus A$, so the matching size is at most $a + \nu(G[K \setminus A]) \leq r$. For fixed r , there are only finitely many core relations, marked sets, and direction choices. $\square$

The restriction on $\nu(G[K \setminus A])$ matters in the converse: the core-size bound alone does not imply matching rank at most r . The normal form reduces any fixed-rank coefficient inequality to finitely many population-polynomial families; it does not assert that arbitrary integer-polynomial positivity is decidable or that higher-degree ULC holds. At specializations where the actual degree drops, the normalization must be changed accordingly.

2.4 Finite enumeration and exact positivity methods

This section specifies what the computer checks and why the checks have the needed quantifiers. The data and source programs in the accompanying archive are part of the certificate.

2.4.1 Covering all finite preorder cores

Every finite preorder is uniquely a poset of equivalence classes with positive block sizes. Choose a linear extension of the quotient poset and label its q elements $0, \dots, q-1$ in that order. Every strict comparison then points forward. We use two complete quotient enumerations:

1. Enumerate all subsets of the $\binom{q}{2}$ forward pairs and keep the transitive relations.
2. Recursively add a last maximal element. Its predecessors must be an order ideal of the previous poset, and each such ideal gives one extension.

For every quotient, enumerate all positive ordered compositions of each allowed total core size into q block sizes. Expand each block to mutually equivalent vertices and each strict comparison to all corresponding directed arcs. Every preorder is represented up to relabeling. Repetitions are allowed; the recorded counts are not counts of isomorphism classes or of labeled preorders.

The naturally labeled quotient-poset counts for $q = 1, \dots, 7$ are

$$1, 2, 7, 40, 357, 4824, 96428.$$

The expanded-core counts for total size $n = 1, \dots, 7$ are

$$1, 3, 12, 68, 568, 7090, 131645. \quad (2.6)$$

For a fixed n , the count equals $\sum_q Q_q \binom{n-1}{q-1}$, where Q_q is the corresponding quotient count.

2.4.2 Counting supports without counting matching multiplicities

For each ordered pair of disjoint equally sized endpoint sets, support feasibility is a Boolean perfect-matching test in the directed incidence relation $S \rightarrow T$. The package cross-checks three implementations: recursive matching search; Hall's condition for every subset of S ; and bottom-up construction of directed matchings with endpoint pairs stored in a set. The last method explicitly discards multiple realizations of the same support.

For a fixed core and exterior types, let q_T be the number of selected exterior vertices of type T . For fixed labeled representatives of those selected vertices, enumerate their source/target roles and the unused/source/target roles of core vertices; retain a role assignment exactly when a realizing matching exists. If this gives $c_{k,q}$ feasible supports, then

$$\gamma_k(\mathbf{n}) = \sum_q c_{k,q} \prod_T \binom{n_T}{q_T}. \quad (2.7)$$

A support selects its exterior subsets uniquely, so this is an exact identity. Every used exterior vertex needs a different attachment in a realizing matching. Thus $\sum_T q_T \leq a$, making the sum finite regardless of the populations. In particular, γ_k has total population degree at most a , and γ_1 has degree at most one.

Multiplication in the binomial basis can be done with integer arithmetic using

$$\binom{n}{a} \binom{n}{b} = \sum_{j=0}^{\min(a,b)} \frac{(a+b-j)!}{j!(a-j)!(b-j)!} \binom{n}{a+b-j}. \quad (2.8)$$

This counts pairs of subsets by the size j of their intersection. Taking products of this identity over the types gives exact binomial-basis expansions for D .

2.4.3 Finite certificates for an infinite population domain

A polynomial with nonnegative coefficients in the multivariate binomial basis is nonnegative on $\mathbb{Z}_{\geq 0}^r$. For other polynomials we partition $\mathbb{Z}_{\geq 0}^r$ according to the set F of positive coordinates and substitute

$$n_i = 0 \ (i \notin F), \quad n_i = 1 + x_i \ (i \in F), \quad x_i \geq 0. \quad (2.9)$$

Every nonnegative integer population vector lies on exactly one such face. A certificate is an exact identity expressing the substituted polynomial as a nonnegative rational combination of nonnegative monomials and terms $x^m Z(x)^2$ with $Z \in \mathbb{Q}[x]$.

Some square terms are stored compactly as three monomials $ux^\alpha + vx^\beta + wx^\eta$, where $u, v > 0$, $w < 0$, $\alpha + \beta = 2\eta$, and $w^2 = 4uv$. Factoring $x^{\min(\alpha, \beta)}$ leaves a square: the remaining endpoint exponents are even. All exponent vectors are nonnegative integers. General square terms include the rational polynomial Z as additional data. The verifier expands every term, checks every sign and exponent, and compares coefficients exactly.

Numerical linear programming and eigenvector-based column generation helped discover some certificates. They are not used to verify positivity. Verification contains no floating-point tolerance, approximate roots, or finite-population extrapolation. The face substitution proves an integer-population statement; it does not assert positivity for coordinates strictly between zero and one.

2.5 No attachments and one attachment

2.5.1 The finite real-rootedness lemma

Lemma 2.7 (Exact finite lemma). *For every preorder on at most seven vertices, Γ_τ has only real negative roots, with multiplicities allowed; the constant polynomial is included.*

Computational proof. The complete enumeration of Section 2.4 gives the core counts (2.6). There are no supports larger than three. A quadratic $1 + bz + cz^2$ is real-rooted exactly when $b^2 - 4c \geq 0$. A cubic $1 + bz + cz^2 + ez^3$, $e > 0$, is real-rooted exactly when

$$\Delta = b^2c^2 - 4c^3 - 4b^3e - 27e^2 + 18bce \geq 0. \quad (2.10)$$

Every enumerated coefficient vector passes the applicable exact integer test. Positivity of the coefficients excludes nonnegative roots. The generator and independent Hall checker are supplied in `code/small-check/`; representatives of all distinct gamma vectors are included. Direct enumeration of all reflexive relations through four vertices gives the known labeled-preorder counts 1, 4, 29, 355 and exactly the same coefficient sets. These checks supplement the coverage proof, rather than replace it. $\square$

The literature already reports computer verification through seven vertices [2, Section 5]; Lemma 2.7 is independently reproduced here, not claimed as a new size range. The supplied final enumerator also verifies every degree-at-most-three preorder on eight vertices. It skips degree-four cases and is not evidence for a claim about their gamma polynomials.

If $a = 0$, there are no edges from D_G to C or between different D_i . The total order of C is at most six, and each nonsingleton D_i has at most seven vertices by the matching budget. Thus every nontrivial connected component has at most seven vertices. Support polynomials multiply over graph components: restricting a realizing matching gives equally sized tail/head sets in each component, and taking unions reverses this restriction. Lemma 2.7 therefore makes the whole polynomial real-rooted. Newton's inequalities imply rank-ULC with its total actual degree. This proves the $a = 0$ branch without a separate ULC convolution argument.

2.5.2 One attachment and the support identity

Suppose $a = 1$, with attachment v . The core has at most seven vertices. After deleting isolated exterior vertices, every exterior vertex is a pendant neighbor of v , and all have the same direction. Reverse all arcs if needed to orient them $v \rightarrow x$. For $m > 0$, transitivity implies that no other core vertex has an arc into v , since $u \rightarrow v \rightarrow x$ would force an additional neighbor of x . Thus v is a singleton minimal element of the core preorder.

If the exterior population is m , a support either avoids the exterior or uses exactly one exterior vertex. In the latter case the matching must use the pendant edge, and deletion of its endpoints gives a support in $K - v$. Conversely, every such support extends in m ways. Consequently

$$\Gamma_m(z) = \Gamma_K(z) + mz\Gamma_{K-v}(z). \quad (2.11)$$

If $m > 0$, a matching in $K - v$ extends by a pendant edge, so $\nu(K - v) \leq 2$. The case $m = 0$ is already Lemma 2.7.

Write $\Gamma_K = 1 + p_1z + p_2z^2 + p_3z^3$ and $\Gamma_{K-v} = 1 + q_1z + q_2z^2$. The two cubic gaps in (2.11) are quadratics in m with coefficient triples (A, B, C) :

$$D_1(m) : \quad (1, 2p_1 - 3q_1, p_1^2 - 3p_2), \quad (2.12)$$

$$D_2(m) : \quad (q_1^2 - 3q_2, 2p_2q_1 - 3p_1q_2 - 3p_3, p_2^2 - 3p_1p_3). \quad (2.13)$$

Every certificate is either $A, B, C \geq 0$, or $A > 0$ and $4AC - B^2 \geq 0$, using

$$4A(Am^2 + Bm + C) = (2Am + B)^2 + (4AC - B^2). \quad (2.14)$$

These prove nonnegativity for every real $m \geq 0$, a strengthening of what the preorder application requires.

The complete marked-core enumeration has 198469 valid marked expansions and 2050 distinct tuples $(p_1, p_2, p_3, q_1, q_2)$. By total core size the marked counts are

$$1, 3, 16, 115, 1160, 16786, 180388.$$

All 2050 first gaps and all 2050 second gaps pass: respectively 1114 and 1697 use nonnegative coefficients, and the remaining 936 and 353 use (2.14). Where $p_3 = q_2 = 0$, the actual degree can be two for all m ; all 136 such tuples also certify $(p_1 + m)^2 - 4(p_2 + mq_1) \geq 0$. At $m = 0$ with $p_3 = 0$, the corresponding quadratic normalization is checked separately.

The independent C++ checker generates quotient posets by ideal extension and constructs supports by endpoint-mask dynamic programming. It reproduces every one of the 2050 tuple multiplicities, not merely one representative or the total count. A separate Python checker validates representatives and all 4236 stored quadratic certificates. This closes $a = 1$.

2.6 Two attachments

Let $a = 2$, so $|K| \leq 5$. Label the attachments v_0, v_1 . Exterior neighborhoods are the nonempty subsets of this pair, encoded as masks 1, 2, 3. For every core preorder, pair of attachments, and two signs, the generator keeps exactly the types allowed by Lemma 2.5. It then identifies templates under permutations of the core respecting the attachment pair, interchange of the two attachments, and simultaneous reversal of all arcs.

Every resulting blow-up has matching number at most three, even when it is not a canonical Gallai–Edmonds instance. Indeed, a matching using r exterior edges has $r \leq 2$ and size at most

$$r + \left\lfloor \frac{5-r}{2} \right\rfloor \leq 3.$$

There are at most three population variables. By (2.7), every gamma coefficient has population degree at most two and D has degree at most four.

Proposition 2.8 (Exact two-attachment certificate). *For every core preorder of order at most five, every pair of attachment vertices, every fixed direction choice, and all nonnegative integer populations of the compatible exterior types, $D \geq 0$.*

Computational proof. The complete generator produces 1084 structural templates, with 5, 23, 141, 915 templates for core orders 2, 3, 4, 5, respectively. They yield 448 distinct full gamma-polynomial families. Grouping is used only when all coefficient polynomials agree in their ordered population variables. Of these, 284 have a coefficientwise nonnegative binomial expansion of D . Each of the other 164 is supplied with all its zero/positive face certificates, totaling 994 faces. Each face identity has the exact nonnegative form described in Section 2.4. Thus every nonnegative integer population vector is covered.

An independent audit enumerates all labeled preorders through core order five, obtaining 4, 29, 355, 6942 at orders 2, 3, 4, 5, respectively. It reproduces the complete set of 1084 canonical templates, reconstructs support polynomials using directed matchings deduplicated by endpoint sets, recomputes D , and checks every rational face identity. The audit does not call the generator's helper routines. An additional independent Hall implementation checks 19084 labeled blow-ups on the grid $\{0, 1, 2, 3\}$ and recovers the binomial coefficients by finite differences; the known degree bound makes this polynomial reconstruction exact. This is an additional support-count check, not the proof of positivity on the unbounded domain. $\square$

The files `templates.json`, `gap2_certificates.json`, the exact verifier, and both independent checking routes are in `code/two-attachment/`. A template with no allowable exterior type contributes only its small core and is covered by Lemma 2.7; omitting it from the nonempty-type export loses no case. If a specialization has actual degree at most two, Proposition 1.5 supplies the correct degree-two normalization. Hence Proposition 2.8 closes the $a = 2$ branch.

2.7 Three attachments and seventeen templates

For $a = 3$, equation (2.4) forces $C = \emptyset$ and every D_i to be a singleton. Thus $K = A$ consists of exactly three vertices. The canonical exterior has singleton equivalence classes by Lemma 2.4. Every attachment has an exterior neighbor because a maximum matching matches all of A into the singleton D_i .

There are 29 labeled three-element preorders. Test all eight active sign vectors and all seven nonempty neighborhood masks. Requiring every attachment to have an allowed neighbor gives 130 labeled templates and 17 classes under core permutations and order reversal. The full list is below. A minus sign denotes core-to-exterior arcs, a plus sign exterior-to-core arcs. Mask t encodes the subset of $\{0, 1, 2\}$ whose bits are one. The listed core arcs exclude loops and include transitive arcs.

Class	Core arcs	Signs	Allowed masks
0	$\emptyset$	---	1, 2, 3, 4, 5, 6, 7
1	$\emptyset$	--+	1, 2, 3, 4
2	01	---	1, 3, 4, 5, 7
3	01	--+	1, 3, 4
4	01	-+-	1, 2, 3, 4, 5
5	01, 02	---	1, 3, 5, 7
6	01, 02	--+	1, 3, 4, 5
7	01, 02	-++	1, 2, 3, 4, 5, 6, 7
8	01, 02	+++	2, 4, 6, 7
9	01, 10	---	3, 4, 7
10	01, 10	--+	3, 4
11	01, 02, 12	---	1, 3, 7
12	01, 02, 12	--+	1, 3, 4, 5, 7
13	01, 02, 10, 12	---	3, 7
14	01, 02, 10, 12	--+	3, 4, 7
15	01, 02, 10, 12	+++	4, 7
16	01, 02, 10, 12, 20, 21	---	7

2.7.1 The bipartite template

Class 0 is a bipartite graph with a source shore of size three and a target shore consisting of the exterior. Its directed disjoint supports are precisely the ordinary bipartite matching supports. The ambient-binomial normalization theorem of Röhrle and Ulirsch [13, Theorem A and Example 2.4] applies: for a relation on disjoint shores of sizes m, n , the support numbers normalized by $\binom{\min(m, n)}{k}$ are log-concave. If the actual degree is three here, the smaller shore has size three, so this is exactly $D \geq 0$. Lower actual degrees are already settled by Proposition 1.5. [*Merge note:* class 0 is also a case of *preorder-root-polytopes*, Theorem `lor:thm:weightedULC` (two-shore ultra-log-concavity normalized by the smaller shore), proved there with the same transversal-matroid device.]

For clarity, the relation support numbers fit the published theorem by assigning an independent indeterminate to each edge of the incidence matrix. A minor is nonzero precisely when its row and column sets support a matching: different matching terms have different monomials and

cannot cancel. These minors define a representable bimatroid. No theorem about normalizing an arbitrary bimatroid by its actual rank is being invoked. This published ambient-normalization theorem is the only non-elementary polynomial-inequality input for the seventeen-template step.

2.7.2 Compact product and extension certificates

In classes 1, 3, 10, the polynomial factors as

$$\Gamma(z) = (1 + Az + Bz^2)(1 + Cz), \quad C = n_4.$$

Consequently

$$D = (B - AC/2)^2 + \frac{3}{4}C^2(A^2 - 4B). \quad (2.15)$$

The remaining expressions are

$$\begin{aligned} \text{class 1: } \quad & A = n_1 + n_2 + 2n_3, \quad B = n_1n_2 + (n_1 + n_2)n_3 + \binom{n_3}{2}, \\ & A^2 - 4B = (n_1 - n_2)^2 + 2n_3^2 + 2n_3; \\ \text{class 3: } \quad & A = n_1 + 2n_3 + 1, \quad B = n_1n_3 + \binom{n_3}{2}, \\ & A^2 - 4B = n_1^2 + 2n_1 + 2n_3^2 + 6n_3 + 1; \\ \text{class 10: } \quad & A = 2n_3 + 2, \quad B = \binom{n_3}{2}, \quad A^2 - 4B = 2n_3^2 + 10n_3 + 4. \end{aligned}$$

These are nonnegative on the required domain. For class 9, subtract the class-10 gap. All coefficients of the difference are nonnegative except possibly those in the pure n_7 contribution. That whole univariate contribution is exactly

$$4\binom{n_7}{1} + 41\binom{n_7}{2} + 57\binom{n_7}{3} + 18\binom{n_7}{4}, \quad (2.16)$$

which is nonnegative for integer $n_7 \geq 0$. The coefficient assertion and identity are verified exactly in `verify_cover3_compact.py`. Classes 11, 13, 15, 16 have nonnegative multivariate-binomial expansions of D directly.

2.7.3 The complete face certificates

Classes 5, 6, 8, 12, 14 have a certificate for every zero/positive face using monomials and monomial-weighted binomial squares. Class 2 has such certificates on 31 of its 32 faces. The remaining face has only n_3, n_4 positive; its gap is exactly that of class 3 at $n_1 = 0$, so (2.15) applies.

Classes 4 and 7 need some general rational square polynomials as well. Class 4 has 23 binomial-square face certificates and 9 general-square certificates, covering all 32 faces. Class 7 has $73 + 55 = 128$ certificates, covering all its faces. The hardest saved class-7 identity has 210 nonzero square/monomial weights; the checker expands its 330 coefficient equations over the rationals.

The independent script `audit_cover3.py` separately enumerates the 29 core preorders, reproduces the 130 signed templates and all 17 canonical keys, and reconstructs every coefficient in (2.7) by Hall's condition on ordered endpoint masks. It recomputes every gap from the gamma coefficients and verifies the face partitions, missing-face reductions, exponent domains, and duplicate guards. The three exact identity verifiers then check all saved rational identities. Thus the certificates are linked to the intended support counts, not merely to an unchecked polynomial export. This proves $D \geq 0$ throughout the $a = 3$ branch.

2.8 Completion of the main theorem

Proof of Theorem 2.3. Lemma 1.3 identifies the degree and excludes internal zeros. Degrees zero and one are immediate; degree two follows from Proposition 1.5. In degree three the same proposition proves the first inequality. Apply the Gallai–Edmonds decomposition to the comparability graph. Equation (2.5) gives exactly four possibilities. The $a = 0$ branch follows from the finite real-rootedness lemma and component multiplication; the $a = 1$ branch from (2.11) and its complete quadratic certificates; the $a = 2$ branch from Proposition 2.8; and the $a = 3$ branch from Section 2.7. These prove the second inequality in every case. All population sizes are unrestricted nonnegative integers throughout these four arguments. $\square$

The reduction is essential to the quantifier. A finite scan of all preorders up to a fixed number of vertices would not prove the theorem. Here the graph structure bounds the core, and algebraic certificates handle the unbounded exterior populations. The finite base computation for $a = 0$ is sufficient because each component is provably small, even though isolated components may occur in arbitrarily large number.

2.9 Real roots and an explicit obstruction

Rank-ULC is strictly weaker than real-rootedness even in actual degree three. Take three minimal elements u_0, u_1, u_2 and three disjoint four-element sets X, Y, Z of maximal elements. Let the strict relations be exactly

$$u_0 < X \cup Y, \quad u_1 < X \cup Z, \quad u_2 < Y \cup Z.$$

This is a height-two poset on fifteen vertices. Its three minimal vertices form a vertex cover, and a matching saturating them exists, so the actual degree is three.

There are 24 arcs. For each choice of two minimal vertices, all pairs of maximal vertices are feasible except pairs contained in either of the two four-element types that see only one chosen minimal vertex. Thus

$$\gamma_2 = 3 \left(\binom{12}{2} - 2 \binom{4}{2} \right) = 162.$$

A triple of maximal vertices fails to match all three minima exactly when all three lie in the same four-element type. Hall’s condition has no other obstruction because every maximal vertex sees two minima. Hence

$$\gamma_3 = \binom{12}{3} - 3 \binom{4}{3} = 208.$$

Therefore

$$\Gamma(z) = 1 + 24z + 162z^2 + 208z^3, \quad \Delta = -2592 < 0. \quad (2.17)$$

Its two cubic Newton gaps are 90 and 11268, both positive, consistent with Theorem 2.3. A direct endpoint-support enumerator independently reproduces (2.17); no population formula is used in that check.

[Merge note: the same example, with the same numbers, is computed in source 22 (Section “A cubic with nonreal zeros”, where it is noted that it lies in the pure three-tail family of source 22’s three-core theorem and needs no role asymmetry), in source 16 (“The preceding research ... exhibited the nonreal-rooted cubic”) and in archive 31. It is printed once, here.]

2.9.1 The exact minimum order in degree three

Theorem 2.9 (Minimum order and uniqueness). *Every finite preorder on at most fourteen vertices whose gamma polynomial has actual degree three is real-rooted. Among fifteen-vertex*

preorders of actual degree three with a nonreal gamma zero, there are exactly two ordinary isomorphism classes, represented by the height-two poset in (2.17) and its dual. Equivalently, there is one class modulo isomorphism and global order reversal.

Computer-assisted proof with global coverage. Apply the canonical four-branch reduction. The $a = 0$ branch is real-rooted at every allowed order by Lemma 2.7 and component multiplication. For $a = 1, 2, 3$, enumerate every population specialization whose effective total order is at most fifteen and whose cubic coefficient is positive. Test the exact integer discriminant (2.10). The complete counts are

Branch	Evaluations	Actual cubics	Negative discriminants
One attachment	19430	17785	0
Two attachments	84191	58800	0
Three attachments	128726	121510	1

The sole negative case has total order fifteen, class 0 of Section 2.7, and populations $(n_1, \dots, n_7) = (0, 0, 4, 0, 4, 4, 0)$. It is exactly (2.17). An independently written scanner regenerates every per-size count and the complete negative-case list using arbitrary-precision integers.

There are two important coverage points. In the two-attachment scan we retain all 1084 structural templates and their actual core sizes, not only the 448 polynomial families. In the one-attachment scan we group by $(p_1, p_2, p_3, q_1, q_2)$ but retain a *minimum-order* realization of every tuple. The complete marked-core generator explicitly minimizes this size over all occurrences; all 2050 stored minima are verified by a fresh enumeration. Replacing a core by this minimum-size representative preserves (2.11) and can only reduce total order. Hence any example through a bound appears at some scanned order through that bound. Isolated vertices may be deleted or added without changing gamma. The displayed counts are therefore representation counts, not a per-order census of polynomials or preorder isomorphism classes.

For uniqueness, any nonreal cubic on at most fifteen vertices must lie in the canonical $a = 3$ branch, after deleting isolates. Up to a permutation of the three core vertices and possibly global duality, its core, signs, and populations occur in the complete seventeen-template scan. The sole negative specialization has effective order fifteen, so no isolates were deleted. Its populations determine the directed relation up to permutations within types. Thus the preorder is the displayed poset or its dual. They are not isomorphic: one has three minimal and twelve maximal elements, and its dual has twelve minimal and three maximal elements. $\square$

[Merge note: this theorem supplies the minimality statement that source 22 explicitly does not make (“No minimal-size claim is made”). It does not conflict with the nonreal gamma polynomials of *preorder-root-polytopes*: the eight-vertex example Θ_3 there has gamma degree four, and the only degree-three member of that family is real-rooted.]

This theorem concerns actual degree three only. It makes no claim about higher-degree preorders on at most fourteen vertices. Its bounded discriminant scan is logically separate from the all-population second-gap certificates used for Theorem 2.3. The scripts and receipts are in `code/minimum/`.

2.10 An infinite family with three negative roots

A complementary positive result holds for a large antichain whose complement has three vertices. Let $\mathcal{A}_m$ be an antichain of size m , and let $\oplus$ denote ordinal sum, with all elements of an earlier summand strictly below all elements of a later one.

Theorem 2.10. *If $m \geq 3$ and P, Q are arbitrary finite preorders with $|P| + |Q| = 3$, allowing either to be empty, then $\Gamma_{P \oplus \mathcal{A}_m \oplus Q}$ has degree three and three distinct negative real roots. Every ordinal sum of antichains of actual gamma degree three also has three distinct negative real roots.*

Proof. Every edge meets the three-element complement of $\mathcal{A}_m$, and its vertices can be matched to three distinct antichain vertices, so the degree is three. Reversal reduces the possible splits to two forms.

For $\mathcal{A}_m \oplus R$, $|R| = 3$, let e be the number of off-diagonal arcs in R and let t be the number of its vertices with at least one outgoing off-diagonal arc. Then

$$(\gamma_1, \gamma_2, \gamma_3) = \left(3m + e, 3 \binom{m}{2} + tm, \binom{m}{3} \right). \quad (2.18)$$

The feasible (e, t) are $(0, 0), (1, 1), (2, 1), (2, 2), (3, 2), (4, 2), (4, 3), (6, 3)$. For $1 \oplus \mathcal{A}_m \oplus R_2$, $|R_2| = 2$, let $j \in \{0, 1, 2\}$ be its number of off-diagonal arcs. Then

$$(\gamma_1, \gamma_2, \gamma_3) = \left(3m + 2 + j, 5 \binom{m}{2} + (j + 1)m, 3 \binom{m}{3} \right). \quad (2.19)$$

To justify (2.18), a size-two support with two antichain points chooses two of the three core heads, while one with a single antichain point chooses a core tail with an outgoing arc. A size-three support uses all three core heads. For (2.19), two chosen antichain points admit five size-two role patterns: one with both as tails, and four with one tail and one head. One antichain point gives $j + 1$ patterns, and three give three choices for the head matched from the lower singleton. These are Boolean support counts.

For all eleven forms, put $x = m - 3$. The coefficients of $12\Delta/(x+3)$ are listed in Appendix 2.12; every one is positive. Thus $\Delta > 0$ for $m \geq 3$. The cubic has three distinct real roots and, by positive coefficients, they are negative.

For an ordinal sum of antichains with total size n and maximum layer size M , its comparability graph is complete multipartite and has matching number $\min(\lfloor n/2 \rfloor, n - M)$. If $M \geq n/2$, match all outside vertices into the largest layer. If $M < n/2$, pairing vertices from two largest remaining parts constructs a matching missing at most one vertex; the invariant that no part exceeds the ceiling of half the remaining total is preserved. Degree three consequently means either $n \in \{6, 7\}$ and $M \leq n - 3$, or $n \geq 8$ and $M = n - 3$. Whenever $M = n - 3$, the preceding result applies. The remaining compositions have $n = 6$ with all layers at most two, or $n = 7$ with all layers at most three: respectively 13 and 44 cases. Their exact cubic discriminants are positive in the supplied table `layered_certificates.json`. An independent layer-quota recurrence and direct support enumeration verify all 57 cases. $\square$

Two specializations are already in the literature. The two-antichain case $(e, t) = (0, 0)$ is covered by Athanasiadis and Chapoton [2, Proposition 9.2]. The case $(e, t) = (6, 3)$, a three-element equivalence block above singleton leaves, is covered by Athanasiadis, Xiao, and Yan [3, Remark 2.3]. Neither specialization is claimed as new. The eleven-template statement is included as a proved complementary family; a complete literature-priority audit for its precise scope has not been performed.

2.11 Reproducibility and scope

The archive contains the editable LaTeX source, this PDF, exact source code, certificate data, checking receipts, and a checksum manifest. It contains no required compiled binaries or numerical-optimizer state. The top-level `README.md` gives the dependency versions and commands. The

primary checks need Python 3 with SymPy and a C++17 compiler. No optimization package is required to verify the saved certificates.

The command `python3 verify_all.py` runs the exact checking routes and writes fresh logs under `receipts/`. It checks the finite-core lemma; the full marked-core enumeration and all one-population quadratic identities; independent two-attachment coverage and all its rational certificates; the seventeen three-cover templates and their exact identities; the ordinal-sum family; the bounded scans proving Theorem 2.9; and the explicit nonreal-root example. The finite generators can also be rerun separately using the commands in the README. The main theorem depends on the published Gallai–Edmonds theorem and the class-0 ambient-normalization theorem as clearly stated in the text, not on a numerical solver.

[Merge note: file names in this section are those of source 25’s delivered package; the files are shipped as `code/25-degree-three-...` and `data/25-degree-three-...`, and the README maps the names and gives rerun instructions. At intake every step of `verify_all.py` reproduced its recorded outputs (on a copy, compared modulo line endings and timing fields) except the order-eight census of `small-check/exhaust`, whose optimized build crashed after finishing its enumeration and was not rerun; the C++ sources also need `-include cassert` with GCC 16. Of the research directions below, the first is answered by source 16 (Section 3) and the fourth, in degree three, by source 22 (9); the second, third and fifth remain open.]

The following directions remain meaningful beyond this result.

1. **Actual degree four.** The same matching-budget argument gives bounded canonical cores, but more exterior types and higher-degree inequalities. A direct certificate approach must control that growth rather than infer higher-rank behavior from the cubic case.
2. **A conceptual second-gap proof.** The rational certificates may reveal common decompositions that replace the finite case analysis. At present the general cubic second inequality genuinely uses the supplied exact certificate computation.
3. **Equality in the second inequality.** The first-gap equality theory in the original report is not automatically a classification for the second gap. Extracting and reconciling the zero sets of the certificates is a separate problem.
4. **Weighted variants.** The present theorem uses unweighted supports. Replacing populations by arbitrary real activities changes the domain and the coefficient model; the integer-face certificates do not by themselves establish such a theorem.
5. **Formal verification.** The small support kernels and rational identities are natural candidates for a proof-assistant checker. Exact arithmetic and independent implementations are valuable checks, but are not a formalized proof.

2.12 Explicit discriminant certificates for the ordinal family

For (2.18) and (2.19), the table lists the coefficients in ascending powers of x of $12\Delta/(x+3)$, with $x = m - 3$. Thus each row gives a standalone integer positivity certificate. The common factor $x + 3$ is positive for $x \geq 0$.

Form	Parameters	x^0	x^1	x^2	x^3	x^4	x^5
Outer	(0, 0)	8640	13248	7884	2286	324	18
Outer	(1, 1)	22484	30336	15577	3789	432	18
Outer	(2, 1)	30148	37836	17888	4014	432	18
Middle	$j = 0$	41412	46188	18456	3378	288	6
Outer	(2, 2)	45376	55296	25472	5490	540	18
Outer	(3, 2)	60804	69480	29613	5877	540	18
Middle	$j = 1$	76356	78552	28221	4473	324	6
Outer	(4, 2)	76880	83328	33268	6174	540	18
Middle	$j = 2$	129156	123780	40644	5718	360	6
Outer	(4, 3)	107300	112980	44092	7938	648	18
Outer	(6, 3)	163620	158076	55080	8766	648	18

The generator `core_templates.py` independently enumerates all 29 labeled three-element preorders and all 82 compatible choices of the core vertices below the antichain, recovering these eleven polynomial forms. It symbolically counts roles with multiplicity $\binom{m}{a}\binom{m-a}{b}$ for a antichain tails and b antichain heads. The layer checker processes antichain layers from bottom to top: if b unmatched earlier tails are available and k tails have been selected so far, choosing a new tails and c new heads is feasible exactly when $c \leq b$. It has multiplicity $\binom{n_i}{a}\binom{n_i-a}{c}$ and updates (b, k) to $(b + a - c, k + a)$. Finishing with $b = 0$ gives the support counts. This recurrence is compared with literal disjoint-support enumeration for all 96 compositions of six and seven.

3 Degree four (source 16, the base)

Source 16 is *Degree four preorder support polynomials* (17 pages as delivered), the four-part package `preorder-degree-four` (archives 16–19); it is the base of this report. Its files are shipped with the prefix `16-degree-four-`, except those that are byte copies of files of sources 20, 22 and 23, which are shipped once under those sources’ prefixes. Its abstract describes, among other things, the role-cover theorem of its Section 3, which is source 20’s article; see Section 3.3.

Abstract of source 16, as delivered. We study the polynomial counting disjoint directed matching supports of a finite preorder, with each feasible pair of endpoint sets counted once. We prove that the unweighted support polynomial of every finite preorder is rank-ultra-log-concave whenever its actual degree is at most four. The proof reduces arbitrary vertex counts to finitely many bounded cores with unbounded integer populations, and certifies the resulting inequalities by exact rational identities. Independently, we prove a structural theorem for every loopless directed relation whose arcs are covered by at most two selected tail vertices and two selected head vertices, with overlap allowed: its signed physical monomer polynomial is real stable, even with arbitrary nonnegative independent tail and head activities. Its support polynomial therefore has only negative real roots. The proof couples two rank-two transversal matroids through a stability-preserving operator and then merges the two copies of each physical vertex without allowing double use. For an unbalanced one-source, three-sink core, an ordinary proof of the last Newton inequality combines Lorentzian polarization with the Rayleigh property of rank-three matroids. A general exterior-only Lorentzian construction, an exact Gallai–Edmonds reduction, and independently checked rational certificates explain both the mathematical scope and the computational component. The final four-attachment ledger covers all 76 canonical templates; its last certificate consists of 2,273 rational square orbits and has been independently replayed. The result is computer-assisted, with ordinary structural proofs and finite algebraic verification distinguished throughout.

3.1 The support polynomial and the questions

A finite preorder is a reflexive transitive relation on a finite set V . Its off-diagonal relation is regarded as a directed graph D : an arc $i \rightarrow j$ means $i \neq j$ and i precedes j . Equivalent elements give opposite arcs. Several structural results below apply to directed relations without transitivity.

Definition 3.1. A feasible support of size k is an ordered pair (S, T) of disjoint subsets of V , each of cardinality k , for which there exists a bijection $\pi : S \rightarrow T$ with $s \rightarrow \pi(s)$ for every $s \in S$. Write

$$\gamma_k(D) = \#\{(S, T) : (S, T) \text{ is feasible of size } k\}, \quad \Gamma_D(t) = \sum_{k \geq 0} \gamma_k(D) t^k.$$

The empty support contributes $\gamma_0 = 1$. A support contributes once, regardless of the number of witnessing bijections.

The same-ground-set disjointness is essential. Passing to independent tail and head copies without subsequently excluding their simultaneous use changes this polynomial. Counting individual matchings also changes it.

For nonnegative tail activities u_i and head activities v_i , replace the contribution of (S, T) by

$$u^S v^T = \prod_{i \in S} u_i \prod_{j \in T} v_j.$$

We use the same notation when the weights are specified. A vertex may have different activities in its two roles. With zero activities, the polynomial degree can decrease.

Preorder polytopes and their lattice-point enumerators were introduced and studied by Chapoton and Athanasiadis [2]. The supplied ProveIt report identifies the unweighted support polynomial with its preorder gamma polynomial [1]. Our proofs concern the explicit support definition, so their validity does not require a particular formalization of that identification.

[Merge note: source 16’s Lemma “degree” is Lemma 1.3, printed once in Section 1.3 with source 16’s own proof.]

Definition 3.2. A polynomial $g(t) = \sum_{k=0}^d g_k t^k$ is *rank-ultra-log-concave*, or rank-ULC, if its coefficients are nonnegative with no internal zeros and

$$\left(\frac{g_k}{\binom{d}{k}} \right)^2 \geq \frac{g_{k-1}}{\binom{d}{k-1}} \frac{g_{k+1}}{\binom{d}{k+1}} \quad (1 \leq k < d),$$

where d is the actual degree.

In degree four the inequalities are

$$3\gamma_1^2 \geq 8\gamma_2, \quad 4\gamma_2^2 \geq 9\gamma_1\gamma_3, \quad 3\gamma_3^2 \geq 8\gamma_2\gamma_4. \quad (3.1)$$

Normalization by the ground-set size, by a larger cover size, or by a presumed degree after some weights vanish is a different assertion.

[Merge note: source 16’s Theorem “Universal first inequality” (if the underlying graph has matching number at most $r \geq 2$, then $\gamma_1^2 \geq \frac{2r}{r-1}\gamma_2$, including nonnegative role activities) is Theorem 1.5, printed once in Section 1.3 with source 16’s Motzkin–Straus proof; it is *preorder-root-polytopes*, Theorem `mr:thm:first`.]

The preceding research established rank-ULC for every finite preorder of actual degree at most three by a separately auditable finite reduction (source 25, Theorem 2.3). It also exhibited the nonreal-rooted cubic $1 + 24t + 162t^2 + 208t^3$, whose discriminant is -2592 (Section 2.9). Thus real-rootedness cannot be assumed in the general preorder problem.

3.2 An exterior-only Lorentzian construction

We use standard facts about Lorentzian polynomials: matroid basis polynomials are Lorentzian; products, nonnegative linear substitutions, coordinate upper truncations, and polarization preserve this class [6, 12]. For a homogeneous bivariate polynomial $\sum a_k z^{r-k} t^k$, the Lorentzian condition gives log-concavity of $a_k / \binom{r}{k}$ with interval support.

Theorem 3.3. *Let P, Q, I be disjoint finite sets, and let the only arcs be $P \rightarrow I$ and $I \rightarrow Q$. Put $r = |P| + |Q|$. With independent nonnegative exterior role activities, the polynomial*

$$H(z, t) = \sum_{k=0}^r F_k z^{r-k} t^k$$

is Lorentzian, where F_k counts disjoint supports of size k . In particular $F_k / \binom{r}{k}$ is log-concave. This is actual-degree ULC when $\deg F = r$.

Proof. For the P side form a transversal matroid on the physical exterior vertices and private dummy elements d_p , one for each $p \in P$. Its presentation has right side P ; an exterior vertex has its given neighbors, and d_p is adjacent only to p . The dummies form a basis, so the rank is exactly $|P|$.

For $P' \subseteq P$ and $R \subseteq I$, the set $R \cup \{d_p : p \in P \setminus P'\}$ is a basis precisely when $|R| = |P'|$ and R can be matched to P' . This encodes an endpoint support once, rather than counting its matchings. Construct the analogous Q matroid with distinct dummies and the same physical variables x_i .

Their squarefree basis polynomials B_P and B_Q are Lorentzian. Multiply them, then discard every monomial in which some x_i has exponent greater than one. Coordinate upper truncation preserves Lorentzianity [12, Proposition 3.3]. It removes exactly the choices using a physical exterior vertex in both roles. The all-dummy monomial survives.

Finally set every dummy variable equal to z and every physical variable equal to t . A surviving pair of bases specifies the support $S = P' \cup L$, $T = Q' \cup R$, where $L, R \subseteq I$ are disjoint. It contributes $z^{r-|L|-|R|} t^{|L|+|R|}$. This is exactly H . Exterior activities are introduced by scaling x_i separately in the two factors before multiplication. Every operation retains homogeneous degree r . $\square$

Remark 3.4. If H has a common factor z , dividing that factor out is not a generally valid Lorentzian closure operation. The theorem does not silently strengthen its order- r normalization to a smaller actual degree. It also does not assert real-rootedness.

For $r = 4$, differentiation of H twice gives the three constants in (3.1). For example $\partial_t^2 H = 2F_2 z^2 + 6F_3 z t + 12F_4 t^2$, whose discriminant yields $3F_3^2 \geq 8F_2 F_4$.

3.3 The two by two role cover

Source 16's Section 3, "Real-rootedness for a complete balanced core" (its Theorems "balanced", "twoedge", "missing" and "rolecover", its Lemmas "quadratic" and "asano", its Corollary "freeprod", the corollary on the gamma expansion, the interpolation remark, and its Appendix A of seven Rayleigh identities for two disjoint core edges), is source 20's article *Real stable monomers under a two by two role cover*: 176 of the 218 sentences of source 20 recur verbatim in source 16, with the same labels. It is printed once, from source 20, in Part II (Section 5; the main theorem is Theorem 5.10). Source 16's text differs in four places, all printed there as marked second routes:

1. where source 20 invokes the stability-preserver theorem of Borcea and Brändén, source 16 proves each operator elementarily: an elementary Asano contraction lemma, a polarized auxiliary quadratic for the rank-two coupling, and the copy merge by diagonalization and differentiation;
2. the two-edge core covers canonical template 6, and the core with one missing edge covers canonical template 21, of the degree-four classification below;
3. the gamma-transfer corollary is stated with $N = |V|$; source 20 allows any $N \geq 2d$;
4. the appendix of source 20 has a typographical slip (“labels0,1”) that source 16 does not.

The degree-four ledger below discharges some templates by ordinary theorems, among them the two-edge core and the core with one missing edge (templates 6 and 21); the package records seven ordinary approvals, all for the last gap, of templates 1, 4, 6, 9, 10, 21 and 34.

3.4 The unbalanced one-source three-sink family

Let p be one source, let Q consist of three sinks, and let I be independent. Allow arbitrary arcs $p \rightarrow I$ and $I \rightarrow Q$, and any subset $p \rightarrow J$ of core arcs, where $J \subseteq Q$. No other arcs are present. All activities in this section are one.

Theorem 3.5. *Every relation in this family, and every reversal of such a relation, satisfies $3\gamma_3^2 \geq 8\gamma_2\gamma_4$.*

Let $Q(t) = 1 + Lt + Bt^2 + Tt^3$ be the exterior-to-three-sink polynomial. Let H be the exterior neighbors of p , $m = |H|$, and set

$$R(t) = \sum_{x \in H} Q_{-x}(t) = m + At + Ct^2 + Dt^3.$$

Let $Q^+(t)$ be Q with one new exterior tail of neighborhood J . Then

$$\Gamma_D(t) = Q^+(t) + tR(t). \quad (3.2)$$

Indeed, supports with no exterior head belong to Q^+ ; otherwise their unique exterior head x is matched from p , leaving exactly a Q_{-x} support. This excludes physical double use and introduces no matching multiplicity.

[Merge note: source 23 (Section 4) treats this family with the internal sink-side arc $q_2 \rightarrow q_0$ added (canonical template 26), where Lemma 3.6 is not available, and replaces it by a deletion-mixture lemma.]

Lemma 3.6. *The homogeneous degree-three polynomial $mz^3 + Az^2t + Czt^2 + Dt^3$ is Lorentzian. In particular $C^2 \geq 3AD$.*

Proof. Use the rank-three transversal matroid with private dummies. Identify dummy variables with z , the m variables indexed by H with s , and the other exterior variables with t . The resulting polynomial is Lorentzian. Polarize s into m slots using the degree bound m [6, Proposition 3.1]. Set one slot equal to zero and the other slots equal to t . A term using k elements of H is multiplied by $\binom{m-1}{k}/\binom{m}{k} = (m-k)/m$. Multiplication by m therefore gives the deletion sum. Vertices of H with no sink neighbor are matroid loops; they still count as unused polarization slots. If $m = 0$ the assertion is immediate. No homogenizing monomial is divided out. $\square$

Lemma 3.7. *For a three-center bipartite support polynomial, adjoining an exterior vertex cannot decrease the ratio of its cubic to quadratic coefficient, wherever that ratio is defined. Without division the assertion is $T_{\text{new}}B - TB_{\text{new}} \geq 0$.*

Proof. Every rank-three matroid is Rayleigh [14, Theorem 1.1]. Adjoin the new element e to the transversal matroid with three private dummy elements d_i . Let f be its basis polynomial. At existing exterior variables equal to one and all dummy variables and e equal to zero, we have

$$f = T, \quad \partial_e f = \Delta T, \quad \sum_i \partial_{d_i} f = B, \quad \sum_i \partial_e \partial_{d_i} f = \Delta B.$$

Summing the Rayleigh inequalities $(\partial_e f)(\partial_{d_i} f) - f \partial_e \partial_{d_i} f \geq 0$ gives $\Delta T B - T \Delta B \geq 0$. Boundary evaluations follow by continuity. An added loop changes neither coefficient. $\square$

Proof of Theorem 3.5. If $D = 0$, the desired inequality is immediate. Otherwise B, T, B^+, T^+, C are positive. For each deleted exterior vertex x , Lemma 3.7 applied when x is restored and when the new core tail is added gives $T_x B \leq B_x T$ and $T B^+ \leq B T^+$. Thus $T_x B^+ \leq B_x T^+$, without dividing by a possibly zero B_x . Summing gives $D B^+ \leq C T^+$. Together with Lemma 3.6 and (3.2),

$$\begin{aligned} 3\gamma_3^2 - 8\gamma_2\gamma_4 &= 3(C + T^+)^2 - 8(A + B^+)D \\ &\geq \frac{C^2}{3} - 2CT^+ + 3(T^+)^2 = \frac{(C - 3T^+)^2}{3} \geq 0. \end{aligned}$$

$\square$

The real-rootedness conclusion of the balanced theorem does not extend to the full unbalanced family. For example the checked polynomial $1 + 40t + 510t^2 + 2380t^3 + 2704t^4$ has discriminant -4112937216 and two real roots. It is realized with four exterior vertices of each of the three two-sink neighborhoods, thirteen source-only exterior vertices, and all three core arcs. Thus there are twenty-five exterior vertices and twenty-nine vertices in total. The package contains the exact population specification and a direct support verifier.

3.5 A finite normal form for bounded matching rank

Let G be the comparability graph of a preorder of matching number r . In the canonical Gallai–Edmonds decomposition, D is the set of vertices missed by some maximum matching, $A = N(D) \setminus D$, and $C = V \setminus (A \cup D)$. The components D_i of $G[D]$ are factor-critical, $G[C]$ has a perfect matching, and

$$r = |A| + \frac{|C|}{2} + \sum_i \frac{|D_i| - 1}{2}. \quad (3.3)$$

These standard facts are proved, for example, by West [17].

Proposition 3.8. *Let I be the union of the singleton components of $G[D]$, and put $K = V \setminus I$, $a = |A|$. Then I is independent, all its neighbors lie in A , and*

$$|K| \leq 3r - 2a.$$

Moreover, $\nu(G[K \setminus A]) = r - a$. Every nonisolated exterior vertex is a singleton preorder equivalence class. Each attachment has one strict orientation toward all its exterior neighbors. Hence at most $2^a - 1$ nonempty exterior types occur, with arbitrary integer populations.

[Merge note: this proposition is source 25’s Theorem 2.6 (its first half), with the same proof: the matching budget (3.3) gives $|K| \leq 3r - 2a$ and $\nu(G[K \setminus A]) = r - a$, and automorphism invariance of the canonical sets together with transitivity gives singleton exterior classes and one orientation per attachment (Lemma 2.4 there). Source 16’s four-paragraph proof is replaced by this pointer.]

For $r = 4$ the core bounds are 10, 8, 6, 4 for $a = 1, 2, 3, 4$. When $a = 0$, each nontrivial component has at most nine vertices. Polynomials multiply over comparability components because every support splits uniquely. Rank-ULC is preserved by products: homogenize each factor at its actual degree, use the bivariate Lorentzian characterization and product closure, then specialize. This uses no change of normalization.

3.6 Exact support kernels and coverage

For a fixed core and a list of legal exterior types, let n_t be the population of type t . If q_t selected exterior labels are prescribed, let $c_{k,q}$ count feasible supports in the induced relation on the core and precisely those prescribed labels, with every prescribed label required to occur. Then

$$\gamma_k(\mathbf{n}) = \sum_{|q| \leq \min(k,a)} c_{k,q} \prod_t \binom{n_t}{q_t}. \quad (3.4)$$

Every selected exterior vertex needs a different core partner, so at most a can occur. The binomial factors choose their physical labels. Feasibility depends only on the type quota and the selected core endpoint sets; unselected vertices cannot help match a fixed support. Thus (3.4) is an exact all-population formula, not interpolation from bounded numerical tests.

Independent kernel checks enumerate disjoint tail and head sets and apply all Hall inequalities to the induced directed bipartite relation. The producer uses a different Boolean matching routine. Both count one feasible support even when it has several witnesses.

Every preorder core is obtained from a quotient poset and positive sizes of its equivalence classes. Posets can be generated by adjoining a maximal element with every possible predecessor ideal. Canonicalization merges only complete adjacency matrices related by an explicitly tried permutation; global reversal exchanges tail and head supports and is also allowed. The four-core audit independently reconstructs the domain from upper-triangular quotient relations, block compositions, and all relabelings.

Exterior legality is checked with two identical copies of a type and with the union of all legal types. Any failed transitivity relation involves at most two exterior vertices, so these tests cover arbitrary multiplicities. For structural pruning, let H be a component of the graph formed with one clone of every legal type. Its largest matching rank over all its populations is

$$|A_H| + \nu((K \cap H) \setminus A_H),$$

where A_H is its set of active attachments. The upper bound follows by removing edges incident with A_H . Enough clones attain it by matching the attachments to distinct exterior vertices before matching the residual core. This is an all-population argument, not a finite sampling rule.

A connected bipartite preorder component with at least three vertices has no equivalence edge and no directed two-edge path on distinct vertices, since either would force a triangle. It therefore has genuine source and sink shores. Its support polynomial is the bipartite shore-support polynomial. The previously established weighted bipartite rank-at-most-four theorem applies [*Merge note*: source 16 cites here a weighted bipartite rank-four package that it embeds as a dependency; that package is batch-73 manuscript 26, which this intake prints in *matching-rank-normalization* as its source 77 (Theorem `mrn:v:rf4:thm:main`). The stronger *matching-rank-normalization*, Corollary `mrn:w:rf:cor:rankfive`, already covers every weighted bipartite graph of matching number at most five.]; an isolated two-element equivalence component contributes $1 + 2t$.

3.7 The bounded branches and the degree four theorem

All activities in the computational classification are one. The term *weighted quotient* below refers only to positive integer equivalence-class sizes, not to arbitrary role activities. The following

finite counts describe exact certificate domains, not random tests.

Branch	Core domain	Exact reduction
$a = 0$	Components of order ≤ 9	Finite degree-four check and lower-degree theorem
$a = 1$	Marked core of order ≤ 10	412622 coefficient pairs and 1237866 quadratic checks
$a = 2$	Core of order ≤ 8	89863 distinct gamma polynomials
$a = 3$	Core of order ≤ 6	6213 templates and 3437 distinct gamma polynomials
$a = 4$	Four attachment vertices	76 canonical nonbipartite templates

The complete ten-element quotient scan checks 5049656 weighted quotient cases, 686481 of actual degree four. Smaller preorders are included by padding with isolated vertices. All actual-degree-four cases satisfy (3.1).

For $a = 1$, transitivity makes the attachment a singleton source or sink. Its m pendant clones give the exact identity

$$\Gamma_{K_m}(t) = \Gamma_K(t) + mt\Gamma_{K-v}(t).$$

Each gap is a quadratic $Am^2 + Bm + C$. The certificate verifies $A, C \geq 0$ and either $B \geq 0$ or $B^2 \leq 4AC$, proving nonnegativity for all real $m \geq 0$. Independent support enumeration reproduces all 412622 coefficient pairs and the complete quadratic ledger.

For $a = 2$, the core condition $\nu(K \setminus A) \leq 2$ bounds the degree by four at every population. The three population variables have coefficient degree at most two. Exact enumeration produces 89863 distinct gamma polynomials and 179726 remaining-gap instances. Independent positivity replay covers 29215 normalized targets and all required face aliases.

For $a = 3$, 90042 legal marked specifications reduce structurally to 6213 templates and 3437 distinct gamma polynomials. Independent Hall recount verifies 188715 quota vectors and 943575 coefficients. All 204388 population faces are covered by exact rational identities, including 53722 normalized nontrivial targets.

For $a = 4$, independent enumeration gives 355 labeled four-element preorders, 4882 legal signed specifications, and 76 canonical retained templates. It checks 32516 quota vectors, 162580 coefficients, and the complete 258153-term gap data. These counts establish coverage and algebra before any positivity claim is made.

Theorem 3.9 (Rank ultra log concavity through degree four). *For every finite preorder whose unweighted support polynomial Γ_D has actual degree $d \leq 4$, the sequence*

$$\left(\gamma_k / \binom{d}{k} \right)_{k=0}^d$$

is log-concave. In particular, if $d = 4$, all three inequalities in (3.1) hold.

Proof. Degrees at most three follow from the separately verified lower-degree theorem (source 25, Theorem 2.3). Suppose $d = 4$. The first inequality follows from Theorem 1.5. The normal form is exhaustive. The $a \leq 3$ sectors satisfy all three gaps by the preceding exact certificates. For the $a = 4$ sector, the complete independently replayed ledger covers both remaining gaps for all 76 templates and every nonnegative integer population, using the structural proofs above and the exact certificate schema below. Structural pruning and product closure handle discarded specifications and disconnected preorders. Every product is normalized by its own actual degree before homogeneous Lorentzian product closure is applied. Thus no vertex-count bound or unproved change of normalization enters the conclusion. $\square$

Theorem 3.9 is computer-assisted. Its proof requires exact replay of a finite, explicitly supplied collection of identities; floating-point optimization is not a logical dependency. The final template, numbered 26, has core arcs $2 \rightarrow 0$, $3 \rightarrow 0$, $3 \rightarrow 1$, $3 \rightarrow 2$, with vertex 3 the unique outward attachment. Its certificate contains 2273 nonnegative rational square orbits. Independent expansion verifies the full unscaled last-gap polynomial, thereby covering its 1785 remaining population faces. The complete ledger has no outstanding template or face obligations.

[Merge note: template 26 has a second, smaller route for its last gap: source 23, Section 4. The original route’s certificate (`global_integer_sextic_shifted_gmp_26.json`, 29.4 MB, the 2273 square orbits above) is not shipped with this report; its provenance record and replay log are (`data/16-degree-four-provenance-...`), and the README says how to retrieve the certificate from the arrival commit. It was replayed exactly at intake.]

3.7.1 A hybrid last-gap certificate

For a four-core template, delete all internal core arcs and let F_k be the exterior-only coefficients. Each size-four support uses all four core vertices on exterior edges, so $\gamma_4 = F_4$. Write $B = \gamma_2 - F_2$ and $C = \gamma_3 - F_3$; these count disjoint support classes using core edges and are nonnegative. Then

$$3\gamma_3^2 - 8\gamma_2\gamma_4 = (3F_3^2 - 8F_2F_4) + E, \quad E = 6F_3C + 3C^2 - 8BF_4. \quad (3.5)$$

The first summand is nonnegative by Theorem 3.3, at cover order four even if the exterior-only degree is smaller. For templates 9,36,37,50 the package proves $E \geq 0$ by exact nonnegative binomial or binomial-square identities. Its degree is at most five, smaller than the full sextic gap. This is a case-specific reduction: E can be negative for other templates, and those cases use full-gap certificates instead.

3.8 Integer population certificates

Write $B_\alpha(\mathbf{n}) = \prod_i \binom{n_i}{\alpha_i}$. A global certificate has the form

$$P(\mathbf{n}) = \sum_j \frac{w_j}{|G|} \sum_{g \in G} B_{m_j}(g\mathbf{n}) \left(\sum_\alpha c_{j,\alpha} B_\alpha(g\mathbf{n}) \right)^2 + \sum_\alpha r_\alpha B_\alpha(\mathbf{n}), \quad (3.6)$$

where $w_j, r_\alpha \geq 0$ are rational and G is a specified coordinate-permutation group. Each summand is nonnegative on $\mathbb{Z}_{\geq 0}^d$. These are identities of rational polynomials, but positivity is asserted on integer populations: for instance $\binom{x}{2}$ is negative for $0 < x < 1$.

A standard-library checker expands every symmetry action and every product using

$$\binom{n}{a} \binom{n}{b} = \sum_{h=0}^{\min(a,b)} \frac{(a+b-h)!}{h!(a-h)!(b-h)!} \binom{n}{a+b-h}.$$

This counts pairs of subsets by their intersection size. Tensor products give the multivariate multiplication rule. Comparing every coefficient, including omitted zeros, proves the identity; checking all rational signs proves the inequality. Solver status and numerical tolerances are irrelevant to this final step.

For some smaller domains, split each coordinate into $n_i = 0$ or $n_i = 1 + y_i$. These finitely many faces cover all integer populations. Each resulting polynomial is certified by nonnegative monomials and nonnegative monomial multiples of exact rational squares. Unused coordinates, permutations, and positive scaling factors are explicitly checked in the face ledger.

3.8.1 Shifted binomial discovery

The successful higher-dimensional search writes a square root accompanying B_m in the basis $B_q(\mathbf{n} - m)$. When an integer population is outside $\mathbf{n} \geq m$, the multiplier vanishes. Within that orthant, shifted coordinates reveal boundary zeros. The final output removes every shift using

$$\binom{n-m}{q} = \sum_{a=0}^q \binom{-m}{q-a} \binom{n}{a},$$

and is verified in the unshifted schema (3.6).

If a target vanishes on the entire lower integer box $m + s$, $0 \leq s \leq q$, any square with multiplier B_m must vanish there. Triangular Newton interpolation then eliminates its coefficients indexed by $a \leq q$. An isolated zero imposes only its evaluation equation, not the deletion of a single basis coefficient. These restrictions aid discovery; only the final full identity is trusted.

3.8.2 Reproducing the finite verification

[Merge note: source 16's package is not shipped in its delivered layout: its files are renamed into `code/` and `data/` with the prefix `16-degree-four-`, the files larger than 2 MB (the 93.7 MB three-attachment certificates among them) stay in the arrival commit, and the two embedded dependency archives are exact copies of source 25's package and of batch-73 manuscript 26. The delivered `verify.py` first checks every one of its 260 manifest entries, so it runs only on the reconstructed package (README, "Reconstructing what is not shipped"). On Windows its path substitution writes backslashes into Python source; run it on Linux or WSL, or in a copy that uses `as_posix().`]

The accompanying directory `reproducibility/` contains the exact data, independent checkers, approved coverage inputs, and a portable driver. From that directory run

```
python verify.py --mode fast
```

This replays the fixed rational identities, gamma-to-gap reconstructions, face aliases, and complete template coverage. Its four-attachment manifest has 50 global identities: 24 middle-gap identities, 23 whole last-gap identities, and 3 identities for a core correction added to a Lorentzian baseline. They contain 15408 square orbits in total. Together with the ordinary proofs and coefficientwise-positive cases, the complete coverage check discharges all 39367 recorded outstanding face tasks and all 76 canonical templates.

For independent exhaustive kernel regeneration as well, run

```
python verify.py --mode full
```

The full mode adds the bounded-core support enumerations and Hall-based coefficient checks; its ten-element enumeration is the expensive step. Python 3.11 or later suffices for fixed certificate replay; full regeneration also requires GNU `g++` with C++17 support (the enumerators use GNU headers and 128-bit integers). Neither mode needs a numerical optimization package. The final package requires no partial-result override: a successful final coverage assertion demands all 76 templates and zero remaining tasks. A separate `--mode hashes` checks integrity only and does not establish the mathematics.

The previous degree-three and bipartite rank-four packages are included as explicit dependencies. The driver verifies their archive hashes; their own documented commands replay their earlier computational components. These are not silently re-enumerated by the new wrapper. Ordinary proof sources are supplied and pinned, but checking a source hash is likewise not a mathematical proof. The wrapper runs fresh copies of the independent checkers in a temporary

directory, removes historical positivity receipts, and derives its final coverage union from the newly replayed identities and the specified ordinary theorem approvals. The supplied README describes the small path-portability adaptations and where fresh logs and results are written.

3.9 Verification boundary and remaining research

The current article separates ordinary structural proofs from finite computer-assisted results. The role-cover theorem includes arbitrary independent role activities; the outward-star last-gap theorem is presently stated for unit activities. The exterior-only theorem has cover-order normalization, which is actual-degree normalization only at full cover rank. None is silently substituted for the general preorder theorem.

The completed degree-four ledger is included in the package. All 76 canonical four-attachment templates have both remaining gaps covered, and the lower-attachment branches and finite support kernels are independently checked. These finite identities, together with the ordinary reductions, establish Theorem 3.9 for arbitrary finite preorders of actual degree at most four. No Lean formalization or global priority claim is made here.

The most useful further questions are:

1. Replace the largest integer-population certificates, particularly the final one-source template with an internal sink-side edge, by ordinary structural proofs. Can its weighted deletion average be proved rank-ultra-log-concave directly?
2. Determine whether arbitrary independent role activities preserve rank-ULC for every degree-three preorder; a separate program proves the one-activity-per-vertex version and all unbounded branches for independent roles, but the remaining finite independent-role cases require their own coverage proof.
3. Determine whether actual-degree ULC holds for every two-by-three role cover. An unrestricted real-rootedness extension is impossible: the rank-three nonreal example, with two independent edges adjoined, already gives a nonreal degree-five product in that cover class.
4. Characterize when the exterior top-coefficient ratio remains monotone in ranks above three. General transversal matroids are not all Rayleigh, so the rank-three proof has a genuine boundary.
5. Classify equality in the degree-four inequalities and determine whether equality forces a factorization into lower-rank components.
6. Formalize the finite support kernel and the rational certificate checker in a proof assistant. This would strengthen the trust boundary without requiring formalization of the numerical discovery algorithm.
7. Find practical degree-five extensions of the bounded-core method, and identify the first true failures under actual-degree normalization rather than cover-size normalization.

[*Merge note:* status of these questions in this report. (1) Source 23 (Section 4) gives a smaller proof of the last gap of the final one-source template, by a deletion-mixture lemma, but it is still certificate-assisted (ten quartic identities) and does not prove the weighted deletion average rank-ULC directly. (2) Answered: source 22 (Part II) proves rank-ULC for every degree-three preorder with arbitrary independent role activities. (3) Open in general; settled for role covers with $P \subseteq Q$ by source 11's corollary, for independent physical cores with saturation by source 15, and by real stability for two by two role covers (source 20); sources 11, 12 and 15 state the general two-tail/three-head case as open. (4)–(6) remain open. (7) Part III proves actual-degree five for five-vertex cores with a universal-sink cloud (source 34); degree five for preorders in general is open.]

4 A second route for the last gap of template 26 (source 23)

Source 23 is *A smaller proof of the last Newton gap for template 26. A certificate assisted supplement* (6 pages as delivered; archive `template26-structural-supplement`; no author line). Its files are shipped with the prefix `23-template26-`.

[*Merge note:* source 23 proves again, by a different and smaller route, the one entry of source 16's ledger with the largest certificate: canonical template 26, last gap (Section 3). Its family and the decomposition $\Gamma = Q^+ + tR$ are those of source 16's Theorem 3.5 with the internal arc $q_2 \rightarrow q_0$ added; its Rayleigh step and its final square are source 16's Lemma 3.7 and the last display of the proof of Theorem 3.5, with an internal-edge correction. It answers source 16's first research question in part. Its dependency "the independently approved vertex-weighted theorem at actual degree at most three" was delivered as archive 31, an earlier version of source 22; the statement is the case $u_i = v_i$ of source 22's theorem, Corollary 9.14. Both routes stay: source 16's certificate for template 26 is a direct binomial-square identity; source 23's route needs ten quartic identities, that theorem and Wagner's rank-three Rayleigh theorem.]

The text below is source 23's, from its opening paragraph.

This supplement proves the last order-four Newton inequality for the integer-population family called template 26. The argument uses a support decomposition, a deletion-mixture lemma, and monotonicity of a ratio of two top coefficients. Its finite algebraic input consists of ten exact quartic identities. The proof also depends on the independently approved vertex-weighted theorem at actual degree at most three [*Merge note:* that theorem is the case $u_i = v_i$ of source 22's Theorem 9.3, recorded as Corollary 9.14.] and Wagner's rank-three Rayleigh theorem. Thus it is a smaller certificate-assisted proof, not a wholly ordinary proof.

4.1 Supports and the precise family

Let P be a finite preorder on a physical ground set V . A size- k support is an *ordered* pair (U, W) of subsets of that same set such that

$$U \cap W = \emptyset, \quad |U| = |W| = k,$$

and some bijection $U \rightarrow W$ uses off-diagonal arcs of P . Count each feasible ordered pair once, regardless of the number of matching witnesses. Distinct ordered pairs with the same union remain distinct. Set

$$\Gamma_P(t) = \sum_{k \geq 0} \gamma_k(P) t^k, \quad \gamma_0(P) = 1.$$

All activities in the template-26 conclusion are one. Diagonal preorder arcs do not contribute to supports because U and W are disjoint.

There are four core vertices q_0, q_1, q_2, p . Their off-diagonal arcs are

$$q_2 \longrightarrow q_0, \quad p \longrightarrow q_i \quad (0 \leq i \leq 2).$$

Exterior vertices are independent. An exterior vertex x may point to any subset $N(x) \subseteq \{q_0, q_1, q_2\}$ satisfying

$$q_2 \in N(x) \implies q_0 \in N(x).$$

Independently, the arc $p \rightarrow x$ may be present. In bit order q_0, q_1, q_2, p , the first three bits describe $x \rightarrow q_i$, while the fourth describes $p \rightarrow x$. The nonisolated types are exactly

$$1, 2, 3, 5, 7, 8, 9, 10, 11, 13, 15. \tag{4.1}$$

Each type has an arbitrary nonnegative integer population. Isolates may be omitted. This is the canonical template with core rows $[0, 0, 1, 7]$ and orientation mask 8.

Theorem 4.1. *For every such population, writing $\Gamma(t) = \sum_k \gamma_k t^k$, one has*

$$3\gamma_3^2 \geq 8\gamma_2\gamma_4. \quad (4.2)$$

At actual degree four, this is the last Newton inequality for $\gamma_k/\binom{4}{k}$. If the degree drops below four, the displayed inequality is automatic.

There can be at most three core heads and at most one exterior head, since only p points to exterior vertices. Thus the actual degree never exceeds four.

4.2 The support decomposition

Delete p and call the induced preorder Q . Its nonisolated exterior types are 1, 2, 3, 5, 7; let their populations be $m = (a, b, c, d, e) \in \mathbb{Z}_{\geq 0}^5$. A former type-8 vertex is now isolated. Write

$$Q(t) = 1 + At + Bt^2 + Tt^3.$$

Let H be the exterior vertices with $p \rightarrow x$, including the possible Q -isolates. For each $x \in H$, let Q_{-x} be the support polynomial after deleting that *physical vertex*, and put

$$R(t) = \sum_{x \in H} Q_{-x}(t) = \sum_{j=0}^3 R_j t^j.$$

Let Q^+ be Q with one new exterior tail adjacent to all three core/head vertices. This is still a preorder, of support degree at most three. Write $B^+ = [t^2]Q^+$ and $T^+ = [t^3]Q^+$.

Proposition 4.2. *The full support polynomial satisfies*

$$\Gamma(t) = Q^+(t) + tR(t). \quad (4.3)$$

Consequently,

$$\gamma_2 = B^+ + R_1, \quad \gamma_3 = T^+ + R_2, \quad \gamma_4 = R_3. \quad (4.4)$$

Proof. If the head set contains no exterior vertex, the support is exactly a support of Q^+ , with its new universal tail identified with p . If it contains an exterior vertex x , then x is unique: only p can match to exterior heads. The pair $p \rightarrow x$ is forced in every matching witness. Removing p and x gives precisely a support of Q_{-x} . Conversely, adjoining $p \rightarrow x$ to such a support recovers a full support. Distinct x give distinct head sets. These are bijections of supports, so no matching multiplicity is introduced. $\square$

For reference, the three coefficients of Q are the following polynomials:

$$A = 1 + a + b + 2c + 2d + 3e, \quad (4.5)$$

$$\begin{aligned} B &= ab + ac + ad + 2ae + bc + 2bd + 2be \\ &\quad + 3cd + 3ce + 3de + \binom{c}{2} + \binom{d}{2} + 3\binom{e}{2} + b + c + e, \end{aligned} \quad (4.6)$$

$$\begin{aligned} T &= b \left((d+e)(a+c) + \binom{d+e}{2} \right) \\ &\quad + a \left(cd + ce + de + \binom{e}{2} \right) + \binom{c+d+e}{3} - \binom{c}{3} - \binom{d}{3}. \end{aligned} \quad (4.7)$$

Here binomial expressions are ordinary binomial polynomials, evaluated at integer populations. The accompanying independent Hall-support checker reconstructs these formulas directly: all 56 exterior quota vectors of total at most three are enumerated, and all 42 nonzero coefficient entries agree. The quota bound follows because every selected exterior vertex needs a distinct core head. The decomposition (4.3) is also independently replayed from direct ordered disjoint support counts, rather than assumed by the algebra checker.

4.3 Reducing deletion mixtures to pairs

For $i \in \{a, b, c, d, e\}$ with $m_i > 0$, write $Q_i = Q(m - e_i)$, where e_i is the corresponding coordinate unit vector, and write $Q_* = Q(m)$. We require the following statement.

Lemma 4.3 (Deletion mixtures). *Every nonnegative real linear combination S of Q_* and the available Q_i satisfies*

$$S_2^2 \geq 3S_1S_3. \quad (4.8)$$

The populations m are nonnegative integers. In particular, (4.8) applies to R from Section 2: same-type deletions coincide, and deletion of a Q -isolate contributes Q_ .*

The two-point principle. Consider any finite list of triples (A_i, B_i, T_i) with $A_i > 0$. For a nonzero nonnegative mixture, normalize by $A = \sum_i w_i A_i$ and let $\lambda_i = w_i A_i / A$. Then

$$\left(\frac{B}{A}, \frac{T}{A}\right) = \sum_i \lambda_i \left(\frac{B_i}{A_i}, \frac{T_i}{A_i}\right), \quad \lambda_i \geq 0, \quad \sum_i \lambda_i = 1.$$

At each fixed horizontal coordinate, a highest point of this finite convex hull is a combination of at most two listed points. To see this, take a maximizing combination with smallest positive support. If it has at least three positive coefficients, a nonzero perturbation δ satisfies $\sum \delta_i = \sum \delta_i (B_i/A_i) = 0$. Both sufficiently small signs are feasible. Maximality forces $\sum \delta_i (T_i/A_i) = 0$; extend one sign until a coefficient vanishes, contradicting minimality. This also covers repeated horizontal coordinates.

It follows that every mixture satisfies $B^2 \geq 3AT$ if every two-point mixture does: all upper boundary points then lie below $y = x^2/3$, as does the entire hull. This does *not* assume that the subgraph of this convex parabola is convex. The zero mixture is immediate. In our family every $A_i \geq 1$, because the internal arc remains.

Undeleted and deleted pairs. Give one marked physical exterior vertex x of type i activity $s \in [0, 1]$, keeping every other vertex activity equal to one. A support has weight equal to the product of the activities of its physical vertices. Therefore its polynomial is exactly

$$Q^{(s)}(t) = sQ_*(t) + (1-s)Q_i(t). \quad (4.9)$$

Indeed, a support either uses x once or avoids it. The independently approved vertex-weighted actual-degree-at-most-three theorem [*Merge note*: that theorem is the case $u_i = v_i$ of source 22's Theorem 9.3, recorded as Corollary 9.14.] gives $B(s)^2 \geq 3A(s)T(s)$ when the surviving degree is three. At lower degree $T(s) = 0$, so the same displayed inequality is automatic, including $s = 0$. This is a change in one vertex's activity, not a substitution of a fractional population into (4.5)–(4.7).

For $u, v \geq 0$ with $u + v > 0$, choose $s = u/(u + v)$ and multiply (4.9) by $u + v$. Homogeneity gives the inequality for $uQ_* + vQ_i$. The same theorem supplies every endpoint gap $B_i^2 - 3A_iT_i \geq 0$. It remains to check the ten distinct-deletion pairs.

4.4 The ten exact quartic identities

Write (A_i, B_i, T_i) for the coefficients of Q_i and define

$$C_{ij}(m) = 2B_iB_j - 3(A_iT_j + A_jT_i) \quad (i < j).$$

Since both deletion types are available, write $m = x + e_i + e_j$ with $x \in \mathbb{Z}_{\geq 0}^5$. Thus the two coefficient triples come from $Q(x + e_j)$ and $Q(x + e_i)$.

Lemma 4.4 (Computer-checked quartic identities). *For each of the ten unordered pairs below, the supplied exact data give an identity*

$$C_{ij}(x + e_i + e_j) = \sum_{\ell} q_{\ell} \binom{x}{\alpha_{\ell}} f_{\ell}(x)^2 + \sum_{\beta} r_{\beta} \binom{x}{\beta}, \quad (4.10)$$

where $q_{\ell}, r_{\beta} > 0$ are rational, $\binom{x}{\alpha} = \prod_{k=1}^5 \binom{x_k}{\alpha_k}$, all exponent vectors belong to $\mathbb{Z}_{\geq 0}^5$, and each f_{ℓ} is an explicitly listed rational polynomial in the same binomial basis. Consequently $C_{ij}(m) \geq 0$ on its integer population domain.

Deleted population coordinates	Square terms	Remainder terms
a, b	15	84
a, c	21	81
a, d	18	83
a, e	22	83
b, c	24	81
b, d	19	88
b, e	21	85
c, d	18	89
c, e	19	86
d, e	19	86
Total	196	846

Certificate format and verification. The data file `distinct_deletion_cross_certificates.json` records, for each pair, its deletion indices ($0, \dots, 4$ correspond to $a, \dots, e$), the population shift, and the two sums in (4.10). A square record contains a rational weight, a multiplier exponent vector, and a list of exponent-coefficient pairs specifying f_{ℓ} . A remainder record is an exponent-coefficient pair. All ten identities are checked coefficientwise with exact rational arithmetic, together with every positivity and integer-exponent condition. Each summand is nonnegative for $x \in \mathbb{Z}_{\geq 0}^5$. Integer scope matters: ordinary binomial polynomials need not be nonnegative at arbitrary nonnegative real arguments.

The independent checker reconstructs the targets from Hall support feasibility and imports no producer code. It also derives binomial multiplication constants by counting pairs of subsets with prescribed union. Thus the finite input is an exact identity certificate, rather than a sample of population values or a floating-point optimization result.

Completion of Lemma 4.3. Write $G(S) = S_2^2 - 3S_1S_3$, so $G_i = G(Q_i) \geq 0$. For $u, v \geq 0$,

$$G(uQ_i + vQ_j) = u^2G_i + uvC_{ij} + v^2G_j \geq 0.$$

Lemma 4.4 settles all distinct-deletion pairs. Section 3 settles the undeleted/deleted pairs; identical deletion types need no extra check. The two-point principle proves (4.8), hence $R_2^2 \geq 3R_1R_3$. $\square$

4.5 The top ratio with the internal edge present

Remove $q_2 \rightarrow q_0$ from Q and call the exterior-only support polynomial F . Set $B_0 = F_2$, $T_0 = F_3$, and let L be the number of exterior vertices adjacent to q_1 . At support level,

$$B = B_0 + L, \quad T = T_0. \quad (4.11)$$

A size-two support using the internal edge must match its sole exterior tail to q_1 , because q_2 is a tail and q_0 a head. A size-three support cannot use the internal edge: it would require three core heads while q_2 is already a tail.

Adjoin one exterior vertex x of any allowed outgoing neighborhood, possibly empty. Set

$$\beta = \Delta B_0, \quad \tau = \Delta T_0, \quad \ell = \Delta L \in \{0, 1\}.$$

We prove the division-free top-ratio inequality

$$T_{\text{new}}B - TB_{\text{new}} = \tau(B_0 + L) - T_0(\beta + \ell) \geq 0. \quad (4.12)$$

Exterior-only contribution. Form the transversal matroid whose elements are the old exterior vertices, x , and three private dummies d_0, d_1, d_2 , each d_i adjacent only to q_i . Its independent sets are those matchable injectively into the three heads; the dummies ensure rank exactly three. Let f be its multiaffine basis polynomial, counting each feasible basis once. Wagner's theorem [14] gives, for distinct elements e, h ,

$$f_e f_h - f_{eh} \geq 0$$

for positive weights and hence, by continuity, for nonnegative weights. Subscripts here mean partial derivatives.

Evaluate with all old exterior weights one and the weights of x, d_0, d_1, d_2 zero. Then

$$f = T_0, \quad f_x = \tau, \quad \sum_{i=0}^2 f_{d_i} = B_0, \quad \sum_{i=0}^2 f_{x d_i} = \beta.$$

For example, a basis using d_i corresponds to an exterior two-tail support on the complementary pair of heads; that omitted head is unique. This preserves support counting even when a tail pair admits several choices of head pair. Summing the three Rayleigh inequalities for x, d_i gives

$$\tau B_0 - T_0 \beta \geq 0. \quad (4.13)$$

Contribution of the internal edge. Let K count exterior two-element supports on the fixed head pair $\{q_0, q_2\}$. Then

$$T_0 \leq LK. \quad (4.14)$$

For every feasible exterior triple choose one matching witness, and record the vertex assigned to q_1 and the remaining pair. These data determine the triple, giving an injection into the pairs counted by LK . If $\ell = 1$, adjoining x to each pair counted by K creates a distinct feasible triple, so $\tau \geq K$ and $\tau L - T_0 \ell \geq LK - T_0 \geq 0$. If $\ell = 0$, this expression is simply $\tau L \geq 0$. Adding this to (4.13) proves (4.12). An isolated new vertex gives equality. No positivity or division is needed here.

4.6 Compatibility and the final square

If $R_3 = 0$, then (4.4) gives $\gamma_4 = 0$, so (4.2) is immediate. Assume $R_3 > 0$. Some Q_{-x} has a positive cubic coefficient. Restoring a vertex preserves its supports; restricting a matching witness gives a quadratic support. Hence $T > 0$ and $B > 0$.

For $x \in H$, write $B_x = [t^2]Q_{-x}$ and $T_x = [t^3]Q_{-x}$. Apply (4.12) first when restoring x , then when adjoining the universal exterior vertex that defines Q^+ :

$$T_x B \leq B_x T, \quad TB^+ \leq BT^+.$$

Multiply by nonnegative factors to obtain

$$T_x B B^+ \leq B_x T B^+ \leq B_x B T^+.$$

Cancel only the strictly positive factor B . Summing over $x \in H$ yields

$$R_3 B^+ \leq R_2 T^+. \quad (4.15)$$

The argument covers $B_x = 0$ or $T_x = 0$ without division by either. An isolate deletion gives equality in the first comparison.

By Lemma 4.3, $R_1 R_3 \leq R_2^2/3$. Combining this with (4.4) and (4.15) proves

$$\begin{aligned} 3\gamma_3^2 - 8\gamma_2\gamma_4 &= 3(R_2 + T^+)^2 - 8(R_1 + B^+)R_3 \\ &\geq 3(R_2 + T^+)^2 - \frac{8}{3}R_2^2 - 8R_2T^+ \\ &= \frac{1}{3}(R_2 - 3T^+)^2 \geq 0. \end{aligned}$$

This completes the proof of Theorem 4.1.

Scope and reproducibility

This supplement proves the last gap. Together with the separately approved middle-gap proof and general first-gap bound, it gives the smaller proof for this family. The original degree-four theorem and frozen certificate archive are unchanged. There is no claim of a general weighted degree-four theorem, fractional-population theorem, real-rootedness, or novelty.

[*Merge note:* the middle-gap proof meant here is source 16's certificate for template 26, gap two (shipped as `data/16-degree-four-a4-gram_za_26.json`), and the first-gap bound is Theorem 1.5. The embedded `weighted-degree-three-package.zip` is an exact copy of archive 31, superseded by source 22, and is not shipped; because source 23's `verify.py` checks it against its checksum list, the delivered wrapper runs only on the reconstructed package (README). The same Windows path problem as in source 16's wrapper applies.]

The tradeoff is explicit: ten quartic certificates replace a much larger direct last-gap certificate, using the already approved vertex-weighted degree-three theorem. The supplement includes the certificates, independent exact checkers, approval receipts, and `weighted-degree-three-package.zip`. From the extracted supplement root, run `python verify.py` for the two local independent replays and the pinned dependency archive. Run `python verify.py --verify-dependency` to extract and replay the full portable verifier of that dependency as well. The local replays check the exact quartic certificates and the support decomposition coefficientwise. The full mathematical assembly has independent approval, including the convex-hull argument, degree-drop endpoints, internal-edge correction, zero cases, and final square.

Part II

Independent role activities and small covers

This Part allows independent nonnegative tail and head activities u_i, v_i (Definition 1.1) on arbitrary loopless directed relations, transitive or not. Five sources contribute.

- Section 5 (source 20): if every arc has its tail in a set P or its head in a set Q with $|P|, |Q| \leq 2$ (overlap allowed), the signed monomer polynomial is real stable, so Γ_D is real-rooted.

- Section 6 (source 12): coefficientwise Rayleigh inequalities in the core monomers when the physical graph is bipartite with a three-vertex shore (computer-assisted boundary inequality).
- Section 7 (source 11): rank-ULC whenever the underlying graph has a vertex cover of at most three vertices, internal core arcs allowed; it uses source 12.
- Section 8 (source 15): a Lorentzian theorem for bipartite physical graphs, and real stability for pendant uncovered heads over a stable transversal seed.
- Section 9 (source 22): rank-ULC for every finite preorder of actual degree at most three under independent role activities, by a Gallai–Edmonds reduction, ordinary structural theorems, a complete ledger of role certificates and half-plane-property side matroids.

Several statements of source 22 are special cases of statements of sources 11, 15 and 20, and source 25’s theorem (Part I) is the unit case of source 22’s; these are recorded where they occur. In the bipartite case, with every vertex in one role, these results specialize to theorems of *preorder-root-polytopes* and *matching-rank-normalization*: two-shore ultra-log-concavity (*preorder-root-polytopes*, `lor:thm:weightedULC`), weighted matching rank three and five (*preorder-root-polytopes*, `lc:cor:rankthree`; *matching-rank-normalization*, `mrn:w:rf:cor:rankfive`), scalar rank-three Rayleigh (*matching-rank-normalization*, `mrn:s:tv:eq:rayleigh`) and stability under a two by two cover (*matching-rank-normalization*, `mrn:w:sb:thm:main`). The directed, overlapping and coefficientwise forms are new.

5 Two by two role covers (source 20)

Source 20 is *Real stable monomers under a two by two role cover* (9 pages as delivered; archive `role-cover-package`). Its files are shipped with the prefix `20-role-cover-`. Source 16’s Section 3 and Appendix A are this article, with the differences printed below as “Second route (source 16)”; exact copies of the whole package were also embedded in source 22 and archive 31.

Abstract of source 20, as delivered. We count ordered disjoint tail and head supports of directed matchings, with every feasible support counted once. For any finite loopless directed relation whose bipartite role graph has a vertex cover comprising at most two tail copies and at most two head copies, we prove real stability of the signed physical monomer polynomial. The selected physical sets may overlap, and arbitrary nonnegative independent tail and head activities are allowed. Consequently the support polynomial has only negative real roots and satisfies Newton’s inequalities at its actual surviving degree. The proof treats every subset of a two-source, two-sink core, using rank-two stable quadratics, stability-preserving operators, and explicit all-real Rayleigh square identities, before merging role copies to enforce physical disjointness. We also obtain a rank-two matroid free-product corollary and an elementary root transfer to gamma expansions. These are ordinary proofs; the accompanying exact algebra and finite Hall-support checks are supplementary. No proof-assistant formalization or literature-wide priority claim is made.

5.1 Support polynomials and normalization

Definition 5.1. A feasible support of size k is an ordered pair (S, T) of disjoint subsets of V , each of cardinality k , for which there exists a bijection $\pi : S \rightarrow T$ with $s \rightarrow \pi(s)$ for every $s \in S$. Write

$$\gamma_k(D) = \#\{(S, T) : (S, T) \text{ is feasible of size } k\}, \quad \Gamma_D(t) = \sum_{k \geq 0} \gamma_k(D) t^k.$$

The empty support contributes $\gamma_0 = 1$. A support contributes once, regardless of the number of witnessing bijections.

The same-ground-set disjointness is essential. Passing to independent tail and head copies without subsequently excluding their simultaneous use changes this polynomial. Counting individual matchings also changes it.

For nonnegative tail activities u_i and head activities v_i , replace the contribution of (S, T) by

$$u^S v^T = \prod_{i \in S} u_i \prod_{j \in T} v_j.$$

We use the same notation when the weights are specified. A vertex may have different activities in its two roles. With zero activities, the polynomial degree can decrease.

[Merge note: source 20's Lemma "degree" is Lemma 1.3, printed once in Section 1.3 with this proof.]

Definition 5.2. A polynomial $g(t) = \sum_{k=0}^d g_k t^k$ is *rank-ultra-log-concave*, or rank-ULC, if its coefficients are nonnegative with no internal zeros and

$$\left(\frac{g_k}{\binom{d}{k}} \right)^2 \geq \frac{g_{k-1}}{\binom{d}{k-1}} \frac{g_{k+1}}{\binom{d}{k+1}} \quad (1 \leq k < d),$$

where d is the actual degree.

Our main theorem concerns directed relations satisfying the two by two role-cover condition in Theorem 5.10; transitivity is not required. Throughout, the monomer variable of a vertex records that the vertex is unused. Real-rootedness is always interpreted at the degree remaining after zero activities are imposed.

5.2 Real-rootedness for a complete balanced core

Let $P = \{p_1, p_2\}$, $Q = \{q_1, q_2\}$, and let I be an independent exterior set. Include all four arcs $P \rightarrow Q$. Each $x \in I$ may have arbitrary incoming neighbors in P and outgoing neighbors in Q , including an empty neighbor set on either side. There are no other arcs. This directed family includes every nonnegative integer population of all fifteen nonempty exterior types.

A real polynomial is *real stable* if it is nonzero whenever every variable lies in the open upper half-plane $\mathbb{H}$. We use the finite-degree stability-preserver theorem of Borcea and Brändén [5]; Wagner's survey gives the same plus-sign algebraic-symbol convention [15, Theorem 5.2].

Theorem 5.3. *For the complete balanced core described above, with arbitrary nonnegative independent tail and head activities, the signed multivariate monomer polynomial*

$$\mu_D(\mathbf{z}) = \sum_{(S,T) \text{ feasible}} (-1)^{|S|} u^S v^T \prod_{i \notin S \cup T} z_i$$

is real stable. Consequently every root of Γ_D is negative real, and Γ_D is rank-ULC at its actual degree. No transitivity assumption is required.

5.2.1 A stable coupling of two rank-two matroids

A loopless rank-two matroid has parallel classes with variable sums $r_1, \dots, r_j$. Its basis polynomial and element sum are $B = e_2(r_1, \dots, r_j)$ and $L = \sum_i r_i$.

Lemma 5.4. *The polynomial $f(s) = s^2 + \sqrt{2}Ls + B$ is real stable.*

Proof. The identity

$$f(s) = (s + L/\sqrt{2})^2 - \frac{1}{2} \sum_i r_i^2$$

shows that its quadratic form in $s, r_1, \dots, r_j$ has exactly one positive direction. It has nonnegative coefficients and is positive on the positive orthant. For an alleged zero at $a + ib$ with b strictly positive, its imaginary part says that a and b are orthogonal for the form. Since $f(b) > 0$, the orthogonal complement of b is nonpositive. Thus $f(a) \leq 0$, contradicting the real-part equation $f(a) = f(b)$. Replacing each r_i by its parallel-class sum preserves stability. $\square$

Apply the lemma to two matroids on disjoint variable sets. On auxiliary monomials of separate degrees at most two define the coefficientwise linear operator

$$\mathcal{T}(1) = 1, \quad \mathcal{T}(st) = -\frac{1}{2}, \quad \mathcal{T}(s^2t^2) = 1,$$

and let it vanish on the other six auxiliary monomials. Its symbol is

$$\mathcal{T}[(s+a)^2(t+b)^2] = (ab-1)^2.$$

This symbol is stable: $ab = 1$ is impossible for $a, b \in \mathbb{H}$, since $1/a$ lies in the lower half-plane. The full symbol, including all untouched variables, acquires only stable factors of the form $(z+w)^\kappa$. Therefore

$$B_P B_Q - L_P L_Q + 1 \tag{5.1}$$

is real stable. The constant term shows that it is not identically zero.

Second route (source 16): elementary operators. Source 16 does not invoke the stability-preserver theorem. It writes: “We use the standard stability closures under products, diagonalization, differentiation, nonnegative linear substitutions, and real specialization (with zero outputs allowed) [15]. The special contraction operators needed here have elementary realizations; they also fit the general stability-preserver framework [5].” It then proves the coupling as follows.

Lemma 5.5 (Elementary Asano contraction; source 16). *For a stable polynomial affine in each of a, b , write $f = abA + aB + bC + D$. The coefficientwise operation $f \mapsto D - A$ preserves stability or gives zero. So does $f \mapsto f - \partial_a \partial_b f$.*

Proof. For $t \in \mathbb{H}$, both t and $-1/t$ lie in $\mathbb{H}$. Therefore

$$tf(t, -1/t) = Bt^2 + (D - A)t - C$$

is stable in t and all untouched variables. Differentiate in t and specialize $t = 0$ to obtain $D - A$. The standard boundary-specialization closure permits the zero polynomial. For the second assertion, first replace a, b by $a + x, b + y$, which preserves stability, and contract the fresh variables x, y . Their constant coefficient is $f(a, b)$ and their product coefficient is $\partial_a \partial_b f$. $\square$

For each rank-two side use the polarized auxiliary quadratic

$$F_P(s_1, s_2) = s_1 s_2 + \frac{L_P}{\sqrt{2}}(s_1 + s_2) + B_P.$$

It is stable by the same one-positive-direction argument as Lemma 5.4: it equals the original auxiliary quadratic at $(s_1 + s_2)/2$, minus $(s_1 - s_2)^2/4$. The extra independent direction is negative, and every coefficient of the expanded polynomial is nonnegative. Construct $F_Q(t_1, t_2)$ similarly.

Apply Lemma 5.5 to the stable product $F_P F_Q$, first in (s_1, t_1) and then in (s_2, t_2) . With $a = L_P/\sqrt{2}$ and $b = L_Q/\sqrt{2}$, the result is $B_P B_Q - 2ab + 1$. Consequently (5.1) is real stable. Its constant coefficient one rules out a zero output at either contraction stage. This proves the coupling without invoking the general classification of stability-preserving linear operators.

Corollary 5.6. *If M and N are loopless rank-two matroids on disjoint ground sets, then the basis polynomial of the free product $M \square N^*$ is real stable.*

Proof. Write the ground sets as S, T , with variables $\mathbf{x}, \mathbf{y}$. By the basis characterization of the free product [8, Proposition 1 and the following paragraph], its bases are $I \cup (T \setminus K)$, where I is independent in M , K is independent in N , and $|I| = |K| \leq 2$. Indeed, $T \setminus K$ spans N^* exactly when K is independent in N , and the total basis size then forces the cardinalities to agree. Consequently its basis polynomial is

$$\left(\prod_{j \in T} y_j \right) \left[1 + L_M(\mathbf{x})L_N(\mathbf{y}^{-1}) + B_M(\mathbf{x})B_N(\mathbf{y}^{-1}) \right].$$

This is (5.1) after substituting $-1/y_j$ for each N -variable and multiplying by $\prod_{j \in T} y_j$. Multi-affinity cancels the denominators. Since $-1/y_j \in \mathbb{H}$ for $y_j \in \mathbb{H}$, the resulting polynomial is real stable. $\square$

5.2.2 The two-copy identity

Temporarily split each exterior vertex into a head copy, adjacent from P , and a tail copy, adjacent to Q . Keep isolated copies. For each two-center side use its rank-two transversal matroid on the nonisolated copies and two private core dummies. Isolated copies contribute only their factors to Z and are omitted from both B and L . The private-neighborhood copies and the corresponding dummy form a parallel class; each common-neighborhood copy forms its own class.

Give a core dummy its monomer variable z_p or z_q , and give an active exterior copy the value $-1/z_x$. These substitutions preserve the upper half-plane. Let Z be the product of all exterior-copy monomer variables. The unweighted signed monomer polynomial of the split relation is

$$\mu_{\text{copy}}(\mathbf{z}) = Z(B_P B_Q - L_P L_Q + 1). \quad (5.2)$$

The possible denominators cancel because each exterior-copy variable is multiaffine before substitution.

Here is the support-count explanation. The number of core arcs in a witnessing matching is determined by its endpoint support. Zero core arcs gives $ZB_P B_Q$. One core arc gives $-ZL_P L_Q$: on each side, the remaining core monomer can be either dummy, while an exterior singleton used together with both core vertices is counted once, not by its neighbor degree. Two core arcs use all four core vertices and no exterior vertex, contributing Z once. Although the complete core admits two perfect matchings, it has only one such endpoint support. Completeness of $P \rightarrow Q$ makes the separate side conditions sufficient.

Formula (5.2) and (5.1) prove split monomer stability. Positive activities w_i on split vertices are introduced by

$$\mu_w(\mathbf{z}) = \left(\prod_i w_i \right) \mu_{\text{copy}}((z_i/w_i)_i).$$

Assign the original tail activity to a tail copy, the head activity to a head copy, u_p to a source core vertex, and v_q to a sink core vertex.

5.2.3 Merging the physical vertices

For the two monomer variables a, b of one physical exterior vertex apply

$$ab \mapsto z, \quad a \mapsto 1, \quad b \mapsto 1, \quad 1 \mapsto 0. \quad (5.3)$$

Both unused copies become one unused physical vertex. Exactly one used copy becomes a used vertex. Two used copies are discarded. Contributions for the two different one-used roles add, as the support definition requires. The algebraic symbol is $\mathcal{M}[(a+r)(b+s)] = z+r+s$, which is stable. Including untouched variables only adds stable factors. Thus every merge preserves stability. After all merges we obtain precisely μ_D .

Second route (source 16): the merge by differentiation. There is also an elementary realization of this operator. For $f = abA + aB + bC + D$, with coefficients independent of a, b , diagonalization gives $f(t, t) = t^2A + t(B+C) + D$. Differentiate in t and substitute $t = z/2$ to obtain $zA + B + C$, exactly (5.3). Diagonalization, differentiation, and positive scaling preserve real stability or give zero. Thus this particular merge needs only these standard closure properties, and the empty-support monomial excludes zero. (Source 15's Lemma 8.4 gives the same elementary realization; source 22 uses the directional-derivative form.)

Zero activities follow by continuity from positive ones. The empty-support monomial has coefficient one throughout, so the limit is nonzero. Finally, with N physical vertices,

$$\mu_D(s, \dots, s) = \sum_k (-1)^k \gamma_k s^{N-2k} = s^N \Gamma_D(-s^{-2}).$$

The left side has only real zeros. A nonreal or positive zero of Γ_D would produce a nonreal solution of $s^2 = -1/\rho$ and hence a nonreal zero of this monomer polynomial. Since $\Gamma_D(0) = 1$, all its roots are negative. Newton's inequalities finish the proof of Theorem 5.3.

5.2.4 A two-edge core extension

The same physical-copy construction also works when the core has only the two arcs $p_2 \rightarrow q_1, p_2 \rightarrow q_2$.

Theorem 5.7. *For this two-edge core, arbitrary exterior neighborhoods and arbitrary nonnegative independent role activities again give a real-stable signed monomer polynomial. The support polynomial is therefore real-rooted and rank-ULC at its actual degree.*

Proof. With the same rank-two side matroids, put $A_P = \partial_{d_{p_2}} B_P$. This is the dummy d_{p_1} plus the sum of all exterior head-copy variables adjacent to p_1 , including common-neighbor copies once. The split kernel is

$$H = B_P B_Q - A_P L_Q.$$

At most one core edge can occur. If it does, its tail is forced to be p_2 ; the remaining P -side support is exactly A_P , and the remaining Q -side support is exactly L_Q . This proves the identity at support level.

For the parallel-class sums of the Q matroid, $B_Q + sL_Q = e_2(r_1, \dots, r_j, s)$ is stable. Fix its original arguments in $\mathbb{H}$. Both B_Q and L_Q are nonzero, and its root $-B_Q/L_Q$ cannot be strictly upper. Hence $-L_Q/B_Q$ has nonnegative imaginary part. Since B_P is affine in d_{p_2} ,

$$H = B_Q B_P(d_{p_2} - L_Q/B_Q),$$

with all other arguments unchanged. The shifted dummy remains strictly upper, so this expression is nonzero. The reciprocal substitutions, activity scalings, physical-copy merges, and monomer-to-support root transfer from Theorem 5.3 now apply unchanged. Zero activities follow by the same nonzero limit argument. $\square$

The proof does not assume that adding a polynomial with positive coefficients preserves stability. [Merge note: source 16 adds: "This includes canonical template 6 of the degree-four classification" (Section 3).]

5.2.5 A core with one missing edge

Theorem 5.8. *Keep the three core arcs $p_1 \rightarrow q_1, p_1 \rightarrow q_2, p_2 \rightarrow q_1$, and omit $p_2 \rightarrow q_2$. With arbitrary exterior neighborhoods and nonnegative independent role activities, the signed monomer polynomial is real stable and the support polynomial is real-rooted at its actual degree.*

The split kernel is

$$H = B_P B_Q - L_P L_Q + 1 + r_P r_Q, \quad (5.4)$$

where r_P is the parallel-class sum containing dummy p_1 and the head copies private to p_1 , and r_Q is defined similarly using q_1 . A one-core-edge support is lost from the complete core exactly when its residual source and sink are forced to be p_2, q_2 . These forced choices contribute $r_P r_Q$; their former signed contribution was negative, so the correction is positive. Common-neighbor copies do not force that residual choice. The two-core-edge endpoint support survives through the off-diagonal matching, and contributes one.

We prove the stability of (5.4) for arbitrary loopless rank-two matroids with a distinguished class on each side. Write their independent class variables as $p_0, \dots, p_a$ and $q_0, \dots, q_b$, with $r = p_0, s = q_0$, and set

$$F = e_2(\mathbf{p}), \quad G = e_2(\mathbf{q}), \quad L = \sum p_i, \quad M = \sum q_j, \quad H = FG - LM + 1 + rs.$$

For a real multiaffine polynomial the all-real Rayleigh criterion says that stability is equivalent to nonnegativity of every $\Delta_{xy} = (\partial_x H)(\partial_y H) - H \partial_x \partial_y H$ at every real assignment [7, Theorem 5.6]; see also [15, Theorem 3.1]. Testing only positive assignments would not suffice. Here five identities cover every pair type, up to exchange of sides and permutation of nondistinguished classes.

One distinguished variable on the same side. For $x = p_0, y = p_i$ with $i > 0$, put $A = \sum_{k \neq 0, i} p_k, U = \sum_{k \neq 0, i} p_k^2, B = \sum_{l > 0} q_l, V = \sum_{l > 0} q_l^2$. Then

$$2\Delta_{xy} = (GA - B)^2 + G^2 U + V.$$

Two nondistinguished variables on the same side. For $x = p_i, y = p_j$ with distinct $i, j > 0$, put $A = \sum_{k \neq i, j} p_k, U = \sum_{k \neq 0, i, j} p_k^2, V = \sum_{l > 0} q_l^2$. Then

$$2\Delta_{xy} = (GA - M)^2 + (Gr - s)^2 + G^2 U + V.$$

Both distinguished variables on opposite sides. For $x = p_0, y = q_0$, put $A = \sum_{k > 0} p_k, U = \sum_{k > 0} p_k^2, B = \sum_{l > 0} q_l, V = \sum_{l > 0} q_l^2$. Then

$$2\Delta_{xy} = A^2 V + B^2 U.$$

One distinguished variable on opposite sides. For $x = p_0, y = q_j$ with $j > 0$, put $A = \sum_{k > 0} p_k, U = \sum_{k > 0} p_k^2, B = \sum_{l \neq j} q_l, V = \sum_{l \neq 0, j} q_l^2$. Here B includes s . Then

$$4\Delta_{xy} = [A(B + s) - 2]^2 + A^2 V + U[(B - s)^2 + V].$$

Neither variable distinguished on opposite sides. For $x = p_i, y = q_j$ with $i, j > 0$, put $A = \sum_{k \neq i} p_k, U = \sum_{k \neq 0, i} p_k^2, B = \sum_{l \neq j} q_l, V = \sum_{l \neq 0, j} q_l^2$. Thus A, B include r, s . Then

$$4\Delta_{xy} = (AB - rs - 2)^2 + (As - rB)^2 + U(B^2 + s^2) + V(A^2 + r^2) + UV.$$

Each identity follows by expansion and $2e_2(\mathbf{z}) = (\sum z_i)^2 - \sum z_i^2$. Every expression is nonnegative for all real variables: U, V are sums of squares, so their displayed products are sums of squares as well. Empty sums are zero. The constant term one makes H nonzero. The Rayleigh criterion proves stability, and substitution of parallel-class sums proves it in the individual matroid variables. The reciprocal, activity, and merge steps of Theorem 5.3 complete the proof of Theorem 5.8. [Merge note: source 16 adds: “This covers canonical template 21.”]

Remark 5.9 (Interpolation). The kernels $H_\theta = FG - LM + 1 + \theta rs$ are stable for $0 \leq \theta \leq 1$. In the independent class variables, every Rayleigh difference of the monomial rs is zero. Thus $\Delta_{xy}(H_\theta)$ is the convex combination of the two endpoint differences. The same linear monomer and copy-merge transformations show that the corresponding interpolated support polynomials remain real-rooted. This is a statement about this particular interpolation, not a general convexity claim for stable polynomials.

5.2.6 Arbitrary cross-core patterns and role covers

For two disjoint core edges $p_1 \rightarrow q_1, p_2 \rightarrow q_2$, the split kernel is

$$H = B_P B_Q - L_P L_Q + 1 + r_{P1} r_{Q2} + r_{P2} r_{Q1},$$

where each r_{Pi}, r_{Qj} is the private-plus-dummy parallel-class sum. Each omitted cross edge removes its uniquely forced residual one-core-edge supports. Those two lost sets are disjoint, and the diagonal perfect support survives. Section 5.4 proves this kernel stable by seven all-real Rayleigh identities. The weighted and physical-merge conclusions follow as before.

The empty core gives the stable product $B_P B_Q$. A single core edge gives

$$(1 - \partial_{d_{p_i}} \partial_{d_{q_j}})(B_P B_Q).$$

On polynomials affine in these two dummy variables, the operator’s symbol is $(a+x)(b+y) - 1$, which is stable because two upper-half-plane numbers cannot have product one. These cases, the two-edge star and its reversal, the disjoint pair, the one-missing-edge case, and the complete core exhaust all sixteen subsets of the four possible core arcs. Thus every such subset has a stable split monomer polynomial.

Second route (source 16). Lemma 5.5, applied after translation by fresh variables, proves that this operator preserves stability or gives zero. The all-dummy monomial excludes zero.

Theorem 5.10 (Two by two role cover). *Let D be any finite loopless directed relation on V . Suppose there are sets $P, Q \subseteq V$, each of size at most two, such that every arc $i \rightarrow j$ has $i \in P$ or $j \in Q$. The two sets may overlap. Then for arbitrary nonnegative independent tail and head activities, the signed physical monomer polynomial is real stable. Its support polynomial has only negative real roots and is rank-ULC at its actual degree.*

Proof. Replace every physical vertex by distinct tail and head copies. The selected tail copies of P and head copies of Q form a bipartite vertex cover. They are disjoint as copies even when their original physical sets overlap. All other copies are exterior; no edge joins two exterior copies. The core is an arbitrary subset of the four possible cross edges, so the preceding classification proves split monomer stability. If a cover side is smaller than two, pad it with isolated auxiliary vertices. Removing their monomer factors preserves nonvanishing in the upper half-plane.

Give every tail copy its tail activity and every head copy its head activity. Merge the two copies of every physical vertex by (5.3), including core/core and core/exterior pairs. An original disjoint support lifts uniquely to its tail and head copies. A split support survives precisely when

it does not use both copies of any physical vertex. Matching feasibility and its product weight are unchanged. Distinct original role assignments can have the same unused-vertex monomial and their coefficients correctly add. This is an exact support bijection, not a count of matching witnesses. Each merge preserves stability, and the empty-support monomial excludes the zero polynomial. The root transfer and zero-activity limit finish the proof. $\square$

The criterion permits arbitrary arcs inside each of two disjoint two-vertex core sides, as well as arbitrary $P \rightarrow Q$ core arcs and $P \rightarrow I \rightarrow Q$ exterior attachments. It does not permit an arbitrary backward $Q \rightarrow P$ arc. It also shows why an actual degree below four causes no normalization problem: real-rootedness applies directly to the surviving polynomial.

[Merge note: for a bipartite graph oriented from one shore to the other, every vertex has one role, a cover with two vertices on each shore is a role cover with $P \cap Q = \emptyset$, vertex activities are role activities, and Γ_D is the matching-support polynomial. That case is *matching-rank-normalization*, Theorem `mrn:w:sb:thm:main`, proved there by a different route (vertex sums, expected determinants and Grace–Walsh–Szegő polarization); batch-73 manuscript 26, printed in *matching-rank-normalization* as its source 77 (Theorem `mrn:v:rf4:thm:mixed`), proves the same undirected statement with the weaker conclusion of real-rootedness by a third route. Theorem 5.10 generalizes it to directed relations, overlapping P and Q , arcs inside P and inside Q , and independent role activities. Source 22 uses it in its role-cover corollary and in the three-attachment branch of its global proof (Section 9); source 16’s degree-four ledger uses its core cases for templates 6 and 21. The free-product corollary has no counterpart in *preorder-root-polytopes* or *matching-rank-normalization*.]

Corollary 5.11 (Transfer to the gamma expansion). *Under Theorem 5.10, let $d = \deg \Gamma_D$ and choose any integer $N \geq 2d$. Put*

$$h_D(t) = \sum_k \gamma_k t^k (1+t)^{N-2k}.$$

Then h_D has only negative real roots. The choice $N = |V|$ is permitted. The same conclusion applies to any polynomial independently identified with this gamma expansion; no geometric convention is assumed here.

Proof. Write $\Gamma_D(z) = \prod_{i=1}^d (1 + \lambda_i z)$ with $\lambda_i > 0$. Since $2d \leq N$,

$$h_D(t) = (1+t)^{N-2d} \prod_{i=1}^d [t^2 + (2 + \lambda_i)t + 1].$$

Each quadratic has discriminant $\lambda_i(\lambda_i + 4) > 0$, negative root sum, and positive root product, so both roots are negative. The remaining roots are -1 . This is the standard elementary root transfer, not a separate novelty claim. $\square$

[Merge note: source 16 states this corollary with $N = |V|$ only and ends: “In particular this applies to a preorder h -polynomial when its gamma expansion is the support expansion identified in the source report” (*preorder-root-polytopes*, Theorem `gam:thm:main`).]

5.3 Proof and verification boundary

The arguments above are ordinary mathematical proofs. The all-real Rayleigh identities hold for any number of parallel classes; their validity is not inferred from a bounded computation. The finite-degree stability-preserver theorem, the all-real multiaffine Rayleigh criterion, and elementary rank-two matroid facts are the external mathematical dependencies.

The supplementary archive contains the approved proof notes and scoped review receipts, two independent exact symbolic checkers for the twelve aggregate-variable identities, and four independent Hall-based support checkers. Each Hall checker compares weighted multivariate monomer coefficients for all 4845 exterior-type multisets through four vertices, including isolated copies and zero activities: 14535 activity-weighted identities per core pattern. These finite checks test the combinatorial formulas; they are not a substitute for the arbitrary-population proofs. The archive includes no numerical optimization or binary executable as a logical dependency. Its portable README describes how to rebuild the article and rerun the exact checks.

[Merge note: source 20's files are shipped with the prefix `20-role-cover-` (proof notes at the top of the report directory, checkers in `code/`, receipts and logs in `data/`). At intake its `verify.py` passed on a copy of the delivered package in 73 s; with GCC 16 its four C++ checkers need `-include cassert`, because they call `assert` without including `<cassert>`.]

5.4 Appendix of source 20: Rayleigh identities for two disjoint core edges

[Merge note: the typographical slip “labels0,1” of source 20 is corrected as in source 16.]

Use independent parallel-class variables $p_0, p_1, p_2, \dots$ and $q_0, q_1, q_2, \dots$. Label the first two Q classes so that the two correction terms are $p_0q_0 + p_1q_1$. Put

$$F = e_2(\mathbf{p}), \quad G = e_2(\mathbf{q}), \quad L = \sum p_i, \quad M = \sum q_j, \quad K = p_0q_0 + p_1q_1, \quad H = FG - LM + 1 + K.$$

The following seven cases cover every pair under side interchange, simultaneous exchange of the partnered labels 0, 1, and permutations of the remaining classes. Every displayed square sum is nonnegative on all real assignments. The identities follow from the elementary symmetric identity used above, with no bound on the number of classes.

Same side with both distinguished. For $x = p_0, y = p_1$, take $A = \sum_{k \geq 2} p_k$, $U = \sum_{k \geq 2} p_k^2$, $B = \sum_{l \geq 2} q_l$, $V = \sum_{l \geq 2} q_l^2$. Then

$$2\Delta_{xy} = (GA - B)^2 + G^2U + V.$$

Same side with one distinguished. For $x = p_0, y = p_i$, $i \geq 2$, take $A = \sum_{k \neq 0, i} p_k$, $U = \sum_{k \geq 2, k \neq i} p_k^2$, $V = \sum_{l \geq 2} q_l^2$. Then

$$2\Delta_{xy} = [GA - (M - q_0)]^2 + (Gp_1 - q_1)^2 + G^2U + V.$$

Same side with neither distinguished. For $x = p_i, y = p_j$, $i, j \geq 2$ and $i \neq j$, take $A = \sum_{k \neq i, j} p_k$, $U = \sum_{k \geq 2, k \neq i, j} p_k^2$, $V = \sum_{l \geq 2} q_l^2$. Then

$$2\Delta_{xy} = (GA - M)^2 + (Gp_0 - q_0)^2 + (Gp_1 - q_1)^2 + G^2U + V.$$

Opposite sides with partnered distinguished variables. For $x = p_0, y = q_0$, take $A = \sum_{k \neq 0} p_k$, $U = \sum_{k \geq 2} p_k^2$, $B = \sum_{l \neq 0} q_l$, $V = \sum_{l \geq 2} q_l^2$. Then

$$2\Delta_{xy} = (Aq_1 - Bp_1)^2 + A^2V + B^2U.$$

Opposite sides with unpartnered distinguished variables. For $x = p_0, y = q_1$, let $u = p_1$, $v = q_0$, $A = \sum_{k \neq 0} p_k$, $U = \sum_{k \geq 2} p_k^2$, $B = \sum_{l \neq 1} q_l$, $V = \sum_{l \geq 2} q_l^2$. Then

$$4\Delta_{xy} = (AB + uB + vA - uv - 2)^2 + U(B - v)^2 + V(A - u)^2 + UV.$$

Opposite sides with one distinguished variable. For $x = p_0, y = q_j, j \geq 2$, let $r = p_1, s = q_1, v = q_0, A = \sum_{k \neq 0} p_k, U = \sum_{k \geq 2} p_k^2, B = \sum_{l \neq j} q_l, V = \sum_{l \geq 2, l \neq j} q_l^2$. Then

$$4\Delta_{xy} = [A(B + v) - rs - 2]^2 + [As - r(B - v)]^2 \\ + V(A^2 + r^2) + U[(B - v)^2 + s^2] + UV.$$

Opposite sides with neither distinguished. For $x = p_i, y = q_j, i, j \geq 2$, take $A = \sum_{k \neq i} p_k, U = \sum_{k \geq 2, k \neq i} p_k^2, B = \sum_{l \neq j} q_l, V = \sum_{l \geq 2, l \neq j} q_l^2, R_2 = p_0^2 + p_1^2, S_2 = q_0^2 + q_1^2$. Then

$$4\Delta_{xy} = (AB - K - 2)^2 + (Aq_0 - Bp_0)^2 + (Aq_1 - Bp_1)^2 \\ + (p_0q_1 - p_1q_0)^2 + U(B^2 + S_2) + V(A^2 + R_2) + UV.$$

The all-real Rayleigh criterion now proves stability. Parallel-class sum substitution transfers it to individual variables. These explicit identities, rather than a finite numerical test, establish the arbitrary-class statement.

6 Coefficientwise Rayleigh inequalities for a three-vertex shore (source 12)

Source 12 is *Coefficientwise Rayleigh inequalities for a three vertex shore* (6 pages as delivered; archive `three-core-rayleigh-result`); source 11 embedded an exact copy of it as its dependency. Its files are shipped with the prefix `12-three-core-rayleigh-`.

Abstract of source 12, as delivered. For a directed relation whose physical graph is bipartite with a three-vertex shore, we prove coefficientwise Rayleigh inequalities for the polynomial of feasible ordered endpoint supports. Opposite arcs and independent tail/head activities are allowed, and each feasible support is counted once. The boundary inequality reduces to 48 role-monomial types and 17,376 finite directed graphs; two independent exact checkers verify every coefficient. A private-head argument then proves coefficientwise positivity of the full Rayleigh differences in the three core monomers. Consequently, use of any two core vertices is negatively correlated in the endpoint-support measure under all nonnegative role activities and positive core monomer activities. We give a four-core example, already admitting a two-tail/three-head role cover, where coefficientwise positivity fails despite an exact positive two-square formula. The boundary theorem is computer-assisted; the corollaries and obstruction are ordinary arguments. No full-stability, internal-core extension, proof-assistant formalization, or priority claim is made.

6.1 Supports and the main inequality

Let D be a finite loopless directed relation on the physical vertex set $V = C \sqcup I$, where $C = \{0, 1, 2\}$, and assume every arc crosses this partition. Both orientations of an edge may be present. Work in the polynomial ring in independent commuting tail and head activities u_v, v_v for $v \in V$.

A *feasible support* of size k is an ordered pair (S, T) of disjoint physical vertex sets of size k admitting a directed perfect matching from S to T . Its weight is $\text{wt}(S, T) = u^S v^T$. It contributes once, regardless of how many bijections witness feasibility. Since all arcs cross the physical bipartition, every size- k support uses exactly k vertices in each shore.

Define a_i as the sum of weights of size-one supports with used core set $\{i\}$, define $b_{ij} = b_{ji}$ similarly for size-two supports with used core set $\{i, j\}$, and let c be the size-three coefficient.

The core activities are included in these polynomials. The polynomial recording unused core vertices is

$$F(\mathbf{z}) = z_0 z_1 z_2 + \sum_{i \in C} a_i z_j z_k + \sum_{i < j} b_{ij} z_k + c, \quad (6.1)$$

where in each summand the omitted indices complete C . Exterior monomer activities are one; exterior role activities remain independent variables.

Write $p \succeq_{\text{coeff}} 0$ when every coefficient of p in the original activity variables is a nonnegative integer.

Theorem 6.1 (Boundary inequality). *For every distinct $i, j, k \in C$,*

$$b_{ij} b_{ik} - a_i c \succeq_{\text{coeff}} 0. \quad (6.2)$$

This holds for every finite exterior set I and every directed relation satisfying the physical bipartition assumption.

Section 6.2 gives the complete localization argument and exact finite certificate for this computer-assisted theorem. Section 6.3 derives the full core-variable Rayleigh property without another enumeration. Rank-three matroid Rayleigh theory [14] provides relevant context, but the proof here works directly with directed endpoint supports and their independent role activities; no identification with an ordinary unweighted matroid basis measure is assumed.

[Merge note: in the scalar case with one orientation (all arcs from the core C to the exterior), the boundary inequality is source 22's inequality $B_{ij} B_{ik} \geq A_i C_0$ (Section 9.4), which source 22 derives from Wagner's rank-three Rayleigh theorem; for undirected graphs its scalar form is *matching-rank-normalization*, Equation `mrn:s:tv:eq:rayleigh`. The coefficientwise form, with both orientations and independent role activities, is new.]

6.2 Complete coefficient localization

By relabeling the core, it suffices to prove $G = b_{01} b_{02} - a_0 c \succeq_{\text{coeff}} 0$.

Lemma 6.2. *Every potentially negative coefficient of G is determined by an induced relation on the three core vertices and either three or four exterior physical vertices. Their exterior physical multiplicities are respectively $(2, 1, 1)$ or $(1, 1, 1, 1)$.*

Proof. A negative coefficient must occur in $a_0 c$. The a_0 factor uses one exterior vertex and the c factor uses three distinct exterior vertices. Their union therefore has exactly one of the two stated patterns. The physical core multiplicities are $(2, 1, 1)$ in either product, and total exterior degree is four. All other monomials have coefficient zero in $a_0 c$, so their gap coefficients are nonnegative automatically.

For a fixed monomial, no support contributing to its coefficient can use a vertex absent from that monomial. Nor can it use an arc whose tail or head role variable is absent. Removing such vertices and arcs preserves the coefficient exactly. The remaining finite graph is determined solely by the roles and physical labels in the monomial. $\square$

Use role bit 1 for a tail and 0 for a head. At doubled core vertex 0, let $h \in \{0, 1, 2\}$ be the exponent of u_0 ; its head exponent is $2 - h$. Let $\epsilon_1, \epsilon_2 \in \{0, 1\}$ be the roles at vertices 1, 2. The allowed ordered allocations of the two roles at a doubled vertex are

$$\mathcal{A}(0) = \{(0, 0)\}, \quad \mathcal{A}(1) = \{(0, 1), (1, 0)\}, \quad \mathcal{A}(2) = \{(1, 1)\}. \quad (6.3)$$

Thus identical-role occurrences have one allocation, not two. The total tail degree in both products is four: the factor sizes are $2 + 2$ or $1 + 3$.

Four distinct exterior vertices. Each exterior vertex has one fixed role, and exactly $4 - h - \epsilon_1 - \epsilon_2$ of them are tails. Up to exterior relabeling, put these tails first and the heads afterward. The twelve choices of $(h, \epsilon_1, \epsilon_2)$ give exactly twelve role-monomial cases.

Three exterior vertices. Label the repeated exterior vertex $0'$ and the other two $1', 2'$. Let $\ell \in \{0, 1, 2\}$ be its tail exponent, and let $\delta_1, \delta_2 \in \{0, 1\}$ be the other exterior roles. Retain precisely the tuples satisfying

$$h + \epsilon_1 + \epsilon_2 + \ell + \delta_1 + \delta_2 = 4. \quad (6.4)$$

There are thirty-six such tuples, including the core choices.

For each role-monomial case, list every cross-partition directed arc whose tail and head roles occur in the monomial, and enumerate all subsets of that list. If both endpoint role types occur, both directions of the same physical edge are separate arc bits. The exact sizes are

Exterior multiplicities	Role types	Most arc bits	Graphs
1, 1, 1, 1	12	12	11 392
2, 1, 1	36	10	5 984
Total	48		17 376

These are complete finite case counts, not a sample of arbitrary larger graphs. Lemma 6.2 explains why they cover every potentially negative coefficient for every exterior size.

Computing one coefficient exactly

For fixed endpoint sets and fixed roles, the producer checker lists the at most six bijections between the selected core and exterior vertices. The feasibility predicate is true if *at least one* bijection uses present, role-compatible directed arcs. The number of successful bijections is never used as a coefficient.

With four exterior vertices, the coefficient in $b_{01}b_{02}$ is obtained by choosing the two exterior vertices assigned to the first factor (six choices), assigning the complement to the second, and summing over the distinct role allocations (6.3) at core 0. A choice contributes one exactly when both endpoint supports are feasible. For a_0c , choose the singleton exterior vertex (four choices), give the remaining three to c , and use the same core-role allocation rule.

With three exterior vertices, both positive factors must contain $0'$. Choose which of $1', 2'$ accompanies it in b_{01} (two choices) and put the other in b_{02} . Sum over the distinct role allocations at core 0 and exterior $0'$. In the negative product, the exterior singleton of a_0 must be $0'$, while c uses all three exterior vertices. Again sum over precisely those two allocation sets. Each choice specifies an ordered pair of endpoint supports uniquely, so subtracting these two counts is the desired coefficient.

All 17 376 cases give nonnegative coefficients; the recorded values in these potentially negative cases are 0, 1, 2, 3. The complete per-case histograms and executable checker are in the package.

Independent reconstruction

A second checker does not import the producer code or use its allocation routine. It constructs actual directed physical matchings by adjoining vertex-disjoint arcs, deduplicates their ordered endpoint sets, and separates them according to their used core subsets. It encodes the formal role monomials in base three: each exponent in a product of two supports is at most two, so

this representation has no carries. It then extracts the positive and negative product coefficients directly.

This independent enumeration checks 228 264 support instances across the same 48 role types and 17 376 graphs. Every coefficient histogram agrees with the producer's output. Together with the complete localization argument, these finite checks prove Theorem 6.1. They are the finite part of the proof, not merely floating-point evidence. No random search is used.

6.3 Full Rayleigh differences and negative correlation

For a multiaffine polynomial F , define

$$\Delta_{jk}(F) = (\partial_{z_j} F)(\partial_{z_k} F) - F \partial_{z_j} \partial_{z_k} F.$$

Theorem 6.3. *For F in (6.1), every $\Delta_{jk}(F)$ is coefficientwise nonnegative in the core monomers and original independent role activities. Consequently, after every nonnegative role-activity specialization, F is Rayleigh as a polynomial of its three core monomers at nonnegative arguments.*

Proof. For distinct i, j, k , direct cancellation gives

$$\begin{aligned} \Delta_{jk}(F) &= (a_j a_k - b_{jk}) z_i^2 \\ &\quad + (a_j b_{ik} + a_k b_{ij} - c - a_i b_{jk}) z_i + (b_{ij} b_{ik} - a_i c). \end{aligned} \quad (6.5)$$

The constant coefficient is nonnegative by Theorem 6.1. Also $a_j a_k \succeq_{\text{coeff}} b_{jk}$: every size-two support using j, k has at least one pair of single-edge witnesses counted by the product, with the same role weight; repeated exterior vertices and additional witnesses only add terms to that product.

To treat the middle coefficient, add a fresh exterior physical vertex x and only the arc $i \rightarrow x$, and put $L = u_i v_x$. Deleting that forced edge is a bijection of endpoint supports; any residual matching-witness multiplicity remains ignored. The enlarged relation has coefficients

$$a'_i = a_i + L, \quad b'_{ij} = b_{ij} + L a_j, \quad b'_{ik} = b_{ik} + L a_k, \quad c' = c + L b_{jk}.$$

It still satisfies the bipartition hypotheses of Theorem 6.1. Its boundary gap is

$$\begin{aligned} b'_{ij} b'_{ik} - a'_i c' &= b_{ij} b_{ik} - a_i c \\ &\quad + L(a_j b_{ik} + a_k b_{ij} - c - a_i b_{jk}) + L^2(a_j a_k - b_{jk}). \end{aligned} \quad (6.6)$$

The fresh variable v_x is absent from the original coefficients. Its coefficient in (6.6) is u_i times the middle coefficient of (6.5). The enlarged boundary gap is coefficientwise nonnegative, and multiplication by the monomial u_i shifts exponents injectively. The middle coefficient is therefore nonnegative as well. $\square$

The statement concerns nonnegative monomer arguments. It is not an all-real Rayleigh criterion, and no real-stability conclusion follows here. It also does not assert Rayleigh inequalities in the exterior or activity variables.

Corollary 6.4. *Fix nonnegative role activities and strictly positive core monomer activities. Give each feasible ordered support probability proportional to*

$$u^S v^T \prod_{i \in C \setminus (S \cup T)} z_i.$$

Then use of any two distinct core vertices is negatively correlated. The same is true of their unused indicators.

Proof. The normalizing constant is $F(\mathbf{z}) > 0$, since the empty support contributes $z_0 z_1 z_2 > 0$. If U_j is the indicator that j is unused, differentiation gives

$$\text{Cov}(U_j, U_k) = \frac{z_j z_k (F \partial_{z_j} \partial_{z_k} F - \partial_{z_j} F \partial_{z_k} F)}{F^2} = -\frac{z_j z_k \Delta_{jk}(F)}{F^2} \leq 0.$$

Replacing each indicator by one minus itself leaves their pairwise covariance unchanged. This proves the assertion for used indicators too. $\square$

This probability measure is on feasible endpoint supports, not on individual matching witnesses. Zero role activities cause no problem: the coefficientwise inequalities specialize directly, and the positive empty-support monomial still normalizes the measure.

6.4 A scalar cubic consequence

Write

$$\Gamma_D(t) = 1 + At + Bt^2 + ct^3, \quad A = a_0 + a_1 + a_2, \quad B = b_{01} + b_{02} + b_{12}.$$

Rank-ultra-log-concavity means log-concavity of $\gamma_k / \binom{d}{k}$ at the actual surviving degree d . Positive-weight submatchings ensure that the coefficient sequence has no internal zeros. At nonnegative activities,

$$B^2 - 3Ac = \frac{1}{2}[(b_{01} - b_{02})^2 + (b_{01} - b_{12})^2 + (b_{02} - b_{12})^2] + 3 \sum_i (b_{ij} b_{ik} - a_i c) \geq 0. \quad (6.7)$$

Moreover, $b_{ij} \leq a_i a_j$ gives $A^2 \geq 3 \sum_{i < j} a_i a_j \geq 3B$.

[*Merge note:* identity (6.7) is printed once, here. Source 11 uses it in its equality proposition, and source 22's identity for its three-core theorem is this identity with b_{ij} , a_i and c replaced by $u_i u_j b_{ij}$, $u_i a_i$ and $u_0 u_1 u_2 C_0$.] Thus the cubic Newton inequalities hold. Equation (6.7) is a scalar nonnegativity identity; its squared differences must not be mistaken for coefficientwise nonnegative polynomials.

[*Merge note:* source 12 continues with a paragraph proving the stronger normalization $A^2 \geq 4B$ after a drop to actual degree two, by Motzkin–Straus compression on the triangle-free graph of positive-weight arcs; this is the case $d = 2$ of Theorem 1.5 and is replaced by that pointer. Degrees zero and one have no inequalities to check.]

Consequently Γ_D is rank-ultra-log-concave at its actual surviving degree, including zero activities. This scalar conclusion is weaker than the main coefficientwise and negative-correlation results.

6.5 An exact obstruction at four cores

The coefficientwise assertion cannot be extended without qualification to a four-vertex shore, even inside the two-tail/three-head role-cover class.

Take tails 0, 1, 2, 3 and five distinct heads a, b, c, d, e , and only the arcs

$$N(0) = \{c, d, e\}, \quad N(1) = \{a, b, e\}, \quad N(2) = \{b, d\}, \quad N(3) = \{e\}. \quad (6.8)$$

The tail roles $\{0, 1\}$ and head roles $\{b, d, e\}$ cover every arc. A matching $0 \rightarrow c, 1 \rightarrow a, 2 \rightarrow d, 3 \rightarrow e$ shows degree four. Set tail activities to one and use the head labels as their activities. Let B_J count head supports matchable to the selected tail set J , once per feasible head set.

Boolean matching tests give

$$\begin{aligned} B_{012} &= abc + abd + abe + acd + ade + bcd + bce + bde + cde, \\ B_{013} &= ace + ade + bce + bde, \\ B_{01} &= ac + ad + ae + bc + bd + be + ce + de, \\ B_{0123} &= abce + abde + acde + bcde. \end{aligned}$$

Their boundary gap is

$$\begin{aligned} B_{012}B_{013} - B_{01}B_{0123} &= e^2(a^2d^2 - abcd + abd^2 + b^2c^2 + b^2cd + b^2d^2) \\ &= e^2 \left[\left(ad + \frac{b(d-c)}{2} \right)^2 + \frac{3}{4}b^2(c+d)^2 \right]. \end{aligned} \quad (6.9)$$

The coefficient of $abcde^2$ is -1 . Restoring tail activities multiplies the entire gap by $u_0^2u_1^2u_2u_3$, so this remains a negative coefficient in the original role-activity ring.

Nevertheless, the second line of (6.9) is nonnegative at every real head-activity assignment. This example refutes the coefficientwise extension only. It is not a counterexample to scalar Rayleigh nonnegativity, real-rootedness, or rank-ULC. No minimality assertion is made.

6.6 Proof status and reproducibility

The boundary theorem consists of a complete all-size localization proof and an exhaustive finite integer check. The full core-variable Rayleigh theorem, negative-correlation corollary, scalar consequence, and four-core obstruction are ordinary arguments once that boundary theorem is established. No proof-assistant verification or literature-wide novelty claim is made.

The package includes the producer's finite checker and all 48 histograms, the independently written direct-support checker and its receipts, exact four-core verifiers, mathematical audits, and editable source. The replay driver first checks file hashes, then runs copied checkers in temporary directories and compares deterministic receipt fields without altering the frozen package. All computations use Python's standard library and exact integer or rational arithmetic.

The physical core must remain independent. No result here permits internal core arcs. The general rank-ULC problem for a cover of at most two tail roles and three head roles, with arbitrary independent activities and physical overlap, remains open. Section 6.5 gives a specific reason that the strongest coefficientwise approach cannot simply be extrapolated to a larger shore.

[Merge note: in this report the two-tail/three-head problem is settled in part: by real stability when $|Q| \leq 2$ (source 20, Theorem 5.10), for $P \subseteq Q$ by source 11's Corollary 7.2, and for an independent physical core at its cover order by source 15's Corollary 8.6. The general case, with internal core arcs and a physical union of four or five vertices, remains open.]

7 A physical vertex cover of size three (source 11)

Source 11 is *Rank ultra log concavity under a three vertex physical cover* (5 pages as delivered; archive `physical-cover-three-result`). Its files are shipped with the prefix `11-physical-cover-`; the copy of source 12 that it embeds is not shipped, because source 12's own files are.

Abstract of source 11, as delivered. We prove actual-degree ultra-log-concavity of the endpoint-support polynomial for every finite loopless directed relation with a physical vertex

cover of size at most three. Internal core arcs are arbitrary, transitivity is unnecessary, and independent nonnegative tail/head activities are allowed. The proof combines a verified bipartite boundary inequality, support-level witness bounds, and an exact two-square decomposition. Each support is counted once, with zero activities normalized at the surviving degree. A separate bipartite-only section characterizes strict core negative correlation and equality in the upper cubic Newton inequality. The ordinary arguments depend on a replayable finite boundary certificate included in the package. No real-rootedness, proof-assistant formalization, or priority claim is asserted.

7.1 Statement and normalization

Let D be a finite loopless directed relation on physical vertex set V . Opposite arcs are allowed. Give every vertex tail activity $u_i \geq 0$ and head activity $v_i \geq 0$. A feasible support of size k is an ordered pair (S, T) of disjoint subsets of size k admitting a directed perfect matching from S to T . It has weight $\text{wt}(S, T) = u^S v^T$ and is counted once, regardless of how many matching witnesses realize it. Set

$$\gamma_k = \sum_{\substack{(S, T) \text{ feasible} \\ |S|=k}} u^S v^T, \quad \Gamma_D(t) = \sum_{k \geq 0} \gamma_k t^k, \quad \gamma_0 = 1.$$

The actual degree d is the matching number of the underlying graph of the positive-weight relation D_+ , which retains precisely the arcs $i \rightarrow j$ with $u_i v_j > 0$. Taking submatchings gives no internal zeros. Rank-ultra-log-concavity, or rank-ULC, means

$$k(d-k)\gamma_k^2 \geq (k+1)(d-k+1)\gamma_{k-1}\gamma_{k+1} \quad (1 \leq k < d), \quad (7.1)$$

at this actual surviving degree.

Theorem 7.1. *If the underlying undirected graph of D has a vertex cover of size at most three, then Γ_D is rank-ULC for every nonnegative choice of independent tail and head activities. The internal core arcs are unrestricted and transitivity is not required.*

[Merge note: Theorem 7.1 contains source 22's three-core theorem (Theorem 9.7: three core tails and no arcs into the core, so the core is a physical vertex cover) and the rank-ULC half of source 22's role-matching corollary (a role cover of size three has at most three physical vertices); source 22's ordinary proof of the former is kept there as a second route. For bipartite graphs with every vertex in one role it is the case $\nu \leq 3$ of *preorder-root-polytopes*, Corollary `lc:cor:rankthree`, itself contained in *matching-rank-normalization*, Corollary `mrn:w:rf:cor:rankfive`.]

Corollary 7.2. *Suppose every arc $i \rightarrow j$ satisfies $i \in P$ or $j \in Q$, with $|P| \leq 2$, $|Q| \leq 3$, and $P \subseteq Q$. Then Γ_D is rank-ULC at its actual surviving degree, with arbitrary independent activities.*

Proof. Every underlying edge meets the physical set Q , so Theorem 7.1 applies. $\square$

This closes the two-tail/three-head role-cover case in which both selected tail vertices also belong to the selected head set. Covers with a physical union of four or five vertices are not settled by this theorem.

[Merge note: source 11's Lemma "first" (Universal first inequality: for any directed relation of actual degree $d \geq 2$, $\gamma_1^2 \geq \frac{2d}{d-1}\gamma_2$), proved by moving weight between nonadjacent positive-weight arcs of the compatibility graph, is Theorem 1.5 with Lemma 1.4; it is replaced by that pointer.]

The lemma supplies the first cubic inequality and the stronger degree-two inequality after a degree drop. Only the last cubic inequality remains.

7.2 The exterior boundary input

Assume now that $C = \{0, 1, 2\}$ is a physical vertex cover and put $I = V \setminus C$. The exterior I is independent. Delete all internal core arcs, obtaining the bipartite physical relation D_0 on $C \sqcup I$. Define, with all original role activities included,

- a_i : the weight sum of size-one supports of D_0 using core $\{i\}$;
- b_{ij} : the weight sum of size-two supports using core $\{i, j\}$;
- c : the weight sum of size-three supports, using all three cores.

Set $A = \sum_i a_i$ and $B = \sum_{i < j} b_{ij}$.

The following is the explicit dependency of the present proof:

Lemma 7.3 (Verified three-core boundary theorem). *For every distinct $i, j, k \in C$,*

$$b_{ij}b_{ik} - a_i c \succeq_{\text{coeff}} 0 \quad (7.2)$$

in the original independent role-activity variables.

This theorem is proved in the accompanying manuscript. [*Merge note:* that manuscript is source 12, printed in Section 6; the statement above is its Theorem 6.1. Source 12's files are shipped once, with its own prefix.] Its complete all-size localization reduces every potentially negative coefficient to three or four exterior physical vertices. There are 48 role-monomial types and 17 376 relevant directed graphs. Two independently written exact checkers reproduce every coefficient histogram. That proof, its audits and the checkers are included and replayed in this package; the dependency is not replaced by a random sample. We use only its valid nonnegative specialization below.

Summing (7.2) gives

$$Ac \leq b_{01}b_{02} + b_{01}b_{12} + b_{02}b_{12}. \quad (7.3)$$

We also need the elementary coefficientwise inequality

$$c \leq a_k b_{ij}. \quad (7.4)$$

For a size-three support, choose a witnessing matching. Its edge at core k is a singleton support, and deleting it leaves a support counted by b_{ij} . The product has the original weight. Additional decompositions and intersecting products only add nonnegative contributions.

7.3 Restoring every internal core arc

For an unordered core pair $\{i, j\}$, with remaining core k , write

$$h_{ij} = \mathbf{1}_{i \rightarrow j \in D} u_i v_j + \mathbf{1}_{j \rightarrow i \in D} u_j v_i, \quad E = h_{01} + h_{02} + h_{12}, \quad w_{ij} = h_{ij} a_k.$$

Let H be the weight sum of size-two supports using all three physical core vertices. These supports use one internal core arc and one exterior arc. The other size-two supports use precisely two core vertices and belong to D_0 . A size-three matching cannot contain an internal core edge, since that would leave only one core vertex for its other two disjoint edges. Thus, exactly at support level,

$$\gamma_1 = A + E, \quad \gamma_2 = B + H, \quad \gamma_3 = c. \quad (7.5)$$

Lemma 7.4 (Internal witness bounds). *At all nonnegative activities,*

$$0 \leq w_{ij} \leq H, \quad w_{01} + w_{02} + w_{12} \leq 2H. \quad (7.6)$$

Both inequalities hold coefficientwise as support-counting statements.

Proof. A product in $h_{ij}a_k$ chooses an internal arc on $\{i, j\}$ and an exterior arc at k . Their physical endpoints are disjoint. For this fixed unordered core pair, every resulting ordered endpoint support is counted exactly once: opposite orientations of the internal arc yield different role assignments, and the exterior edge is then forced. Hence $w_{ij} \leq H$.

The sum of the three w 's counts the internal-edge witnesses of each H -support. Such a support has three core vertices and one exterior vertex. If two core vertices are tails and one is a head, the internal edge must run from one of those two tails to that head. There are at most two choices. The case of one core tail and two core heads is the reverse case. Each witness has the same prescribed support weight, so the total is at most $2H$. In particular, fork-induced duplicate witnesses are retained in this bound rather than silently counted as distinct supports in H . $\square$

Multiplying (7.4) by h_{ij} and summing yields

$$Ec \leq \sum_{i < j} w_{ij} b_{ij}. \quad (7.7)$$

At the fixed activity specialization, order the three b -values as $x \geq y \geq z \geq 0$, and let w_x, w_y, w_z be their corresponding w -values. The witness bounds imply

$$xw_x + yw_y + zw_z \leq H(x + y), \quad (7.8)$$

because its slack is exactly

$$(x - y)(H - w_x) + y(2H - w_x - w_y - w_z) + (y - z)w_z \geq 0. \quad (7.9)$$

Combining (7.3), (7.7) and (7.8) gives

$$\gamma_1 \gamma_3 \leq xy + xz + yz + H(x + y), \quad \gamma_2 = x + y + z + H.$$

Consequently

$$\begin{aligned} \gamma_2^2 - 3\gamma_1 \gamma_3 &\geq (x + y + z + H)^2 - 3[xy + xz + yz + H(x + y)] \\ &= \left(z + H - \frac{x + y}{2}\right)^2 + \frac{3}{4}(x - y)^2 \geq 0. \end{aligned} \quad (7.10)$$

Together with Lemma 1.5, this proves Theorem 7.1 at actual degree three. Degrees at most two were already handled by that lemma, so zero-weight degree drops cause no normalization gap.

For a completely explicit accounting of the slack, the proof gives the exact identity

$$\begin{aligned} \gamma_2^2 - 3\gamma_1 \gamma_3 &= \left(z + H - \frac{x + y}{2}\right)^2 + \frac{3}{4}(x - y)^2 \\ &\quad + 3 \sum_i (b_{ij} b_{ik} - a_i c) + 3 \sum_{i < j} h_{ij} (a_k b_{ij} - c) \\ &\quad + 3[(x - y)(H - w_x) + y(2H - \sum w) + (y - z)w_z]. \end{aligned} \quad (7.11)$$

Every term is nonnegative. The ordering of x, y, z is numerical, after activity specialization; (7.11) is not a coefficientwise positivity claim for the full Newton gap.

7.4 Bipartite only strictness results

The remainder of this section assumes that *there are no internal core arcs*. It must not be applied to the unrestricted internal-core theorem. Let D_+ denote the positive-weight relation, and give the three core monomers strictly positive activities z_0, z_1, z_2 . The core polynomial is

$$F = z_0 z_1 z_2 + \sum_i a_i z_j z_k + \sum_{i < j} b_{ij} z_k + c.$$

The boundary theorem and a private-head insertion prove (source 12, Theorem 6.3) that all three coefficients of

$$\begin{aligned} \Delta_{jk}(F) = & (a_j a_k - b_{jk}) z_i^2 \\ & + (a_j b_{ik} + a_k b_{ij} - c - a_i b_{jk}) z_i + (b_{ij} b_{ik} - a_i c) \end{aligned} \quad (7.12)$$

are coefficientwise nonnegative. Here $\Delta_{jk} = (\partial_j F)(\partial_k F) - F \partial_j \partial_k F$.

Proposition 7.5. *Give each endpoint support probability proportional to*

$$u^S v^T \prod_{i \in C \setminus (S \cup T)} z_i.$$

Then the covariance of the used indicators of two distinct core vertices is strictly negative exactly when they are in the same connected component of the underlying graph of D_+ . Otherwise it is zero. The unused indicators have the same covariance.

Proof. Supports and weights factor across different physical components, so those components are independent. If j, k are connected, a simple alternating path between them has length two or four, since there are only three core vertices.

For a path $j - x - k$, choose a positive available arc on each physical edge. Their product is a positive monomial of $a_j a_k$ with exterior vertex x used twice. It cannot occur in b_{jk} , whose supports require two distinct exterior vertices. Thus the quadratic coefficient of (7.12) is strictly positive at these activities.

For a path $j - x - i - y - k$, the two supports formed by edges $\{j - x, i - y\}$ and $\{i - x, k - y\}$ contribute a positive monomial to $b_{ij} b_{ik}$ on only the two exterior vertices x, y . It cannot occur in $a_i c$, because c requires three distinct exterior vertices. The constant coefficient is therefore strictly positive. Each support separately is physically disjoint; role reuse across the product is allowed.

In either case, coefficientwise nonnegativity prevents cancellation and $\Delta_{jk}(F) > 0$. The empty support makes $F > 0$, and the unused indicators U_j, U_k satisfy

$$\text{Cov}(U_j, U_k) = -\frac{z_j z_k \Delta_{jk}(F)}{F^2} < 0.$$

Replacing both indicators by one minus themselves preserves covariance. $\square$

Proposition 7.6. *Assume actual degree three, so $c > 0$, and still no internal core arcs. Then $B^2 = 3Ac$ if and only if, after deleting zero-weight arcs and isolates, the physical graph is a union of three separate stars centered at the core vertices, with equal positive total directed arc activities. In that case $\Gamma_D(t) = (1 + at)^3$ for their common activity total a .*

Proof. The exact identity (6.7) of source 12, which source 11 prints again, uses the three unordered core pairs as e, f . Equality forces the three b -values to agree and all boundary gaps to vanish.

Suppose an exterior vertex x has at least two core neighbors, say j, k . If the remaining core i has a neighbor $y \neq x$, the supports $\{i - y, j - x\}$ and $\{i - y, k - x\}$ give a positive monomial of the boundary gap for i on only two exterior vertices. It cannot occur in $a_i c$. If i has no neighbor other than x , then x is adjacent to all three cores. A saturating matching exists because $c > 0$; it uses $i - x$ and must match j to some $y \neq x$. Use j as the doubled core in the same two-support construction. In either case, one boundary gap is strictly positive, contradicting equality.

Thus no exterior vertex is shared by different cores. The graph is the stated union of stars and supports factor: $b_{ij} = a_i a_j$, $c = a_0 a_1 a_2$, and $\Gamma_D = \prod_i (1 + a_i t)$. Each $a_i > 0$, so equality of the three b -values forces $a_0 = a_1 = a_2$. The converse is immediate. Opposite arcs simply contribute their two distinct singleton-support weights to a star's total. $\square$

7.5 Proof status and independent replay

The only computational dependency is the three-core bipartite boundary theorem. The internal-core assembly and bipartite-only strictness results are ordinary arguments from that input. No real-rootedness or signed-monomer stability is asserted for arbitrary internal core arcs.

The independent audit checks all 4 096 one-exterior graphs for witness bounds, 1 600 three-exterior cases for (7.4), and 10 080 exact activity evaluations for the assembly and actual-degree inequalities. The producer checks 10 496 weighted cases, including all 64 core patterns, 6 478 zero-activity cases and 504 larger activity-induced degree drops, and verifies (7.11). These checks corroborate the ordinary all-size proof; they impose no exterior-population bound.

The archive includes the full boundary manuscript, source, audits and both finite checkers, plus the new independent and producer assembly checks. The replay driver verifies hashes and deterministic receipts using temporary copies. All code uses Python's standard library and exact arithmetic.

The general two-tail/three-head problem with four or five physical core vertices remains open. No proof-assistant formalization or priority claim is made.

8 Bipartite physical graphs and pendant heads (source 15)

Source 15 is *Lorentzian and real stable directed support polynomials* (6 pages as delivered; archive `bipartite-and-pendant-support-result`). Its files are shipped with the prefix `15-pendant-support-`.

Abstract of source 15, as delivered. We count ordered disjoint endpoint supports of directed matchings once per feasible support, with independent nonnegative tail and head activities. For any directed relation whose physical graph is bipartite, we prove a multiaffine Lorentzian theorem at the physical shore order. It gives actual-degree ultra-log-concavity when a positive-weight matching saturates a shore, and accounts for overlap among selected core roles. We also prove that a stable selected-head transversal seed remains stable after adding pendant heads at arbitrarily many tail hubs and merging physical copies. This yields negative real roots and actual-degree ultra-log-concavity even on zero-weight degree drops. Singleton-or-universal neighborhoods give an explicit stable seed in arbitrary rank. The proofs are ordinary arguments; exact independent checks are supplementary. The general two-tail/three-head question remains open. No proof-assistant formalization or priority claim is made.

8.1 Supports and the normalization issue

Let D be a finite loopless directed relation on V . Opposite arcs are allowed; transitivity is not assumed. Assign tail activities $u_i \geq 0$ and head activities $v_i \geq 0$.

Definition 8.1. A *feasible support* of size k is an ordered pair (S, T) of disjoint subsets of V , with $|S| = |T| = k$, admitting a directed perfect matching from S to T . Its weight is

$$\text{wt}(S, T) = u^S v^T = \prod_{i \in S} u_i \prod_{j \in T} v_j.$$

Every feasible pair contributes once, regardless of how many matching witnesses it has. Define

$$\gamma_k = \sum_{\substack{(S, T) \text{ feasible} \\ |S|=k}} \text{wt}(S, T), \quad \Gamma_D(t) = \sum_{k \geq 0} \gamma_k t^k.$$

The empty support has weight one, so $\gamma_0 = 1$.

The two activities of the same physical vertex can differ. Replacing physical vertices by independent role copies without later excluding their simultaneous use changes the polynomial.

Let D_+ retain exactly the arcs $i \rightarrow j$ for which $u_i v_j > 0$. The actual degree d of Γ_D is the matching number of the underlying undirected graph of D_+ . Indeed, every positive-weight support supplies such a matching; conversely, orient each edge of such a matching in an available direction. Taking submatchings also shows that $\gamma_k > 0$ for $0 \leq k \leq d$.

Definition 8.2. For an integer $r \geq d$, the coefficients satisfy *order- r ultra-log-concavity* if

$$k(r-k)\gamma_k^2 \geq (k+1)(r-k+1)\gamma_{k-1}\gamma_{k+1} \quad (1 \leq k < r), \quad (8.1)$$

with coefficients beyond degree d set to zero. *Rank-ULC* means (8.1) with $r = d$, the actual surviving degree.

These distinctions matter. The binary polynomial

$$s^3 + 3s^2t + 3st^2 = s(s^2 + 3st + 3t^2)$$

is Lorentzian, but its degree-two quotient is not: $3^2 < 4 \cdot 1 \cdot 3$. Thus division by a common monomial cannot be used to upgrade an order bound to actual-degree ULC.

Write $\mathbb{H} = \{z \in \mathbb{C} : \text{Im } z > 0\}$. A real polynomial is *real stable* if it is nonzero whenever every variable lies in $\mathbb{H}$.

We use the classical Lorentzian theory of Brändén and Huh [6]. In particular, matroid basis polynomials are Lorentzian; products, nonnegative linear substitutions and multiaffine-part extraction preserve this class. For a nonzero binary homogeneous polynomial $\sum_{k=0}^r a_k s^{r-k} t^k$, Lorentzianity is equivalent to the log-concavity of $a_k / \binom{r}{k}$, with interval support [6, Example 2.26]. This paragraph invokes that theory, rather than reproving it.

8.2 A polynomial for bipartite physical graphs

Theorem 8.3. Suppose $V = C \sqcup I$ is a bipartition of the underlying undirected graph of D . Every arc crosses the partition, but may have either orientation and may coexist with its reverse. Then

$$F_C(\mathbf{z}, \mathbf{x}) = \sum_{(S,T) \text{ feasible}} \text{wt}(S,T) \prod_{c \in C \setminus (S \cup T)} z_c \prod_{i \in I \cap (S \cup T)} x_i \quad (8.2)$$

is a nonzero multiaffine Lorentzian polynomial, homogeneous of degree $|C|$.

Lemma 8.4 (Core merge). On polynomials affine in a, b , the coefficientwise operation

$$ab \mapsto z, \quad a \mapsto 1, \quad b \mapsto 1, \quad 1 \mapsto 0 \quad (8.3)$$

preserves Lorentzianity, allowing zero output. It lowers homogeneous degree by one. It also preserves real stability, allowing zero output.

Proof. The Lorentzian assertion is precisely [6, Lemma 3.3], after renaming its surviving variable. Alternatively, the operator is homogeneous of degree -1 , preserves coefficient nonnegativity, and has algebraic symbol

$$T[(a+r)(b+s)] = z + r + s.$$

This symbol is homogeneous and real stable, hence Lorentzian; the full symbol includes only additional factors $(w_j + y_j)^{\kappa_j}$ for untouched variables. The finite-degree symbol theorem [6, Theorem 3.2] applies.

For an elementary stability proof, write $f = Aab + Ba + Cb + D$. Then the image $Az + B + C$ equals

$$\left. \frac{\partial}{\partial t} f(t, t) \right|_{t=z/2}.$$

Diagonalization, differentiation and positive rescaling preserve real stability or give zero; see [5, 15]. Every surviving monomial in (8.3) loses exactly one degree. $\square$

[Merge note: this is the copy merge of source 20, (5.3), with source 16's elementary realization (Section 5). The same operation glues two undirected graphs at a vertex in *preorder-root-polytopes*, Lemma `ade:mat:lem:vertex-sum` (support-exact vertex sum).]

Proof of Theorem 8.3. For the outgoing side, form a transversal matroid M_+ with matching positions indexed by C , and ground set consisting of an exterior head copy for each $i \in I$ and a private dummy a_c for each $c \in C$. The exterior copy of i is adjacent to c exactly when $c \rightarrow i$ is an arc; a_c is adjacent only to c . The dummies form a basis, so the rank is exactly $|C|$.

A set consisting of exterior copies $R \subseteq I$ and dummies $C \setminus U$, for $U \subseteq C$, is a basis precisely when $|U| = |R|$ and U can be matched to R . Its basis monomial records the endpoint sets once, not their matching witnesses. Similarly form M_- from arcs $i \rightarrow c$, using exterior tail copies and private dummies b_c . Its bases encode (L, W) with $L \subseteq I$, $W \subseteq C$, and a matching $L \rightarrow W$.

The squarefree basis polynomials B_+, B_- are Lorentzian [6, Theorem 3.10]. First assume positive activities. In B_+ substitute a_c/u_c for a_c and $v_i x_i$ for the exterior variable, and multiply by $\prod_{c \in C} u_c$. A basis (U, R) now has coefficient $u^U v^R$. In B_- substitute b_c/v_c and $u_i y_i$, and multiply by $\prod_{c \in C} v_c$. Its coefficient for (L, W) is $u^L v^W$.

Multiply the two polynomials, identify $x_i = y_i$, and take the multiaffine part. Squared exterior variables correspond exactly to vertices selected in both exterior roles, so this discards precisely $R \cap L \neq \emptyset$. All other variables were already multiaffine. The result remains Lorentzian by [6, Theorem 2.10, Corollaries 2.32 and 3.5], and has degree $2|C|$.

For each $c \in C$, merge a_c, b_c by Lemma 8.4. If both dummies occur, c is unused in both roles and becomes one physical monomer z_c . If exactly one dummy occurs, c is used in exactly one role and the dummy disappears. Neither dummy occurs exactly when c is used in both roles, and that term is discarded. The degree falls by one at each of the $|C|$ merges.

A surviving pair of bases determines

$$S = U \cup L, \quad T = R \cup W, \quad U \cap W = \emptyset, \quad R \cap L = \emptyset.$$

Conversely, intersecting a feasible ordered support with C and I determines these four sets uniquely. Every witnessing matching splits into the two orientations across the bipartition, which establish the two basis conditions. This is a bijection of supports and surviving pairs of bases. Distinct role assignments may have the same physical-variable monomial, but their weights should add, because they are distinct ordered supports.

A support of size k uses exactly k vertices in each shore. It therefore contributes x -degree k and z -degree $|C| - k$, as required in (8.2). The all-dummy term survives as $\prod_{c \in C} z_c$, with coefficient one. Finally, nonnegative activities follow by limits in the closed Lorentzian cone; the same coefficient-one term prevents zero output. $\square$

[Merge note: with every arc oriented from C to I this is the two-shore setting of *preorder-root-polytopes*, Theorem `lor:thm:weightedULC`, normalized by the shore order. Corollary 8.5 is an order- $|C|$ statement; it gives rank-ULC only under the saturation hypothesis stated there.]

Corollary 8.5. *The coefficients of Γ_D satisfy order- $|C|$ ULC, and likewise order- $|I|$ ULC. If the positive-weight relation admits a matching saturating either shore, they satisfy actual-degree rank-ULC.*

Proof. Set every $z_c = s$ and every $x_i = t$ in (8.2). The result is $\sum_k \gamma_k s^{|C| - k} t^k$, which remains Lorentzian. Apply the binary characterization, and interchange the shores for the second statement. $\square$

For a disconnected graph, each component's bipartition may be reversed independently. Placing the smaller shore of each nontrivial component in C , and isolated vertices in I , gives the order

$$r = \sum_{K \text{ nontrivial component}} \min\{|C_K|, |I_K|\}. \quad (8.4)$$

The theorem gives actual-degree ULC if every chosen smaller shore is saturated in D_+ . The same observation can be applied after zero-weight arcs are removed.

Corollary 8.6 (Overlapping independent role covers). *Suppose every arc $i \rightarrow j$ satisfies $i \in P$ or $j \in Q$, where $P, Q \subseteq V$ may overlap. If the physical core $C = P \cup Q$ is independent in the underlying graph, then Γ_D satisfies order- $|P \cup Q|$ ULC. A positive-weight matching saturating C makes this actual-degree rank-ULC.*

Proof. There are no exterior-to-exterior arcs by the role-cover condition and no core-to-core arcs by independence. Thus C and $V \setminus C$ are a physical bipartition. Apply Corollary 8.5. $\square$

For a cover of two tail roles and three head roles, this improves the cover order from five to four or three when the selected physical sets intersect in one or two vertices. It does not allow internal physical core arcs.

8.3 Pendant heads attached to a stable transversal seed

Theorem 8.7. *Fix $Q \subseteq V$, and suppose every vertex $j \notin Q$ has at most one incoming neighbor in D . Form the transversal matroid M_Q with positions Q , all tail copies i^+ as ground elements adjacent to $N_D^+(i) \cap Q$, and a private dummy d_q adjacent only to each $q \in Q$. If its ordinary basis polynomial B_{M_Q} is real stable, then the signed physical monomer polynomial*

$$\mu_D(\mathbf{z}) = \sum_{(S,T) \text{ feasible}} (-1)^{|S|} \text{wt}(S,T) \prod_{i \notin S \cup T} z_i \quad (8.5)$$

is real stable for every choice of nonnegative independent activities. Consequently Γ_D has only negative real zeros and is rank-ULC at its actual surviving degree.

[Merge note: source 22's one-tail side criterion (Theorem 9.11) is the case of this theorem with a single tail hub p : there every head outside Q has at most the one in-neighbour p , and the matroid is the same. Source 22's proof (seed identity, pendant insertion, physical merge) is the proof below and is replaced there by a pointer. The seed construction is also used by batch-73 manuscript 08, printed in *matching-rank-normalization*.]

There is no bound on $|Q|$, on the number of distinct tail hubs, or on physical overlap. If a role cover $P^+ \cup Q^-$ is specified, the condition says that every uncovered head has degree at most one; the unique parents may be different elements of P . Internal physical core arcs are permitted.

Proof. Split every physical vertex into tail and head copies. First retain all tail copies and only the head copies in Q , with the specified arcs into Q . Write a_i for tail-copy monomers and b_q for these head-copy monomers. For positive activities, the split seed has signed monomer polynomial

$$\mu_0(\mathbf{a}, \mathbf{b}) = \left(\prod_{i \in V} a_i \right) \left(\prod_{q \in Q} v_q \right) B_{M_Q} \left(x_i = -\frac{u_i}{a_i}, d_q = \frac{b_q}{v_q} \right). \quad (8.6)$$

A basis consists of selected tails S and dummies for the unused heads $Q \setminus T$; its basis condition says that the selected tail copies S can be matched to the head copies T in the split role graph. No physical disjointness is imposed yet. Its transformed monomial is

$$(-1)^{|S|} u^S v^T \prod_{i \notin S} a_i \prod_{q \in Q \setminus T} b_q.$$

Thus (8.6) counts supports once. All denominators cancel by multiaffinity. For upper-half-plane monomer arguments, both $-u_i/a_i$ and b_q/v_q lie in the upper half-plane. Stability of the basis polynomial therefore gives stability of μ_0 .

Now insert each missing head copy j^- , $j \notin Q$. An isolated head contributes the factor b_j . Otherwise let i^+ be its unique parent and put $w = u_i v_j$. The exact support recurrence is

$$\mu_{\text{new}} = b_j \mu_{\text{old}} - w \partial_{a_i} \mu_{\text{old}} = b_j \mu_{\text{old}} \left(a_i - \frac{w}{b_j} \right), \quad (8.7)$$

with other arguments unchanged. Indeed, a support using the new head must contain the forced edge $i^+ \rightarrow j^-$; removing its endpoints leaves exactly a support in which i^+ is unused. This proves the recurrence without matching multiplicity. Its second equality uses affinity in a_i .

For $b_j \in \mathbb{H}$ and $w \geq 0$, the number $-w/b_j$ has nonnegative imaginary part, so the shifted a_i stays strictly upper. Equation (8.7) proves stability after each insertion, at any sequence of tail hubs. Multiaffinity prevents a previously used hub from being used again.

Once all head copies are present, merge each physical pair by (8.3). Exactly the supports with disjoint physical endpoint sets survive, giving (8.5). The coefficient-one all-unused monomial survives every stage. Nonnegative activities follow by limits and this monomial excludes an identically zero limit.

With $n = |V|$, diagonal specialization yields

$$\mu_D(s, \dots, s) = s^n \Gamma_D(-s^{-2}). \quad (8.8)$$

The left side has only real zeros. A zero ρ of Γ_D that is not negative real would give a nonreal solution of $s^2 = -1/\rho$, contradicting (8.8). Since $\Gamma_D(0) = 1$, all zeros are strictly negative. Newton's inequalities apply at the actual remaining degree, including after weights vanish. $\square$

8.4 An explicit stable seed in arbitrary rank

Corollary 8.8. *Under the pendant-head assumption of Theorem 8.7, suppose every nonempty set $N_D^+(i) \cap Q$ is a singleton or all of Q . Then its conclusions hold, without any additional matroid hypothesis.*

Proof. Write $r = |Q| \geq 2$. For $q \in Q$, let

$$R_q = d_q + \sum_{\{i: N_D^+(i) \cap Q = \{q\}\}} x_i,$$

and let $g_1, \dots, g_m$ be the separate variables for tails whose neighborhood is all of Q . Empty-neighborhood elements are loops. The basis polynomial is

$$B_{M_Q} = e_r((R_q)_{q \in Q}, g_1, \dots, g_m). \quad (8.9)$$

To see this, singleton elements with the same neighbor, together with their private dummy, form one parallel class. An r -element set can be matched precisely when it uses at most one element from each such class. Its remaining universal elements can then be assigned to the

unused positions. This is exactly (8.9); distinct universal elements must not be merged into a single parallel class.

The elementary symmetric polynomial is stable: differentiate the stable product $\prod_j (s + y_j)$ sufficiently many times in s , then specialize $s = 0$. Replacing variables by nonnegative class sums preserves stability. For $r = 1$ the basis polynomial is linear, with no separate universal class; for $r = 0$ the seed is edgeless and the pendant construction starts from the product of tail monomers. $\square$

If a physical tail belongs to Q , a universal neighborhood would include a loop and is therefore disallowed by the original loopless hypothesis. There is no conflict with the corollary: the condition is checked on the actual relation.

An overlapping non-bipartite example. Let $V = \{0, \dots, 8\}$, $P = \{0, 1\}$, and $Q = \{1, 2, 3\}$. Include only

$$0 \rightarrow 1, 2, 3, 7; \quad 1 \rightarrow 0, 2, 8; \quad i \rightarrow 1, 2, 3 \quad (i = 4, 5, 6).$$

The physical graph contains the triangle $0, 1, 2$, so Theorem 8.3 does not apply. The selected role cover overlaps at vertex 1. Every uncovered head has at most one parent, and every Q -neighborhood is empty, singleton or universal. Corollary 8.8 therefore gives real-rootedness for all independent nonnegative activities. At unit activities,

$$\Gamma_D(t) = 1 + 16t + 47t^2 + 31t^3 + 4t^4. \quad (8.10)$$

There are 99 feasible supports, but 163 directed matching witnesses. The independent checkers recover (8.10) by deduplicating endpoint sets.

8.5 Reproducibility and the remaining question

The proofs above are unbounded ordinary mathematical arguments. The package includes two independently written audit programs, their exact-check receipts, and the producer's separate checkers. They corroborate support identities, boundary cases and examples; finite tests are not the premise of either theorem.

For the bipartite construction, the independent audit checks 5 120 graph-weight instances, including all 4 096 directed subrelations of $K_{2,3}$. It compares direct physical support enumeration with Hall-based transversal constructions, tests symmetric basis exchange, and performs 13 152 exact rational Hessian-inertia checks. The separate producer check covers 12 800 graph-weight-shore tests, including 4 345 zero-activity cases.

For the pendant criterion, the independent audit checks 1 884 weighted cases, including all 1 024 eligible loopless four-vertex relations with $|Q| = 2$, and random seed ranks up to five. These include 1 734 zero-activity cases and 880 activity-induced degree drops. It checks exact operator/support identities, the elementary-symmetric Hall identity, negative real roots at the surviving degree, and 70 752 exact Rayleigh evaluations. The producer separately checks 480 random weighted cases and (8.10) by rational Sturm sequences. Rayleigh evaluations at finitely many points are supplementary checks, not stability certificates.

The two structural conclusions have different normalization guarantees. The bipartite theorem gives the shore order, or (8.4); saturation is required to identify that order with the actual degree. The stable-seed theorem instead gives real-rootedness and hence actual-degree ULC directly, even on degree-dropping zero-weight faces.

Open question. Does every finite directed relation whose bipartite role graph has a vertex cover comprising at most two tail copies and three head copies have a rank-ULC support polynomial

for all nonnegative independent activities, with physical overlap allowed? The present results cover independent physical cores with suitable saturation, and stable transversal seeds with pendant uncovered heads. They do not resolve the general internal-core case or all deficient-rank cases. Real-rootedness is a stronger property and is not asserted for the entire class.

[*Merge note:* the case of two tail and two head roles is settled in the strongest sense by source 20 (Theorem 5.10), which source 15 does not cite; the case $P \subseteq Q$ of the question by source 11 (Corollary 7.2). The question as stated, with $|Q| = 3$, internal core arcs and a physical union of four or five vertices, remains open.]

For comparison, the bimatroid ULC theorem of Röhrle and Ulirsch [13, Theorem A] is normalized by the smaller ground-side size, not by the actual bimatroid rank. Likewise, the current revision of Alexandersson and Jal [4, Corollary 4.1] uses the number of board rows. Neither is used here as an unproved actual-degree shortcut. All stability assumptions in Theorem 8.7 are explicit; arbitrary transversal matroids are not assumed to have the half-plane property.

9 Independent role activities through actual degree three (source 22)

Source 22 is *Independently weighted preorder support polynomials of actual degree at most three* (21 pages as delivered; archive `independent-role-degree-three-package`). Its files are shipped with the prefix `22-independent-role-`; its nested audit package is shipped as the delivered zip, the only carrier of its role certificates. Archive 31, *Vertex-weighted preorder support polynomials of actual degree at most three*, is an earlier version of this manuscript, with one activity per vertex; its theorem is Corollary 9.14 below, and it is not printed separately.

Abstract of source 22, as delivered. For a finite preorder, we count ordered disjoint tail and head supports admitting a directed matching, counting each support once. Assign arbitrary nonnegative activities independently to the tail and head role of every vertex. We prove ultra-log-concavity with binomial normalization at the actual surviving degree whenever that degree is at most three, with no bound on the number of vertices. The proof combines the Gallai–Edmonds decomposition with ordinary structural inequalities and a complete exact verification through seven vertices. Its main ingredients are Motzkin–Straus, an all-rank articulation theorem, rank-three matroid Rayleigh inequalities, two-attachment compression and moment cones, and rational nonnegative identities. A half-plane-property side-matroid criterion resolves the final finite cases. One ten-element side matroid is certified by twenty half-plane-property minors and an exact rank-32 rational Gram identity for a Rayleigh difference, using Wagner–Wei’s criterion. Published certified matroid data are explicit dependencies. Portable exact checkers and all local verification inputs are included. General real-rootedness fails already in degree three, as an explicit example shows. This is a computer-assisted proof, not a proof-assistant formalization or a literature-wide priority claim.

9.1 Definitions and the main theorem

Let R be a preorder on a finite set V : it is reflexive and transitive, but need not be antisymmetric. Its directed arcs are the *off-diagonal* pairs $i \rightarrow j$ with $i \neq j$ and iRj . Reflexive loops never participate in a matching.

Definition 9.1. A feasible support of size k is an ordered pair (S, T) of subsets of V such that $S \cap T = \emptyset$, $|S| = |T| = k$, and some bijection $\pi : S \rightarrow T$ satisfies $i \rightarrow \pi(i)$ for every $i \in S$. For

independent nonnegative tail and head activities $\mathbf{u} = (u_i)_{i \in V}$ and $\mathbf{v} = (v_i)_{i \in V}$, put

$$\gamma_k = \sum_{(S,T) \text{ feasible, } |S|=k} u^S v^T, \quad \Gamma_R(z; \mathbf{u}, \mathbf{v}) = \sum_{k \geq 0} \gamma_k z^k, \quad (9.1)$$

where $u^S v^T = \prod_{i \in S} u_i \prod_{j \in T} v_j$. The empty support contributes $\gamma_0 = 1$.

A support is counted once, irrespective of the number of witnessing bijections. However, *different ordered pairs with the same physical union are different summands*. Thus the object is neither an unordered endpoint-set enumerator nor a matching-witness enumerator. Physical disjointness $S \cap T = \emptyset$ is essential.

The single-vertex-activity model is the specialization $u_i = v_i = x_i$. It weights a support by $\prod_{i \in S \cup T} x_i$. The theorem below includes this specialization but does not require it.

Definition 9.2. For a nonnegative sequence $(g_0, \dots, g_d)$ with $g_0, g_d > 0$ and no internal zeros, *rank-ultra-log-concavity*, abbreviated rank-ULC, means

$$\left(\frac{g_k}{\binom{d}{k}} \right)^2 \geq \frac{g_{k-1}}{\binom{d}{k-1}} \frac{g_{k+1}}{\binom{d}{k+1}} \quad (1 \leq k < d). \quad (9.2)$$

Here d is the actual polynomial degree, not a degree obtained by adding zero coefficients.

Theorem 9.3 (Independent-role degree-three theorem). *For every finite preorder R and every assignment $u_i, v_i \geq 0$, if the actual surviving degree $d = \deg \Gamma_R(z; \mathbf{u}, \mathbf{v})$ is at most three, then $\Gamma_R(z; \mathbf{u}, \mathbf{v})$ is rank-ULC. In particular,*

$$d = 2 : \quad \gamma_1^2 \geq 4\gamma_2, \quad (9.3)$$

$$d = 3 : \quad \gamma_1^2 \geq 3\gamma_2, \quad \gamma_2^2 \geq 3\gamma_1\gamma_3. \quad (9.4)$$

The conditions in degrees zero and one are vacuous.

9.1.1 What the theorem does and does not say

The number of vertices is unrestricted. In particular, this is not merely an enumeration of preorders on seven vertices. The reduction to bounded cores, with arbitrary exterior populations and heterogeneous activities, is part of the proof.

Several structural ingredients below hold even for directed relations that are not transitive. Their scopes are stated separately. The global theorem uses transitivity in its bounded-core reduction. It does not imply real-rootedness; see Section 9.10. In particular, stability of the auxiliary matroids used for certain finite targets is a sufficient local tool, not an assertion about every preorder support polynomial.

The accompanying package separates ordinary proofs, finite polynomial identities, exhaustive finite-domain coverage, and audit receipts. An audit receipt identifies exactly what was checked; it is not a substitute for the mathematical reduction. Numerical optimization was used in discovery, but no numerical solver is needed to verify any certificate.

9.2 Degree, zero activities, and the universal first gap

Let G_R be the simple undirected graph whose edge ij is present when at least one of $i \rightarrow j$ and $j \rightarrow i$ is present. Write $\nu(G_R)$ for its matching number.

Lemma 9.4. *If all relevant activities are positive, then $\deg \Gamma_R = \nu(G_R)$ and the coefficients up to the actual degree are positive. With nonnegative independent role activities, there is a preorder with positive activities and the same support polynomial, whose physical matching number equals the original surviving degree.*

Proof. A witnessing bijection gives an undirected matching. Conversely, orient each edge of an undirected matching along a direction available in R . Its disjoint tail and head sets form a feasible support with positive weight. Submatchings give positivity at every lower degree.

For the second statement let $U = \{i : u_i > 0\}$ and $W = \{j : v_j > 0\}$. Retain exactly the original off-diagonal arcs $i \rightarrow j$ with $i \in U$, $j \in W$, and then add all reflexive loops. This relation is transitive. Indeed, a surviving nontrivial path $i \rightarrow j \rightarrow k$ has $i \in U$, $k \in W$ and the original transitive relation iRk ; its shortcut survives when $i \neq k$, and otherwise the required loop was added. Steps containing loops are harmless.

A positive-weight original support uses only retained arcs. Conversely every support of the retained relation uses only previously positive role activities. Set a previously zero activity to 1 in its now-unused role. No new support is created, so the polynomial is unchanged and all activities are positive. The first statement now identifies its matching number with the original surviving degree. $\square$

This filtering argument is stronger than vertex deletion: a vertex may still be usable in one role when its other role has zero activity. It proves the actual-degree scope even if the original, unfiltered comparability graph has matching number larger than three. In the vertex specialization ordinary deletion of zero-activity vertices is enough.

[*Merge note:* the first statement of this lemma is Lemma 1.3 (*preorder-root-polytopes*, Theorem `gam:thm:deficiency`, for preorders). The second, the transitivity-preserving filter for zero role activities, is source 22's own; source 22's earlier version (archive 31) mentioned it only as a remark.]

[*Merge note:* source 22's Proposition "Universal first inequality" (for any finite loopless directed relation with positive independent role activities and actual degree $d \geq 2$, $\gamma_1^2 \geq \frac{2d}{d-1}\gamma_2$, and the same at the surviving degree for nonnegative activities) is Theorem 1.5; its proof by Motzkin–Straus on the graph of arcs with four distinct endpoints is the proof printed there.]

For degree two the quadratic $1 + \gamma_1 z + \gamma_2 z^2$ thus has negative real roots. Degree-one polynomials do as well. This observation will handle disconnected graphs without needing a general convolution theorem for ULC sequences.

Lemma 9.5 (Products over components). *If R has underlying connected components $R_1, \dots, R_m$, then $\Gamma_R = \prod_i \Gamma_{R_i}$. At total degree at most three, rank-ULC follows if the unique degree-three component, when present, is rank-ULC; otherwise the product is real-rooted.*

Proof. Every feasible support splits uniquely among components, and witnesses can be combined. If a component has degree three, all others have degree zero. Otherwise every component has degree at most two and is real-rooted by Proposition 1.5. Newton's inequalities apply to their product at its actual degree. $\square$

9.3 An ordinary all-rank articulation argument

This structural result does not require transitivity and is useful beyond the main theorem.

Theorem 9.6. *Let D be a loopless directed relation and h a vertex such that every component of the underlying graph of $D - h$ has matching number at most one. For arbitrary nonnegative independent role activities, Γ_D has only negative real zeros. In particular it is rank-ULC at its actual degree.*

Proof. Let the components after deleting h be $C_1, \dots, C_m$. Write

$$Q_i(z) = \Gamma_{C_i}(z) = 1 + a_i z, \quad P_i(z) = \Gamma_{C_i \cup \{h\}}(z) - Q_i(z) = c_i z + d_i z^2.$$

All coefficients are nonnegative, and

$$d_i \leq a_i c_i. \quad (9.5)$$

Indeed, a size-two support counted by d_i has a witness consisting of an arc incident to h and a disjoint arc inside C_i . The product $a_i c_i$ contains that support's weight at least once. Extra witnesses and intersecting arc pairs only add nonnegative terms. This is a coefficientwise inequality in the role activities.

At support level, the exact identity is

$$\Gamma_D = \prod_i Q_i + \sum_i P_i \prod_{j \neq i} Q_j. \quad (9.6)$$

A support omitting h contributes to the product. If it uses h , precisely one component contains an odd number of its selected endpoints: the component paired with h . Every other component has an even number. Thus that component is determined by the support itself, independent of any choice of witness; there is no duplicate counting in the sum.

For $a_i > 0$,

$$\frac{P_i(z)}{Q_i(z)} = \frac{d_i}{a_i} z + \left(c_i - \frac{d_i}{a_i} \right) \frac{z}{1 + a_i z}. \quad (9.7)$$

Both coefficients are nonnegative by (9.5). If $a_i = 0$, then $d_i = 0$ and the quotient is $c_i z$. For $z \in \mathbb{H} = \{z : \operatorname{Im} z > 0\}$,

$$\operatorname{Im} \frac{z}{1 + az} = \frac{\operatorname{Im} z}{|1 + az|^2} > 0 \quad (a \geq 0).$$

Hence $1 + \sum_i P_i/Q_i$ is either the constant 1 or has positive imaginary part on $\mathbb{H}$. It has no zero there, and neither does any Q_i . Equation (9.6) excludes upper-half-plane zeros of Γ_D ; conjugation excludes lower-half-plane zeros. Its nonnegative coefficients and constant term 1 exclude nonnegative real roots. The argument includes zero activities directly. $\square$

The odd-endpoint argument in (9.6) is important: simply adding local matching polynomials would generally count witnesses instead of supports.

9.4 Three active tails: a rank-three Rayleigh proof

Theorem 9.7. *Let a loopless directed relation have a three-vertex set $C = \{0, 1, 2\}$ and an independent exterior X . Allow arbitrary internal arcs in C and arbitrary arcs from C to X , but no arcs from X to C . Then its support polynomial is rank-ULC for arbitrary nonnegative independent role activities. The order-dual assertion also holds.*

Only the second cubic inequality needs proof, by Proposition 1.5. Transitivity is not assumed in this theorem.

[Merge note: this theorem is a special case of source 11's Theorem 7.1: the three core vertices form a physical vertex cover, since there are no exterior-to-exterior arcs. Source 11's proof rests on source 12's computer-assisted boundary inequality; the proof below is a second route for this special case, ordinary apart from Wagner's rank-three Rayleigh theorem. Its inequality $B_{ij}B_{ik} \geq A_i C_0$ is the scalar one-orientation case of source 12's Theorem 6.1.]

9.4.1 Exterior head sets and private dummies

Let $N_i \subseteq X$ be the exterior heads available to tail i . Define

$$\begin{aligned} A_i &= \sum_{x \in N_i} v_x, \\ B_{ij} &= \sum_{\{x,y\} \subseteq X \text{ matchable from } \{i,j\}} v_x v_y, \\ C_0 &= \sum_{\{x,y,z\} \subseteq X \text{ matchable from } \{0,1,2\}} v_x v_y v_z. \end{aligned}$$

Each head set is counted once. For distinct i, j, k we use

$$B_{ij}B_{ik} \geq A_i C_0, \quad A_k B_{ij} \geq C_0. \quad (9.8)$$

For the first inequality, adjoin private dummy heads d_i , each adjacent only to tail i . The resulting transversal matroid on heads and dummies has rank exactly three. Its basis polynomial, after substituting the exterior activities, is

$$M(\mathbf{d}) = d_0 d_1 d_2 + \sum_i A_i d_j d_k + \sum_{i < j} B_{ij} d_k + C_0. \quad (9.9)$$

Wagner's theorem states that every rank-three matroid is Rayleigh [14, Theorem 1.1]. Thus $\partial_j M \partial_k M - M \partial_{jk} M \geq 0$ on the nonnegative orthant. Setting the three dummy variables to zero, by continuity, gives $B_{ij}B_{ik} \geq A_i C_0$.

The second inequality is coefficientwise. For each feasible three-head set choose a witness and its head paired with k . The remaining pair is feasible for $\{i, j\}$ and gives the same monomial in $A_k B_{ij}$. Different three-head sets give different squarefree monomials; repeated-head monomials or extra witnesses cause no difficulty.

9.4.2 Restoring every internal arc

Let $\epsilon_{ij} = 1$ when $i \rightarrow j$ is an internal arc, and 0 otherwise. For distinct i, j, k , let

$$H_j = \sum_{x \in (N_k \text{ if } \epsilon_{ij}=1) \cup (N_i \text{ if } \epsilon_{kj}=1)} v_x.$$

An absent set in this notation means the empty set. This union, rather than a sum with possible duplication, counts exterior heads which can be used alongside core head j . Set

$$a_i = A_i + \epsilon_{ij} v_j + \epsilon_{ik} v_k, \quad b_{ij} = B_{ij} + v_k H_k.$$

Every tail lies in C , so direct support counting gives

$$\gamma_1 = \sum_i u_i a_i, \quad \gamma_2 = \sum_{i < j} u_i u_j b_{ij}, \quad \gamma_3 = u_0 u_1 u_2 C_0. \quad (9.10)$$

In particular a size-three support uses every core vertex as a tail and every head in the exterior.

The Rayleigh-type inequalities survive the restoration:

$$\begin{aligned} b_{ij}b_{ik} - a_i C_0 &= (B_{ij}B_{ik} - A_i C_0) \\ &\quad + v_j(B_{ij}H_j - \epsilon_{ij}C_0) + v_k(B_{ik}H_k - \epsilon_{ik}C_0) + v_j v_k H_j H_k \geq 0. \end{aligned} \quad (9.11)$$

If $\epsilon_{ij} = 0$ its bracket is nonnegative. If it is 1, then $H_j \geq A_k$, and the second inequality of (9.8) proves the same conclusion. The other bracket is symmetric; all remaining terms are nonnegative.

Put $U = u_0 u_1 u_2$ and $x_{ij} = u_i u_j b_{ij}$. Expansion yields the exact identity

$$\gamma_2^2 - 3\gamma_1\gamma_3 = \frac{1}{2}[(x_{01} - x_{02})^2 + (x_{01} - x_{12})^2 + (x_{02} - x_{12})^2] + 3U \sum_i u_i (b_{ij} b_{ik} - a_i C_0).$$

[Merge note: this is the three-squares identity (6.7) of source 12 with b_{ij} , a_i , c replaced by x_{ij} , $u_i a_i$, $U C_0$; source 22's label for it is redirected there.] Equations (9.11)–(6.7) prove Theorem 9.7. If the degree drops below three, the stronger surviving-degree inequality is supplied by Proposition 1.5, not by padding the cubic formula.

9.5 Role covers and a structural corollary

The *bipartite role graph* has a tail copy and a head copy of every physical vertex, with an edge $i_{\text{tail}} j_{\text{head}}$ for every arc $i \rightarrow j$. Its matching number can exceed the physical matching number: a role matching can use the two copies of a physical vertex simultaneously.

We use the following independently verified companion theorem, whose full ordinary proof and portable algebraic checker are included as a separate dependency (source 20's package; [Merge note: source 22 restates here, as its Theorem “Two-tail/two-head cover”, source 20's Theorem 5.10 (Section 5); the statement is not repeated, and source 22's references to it point there. Its embedded copy of source 20's package is not shipped.]). Here real stability means nonvanishing whenever all variables lie in $\mathbb{H}$. The companion proof first splits roles, treats every subset of the four possible core arcs, and then merges the copies of each physical vertex. In monomer variables the local merge sends $ab \mapsto z$, $a, b \mapsto 1$, $1 \mapsto 0$. Its stable symbol is $z + r + s$. Thus simultaneous use of both copies is removed rather than silently allowed. Positive activities enter by scaling before merging. Zero activities follow by continuity; the empty-support monomial prevents a zero-polynomial limit. No assertion that an arbitrary transversal matroid is stable is used.

To pass from monomer stability to support real-rootedness, diagonalize: $\mu(t, \dots, t) = t^n \Gamma(-t^{-2})$. The diagonal is real-rooted. A nonzero nonreal or nonnegative root of Γ would yield a forbidden nonreal root in this identity. The constant coefficient is 1, so zero is not a root of Γ . This also handles surviving-degree losses.

Corollary 9.8. *Any finite loopless directed relation whose bipartite role graph has matching number at most three has a rank-ULC support polynomial under arbitrary nonnegative independent role activities.*

Proof. By König's matching-cover theorem, choose a role-vertex cover of size at most three. If both shores occur, each contributes at most two vertices and Theorem 5.10 applies, even if their physical sets overlap. If just one shore occurs, there are at most three active tails or at most three active heads. Theorem 9.7 or its dual applies. Missing active vertices can be supplied as isolated, zero-activity placeholders. Lower surviving degrees are covered by Proposition 1.5. $\square$

[Merge note: the rank-ULC conclusion of this corollary is contained in source 11's Theorem 7.1, since the physical images of a role cover of size three cover every underlying edge. When both shores of the role cover occur, the proof gives real-rootedness through source 20, which is more. For bipartite relations with one role per vertex the corollary is *preorder-root-polytopes*, Theorem `mr:thm:rankthree` and Corollary `lc:cor:rankthree`.]

This corollary does not by itself prove Theorem 9.3: the bipartite role matching number may exceed three even when the physical matching number is three.

9.6 The finite theorem through seven vertices

Theorem 9.9 (Finite independent-role theorem). *Every preorder on at most seven vertices has a rank-ULC support polynomial under arbitrary nonnegative independent role activities, at its actual surviving degree.*

Since these orders have physical matching number at most three, only the second cubic gap requires new work. We describe both the exact finite domain and the scope of its certificates.

9.6.1 A complete enumeration rather than a sample

The equivalence relation $i \sim j$ defined by iRj and jRi partitions any preorder into equivalence classes. The quotient is a poset. Choose a linear extension and label its classes $1, \dots, m$. Every quotient arc then goes forward. Conversely, a transitive forward relation on m labels and a positive composition $n = n_1 + \dots + n_m$ determine a preorder on n vertices by expanding each class into n_i equivalent vertices.

The verifier enumerates every forward relation by finite bit masks, tests transitivity, and enumerates every positive composition. This is redundant, but completeness is immediate from the quotient construction. Isomorphisms and global order duality are then checked explicitly; weights are transported with their vertices, and duality exchanges the tail and head activity families.

For the connected, nonbipartite, nontotal, physical-rank-three seven-vertex domain there are

Quotient/composition configurations	131 645
Relevant configurations	75 160
Isomorphism/duality representatives	1 686

The entire catalog and all maps are retained. The independent replay regenerates the domain; it does not infer completeness from the number of files in a certificate directory.

9.6.2 Branches outside the residual catalog

Degree at most two is universal. Disconnected cases are handled by Lemma 9.5. A connected bipartite preorder with more than two vertices is a height-two poset: a directed path on three distinct vertices forces a triangle, and a nontrivial equivalence pair with a third neighbor also forces a triangle. Its physical and role matching numbers therefore agree. Corollary 9.8 applies.

A total preorder has complete underlying graph, so positive-activity degree three means $n = 6$ or 7 . The $32 + 64 = 96$ positive compositions reduce, by the extremal-block identities below and duality, to 24 role-weighted targets. Exact rational identities verify every target. For the nontotal six-vertex residual domain, the same quotient procedure gives 7 090 configurations, 3 059 relevant configurations, and 223 representatives. Their independent-role identities are all included.

9.6.3 Exact certificates and independent support reconstruction

Let w denote all independent role variables. A typical certificate for the second cubic gap is

$$p(w) := \gamma_2(w)^2 - 3\gamma_1(w)\gamma_3(w) = \sum_{\ell} q_{\ell} w^{\alpha_{\ell}} \left(\sum_{\beta} c_{\ell\beta} w^{\beta} \right)^2 + \sum_{\eta} r_{\eta} w^{\eta}, \quad (9.12)$$

where $q_{\ell} > 0$, $c_{\ell\beta} \in \mathbb{Q}$, $r_{\eta} \geq 0$, and all exponents are nonnegative integers. The remainder is reconstructed by exact subtraction and checked coefficientwise, rather than accepted from a numerical residual.

The independent checkers reconstruct the feasible supports by two complementary methods. The main hybrid replay and the two-attachment auditors enumerate physical directed matchings and deduplicate their ordered support pairs. The separate six-vertex and total-preorder routines use Hall's criterion

$$|N(A) \cap T| \geq |A| \quad \text{for every } A \subseteq S$$

for disjoint equal-sized sets (S, T) , with explicit bijection cross-checks. In either method, a feasible pair contributes its monomial once. In the vertex specialization, different pairs that specialize to the same monomial have their coefficients *added*. This is where losing ordered-support multiplicity would invalidate the computation. Matching enumeration is used only to construct the set of supports, never to retain the number of matching witnesses.

All arithmetic in the replay uses integers and rational numbers. Floating-point optimization proposed square directions during discovery; those optimization programs and their binaries are not part of the proof package or needed by the verifier.

9.6.4 Two structural reductions used in the selected ledger

The finite ledger uses several ordinary theorems already proved above, as well as the two-attachment theorem of Section 9.8. Two other reductions are worth stating explicitly.

A marked five-vertex deletion lemma. Let H be a preorder on five vertices with factor-critical underlying graph, and mark h . Write

$$Q = \Gamma_{H-h} = 1 + az + bz^2, \quad P = \Gamma_H - Q = cz + dz^2.$$

The finite role-weighted identity is

$$acd - bc^2 - d^2 \geq 0. \tag{9.13}$$

There are 96 marked isomorphism/duality types; 48 margins are coefficientwise nonnegative and 48 have rational square certificates. The collection contains 258 square terms and 6 971 positive remainder monomials. The independent enumeration also checks the factor-critical condition, rather than presupposing it from a label.

For positive activities $b, c, d > 0$ and Q has negative real roots by Proposition 1.5. Equation (9.13) says $Q(-c/d) \leq 0$, so the root of $c + dz$ lies between the roots of Q . Thus $(c + dz)/Q$ has nonnegative residues at its negative poles, and $P/Q = z(c + dz)/Q$ maps $\mathbb{H}$ into its closure. Repeated roots are handled by cancellation and limits.

Consequently the decomposition (9.6) proves real-rootedness whenever every block $C_i \cup \{h\}$ is a factor-critical preorder on three or five vertices. This is an all-rank gluing statement. In a seven-vertex factor-critical graph with a cut vertex h , the components after deleting h have even sizes $2 + 4$ or $2 + 2 + 2$. Each block is factor-critical: for $v \in C_i$, parity forces h to be matched into $C_i \setminus \{v\}$ in a perfect matching of $G - v$. Restricting that matching to $(C_i \setminus \{v\}) \cup \{h\}$ proves the required deletion property; deleting h itself leaves the perfectly matched C_i . Hence every such seven-vertex cut-vertex case is covered.

Refining small extremal equivalence classes. A minimal equivalence class has identical outgoing neighborhoods outside the class and no incoming outside arcs. For a class of size two, replacing equivalence by the chain $1 < 2$, keeping tails u_1, u_2 , and setting

$$v'_2 = v_2 + u_2 v_1 / u_1 \tag{9.14}$$

preserves the support polynomial. The first head activity in the chain is unused.

For a class of size three, sort the tail activities as $x \leq y \leq z$, carrying their head activities v_1, v_2, v_3 with them. Put

$$a = \sum_{i \neq j} u_i v_j, \quad c = yz v_1 + xz v_2 + xy v_3, \quad \Delta = z(x + y) - xy.$$

Replace the class by a strict chain with the same tails and effective heads

$$v'_2 = \frac{(x + y)c - xya}{x\Delta}, \quad v'_3 = \frac{za - c}{\Delta}. \quad (9.15)$$

These are nonnegative since

$$\begin{aligned} (x + y)c - xya &= y^2(z - x)v_1 + x^2(z - y)v_2, \\ za - c &= z^2(v_1 + v_2) + \Delta v_3, \end{aligned}$$

and $\Delta > 0$ for positive tails. They satisfy

$$xv'_2 + (x + y)v'_3 = a, \quad xzv'_2 + xyv'_3 = c.$$

These are precisely the local weights with one selected head and respectively one or two selected tails. Pure-tail states are unchanged. No other local head state is possible in a three-element minimal class. The surplus tails have identical outside neighborhoods, so equality of these local states preserves every continuation. Maximal classes are handled by duality. Disjoint chosen classes can be refined successively; each is refined once, and other classes' positive denominator activities are unchanged.

This transfer holds for any minimal or maximal class of size two or three, not just a boundary class of a total preorder. The full seven-vertex refinement manifest has 648 checked relational maps, each strictly reducing the number of equivalent pairs. Dependencies are therefore acyclic.

Remark 9.10 (Activity and dependency safeguards). The transformations (9.14)–(9.15) use strictly positive original denominator activities. They may create zero effective heads, which are permitted by the closed-orthant target theorems. Original zero activities are handled by Lemma 9.4, or by continuity of the original gap; no finite sorted-chain replacement at arbitrary zero denominators is asserted. The transformations generally destroy $u = v$. Every refinement here terminates in a proof valid for fully independent role activities. No vertex-only target is used. Strict decrease of equivalent pairs makes the dependency graph acyclic, and the verifier requires a proved terminal target.

9.6.5 The complete selected independent-role ledger

The final manifest partitions the 1686 residual targets as follows. Labels identify the proof selected, rather than every applicable proof.

Selected justification	Number
Articulation real-rootedness	168
Two-tail/two-head role-cover real-rootedness	478
Three-core role-ULC	28
Factor-critical cut-vertex real-rootedness	83
Direct rational role certificate	812
Two-attachment role-ULC	43
HPP side matroid by small-matroid classification	27
HPP side matroid by published nine-element list	6
HPP side matroid by the final exact Rayleigh identity	1
Fully role-justified extremal refinement	40
Total	1686

The HPP branches are proved in Section 9.7 below. All refinements end in unconditional role-valid proofs; there are no remaining conditional sources or independent obligations.

Four selected direct certificates, with catalog IDs 722, 736, 756, 758, use a positive multiplier. They certify (9.12) for $H(w)p(w)$ rather than $p(w)$, where

$$H = (u_4 + u_5)(v_0 + v_1) \quad \text{for 722, 736, 756,} \quad H = (u_5 + u_6)(v_0 + v_1) \quad \text{for 758.}$$

The exact multiplier is reconstructed and checked. It is strictly positive when all role variables are positive, so division proves $p \geq 0$ there; the original polynomial gap extends to the closed orthant by continuity. This is a declared positive-orthant argument and is distinct from the all-real Rayleigh identity in Section 9.7.3.

For reference, the six-vertex role certificates contain 586 square terms and 31 094 positive remainders. The 24 total-preorder role targets contain 793 squares and 50 660 positive remainders. The full local catalog, exact identities and all selected terminal witnesses are included; counts describe proof data rather than sampled assignments. With the HPP arguments below, this establishes Theorem 9.9.

9.7 Half-plane-property side matroids and the final cases

The final finite targets have a useful structural presentation. It leads to a general real-rootedness criterion, valid in every degree and without transitivity. The criterion is conditional on a matroid half-plane property; we verify that condition exactly in every application.

9.7.1 The one-tail side criterion

Theorem 9.11. *Let D be a finite loopless directed relation. Suppose a physical vertex p and a set Q satisfy*

$$i \rightarrow j \implies i = p \text{ or } j \in Q.$$

The vertex p may belong to Q . Form a transversal matroid M with centers Q , one ground element for every tail copy i with neighborhood $N(i) \cap Q$, and one private dummy element d_j with neighborhood $\{j\}$ for each $j \in Q$. If the basis-generating polynomial F_M is real stable, then the signed physical monomer polynomial of D is real stable for arbitrary nonnegative independent role activities. Hence Γ_D has only negative real zeros, with rank-ULC at its actual degree.

The private dummies are an independent basis, so M has rank $|Q|$. A basis is counted once, not once for each matching that witnesses it. For homogeneous real polynomials, the half-plane property (nonvanishing when all variables have positive real parts) is equivalent to real stability by a simultaneous rotation of variables. We use the abbreviation HPP.

Proof (pointer). Every head copy outside Q has no possible predecessor except p , so the hypothesis of source 15's Theorem 8.7 holds with the single hub p , and the matroid M is the matroid M_Q there. That theorem gives the conclusion. $\square$

[Merge note: source 22's own proof is source 15's proof in this special case: the split seed identity $\mu_B = (\prod_i a_i)(\prod_{j \in Q} v_j)F_M(d_j = b_j/v_j, t_i = -u_i/a_i)$ (source 15's (8.6)), the forced pendant insertion $f \mapsto b_x f - u_p v_x \partial_{a_p} f$ ((8.7)), the physical merge, here realized as the directional derivative $T_i f = (\partial_{a_i} + \partial_{b_i})f|_{a_i=b_i=z_i/2}$, and the diagonal root transfer. It is replaced by this pointer. Source 22 notes that successive pendant additions cannot reuse p , and that the evolving empty-support term keeps coefficient one through all merges.]

Reversing every arc and exchanging u with v gives the dual one-covered-head criterion.

9.7.2 Simplification and the published HPP input

An empty-neighborhood tail element is a loop. A singleton-neighborhood element with neighbors $\{j\}$ is parallel to the dummy d_j . These are the only parallel classes: two nonempty columns are a dependent pair exactly when their neighborhood union has size one. Repeated multi-neighbor columns must therefore *not* be merged.

The side simplification consists of one private representative for each center and one separate element for each tail whose neighborhood has size at least two. HPP lifts back from the simplification by replacing a private representative’s variable with the sum over its true parallel class; sums of upper-half-plane variables stay in that half-plane. Loops cause no difficulty.

We use two precisely bounded published inputs from Kummer–Sert [9, 10]:

1. The complete exceptions on at most seven elements are the five rank-three matroids $F_7, F_7^-, F_7^{--}, F_7^{-3}, M(K_4) + e$ and their rank-four duals. Every new eight-element excluded minor is rank-four sparse paving. Thus an eight-element matroid whose one-element deletions and contractions are HPP is HPP if its rank is not four or it is not sparse paving [9, Section 5, Theorem 5.2].
2. The published file `n9r4Hpp.txt` contains 4 125 certified-positive rank-four nine-element matroids. Section 6 of [9] explains their rational positive-semidefinite Gram certifications. The separate lists of candidates and undetected cases are not used.

We do not assume every small transversal matroid has HPP. In particular, the rank-four dual of F_7^{-3} is transversal and is an exception. Our higher-rank side matroids are checked against the full exception/minor tests, including duality.

The small-matroid side screen supplies 27 selected catalog targets. It checks 58 candidate presentations, including 47 eight-element sides and 752 one-element minors. Each witness stores the working relation, direction reversal, p, Q , every tail-neighborhood mask, retained simplified columns, and the complete basis set. The independent audit reconstructs these objects and checks exact matroid isomorphisms.

Six further targets use the published nine-element positive list. Its basis fingerprints use increasing colex order on four-element subsets, equivalently increasing subset bit masks; an asterisk marks a basis. Rank-five sides are dualized to rank four before comparison. The selected applications are

Catalog targets	One-based published line
1030, 1474, 1480	1975
1057, 1486	1536
1495	1751

The data retain explicit ground-element permutations, not just invariant matches. All 126 basis indicators are compared after the permutation. Both possible side presentations were independently recounted, giving twelve comparisons of complete side basis systems.

The pinned official file has SHA256

`a3c7a6eaebe02ef50998710282542be2751b218a6713853cdd2eecdabe1f1f86`.

[*Merge note:* the published file `n9r4Hpp.txt` of Kummer and Sert is distributed under CC BY 4.0 (Zenodo record 6108027) and is shipped unmodified, four times, inside source 22’s nested audit zip; the report README carries the attribution and licence notice.]

Its MD5 matches the publisher’s `c73d80267112a960df5485f159ab56e0`. The exact data file is included for offline membership replay. This is an application of the published certified-positive list, not a local rerun of all of its original Gram certificates.

9.7.3 The final ten-element side matroid

The final independent target, catalog ID 1565, has seven off-diagonal row masks

$$(0, 1, 3, 7, 39, 23, 63). \quad (9.16)$$

Choose $p = 6$ and $Q = \{0, 1, 2, 4, 5\}$ in this order. Its side simplification is the rank-five transversal matroid M on ten elements whose center-neighborhood masks are

$$(1, 2, 4, 8, 16, 3, 7, 23, 15, 31). \quad (9.17)$$

We index these elements $0, \dots, 9$. The first five private columns form a basis, and exact matching enumeration gives 198 bases in total.

All twenty one-element deletions and contractions have HPP. Loops, coloops, genuine parallel classes and series classes are reduced with exact basis checks. HPP is restored by the standard loop/coloop and parallel operations, with series reduction obtained by duality. After reduction, the dependencies group as follows; “small” means the previously stated at-most-eight criterion, and a number is a one-based line of the positive nine-element file, with dualization when the reduced rank is five.

Elements	Deletion	Contraction
0, 1, 5	3368	small
2, 6	1751	2020
3, 4, 7, 8	small	1536
9	small	2385

Eight minors use the small criterion and twelve use the published list. The independent replay verifies 10 240 minor rank-function identities, every reduction, and all explicit published-basis comparisons. These checks establish the proper-minor hypothesis only; one further all-real inequality is needed.

Let F be the basis polynomial of M , and write F_{ab} for the coefficient of $x_0^a x_9^b$, $a, b \in \{0, 1\}$. Its Rayleigh difference is

$$q = \Delta_{09}F = F_{10}F_{01} - F_{00}F_{11}. \quad (9.18)$$

It is independent of x_0, x_9 . In the remaining element order $1, \dots, 8$, it is homogeneous of degree eight and has 397 nonzero monomials. Let $\mathcal{B}$ be the sorted set of the 37 squarefree degree-four exponent vectors a for which the coefficient of x^{2a} in q is nonzero. Define

$$N_e = \#\{(a, b) \in \mathcal{B}^2 : a + b = e\}, \quad G_{ab} = \frac{[x^{a+b}]q}{N_{a+b}}. \quad (9.19)$$

The exact support check gives $\text{supp}(q) \subseteq \mathcal{B} + \mathcal{B}$. With $h = (x^a)_{a \in \mathcal{B}}$, the identity $q = h^T G h$ therefore follows by collecting coefficients. The matrix entries are only

$$0, \quad \frac{1}{8}, \quad \frac{1}{6}, \quad \frac{1}{4}, \quad \frac{1}{3}, \quad \frac{1}{2}, \quad 1.$$

Lemma 9.12 (Exact final Gram certificate). *The rational matrix in (9.19) is positive semidefinite of rank 32. Equivalently, the supplied exact vectors $c_1, \dots, c_{32} \in \mathbb{Q}^{37}$ and positive rational weights $w_1, \dots, w_{32}$ satisfy*

$$G = \sum_{\ell=1}^{32} w_\ell c_\ell c_\ell^T, \quad q = \sum_{\ell=1}^{32} w_\ell (c_\ell^T h)^2. \quad (9.20)$$

In particular q is nonnegative on all of $\mathbb{R}^8$.

Exact certificate verification. The supplied complementary standard-library checker reconstructs all 198 bases by explicit assignment to the five centers, forms the four coefficient polynomials F_{ab} , and expands (9.18). It independently reconstructs $\mathcal{B}$, every multiplicity N_e and the uniform Gram formula. Exact expansion of both identities in (9.20) has zero remainder. Finally it performs rational Schur elimination, obtaining 32 strictly positive pivots and a zero residual matrix, independently confirming the rank and positive semidefiniteness. All vectors and weights are included in the machine-readable certificate. The independent audit separately reconstructs the full ten-variable derivatives, verifies both the Gram expansion and the positive-square factorization, and confirms rank 32 by exact row elimination.

This is an all-real sum of ordinary polynomial squares. There is no uncontrolled-sign monomial prefactor and no merely coefficientwise-positive remainder. Thus it supplies precisely the all-real domain required below. Numerical projection found the matrix, but no floating-point output or tolerance is used in its exact verification. $\square$

Wagner–Wei’s criterion [16, Theorem 3(c)] says that a multiaffine polynomial with positive coefficients is stable if every one-variable deletion and contraction is strongly Rayleigh and one Rayleigh difference is nonnegative on all real values of the remaining variables. For homogeneous matroid basis polynomials, HPP, real stability and the strong Rayleigh property are equivalent. The twenty proper minors and Lemma 9.12 satisfy these hypotheses. Hence M has HPP, and Theorem 9.11 proves real-rootedness for target 1565 under every nonnegative independent role assignment.

The existing approved extremal-class refinement maps the last conditional source, ID 1611, to this target. Effective zero activities are permitted by Theorem 9.11. This closes the final two entries in the selected finite ledger.

9.8 Two attachments and arbitrary exterior populations

The following theorem supplies the two-attachment infinite-family input to the global reduction.

Theorem 9.13. *Let a preorder of physical matching number at most three have a retained core K of at most five vertices, with two marked attachments i, j , and an independent exterior whose neighbors lie in $\{i, j\}$. Suppose each attachment has a fixed direction toward all of its exterior neighbors and no exterior vertex belongs to a nontrivial equivalence class. In the canonical Gallai–Edmonds core family, the support polynomial is rank-ULC under arbitrary nonnegative independent role activities and arbitrary heterogeneous exterior populations.*

The canonical-core qualification fixes the finite catalog used in the proof. We do not replace it by an unchecked assertion about arbitrary marked five-vertex directed cores. The catalog has 1 084 structural templates, including harmless overcoverage. Its generation and coverage are independently checked in the included two-attachment dependency. Degree at most two and the first cubic gap are settled by Proposition 1.5; below we prove the second cubic gap.

9.8.1 Opposite signs: exact two-vertex compression

Assume exterior arcs at i point outward and those at j inward. The exterior types are head-only neighbors of i , tail-only neighbors of j , and shared vertices $i \rightarrow s \rightarrow j$. Put

$$H = \sum_{s: i \rightarrow s} v_s, \quad T = \sum_{s: s \rightarrow j} u_s, \quad W = \sum_{s: i \rightarrow s \rightarrow j} u_s v_s.$$

The exact support identity is

$$\begin{aligned} \Gamma_R = \Gamma_K + T v_j z \Gamma_{K-j} + H u_i z \Gamma_{K-i} \\ + (HT - W) u_i v_j z^2 \Gamma_{K-\{i,j\}}. \end{aligned} \tag{9.21}$$

A selected exterior tail must be matched to j and a selected exterior head to i . Two selected exterior vertices must have distinct physical identities; $HT - W$ removes exactly the forbidden shared-vertex collisions. Removing their forced matched endpoints leaves a core support. This proves (9.21) without counting witnessing matchings.

We have $0 \leq W \leq HT$. If the shared type is legal, so are both private types: after deleting either its incoming or its outgoing incidence, no two-step transitivity requirement through that exterior vertex remains, and core closure of the retained neighborhood is inherited. Legality is also checked directly in the finite catalog. For $T > 0$, use one shared vertex with

$$u_s = T, \quad v_s = W/T,$$

and one private head at i of activity $H - W/T$. This preserves H, T, W and hence the whole polynomial. If $T = 0$, then $W = 0$ and only the head sum remains. If the shared type is not legal, sum the two private clouds separately. The replacement uses at most two exterior vertices, so has at most seven vertices.

Crucially, this compression need not preserve $u = v$. Its finite targets are proved under the stronger role weighting, independently of the global assembly. The finite certificate targets are replayed directly, so this use creates no circular dependence on the two-attachment branch of the selected ledger. Among the 494 mixed-sign templates, the checked compressed branches are

At most six vertices	366
Physical matching rank at most two	19
Disconnected	49
Connected bipartite	3
Residual seven-vertex references	57

The last 57 references use 56 distinct direct role certificates, with 1 135 square terms and 6 445 positive remainders. Explicit relabeling/duality witnesses and all compression identities are included. The three bipartite cases can use Corollary 9.8; no higher-rank bipartite theorem is required here.

9.8.2 Equal signs: the exact moment cone

By duality suppose both attachments point to exterior heads. Let A, B be the private head-activity sums at i, j . For the shared cloud define

$$U = \sum_s v_s, \quad E = \sum_{s < t} v_s v_t.$$

Put

$$F_i = zu_i \Gamma_{K-i}, \quad F_j = zu_j \Gamma_{K-j}, \quad G = z^2 u_i u_j \Gamma_{K-\{i,j\}}.$$

Let F_* be the contribution of supports using one newly adjoined common head, whose activity is set to 1. This is a genuine support kernel: if either attachment can use the same head, a final support is still counted once. In general $F_* \neq F_i + F_j$.

There are at most two exterior vertices in a support, and direct counting gives

$$\Gamma_R = \Gamma_K + AF_i + BF_j + UF_* + [AB + (A + B)U + E]G. \quad (9.22)$$

Two heads from one private cloud cannot be used. The remaining pairs of types contribute AB, AU, BU, E , each at support multiplicity one. With two exterior heads selected, the two attachments are used, leaving exactly the core support in $K - \{i, j\}$.

The moment range satisfies

$$U \geq 0, \quad 0 \leq E \leq U^2/2,$$

because $2E = U^2 - \sum_s v_s^2$. The whole closed cone is parameterized by

$$U = \xi + \eta, \quad E = 2\xi\eta, \quad \xi, \eta \geq 0. \quad (9.23)$$

Indeed the inverse values are $(U \pm \sqrt{U^2 - 2E})/2$. This is a positivity reduction on a cone, not an assertion that two ordinary heads realize every shared cloud. In the formal support generator, two common heads of weights ξ, η are used and a support selecting both is assigned coefficient 2. That formal convention produces exactly (9.22) after (9.23); it is not confused with an ordinary two-clone preorder.

If the common type is not legal, at most one private type is legal: a head's incoming neighborhood must be predecessor-closed in the core, and the union of two such legal neighborhoods is again predecessor-closed. Therefore the private-only cases compress to at most six vertices.

There are 590 same-sign catalog templates: 333 private-only reductions and 257 shared-cone cases. Every shared-cone substituted cubic gap has an exact certificate of the form (9.12), in the core role activities and A, B, ξ, η . Together these contain 7 866 squares and 32 729 positive remainder monomials. The verifier independently constructs F_* from deduplicated ordered supports, checks the symbolic moment identity, tests heterogeneous and zero-face substitutions, and checks every rational identity. A separate local Hall-based reconstruction provides an additional check. No unfinished chamber split or numerical multiplier is needed for these 257 certificates.

9.8.3 Degree and zero-activity safeguards

Both formulas preserve the original weighted polynomial, not merely its first few coefficients. Effective activities can be zero; all target polynomial inequalities hold on the closed orthant. Original degree at most two is handled before applying a cubic certificate, with its factor 4. At original degree three, equality of support polynomials preserves that degree. Thus the argument never deduces degree-two rank-ULC from a degree-three zero-padding convention. This completes Theorem 9.13.

9.9 The Gallai–Edmonds reduction and the global proof

We now pass from bounded cores to arbitrarily many vertices. The graph-theoretic input is the classical Gallai–Edmonds structure theorem; an accessible primary proof is West [17, Theorem 5].

For a graph G , let D be the vertices omitted by some maximum matching, let $A = N(D) \setminus D$, and let $C = V(G) \setminus (A \cup D)$. Each component of $G[D]$ is factor-critical, $G[C]$ has a perfect matching, and every maximum matching pairs every vertex of A into a distinct component of D , with near-perfect matchings inside those components. Hence

$$\nu(G) = |A| + \frac{|C|}{2} + \sum_{D_t} \frac{|D_t| - 1}{2}. \quad (9.24)$$

9.9.1 The canonical exterior and its size bound

Assume positive activities and $\nu(G_R) = 3$. Let I be the union of the singleton components of $G_R[D]$ and let $K = V \setminus I$. Distinct vertices of I are nonadjacent and all their neighbors lie in A . Write $a = |A|$, $|C| = 2c$, and $|D_t| = 2d_t + 1$ for the nonsingleton components, with $d_t \geq 1$. Then

$$a + c + \sum_t d_t = 3, \quad |K| = a + 2c + \sum_t (2d_t + 1) \leq a + 3 \left(c + \sum_t d_t \right) = 9 - 2a.$$

Thus $a \leq 3$, with core bounds 7, 5, 3 when $a = 1, 2, 3$.

The preorder adds two essential restrictions. Equivalent vertices are adjacent true twins, so their transposition is a graph automorphism. The canonical sets D, A, C are invariant under every automorphism. An exterior vertex in a singleton D component therefore cannot be equivalent to another vertex: such a vertex would also lie in D and be adjacent to it. In particular, exterior-to-core arcs are not bidirectional.

At each attachment, all exterior arcs have one direction. Otherwise $x \rightarrow h \rightarrow y$ with distinct $x, y \in I$ would make two exterior vertices adjacent by transitivity. The same exterior vertex cannot create the exception because bidirectionality has just been excluded. Each exterior type is thus determined by its nonempty neighborhood subset of A and the fixed signs of those attachments. Isolated exterior points have no effect.

9.9.2 The four attachment cases

$a = 0$. There are no attachments. Every nontrivial component is either a factor-critical D component on at most seven vertices or a component in C , whose total order is at most six. Apply Theorem 9.9 to each and Lemma 9.5 to their product.

$a = 1$. All nonisolated exterior vertices are equally oriented pendant twins at the sole attachment h . At most one occurs in a support. Sum their relevant activities into S . Replacing them by one pendant preserves the support polynomial. Assign the new pendant its relevant role activity S and any positive activity in its unused role. Core activities are unchanged. The full independent-role polynomial is preserved; in the vertex specialization both new role activities may be set to S .

If $|K| \leq 6$, the result has at most seven vertices and Theorem 9.9 applies. If $|K| = 7$, equality in the budget bound forces $c = 0$ and precisely two nonsingleton factor-critical D components, each with $d_t = 1$ and hence three vertices. Deleting h leaves these two rank-one components and the isolated exterior points. Theorem 9.6 applies directly to every population. If the exterior is empty, the seven-vertex finite theorem is already sufficient.

$a = 2$. The core has at most five vertices and the canonical exterior has only the three possible nonempty neighborhood types. All hypotheses of Theorem 9.13 hold. Its independent-role conclusion applies directly.

$a = 3$. Equation (9.24) forces $K = A$, with three attachments and no other core vertices. Every attachment has an exterior neighbor, since $A = N(D) \setminus D$ and now every D component is a singleton. Split $A = P \sqcup Q$, with exterior arcs outward at P and inward at Q .

There is no core arc $q \rightarrow p$ with $q \in Q$ and $p \in P$. Choose exterior neighbors $x \rightarrow q$ and $p \rightarrow y$. If $x \neq y$, the path $x \rightarrow q \rightarrow p \rightarrow y$ contradicts independence of I . If $x = y$, the directed cycle makes that exterior vertex equivalent to a core vertex, contradicting the singleton-equivalence property above. Every core arc is therefore covered by its tail in P or its head in Q : arcs within P are covered on the tail side, arcs within Q on the head side, and $P \rightarrow Q$ on both. The exterior arcs are covered too.

The role graph consequently has a cover of size three, and Corollary 9.8 proves the result. This gives an ordinary conceptual replacement for following all seventeen three-attachment classes individually. The previously checked catalog of nine same-sign and eight mixed-sign classes remains in the supplementary dependency record.

These cases exhaust the Gallai–Edmonds decomposition. Together with Proposition 1.5 and Lemma 9.4, they prove Theorem 9.3, including arbitrary vertex count and all zero-activity

specializations at surviving degree.

Corollary 9.14 (Vertex activities; archive 31). *Every finite preorder with nonnegative vertex activities x_i , that is $u_i = v_i = x_i$, whose support polynomial has actual degree at most three, is rank-ULC at that degree.*

Proof. The case $u_i = v_i = x_i$ of Theorem 9.3. □

[Merge note: this is the main theorem of archive 31, the earlier, vertex-activity version of this manuscript (same skeleton; eleven proof notes and the role-cover dependency byte-identical). Archive 31 closed its finite seven-vertex domain with a selected ledger of 97 direct vertex-only certificates (100 shipped) after the specialization $u = v$; that ledger is redundant given the role ledger above and survives only in the arrival commit (README). Source 23 (Section 4) uses exactly this corollary. With $x_i = 1$ it is source 25's Theorem 2.3, whose proof in Part I is an independent route.]

9.10 A cubic with nonreal zeros

Source 22 computes the fifteen-vertex example of Section 2.9 (three lower vertices, twelve upper vertices, four of each pair-neighborhood type; all activities one) with the same counts, $\Gamma(z) = 1 + 24z + 162z^2 + 208z^3$, discriminant -2592 , gaps 90 and 11268; that computation is printed once, in Part I.

It is rank-ULC, consistently with the theorem. This example requires no independent-role asymmetry and lies in the pure three-tail family of Section 9.4. No minimal-size claim is made.

[Merge note: source 25's Theorem 2.9 makes the minimality claim: fifteen vertices is the least order of a preorder of actual degree three with a nonreal-rooted gamma polynomial, and this example and its dual are the only ones at that order.]

9.11 Verification package and dependency closure

The deliverable contains this TeX source and PDF, a portable exact checker entry point, frozen certificate archives, the companion role-cover paper and its checker package, and proof notes recording the precise scopes of structural inputs. All required data are local. The full replay uses Python 3 and a C++17 compiler for supplementary role-cover support checks; it needs no network connection, numerical solver, precompiled binary, or third-party Python library.

[Merge note: in this report source 22's package is shipped renamed: its proof notes at the top of the report directory, its checkers in `code/`, its nested audit package as `data/22-independent-role-checker-role-degree-three-audit-package.zip`. Its embedded copies of source 20's package and article are not shipped (source 20's files are). Because `verify.py` first checks every listed file, the delivered wrapper runs only on the reconstructed package (README). At intake its sub-checkers passed on copies: the final Rayleigh certificate (198 bases, rank 32), the coverage and aggregate phases of the complete hybrid ledger, the 39 role certificates new relative to archive 31, the two-attachment and marked-five modules and the side-1565 package; the whole wrapper was not run in one process.]

9.11.1 How to run the check

With Python 3 and a C++17 compiler available, from any working directory run

```
python /path/to/package/verify.py
```

The wrapper verifies pinned dependency archives, extracts them into temporary directories, and runs their exact checkers. It verifies the complete role catalog, its rational certificates and multipliers, every structural witness and refinement terminal, all 223 six-vertex role certificates and all 24 total-preorder targets. Both two-attachment certificate modules and the marked-five deletion/gluing module are replayed as well. The HPP modules verify the side-matroid bases, explicit published-list isomorphisms, proper-minor reductions and final all-real Gram identity. The companion role-cover algebra and the nonreal example are checked separately.

The command writes a fresh local replay receipt while preserving the original approval receipts. A separate build script compiles the paper with a standard LaTeX installation; the mathematical checker does not require LaTeX. Python checks use the standard library, and the companion role-cover checker compiles included exact C++ sources into temporary storage.

9.11.2 What is checked exactly

The independent checkers reconstruct support polynomials from relations, expand all rational squares, and compare coefficients exactly. They verify template generation, structural witnesses, maps under relabeling and duality, population-compression formulas, moment substitutions, and terminal dependency scope. Finite point evaluations supplement these symbolic checks but do not prove positivity.

Ordinary theorems such as Gallai–Edmonds and Wagner’s rank-three Rayleigh theorem are cited mathematical inputs, not re-proved by running a Python script. The support-level structural proofs are given in this paper and the included companion. The certificate and coverage layers are independently audited, with their frozen source hashes recorded. These distinctions make the boundary of the computer-assisted proof explicit.

9.11.3 Dependency closure and published data

The selected ledger is complete for independent roles. Each extremal refinement has a proved terminal; positive-multiplier division is confined to the positive orthant and followed by continuity. The final Rayleigh certificate instead has an exact sum of squares valid on all real inputs. These different domains are never interchanged.

The classification results of Kummer–Sert and the certified-positive matroid list are cited mathematical dependencies. The local replay verifies membership in the published list by exact basis isomorphism, with the official file hash pinned. It does not purport to rerun the authors’ original Gram certificates for every published entry. The new ten-element case has its own fully local rational Gram and square identity. Ordinary results such as Wagner–Wei’s criterion and stability preservation under directional differentiation are proved or cited, not inferred from test points.

Some provenance notes retain draft-status wording from the time their mathematical text was frozen. The approval receipts and complete selected ledger specify the final scope. Numerical discovery code is excluded. The earlier vertex-only release is a separate valid specialization and is not a proof dependency of this stronger theorem.

9.12 Further questions and mathematical boundaries

Beyond degree three. The independent-role extension in degree three is now proved. Degree four requires two interior coefficient comparisons beyond the universal first inequality and has larger retained cores. An unweighted theorem does not imply its weighted counterpart; separate role-sensitive inequalities or exact certificates are needed. The HPP side-matroid criterion applies in every degree, but it is only a sufficient structural condition, not a general solution.

A conceptual replacement for the finite certificates. The Rayleigh identity (6.7) and the upper-half-plane articulation proof are short structural explanations. It would be useful to find similarly uniform inequalities for the remaining two-connected factor-critical cores, eliminating some or all of the finite square certificates. The marked five-vertex deletion inequality suggests that interlacing under deletion may be stronger and more useful than ULC alone in certain small cores.

Real-rootedness versus rank-ULC. The example in Section 9.10 rules out general real-rootedness already in degree three. A useful separate problem is to characterize the preorder or role-cover structures that force real-rootedness for every positive activity assignment. The articulation and two-tail/two-head cover theorems give substantial sufficient families but no general characterization.

Equality and higher degree. Classifying equality in the second cubic gap could clarify why weighted positivity is robust in some structures and difficult in others. The square identity identifies several simultaneous equality requirements in the pure three-tail case. Extending arbitrary independent-role-weighted rank-ULC to degree four or higher requires additional inequalities and is not implied by this paper. Even a complete unweighted higher-degree theorem would not justify the weighted extension without a new argument.

[*Merge note:* in this report, unit-activity degree four is settled (Part I), real-rootedness is forced by the role-cover and articulation theorems and by source 15’s pendant criterion in their families, and Part III settles weighted degree four for four-vertex cores with universal sinks and for the five-vertex boundary. Weighted degree four in general remains open.]

Attribution. The first-gap method is an application of Motzkin–Straus; the bounded-core reduction uses Gallai–Edmonds; the pure three-tail proof applies Wagner’s rank-three Rayleigh theorem; the role-cover companion uses classical real-stability preservers; the side-matroid applications use Kummer–Sert’s published classification and certified data; and the final ten-element case applies Wagner–Wei’s criterion. The new exact artifacts establish the statements and scopes recorded here. We make no claim that a literature-wide novelty search or a proof-assistant formalization has been completed.

Appendix

A Provenance and merge record

A.1 The archives

All eighteen archives arrived in ProveIt commit `f8c3a392a` on 1 October 2026 and were placed in commit `8f5c53106` (batch 73, cluster C2); archives 11, 12 and 15 were handed over to this cluster by the cluster that placed batch-73 manuscript 26 in *matching-rank-normalization*. Only sources 16 and 25 cite a ProveIt commit: `d5e863bba` (*preorder-root-polytopes* at “Write batch 70 (4/5)”); the labels they cite are unchanged since. All other 40-digit hexadecimal strings in the packages are SHA-256 digests of package files. Every package’s checksum ledger verified at intake.

Src.	Archive(s)	Manuscript (pages as delivered)	Pin	Printed
25	preorder-gamma-degree-three-package	Rank ultra log concavity for preorder gamma polynomials of degree at most three (14)	d5e863bba	Section 2
16	archives 16–19, preorder-degree-four-part- k -of-4, $k = 4, 3, 2, 1$	Degree four preorder support polynomials (17); the base	d5e863bba	Section 3
23	template26-structural-supplement	A smaller proof of the last Newton gap for template 26 (6)	—	Section 4
20	role-cover-package	Real stable monomers under a two by two role cover (9)	—	Section 5
12	three-core-rayleigh-result	Coefficientwise Rayleigh inequalities for a three vertex shore (6)	—	Section 6
11	physical-cover-three-result	Rank ultra log concavity under a three vertex physical cover (5)	—	Section 7
15	bipartite-and-pendant-support-result	Lorentzian and real stable directed support polynomials (6)	—	Section 8
22	independent-role-degree-three-package	Independently weighted preorder support polynomials of actual degree at most three (21)	—	Section 9
14	universal-sink-last-gap-package	Sharp final coefficient bounds for directed cores with universal sinks (4)	—	Section in Part III
10	universal-sink-all-degrees-package	Rank ultra log concavity for four vertex cores with universal sinks (5)	—	Section in Part III
34	universal-sink-five-core-package	Rank five ultra log concavity for five vertex cores with universal sinks (6)	—	Section in Part III
06	archives 07 and 06, universal-sink-five-core-boundary-package-part-1, -part-2	The rank four boundary for five vertex cores with universal sinks (7)	—	Section in Part III
31	weighted-degree-three-package	Vertex-weighted preorder support polynomials of actual degree at most three (17)	—	superseded by 22
13	universal-sink-degree-four-package	Weighted degree four rank ultra log concavity for universal sink extensions (5)	—	superseded by 10

A.2 What each source contributed

- 25: unit-activity rank-ULC through actual degree three; the bounded-core normal form at every degree; the minimum order fifteen of a nonreal-rooted degree-three gamma polynomial and its uniqueness; the ordinal-sum family. Its first inequality is *preorder-root-polytopes*'s.
- 16 (base): unit-activity rank-ULC at actual degree four; the exterior-only Lorentzian construction; the outward-star last gap; the kernel formula and coverage, the bounded branches and the integer-population certificate schema. Its Section 3 and Appendix A are source 20's article (printed from source 20); its first inequality is *preorder-root-polytopes*'s.
- 23: a second, smaller route for the last gap of template 26.
- 20: real stability of the signed monomer polynomial under every two by two role cover, with independent activities; the free-product corollary; the gamma transfer. It generalizes

matching-rank-normalization, Theorem `mrn:w:sb:thm:main` (the bipartite case with disjoint cover sides) by a different proof.

- 12: the coefficientwise three-core boundary inequality (computer-assisted), the full coefficient-wise Rayleigh property, negative correlation, and a four-core obstruction.
- 11: rank-ULC under every physical vertex cover of size at most three, with arbitrary internal core arcs; bipartite strictness and equality.
- 15: the bipartite physical Lorentzian theorem, overlapping role covers with an independent core, the pendant-head stability criterion and its elementary seed.
- 22: rank-ULC of every finite preorder of actual degree at most three with independent role activities; the articulation theorem; the zero-activity role filter; the half-plane-property side-matroid criterion and the final rank-32 Gram certificate. Its three-core and one-tail theorems are special cases of sources 11 and 15. It supersedes archive 31.
- 14: the sharp last-coefficient bound $K(r, m)$ for every core size with universal sinks.
- 10: rank-ULC at every actual degree for four-vertex cores with universal sinks, through the certified inequality $\gamma_2^2 \geq 3\gamma_1\gamma_3$. It supersedes archive 13.
- 34: rank-ULC at actual degree five for five-vertex cores with universal sinks, and a counterexample to the constant three there.
- 06: the actual-degree-four boundary of the five-vertex family, and the strengthened ordinary core comparisons.

A.3 Where the merge had to choose

1. *One report, three Parts.* All sources answer *preorder-root-polytopes*'s Research question `mr:q:gamma`; they would be additions to *preorder-root-polytopes*, which already has thirteen Parts and whose batch-72 continuation was already split off into *matching-rank-normalization*. They form one new report beside *matching-rank-normalization*.
2. *Base.* Source 16, the farthest-reaching unit-activity result. Its Section 3 and Appendix A are printed once, from source 20, in Part II; source 16's different operator proofs are kept there as second routes (Section 3.3).
3. *Common definitions and the first inequality* are printed once (Section 1.3); the sources' copies are pointers.
4. *Same theorem, two sources.* Source 25's theorem is the unit case of source 22's: both proofs are printed, in Parts I and II. Source 23 re-proves one entry of source 16's ledger: both routes are described, source 23's printed. Source 14's sharp bound contains the last inequalities of sources 10 and 34 (and of archive 13): it is printed once, and those proofs, which are the same route at $r = 4, 5$, are pointers. Sources 11 and 15 contain source 22's three-core and one-tail theorems: source 22's three-core proof, an ordinary argument through Wagner's rank-three theorem, is kept as a second route; its one-tail proof is the pendant proof of source 15 and is a pointer. The three-squares identity of sources 11, 12 and 22 is printed once (in source 12's section); the moment envelope of sources 10, 34 and 06 once (in source 10's); the core comparisons of sources 34 and 06 once, in source 06's stronger form; the nonreal cubic of sources 16, 22, 25 and archive 31 once (in source 25's).

5. *Superseded versions.* Archive 31 (vertex activities) is an earlier version of source 22: same skeleton, eleven byte-identical proof notes; its theorem is printed as the case $u_i = v_i$ of source 22's. Its only own material, a ledger of vertex-only certificates for a finite special case, survives in the arrival commit. Archive 13 is an earlier version of source 10 with a weaker middle inequality, $4\gamma_2^2 \geq 9\gamma_1\gamma_3$, certified by binomial squares only, and an alternative first-gap argument valid in degree four; it survives in the arrival commit.
6. *Source numbers* of multi-archive packages are their lowest archive numbers (16 and 06), as are their file prefixes.
7. *Not shipped* (README): embedded exact copies of other archives, PDFs, delivery READMEs, checksum ledgers, rendered page images, build logs and scripts of the delivered PDFs, the files of source 16 larger than 2 MB, source 06's two bounded-sink certificate archives and their manifest, and the regenerable outputs listed in the README. Source 34's 9,610 certificate files are shipped as one derived container whose members are byte-identical to the delivered files.

A.4 Shipped files

Each source's code, data, audits and proof notes are shipped as delivered, renamed with its prefix (25-degree-three-, 16-degree-four-, 23-template26-, 20-role-cover-, 12-three-core-rayleigh-, 11-physical-cover-, 15-pendant-support-, 22-independent-role-, 14-sink-last-gap-, 10-sink-all-degrees-, 34-five-core-, 06-five-core-boundary-): code in `code/`, data and records in `data/`, audits and proof notes at the top of the report directory. The report README lists them, maps the delivered names, lists what is not shipped and how to retrieve it from the arrival commit, and gives rerun instructions that work on copies. The text of delivered files still uses delivered paths.

References

- [1] ProveIt project report, *A Reflexive Root-Polytope Model for Preorder h -Polynomials* (*preorder-root-polytopes*), cited by sources 16 and 25 at the pinned source revision `d5e863bba6b40738af292b45005a0cf0495903e4`, Parts IV and XII. <https://github.com/VladimirReshetnikov/ProveIt/blob/d5e863bba6b40738af292b45005a0cf0495903e4/SetTheory/Cardinals/docs/reports/enumerative-combinatorics/preorder-root-polytopes/article.tex>. Current version: `SetTheory/Cardinals/docs/reports/enumerative-combinatorics/preorder-root-polytopes/`.
- [2] C. A. Athanasiadis and F. Chapoton, *Polytopes and posets associated to preorders*, arXiv:2605.26916v1 (2026), Section 5 and Proposition 9.2. <https://arxiv.org/abs/2605.26916>. (Source 16 lists the authors as F. Chapoton and C. A. Athanasiadis.)
- [3] C. A. Athanasiadis, Q. Xiao, and X. Yan, *Lattice point enumeration of some arbor polytopes*, arXiv:2603.11654v1 (2026), Remark 2.3. <https://arxiv.org/abs/2603.11654>.
- [4] P. Alexandersson and A. Jal, *Rook matroids and log-concavity of P -Eulerian polynomials*, arXiv:2410.00127v3, June 30, 2026.
- [5] J. Borcea and P. Brändén, *The Lee–Yang and Pólya–Schur programs. I. Linear operators preserving stability*, *Inventiones Mathematicae* **177** (2009), 541–569. <https://arxiv.org/abs/0809.0401>.
- [6] P. Brändén and J. Huh, *Lorentzian polynomials*, *Annals of Mathematics* **192** (2020), 821–891. doi:10.4007/annals.2020.192.3.4; arXiv:1902.03719.
- [7] P. Brändén, *Polynomials with the half-plane property and matroid theory*, *Advances in Mathematics* **216** (2007), no. 1, 302–320. <https://arxiv.org/abs/math/0605678>.

- [8] H. Crapo and W. Schmitt, *The free product of matroids*, European Journal of Combinatorics **26** (2005), no. 7, 1060–1065. <https://arxiv.org/abs/math/0409080>.
- [9] M. Kummer and B. Sert, *Matroids on Eight Elements with the Half-plane Property and Related Concepts*, arXiv:2111.09610v4, Sections 5 and 6, especially Theorem 5.2. <https://arxiv.org/abs/2111.09610>.
- [10] M. Kummer and B. Sert, supplementary matroid data, Zenodo record 6108027 (CC BY 4.0). Certified-positive nine-element list `n9r4Hpp.txt`. <https://zenodo.org/records/6108027>.
- [11] T. S. Motzkin and E. G. Straus, *Maxima for graphs and a new proof of a theorem of Turán*, Canadian Journal of Mathematics **17** (1965), 533–540, Theorem 1. doi:10.4153/CJM-1965-053-6.
- [12] J. Ross, H. Süß and T. Wannerer, *Dually Lorentzian polynomials*, Monatshefte für Mathematik **208** (2025), 495–524. <https://doi.org/10.1007/s00605-025-02134-6>.
- [13] F. Röhrle and M. Ulirsch, *Logarithmic concavity of bimatroids*, Annals of Combinatorics **30** (2026), 501–523. Theorem A and Example 2.4. <https://doi.org/10.1007/s00026-025-00780-z>; <https://arxiv.org/abs/2402.15317>.
- [14] D. G. Wagner, *Rank-three matroids are Rayleigh*, Electronic Journal of Combinatorics **12** (2005), Note N8, Theorem 1.1. doi:10.37236/1975; <https://arxiv.org/abs/math/0403216>. (Source 12 cites the 2004 arXiv version as a background reference only: its proof of the boundary theorem uses the directed-support classification.)
- [15] D. G. Wagner, *Multivariate stable polynomials: theory and applications*, Bulletin of the American Mathematical Society **48** (2011), 53–84. <https://arxiv.org/abs/0911.3569>; https://www.math.uwaterloo.ca/~dgwagner/ceb_wagner.pdf.
- [16] D. G. Wagner and Y. Wei, *A criterion for the half-plane property*, arXiv:0709.1269 (2007), Theorem 3. <https://arxiv.org/abs/0709.1269>.
- [17] D. B. West, *A short proof of the Berge–Tutte Formula and the Gallai–Edmonds Structure Theorem*, author manuscript, Theorem 5. <https://dwest.web.illinois.edu/pubs/galledm.pdf>.