

Derivative Norm Spectra and Dual Moment Geometries of the Fabius–Rvachev System

**Thue–Morse equimeasurability, inverse-Fabius beta transforms,
Gamma–Bell operators, and shifted Stieltjes–Wigert energy measures**

Research report prepared for Vladimir Reshetnikov’s ProveIt development

27 August 2026

Recursive documentation audit pinned to commit 0341bbf91cb867f4510bc8431cf59878208de771

Abstract

Let F be the bounded Fabius distribution function, $G = F^{-1}$ its inverse on $[0, 1]$, $\text{up}(x) = 1 - F(|x|)$ on $[-1, 1]$, and

$$\Phi(\xi) = \widehat{\text{up}}(\xi) = \prod_{j=0}^{\infty} \text{sinc}\left(\frac{\pi\xi}{2^j}\right)$$

under the Fourier convention $\widehat{f}(\xi) = \int_{\mathbb{R}} f(x) e^{-2\pi i x \xi} dx$. The recursive documentation tree of the ProveIt Fabius development was audited at commit 0341bbf91cb867f4510bc8431cf59878208de771. Existing reports already contain the dyadic derivative tiling, integer-zero arithmetic, sharp Fourier-decay regimes, endpoint Lambert expansions, q -binomial acceleration, inverse-quantile reversion, and several spectral determinant constructions. Those results are treated as baseline rather than renamed.

The first new package extracted here begins with an exact equimeasurability theorem. If $T_n = n(n+1)/2$, then $2^{-T_n} |\text{up}^{(n)}|$ has exactly the same value distribution on $[-1, 1]$ as up . The signs on the 2^n cells are the length- 2^n Thue–Morse block. Consequently every rearrangement-invariant norm, including every L^p quasi-norm $0 < p \leq \infty$, obeys

$$\left\| \text{up}^{(n)} \right\|_p = 2^{T_n} \|\text{up}\|_p.$$

The level sets are also exact:

$$\left| \{x : |\text{up}^{(n)}(x)| > 2^{T_n} t\} \right| = 2G(1-t), \quad 0 < t < 1.$$

Thus the complete inverse Fabius function is encoded in the value statistics of every nonzero derivative.

Writing $A_p = \frac{1}{2} \|\text{up}\|_p^p$, one obtains the exact beta transform

$$A_p = p \int_0^1 (1-y)^{p-1} G(y) dy.$$

For $\mathcal{D} = y \, d/dy$, the dilation variable $Z_p = pY_p$ with $Y_p \sim \text{Beta}(1, p)$ has Mellin symbol

$$\mathbb{E} Z_p^s = p^s \frac{\Gamma(p+1)\Gamma(1+s)}{\Gamma(p+1+s)}.$$

This yields a new Euler–Gamma differential calculus

$$A_p \sim G - \gamma \mathcal{D}G + \frac{\gamma^2 + \pi^2/6}{2} \mathcal{D}^2 G - \frac{\gamma^3 + \gamma\pi^2/2 + 2\zeta(3)}{6} \mathcal{D}^3 G + \dots \quad \text{at } y = p^{-1}.$$

The coefficients are complete Bell polynomials in log-exponential cumulants; finite- p corrections are generated by Bernoulli polynomials through the gamma-ratio expansion. Combining this operator with the periodic inverse-Fabius endpoint expansion gives

$$\|\text{up}\|_p^p = 2G(p^{-1}) \left(1 - \frac{\gamma}{\rho} + \mathcal{O}(\rho^{-2})\right), \quad \rho = \sqrt{\frac{2 \log p}{\log 2}},$$

and, more sharply, shows that Euler’s constant subtracts from the first periodic inverse correction. In particular

$$1 - \|\text{up}\|_p \sim \frac{\sqrt{2 \log 2 \log p}}{p}.$$

A phase-locked sequence of p -norms recovers the nonconstant Gamma-zeta endpoint fluctuation itself.

Parseval turns the same norm law into a second, frequency-side geometry. The pushforward of $|\Phi(\xi)|^2 d\xi / \|\text{up}\|_2^2$ by $z = (2\pi\xi)^2$ has moments

$$m_n = 2^{n(n+1)} = q^{-n(n+1)/2}, \quad q = \frac{1}{4}.$$

It is therefore an explicit nonclassical representing measure for a shifted Stieltjes–Wigert moment problem. A lognormal measure has the same moments but a different density, proving indeterminacy directly. We calculate the Hankel determinants, monic orthogonal polynomials, recurrence coefficients, and Gaussian- q -binomial formulae, and derive the spectral q -Pearson law

$$w_{\text{sp}}(4z) = \frac{1}{2} \text{sinc}^2(\sqrt{z}) w_{\text{sp}}(z).$$

The report closes with numerical checks, explicit conjectures, a phase-locked confluent q -Richardson program, and a Lean formalization roadmap. Claims labelled “new” are new relative to the pinned repository corpus after a formula- and terminology-level collision audit; no universal publication-priority claim is made.

Contents

1	Scope, corpus audit, and novelty boundary	3
1.1	What was inspected	3
1.2	Status convention	3
2	Normalization and established structural identities	4
3	Exact equimeasurability and the complete derivative norm spectrum	5
3.1	Absolute and signed value distributions	5
3.2	Every rearrangement-invariant norm scales exactly	6
3.3	Thue–Morse extremizers and exact level sets	6
4	The norm spectrum determines the Fabius function	7
5	An exact inverse-Fabius beta transform	8
6	The Euler–Gamma differential operator	9
6.1	Exact Mellin symbol	9
6.2	Cumulants, Bell polynomials, and finite- p corrections	9
7	Large-p norms, Lambert coordinates, and phase tomography	11
7.1	Imported inverse endpoint expansion	11
7.2	Euler’s constant shifts the inverse correction	11
7.3	Recovering the endpoint periodic function from global norms	12

8	Fourier energy and a shifted Stieltjes–Wigert moment problem	12
8.1	Exact spectral moments	12
8.2	A direct proof of moment indeterminacy	13
8.3	A nonclassical q -Pearson equation	13
9	Hankel determinants, q-binomial polynomials, and Lagrange quadrature	14
9.1	Gaussian quadrature as q -binomial Lagrange interpolation	15
10	The optimal derivative weight and a q-Legendre Fourier envelope	16
10.1	Exact Denjoy–Carleman weight	16
10.2	Exact associated function and periodic lattice correction	16
11	Numerical experiments	17
11.1	Derivative tiling and norm checks	17
11.2	Spectral moments	17
11.3	Euler–Gamma operator convergence	18
11.4	Symbolic Hankel checks	18
12	Conjectures and promising research directions	18
12.1	A full norm transseries	19
12.2	Nevanlinna parameter of the sinc-square measure	19
12.3	Orthogonal complement and spectral zeros	20
12.4	Confluent q -Richardson phase tomography	20
12.5	Base- b and Pascal–Rvachev extensions	20
12.6	Formalization roadmap	21
13	Conclusion	21
A	Recursive TeX corpus ledger	22
B	Reproducibility files	24

1 Scope, corpus audit, and novelty boundary

1.1 What was inspected

The recursive tree under <https://github.com/VladimirReshetnikov/ProveIt/tree/0341bbf91cb867f4510bc8431cf59878208de771/Analysis/FabiusFunction/docs> contained 68 *.tex artifacts at the pinned commit. Every path was included in the audit. The living expositions, active frontier reports, Fourier-decay chain, imported paper transcriptions, and mathematically distinctive archival deltas were read. The archive deliberately preserves first versions and provenance; its own ledger explains that byte-identical or superseded descendants should not be treated as independent contemporary claims. Accordingly, archived variants were checked for distinctive content but deduplicated when building the novelty map. The complete path ledger appears in Section A.

The most important collision checks were against the following current reports:

- the primary Fabius–Rvachev exposition and mathematical glossary;
- the consolidated non-formalized frontier volume;
- the inverse-frontier report, including periodic Lambert–Bell quantile reversion;
- the complete Fourier-decay synthesis and its comparative audits;
- the finite-block/q-Richardson/deconvolution report;
- the half-integer, arithmetic-ray, spectral determinant, Pascal–Rvachev, and Gamma-duality frontier reports;
- all archived q-calculus, dyadic-asymptotic, Thue–Morse, and repeated-integration studies; and
- the seven imported paper sources.

Targeted searches were then run for the exact concepts and formula shapes used below: “equimeasurable derivative”, “all- L^p derivative norm”, “Beta(1, p) inverse Fabius”, “ $\Gamma(1 + \mathcal{D})$ ”, “Stieltjes–Wigert”, “Denjoy–Carleman”, and the exact moment sequence $2^{n(n+1)}$. No collision was found for the theorem package developed here. The underlying derivative tiling itself is present in the corpus and is explicitly marked as baseline.

1.2 Status convention

Three proof-status labels. *Corpus baseline* means a statement already present, perhaps in another normalization, in the inspected repository or one of its imported papers. *New deduction* means a statement and proof not located in that corpus after the collision audit. *Conjecture* means a precise, numerically or structurally motivated statement for which this report does not claim a proof. None of these labels asserts global publication priority.

The following are corpus baseline and are not claimed as new:

- (i) the identities relating F , up, and the signed global extension;
- (ii) the differential-refinement law for up and its finite Thue–Morse derivative tiling;
- (iii) the infinite sinc product and its integer zero multiplicities;
- (iv) the exact dyadic values of F , their q -Pochhammer and Gaussian-binomial forms;
- (v) the corrected endpoint expansion with a nonconstant periodic Gamma–zeta term;
- (vi) the inverse endpoint expansion in the lower-Lambert coordinate, including Bell reversion and quantile elasticity;

- (vii) the distinct pointwise, metric, L^1 , L^2 , and extremal Fourier-decay exponents;
- (viii) the existing finite-block, q-Richardson, convolution-tail, spectral determinant, and Pascal–Rvachev constructions.

New deductions proved in this report. The main new results are: exact derivative equimeasurability; scaling in every rearrangement-invariant norm; exact inverse-Fabius level-set laws; a Hausdorff reconstruction of F from integer derivative norms; the inverse-Fabius beta transform; an exact Mellin symbol and asymptotic Euler–Gamma/Bell/Bernoulli differential calculus; a large- p Lambert law with an Euler-constant shift and phase tomography; a shifted Stieltjes–Wigert spectral-energy measure; explicit Hankel determinants and q-binomial monic orthogonal polynomials; a nonclassical sinc-square q-Pearson equation; and the exact Denjoy–Carleman weight and associated periodic Legendre envelope generated by the derivative norms.

2 Normalization and established structural identities

Let $F : \mathbb{R} \rightarrow [0, 1]$ be the bounded Fabius function, with $F = 0$ on $(-\infty, 0]$, $F = 1$ on $[1, \infty)$, symmetry

$$F(1 - x) = 1 - F(x), \quad 0 \leq x \leq 1, \quad (2.1)$$

and differential law $F'(x) = 2F(2x)$ on $0 < x < 1/2$, continued by symmetry. Let

$$\text{up}(x) = \begin{cases} 1 - F(|x|), & |x| \leq 1, \\ 0, & |x| \geq 1. \end{cases} \quad (2.2)$$

Then up is even, nonnegative, C^∞ , supported on $[-1, 1]$, satisfies $\text{up}(0) = 1$ and $\int \text{up} = 1$, and obeys

$$\text{up}'(x) = 2(\text{up}(2x + 1) - \text{up}(2x - 1)). \quad (2.3)$$

Its Fourier transform is

$$\Phi(\xi) := \widehat{\text{up}}(\xi) = \prod_{j=0}^{\infty} \text{sinc}\left(\frac{\pi \xi}{2^j}\right), \quad \text{sinc } z = \frac{\sin z}{z}. \quad (2.4)$$

Write

$$\varepsilon_k := (-1)^{s_2(k)}, \quad T_n := \binom{n+1}{2} = \frac{n(n+1)}{2}, \quad (2.5)$$

where $s_2(k)$ is the binary digit sum. The identities $\varepsilon_{2k} = \varepsilon_k$ and $\varepsilon_{2k+1} = -\varepsilon_k$ will be used repeatedly.

Theorem 2.1 (Corpus baseline: exact derivative tiling). *For every $n \geq 0$ and $x \in \mathbb{R}$,*

$$\boxed{\text{up}^{(n)}(x) = 2^{T_n} \sum_{k=0}^{2^n-1} \varepsilon_k \text{up}(2^n x + 2^n - 1 - 2k).} \quad (2.6)$$

Equivalently, for $0 \leq k < 2^n$ and $-1 \leq y \leq 1$,

$$\boxed{\text{up}^{(n)}\left(-1 + \frac{2k+1+y}{2^n}\right) = 2^{T_n} \varepsilon_k \text{up}(y).} \quad (2.7)$$

Proof. The case $n = 0$ is immediate. Differentiate (2.6). Each derivative of an inner copy contributes 2^n , and (2.3) contributes another factor 2. The positive branch has index $2k$ and the negative branch index $2k + 1$; the Thue–Morse identities above give exactly the signs ε_{2k} and ε_{2k+1} . The scalar grows from 2^{T_n} to $2^{T_n+n+1} = 2^{T_{n+1}}$.

For the cell form, the support condition $|2^n x + 2^n - 1 - 2j| \leq 1$ partitions $[-1, 1]$ into the 2^n closed cells

$$I_{n,k} = \left[-1 + \frac{2k}{2^n}, -1 + \frac{2k+2}{2^n} \right].$$

On the interior of $I_{n,k}$ only the k th summand survives. At common boundaries every potentially overlapping copy is evaluated at ± 1 and hence vanishes to infinite order. Solving for the local coordinate gives (2.7). $\square$

Remark 2.2. The corpus already exploits this tiling for derivative values, zeros, and discrete Thue–Morse structure. The new step is to view (2.7) as a measure-preserving value-distribution identity and push that observation through normed function spaces, inverse quantiles, Mellin operators, and spectral moments.

3 Exact equimeasurability and the complete derivative norm spectrum

3.1 Absolute and signed value distributions

Let U be uniform on $[-1, 1]$ and let R be an independent Rademacher variable, $\mathbb{P}(R = 1) = \mathbb{P}(R = -1) = 1/2$.

Theorem 3.1 (Derivative equimeasurability). *For every $n \geq 0$,*

$$2^{-T_n} \left| \text{up}^{(n)}(U) \right| \stackrel{d}{=} \text{up}(U). \quad (3.1)$$

For every $n \geq 1$,

$$2^{-T_n} \text{up}^{(n)}(U) \stackrel{d}{=} R \text{up}(U). \quad (3.2)$$

Equivalently, for every Borel function H for which the integrals exist,

$$\frac{1}{2} \int_{-1}^1 H\left(2^{-T_n} \left| \text{up}^{(n)}(x) \right| \right) dx = \frac{1}{2} \int_{-1}^1 H(\text{up}(y)) dy, \quad (3.3)$$

$$\frac{1}{2} \int_{-1}^1 H\left(2^{-T_n} \text{up}^{(n)}(x)\right) dx = \frac{1}{4} \int_{-1}^1 (H(\text{up}(y)) + H(-\text{up}(y))) dy \quad (n \geq 1). \quad (3.4)$$

Proof. Integrate cell by cell and use (2.7). The affine map from each cell to $[-1, 1]$ has Jacobian 2^{-n} , hence

$$\int_{I_{n,k}} H\left(2^{-T_n} \left| \text{up}^{(n)}(x) \right| \right) dx = 2^{-n} \int_{-1}^1 H(\text{up}(y)) dy.$$

Summing 2^n equal terms proves (3.3). For $n \geq 1$, exactly half of the length- 2^n Thue–Morse block has sign $+1$ and half sign -1 , proving (3.4). $\square$

Corollary 3.2 (All absolute moments and signed moments). *For every real $p > 0$,*

$$\int_{-1}^1 \left| \text{up}^{(n)}(x) \right|^p dx = 2^{pT_n} \int_{-1}^1 \text{up}(x)^p dx. \quad (3.5)$$

For every integer $m \geq 0$ and $n \geq 1$,

$$\int_{-1}^1 \text{up}^{(n)}(x)^m dx = \begin{cases} 0, & m \text{ odd,} \\ 2^{mT_n} \int_{-1}^1 \text{up}(x)^m dx, & m \text{ even.} \end{cases} \quad (3.6)$$

3.2 Every rearrangement-invariant norm scales exactly

A rearrangement-invariant quasi-norm on a finite measure space depends only on the distribution of $|f|$. This includes Lebesgue, Lorentz, Lorentz–Zygmund, Orlicz, and many interpolation spaces.

Theorem 3.3 (Universal norm ladder). *Let $X[-1, 1]$ be any absolutely homogeneous rearrangement-invariant quasi-normed function space. Then*

$$\left\| \text{up}^{(n)} \right\|_X = 2^{T_n} \left\| \text{up} \right\|_X, \quad n \geq 0. \quad (3.7)$$

In particular, for every $0 < p \leq \infty$,

$$\left\| \text{up}^{(n)} \right\|_{L^p(\mathbb{R})} = 2^{n(n+1)/2} \left\| \text{up} \right\|_{L^p(\mathbb{R})}. \quad (3.8)$$

Proof. The functions vanish outside $[-1, 1]$. By Theorem 3.1, $2^{-T_n} |\text{up}^{(n)}|$ and up have the same decreasing rearrangement. Rearrangement invariance and absolute homogeneity give (3.7); the L^p case follows. $\square$

Corollary 3.4 (Exact sup norms, total variations, and derivative ratios). *Since $\|\text{up}\|_1 = \|\text{up}\|_\infty = 1$,*

$$\left\| \text{up}^{(n)} \right\|_1 = 2^{T_n}, \quad (3.9)$$

$$\left\| \text{up}^{(n)} \right\|_\infty = 2^{T_n}, \quad (3.10)$$

$$\text{TV}(\text{up}^{(n)}) = \left\| \text{up}^{(n+1)} \right\|_1 = 2^{T_{n+1}}. \quad (3.11)$$

For every $0 < p \leq \infty$,

$$\frac{\left\| \text{up}^{(n+1)} \right\|_p}{\left\| \text{up}^{(n)} \right\|_p} = 2^{n+1}. \quad (3.12)$$

3.3 Thue–Morse extremizers and exact level sets

Corollary 3.5 (Midpoint extrema). *For $0 \leq k < 2^n$, put*

$$x_{n,k} := -1 + \frac{2k+1}{2^n}. \quad (3.13)$$

Then

$$\text{up}^{(n)}(x_{n,k}) = 2^{T_n} \varepsilon_k. \quad (3.14)$$

These are precisely the points at which $|\text{up}^{(n)}|$ attains its global maximum. Thus the signs of the complete extremal set of the n th derivative form the first 2^n terms of the signed Thue–Morse sequence.

Proof. Set $y = 0$ in (2.7) and use $\text{up}(0) = 1$. Strict decrease of up on $[0, 1]$ makes $y = 0$ the unique maximum inside each cell. $\square$

Let $G = F^{-1} : [0, 1] \rightarrow [0, 1]$. Since $\text{up}(y) = 1 - F(|y|)$,

$$\{y \in [-1, 1] : \text{up}(y) > t\} = (-G(1-t), G(1-t)), \quad 0 < t < 1. \quad (3.15)$$

Theorem 3.6 (Exact inverse-Fabius level-set law). *For every $n \geq 0$ and $0 < t < 1$,*

$$\left| \left\{ x \in \mathbb{R} : \left| \text{up}^{(n)}(x) \right| > 2^{T_n} t \right\} \right| = 2G(1-t). \quad (3.16)$$

Consequently, if $V_n = 2^{-T_n} |\text{up}^{(n)}(U)|$, then

$$\mathbb{P}(V_n > t) = G(1 - t), \quad (3.17)$$

$$\mathbb{P}(V_n \leq t) = 1 - G(1 - t), \quad (3.18)$$

$$f_{V_n}(t) = G'(1 - t), \quad 0 < t < 1. \quad (3.19)$$

For $n \geq 1$, the signed variable $2^{-T_n} \text{up}^{(n)}(U)$ has density

$$f_n(v) = \frac{1}{2} G'(1 - |v|), \quad -1 < v < 1. \quad (3.20)$$

Proof. Combine equimeasurability with (3.15). The density follows by differentiating the distribution function on the open interval. The signed density is the symmetric Rademacher mixture. $\square$

Remark 3.7 (A quantile coupling). If U is uniform on $[-1, 1]$, then $|U|$ is uniform on $[0, 1]$ and

$$G(1 - \text{up}(U)) = |U|. \quad (3.21)$$

Thus the inverse Fabius function is not merely asymptotically related to derivative values; it is their exact probability integral transform.

4 The norm spectrum determines the Fabius function

Define the normalized norm spectrum

$$A_p := \frac{1}{2} \|\text{up}\|_p^p = 2^{-1-pT_n} \left\| \text{up}^{(n)} \right\|_p^p, \quad p > 0. \quad (4.1)$$

The second expression is independent of n by (3.8). Set $A_0 := 1$. If $W = \text{up}(U)$, then $A_p = \mathbb{E}W^p$ for every $p \geq 0$.

Proposition 4.1 (Complete monotonicity and log-convexity). *The function $p \mapsto A_p$ is strictly completely monotone on $(0, \infty)$:*

$$(-1)^r A_p^{(r)} = \mathbb{E}[W^p (-\log W)^r] > 0, \quad r \geq 0. \quad (4.2)$$

It is strictly log-convex. Its integer samples are a completely monotone Hausdorff moment sequence:

$$(-1)^r \Delta^r A_m = \mathbb{E}[W^m (1 - W)^r] > 0, \quad m, r \geq 0, \quad (4.3)$$

where $\Delta A_m = A_{m+1} - A_m$.

Proof. Write $W^p = \exp(-pX)$ with $X = -\log W \geq 0$ and differentiate under the integral. Strict log-convexity follows from Hölder's inequality because W is nonconstant. The finite-difference formula follows by expanding $(1 - W)^r$. $\square$

Theorem 4.2 (Hausdorff reconstruction from integer derivative norms). *The sequence $\{A_m\}_{m \geq 0}$ uniquely determines G , hence F and up . More explicitly, define*

$$b_{N,k} := \binom{N}{k} \sum_{j=0}^{N-k} (-1)^j \binom{N-k}{j} A_{k+j}, \quad 0 \leq k \leq N. \quad (4.4)$$

Then $b_{N,k} \geq 0$, $\sum_k b_{N,k} = 1$, and at every $0 < t < 1$,

$$G(1 - t) = 1 - \lim_{N \rightarrow \infty} \sum_{k=0}^{\lfloor Nt \rfloor} b_{N,k}. \quad (4.5)$$

The same reconstruction applies to the normalized norms of any fixed derivative $\text{up}^{(n)}$ through (4.1).

Proof. By (4.3),

$$b_{N,k} = \binom{N}{k} \mathbb{E}[W^k(1-W)^{N-k}].$$

Conditioned on $W = w$, these are the probabilities of a binomial random variable $K_N \sim \text{Bin}(N, w)$. Therefore

$$\sum_{k=0}^{\lfloor Nt \rfloor} b_{N,k} = \mathbb{EP}(K_N/N \leq t \mid W).$$

The conditional law of large numbers makes the inner probability converge to $\mathbf{1}_{\{W \leq t\}}$ away from $w = t$. The distribution of W is continuous, so the expectation converges to $\mathbb{P}(W \leq t) = 1 - G(1-t)$ by (3.18). Uniqueness is also the classical determinacy of the Hausdorff moment problem on $[0, 1]$. $\square$

Remark 4.3. This theorem gives a perhaps unexpected completeness statement: the scalar sequence $\|\text{up}^{(n)}\|_m$, $m = 1, 2, \dots$, for any one derivative order n , contains the entire Fabius function. The Thue–Morse oscillation disappears under absolute moments, yet no information about the underlying profile is lost.

5 An exact inverse-Fabius beta transform

Theorem 5.1 (Beta transform of the inverse Fabius function). *For every real $p > 0$,*

$$A_p = p \int_0^1 (1-y)^{p-1} G(y) \, dy. \quad (5.1)$$

Equivalently, if $Y_p \sim \text{Beta}(1, p)$, then

$$A_p = \mathbb{E}G(Y_p). \quad (5.2)$$

Proof. Using evenness and (2.2),

$$A_p = \int_0^1 (1-F(x))^p \, dx.$$

The monotone substitution $y = F(x)$, interpreted as a Lebesgue–Stieltjes substitution at the flat endpoints, gives

$$A_p = \int_0^1 (1-y)^p \, dG(y).$$

Integration by parts has no boundary term because $G(0) = 0$ and $(1-y)^p = 0$ at $y = 1$. This proves (5.1); the beta density is $p(1-y)^{p-1}$. $\square$

Corollary 5.2 (Layer-cake form). *For every $p > 0$ and every $n \geq 0$,*

$$\left\| \text{up}^{(n)} \right\|_p^p = 2^{pT_n+1} p \int_0^1 (1-y)^{p-1} G(y) \, dy. \quad (5.3)$$

The beta kernel concentrates at $y \asymp p^{-1}$. Because G is slowly varying at zero, (5.1) already implies $A_p \sim G(1/p)$. The next section replaces that first-order Abelian statement by an all-orders differential operator.

6 The Euler–Gamma differential operator

6.1 Exact Mellin symbol

Let

$$\mathcal{D} := y \frac{d}{dy} \quad (6.1)$$

be the Euler dilation derivative. Put $Z_p = pY_p$.

Proposition 6.1 (Mellin symbol of the beta dilation). *For $\Re s > -1$,*

$$M_p(s) := \mathbb{E} Z_p^s = p^s \frac{\Gamma(p+1)\Gamma(1+s)}{\Gamma(p+1+s)}. \quad (6.2)$$

For any bounded extension of G to $(1, \infty)$,

$$A_p = [M_p(\mathcal{D})G](p^{-1}), \quad (6.3)$$

where the operator is rigorously interpreted by the Bochner average

$$[M_p(\mathcal{D})H](y) := \mathbb{E}[H(Z_p y)]. \quad (6.4)$$

The value in (6.3) is independent of how G is extended, because $0 < Z_p/p = Y_p < 1$.

Proof. The beta integral gives

$$\mathbb{E} Z_p^s = p^s p \int_0^1 y^s (1-y)^{p-1} dy = p^s \frac{\Gamma(p+1)\Gamma(1+s)}{\Gamma(p+1+s)}.$$

The dilation semigroup satisfies $\exp(t\mathcal{D})H(y) = H(e^t y)$. Averaging with $t = \log Z_p$ and evaluating at $y = p^{-1}$ gives $\mathbb{E} G(Z_p/p) = \mathbb{E} G(Y_p) = A_p$. $\square$

As $p \rightarrow \infty$, Z_p converges to an exponential random variable Z of mean one, and $M_p(s) \rightarrow \Gamma(1+s)$. Thus $\Gamma(1+\mathcal{D})$ is not decorative notation: it is the Mellin symbol of the limiting random dilation.

6.2 Cumulants, Bell polynomials, and finite-p corrections

Let

$$K_p(s) := \log M_p(s). \quad (6.5)$$

The cumulants of $\log Z_p$ are

$$\kappa_{1,p} = \log p - \psi(p+1) - \gamma, \quad (6.6)$$

$$\kappa_{m,p} = \psi^{(m-1)}(1) - \psi^{(m-1)}(p+1), \quad m \geq 2. \quad (6.7)$$

If B_m denotes the complete exponential Bell polynomial, then

$$\mu_{m,p} := \mathbb{E}(\log Z_p)^m = B_m(\kappa_{1,p}, \dots, \kappa_{m,p}). \quad (6.8)$$

As $p \rightarrow \infty$,

$$\kappa_1 = -\gamma, \quad (6.9)$$

$$\kappa_m = (-1)^m (m-1)! \zeta(m), \quad m \geq 2, \quad (6.10)$$

which are the cumulants of $\log Z$ for $Z \sim \text{Exp}(1)$.

The uniform gamma-ratio expansion on compact s -sets is

$$K_p(s) \sim \log \Gamma(1+s) + \sum_{r=1}^{\infty} \frac{(-1)^{r+1} (B_{r+1}(1) - B_{r+1}(1+s))}{r(r+1)p^r}. \quad (6.11)$$

Here $B_m(x)$ are Bernoulli polynomials. The first correction is $-s(s+1)/(2p)$.

Theorem 6.2 (Local Gamma–Bell expansion). *Fix $N \geq 1$. As $p \rightarrow \infty$,*

$$A_p = \sum_{m=0}^{N-1} \frac{\mu_m}{m!} \mathcal{D}^m G(p^{-1}) + \mathcal{O}_N \left(G(p^{-1}) \frac{(\log \rho)^{c_N}}{\rho^N} \right), \quad (6.12)$$

where

$$\rho = \sqrt{\frac{2 \log p}{\log 2}}, \quad \mu_m = B_m(-\gamma, \zeta(2), -2\zeta(3), 6\zeta(4), \dots), \quad (6.13)$$

and c_N is a finite integer depending only on N . A finite- p expansion with corrections to every power of p^{-1} is obtained by replacing μ_m with (6.8) or, equivalently, substituting (6.11) into $\exp K_p(\mathcal{D})$.

Proof. Write $y = p^{-1}$ and $t = \log Z_p$. On a logarithmic neighborhood of y ,

$$G(e^t y) = \sum_{m=0}^{N-1} \frac{t^m}{m!} \mathcal{D}^m G(y) + R_N(t, y).$$

The all-orders inverse endpoint expansion in the corpus implies, for each fixed m , $\mathcal{D}^m G(y)/G(y)$ is bounded by a power of $\log \rho$ times ρ^{-m} . Fix a constant $C > L + 1$ and split the expectation into $|t| \leq C\rho$ and its complement. On the central range, every point $e^u y$ between y and $e^t y$ has endpoint coordinate $\rho + \mathcal{O}_C(1)$. Integrating the elasticity shows $G(e^u y)/G(y) = \mathcal{O}_C(1)$ there, so Taylor’s integral remainder is

$$|R_N(t, y)| \leq C_N G(y) \frac{(\log \rho)^{c_N}}{\rho^N} |t|^N.$$

The moments $\mathbb{E}|\log Z_p|^N$ are bounded uniformly for $p \geq 2$: near zero the density of Z_p is at most one, while for $z \geq 1$ it is bounded by $e^{-z/2}$. For the omitted negative tail, $\mathbb{P}(Z_p < e^{-C\rho}) \leq e^{-C\rho}$; for the positive tail, $\mathbb{P}(Z_p > e^{C\rho}) \leq \exp(-e^{C\rho})$ once p is large. Since $G(p^{-1}) \asymp \rho e^{-L\rho}$, both tail contributions are $o(G(p^{-1})\rho^{-N})$. Taking expectations gives coefficients $\mu_{m,p}$; replacing them by μ_m changes the result by $\mathcal{O}(p^{-1}G(p^{-1}))$, which is beyond every power of ρ^{-1} relative to $G(p^{-1})$. The Bell and Bernoulli descriptions follow from differentiation of (6.5) and the standard gamma-ratio expansion. $\square$

The first five log-exponential moments are

$$\begin{aligned} \mu_0 &= 1, \\ \mu_1 &= -\gamma, \\ \mu_2 &= \gamma^2 + \zeta(2), \\ \mu_3 &= -\gamma^3 - 3\gamma\zeta(2) - 2\zeta(3), \\ \mu_4 &= \gamma^4 + 6\gamma^2\zeta(2) + 8\gamma\zeta(3) + 3\zeta(2)^2 + 6\zeta(4). \end{aligned} \quad (6.14)$$

Thus

$$\begin{aligned} A_p &= G - \gamma \mathcal{D}G + \frac{\gamma^2 + \pi^2/6}{2} \mathcal{D}^2 G \\ &\quad - \frac{\gamma^3 + \gamma\pi^2/2 + 2\zeta(3)}{6} \mathcal{D}^3 G \\ &\quad + \frac{\gamma^4 + \gamma^2\pi^2 + 8\gamma\zeta(3) + 3\pi^4/20}{24} \mathcal{D}^4 G + \dots, \quad y = p^{-1}. \end{aligned} \quad (6.15)$$

Remark 6.3 (Why Bell and Bernoulli polynomials appear). Bell polynomials convert the cumulants of the random logarithmic dilation into its ordinary moments, hence into powers of $\mathcal{D}$. Bernoulli polynomials govern the finite- p deviation of the beta dilation from the limiting exponential dilation through the quotient $\Gamma(p+1)/\Gamma(p+1+s)$. The two families enter for different structural reasons and then compose automatically.

7 Large-p norms, Lambert coordinates, and phase tomography

7.1 Imported inverse endpoint expansion

Put

$$L := \log 2, \quad T := \log p, \quad \rho := \sqrt{\frac{2T}{L}}, \quad \beta := \frac{1}{L} + \frac{1}{2}, \quad \theta := \rho + \beta. \quad (7.1)$$

The inverse-frontier report in the corpus proves

$$G_0(p^{-1}) := \rho e^{-L\rho - L\beta} = \frac{\sqrt{T}}{e\sqrt{L}} e^{-\sqrt{2LT}}, \quad (7.2)$$

and

$$\frac{G(p^{-1})}{G_0(p^{-1})} = 1 + \frac{\frac{1}{2} \log \rho + \kappa_0 - \Psi(\theta)}{\rho} + \mathcal{O}\left(\frac{(\log \rho)^2}{\rho^2}\right), \quad (7.3)$$

where Ψ is a nonconstant real-analytic one-periodic Gamma-zeta function. It also proves the endpoint elasticity

$$\frac{yG'(y)}{G(y)} = \frac{1}{\rho} + \frac{\Psi'(\rho + \beta) - 1}{L\rho^2} + \mathcal{O}\left(\frac{\log \rho}{\rho^3}\right), \quad y \downarrow 0. \quad (7.4)$$

These are baseline inputs; the norm consequences below are new.

7.2 Euler's constant shifts the inverse correction

Theorem 7.1 (Two-term norm-to-quantile ratio). *As $p \rightarrow \infty$,*

$$\boxed{\frac{A_p}{G(p^{-1})} = 1 - \frac{\gamma}{\rho} + \frac{1}{\rho^2} \left[\frac{\gamma^2 + \pi^2/6}{2} - \frac{\gamma(\Psi'(\theta) - 1)}{L} \right] + \mathcal{O}\left(\frac{\log \rho}{\rho^3}\right)}. \quad (7.5)$$

In particular,

$$\boxed{A_p \sim G(p^{-1})}. \quad (7.6)$$

Proof. By (6.15),

$$\frac{A_p}{G} = 1 - \gamma \frac{\mathcal{D}G}{G} + \frac{\gamma^2 + \pi^2/6}{2} \frac{\mathcal{D}^2G}{G} + \mathcal{O}(\rho^{-3} \log \rho).$$

Equation (7.4) gives

$$\frac{\mathcal{D}G}{G} = \frac{1}{\rho} + \frac{\Psi'(\theta) - 1}{L\rho^2} + \mathcal{O}\left(\frac{\log \rho}{\rho^3}\right).$$

If $a = \mathcal{D} \log G$, then $\mathcal{D}^2G/G = a^2 + \mathcal{D}a = \rho^{-2} + \mathcal{O}((\log \rho)/\rho^3)$. Substitution proves the formula. $\square$

Corollary 7.2 (First periodic norm expansion and Euler shift). *As $p \rightarrow \infty$,*

$$\boxed{\frac{\|\text{up}\|_p^p}{2G_0(p^{-1})} = 1 + \frac{\frac{1}{2} \log \rho + \kappa_0 - \Psi(\theta) - \gamma}{\rho} + \mathcal{O}\left(\frac{(\log \rho)^2}{\rho^2}\right)}. \quad (7.7)$$

Thus the first correction for the global L^p norm is the first inverse-quantile correction with the universal subtraction γ .

Proof. Multiply (7.5) by (7.3) and use $\|\text{up}\|_p^p = 2A_p$. $\square$

Corollary 7.3 (Approach to the sup norm). *As $p \rightarrow \infty$,*

$$1 - \|\text{up}\|_p \sim \frac{\sqrt{2 \log 2 \log p}}{p}. \quad (7.8)$$

More precisely,

$$-p \log \|\text{up}\|_p = L\rho - \log \rho + 1 - \frac{L}{2} - \frac{\frac{1}{2} \log \rho + \kappa_0 - \Psi(\theta) - \gamma}{\rho} + \mathcal{O}\left(\frac{(\log \rho)^2}{\rho^2}\right). \quad (7.9)$$

Proof. Take logarithms in (7.7) and use $G_0 = \rho \exp(-L\rho - L\beta)$ with $L\beta = 1 + L/2$. The leading term of $-\log \|\text{up}\|_p$ is $L\rho/p$; since it tends to zero, $1 - \|\text{up}\|_p \sim -\log \|\text{up}\|_p$. $\square$

7.3 Recovering the endpoint periodic function from global norms

Theorem 7.4 (Phase tomography by L^p norms). *Fix a phase $\vartheta \in \mathbb{R}/\mathbb{Z}$. For all sufficiently large integers j , define*

$$\rho_j := j + \vartheta - \beta, \quad p_j := \exp\left(\frac{L}{2}\rho_j^2\right). \quad (7.10)$$

Then

$$\lim_{j \rightarrow \infty} \left[\rho_j \left(\frac{\|\text{up}\|_{p_j}^{p_j}}{2G_0(p_j^{-1})} - 1 \right) - \frac{1}{2} \log \rho_j \right] = \kappa_0 - \gamma - \Psi(\vartheta). \quad (7.11)$$

Hence the nonconstant endpoint fluctuation Ψ can be reconstructed from a family of global norm measurements; pointwise evaluation or inversion of F is not required.

Proof. For (7.10), the parameter in (7.1) is exactly ρ_j and $\theta_j = \rho_j + \beta = j + \vartheta$. Periodicity gives $\Psi(\theta_j) = \Psi(\vartheta)$. Multiply (7.7) by ρ_j and subtract $\frac{1}{2} \log \rho_j$. $\square$

Remark 7.5 (Lambert interpretation). The Gaussian-logarithmic sampling law $\log p_j = (L/2)\rho_j^2$ is the inverse-side analogue of the lower-Lambert phase coordinate used in the endpoint expansion. It is precisely the sampling geometry that freezes the periodic saddle phase while sending the asymptotic scale to zero.

8 Fourier energy and a shifted Stieltjes–Wigert moment problem

8.1 Exact spectral moments

Let

$$C_2 := \|\text{up}\|_2^2 = \int_{\mathbb{R}} |\Phi(\xi)|^2 d\xi. \quad (8.1)$$

Numerically, $C_2 = 0.8088735359679707\dots$, but no decimal evaluation is needed below.

Theorem 8.1 (Shifted Stieltjes–Wigert spectral moments). *Define the probability measure*

$$d\nu(\xi) := \frac{|\Phi(\xi)|^2}{C_2} d\xi \quad (8.2)$$

and let μ_{sp} be its pushforward by $z = (2\pi\xi)^2$. Then, for every $n \geq 0$,

$$\int_0^\infty z^n d\mu_{\text{sp}}(z) = 2^{n(n+1)}. \quad (8.3)$$

With $q = 1/4$, this is

$$m_n = q^{-n(n+1)/2}. \quad (8.4)$$

Proof. Parseval and the Fourier differentiation rule give

$$\int_{\mathbb{R}} (2\pi\xi)^{2n} |\Phi(\xi)|^2 d\xi = \left\| \text{up}^{(n)} \right\|_2^2.$$

By (3.8) with $p = 2$, the right side is $2^{2T_n} C_2 = 2^{n(n+1)} C_2$. Divide by C_2 and push forward. $\square$

The measure has the explicit density

$$w_{\text{sp}}(z) = \frac{|\Phi(\sqrt{z}/(2\pi))|^2}{2\pi C_2 \sqrt{z}}, \quad z > 0. \quad (8.5)$$

8.2 A direct proof of moment indeterminacy

Let $L = \log 2$ and define the lognormal density

$$w_{\text{ln}}(z) := \frac{1}{z\sqrt{4\pi L}} \exp\left(-\frac{(\log z - L)^2}{4L}\right), \quad z > 0. \quad (8.6)$$

This is the law of $\exp N(L, 2L)$ and therefore

$$\int_0^\infty z^n w_{\text{ln}}(z) dz = \exp(nL + n^2 L) = 2^{n(n+1)}. \quad (8.7)$$

Corollary 8.2 (Explicit indeterminacy). *The Stieltjes moment sequence (8.3) is indeterminate. Indeed, $w_{\text{sp}}(z) dz$ and $w_{\text{ln}}(z) dz$ are distinct probability measures with all moments equal. More generally,*

$$w_t := (1-t)w_{\text{ln}} + tw_{\text{sp}}, \quad 0 \leq t \leq 1, \quad (8.8)$$

forms an explicit continuum of representing densities.

Proof. The moment equality is (8.3)–(8.7). The densities are different because w_{sp} vanishes whenever $z = (2\pi m)^2$, $m \in \mathbb{N}$, whereas w_{ln} is strictly positive on $(0, \infty)$. Convex combinations preserve every moment. $\square$

8.3 A nonclassical q-Pearson equation

The Fourier product satisfies

$$\Phi(2\xi) = \text{sinc}(2\pi\xi)\Phi(\xi). \quad (8.9)$$

Proposition 8.3 (Sinc-square q-Pearson law). *The spectral density (8.5) obeys*

$$w_{\text{sp}}(4z) = \frac{1}{2} \text{sinc}^2(\sqrt{z}) w_{\text{sp}}(z). \quad (8.10)$$

By contrast, the lognormal representative satisfies

$$w_{\text{ln}}(4z) = \frac{1}{4z} w_{\text{ln}}(z). \quad (8.11)$$

Proof. In (8.5), replacing z by $4z$ doubles the Fourier argument and multiplies the square-root denominator by 2. Apply (8.9) and note that $2\pi(\sqrt{z}/(2\pi)) = \sqrt{z}$. The lognormal identity follows by direct substitution in (8.6). $\square$

Remark 8.4. Classical Stieltjes–Wigert measures satisfy q-Pearson equations with monomial or rational multipliers. Equation (8.10) replaces that multiplier by an entire sinc-square factor carrying the complete integer-zero divisor. Determining its position in the Nevanlinna family is one of the principal open directions in Section 12.




Figure 1: The sinc-square spectral density and a lognormal density have exactly the same nonnegative integer moments but visibly different local structures. The spectral zeros are inherited from the integer zeros of the infinite sinc product.

9 Hankel determinants, q-binomial polynomials, and Lagrange quadrature

Fix $0 < q < 1$ and the moment sequence

$$m_n = q^{-n(n+1)/2}. \quad (9.1)$$

The Fabius–Rvachev spectral case is $q = 1/4$.

Theorem 9.1 (Hankel determinants). *For $N \geq 0$,*

$$\Delta_N := \det[m_{i+j}]_{i,j=0}^N = q^{-N(N+1)(4N+5)/6} \prod_{k=1}^N (q; q)_k, \quad (9.2)$$

where $(q; q)_k = \prod_{r=1}^k (1 - q^r)$.

Proof. Factor

$$m_{i+j} = q^{-i(i+1)/2} q^{-j(j+1)/2} q^{-ij}.$$

The row and column factors contribute $q^{-\sum_{i=0}^N i(i+1)}$. The remaining determinant is the Vandermonde determinant in q^{-i} :

$$\det[(q^{-i})^j]_{i,j=0}^N = \prod_{0 \leq i < k \leq N} (q^{-k} - q^{-i}) = q^{-\sum_{k=1}^N k^2} \prod_{k=1}^N (q; q)_k.$$

Finally

$$\sum_{i=1}^N i(i+1) + \sum_{i=1}^N i^2 = \frac{N(N+1)(4N+5)}{6}.$$

□

Theorem 9.2 (Monic shifted Stieltjes–Wigert polynomials). *The monic orthogonal polynomial of degree n for any representing measure of (9.1) is*

$$P_n(z) = \sum_{k=0}^n (-1)^{n-k} q^{k^2-n^2} \begin{bmatrix} n \\ k \end{bmatrix}_q z^k. \quad (9.3)$$

It satisfies

$$\int z^j P_n(z) d\mu(z) = 0, \quad 0 \leq j < n, \quad (9.4)$$

for every measure μ with moments (9.1), and

$$h_n := \int P_n(z)^2 d\mu(z) = q^{-(2n^2+n)} (q; q)_n. \quad (9.5)$$

The three-term recurrence is

$$P_{n+1}(z) = (z - a_n)P_n(z) - b_n P_{n-1}(z), \quad (9.6)$$

with

$$a_n = q^{-2n-1}(1 + q - q^{n+1}), \quad (9.7)$$

$$b_n = q^{-(4n-1)}(1 - q^n), \quad n \geq 1. \quad (9.8)$$

Proof. Let c_k be the coefficient of z^k in (9.3). For $j < n$,

$$\begin{aligned} \sum_{k=0}^n c_k m_{k+j} &= (-1)^n q^{-n^2-j(j+1)/2} \sum_{k=0}^n (-1)^k q^{k(k-1)/2} \begin{bmatrix} n \\ k \end{bmatrix}_q q^{-jk} \\ &= (-1)^n q^{-n^2-j(j+1)/2} (q^{-j}; q)_n = 0, \end{aligned}$$

by the finite q-binomial theorem; one factor is $1 - q^0$. The leading coefficient is one. The norm is the standard Gram-determinant quotient $h_n = \Delta_n / \Delta_{n-1}$, which gives (9.5). Hence $b_n = h_n / h_{n-1}$. Comparing the coefficients of z^n in the monic recurrence gives a_n ; the coefficient of z^{n-1} in P_n is $-q^{-2n+1}(1 - q^n)/(1 - q)$, yielding (9.7). $\square$

For $q = 1/4$, the first polynomials are

$$\begin{aligned} P_0(z) &= 1, \\ P_1(z) &= z - 4, \\ P_2(z) &= z^2 - 80z + 256, \\ P_3(z) &= (z - 64)(z^2 - 1280z + 4096). \end{aligned} \quad (9.9)$$

9.1 Gaussian quadrature as q-binomial Lagrange interpolation

Let $\lambda_{n,1}, \dots, \lambda_{n,n}$ be the zeros of P_n and let $\ell_{n,j}$ be the Lagrange cardinal polynomial at these nodes. The Gaussian weights are

$$\omega_{n,j} := \int \ell_{n,j}(z) d\mu(z) = \left(\sum_{r=0}^{n-1} \frac{P_r(\lambda_{n,j})^2}{h_r} \right)^{-1}. \quad (9.10)$$

Because the polynomials and h_r are determined by the moment sequence, the nodes and weights are identical for μ_{sp} , the lognormal measure, and every other representing measure.

Corollary 9.3 (Measure-independent Gaussian quadrature). *For every polynomial Q of degree at most $2n - 1$,*

$$\int_0^\infty Q(z) d\mu_{\text{sp}}(z) = \sum_{j=1}^n \omega_{n,j} Q(\lambda_{n,j}) = \int_0^\infty Q(z) w_{\text{ln}}(z) dz. \quad (9.11)$$

The q -binomial coefficients in (9.3) therefore generate exact finite Lagrange quadrature rules for polynomial spectral energies of the infinite sinc product.

Remark 9.4. Moment indeterminacy is invisible to every finite polynomial test: all representing measures produce the same Gaussian quadrature. The distinction lives in the orthogonal complement of the polynomial closure, exactly where the integer-zero geometry of Φ can enter.

10 The optimal derivative weight and a q-Legendre Fourier envelope

10.1 Exact Denjoy–Carleman weight

Define

$$M_n := 2^{T_n} = 2^{n(n+1)/2}. \quad (10.1)$$

By (3.10), up saturates this derivative weight exactly:

$$\|\text{up}^{(n)}\|_\infty = M_n. \quad (10.2)$$

The sequence is logarithmically convex and

$$\frac{M_n}{M_{n+1}} = 2^{-(n+1)}, \quad \sum_{n=0}^\infty \frac{M_n}{M_{n+1}} = 1 < \infty. \quad (10.3)$$

Thus the direct-weight Denjoy–Carleman class generated by M is nonquasianalytic, as the existence of the compactly supported nonzero function up requires.

Proposition 10.1 (Outside every finite Gevrey class). *For no finite $s > 0$ do constants $A, C > 0$ satisfy*

$$M_n \leq CA^n (n!)^s \quad (n \geq 0). \quad (10.4)$$

Hence up belongs to no finite-order Gevrey class on any interval containing its support.

Proof. The logarithm of M_n is $(\log 2)n^2/2 + \mathcal{O}(n)$, whereas $\log((n!)^s A^n) = sn \log n + \mathcal{O}(n)$. The quadratic term wins. $\square$

10.2 Exact associated function and periodic lattice correction

Define the associated function

$$\omega_M(t) := \sup_{n \geq 0} \log \frac{t^n}{M_n}, \quad t > 0. \quad (10.5)$$

Let $r = \log_2 t$.

Theorem 10.2 (Exact q-Legendre transform). *For $r \geq 1/2$,*

$$\omega_M(t) = \frac{\log 2}{2} \left[\left(r - \frac{1}{2} \right)^2 - \text{dist} \left(r - \frac{1}{2}, \mathbb{N}_0 \right)^2 \right]. \quad (10.6)$$

For $r \leq 1/2$, $\omega_M(t) = 0$. Thus the continuous log-square Legendre transform carries an exact bounded one-periodic lattice correction in $\log_2 t$.

Proof. For each integer $n \geq 0$,

$$\log \frac{t^n}{M_n} = \log 2 \left(nr - \frac{n(n+1)}{2} \right) = \frac{\log 2}{2} \left[\left(r - \frac{1}{2} \right)^2 - \left(n - r + \frac{1}{2} \right)^2 \right].$$

Maximizing means choosing the nearest nonnegative integer to $r - 1/2$. If $r \leq 1/2$, the maximum is attained at $n = 0$ and equals zero. $\square$

Corollary 10.3 (Optimized integration-by-parts envelope). *For every $\xi \neq 0$,*

$$|\Phi(\xi)| \leq \inf_{n \geq 0} \frac{M_n}{(2\pi|\xi|)^n} = \exp(-\omega_M(2\pi|\xi|)). \quad (10.7)$$

If $r = \log_2(2\pi|\xi|) \geq 1/2$, this is

$$|\Phi(\xi)| \leq 2^{-\frac{1}{2}(r-1/2)^2 + \frac{1}{2} \text{dist}(r-1/2, \mathbb{N}_0)^2}. \quad (10.8)$$

Proof. Integrating the Fourier transform by parts n times and using (3.9) gives $|\Phi(\xi)| \leq M_n/(2\pi|\xi|)^n$. Optimize with Theorem 10.2. $\square$

Warning 10.4. The envelope (10.8) is the best one obtainable from the exact L^1 derivative norms alone; it is not the sharp pointwise or annular Fourier envelope. The Fourier-decay reports in the corpus exploit the lacunary sine cocycle and obtain stronger linear-in-log $|\xi|$ corrections. The value of (10.8) is its exact functional-analytic duality with the derivative weight, not a replacement for those sharper results.

11 Numerical experiments

All code used here is in `numerical_experiments.py`. It is included with detailed comments in the accompanying archive. The script reconstructs `up` by inverse FFT on a periodic box of length 4 from 55 dyadic sinc factors, using 2^{19} samples. Since `up` is supported on $[-1, 1]$, the box supplies exact geometric zero padding. The run used

$$\Delta x = 2^{-17} = 7.62939453125 \times 10^{-6}, \quad \Delta \xi = \frac{1}{4}.$$

The reconstructed density had integral 1 and value `up(0) = 1` to displayed precision; the largest value outside the support was 3.33×10^{-16} .

11.1 Derivative tiling and norm checks

For $n = 0, \dots, 6$ and $p \in \{1, 2, 4, 8, \infty\}$, the script formed the derivative from the exact cell formula and compared its numerical norm with $2^{T_n} \|\text{up}\|_p$. The largest absolute deviation of the ratio from 1 was 2.22×10^{-16} .

11.2 Spectral moments

n	numerical $\int (2\pi\xi)^{2n} \Phi ^2$	predicted $2^{n(n+1)} C_2$	relative error
0	$8.088735359679706 \times 10^{-1}$	$8.088735359679707 \times 10^{-1}$	-1.11×10^{-16}
1	3.235494143871882	3.235494143871883	-2.22×10^{-16}
2	$5.176790630195011 \times 10^1$	$5.176790630195013 \times 10^1$	-2.22×10^{-16}
3	$3.313146003324810 \times 10^3$	$3.313146003324808 \times 10^3$	4.44×10^{-16}
4	$8.481653768511517 \times 10^5$	$8.481653768511509 \times 10^5$	8.88×10^{-16}
5	$8.685213458955796 \times 10^8$	$8.685213458955785 \times 10^8$	1.33×10^{-15}
6	$3.557463432788293 \times 10^{12}$	$3.557463432788290 \times 10^{12}$	8.88×10^{-16}
7	$5.828548088280333 \times 10^{16}$	$5.828548088280334 \times 10^{16}$	-1.11×10^{-16}

The near-machine equality is a stringent joint check of the FFT normalization, derivative scale, Parseval convention, and moment exponent.




Figure 2: Normalized derivatives $2^{-T_n} \text{up}^{(n)}$, vertically offset. Each row is a concatenation of copies of the original bump; their signs are the length- 2^n Thue–Morse block.

11.3 Euler–Gamma operator convergence

The inverse interpolant was obtained directly from $G(1 - \text{up}(x)) = x$ on $0 \leq x \leq 1$. Logarithmic finite differences estimated $\mathcal{D}^k G(1/p)$ for $k \leq 3$. The following table lists relative errors of successive truncations of (6.15).

p	$A_p/G(1/p)$	G only	through $\mathcal{D}$	through $\mathcal{D}^2$	through $\mathcal{D}^3$
256	0.92618125	-7.38×10^{-2}	4.84×10^{-2}	-1.47×10^{-2}	6.46×10^{-3}
512	0.92793214	-7.21×10^{-2}	4.32×10^{-2}	-1.20×10^{-2}	6.37×10^{-3}
1024	0.92942377	-7.06×10^{-2}	3.91×10^{-2}	-9.78×10^{-3}	4.87×10^{-3}
2048	0.93089393	-6.91×10^{-2}	3.60×10^{-2}	-8.37×10^{-3}	3.35×10^{-3}
4096	0.93235311	-6.76×10^{-2}	3.33×10^{-2}	-7.55×10^{-3}	2.53×10^{-3}
8192	0.93370939	-6.63×10^{-2}	3.09×10^{-2}	-6.86×10^{-3}	2.39×10^{-3}
16384	0.93492704	-6.51×10^{-2}	2.87×10^{-2}	-6.10×10^{-3}	2.29×10^{-3}
32768	0.93604971	-6.40×10^{-2}	2.69×10^{-2}	-5.39×10^{-3}	1.97×10^{-3}
65536	0.93712995	-6.29×10^{-2}	2.53×10^{-2}	-4.83×10^{-3}	1.60×10^{-3}

11.4 Symbolic Hankel checks

SymPy evaluated the determinants in (9.2) exactly for $N = 0, \dots, 6$ at $q = 1/4$; every quotient of the direct determinant by the claimed closed form simplified to 1. The base-10 logarithms of the determinants for $N = 1, \dots, 6$ were

$$1.68124, 7.54887, 20.03233, 41.54498, 74.49635, 121.29499,$$

illustrating the rapid $q^{-\frac{2}{3}N^3 + \mathcal{O}(N^2)}$ growth.

12 Conjectures and promising research directions




Figure 3: Absolute relative errors of the first four Euler–Gamma operator truncations. The slow scale $\rho^{-1} \asymp (\log p)^{-1/2}$ explains why very large p is needed before successive asymptotic orders separate cleanly.

12.1 A full norm transseries

The inverse report gives an all-orders expansion of G in the differential algebra generated by periodic saddle jets. The exact operator (6.3) should transport that algebra to the norm spectrum without introducing any new transcendental data beyond the log-exponential cumulants.

Conjecture 12.1 (All-orders Gamma–Lambert norm transseries). *There exist one-periodic real-analytic coefficient functions $C_{m,\ell}(\theta)$ such that*

$$\frac{\|\text{up}\|_p^p}{2G_0(p^{-1})} \sim 1 + \sum_{m=1}^{\infty} \rho^{-m} \sum_{\ell=0}^m C_{m,\ell}(\theta) (\log \rho)^\ell, \quad (12.1)$$

with $C_{1,1} = 1/2$ and $C_{1,0} = \kappa_0 - \gamma - \Psi(\theta)$. Every $C_{m,\ell}$ belongs to the differential algebra generated by the inverse saddle jets and the constants $\gamma, \zeta(2), \dots, \zeta(m)$; no additional periodic functions occur.

A proof should combine the formal inverse recursion already in the corpus with a tame remainder theorem for the operator $M_p(\mathcal{D})$. The key analytic task is uniform control of $\mathcal{D}^m G$ on logarithmic windows whose width grows slowly with ρ .

12.2 Nevanlinna parameter of the sinc-square measure

Problem 12.2 (Locate μ_{sp} in the Stieltjes–Wigert Nevanlinna family). Determine the Nevanlinna parameter corresponding to the density (8.5). Equivalently, calculate the Cauchy transform

$$S_{\text{sp}}(z) = \int_0^\infty \frac{d\mu_{\text{sp}}(t)}{z - t} \quad (12.2)$$

relative to the classical entire functions A, B, C, D of the shifted Stieltjes–Wigert moment problem.

The exact q-Pearson law (8.10) supplies unusually rigid extra data. One promising route is to Mellin-transform it, obtaining a functional equation coupling the spectral Mellin transform to the Mellin transform of $\text{sinc}^2(\sqrt{z})$. Another is to use the canonical product of Φ to express S_{sp} as a convergent sum over integer-zero lobes.

Conjecture 12.3 (Sinc-Pearson uniqueness). *Among absolutely continuous representing measures of $m_n = 2^{n(n+1)}$ whose density is locally bounded after multiplication by $\sqrt{z}$, the normalization, moment conditions, and q -Pearson equation (8.10) determine w_{sp} uniquely.*

12.3 Orthogonal complement and spectral zeros

For an indeterminate moment problem, the polynomials are not dense in every representing L^2 space. The spectral measure should possess an orthogonal complement reflecting the integer-zero structure of Φ .

Problem 12.4 (Explicit nonpolynomial orthogonal basis). Construct functions $Q_j(z)$ such that

$$\int_0^\infty z^n Q_j(z) w_{\text{sp}}(z) dz = 0 \quad (n \geq 0)$$

and such that the Q_j span the orthogonal complement of the shifted Stieltjes–Wigert polynomials in $L^2(\mu_{\text{sp}})$. Dyadic rescalings of sinc factors, integer-lobe indicator functions, and q -Bessel functions from the classical theory are the three natural candidate families.

12.4 Confluent q -Richardson phase tomography

The existing corpus develops Gaussian- q Richardson weights for pure power error germs. The norm expansion (12.1) contains powers multiplied by logarithms. Choose a fixed phase ϑ and a dyadic phase-locked sequence

$$\rho_j = M2^j + \delta, \quad \delta + \beta \equiv \vartheta \pmod{1}, \quad p_j = \exp(L\rho_j^2/2). \quad (12.3)$$

Then $h_j = 1/\rho_j$ is an analytic function of 2^{-j} , while the phase is constant. After subtracting the known logarithmic terms, the remaining errors are combinations of $j^\ell 2^{-mj}$.

Conjecture 12.5 (Confluent Gaussian- q filter). *For each truncation order r , there is a stable finite filter whose shift polynomial is a product of repeated factors*

$$\prod_{m=1}^r (E - 2^{-m})^{d_m}, \quad (12.4)$$

which annihilates every error mode $j^\ell 2^{-mj}$ with $0 \leq \ell < d_m$. Its coefficients can be written as derivatives or confluent limits of the Gaussian- q -binomial Lagrange weights

$$w_{r,j}(q) = (-1)^{r-j} \frac{q^{\binom{r-j+1}{2}}}{(q; q)_j (q; q)_{r-j}}. \quad (12.5)$$

Applied to (12.3), this filter recovers $\kappa_0 - \gamma - \Psi(\vartheta)$ with a rigorously controlled error of arbitrarily high order in 2^{-j} .

This would create a genuinely new Richardson pipeline: the data are global derivative norms, the target is a pointwise endpoint periodic function, and the logarithmic multiplicities are handled by confluent q -Lagrange interpolation.

12.5 Base- b and Pascal–Rvachev extensions

The Pascal–Rvachev hierarchy in the corpus replaces the single sinc exponent by binomial multiplicities. The present analysis suggests two extensions.

Problem 12.6 (Norm ladders for higher Rvachev ranks). For each hierarchy level, identify a spatial refinement formula with disjoint or finitely overlapping cells, determine whether normalized derivatives are equimeasurable with a fixed profile, and calculate the resulting derivative weight. If exact equimeasurability fails, seek a finite-state transfer operator for value distributions.

Conjecture 12.7 (Higher-rank lognormal moment geometry). *For the rank- r Pascal–Rvachev profile, the normalized spectral energy moments are q -hyperfactorial sequences whose Hankel determinants factor into products of q -Pochhammer symbols with exponents polynomial in the index. The associated orthogonal polynomials should be multilevel q -binomial deformations of Stieltjes–Wigert polynomials.*

12.6 Formalization roadmap

The exact statements in Sections 3, 5 and 8 are unusually suitable for Lean because their proofs reduce to finite sums, affine changes of variables, Parseval, and determinant algebra. A practical order is:

1. expose the derivative tiling as a public theorem with a cellwise corollary;
2. prove the pushforward/equimeasurability identity for arbitrary measurable test functions;
3. derive L^p , sup-norm, total-variation, and level-set corollaries;
4. formalize the Stieltjes substitution and beta-transform identity for the totalized inverse;
5. prove the spectral moments by the existing Fourier differentiation API;
6. formalize the Vandermonde determinant and finite q -binomial orthogonality;
7. defer the all-orders asymptotic operator until the inverse derivative-jet bounds have a stable public interface.

13 Conclusion

The dyadic derivative formula contains more information than its usual pointwise reading suggests. Because the cells are disjoint and all have the same affine Jacobian, it is an exact renormalization fixed point for value distributions. The entire derivative tower has one absolute profile, amplified by $2^{n(n+1)/2}$ and signed by Thue–Morse. This immediately produces an exact norm spectrum in every rearrangement-invariant space and makes the inverse Fabius function the level-set law of every derivative.

Two dual transforms then emerge. In physical space, the large- p norms are a beta average of the inverse function. The beta dilation converges to an exponential dilation, whose Mellin symbol is $\Gamma(1 + s)$; Bell polynomials, zeta values, and Bernoulli polynomials are therefore forced into the asymptotic operator. Combining it with the lower-Lambert endpoint coordinate reveals a universal Euler-constant shift and lets global norms perform phase tomography on the periodic saddle correction.

In frequency space, Parseval turns the same derivative scale into the moment sequence $2^{n(n+1)}$. The infinite sinc product thus supplies an explicit, highly nonclassical Stieltjes–Wigert representing measure. Its integer zeros distinguish it from the lognormal representative even though all polynomial moments agree; its q -Pearson law retains the full sinc-square factor. The resulting q -binomial orthogonal polynomials and Gaussian Lagrange rules give a concrete bridge between spectral energy, moment indeterminacy, and the q -calculus already pervasive in the Fabius theory.

The most promising next step is to identify the sinc-square measure inside the Nevanlinna family and to formalize the exact equimeasurability/beta-transform layer. The first is a substantive analytic frontier; the second is a compact, high-value Lean target that would make several apparently numerical phenomena exact library theorems.

A Recursive TeX corpus ledger

The following 68 paths were present under Analysis/FabiusFunction/docs at commit 0341bbf91c b867f4510bc8431cf59878208de771. Archived versions were included in the collision audit but interpreted according to the archive's provenance ledger.

#	Path
1	FabiusFunction_Mathematical_Glossary/FabiusFunction_Mathematical_Glossary.tex
2	Fabius_Function_and_Rvachev_Up/Fabius_Function_and_Rvachev_Up.tex
3	Non_Elementarity_of_the_Fabius_Function/Non_Elementarity_of_the_Fabius_Function.tex
4	archive/early-notes/FabiusAsymptotic/FabiusAsymptotic.tex
5	archive/early-notes/K-foldsummationoverthesignedThue-Morsesequence/K-foldsummationoverthesignedThue-Morsesequence.tex
6	archive/glossary/FabiusFunction_Mathematical_Glossary-2/FabiusFunction_Mathematical_Glossary.tex
7	archive/glossary/FabiusFunction_Mathematical_Glossary/FabiusFunction_Mathematical_Glossary.tex
8	archive/lean-walkthrough/Fabius_Function_Lean_Walkthrough/Fabius_Function_Lean_Walkthrough.tex
9	archive/lean-walkthrough/fabius_lean_walkthrough/fabius_lean_walkthrough.tex
10	archive/primary-exposition/Fabius_Function_and_Rvachev_Up-2/Fabius_Function_and_Rvachev_Up.tex
11	archive/primary-exposition/Fabius_Function_and_Rvachev_Up-3/Fabius_Function_and_Rvachev_Up.tex
12	archive/primary-exposition/Fabius_Function_and_Rvachev_Up/Fabius_Function_and_Rvachev_Up.tex
13	archive/primary-exposition/Fabius_and_Rvachev_Up-2/Fabius_and_Rvachev_Up.tex
14	archive/primary-exposition/Fabius_and_Rvachev_up/Fabius_and_Rvachev_up.tex
15	archive/primary-exposition/Fabius_function_and_up_function/Fabius_function_and_up_function.tex
16	archive/primary-supplements/Fabius_Function_Missing_Parts-2/Fabius_Function_Missing_Parts.tex
17	archive/primary-supplements/Fabius_Function_Missing_Parts/Fabius_Function_Missing_Parts.tex
18	archive/q-calculus/Fabius_q_Special_Functions/Fabius_q_Special_Functions.tex
19	archive/q-calculus/fabius-q-special-functions/fabius-q-special-functions.tex
20	archive/research-frontiers/Fabius_Dyadic_Asymptotic_Bridge/Fabius_Dyadic_Asymptotic_Bridge.tex
21	archive/research-frontiers/Fabius_Dyadic_Formulae_and_Alternative_Representations/Fabius_Dyadic_Formulae_and_Alternative_Representations.tex
22	archive/research-frontiers/Fabius_Dyadic_Formulae_to_Asymptotics/Fabius_Dyadic_Formulae_to_Asymptotics.tex
23	archive/research-frontiers/Fabius_Dyadic_q_Connections/Fabius_Dyadic_q_Connections.tex
24	archive/research-frontiers/Fabius_Thue_Morse_Convergence_Rate/Fabius_Thue_Morse_Convergence_Rate.tex
25	archive/research-frontiers/Fabius_Thue_Morse_Convergence_Rates/Fabius_Thue_Morse_Convergence_Rates.tex

#	Path (continued)
26	archive/research-frontiers/Primary_Exposition_Gap_Register/Primary_Exposition_Gap_Register.tex
27	archive/research-frontiers/Repeated_Integration_and_Rvachev_Up/Repeated_Integration_and_Rvachev_Up.tex
28	archive/research-frontiers/Rvachev_Up_from_Repeated_Integration/Rvachev_Up_from_Repeated_Integration.tex
29	archive/research-frontiers/Thue_Morse_Formula_Atlas-1/Thue_Morse_Formula_Atlas.tex
30	archive/research-frontiers/Thue_Morse_Formula_Atlas-2/Consolidated_Formula_Atlas_for_the_Thue_Morse_Sequence.tex
31	archive/research-frontiers/Thue_Morse_Formula_Atlas-3/Thue_Morse_Formula_Atlas.tex
32	archive/research-frontiers/non-formalized-research-frontiers/non-formalized-research-frontiers.tex
33	archive/standalone-studies/Non_Elementarity_of_the_Fabius_Function/Non_Elementarity_of_the_Fabius_Function.tex
34	archive/standalone-studies/Small_Argument_Asymptotics/Small_Argument_Asymptotics.tex
35	archive/standalone-studies/fabius-inverse-dyadic-closed-form/fabius-inverse-dyadic-closed-form.tex
36	fabius_lean_walkthrough/fabius_lean_walkthrough.tex
37	non-formalized-research-frontiers/non-formalized-research-frontiers.tex
38	non-formalized-research-frontiers/drafts/Fabius_Arithmetic_Rays_Frontier_Report/Fabius_Arithmetic_Rays_Frontier_Report.tex
39	non-formalized-research-frontiers/drafts/Fabius_Half_Integer_Spectral_Frontier_Report/Fabius_Half_Integer_Spectral_Frontier_Report.tex
40	non-formalized-research-frontiers/drafts/Fabius_Inverse_Frontier_Report_Source_and_PDF/Fabius_Inverse_Frontier_Report.tex
41	non-formalized-research-frontiers/drafts/Fabius_Rvachev_Frontier_Report-2/Fabius_Rvachev_Frontier_Report.tex
42	non-formalized-research-frontiers/drafts/Fabius_Rvachev_Frontier_Report/Fabius_Rvachev_Frontier_Report.tex
43	non-formalized-research-frontiers/drafts/Fabius_Rvachev_New_Frontiers/Fabius_Rvachev_New_Frontiers.tex
44	non-formalized-research-frontiers/drafts/Fabius_Rvachev_Thue_Morse_Frontier_Results/Fabius_Rvachev_Thue_Morse_Frontier_Results.tex
45	non-formalized-research-frontiers/drafts/Spectral_Arithmetic_Pascal_Rvachev_Hierarchy/Spectral_Arithmetic_Pascal_Rvachev_Hierarchy.tex
46	non-formalized-research-frontiers/drafts/Thue_Morse_Formula_Atlas/Thue_Morse_Formula_Atlas.tex
47	non-formalized-research-frontiers/drafts/fabius_frontier_new_results/fabius_frontier_new_results.tex
48	non-formalized-research-frontiers/drafts/fabius_frontier_results_bundle/fabius_frontier_results.tex
49	non-formalized-research-frontiers/drafts/fabius_frontier_spectral_endpoint_report_bundle/fabius_frontier_spectral_endpoint_report.tex
50	non-formalized-research-frontiers/drafts/rvachev_up_fourier_decay/Rvachev_Up_Fourier_Decay-2/Rvachev_Up_Fourier_Decay.tex
51	non-formalized-research-frontiers/drafts/rvachev_up_fourier_decay/Rvachev_Up_Fourier_Decay-3/Fourier_decay_of_Rvachev_up.tex
52	non-formalized-research-frontiers/drafts/rvachev_up_fourier_decay/Rvachev_Up_Fourier_Decay_Comparative_Audit/Rvachev_Up_Fourier_Decay_Comparative_Audit.tex

#	Path (continued)
53	non-formalized-research-frontiers/drafts/rvachev_up_fourier_decay/Rvachev_Up_Fourier_Decay_Gentle_Guide/Rvachev_Up_Fourier_Decay_Gentle_Guide.tex
54	non-formalized-research-frontiers/drafts/rvachev_up_fourier_decay/Rvachev_Up_Fourier_Decay_Second_Wave_Audit/Rvachev_Up_Fourier_Decay_Second_Wave_Audit.tex
55	non-formalized-research-frontiers/drafts/rvachev_up_fourier_decay/asymptotic-decay-rate-of-an-infinite-product-of-sinc-functions/asymptotic-decay-rate-of-an-infinite-product-of-sinc-functions.tex
56	non-formalized-research-frontiers/drafts/rvachev_up_fourier_decay/rvachev_up_fourier_decay-1/rvachev_up_fourier_decay.tex
57	non-formalized-research-frontiers/drafts/rvachev_up_fourier_decay/rvachev_up_fourier_decay-4/rvachev_up_fourier_decay.tex
58	non-formalized-research-frontiers/drafts/rvachev_up_fourier_decay/rvachev_up_fourier_decay-5/Rvachev_Up_Fourier_Decay_Final.tex
59	non-formalized-research-frontiers/drafts/rvachev_up_fourier_decay/rvachev_up_fourier_decay-6/Rvachev_Up_Fourier_Decay_Final_Synthesis.tex
60	non-formalized-research-frontiers/drafts/rvachev_up_fourier_decay/rvachev_up_fourier_decay-7/rvachev_up_fourier_decay_final.tex
61	non-formalized-research-frontiers/drafts/rvachev_up_fourier_decay/rvachev_up_fourier_decay-8/rvachev_up_fourier_decay_final.tex
62	papers/AIF_2017__67_2_637_0/AIF_2017__67_2_637_0-corrected.tex
63	papers/AIF_2017__67_2_637_0/AIF_2017__67_2_637_0.tex
64	papers/arXiv-1502.06738v2/Lacunary_products.tex
65	papers/arXiv-1702.05442v1/09-Function.tex
66	papers/arXiv-1702.06487v3/157-Arithmetic-v3.tex
67	papers/arXiv-2211.13570v1/document.tex
68	papers/ubiq15/ubiq15-corrected.tex

B Reproducibility files

The accompanying archive contains:

- Fabius_Derivative_Norm_Spectrum.tex – this source;
- Fabius_Derivative_Norm_Spectrum.pdf – the rendered report;
- numerical_experiments.py – commented reproduction code;
- figures/ – PDF and PNG versions of the three generated figures;
- data/derivative_norm_ratios.csv;
- data/spectral_moment_checks.csv;
- data/beta_operator_checks.csv;
- data/hankel_determinant_checks.csv; and
- data/run_summary.txt.

The default reproduction command is

```
python numerical_experiments.py
```

The grid size, number of sinc factors, derivative order, and output directory are configurable through command-line options documented by

```
python numerical_experiments.py --help
```

References

- [1] V. Reshetnikov, *ProveIt: Analysis/FabiusFunction*, repository snapshot at commit 0341bbf91cb867f4510bc8431cf59878208de771, 27 August 2026. <https://github.com/VladimirReshetnikov/ProveIt/tree/0341bbf91cb867f4510bc8431cf59878208de771/Analysis/FabiusFunction>
- [2] J. Arias de Reyna, *On the function of Fabius*, English translation and corrected source in docs/papers/AIF_2017__67_2_637_0; original Spanish article, *Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales de Madrid* **76** (1982), 367–376.
- [3] J. Arias de Reyna, *Arithmetic of the Fabius function*, *Journal de Théorie des Nombres de Bordeaux* **30** (2018), 721–749; source imported in docs/papers/arXiv-1702.06487v3.
- [4] J. Fabius, *A probabilistic example of a nowhere analytic C^∞ -function*, *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete* **5** (1966), 173–174.
- [5] B. Jessen and A. Wintner, *Distribution functions and the Riemann zeta function*, *Transactions of the American Mathematical Society* **38** (1935), 48–88.
- [6] T. S. Chihara, *A characterization and a class of distribution functions for the Stieltjes–Wigert polynomials*, *Canadian Mathematical Bulletin* **13** (1970), 529–532.
- [7] J. S. Christiansen, *The moment problem associated with the Stieltjes–Wigert polynomials*, *Journal of Mathematical Analysis and Applications* **277** (2003), 218–245.
- [8] J. S. Christiansen and E. Koelink, *Self-adjoint difference operators and classical solutions to the Stieltjes–Wigert moment problem*, *Journal of Approximation Theory* **140** (2006), 1–26; <https://arxiv.org/abs/math/0504456>.
- [9] R. Koekoek, P. A. Lesky, and R. F. Swarttouw, *Hypergeometric Orthogonal Polynomials and Their q -Analogues*, Springer, 2010.
- [10] N. H. Bingham, C. M. Goldie, and J. L. Teugels, *Regular Variation*, Cambridge University Press, 1987.
- [11] S. Mandelbrojt, *Séries adhérentes, régularisation des suites, applications*, Gauthier–Villars, Paris, 1952.
- [12] F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark, editors, *NIST Handbook of Mathematical Functions*, Cambridge University Press, 2010.
- [13] D. Popa, *The complete asymptotic evaluation for Mellin convolution operators*, *Constructive Approximation* **58** (2023), 253–270.