

Arithmetic Classification of Geometric Observation Masks

Research note prepared for Vladimir Reshetnikov with OpenAI

1 October 2026

Abstract

We classify universal comparison of arbitrary finite or countable observation masks for every geometric ratio $0 < q < 1$. Nontrivial comparisons beyond discarding observations occur only when a positive integer power of q^{-1} is an integer. If d is the least such power, the exact order is prefix-count dominance separately in the residue classes modulo d ; deterministic residue maps realize every allowed comparison. Outside this countable exceptional set of ratios, universal dominance is exactly mask inclusion. For any fixed comparison not arising from inclusion, its admissible ratios form a locally finite subset of $(0, 1)$. The theorem permits overlapping masks, unequal cardinalities, arbitrary randomized kernels combining observations, and arbitrary independent additive noise.

1 The experiment and the integer-power distances

Fix $q \in (0, 1)$, $c > 0$, and $\beta \in \mathbb{R}$. Let I be a finite initial segment of $\mathbb{N}$ or all of $\mathbb{N}$. Under Q , the independent coordinates X_i have caps and densities

$$a_i = cq^i, \quad q_{\beta, a_i}(x) = \frac{e^{-\beta x}}{Z_{\beta}(a_i)} \mathbf{1}_{(0, a_i)}(x), \quad Z_{\beta}(a) = \int_0^a e^{-\beta x} dx.$$

Let Z be arbitrary independent real noise, and put $T = \sum_{i \in I} X_i + Z$. The noise may be zero. For every nonnegative Borel w with $0 < \mathbb{E}_Q w(T) < \infty$, let P_w have density $w(T)/\mathbb{E}_Q w(T)$ relative to Q .

A mask $E \subseteq I$ observes $V_E = (X_i)_{i \in E}$. Masks may overlap, be empty or countably infinite, and have different cardinalities. We write $E \succeq_q L$ when one Borel Markov kernel from V_E to V_L works under every P_w . This is the universal Blackwell order of these experiments. The same kernel must work throughout the weight family; it does not observe T or Z .

For the observed sum $U_E = \sum_{i \in E} X_i$, set $\mu_E = \text{Law}_Q(U_E)$. The general mask theorem [1] gives

$$E \succeq_q L \iff \mu_E = \mu_L * \nu \text{ for some probability measure } \nu. \quad (1)$$

The factor is compactly supported. The theorem also shows that requiring one kernel for all positive-probability lower thresholds $T \leq t$ is equivalent to requiring every weight. Its proof reduces each vector experiment to its sum, then identifies an independent difference from compact-transform identities; arbitrary noise cancels through its characteristic function near zero. We use this criterion for necessity and verify the realizing maps directly below.

Define

$$H_q = \{k \geq 1 : q^{-k} \in \mathbb{N}\}. \quad (2)$$

Lemma 1.1. *Either H_q is empty or $H_q = d\mathbb{N}$ for its least element d . In the second case $M = q^{-d}$ is an integer at least two.*

Proof. If $a, b \in H_q$ and $g = \gcd(a, b)$, Bézout's identity gives

$$q^{-g} = (q^{-a})^u (q^{-b})^v \in \mathbb{Q}_{>0}$$

for suitable integers u, v . A positive integer power of this rational number is an integer; its reduced denominator must therefore be one. Hence $g \in H_q$. If d is the least positive element, $\gcd(d, k) \in H_q$ forces $d \mid k$ for every $k \in H_q$. Conversely every positive multiple of d belongs. Since $q < 1$, its integer power M is at least two. $\square$

2 The complete classification

Theorem 2.1. *For the model above, the following classifications hold.*

1. If $H_q = \emptyset$, then $E \succeq_q L$ if and only if $L \subseteq E$.
2. If $H_q = d\mathbb{N}$, then $E \succeq_q L$ if and only if, for every $j \in I$,

$$|\{i \in E : i \leq j, i \equiv j \pmod{d}\}| \geq |\{i \in L : i \leq j, i \equiv j \pmod{d}\}|. \quad (3)$$

In case 2, pair the r th target index ℓ in each residue class with the r th source index e in that class. Condition (3) ensures $e \leq \ell$. A realizing map takes $X_e \bmod a_\ell$ for that target and discards unmatched source coordinates. In unit coordinates $U_i = X_i/a_i$, it is

$$U_e \mapsto \{M^{(\ell-e)/d} U_e\}, \quad M = q^{-d}, \quad (4)$$

independent of c and β . Case 1 is realized by coordinate projection.

2.1 Zero multiplicities give necessity

The coordinate transform is the entire function

$$F_i(z) = \frac{e^{(z-\beta)a_i} - 1}{(z - \beta)Z_\beta(a_i)},$$

with its removable value at $z = \beta$. At

$$z_j = \beta + \frac{2\pi i}{a_j},$$

it vanishes precisely when $i \leq j$ and $q^{-(j-i)}$ is an integer, including the ratio one at $i = j$. Each such zero is simple. For $i > j$ the cap ratio lies strictly between zero and one and cannot give a zero.

For an infinite mask, the sum transform F_E is the locally uniformly convergent product of coordinate transforms. Indeed, on every compact disk,

$$|F_i(z) - 1| \leq |z| a_i e^{|z|a_i}, \quad \sum_i a_i < \infty. \quad (5)$$

The product agrees with the sum transform by uniform convergence of the coordinate sums. Near a fixed z_j , sufficiently late factors lie uniformly within distance $1/2$ of one. Their analytic logarithms converge uniformly by (5); hence their product is nonzero. The finitely many other nonvanishing factors are also nonzero on a small neighborhood. Consequently the multiplicity of F_E at z_j is exactly the finite number of vanishing coordinate factors.

Under dominance, (1) gives $F_E = F_L F_\nu$ with F_ν entire. Thus the source zero multiplicity is at least the target multiplicity. If H_q is empty, those multiplicities are simply $\mathbf{1}_{\{j \in E\}}$ and $\mathbf{1}_{\{j \in L\}}$, proving $L \subseteq E$. If $H_q = d\mathbb{N}$, Lemma 1.1 identifies them with the residue-class prefix counts in (3). This proves necessity for arbitrary randomized kernels combining the entire source observation.

2.2 Independent residues give sufficiency

Suppose (3) holds. By the r th target index in any residue class there are at least r source indices in that class. The pairing stated in the theorem therefore uses distinct source indices and covers every target. For each pair $e \rightarrow \ell$,

$$\frac{a_e}{a_\ell} = M^{(\ell-e)/d} \in \mathbb{N}, \quad X_e = a_\ell J_e + R_\ell, \quad R_\ell = X_e \bmod a_\ell.$$

Factoring the density on the residue cells shows that R_ℓ has the target coordinate law and is independent of J_e . Independence of source coordinates implies independence of the entire residue vector from all digits and unmatched source coordinates. In the countable case this follows on finite cylinder sets.

The nonnegative sum

$$D = \sum_{e \rightarrow \ell} a_\ell J_e + \sum_{e \in E \text{ unmatched}} X_e$$

converges and is bounded by $\sum_{e \in E} a_e$. It satisfies

$$U_E = \sum_{\ell \in L} R_\ell + D, \quad D \text{ independent of } R_L, \quad \text{Law}_Q(R_L) = \text{Law}_Q(V_L).$$

Thus $\mu_E = \mu_L * \text{Law}(D)$, proving dominance by (1).

The stated deterministic map itself works. The total-law identity and compact-transform cancellation give

$$\mu_{L^c} = \text{Law}(D) * \mu_{E^c}.$$

Since the source complement is independent of the source vector, $D + U_{E^c}$ is independent of R_L and has the target-complement law. Adding the independent noise shows that the joint laws of (T, R_L) and (T, V_L) agree. Multiplying this identity by any $w(T)$ proves the required pushforward under every reweighting. This argument permits overlap and infinitely many observations without an infinite sequence of swaps. It is a distributional identity, not a pathwise identification of target coordinates in the original sample. If H_q is empty and $L \subseteq E$, projection proves sufficiency. Theorem 2.1 follows.

3 Exceptional ratios and isolated comparisons

Corollary 3.1. *The ratios with $H_q \neq \emptyset$ are exactly*

$$\mathcal{Q} = \{M^{-1/d} : M \geq 2, d \geq 1 \text{ integers}\}. \quad (6)$$

This set is countable and dense in $(0, 1)$. Outside it, universal dominance is exactly inclusion. Among rational ratios, its elements are precisely the reciprocal integers; every transcendental ratio lies outside it.

Proof. The set identity is the definition of an integer inverse power. If $q = p/s$ in lowest terms with $p > 1$, no power $(s/p)^k$ is integral. A ratio in (6) is algebraic, excluding transcendental q . For density, fix $\alpha > 0$ and choose integers M_d nearest to $e^{\alpha d}$. Then $M_d^{-1/d} \rightarrow e^{-\alpha}$, and $M_d \geq 2$ eventually. $\square$

The least d must be used: $q = 4^{-1/2} = 1/2$ has $d = 1$, whereas $q = 1/\sqrt{2}$ has $d = 2$. In the latter case the order separates into odd and even chains. For instance, source mask $\{1, 2\}$ dominates target mask $\{2, 3\}$ by retaining index 2 and taking the residue of index 1 modulo the cap at index 3. This is a classwise pairing, not the globally sorted pairing across both classes.

Corollary 3.2 (Local finiteness). *Fix masks with $L \not\subseteq E$. The set of ratios $q \in (0, 1)$ for which $E \succeq_q L$ is locally finite. Thus only inclusion comparisons can persist throughout a nonempty open interval of ratios.*

Proof. Choose $j \in L \setminus E$. At z_j , the target transform vanishes. Necessity forces a source zero, which must come from an index $i \in E$ with $i < j$. Hence every admissible ratio belongs to

$$\bigcup_{\substack{i \in E \\ i < j}} \{M^{-1/(j-i)} : M \geq 2 \text{ an integer}\}. \quad (7)$$

There are finitely many source indices below j . On any compact subinterval $[a, b] \subset (0, 1)$, each corresponding integer satisfies $b^{-(j-i)} \leq M \leq a^{-(j-i)}$, so only finitely many are possible. A locally finite set contains no open interval. Inclusion works for every q by projection. $\square$

If $j = 1$, the union in (7) is empty. For $j \geq 2$, it also gives the bound $q \leq 2^{-1/(j-1)}$. In particular the admissible ratios of a fixed non-inclusion comparison can accumulate only at zero, never at one.

Corollary 3.3 (Antisymmetry). *For any fixed $q \in (0, 1)$, universal equivalence in both directions holds if and only if $E = L$.*

Proof. Outside $\mathcal{Q}$, use mutual inclusion. Inside it, two-way dominance makes every classwise prefix count equal. Successive differences within each residue class recover mask membership, so the masks coincide. $\square$

4 Scope and source context

Universal comparison requires one kernel for every total-sum weight or lower threshold. It does not preclude a comparison for one fixed conditioning rule or a specified pair of hypotheses when the ratio is nonexceptional. Distinct comparable masks may give equal divergence values for particular weights, including $w = 1$.

The preceding mask report [1] established the convolution criterion and reciprocal-integer case. This note gives the complete arithmetic consequence for all geometric ratios. The repository’s arbitrary-mask question [4] also asks about finer fixed-conditioning behavior; those asymptotic questions remain distinct from the universal order.

Editorial note (ProveIt, 2026-10-01). For ratios q no positive power of which is rational, case 1 of Theorem 2.1 also follows from Theorem `thm:separated` of the repository article `../../spectra-and-arithmetic/Arithmetic_Rigidity_off_Resonance_Geometric_Uniform_Laws/` (filed 2026-09-30, batch 69 of `docs/incoming/`; unreviewed), which this note does not cite. The caps cq^i , $i \in E$, are then pairwise rationally incommensurable, and that theorem says that a sum of independent uniforms of widths a_j divides their sum exactly when $a_j = cq^{\kappa(j)}/n_j$ for an injection κ into E and integers $n_j \geq 1$; for $a_j = cq^j$, $j \in L$, this forces $\kappa(j) = j$ and $n_j = 1$, that is, $L \subseteq E$. It is stated for centred uniforms without tilt, but translation does not affect divisibility, and a common exponential tilt carries a factorization $\mu_E = \mu_L * \nu$ to one of the tilted laws and back. The present note adds the ratios with a rational but no integer power of $1/q$, such as $q = 2/3$. Its exceptional set $\mathcal{Q}$ is the part of the countable dense set of ratios with a rational power (that article’s Proposition `prop:full`) on which some power of $1/q$ is an integer, and the residue classes of case 2 are a sub-mask analogue of the rationality classes in that article’s Theorem `thm:channels` ($q = p^{-e/h}$, p prime), which classifies all uniform divisors of the whole series.

Blackwell comparison and convolution methods have classical antecedents [2, 3]. The proofs here make no general priority claim. They are ordinary mathematical arguments, not formalized or externally refereed results.

Editorial note (ProveIt, 2026-10-01). This note is the fifth of eight notes written in one series and filed together on 2026-10-01 in the repository’s Fabius drafts tree (batch 72 of its docs/incoming/; unreviewed). In the order written they are `../Exact_Fabius_Mask_Order/`, `../Universal_Garbling_Classification/`, `../Universal_Fabius_Mask_Criterion/`, `../Uniform_Smoothing_Mask_Stabilization/`, `../Arithmetic_Geometric_Mask_Order/`, `../Exact_Fixed_Conditioning_Order/`, `../Strict_Fabius_Conditioning_Order/` and `../Proportional_Fabius_Mask_Edgeworth/`. The preceding report [1], source `universal_fabius_mask_criterion.tex`, is filed as `../Universal_Fabius_Mask_Criterion/`, and [4] is `../Critical_Complements_Sharp_Information_Loss/`. At $d = 1$, Theorem 2.1 is Theorem 2.1 of [1], re-proved here with the same tail bound and zero-multiplicity argument; read as one series, Theorem 2.1 is the primary statement of the universal order. Its residue maps contain those of the first note, `../Exact_Fabius_Mask_Order/`, for geometric caps.

References

- [1] *Universal Comparison of Fabius Observation Masks*, research companion prepared for Vladimir Reshetnikov with OpenAI, 1 October 2026. Source `universal_fabius_mask_criterion.tex`, Theorems 1.1 and 2.1.
- [2] D. Blackwell, *Equivalent comparisons of experiments*, *Annals of Mathematical Statistics* **24** (1953), 265–272. doi:10.1214/aoms/1177729032.
- [3] E. N. Torgersen, *Comparison of translation experiments*, *Annals of Mathematical Statistics* **43** (1972), 1383–1399. doi:10.1214/aoms/1177692372.
- [4] V. Reshetnikov, ProveIt, *Critical Complements in Fabius Conditioning*, source at commit `7421a4ca60fd`, rechecked 1 October 2026. Pinned source on GitHub. Full identifiers are included in the package.