

Positivity at the First Feedback Boundary in Thue Morse Pressure

Research note prepared for Vladimir Reshetnikov with OpenAI

1 October 2026

Abstract

We study the coefficient of degree $4m$ in integer Thue–Morse pressure, where nonlinear eigenvalue feedback first enters. An exact regularization separates the boundary eigenfunction response into a strictly positive finite resolvent and an absolutely convergent mixed inverse-orbit sum. A strict dyadic tangent comparison and a lattice local limit argument give a positive diagonal asymptotic. An explicit analytic estimate proves positivity for every $m \geq 70$, and exact integer enclosure certificates cover $2 \leq m \leq 69$. Together they prove that the degree- $4m$ pressure coefficient is positive for every integer $m \geq 2$. This continues the positive triangle strictly below degree $4m$ without changing the earlier reports.

Status. This is unrefereed ordinary mathematics with exact arithmetic certificates, not a Lean formalization. No worldwide priority claim or general formula for the first later negative coefficient is made. The proofs distinguish the infinite-order estimates from their finite verification complement.

1 The boundary result

Use the doubling-map pressure $p_m(c)$ of $m \log \cos^2 \pi(x - c)$ in the sum-over-preimages convention. Set $a = \tan t$ and

$$(V_a f)(x) = \sum_{2y=x \bmod 1} (\cos \pi y + a \sin \pi y)^{2m} f(y).$$

Let $h_a(0) = 1$ normalize its local analytic Perron eigenfunction. Write

$$R_m = \frac{2(2^{2m} - 1)\zeta(2m)}{\pi^{2m}}, \quad H_{m,r} = [a^{2r}]h_a(1/2), \quad E_{m,r} = [t^{2m+2r}]p_m(t/\pi).$$

The preceding companion proves $H_{m,r}, E_{m,r} > 0$ whenever $1 \leq r < m$ [2]. Here the new response is $H_{m,m}$ and the new pressure coefficient is $E_{m,m} = [t^{4m}]p_m(t/\pi)$.

Theorem 1.1. *For every integer $m \geq 2$,*

$$H_{m,m} > 0, \quad [t^{4m}]p_m(t/\pi) > 0.$$

Moreover, put $b_j = \tan(\pi/2^{j+1})$ and $B(a) = \prod_{j \geq 1} (1 + ab_j)$. There is a unique $a_ > 0$ satisfying*

$$\sum_{j \geq 1} \frac{a_* b_j}{1 + a_* b_j} = 1.$$

With

$$C = \frac{B(a_*)}{a_*}, \quad p_j = \frac{a_* b_j}{1 + a_* b_j}, \quad \sigma^2 = \sum_{j \geq 1} p_j(1 - p_j),$$

we have the asymptotic

$$H_{m,m} \sim \frac{2(2C/\pi)^{2m}}{\sqrt{4\pi m \sigma^2}} \quad (m \rightarrow \infty). \quad (1)$$

The constants are defined by convergent products and sums; approximate values are $a_* \approx 0.822918569054$, $C \approx 4.046860022$, and $\sigma^2 \approx 0.7041449521$. These decimals are orientation only and are not used in the sign proof. The finite-complement certification is described in Section 7. Combined with the earlier triangle, this proves $H_{m,r}, E_{m,r} > 0$ for every $1 \leq r \leq m$. The response range is sharp as a universal statement: $H_{2,3} = -488/27 < 0$.

Editorial note (ProveIt, 2026-10-01). The pressure statement $E_{m,m} > 0$ is re-proved, by a separate large-order bound and separate finite certificates, as the case $r = m$ of Theorem 1.1 of `../Thue_Morse_Integer_Pressure_Full_Range/` ($E_{m,r} > 0$ for every $m \geq 2$ and $1 \leq r < 2m$), so it does not depend on the trace archives discussed in the note in Section 7. The response statement $H_{m,m} > 0$ and the asymptotic (1) are this note's own; `../Thue_Morse_Integer_Pressure_Higher_Positive/` had proved the case $m = 2$ ($H_{2,2} = 536/27$), and `../Thue_Morse_Integer_Pressure_Beyond_Boundary/` extends the asymptotic to every fixed offset s (the case $s = 0$ is (1)).

2 The exact boundary regularization

Put $d = 2m$, $A = V_0$, and $h = F_{0,m}$. The atomic basis is

$$F_{j,m}(x) = \sin^{2m}(\pi x) \sum_{n \in \mathbb{Z}} (\pi(x+n))^{-(2m-j)}, \quad 0 \leq j \leq 2m-2.$$

It spans the invariant Fourier space with indices $1-m, \dots, m-1$ and satisfies $AF_{j,m} = 2^{-j}F_{j,m}$. In particular $h(0) = 1$, $h(1/2) = R_m$, and the rank-one spectral projection is $Pf = f(0)h$. These facts follow by splitting the periodization into even and odd integers; the Fourier model and its local pressure interpretation are reviewed in [1, 2]. Let $Q = I - P$ and let $\text{ev } f = f(1/2)$.

The eigenvalue and pressure satisfy

$$\lambda_m(a) = 1 + a^d h_a(1/2), \quad p_m(t/\pi) = 2m \log \cos t + \log \lambda_m(\tan t). \quad (2)$$

At degree d , the raw inverse-orbit coefficient no longer has the absolute-convergence estimate used below the boundary. Its exact regularization is as follows.

Lemma 2.1. *If $C_N = [a^d](V_a^N h)(1/2)$, then*

$$H_{m,m} = \lim_{N \rightarrow \infty} (C_N - NR_m^2). \quad (3)$$

For every finite N ,

$$(V_a^N h)(1/2) = \sum_{w \in \mathbb{Z} + 1/2} (\pi w)^{-d} \prod_{j=1}^N \left(1 + a \tan \frac{\pi w}{2^j} \right)^d. \quad (4)$$

Proof. Evaluation at zero gives $(V_a f)(0) = f(0) + a^d f(1/2)$. Since $A^N h = h$,

$$(V_a^N h)(0) = 1 + NR_m a^d + O(a^{d+1}).$$

The normalized iterates converge locally uniformly analytically to h_a , by the simple dominant eigenvalue and finite-dimensional spectral projection. Dividing by the last display proves (3) by Cauchy's coefficient formula.

For (4), first use centered inverse representatives $|w| < 2^{N-1}$. The cosine product contributes $(2^N \sin(\pi w/2^N))^{-d}$. Substituting the periodization defining $h(w/2^N)$ cancels its sine factor and unfolds the representatives into all half-integers. The finite tangent product is periodic in w with period 2^N and has no pole there, so the resulting series is absolutely convergent. $\square$

Let $t_j(w) = \tan(\pi w/2^j)$ and define

$$K_m(w) = [a^d] \prod_{j \geq 1} (1 + at_j(w))^d - \sum_{j \geq 1} t_j(w)^d.$$

The subtracted monomials choose all d powers from just one tangent factor. Every remaining monomial uses at least two factors, with each selected exponent below d .

Proposition 2.2. *The mixed orbit series is absolutely convergent, and*

$$H_{m,m} = P_m + \sum_{w \in \mathbb{Z} + 1/2} (\pi w)^{-d} K_m(w), \quad (5)$$

where the finite resolvent is

$$P_m = \sum_{n \geq 0} (\text{ev } A^n V_d h - R_m^2) = \text{ev}(I - A)^{-1} Q V_d h.$$

Here $V_a = \sum_{k=0}^d a^k V_k$, and the inverse is restricted to Q .

Proof. Let $g(x) = \sin \pi x$ on $[0, 1]$, extended periodically, and let W_k replace each weight of V_k by its absolute value. With $s = \sin(\pi x/2)$ and $c = \cos(\pi x/2)$,

$$Ag = sc(c^{d-1} + s^{d-1}) \leq \frac{1}{2}g \quad (d \geq 4).$$

For $1 \leq k < d$ and bounded f ,

$$W_k |f| \leq \frac{1}{2} \binom{d}{k} \|f\|_\infty g. \quad (6)$$

Indeed, after factoring sc from $c^{d-k}s^k + s^{d-k}c^k$, the remaining sum is at most $c^{d-2} + s^{d-2} \leq 1$, by weighted AM-GM.

Fix a composition $k_1 + \dots + k_s = d$ with $s \geq 2$ and all $k_i \geq 1$. Its absolute insertion words have the form

$$\text{ev } A^{n_0} W_{k_1} A^{n_1} \dots A^{n_{s-1}} W_{k_s} h.$$

The final power of A after the last insertion fixes h . Inequality (6) and $Ag \leq g/2$ bound each word by a constant times $2^{-\sum n_i}$. Summing all zero-degree gaps is finite, and there are only finitely many compositions of d . Tonelli's theorem therefore proves absolute convergence of the mixed orbit monomials, including convergence of their finite truncations.

The pure terms in C_N are exactly $\sum_{n=0}^{N-1} \text{ev } A^n V_d h$. As $(V_d h)(0) = R_m$, their spectral projection is $R_m h$, giving the secular part NR_m^2 . Subtraction in (3) yields (5). $\square$

3 A strictly positive pure correction

Proposition 3.1. *For every $m \geq 2$,*

$$R_m(1 - R_m) < P_m \leq \frac{4}{3} R_m(1 - R_m). \quad (7)$$

The upper equality holds only at $m = 2$.

Proof. Write $F_{0,m}(x + 1/2) = q_m(u)$, where $u = \sin^2 \pi x$. The polynomial q_m has degree $m - 1$ and all coefficients strictly positive. A direct induction proves this: $q_1 = 1$, and twice differentiating the periodized secant expression gives

$$\begin{aligned} q_{m+1}(u) = \frac{1}{2m(2m+1)} [& 4u(1-u)^2 q_m''(u) \\ & + 2(1-u)(1+(4m-2)u) q_m'(u) + 2m(1+2mu) q_m(u)]. \end{aligned} \quad (8)$$

Applied to u^k , its numerator has coefficients

$$2k(2k-1), \quad 2m+2k(4m+1-4k), \quad 4(m-k)^2$$

at u^{k-1}, u^k, u^{k+1} . They are nonnegative for $0 \leq k \leq m-1$ and propagate strict positivity. Formula (8) follows from differentiating $(1-u)^{-m}q_m(u)$ and using $u' = 2\sin z \cos z$, $u'' = 2(1-2u)$, with $z = \pi x$. Also $q_m(0) = R_m$ and $q_m(1) = 1$, proving $R_m \leq h \leq 1$.

The monomials have positive atomic expansions

$$u^k = \sum_{j=k}^{m-1} \beta_{m-k,j-k} F_{2j,m}(x), \quad \beta_{p,\ell} = [z^{2\ell}](z/\sin z)^{2p} > 0.$$

This is the principal-part decomposition of $\csc^{2(m-k)} z$ after division by $\sin^{2m} z$. The remaining entire periodic difference vanishes at imaginary infinity. Positivity of $\beta_{p,\ell}$ follows from the Euler product for $z/\sin z$. Thus

$$F_{0,m}(x+1/2) = \sum_{j=0}^{m-1} d_{m,j} F_{2j,m}(x), \quad d_{m,j} > 0, \quad d_{m,0} = R_m.$$

The exact identity $V_d = AT_{1/2}$ now gives

$$P_m = \sum_{j=1}^{m-1} \frac{d_{m,j} R_{m-j}}{4^j - 1} > 0.$$

At $x = 1/2$ the sine and cosine weights agree, so $\text{ev } V_d h = \text{ev } Ah = R_m$. Consequently

$$\sum_{j=1}^{m-1} 4^{-j} d_{m,j} R_{m-j} = R_m - R_m^2.$$

Comparing termwise with $1 < (1-4^{-j})^{-1} \leq 4/3$ proves (7) and its equality statement. $\square$

4 A strict tangent comparison and an exponential tail

For positive odd n , put $x_j(n) = |\tan(n\pi/2^{j+1})|/n$. The tangent theorem in [2] proves that its decreasing rearrangement is weakly log-majorized by $(b_1, b_2, \dots)$, where $b_1 = 1$. We need the following strict refinement.

Lemma 4.1. *For every odd $n \geq 3$, $x(n)$ is weakly log-majorized by*

$$(\theta, b_2, b_3, \dots), \quad \theta = \frac{1+\sqrt{2}}{3} < 1.$$

Hence, for every $a > 0$,

$$B_{x(n)}(a) := \prod_j (1 + ax_j(n)) \leq \frac{1+\theta a}{1+a} B(a). \quad (9)$$

Proof. We recall the finite structure of the earlier tangent proof. Set $L = \lfloor \log_2 n \rfloor$ and $s_j = |\sin(n\pi/2^{j+1})|/(n \sin(\pi/2^{j+1}))$. The sorted ratios $x_j^\downarrow/b_j$ cross 1 only once, from below to above. After that crossing, the largest k terms are exactly the first k original terms, with product ratio

$$s_k^2 \prod_{j=2}^{k-1} s_j \quad (k \geq 2).$$

These statements follow from the initial cotangent bounds and the tail counting argument in [2]; all s_j lie in $(0, 1]$.

The largest normalized tangent is at most θ . For $n = 3$, its value is $\tan(3\pi/8)/3 = \theta$. For $n \geq 5$, write $q = 2^{L+1} \geq 8$. The original early terms are below $2/\pi$ and the late terms below $1/n$; the only remaining term obeys

$$x_{L+1} \leq \frac{\cot(\pi/(2q))}{q-1} < \frac{2q}{\pi(q-1)} \leq \frac{16}{7\pi} < \theta.$$

Also $s_2 \leq \theta$: it is equal to θ for $n = 3$, and for $n \geq 5$ it is below $1/(5 \sin(\pi/8)) < 4/5 < \theta$.

Before the sorted crossing, the first ratio is at most θ and every later ratio is at most 1. After the crossing, the original-prefix identity is at most θ because it contains s_2 , with the case $k = 2$ giving s_2^2 . Every sorted prefix product is therefore at most θ times its b -prefix product. The comparison sequence is decreasing because $\theta > b_2$.

Weak log-majorization implies all elementary-symmetric coefficient inequalities: apply the increasing symmetric convex function $z \mapsto \sum_{|I|=r} \exp(\sum_{i \in I} z_i)$ to the logarithms of finite prefixes. This elementary majorization step is proved in [2]. Geometric tails justify the monotone limit to infinite products and give (9). $\square$

Define the mixed central coefficient

$$M_d(x) = [a^d] \prod_j (1 + ax_j)^d - \sum_j x_j^d.$$

For a nonnegative summable sequence x and any $a > 0$, marking two distinct factor indices gives

$$M_d(x) \leq \frac{d^2}{d-1} a^{2-d} e_2(x) B_x(a)^d. \quad (10)$$

Indeed, a mixed degree- d monomial with exponents k_j is marked $\sum_{i < j} k_i k_j \geq d-1$ times. The marked generating function is

$$d^2 a^2 B_x(a)^d \sum_{i < j} \frac{x_i x_j}{(1 + ax_i)(1 + ax_j)}.$$

Each canceled product has nonnegative coefficients, so evaluation at $a > 0$ bounds its degree- d coefficient and proves (10).

The crucial summability constant is

$$T := \sum_{\substack{n \geq 3 \\ n \text{ odd}}} e_2(x(n)) \leq \frac{\pi^2(4 + 3\sqrt{2})}{8} < 11. \quad (11)$$

To verify it, repeat the absolute mixed-word argument at $m = 1, d = 2$, where the atomic eigenfunction is 1. Now $Ag \leq g/\sqrt{2}$, $W_1 1 = 2g$, and $W_1 g \leq \sqrt{2}g$. Summing the two zero-degree gaps bounds the absolute mixed sum by $2\sqrt{2}/(1 - 1/\sqrt{2})^2$. Its orbit expression is exactly $(32/\pi^2) \sum_{n \geq 1, n \text{ odd}} e_2(x(n))$, including the nearest term. This proves (11).

At the saddle a_* , set $q = (1 + \theta a_*)/(1 + a_*) < 1$. Equations (9)–(11) show that the entire absolute remote mixed tail in the normalized tangent scale is at most

$$\frac{d^2}{d-1} a_*^2 T C^d q^d = O(d C^d q^d). \quad (12)$$

5 The positive diagonal asymptotic

The function $aB'(a)/B(a) = \sum_j ab_j/(1 + ab_j)$ increases continuously from 0 to infinity, so a_* exists uniquely. Let X be the countable sum of independent Bernoulli variables with success probabilities $p_j = a_*b_j/(1 + a_*b_j)$. Its mean is 1, variance is $\sigma^2 \in (0, \infty)$, and probability generating function is $B(a_*z)/B(a_*)$. The sum is finite almost surely, and all exponential moments exist because B is entire. If S_d is a sum of d independent copies, then

$$g_d := [a^d]B(a)^d = C^d \Pr(S_d = d).$$

The elementary lattice local limit argument gives

$$\Pr(S_d = d) \sim \frac{1}{\sqrt{2\pi d\sigma^2}}. \quad (13)$$

For completeness, X assigns positive mass to both 0 and 1, so its characteristic function has modulus strictly below 1 away from $2\pi\mathbb{Z}$. Near zero, the centered characteristic function is $1 - \sigma^2 u^2/2 + O(u^3)$. In Fourier inversion split at a small fixed neighborhood of zero. Outside it the integral decays exponentially in d . Inside it, the substitution $u = v/\sqrt{d}$, a Gaussian majorant, and dominated convergence give (13).

The nearest mixed coefficient is $g_d - \sum_j b_j^d$, where $\sum_j b_j^d < 2$ for $d \geq 2$. Also $C > 2$, using only the first two factors. The remote mixed contribution in (12) is exponentially smaller than (13), and $P_m = O((2/\pi)^d)$ by (7). Substitution in (5), pairing positive and negative half-integers, gives (1).

6 An effective analytic cutoff

The integer-mean mode theorem for countable Bernoulli sums states that the integer mean is a mode; see Gneden's extension of Darroch's rule, Theorem 2 in [3]. It applies to S_d , whose mean is d and variance is at most d . Chebyshev therefore gives

$$\Pr(S_d = d) \geq \frac{3/4}{4\sqrt{d} + 1} \geq \frac{1}{8\sqrt{d}}, \quad g_d \geq \frac{C^d}{8\sqrt{d}}.$$

Here the interval $|S_d - d| < 2\sqrt{d}$ contains at most $4\sqrt{d} + 1$ integers and has probability at least $3/4$.

The exact half-angle certificate in the package proves

$$\frac{4}{5} < a_* < \frac{5}{6}.$$

It encloses the first twelve b_j with rational bounds using $b_{j+1} = b_j/(1 + \sqrt{1 + b_j^2})$, and bounds the remaining mean tail geometrically. Specifically it proves the upper bound 981/1000 for the mean at $a = 4/5$ and the lower bound 1008/1000 at $a = 5/6$. Since $\theta < 81/100$ and q decreases with a_* ,

$$q < \frac{206}{225}.$$

The tail in (12), divided by C^d , is consequently below

$$\frac{275}{36}(d+2) \left(\frac{206}{225}\right)^d.$$

For $d \geq 140$ this is less than $1/(16\sqrt{d})$. At $d = 140$ the exact rational certificate checks

$$\left(\frac{275 \cdot 16}{36}\right)^2 142^2 \cdot 140 \left(\frac{206}{225}\right)^{280} < 1.$$

The expression $(d+2)\sqrt{d}(206/225)^d$ decreases thereafter: its consecutive ratio is at most $(143/142)(281/280)(206/225) < 1$.

Finally $\sum_j b_j^d < 2 < C^d/(16\sqrt{d})$ for $d \geq 140$, using $C > 2$ and $2^d > 32d$. The nearest mixed coefficient therefore strictly exceeds the absolute remote mixed tail. The positive P_m proves $H_{m,m} > 0$ for $m \geq 70$.

At degree $4m$, the feedback formula is exactly

$$E_{m,m} = \sum_{j=0}^m H_{m,j} [t^{4m}] \tan^{2m+2j} t - \frac{R_m^2 + R_{2m}}{2}. \quad (14)$$

The preceding triangle supplies $H_{m,j} > 0$ for $j < m$. The $j = 0$ term is at least $R_m \binom{2m}{m}/3^m \geq 2R_m/3$; the binomial ratio starts at $2/3$ for $m = 2$ and its successive ratio is $(4m+2)/(3m+3) \geq 1$. In contrast, $R_{2m} \leq R_m^2$ and $R_m \leq R_2 = 1/3$, so the two negative terms total at most $R_m/3$. Thus $H_{m,m} \geq 0$ implies $E_{m,m} > 0$, completing the analytic part of Theorem 1.1 for $m \geq 70$.

7 Exact certification of the finite complement

The package uses a fixed-grid enclosure computation whose arithmetic consists solely of exact integers. It never decides a sign from an unbounded numerical approximation. Let $D = 2^{2m-1}$, with Fourier indices $1 - m, \dots, m - 1$. Define integer matrices

$$(B_j)_{k,\ell} = \binom{2m}{m+2k-\ell} [z^j] (1-z)^{m+2k-\ell} (1+z)^{m-2k+\ell}.$$

An entry is defined as zero when $m+2k-\ell$ lies outside $[0, 2m]$. Then $V_a = D^{-1} \sum_j i^j B_j a^j$. The atomic vector is the Eulerian row of order $2m-1$, divided by $(2m-1)!$. Its normalization and eigenvector equation are checked exactly.

Write $h_a = \sum_n i^n v_n a^n$. Below order $2m$, the eigenvalue has no nonconstant term, so

$$(DI - B_0)v_n = \sum_{j=1}^n B_j v_{n-j}, \quad \mathbf{1}^\top v_n = 0.$$

At $n = 2m$, subtract $D(-1)^m R_m v_0$ from the right side. Reflection gives $(v_n)_{-k} = (-1)^n (v_n)_k$, permitting separate even and odd blocks. In the even block, the zero-index equation is replaced by normalization with row $(1, 2, \dots, 2)$; in the odd block normalization is automatic.

Both block inverses are computed fraction-free, with every division checked exact and the final integer identity $CN = qI$ verified. The grid denominator is $S = 2^{256} f^2$, where f is the reduced denominator of v_0 . This makes both v_0 and the feedback forcing exact on the grid. Every later inverse product is rounded downward to the grid, with the rounding inequality checked by integers.

Here is the rigorous error propagation. Use the full-vector ℓ^1 norm on compressed coordinates: weight 1 at index zero and weight 2 at positive indices. Let α_n be the induced norm of the appropriate block inverse, excluding the normalization column in the even case. This is computed exactly from the integer inverse. Since

$$|B_j| \leq \binom{2m}{j} B_0, \quad \|B_j\|_1 \leq D \binom{2m}{j},$$

an integer bound $U_n \geq S\|v_n - \tilde{v}_n/S\|_1$ is propagated by

$$U_n = \left\lceil \alpha_n D \sum_{j=1}^n \binom{2m}{j} U_{n-j} \right\rceil + \begin{cases} 2m-1, & n \text{ even}, \\ 2m-2, & n \text{ odd}, \end{cases} \quad U_0 = 0.$$

The last term bounds the total weighted rounding error. The half-period evaluation has norm one, so its computed value plus or minus U_{2m}/S rigorously encloses $H_{m,m}$. All intermediate vectors and error bounds are saved in compressed replay traces. Independent full rational calculations and full-vector enclosures check selected cases.

All 68 moments $m = 2, \dots, 69$ yield strictly positive rational lower endpoints. Their 4,828 response-order enclosures, baseline vectors, propagated errors, final intervals, and available full-rational comparisons were independently checked. This finite verification proves $H_{m,m} > 0$ throughout the complement of the analytic range. Equation (14) then proves $E_{m,m} > 0$, completing Theorem 1.1. The exact sources, every per-order vector, and the certificate checker are included; the finite computation is a distinct part of the proof.

Editorial note (ProveIt, 2026-10-01). As filed, the per-order traces are missing: this package is the source archive of a split delivery whose five trace archives (the 68 compressed files `data/certificates/parity_mNNN.json.trace.gz`; see `DELIVERY_README.txt`) were not delivered. The trace audit `code/check_trace_coverage.py` and the full replay `code/reproduce_all.py` therefore cannot run on the filed files, and the baseline vectors, the 4,828 propagated error bounds and the final vectors cannot be rechecked. What the filed files show (checked on filing): the 68 interval records `data/certificates/parity_mNNN.json` have strictly positive lower endpoints; exact rational values of $H_{m,m}$ filed with the package lie inside them for the 35 orders $2 \leq m \leq 35$ and $m = 50$; for $m = 68, 69$ an independent full-vector enclosure overlaps them; for $36 \leq m \leq 49$ and $51 \leq m \leq 67$ the filed intervals are the only finite evidence for $H_{m,m} > 0$. The pressure coefficient $E_{m,m}$ is proved again in `../Thue_Morse_Integer_Pressure_Full_Range/` (whose own finite intervals were not delivered either), and its positivity for $2 \leq m \leq 128$ is certified by the filed intervals of `../Thue_Morse_Integer_Pressure_First_Negative_Data/`.

Editorial note (ProveIt, 2026-10-01). The missing traces can be regenerated, byte for byte, with the filed producer `code/check_boundary_parity.cpp`: see the section “Reconstructing the trace archives” of `README.md`. On filing, the traces for $m = 2, 3, 10, 20, 35, 50, 69$ were regenerated on a copy: the interval records agree with the filed ones, the compressed traces reproduce the hashes recorded in `data/trace_checks.json`, and the trace audit passes on them. The full set (68 orders, about 45 MB compressed, a few minutes of computation) was not regenerated.

8 Later negative coefficients and open directions

The response theorem cannot uniformly extend to $r = m + 1$. For $m = 2$, set $x = a^2$ and write $\Lambda(a) = \lambda_2(a)$. The pressure cubic in $[1]$, under $\rho = \Lambda/(1 + a^2)^2$ and $\cos 2t = (1 - x)/(1 + x)$, gives

$$8\Lambda^3 - 2(7 - x)(1 + x)\Lambda^2 + (7 - 3x)(1 + x)^3\Lambda - (1 + x)^6 = 0.$$

Its derivative in Λ at $(\Lambda, x) = (1, 0)$ is 3, so its local branch is unique. Exact substitution and a separate Fourier recursion agree on

$$\Lambda = 1 + \frac{x^2}{3} + 8x^3 + \frac{536}{27}x^4 - \frac{488}{27}x^5 - \frac{90472}{243}x^6 + O(x^7).$$

Since $\Lambda = 1 + a^4 h_a(1/2)$, this proves $H_{2,3} = -488/27 < 0$. Pressure at the corresponding degree ten remains positive: $E_{2,3} = 105448/2025$. Thus response and pressure sharpness are different questions.

A separate direct rational phase-eigenvalue recursion locates the first negative pressure degree *above* $2m$ for the following small moments:

m	2	3	4	5	6	7	8	9	10	11	12	13	14
degree	12	18	24	32	38	44	52	58	64	72	78	84	92

The lower even coefficients are already negative, so the qualification “above $2m$ ” matters. These are exact finite observations, not an all-order law. In particular, neither a limiting ratio nor a rounding formula for the first negative degree is asserted. Whether pressure alone stays positive throughout $1 \leq r < 2m$ remains open here; the finite data are compatible with that extension. Positivity of each individual atomic-basis response coefficient throughout $r \leq m$ is also a separate, unproved question.

Editorial note (ProveIt, 2026-10-01). Pressure positivity throughout $1 \leq r < 2m$ is proved for every $m \geq 2$ in `../Thue_Morse_Integer_Pressure_Full_Range/`. The table above agrees with the certified first negative degrees of `../Thue_Morse_Integer_Pressure_First_Negative_Data/` ($2 \leq m \leq 128$) and `../Thue_Morse_Integer_Pressure_Sign_Changes/` ($2 \leq m \leq 20$), and `../Thue_Morse_Integer_Pressure_First_Negative/` proves $N_m/m \rightarrow \gamma \in (6.662966, 6.662968)$, with a $\log \log m$ correction and a rounding rule valid away from the even lattice.

Editorial note (ProveIt, 2026-10-01). This note is the fourth of thirteen packages written in one series and filed together on 2026-10-01 beside the repository manuscript `../Thue_Morse_Integer_Pressure/` in the Fabius drafts tree (batch 72 of the repository’s docs/incoming/; unreviewed). In logical order they are the positivity series `../Thue_Morse_Integer_Pressure_First_Positive/`, `../Thue_Morse_Integer_Pressure_Higher_Positive/`, `../Thue_Morse_Integer_Pressure_Positive_Triangle/`, `../Thue_Morse_Integer_Pressure_Feedback_Boundary/`, `../Thue_Morse_Integer_Pressure_Beyond_Boundary/`, `../Thue_Morse_Integer_Pressure_Linear_Region/` and `../Thue_Morse_Integer_Pressure_Full_Range/`, and the sign series, which imports the full-range theorem: `../Thue_Morse_Integer_Pressure_Sign_Changes/`, `../Thue_Morse_Integer_Pressure_Sign_Densities/`, `../Thue_Morse_Integer_Pressure_Negative_Bound/`, `../Thue_Morse_Integer_Pressure_First_Negative/` and `../Thue_Morse_Integer_Pressure_Cluster_Asymptotics/`, with the dataset `../Thue_Morse_Integer_Pressure_First_Negative_Data/` (no article). An earlier version of `../Thue_Morse_Integer_Pressure_Full_Range/`, *The Full Pressure Positivity Range for Large Integer Orders* (orders $m \geq 4096$), delivered in the same batch, is superseded by it and was not filed. Here [1] is `../Thue_Morse_Integer_Pressure/` (the version cited differs from the filed manuscript only by dated editorial changes, none of them mathematical), and [2] is `../Thue_Morse_Integer_Pressure_Positive_Triangle/`.

References

- [1] *Integer Pressure and a Missing Taylor Coefficient*. ProveIt research manuscript, 28 September 2026. Inspected version: commit 6a98f89bac85cc6b4ba5d5c3b63996037d9824e1, blob 1973fd17245864bb8a83e1b1d90416ac032ba7c1. Versioned repository source.
- [2] *The Full Positive Triangle in Integer Thue Morse Pressure*. Separate research note prepared for Vladimir Reshetnikov with OpenAI, 1 October 2026. The tangent single-crossing theorem, full $r < m$ response and pressure positivity, and exact certificates.
- [3] A. Gnedin, *Cross Modality of the Extended Binomial Sums*, 2024, Theorem 2 (integer-mean mode rule for finite and countable Bernoulli sums). <https://arxiv.org/html/2408.06477v1>