

Exponential Accuracy and Stokes Transport for Inverse Harmonic Transseries

Large-order coefficients, exact-rational certificates,
and a convergent inverse jump expansion

Research note prepared for Vladimir Reshetnikov

29 September 2026

Abstract

The *ProveIt* companion on combinatorial transseries gives every coefficient of the inverse harmonic function and proves a remainder at each fixed algebraic order. This article develops the missing large-order and exponentially accurate layers for that particular family. Writing

$$H(x) = \psi(x+1) + \gamma, \quad X = e^{y-\gamma}, \quad H^{-1}(y) = W(X) - \frac{1}{2},$$

we prove a finite formula for the complete large-order expansion of the coefficients of $W(X) \sim X + \sum_{n \geq 1} h_n X^{1-2n}$. In particular, $h_n \sim 2(-1)^n \Gamma(2n)/(2\pi)^{2n}$. A general factorial-convolution theorem supplies the proof rather than a fit to computed coefficients.

An exact Fermi–Dirac integral yields signed forward remainders and rationally checkable inverse enclosures. If u_M is the local inverse of the forward expansion truncated after M terms, and $M - \pi X$ stays bounded, we prove the sharp law $u_M - W(X) = (-1)^M \sqrt{X} e^{-2\pi X} (1 + \mathcal{O}(X^{-1}))$, including its first correction and an all-orders construction. The exact Borel kernel also determines the forward Stokes jump. Transporting that jump through inversion gives an explicit convergent expansion of every inverse exponential sector. Known nonlinear resurgence theorems identify the formal inverse as a Borel-summable resurgent series. Exact symbolic checks, 190-digit experiments, and independent exact-rational enclosure checks accompany the proofs.

Scope and status. These are proved, explicitly scoped extensions of the inspected repository material, not a claim to have settled all of its transseries questions. Binet’s formulas and general resurgence closure are existing results, credited below. Global priority for the specialized theorems has not been established. No new Lean formalization is claimed.

Repository snapshot: 04e06e032dff1966513bfba316966d52db3646b4.

Source, PDF, verification programs, exact certificates, and recorded outputs are included in the accompanying archive.

Contents

1	The research question and the repository boundary	2
1.1	Why this family is a useful test case	2
1.2	What is established here	2
1.3	Relation to existing work	2
2	Normalization and the formal inverse	3
3	A factorial endpoint-dominance theorem	4
4	The complete large-order expansion of the inverse coefficients	5
5	An exact integral and explicit inverse certificates	6
5.1	The centered Binet formula	6
5.2	A bracket that works before computing any correction	7
6	Sharp exponential accuracy of forward-truncation inversion	8
6.1	The main error law	8
6.2	Concentration of the exact remainder	9
6.3	Transport from the forward remainder to the inverse error	10
7	Exact-rational numerical certification	11
7.1	Removing the floating-point premise	11
7.2	A concrete certified interval	12
7.3	An implementation-level algorithm	12
8	Borel summation and the resurgent inverse	12
8.1	The exact Borel kernel	12
8.2	Passing through nonlinear inversion	13
9	Every inverse Stokes sector, with a convergence theorem	14
9.1	Orientation and the exact forward jump	14
9.2	The inverse transport formula	14
9.3	The first three nonlinear sectors	16
9.4	Why the positive-real error is not a Stokes jump	16
10	Reproducible checks and their logical status	16
10.1	Late coefficients	17
10.2	The sharp inverse error	17
10.3	Complex-sector checks	17
11	Further research questions and concrete next steps	18
12	Synthesis	20
A	Finite coefficient algorithms	21
A.1	Exact coefficients without expanding an exponential symbolically	21
A.2	Computing a prescribed sharp-error coefficient	21
A.3	Computing inverse jump coefficients	21
B	Reproduction, provenance, and audit boundary	21
B.1	Contents of the archive	21
B.2	Exact repository checkpoint	22
B.3	What the verification does and does not establish	22
	References	23

1 The research question and the repository boundary

1.1 Why this family is a useful test case

A transseries calculation can answer several quite different questions. It can determine a formal coefficient, prove an asymptotic expansion to a fixed order, locate the least term of a divergent expansion, assign a sum to the expansion, or identify the exponentially small difference between two such sums. None of these steps should be silently substituted for another.

The harmonic inverse is particularly suitable for separating these tasks. Its coefficients have a compact exact formula, its forward function has an explicit integral representation, and its Borel singularities lie on an arithmetic lattice. Yet nonlinear inversion already changes the large-order coefficients and creates arbitrarily many exponential sectors. Thus it is a nontrivial example in which the formal, analytic, and computational layers can all be made explicit.

The inspected starting point is the companion volume *Combinatorial Transseries and Their Inverses: Gamma quotients, finite exponential sums, moment sectors, moving saddles, arithmetic sheets, and q-products* in the *ProveIt* repository [1]. Its theorem labelled `t2:thm:harmonic-inverse` gives

$$h_n = -\frac{1}{2n-1} [z^n] \exp((2n-1)\mathcal{C}(z)), \quad \mathcal{C}(z) = -\sum_{j \geq 1} \frac{B_{2j}(1/2)}{2j} z^j, \quad (1.1)$$

and a fixed-order inverse remainder. The same volume explicitly distinguishes such results from global resurgence or Borel summation. It also records that formal inversion cannot supply Stokes constants not already determined by the original function. The group's README describes the distinction between its formal-calculus, analytic-inversion, and Lean interfaces [2].

This article takes those distinctions seriously. We retain the repository's normalization and prove the additional statements needed in this one family. The result is a scoped answer to its summation and large-order gap, not a claim that every family in the companion has the same solution.

1.2 What is established here

The central results are as follows. [Theorem 3.2](#) is a general endpoint-dominance theorem for coefficients of $\exp(N_n \mathcal{C})$ when $N_n = O(n)$ and the input has $\Gamma(2n)$ growth. [Theorem 4.1](#) applies it to obtain every large-order correction to h_n . [Theorem 5.2](#) gives a completely explicit inverse certificate, and [theorem 6.1](#) determines the sharp error of an optimally scaled *forward-truncation inverse*. [Theorems 8.2](#) and [9.1](#) identify the summation class and all inverse Stokes sectors.

There are three deliberate boundaries. First, the exponentially accurate approximation u_M is not the direct partial sum $X + \sum_{n \leq M} h_n X^{1-2n}$. We do not transfer a sharp error statement between them without proof. Second, a positive-real least-term error $e^{-2\pi X}$ is not a positive-real Stokes jump: the actual Borel singularities are imaginary. Third, the exact-rational computations below certify numerical enclosures but are not kernel-checked Lean proofs.

1.3 Relation to existing work

General asymptotic reversion is classical; the distinction between formal inversion and remainder control is discussed, for example, by Fabijonas and Olver [4]. Borinsky develops a powerful algebraic calculus for factorially divergent series and their inverses [5]. Our coefficient proof is self-contained and tailored to the growing multiplier $2n-1$ and the growth $\Gamma(2n)$; it should be read in that broader context, not as a claim that factorial asymptotics under inversion is a newly discovered subject.

For resurgence, the nonlinear closure results used here are Sauzin's: closure under convergent substitution and under composition/inversion at infinity, together with the corresponding Borel-summability statements

[6]. Our contribution in that part is the explicit specialization to the centered harmonic inverse and the finite all-sector jump formula. A targeted literature search and repository inspection support this positioning, but are not an exhaustive priority investigation.

2 Normalization and the formal inverse

Put

$$F(w) = \psi(w + \tfrac{1}{2}), \quad C(w) = F(w) - \log w, \quad F(W(X)) = \log X. \quad (2.1)$$

Initially $X \geq 1$ and $w > 0$ are real. Then

$$H^{-1}(y) = W(e^{y-\gamma}) - \tfrac{1}{2}. \quad (2.2)$$

The complex branches needed for Stokes analysis will be specified separately. We reserve $C(w)$ for an exact function and $\mathcal{C}(z)$ for a formal power series:

$$C(w) \sim \mathcal{C}(w^{-2}) = \sum_{j \geq 1} c_j w^{-2j}, \quad c_j = -\frac{B_{2j}(1/2)}{2j}. \quad (2.3)$$

Thus

$$c_1 = \frac{1}{24}, \quad c_2 = -\frac{7}{960}, \quad c_3 = \frac{31}{8064}, \quad c_4 = -\frac{127}{30720}. \quad (2.4)$$

The relevant centered expansion follows from the classical gamma-function formulas [3]; a direct integral proof will be given in [section 5](#).

Proposition 2.1 (Formal inverse and fixed-order meaning). *There is a unique formal Laurent series*

$$W(X) = X + \sum_{n \geq 1} h_n X^{1-2n} \quad (2.5)$$

satisfying $\log(W/X) + \mathcal{C}(W^{-2}) = 0$. Its coefficients are given by (1.1). For every fixed $M \geq 0$, the actual positive inverse satisfies

$$W(X) = X + \sum_{n=1}^M h_n X^{1-2n} + O(X^{-2M-1}) \quad (X \rightarrow +\infty). \quad (2.6)$$

The constants in this assertion may depend on M .

Proof. Set $v = W^{-1}$ and $t = X^{-1}$. The formal equation becomes $v = t \exp(\mathcal{C}(v^2))$. Formal Lagrange inversion gives a unique odd series $v(t) = t + O(t^3)$. For $n \geq 1$, its Laurent version gives

$$[t^{2n-1}] \frac{1}{v(t)} = -\frac{1}{2n-1} [u^{2n}] \exp((2n-1)\mathcal{C}(u^2)),$$

which is (1.1). This Laurent identity also follows directly by the change of variable $t = v \exp(-\mathcal{C}(v^2))$ in formal residues.

For fixed M , substitute the finite candidate into a sufficiently long finite expansion of F . Coefficient cancellation makes the forward residual $O(X^{-2M-2})$. Near the candidate, $F'(w) \asymp X^{-1}$, so the mean value theorem gives (2.6). The integral bounds below justify both the finite expansion and this slope estimate independently of formal manipulations. $\square$

The first coefficients, already present in the repository, are

$$h_1 = -\frac{1}{24}, \quad h_2 = \frac{3}{640}, \quad h_3 = -\frac{1525}{580608}, \quad h_4 = \frac{615881}{199065600}, \quad h_5 = -\frac{3058641}{504627200}. \quad (2.7)$$

It is important that (2.6) alone says nothing about choosing M proportional to X . The next sections supply information uniform in that different regime, using a different approximation.

3 A factorial endpoint-dominance theorem

The coefficient extraction in (1.1) has a growing parameter. A product with one large index and several small indices is much larger than a balanced product, but the number of factors is not fixed in the exponential. Both facts must be controlled.

Lemma 3.1 (Uniform factorial convolution). *Let $b_j = \Gamma(2j) = (2j - 1)!$ for $j \geq 1$. For $n \geq 2$,*

$$\sum_{j=1}^{n-1} b_j b_{n-j} \leq 3b_{n-1}. \quad (3.1)$$

Consequently, for $1 \leq \ell \leq n$,

$$\sum_{\substack{i_1 + \dots + i_\ell = n \\ i_r \geq 1}} b_{i_1} \cdots b_{i_\ell} \leq 3^{\ell-1} b_{n-\ell+1}. \quad (3.2)$$

Proof. For $n \geq 4$, the two endpoint terms in (3.1) contribute $2b_{n-1}$. Every remaining normalized term is

$$\frac{b_j b_{n-j}}{b_{n-1}} = \frac{2n-2}{\binom{2n-2}{2j-1}} \leq \frac{2n-2}{\binom{2n-2}{3}}.$$

Their sum is at most $6(n-3)/((2n-3)(2n-4)) \leq 1$. The cases $n = 2, 3$ are immediate. Induct on ℓ , applying the binary bound to the final convolution, to obtain (3.2). $\square$

Theorem 3.2 (Endpoint extraction with a growing exponential parameter). *Let $A, D, B > 0$, and let $\mathcal{C}(z) = \sum_{j \geq 1} c_j z^j$ be a complex formal series satisfying*

$$|c_j| \leq AD^j \Gamma(2j) \quad (j \geq 1). \quad (3.3)$$

Let $0 < N_n \leq Bn$, and define

$$d_n = \frac{1}{N_n} [z^n] \exp(N_n \mathcal{C}(z)), \quad E_k(N) = [z^k] \exp(N \mathcal{C}(z)).$$

For every fixed integer $K \geq 0$,

$$d_n = \sum_{k=0}^K c_{n-k} E_k(N_n) + O\left(D^n \Gamma(2n) n^{-K-1}\right). \quad (3.4)$$

The implied constant may depend on A, D, B, K , but not on n or on the particular choices $N_n \leq Bn$.

Proof. Write $N = N_n$. Expansion by the number of factors gives

$$d_n = \sum_{\ell=1}^n \frac{N^{\ell-1}}{\ell!} \sum_{\substack{i_1 + \dots + i_\ell = n \\ i_r \geq 1}} c_{i_1} \cdots c_{i_\ell}. \quad (3.5)$$

Assume $n > 2K$. A composition containing a part $n - k$ with $0 \leq k \leq K$ has exactly one such part. Selecting it in the exponential, or selecting its position in (3.5), shows that its total contribution is exactly $c_{n-k} E_k(N)$. Thus the displayed sum in (3.4) is exact for all compositions with one of those large parts. We bound the rest in two groups.

Many factors. By theorem 3.1, the absolute contribution of length $\ell = t + 1$ divided by $D^n b_n$ is at most

$$A \frac{(3AN)^t}{(t+1)!} \frac{b_{n-t}}{b_n}. \quad (3.6)$$

For $t \leq n/2$, the $2t$ factors in b_n/b_{n-t} are each at least n , so this is bounded by $A(3AB/n)^t/(t+1)!$. Summing over $K+1 \leq t \leq n/2$ gives $O(n^{-K-1})$. For $t > n/2$,

$$\frac{b_{n-t}}{b_n} \leq \frac{\Gamma(n)}{\Gamma(2n)} \leq n^{-n}.$$

Summing (3.6) over this range gives at most $An^{-n} \exp(3ABn)$, smaller than every fixed negative power of n .

A bounded number of factors, but no sufficiently large part. Fix $2 \leq \ell \leq K+1$. Every remaining composition has a largest part j with $n/\ell \leq j \leq n-K-1$. There are at most ℓ choices for its position. Applying (3.2) to the other $\ell-1$ parts bounds the unsigned factorial sum by

$$\ell 3^{\ell-2} \sum_{\lceil n/\ell \rceil \leq j \leq n-K-1} b_j b_{n-j-\ell+2}. \quad (3.7)$$

The summand is log-convex in j , so its maximum is at an endpoint. At the right endpoint it equals $b_{n-K-1} b_{K-\ell+3}$. At the left endpoint, Stirling's formula gives, relative to b_n , an exponentially small quantity times a fixed power of n : its exponential factor is

$$\exp\{-2n\mathcal{H}(1/\ell)\}, \quad \mathcal{H}(a) = -a \log a - (1-a) \log(1-a) > 0.$$

It is therefore $O_K(b_{n-K-1})$ as well. There are at most n summands in (3.7). After restoring $A^\ell D^n N^{\ell-1}/\ell!$ and dividing by $D^n b_n$, the bound is

$$O_K\left(n^\ell \frac{b_{n-K-1}}{b_n}\right) = O_K(n^{\ell-2K-2}) = O_K(n^{-K-1}).$$

There are only finitely many such ℓ . Length one has no omitted contribution. Combining the two groups proves the theorem. $\square$

Remark 3.3. The argument is insensitive to signs or complex phases. It also does not assume convergence of $\mathcal{C}$. At any specified order K , it replaces a coefficient involving the entire prefix $c_1, \dots, c_n$ by $K+1$ large-index coefficients and finite polynomials in $c_1, \dots, c_K$. This is the mechanism behind the complete large-order expansion below.

4 The complete large-order expansion of the inverse coefficients

For $s > 1$, write $\eta(s) = (1 - 2^{1-s})\zeta(s)$. The centered Bernoulli coefficients satisfy the exact identity

$$c_n = 2(-1)^{n+1} \frac{\Gamma(2n)}{(2\pi)^{2n}} \eta(2n). \quad (4.1)$$

It also follows from the moment integral in section 5. Notice that $\eta(2n) = 1 + O(4^{-n})$.

Theorem 4.1 (All orders of late-coefficient asymptotics). *Let h_n be defined by (1.1). For fixed $K \geq 0$, set*

$$\mathcal{R}_K(n) = \sum_{k=0}^K \frac{(-1)^k (2\pi)^{2k} E_k(2n-1)}{\prod_{r=1}^{2k} (2n-r)}, \quad E_k(N) = [z^k] \exp(N\mathcal{C}(z)), \quad (4.2)$$

where the empty product is one. Then

$$h_n = \frac{2(-1)^n \Gamma(2n)}{(2\pi)^{2n}} \left(\mathcal{R}_K(n) + O(n^{-K-1}) \right). \quad (4.3)$$

In particular,

$$\begin{aligned} \frac{h_n(2\pi)^{2n}}{2(-1)^n\Gamma(2n)} &= 1 - \frac{\pi^2}{12n} + \frac{\pi^4/288 - \pi^2/12}{n^2} \\ &\quad + \frac{-\pi^6/10368 - \pi^4/1440 - \pi^2/12}{n^3} + O(n^{-4}). \end{aligned} \quad (4.4)$$

Every coefficient of the expansion in $1/n$ is a finite, effectively computable polynomial in π^2 with rational coefficients.

Proof. Use [theorem 3.2](#) with $D = (2\pi)^{-2}$, any fixed upper bound for $2\eta(2j)$ as A , and $N_n = 2n - 1$. Since $h_n = -d_n$, it gives

$$h_n = - \sum_{k=0}^K c_{n-k} E_k(2n-1) + O\left((2\pi)^{-2n}\Gamma(2n)n^{-K-1}\right).$$

Divide by $2(-1)^n\Gamma(2n)/(2\pi)^{2n}$ and use (4.1). For each fixed k , the factor $\eta(2n-2k)$ differs from one exponentially little. The gamma quotient is precisely the reciprocal product in (4.2). This proves (4.3).

The polynomial $E_k(2n-1)$ has degree at most k in n . Its contribution in (4.2) is therefore $O(n^{-k})$. Expanding the finite rational expression to the requested power of $1/n$ proves existence and computability of the complete expansion. The first three polynomials are

$$E_1(N) = Nc_1, \quad E_2(N) = Nc_2 + \frac{1}{2}N^2c_1^2, \quad E_3(N) = Nc_3 + N^2c_1c_2 + \frac{1}{6}N^3c_1^3.$$

Substitution and expansion give (4.4). □

Corollary 4.2 (Divergence, exact Gevrey order, and least-term scale). *The series $\sum_{n \geq 1} h_n z^n$ has radius of convergence zero and has exact Gevrey order two. The inverse displacement series $\sum_{n \geq 1} h_n X^{1-2n}$ has exact Gevrey order one in X^{-1} , and its ordinary Borel transform has radius of convergence exactly 2π . Moreover, uniformly when $n - \pi X$ stays bounded,*

$$|h_n|X^{1-2n} = 2\sqrt{X}e^{-2\pi X} \left(1 + O(X^{-1})\right). \quad (4.5)$$

In particular, the signs of h_n are eventually $(-1)^n$.

Proof. The leading asymptotic in [theorem 4.1](#) proves eventual nonzero alternating sign and factorial growth. Stirling's formula identifies $\Gamma(2n)$, up to an exponential factor and a power of n , with $(n!)^2$. No smaller Gevrey order can dominate that growth.

Under the convention $\mathcal{B}(X^{-m}) = \xi^{m-1}/\Gamma(m)$, the coefficient of ξ^{2n-2} in the Borel transform of the displacement is $h_n/\Gamma(2n-1)$. Its magnitude is asymptotic to $2(2n-1)/(2\pi)^{2n}$, proving the radius assertion by the root test. Finally, insert $n = \pi X + O(1)$ in Stirling's formula for the leading term of (4.3); it gives (4.5). □

This result describes large-order coefficients. It does *not*, by itself, prove that a direct inverse partial sum has error equal to its least term. Such an assertion requires additional analytic information; the next two sections obtain a sharp result for forward-truncation inverses instead.

5 An exact integral and explicit inverse certificates

5.1 The centered Binet formula

The half-integer centering has a useful analytic consequence: the Bose kernel in Binet's formula becomes a positive Fermi kernel. This gives more than the signs of the Bernoulli coefficients; it gives the sign and size of the exact remainder.

Lemma 5.1 (Exact centered integral and signed remainders). *For $w > 0$,*

$$C(w) = 2 \int_0^\infty \frac{t \, dt}{(t^2 + w^2)(e^{2\pi t} + 1)}. \quad (5.1)$$

Put $a_j = |c_j|$, and, for $M \geq 0$, define

$$P_M(w) = \log w + \sum_{j=1}^M c_j w^{-2j}, \quad E_M(w) = a_{M+1} w^{-2M-2}. \quad (5.2)$$

Then $R_M(w) = F(w) - P_M(w)$ has the exact representation

$$R_M(w) = 2(-1)^M w^{-2M} \int_0^\infty \frac{t^{2M+1} \, dt}{(w^2 + t^2)(e^{2\pi t} + 1)}, \quad (5.3)$$

and consequently

$$0 < (-1)^M R_M(w) < E_M(w), \quad (5.4)$$

$$|R'_M(w)| \leq \frac{2M+2}{w} |R_M(w)| < \frac{2M+2}{w} E_M(w). \quad (5.5)$$

Proof. The digamma duplication identity and Binet formula [3, §§5.5, 5.9] give

$$\psi(w + \tfrac{1}{2}) = 2\psi(2w) - \psi(w) - 2\log 2, \quad \psi(w) = \log w - \frac{1}{2w} - 2 \int_0^\infty \frac{t \, dt}{(t^2 + w^2)(e^{2\pi t} - 1)}.$$

In the integral with argument $2w$, substitute $t = 2s$. The elementary identity

$$\frac{1}{e^u - 1} - \frac{2}{e^{2u} - 1} = \frac{1}{e^u + 1}$$

then proves (5.1). Finite geometric expansion of $(w^2 + t^2)^{-1}$ proves (5.3), with

$$a_j = 2 \int_0^\infty \frac{t^{2j-1} \, dt}{e^{2\pi t} + 1} = \frac{2\Gamma(2j)}{(2\pi)^{2j}} \eta(2j). \quad (5.6)$$

For the last identity, expand the Fermi factor in its alternating exponential series and integrate. Absolute convergence of the resulting integrated series holds since $2j > 1$. Thus the moments agree with the centered Bernoulli coefficients in (4.1).

Positivity of the integrand gives the sign. Replacing $(w^2 + t^2)^{-1}$ by w^{-2} gives the strict bound in (5.4). Differentiating its positive integral representation, the factor w^{-2M} contributes $2M/w$, and the denominator contributes at most $2/w$ relative to its integrand. This proves (5.5). All differentiations are dominated on compact positive w -intervals by an exponentially integrable function. $\square$

5.2 A bracket that works before computing any correction

The trigamma series gives

$$F'(w) = \sum_{k=0}^\infty (w + \tfrac{1}{2} + k)^{-2} > \frac{1}{w + 1/2} > 0. \quad (5.7)$$

The strict inequality follows by comparison with the integral of the decreasing summand. In particular, the forward equation has at most one positive solution.

Theorem 5.2 (Explicit inverse bracket and a posteriori certificate). *For $X \geq 1$, put*

$$a = X - \frac{1}{12X}, \quad I_X = [a, X], \quad d_X = \frac{1}{12X^2} - \frac{1}{24a^2}, \quad \ell_X = \frac{1}{24X^2} - \frac{7}{960X^4}. \quad (5.8)$$

Both d_X and ℓ_X are positive. The equation $F(w) = \log X$ has a unique root $W(X)$ in (a, X) . For every $v \in I_X$ and every $M \geq 0$,

$$|v - W(X)| \leq (X + \frac{1}{2}) (|P_M(v) - \log X| + E_M(v)). \quad (5.9)$$

Suppose, in addition, that

$$E_M(a) < \min(d_X, \ell_X), \quad \frac{(2M+2)E_M(a)}{a} < \frac{1}{X+1/2}. \quad (5.10)$$

Then $P_M(u_M) = \log X$ has a unique root u_M in I_X , and

$$|u_M - W(X)| < (X + \frac{1}{2})E_M(a), \quad \text{sgn}(u_M - W(X)) = (-1)^M. \quad (5.11)$$

Proof. Since $a \geq 11X/12$, the first two quantities in (5.8) immediately give $d_X > 0$; positivity of ℓ_X is also immediate for $X \geq 1$. Applying theorem 5.1 with $M = 0$ gives

$$F(a) - \log X < \log(a/X) + \frac{1}{24a^2} \leq -\frac{1}{12X^2} + \frac{1}{24a^2} = -d_X.$$

The same lemma with $M = 2$ gives $F(X) - \log X > \ell_X$. Continuity and (5.7) prove the existence and uniqueness of $W(X)$ in the open interval.

For $v \in I_X$, the mean value theorem and (5.7) imply

$$|v - W(X)| \leq (X + \frac{1}{2})|F(v) - \log X|.$$

Splitting $F = P_M + R_M$ proves (5.9).

Under (5.10), the endpoint signs persist when F is replaced by P_M . Moreover, throughout I_X ,

$$P'_M(w) = F'(w) - R'_M(w) > \frac{1}{X+1/2} - \frac{(2M+2)E_M(a)}{a} > 0.$$

Thus u_M exists and is unique there. Applying the residual estimate at u_M proves its error bound. Finally,

$$F(u_M) - F(W(X)) = R_M(u_M),$$

whose sign is $(-1)^M$. Strict monotonicity of F proves the asserted sign of the inverse error. $\square$

The uniqueness in this theorem is local to the specified interval. A high-degree truncated forward expression need not be globally monotone on the whole positive axis. The interval and slope conditions are therefore part of the result, not optional numerical details. Conversely, (5.9) needs no monotonicity assumption on P_M and applies to any proposed approximation in I_X .

6 Sharp exponential accuracy of forward-truncation inversion

6.1 The main error law

The finite-order estimates above remain valid when the truncation order depends on the large parameter. Their exact integral representation makes it possible to analyze the important regime $M \asymp X$ without extrapolating a fixed- M big-O constant.

Theorem 6.1 (Sharp optimal-scale inverse error). *Let $X \rightarrow +\infty$ and let $M = M(X)$ be a nonnegative integer such that $M - \pi X$ stays bounded. Define*

$$\sigma = M + 1 - \pi X. \quad (6.1)$$

For all sufficiently large X , the root u_M in theorem 5.2 exists. Uniformly for bounded σ ,

$$u_M - W(X) = (-1)^M \sqrt{X} e^{-2\pi X} \left(1 + \frac{B_1(\sigma)}{X} + O(X^{-2}) \right), \quad (6.2)$$

$$B_1(\sigma) = \frac{\pi}{12} + \frac{2\sigma^2 - 3\sigma + 7/12}{2\pi}. \quad (6.3)$$

More generally, there are effectively computable polynomials $B_j(\sigma)$, of degree at most $2j$, with $B_0 = 1$, such that, for each fixed $J \geq 1$,

$$u_M - W(X) = (-1)^M \sqrt{X} e^{-2\pi X} \left(\sum_{j=0}^{J-1} \frac{B_j(\sigma)}{X^j} + O_J(X^{-J}) \right). \quad (6.4)$$

The constants in the remainders may be chosen uniformly for σ in any prescribed compact interval.

The leading constant in (6.2) is one, whereas the magnitude of an inverse-series term in (4.5) has leading constant two. These are statements about different objects; their difference is not a contradiction. The theorem identifies the actual error of the root of a truncated *forward* equation.

Editorial note (ProveIt, 2026-09-29). The canonical volume `Transseries_And_Inversion/transseries_and_inversion.tex` of this series (directory `Analysis/Transseries/docs/series-and-transseries/`) proves a general least-term theorem, `p0:thm:optimal-truncation`, whose second part bounds the error of the root of an optimally truncated forward equation by a constant times $x^{\beta-1/2}e^{-Ax}$ under a Gevrey-1 remainder hypothesis. [Theorem 6.1](#) is a sharp instance of that kind of statement for the present family: it identifies the leading constant and every algebraic correction. This article does not cite that theorem; the hypotheses were not matched constant by constant. The residual-to-root bracket behind the certificates of [theorem 5.2](#) is related to the machine-checked lemma `Fabius.exists_eq_in_residual_interval` (`Analysis/FabiusFunction/Lean/FabiusFunction/MeanValueBracket.lean`), which this article does not use. The three later packages cited in the note after [research problem 1](#) re-derive the coefficient B_1 of (6.3), with the sign $+\pi/12$, for the same forward-truncation root u_M .

6.2 Concentration of the exact remainder

Lemma 6.2 (The transition at the least forward term). *Let $n = M + 1$, $b = 2\pi w$, and $\tau = n - \pi w$. As $w \rightarrow +\infty$, uniformly for bounded τ ,*

$$R_M(w) = (-1)^M w^{-1/2} e^{-2\pi w} \left(1 + \frac{2\tau^2 - 3\tau + 7/12}{2\pi w} + O(w^{-2}) \right). \quad (6.5)$$

There is a full expansion of the parenthesis in powers of w^{-1} with polynomial coefficients in τ .

Proof. In (5.3), substitute $u = 2\pi t$ and put $s = 2n = b + 2\tau$. If U has the gamma density $u^{s-1}e^{-u}/\Gamma(s)$ on $u > 0$, then

$$R_M(w) = (-1)^M \frac{2\Gamma(s)}{b^s} \left\{ \mathbb{E} \frac{1}{1 + (U/b)^2} + O(2^{-s}) \right\}. \quad (6.6)$$

Indeed, $0 < e^{-u} - (e^u + 1)^{-1} < e^{-2u}$ and the rational factor is at most one. The normalized error is bounded by 2^{-s} exactly.

For $f(v) = (1 + v^2)^{-1}$, direct differentiation gives

$$f(1) = \frac{1}{2}, \quad f'(1) = -\frac{1}{2}, \quad f''(1) = \frac{1}{2}, \quad f'''(1) = 0.$$

Since $s = b + 2\tau$, the gamma moments give

$$\mathbb{E}(U/b - 1) = \frac{2\tau}{b}, \quad \mathbb{E}(U/b - 1)^2 = \frac{1}{b} + O(b^{-2}), \quad \mathbb{E}(U/b - 1)^4 = O(b^{-2}).$$

The fourth derivative of f is bounded on the positive real axis. Taylor's theorem therefore yields

$$\mathbb{E}f(U/b) = \frac{1}{2} + \frac{1/4 - \tau}{b} + O(b^{-2}). \quad (6.7)$$

Stirling's formula with bounded shift gives

$$\frac{\Gamma(b + 2\tau)}{b^{b+2\tau}} = \sqrt{2\pi} b^{-1/2} e^{-b} \left(1 + \frac{2\tau^2 - \tau + 1/12}{b} + O(b^{-2}) \right). \quad (6.8)$$

Multiplication of (6.7) and (6.8) in (6.6) proves (6.5).

For all orders, use Taylor's theorem through degree $2J - 1$. Every derivative of f is bounded on $[0, \infty)$, and the even centered gamma moment gives a remainder $O(b^{-J})$, uniformly for bounded τ . Every moment used in the finite Taylor polynomial is explicit:

$$\mathbb{E}(U/b - 1)^j = b^{-j} \sum_{\ell=0}^j \binom{j}{\ell} (-b)^{j-\ell} (s)_\ell. \quad (6.9)$$

For the other factor, the shifted logarithmic Stirling expansion is

$$\log \frac{\Gamma(b + d)}{\sqrt{2\pi} e^{-b} b^{b+d-1/2}} \sim \sum_{r \geq 1} \frac{(-1)^{r+1} B_{r+1}(d)}{r(r+1)b^r}, \quad d = 2\tau. \quad (6.10)$$

This classical expansion has a uniform fixed-order remainder for bounded d [3]. Expanding its exponential and using (6.9) is a finite algorithm at every requested order. The coefficient at b^{-j} has degree at most $2j$ in τ . The exponentially small Fermi-to-gamma replacement affects none of these coefficients. $\square$

6.3 Transport from the forward remainder to the inverse error

Proof of theorem 6.1. For bounded $M - \pi X$, Stirling's formula in (5.6) gives

$$E_M(a) = O(X^{-1/2} e^{-2\pi X}).$$

Here $a = X - 1/(12X)$, so replacing X by a changes this estimate by a bounded factor tending to one. The quantities d_X, ℓ_X are bounded below by positive constants times X^{-2} . Both conditions (5.10) therefore hold for sufficiently large X .

Write $w = W(X)$ and $\delta = u_M - w$. The certificate first gives $\delta = O(\sqrt{X} e^{-2\pi X})$. The equation $P_M(u_M) = F(w)$ reads

$$0 = F(w + \delta) - F(w) - R_M(w + \delta).$$

On I_X , $F'(w) \asymp X^{-1}$, $F''(w) = O(X^{-2})$, and (5.5) gives $R'_M = O(X^{-1/2} e^{-2\pi X})$. Taylor's theorem, followed by division by $F'(w)$, proves

$$\delta = \frac{R_M(w)}{F'(w)} + O(X e^{-4\pi X}). \quad (6.11)$$

This exponentially smaller error affects none of the coefficients in (6.4).

The fixed-order inverse expansion and the forward derivative give

$$w = X - \frac{1}{24X} + O(X^{-3}), \quad \frac{1}{F'(w)} = w \left(1 + O(w^{-2})\right). \quad (6.12)$$

Thus $\tau = M + 1 - \pi w = \sigma + \pi/(24X) + O(X^{-3})$ and

$$\sqrt{w} e^{-2\pi w} = \sqrt{X} e^{-2\pi X} \left(1 + \frac{\pi}{12X} + O(X^{-2})\right).$$

Substitute [theorem 6.2](#) in [\(6.11\)](#) and use these identities. This gives [\(6.2\)](#) and [\(6.3\)](#).

For any higher fixed order, take sufficiently many terms of the same expansions. The fixed-order expansion of w is legitimate inside these finite Taylor calculations, because its omitted powers can be made arbitrarily small compared with the requested relative power of X^{-1} . Equations [\(6.9\)](#) and [\(6.10\)](#), the finite coefficient formula for W , and reciprocal series for F' give an algorithm for every B_j . The degree bound $2j$ is preserved by these operations: the only dependence on σ comes from the bounded shift in the gamma factor and its moments. This proves [\(6.4\)](#). $\square$

Remark 6.3 (What “optimal-scale” means). The theorem applies to every integer choice with bounded $M - \pi X$, not only the unique best integer at a given X . Minimizing the first correction B_1 as a polynomial in a continuous σ suggests $\sigma = 3/4$, followed by a nearest-integer choice of M . This is an asymptotic guide, not a claim that it determines the exact minimizing integer for every X . The readily implemented choice $M = \lfloor \pi X \rfloor$ is covered uniformly by the theorem.

7 Exact-rational numerical certification

7.1 Removing the floating-point premise

All the coefficients c_j are rational. When X and a candidate v are rational, the sole nonrational term in the residual of P_M is $\log(v/X)$. It can be enclosed by elementary rational arithmetic. For $r > 0$, let $t = (r - 1)/(r + 1)$. Then $|t| < 1$ and

$$\log r = 2 \sum_{k=0}^{N-1} \frac{t^{2k+1}}{2k+1} + \rho_N, \quad (7.1)$$

$$|\rho_N| \leq \frac{2|t|^{2N+1}}{(2N+1)(1-t^2)}. \quad (7.2)$$

This follows by integrating the geometric expansion of $(1 - t^2)^{-1}$, or directly by bounding the tail of the absolutely convergent arctangent-hyperbolic series. For rational r , both endpoints of this enclosure are rational.

Apply [\(7.1\)](#) with $r = v/X$ and add the exact rational sum $\sum_{j=1}^M c_j v^{-2j}$. The signed remainder interval from [\(5.4\)](#) then gives an exact rational enclosure for $F(v) - \log X$. There are two independent ways to finish. One can apply the bound [\(5.9\)](#); alternatively, at two rational endpoints v_- , v_+ one can verify

$$F(v_-) < \log X < F(v_+).$$

Strict monotonicity proves that the true inverse is enclosed. Both checks are implemented in the accompanying certificate program.

The floating-point calculation only proposes a useful candidate. It is not an assumption in the final sign comparisons: an inaccurate candidate would simply fail the exact checks. The proof of each exported interval consists of rational inequalities together with [theorems 5.1](#) and [5.2](#) and [\(7.2\)](#).

7.2 A concrete certified interval

At $X = 10$, so that the exact target is $y = \gamma + \log 10$, the program certifies

$$\begin{aligned} 9.495837994871258428840380076882531132 &< H^{-1}(\gamma + \log 10) \\ &< 9.495837994871258428840380083938321964. \end{aligned} \tag{7.3}$$

Every displayed endpoint is an exact finite decimal, and the rounding was directed outwards. The two strict inequalities were checked with Python's exact `Fraction` arithmetic, using rational Bernoulli coefficients generated by SymPy. Thus (7.3) is not an enclosure inferred from the number of displayed floating-point digits.

The file `verification/certificates.json` also records enclosures for $X = 5, 20, 40$, the truncation orders, the numbers of logarithmic terms used, and the outcomes of both exact sign checks. The corresponding targets are exactly $\gamma + \log X$. To handle an independently prescribed arbitrary real y , one must also enclose $X = e^{y-\gamma}$, or work with a certified enclosure for $y - \gamma$ in the residual. Treating a rounded value of X as exact would solve a slightly different problem.

7.3 An implementation-level algorithm

Choose a truncation order near πX , verify (5.10), and approximate the root of $P_M(v) = \log X$ inside I_X . Convert that approximation to a rational candidate. Increase the number of terms in (7.1) until the logarithmic uncertainty is safely below the analytic remainder. Use (5.9) to obtain rational endpoints, and check their forward signs directly. A failed check calls for a wider interval or more arithmetic precision, not an unproved rounding rule.

This algorithm separates three error sources: root-finding error, logarithm-evaluation error, and the exact forward truncation remainder. For v close to X , the logarithm argument is especially favorable: $t = O(X^{-2})$, so only a modest number of rational logarithmic terms is required, even when the desired absolute enclosure is exponentially narrow.

8 Borel summation and the resurgent inverse

8.1 The exact Borel kernel

We use the ordinary Borel transform at infinity,

$$\mathcal{B} \left(\sum_{m \geq 1} a_m w^{-m} \right) (\xi) = \sum_{m \geq 1} \frac{a_m}{\Gamma(m)} \xi^{m-1}.$$

A Gevrey-one series has a locally convergent transform. A series is called 1-summable in direction θ when that transform continues to a neighborhood of the ray of angle θ , has at most exponential growth there, and its directional Laplace integral exists for sufficiently large w in the dual sector. A resurgent germ has analytic continuation along paths avoiding a specified discrete singular set; here the permitted set is an additive lattice. These definitions distinguish local Borel convergence from continuation and from summability.

Proposition 8.1 (Borel kernel of the centered correction). *The Borel transform of $\mathcal{C}(w^{-2})$ is*

$$\widehat{C}(\xi) = \frac{1}{\xi} - \frac{1}{2 \sinh(\xi/2)}. \tag{8.1}$$

The origin is removable. The other singularities are simple poles at $\omega_k = 2\pi i k$, $k \in \mathbb{Z} \setminus \{0\}$, with

$$\operatorname{Res}_{\xi=\omega_k} \widehat{C}(\xi) = (-1)^{k+1}. \quad (8.2)$$

For $\Re w > 0$,

$$C(w) = \int_0^\infty e^{-w\xi} \widehat{C}(\xi) \, d\xi. \quad (8.3)$$

Proof. The Bernoulli generating function at $1/2$ is

$$\frac{\xi e^{\xi/2}}{e^\xi - 1} = \frac{\xi}{2 \sinh(\xi/2)} = \sum_{m \geq 0} \frac{B_m(1/2)}{m!} \xi^m.$$

Subtract the constant, divide by $-\xi$, and use the vanishing of the odd Bernoulli polynomials at $1/2$. The resulting coefficient of ξ^{2j-1} is $-B_{2j}(1/2)/(2j)! = c_j/\Gamma(2j)$, proving (8.1). The derivative of $2 \sinh(\xi/2)$ at $2\pi i k$ is $(-1)^k$, proving (8.2).

For the exact Laplace identity, use the classical digamma integral

$$\psi(z) = \int_0^\infty \left(\frac{e^{-t}}{t} - \frac{e^{-zt}}{1 - e^{-t}} \right) dt$$

and the integral for $\log w$ obtained by subtracting exponentials divided by t [3, §5.9]. Subtracting the two formulas at $z = w + 1/2$ gives (8.3). The subtraction is made inside the integral: the apparent singularity at zero cancels. At infinity, $\Re w > 0$ gives absolute convergence. This also proves directly that the formal correction has the stated positive-direction Borel sum. $\square$

8.2 Passing through nonlinear inversion

Theorem 8.2 (Resurgence and summation of the harmonic inverse). *Let $\Omega = 2\pi i\mathbb{Z}$. The formal map*

$$\widetilde{G}(w) = w \exp(\mathcal{C}(w^{-2})) = w + O(w^{-1}) \quad (8.4)$$

and its formal compositional inverse $\widetilde{W}(X) = X + \sum_{n \geq 1} h_n X^{1-2n}$ are Ω -resurgent maps tangent to the identity at infinity. They are 1-summable in every direction other than the two imaginary Borel directions. Their directional sums are inverse maps wherever the corresponding local inverses are defined near infinity.

In the positive direction, the sum of $\widetilde{W}$ is the real branch $W(X)$ considered above, for sufficiently large X . It continues uniquely to that branch for all $X \geq 1$.

Proof. The meromorphic function (8.1) continues outside Ω , which is closed, discrete, and stable under addition. On every closed angular sector avoiding the imaginary rays, its growth at infinity is bounded by an exponential, in fact much more mildly. Thus the formal correction is Ω -resurgent and 1-summable in each nonsingular direction.

Apply the nonlinear closure theorems of Sauzin [6, Theorems 3, 5 and 5']: convergent substitution preserves this resurgent class; maps tangent to the identity are closed under composition and inversion; and the corresponding statements hold for 1-summability, with summation commuting with the operations. The hypotheses apply to the exponential in (8.4), and then to its inverse. This proves the first assertions. Multiplication by w only shifts the formal powers and introduces no new nonzero Borel singular locations.

In the positive direction, (8.3) identifies the sum of $\widetilde{G}$ as $G(w) = w e^{C(w)} = e^{F(w)}$. Its positive local inverse satisfies $F(w) = \log X$ and is therefore $W(X)$ by uniqueness in theorem 5.2. The same monotonicity supplies its continuation along $X \geq 1$. $\square$

The statement about Ω is a containment and continuation statement. It does not say that every nonzero point of Ω must be an actual singularity on every continuation sheet of the inverse Borel transform. Conversely, [theorem 4.2](#) proves independently that its initial Borel disk has radius exactly 2π , so the inverse is not made convergent by cancellation of all its nearest singularities.

A general closure theorem establishes membership in a function class, not explicit Stokes data. Those data are accessible here because the original kernel and all its residues are known. We now transport them through inversion without introducing any undetermined constants.

9 Every inverse Stokes sector, with a convergence theorem

9.1 Orientation and the exact forward jump

Consider the positive imaginary Borel ray. Let C_- be the Laplace continuation along a ray with angle just below $\pi/2$, and let C_+ be the continuation along a ray just above $\pi/2$. The subscripts refer to these Borel-ray angles, not to signs of $\Im w$. Use compatible logarithm branches and write

$$F_{\pm}(w) = \log w + C_{\pm}(w).$$

Their common region includes large w in a closed sector strictly inside the lower half-plane. In this region put

$$\lambda = 2\pi i, \quad q(w) = e^{-\lambda w}, \quad |q(w)| < 1. \quad (9.1)$$

The lower Borel contour directed outwards, followed by the upper contour directed inwards, has positive orientation around the enclosed poles. Hence (8.2) gives

$$F_-(w) - F_+(w) = 2\pi i \sum_{k \geq 1} (-1)^{k+1} e^{-2\pi i k w} = \frac{\lambda q(w)}{1 + q(w)}. \quad (9.2)$$

For a finite wedge this is the residue theorem. Its infinite limit is justified by exponential decay of the Laplace factor, using radii separated from the lattice poles. The final residue series converges absolutely for $|q(w)| < 1$.

With the lower-angle continuation from the positive Borel direction, F_- is the continuation of $\psi(w + 1/2)$ from the right half-plane. Thus the numerical tests may evaluate $F_+ = \psi(w + 1/2) - \lambda q/(1 + q)$ directly. This convention also fixes every sign in the inverse formula below.

9.2 The inverse transport formula

Let L be a target in a compatible logarithmic coordinate and let

$$g(L) = F_+^{-1}(L), \quad \widetilde{g}(L) = F_-^{-1}(L), \quad p(L) = g'(L), \quad D = \frac{d}{dL}, \quad q_0 = e^{-\lambda g(L)}. \quad (9.3)$$

The inverse branches are the nearby branches asymptotic to e^L in the specified sector. These are locally defined analytic inverses; no claim of global univalence on the entire complex plane is needed.

Theorem 9.1 (Explicit, convergent inverse jump expansion). *For sufficiently large $g(L)$ in any fixed closed sector strictly inside the lower half-plane,*

$$\widetilde{g}(L) - g(L) = \sum_{n \geq 1} A_n(L) q_0^n, \quad (9.4)$$

where

$$A_n(L) = (-1)^n \sum_{m=1}^n \frac{\lambda^m}{m!} \binom{n-1}{m-1} (D - n\lambda p)^{m-1} p. \quad (9.5)$$

The power denotes repeated application of the differential operator to the function on its right; p is not held constant.

There are constants $r, c > 0$, independent of sufficiently large $w_0 = g(L)$ in the chosen sector, such that the series in the auxiliary variable z with coefficients $A_n(L)$ converges for $|z| < c/|w_0|$. In particular, if

$$\eta_0 = \frac{|w_0 q_0|}{c} < 1, \quad (9.6)$$

then the tail after N sectors obeys

$$\left| \tilde{g} - g - \sum_{n=1}^N A_n q_0^n \right| \leq r \frac{\eta_0^{N+1}}{1 - \eta_0}. \quad (9.7)$$

The condition (9.6) holds automatically sufficiently far out in the specified sector.

Proof. We first identify the coefficients. For an analytic locally invertible F_+ and a perturbation $r_0(w)$, the inverse of $F_+(w) + tr_0(w)$ satisfies the classical inverse-perturbation identity

$$w(L, t) = g(L) + \sum_{m \geq 1} \frac{(-t)^m}{m!} D^{m-1} (p(L) r_0(g(L))^m). \quad (9.8)$$

For completeness, put $v = F_+(w)$, so that $v = L - tr_0(g(v))$ and $w = g(v)$. Apply the Lagrange–Bürmann formula to this local equation for v , with observable g . It gives (9.8). Thus the identity is a local analytic coefficient statement, independent of the particular special function.

In the present case, $r_0(w) = \lambda q(w)/(1 + q(w))$. For each $m \geq 1$,

$$r_0(g)^m = \lambda^m \sum_{n \geq m} (-1)^{n-m} \binom{n-1}{m-1} q_0^n.$$

Moreover, $Dq_0 = -\lambda p q_0$, so induction gives

$$D^{m-1}(p q_0^n) = q_0^n (D - n\lambda p)^{m-1} p.$$

To make coefficient extraction independent of the exact coefficient functions, first replace $q(w)$ in the perturbation by $\varepsilon q(w)$, keeping F_+ fixed. The coefficient of ε^n uses only $1 \leq m \leq n$ in (9.8). It is $A_n(L) q_0^n$, with A_n given by (9.5). This argument is finite at each sector and does not assume that a rearrangement of two unbounded sums is valid.

To prove actual convergence and the tail bound, freeze $w_0 = g(L)$ and introduce an independent variable z . Consider

$$F_+(w_0 + \delta) - F_+(w_0) + \frac{\lambda z e^{-\lambda \delta}}{1 + z e^{-\lambda \delta}} = 0. \quad (9.9)$$

At $(\delta, z) = (0, 0)$ its derivative in δ is $F'_+(w_0) \neq 0$. The analytic implicit function theorem therefore gives an analytic solution $\delta(z)$ with $\delta(0) = 0$. Its Taylor coefficients are the previously obtained A_n by uniqueness of formal implicit solutions.

Here the convergence disk can be controlled uniformly. Choose a fixed small $r > 0$. The sectorial expansion gives

$$F_+(w_0 + \delta) - F_+(w_0) = \frac{\delta}{w_0} + O(|w_0|^{-2}) \quad (|\delta| = r),$$

uniformly as $|w_0| \rightarrow \infty$ in the closed sector. Consequently its modulus on that circle is at least $r/(2|w_0|)$ for sufficiently large w_0 , and it has precisely one zero inside. For $|z| \leq c/|w_0|$, with fixed sufficiently small $c > 0$, the rational perturbation in (9.9) is bounded by $2|\lambda|e^{|\lambda|r}|z| < r/(2|w_0|)$; its denominator has no zero there. Rouché's theorem gives exactly one root, counted with multiplicity, inside $|\delta| < r$. The root is simple and varies analytically with z . Thus $|\delta(z)| < r$ on the stated disk.

Cauchy's estimate, with radii tending to $c/|w_0|$, yields $|A_n(L)| \leq r(|w_0|/c)^n$. The geometric tail proves (9.7). Substituting $z = q_0$ in (9.9) recovers the exact equation $F_-(w_0 + \delta) = L$, so the sum is the required inverse difference. Finally, $|q_0| = \exp(2\pi\Im w_0)$ decreases exponentially in the sector, whereas $|w_0|$ grows only algebraically. This proves the last assertion. $\square$

Editorial note (ProveIt, 2026-09-29). The later package `Nonlinear_Stokes_Transport_Logarithmic_Inversion/article.tex` of this series (directory `Analysis/Transseries/docs/series-and-transseries/`) proves, independently and without citing this article, a multi-action form of the same Lagrange–Bürmann inverse-sector expansion: its `thm:transport` gives every mixed coefficient of an analytic map perturbed by finitely many exponential terms, with a disk certificate and the tail bound $\rho\chi^{M+1}/((M+1)(1-\chi))$, and `cor:countable` extends it to weighted summable countable families. For one action it has the form of [theorem 9.1](#). Both articles obtain inverse resurgence by verifying Sauzin's closure hypotheses.

9.3 The first three nonlinear sectors

Writing primes for derivatives with respect to L , (9.5) gives

$$A_1 = -\lambda p, \tag{9.10}$$

$$A_2 = \lambda p + \frac{\lambda^2}{2}(p' - 2\lambda p^2), \tag{9.11}$$

$$A_3 = -\lambda p - \lambda^2(p' - 3\lambda p^2) - \frac{\lambda^3}{6}(p'' - 9\lambda p p' + 9\lambda^2 p^3). \tag{9.12}$$

For evaluating them directly from the forward branch, at $w_0 = g(L)$ one has

$$p = \frac{1}{F'_+(w_0)}, \quad p' = -\frac{F''_+(w_0)}{F'_+(w_0)^3}, \quad p'' = \frac{3F''_+(w_0)^2 - F'_+(w_0)F'''_+(w_0)}{F'_+(w_0)^5}. \tag{9.13}$$

The p' and p'' terms are essential. Treating p as a constant would erase part of the second sector and several terms of the third. Likewise, the factor $q(w_0 + \delta)$ must be shifted during inversion; replacing it by $q(w_0)$ would lose the terms in p^2 and p^3 .

The expansion is convergent in its outer exponential marker, with exact analytic coefficient functions $A_n(L)$. Replacing those functions by truncated asymptotic series introduces an additional, separate error. This is the same distinction between exact sector amplitudes and their formal expansions emphasized in the repository.

9.4 Why the positive-real error is not a Stokes jump

For positive real X , the quantity $e^{-2\pi i X}$ has modulus one, so (9.4) is not a small-sector expansion on that ray. There is no positive-real singularity in the Borel kernel (8.1), and the positive-direction Borel sum is unambiguous. The real error $e^{-2\pi X}$ in [theorem 6.1](#) is governed by the distance 2π to the nearest complex Borel singularities and by optimal truncation; it is not a freely adjustable real exponential addition to W .

This distinction also explains why a formal inverse coefficient list alone could not have determined (9.2): its normalization comes from the exact forward function and its lateral continuations. Once that normalization is specified, all nonlinear inverse sectors follow from the finite formula (9.5).

10 Reproducible checks and their logical status

The accompanying scripts were run for this article. They use exact rational arithmetic for the finite coefficient calculations, SymPy for symbolic residual identities, and mpmath at 190 decimal digits for the

analytic comparisons. The separate certificate program uses 200-digit arithmetic only to propose candidate roots and then checks all exported interval assertions with exact fractions. The distinction matters: the floating-point tables test the formulas, whereas the exact interval checks instantiate a proved certificate.

10.1 Late coefficients

Let

$$T_n = \frac{h_n(2\pi)^{2n}}{2(-1)^n\Gamma(2n)}.$$

The h_n in the following table were computed as exact rational numbers, not fitted from the asymptotic expression. The columns compare T_n with the finite rational-in- n expressions $\mathcal{R}_K$ in (4.2), rather than with their further expansions in $1/n$.

n	T_n	$ T_n - \mathcal{R}_1 $	$ T_n - \mathcal{R}_3 $	$ T_n - \mathcal{R}_5 $
20	0.957519636599	$8.07375 \cdot 10^{-4}$	$2.97309 \cdot 10^{-8}$	$1.36284 \cdot 10^{-9}$
40	0.979119023315	$2.07922 \cdot 10^{-4}$	$6.00223 \cdot 10^{-9}$	$3.38346 \cdot 10^{-14}$
80	0.989641503387	$5.24785 \cdot 10^{-5}$	$4.17995 \cdot 10^{-10}$	$1.21710 \cdot 10^{-15}$
160	0.994840420773	$1.31694 \cdot 10^{-5}$	$2.75099 \cdot 10^{-11}$	$2.19491 \cdot 10^{-17}$

Table 1: Actual large-order coefficient checks. Theorem 4.1, not the finite table, proves the all-orders conclusion.

10.2 The sharp inverse error

For $M = \lfloor \pi X \rfloor$, the normalized error is $Q_X = (-1)^M(u_M - W)/(\sqrt{X}e^{-2\pi X})$. The values below compare it with the first-correction prediction $1 + B_1(\sigma)/X$.

X	M	$ u_M - W $	Q_X	$1 + B_1(\sigma)/X$
5	15	$5.33350 \cdot 10^{-14}$	1.050244866478	1.048469941131
10	31	$1.66230 \cdot 10^{-27}$	1.019148786413	1.018435404563
20	62	$1.20674 \cdot 10^{-54}$	1.014267790151	1.014167761314
40	125	$4.50221 \cdot 10^{-109}$	1.005782589474	1.005751752002

Table 2: Actual checks of the leading constant and first correction in theorem 6.1.

For these four values, the analytic bound $(X + 1/2)E_M(a)$ evaluates respectively to approximately $1.24171 \cdot 10^{-13}$, $3.62340 \cdot 10^{-27}$, $2.50333 \cdot 10^{-54}$, and $9.18334 \cdot 10^{-109}$. These displayed decimal evaluations are floating-point diagnostics of an analytic bound. The exported intervals in `certificates.json`, by contrast, are checked by exact rational inequalities.

10.3 Complex-sector checks

The Stokes formula was tested by taking $w_0 = 2 - iR$, setting $F_+(w) = \psi(w + 1/2) - \lambda q/(1 + q)$, choosing $L = F_+(w_0)$, and solving the independent exact equation $\psi(w + 1/2) = L$ near w_0 . The first three coefficients were evaluated from (9.13), not obtained by fitting that root.

The symbolic checks are logically separate. Substitution of the first six exact h_n into the formal inverse equation cancels all residual coefficients through that order. Substitution of equations (9.10) to (9.12) into a generic cubic Taylor model of (9.9) cancels its first three powers of z identically, with the forward derivatives treated as independent symbols. The exact Fermi integral was also compared numerically with the centered digamma function at $w = 3/4, 2, 5$. All assertions in the supplied checks passed. None of these finite checks replaces the proofs for arbitrary order.




Figure 1: At $X = 10$, increasing the number of retained forward coefficients first reduces and then increases the inverse error. The actual error is measured by high-precision root finding; the dashed curve evaluates the analytic bound from [theorem 5.2](#). The vertical line marks the scale $M = \pi X$, not an asserted exact integer minimizer.

R	No sectors	Through $A_1 q_0$	Through $A_2 q_0^2$	Through $A_3 q_0^3$
2	$6.19997 \cdot 10^{-5}$	$2.38521 \cdot 10^{-8}$	$1.37686 \cdot 10^{-11}$	$9.42072 \cdot 10^{-15}$
3	$1.47188 \cdot 10^{-7}$	$1.34191 \cdot 10^{-13}$	$1.83514 \cdot 10^{-19}$	$2.97445 \cdot 10^{-25}$

Table 3: Absolute errors of successive inverse Stokes approximations.

11 Further research questions and concrete next steps

The results isolate several sharply stated next problems. Their status here is “not resolved in this article”; the list is not an assertion that every formulation has already been recognized as an open problem in the wider literature.

Research problem 1 (Direct inverse partial sums and an enveloping law). For $S_M(X) = X + \sum_{n=1}^M h_n X^{1-2n}$, determine whether a useful domain admits a signed first-omitted-term bound for $W(X) - S_M(X)$. In particular, determine the sharp remainder when $M - \pi X$ stays bounded.

The forward expansion is enveloping by positivity of an exact Stieltjes-type integral. That property does not automatically survive nonlinear inversion. [Theorem 6.1](#) therefore cannot be reused as a proof for S_M . A promising route is to derive a contour representation for the inverse displacement itself and analyze the discontinuities of its continuation. Enveloping results for reverted Airy and cylinder-zero expansions, proved in a different setting by Nemes [7], show that an analytic inverse representation can be decisive. They do not settle the present function. A counterexample to a global sign claim would also be useful: it would specify the right local or asymptotic formulation before attempting a general theorem.

Editorial note (ProveIt, 2026-09-29). Three later packages of this series (directory `Analysis/Transseries/docs/series-and-transseries/`) answer the bounded-offset part of this problem, independently of one another and by three methods: `Direct_Optimal_Truncation_Inverse_Ha`

rmmonic/direct_inverse_harmonic.tex (thm:sharp, from an exterior dispersion representation with a positive tail density), Inverse_Digamma_Spectral_Representation/inverse_digamma_spectral.tex (thm:optimal, from a spectral representation, the route suggested above) and Optimal_Truncation_After_Nonlinear_Reversion/article.tex (thm:direct, from a general cutoff-transfer theorem). With $\sigma = M + 1 - \pi X$ in the normalization of this article, all three prove $S_M - W = (-1)^M \sqrt{X} e^{-2\pi X} (1 + B_1(\sigma)/X + B_2(\sigma)/X^2 + O(X^{-3}))$ to every fixed order, with $B_1(\sigma) = (2\sigma^2 - 3\sigma + 7/12)/(2\pi) - \pi/12$ and the same B_2 ; the three expressions for B_2 were checked to agree term by term. Hence $S_M - u_M = (-1)^{M+1} (\pi/6) X^{-1/2} e^{-2\pi X} (1 + O(X^{-1}))$. The first two also prove eventual strict enveloping, for all sufficiently large X and M with existential thresholds (thm:enveloping in each); the third proves it only on windows $|M - \pi X| \leq K$ (cor:enveloping). A signed bound for every M at a given X , and explicit thresholds, are established by none of them.

Research problem 2 (Explicit local singularities of the inverse Borel transform). Determine the full local expansions of $\mathcal{B}(\tilde{W} - X)$ at $\pm 2\pi i$, including pole and logarithmic terms, and propagate them to subsequent lattice points on specified continuation sheets.

The radius 2π and all inverse jump coefficients are already known here, but they do not constitute a complete description of local Borel singular germs on every sheet. The factorial asymptotics should constrain the nearest singular expansions, while (9.5) supplies exact nonlinear consistency conditions. A constructive proof should match these two descriptions and state precisely which lattice points are actual singularities, rather than merely permitted singular locations.

Editorial note (ProveIt, 2026-09-29). The later package Inverse_Digamma_Spectral_Representation/inverse_digamma_spectral.tex (directory Analysis/Transseries/docs/series-and-transseries/) answers the first part of this problem at the nearest points only: its thm:borel continues the Borel transform to the strip $|\Im \xi| < 2\pi$ by an inverse cosine representation, and its thm:boundary gives the complete singular expansion at $\pm 2\pi i$, with every logarithmic coefficient, for inward approach from that strip. Propagation to further lattice points and continuation sheets is not claimed there and remains open.

Research problem 3 (A certified hyperasymptotic inverse). Use the exact gamma-expectation remainder to improve u_M beyond its entire first exponential block, with a rigorously quantified next exponential scale.

Subtracting finitely many B_j/X^j from (6.4) improves the algebraic amplitude of $e^{-2\pi X}$; it does not by itself produce an error of order $e^{-4\pi X}$. To reach a second exponential layer, one must retain an exact or exponentially accurate first remainder block and then control both its own error and the quadratic inverse transport in (6.11). The exact expectation (6.6), together with a certified evaluation method for its rational gamma average, provides a concrete starting point.

Research problem 4 (Uniform passage from generalized harmonic inverses to the harmonic inverse). Develop a uniform inversion and exponentially improved remainder theory for $H_s(x) = \zeta(s) - \zeta(s, x+1)$ as $s \downarrow 1$ and $x \rightarrow \infty$ jointly.

The repository gives fixed- $s > 1$ endpoint inverse coefficients in terms of $T = ((s-1)(\zeta(s)-y))^{-1/(s-1)}$. The singular normalization as $s \rightarrow 1$ requires a renormalized target and control of the joint parameter $(s-1) \log x$. A useful theorem should recover the harmonic result in one limit and the fixed- s endpoint result in another, with common remainder constants on an explicitly stated transition region. Merely substituting $s = 1$ into a fixed- s big-O estimate is insufficient.

Research problem 5 (Other inverse Bernoulli families and competing actions). Apply the endpoint-extraction theorem to inverse gamma quotients and related exactly centered phases, then determine how several equally near complex actions interact under inversion.

The proof of theorem 3.2 accommodates complex coefficients and does not require positivity. It can therefore extract late-coefficient contributions from several oscillatory factorial sources. What changes is

the analytic normalization and the possibility of cancellation. A useful general result would identify when such cancellation changes the effective Gevrey scale or removes a nearest Borel singularity, and would connect those algebraic criteria to an exact remainder or contour representation for the original function.

Research problem 6 (Arithmetic action accumulation rather than a discrete Borel lattice). Construct a summable inverse calculus for divisor-dependent sectors whose positive actions have a finite accumulation point, as in the repository’s necklace and irreducible-polynomial examples.

This is a separate frontier explicitly identified in the companion [1]. The lattice $2\pi i\mathbb{Z}$ used here is closed and discrete, so the standard nonlinear resurgence closure theorem applies. A divisor-action set accumulating at a finite location violates that hypothesis. Finite divisor cutoffs are well defined, but an infinite calculus must specify a topology or summability condition, retain the divisibility indicators, and justify products and inverse operations in that setting. A formal Hahn-support condition alone is not an analytic convergence argument.

Editorial note (ProveIt, 2026-09-29). The later package `Arithmetic_Transseries_Beyond_Accumulation_Cut/arithmetric_transseries.tex` (directory `Analysis/Transseries/docs/series-and-transseries/`) answers this problem, without citing it, in the form the companion’s third research direction states: it retains the divisibility indicators, constructs a weighted Banach algebra over a faithful family of analytic sheets (`thm:banach`), characterizes which elementary blocks are admissible (`thm:admission`), and proves convergent inverses on each fixed arithmetic sheet (`thm:inverse`, `thm:range`). It proves no canonical global interpolation and no resurgence, Borel summation or Stokes automorphism in that setting. The package `Action_Accumulation_Nonlinear_Inversion/` gives a label-free analytic inversion calculus for accumulating actions that drops the divisibility indicators.

Research problem 7 (Mechanization and bit complexity). Formalize the finite factorial estimates and the exact-rational certificate first; then obtain an explicit bit-complexity bound for a certified inverse evaluation to a prescribed accuracy.

The most accessible initial targets are [theorem 3.1](#), the coefficient extraction underlying [theorem 3.2](#), the rational logarithm bound, and the mean-value enclosure in [theorem 5.2](#). They connect naturally to the repository’s existing asymptotic-scale and backward-error interfaces [2]. The full sectorial resurgence machinery is a much larger dependency and is not needed to certify positive-real values. The current scripts prioritize transparent arithmetic over optimal complexity; a faster implementation should use truncated-series algorithms and binary splitting for exact sums while preserving the same small certificate boundary.

12 Synthesis

For this specific inverse, the gap between formal transseries and quantitative analysis can be crossed completely along two complementary routes. On the positive axis, an exact positive integral yields signed remainders, explicit inverse brackets, and a sharp exponential error with all algebraic corrections. In the complex domain, an exact meromorphic Borel kernel fixes the forward jump, and a finite nonlinear coefficient formula transports it to a convergent inverse-sector expansion. Large-order coefficient asymptotics provide an independent connection between the two descriptions.

The logical dependencies remain visible. The all-coefficient inverse formula comes from the repository. The factorial endpoint estimate and its specialization prove the additional late-coefficient result. The exact centered integral is derived from classical Binet and duplication formulas. General resurgent closure is supplied by existing theory. The specific sharp inverse error and all-sector transport are then proved using those inputs. Exact-rational certificates turn the real error bounds into independently reproducible numerical enclosures. The next questions concern direct inverse truncation, finer Borel singular structure, higher exponential accuracy, uniform parameters, and action accumulation—not missing premises of the results proved here.

A Finite coefficient algorithms

A.1 Exact coefficients without expanding an exponential symbolically

For fixed N and degree K , set $E_0 = 1$ and use

$$E_k = \frac{N}{k} \sum_{j=1}^k j c_j E_{k-j}, \quad 1 \leq k \leq K. \quad (\text{A.1})$$

This is coefficient comparison in $(\exp(N\mathcal{C}))' = N\mathcal{C}' \exp(N\mathcal{C})$. With $N = 2n - 1$ and $K = n$, it gives $h_n = -E_n/(2n - 1)$ exactly. With K fixed, it computes (4.2) using only $c_1, \dots, c_K$, regardless of the size of n . The current exact implementation takes quadratically many rational arithmetic operations for one such truncated exponential. The cost in bits also depends on the growth of numerator and denominator sizes.

For reference, the next two coefficients of the normalized late expansion after (4.4), with $p = \pi^2$, are

$$[n^{-4}]T_n = \frac{p^4}{497664} + \frac{p^3}{2880} - \frac{37p^2}{1440} - \frac{p}{12}, \quad (\text{A.2})$$

$$[n^{-5}]T_n = -\frac{p^5}{29859840} - \frac{23p^4}{1244160} - \frac{403p^3}{362880} - \frac{29p^2}{288} - \frac{p}{12}. \quad (\text{A.3})$$

Here the bracket notation denotes the coefficient in the asymptotic expansion in powers of $1/n$, not a power-series coefficient of a convergent function T_n .

A.2 Computing a prescribed sharp-error coefficient

To calculate B_j in (6.4), take Taylor coefficients of $f(v) = (1 + v^2)^{-1}$ at one through a sufficiently high even order. Replace powers of $v - 1$ by the finite moment polynomial (6.9), put $s = b + 2\tau$, and expand to order b^{-j} . Multiply by the exponential of the finite shifted-Stirling series (6.10). Finally substitute $w = W(X)$ to the necessary finite order, use $\tau = \sigma + \pi(X - w)$, and multiply by $1/F'(w)$. At every fixed order this uses finitely many rational coefficients and powers of π .

The gamma-moment proof provides the error control for this algorithm. It is not merely a formal prescription for guessing B_j from a numerical fit. Higher derivative estimates and central moments are needed only to the finite order requested.

A.3 Computing inverse jump coefficients

For each desired sector n , start with the function $p(L)$ and repeatedly apply $D - n\lambda p$ to obtain the n expressions needed in (9.5). Combine them with the displayed finite binomial weights. This keeps the dependence on the exact dominant inverse visible and avoids expanding its internal asymptotic series prematurely. For evaluation, the derivatives of p can instead be generated from derivatives of F_+ by implicit differentiation, as illustrated in (9.13).

B Reproduction, provenance, and audit boundary

B.1 Contents of the archive

The archive contains this editable TeX source, its compiled PDF, the figure in PDF and PNG form, and three Python programs with their actual outputs. Run

```
python verification/verify.py
python verification/certify.py
python verification/make_figure.py
pdflatex -interaction=nonstopmode -halt-on-error \
  inverse_harmonic_transseries.tex
pdflatex -interaction=nonstopmode -halt-on-error \
  inverse_harmonic_transseries.tex
```

The Python programs need Python 3.10 or later, SymPy, mpmath, and, for the figure, matplotlib. The PDF needs a normal LaTeX installation with the packages listed in the preamble, including `newtxtext` and `newtxmath`. No repository-relative style file is required. The stored figure permits rebuilding the article without rerunning the numerical computation.

B.2 Exact repository checkpoint

The inspection used commit

04e06e032dff1966513bfba316966d52db3646b4.

The principal companion source is

```
Analysis/FabiusFunction/docs/semi-formalized-research-frontiers/drafts/series-and-transseries/Combinatorial_Transseries_Inverses/Combinatorial_Transseries_Inverses.tex.
```

Editorial note (ProveIt, 2026-09-29). The path above is correct at the commit shown. The series has since moved; the companion is now `Analysis/Transseries/docs/series-and-transseries/Combinatorial_Transseries_Inverses/Combinatorial_Transseries_Inverses.tex`.

The relevant checkpoint source ranges were 4220–4500 for harmonic and generalized-harmonic inversion, 4500–4870 for certification and numerical distinctions, and 4870–5160 for support closure, status boundaries, and research questions. The group README and canonical transseries README were also inspected. These source ranges, paths, and commit identify a reproducible starting point; they are not a claim that every file in the repository was read.

B.3 What the verification does and does not establish

Exact symbolic checks. The displayed finite inverse and Stokes coefficients satisfy their defining residual identities through the checked orders.

Numerical checks. The large-order, optimal-scale, integral, and complex-sector tests agree with the theorems at the recorded inputs and precision. These are finite floating-point tests, not universal proofs.

Exact-rational certificates. The exported real inverse intervals have independently checked rational endpoint signs, using the proved analytic remainder inequality.

Human proofs. The all-orders and uniform asymptotic statements are established in the arguments of the article, independently of the finite tests.

Not asserted. No new Lean file is supplied; no global priority claim or complete treatment of all repository transseries families is made; and no sharp direct-inverse-partial-sum remainder is inferred from the forward-truncation theorem.

References

- [1] Vladimir Reshetnikov, *ProveIt*, companion manuscript *Combinatorial Transseries and Their Inverses: Gamma quotients, finite exponential sums, moment sectors, moving saddles, arithmetic sheets, and q -products*, repository snapshot 04e06e032dff1966513bfba316966d52db3646b4, inspected 29 September 2026. Exact path and inspected ranges in section B. https://github.com/VladimirReshetnikov/ProveIt/tree/04e06e032dff1966513bfba316966d52db3646b4/Analysis/FabiusFunction/docs/semi-formalized-research-frontiers/drafts/series-and-transseries/Combinatorial_Transseries_Inverses.
Editorial note (ProveIt, 2026-09-29): now at Analysis/Transseries/docs/series-and-transseries/Combinatorial_Transseries_Inverses/.
- [2] Vladimir Reshetnikov, *ProveIt*, “Series and transseries” group README and “Transseries: the polynomial–logarithmic calculus and its inversions” README, same snapshot. <https://github.com/VladimirReshetnikov/ProveIt/tree/04e06e032dff1966513bfba316966d52db3646b4/Analysis/FabiusFunction/docs/semi-formalized-research-frontiers/drafts/series-and-transseries>.
Editorial note (ProveIt, 2026-09-29): now at Analysis/Transseries/docs/series-and-transseries/README.md and Analysis/Transseries/docs/series-and-transseries/Transseries_And_Inversion/README.md.
- [3] NIST Digital Library of Mathematical Functions, Chapter 5, especially §5.5 (functional relations), §5.9 (integral representations), and §5.11 (asymptotic expansions), accessed 29 September 2026. <https://dlmf.nist.gov/5.5>; <https://dlmf.nist.gov/5.9>; <https://dlmf.nist.gov/5.11>.
- [4] B. R. Fabijonas and F. W. J. Olver, “On the reversion of an asymptotic expansion and the zeros of the Airy functions,” *SIAM Review* **41**(4) (1999), 762–773. <https://doi.org/10.1137/S0036144598349538>.
- [5] M. Borinsky, “Generating asymptotics for factorially divergent sequences,” *Electronic Journal of Combinatorics* **25**(4) (2018), P4.1. <https://doi.org/10.37236/5999>. Preprint: <https://arxiv.org/abs/1603.01236>.
- [6] D. Sauzin, “Nonlinear analysis with resurgent functions,” *Annales scientifiques de l’École normale supérieure*, series 4, **48**(3) (2015), 667–702. <https://doi.org/10.24033/asens.2255>. Preprint: <https://arxiv.org/abs/1212.4477>.
- [7] G. Nemes, “Proofs of two conjectures on the real zeros of the cylinder and Airy functions,” *SIAM Journal on Mathematical Analysis* **53**(4) (2021), 4328–4349. <https://doi.org/10.1137/21M1396794>. Preprint: <https://arxiv.org/abs/2010.13069>.