

A MERGED RESEARCH CONTINUATION OF THE SURREAL REPOSITORY

Nonabelian Support Obstructions

A Matrix Cousin Criterion
and the Surcomplex
Riemann–Hilbert Problem

September 21, 2026; Part III added September 22, 2026;
Part V added September 23, 2026

Abstract

We study one coefficient category from two sides. Its objects are matrices congruent to the identity modulo positive Hahn exponents, $I_r + \text{Mat}_r(\mathcal{J}_\Gamma)$, over a set-sized and possibly non-Archimedean ordered value group Γ , with ordinary holomorphic coefficient functions on a single fixed ordinary complex domain; when $\Gamma \subseteq \mathbf{No}$, the substitution $t^\gamma = \omega^{-\gamma}$ gives the surcomplex reading.

Part I solves the multiplicative gluing problem. On a puncture-and-disk cover of a connected open Riemann surface we prove an exact criterion: factor each transition matrix *independently* into a normalized polar factor and a factor holomorphic on the full disk, and test whether the union of the supports of the *polar* factors is well ordered. The criterion is invariant under arbitrary local holomorphic right gauges and under ordered value-group extension, and it produces genuinely nonabelian counterexamples: a rank-three family whose determinant is one, whose only raw singular exponent is 1, and whose abelianized problem splits, yet whose integral vector bundle is nontrivial; a repair of that family by a term which *worsens* its raw singular support; and, for every $r \geq 3$, an example trivial modulo the last central subgroup of the upper-unitriangular group but nonsplit in full positive rank r . We also prove support-controlled Stein splitting and deformation rigidity, with a cyclic-versus-noncyclic dichotomy.

Part II solves the inverse monodromy problem in the same category. A near-identity representation of $\pi_1(\mathbb{C} \setminus D)$ is the monodromy of a positive globally common-domain Hahn connection exactly when the union over punctures of the supports of $\rho(\ell_d) - I_r$ is well ordered; the realizer may then be taken coefficientwise logarithmic, and a fixed linear Mittag–Leffler period section selects a unique normalized realizer. We prove framed gauge classification, entire extension of logarithmic comparison gauges, a positive Frobenius form, a determinant-one refinement, calibrating examples on both sides of the criterion, an explicit mixed commutator correction in the value group $\mathbb{Q} + \mathbb{Q}\omega$, and the impossibility of a continuous linear period right inverse for infinitely many punctures.

Part III, from a third source, asks what survives when integrality is given up. Placing a strictly decreasing sequence of positive exponents in front of ordinary essential singularities, it constructs determinant-one unipotent bundles on the plane, trivial on every bounded domain, that stay indecomposable and not stably trivial after all Hahn monomials are inverted and even over the coefficientwise meromorphic sheaf $\text{Mer}((t^\Gamma))$. It computes all their sections and morphisms, classifies them exactly, and proves a localization-stable cyclic-versus-noncyclic dichotomy, continuum many classes, a tensor rule, and the failure of semidecidability of isomorphism. It does not settle whether Part I’s pole-only examples survive monomial localization; over $\text{Mer}((t^\Gamma))$ they become trivial.

Part V, from a fourth source, settles it negatively: inverting one Hahn monomial trivializes Part I’s hidden and deep examples and every descending-scale pole cocycle with one fixed nilpotent direction. It also gives free-middle nonsplit extensions, an omnific-valued sine family, a proper locally unit ideal of projective dimension two, and that family’s zeros on finite halos.

The two headline criteria are different criteria on different objects. Part I tests the supports of normalized polar factors, not the raw transition supports; Part II tests the raw meridian monodromy supports. [Section 12](#) states which object each is a criterion on, and [section 8](#) contains the example that makes the distinction unavoidable.

Status and scope. This report merges four independently written manuscripts; see [appendix C](#). The exact global criteria and the explicit obstruction families are proposed new results in the specified Hahn-coherent categories. The local factorization mechanisms, the Neumann support calculus and every ordinary analytic input are established mathematics and are credited as such. The literature checks were targeted and certify no priority; no named published conjecture is claimed solved. Nothing here is refereed or machine checked. Nontriviality in Part I is proved over the *integral* coefficient sheaf only, and Part II assumes positive support and near-identity monodromy throughout: neither theorem is restated over a coefficient field in which negative-valuation changes of frame are admitted, and neither is the unrestricted classical Cousin or Riemann–Hilbert theorem. Part III proves nontriviality after localization for a different, essential-singularity family only, and Part V proves that Part I’s pole-only examples become trivial after localization. All infinite algebra is by well-ordered supports and finite coefficient multiplicity, never by convergence in the fine surreal topology.

Contents

1	Two questions, one category	1
1.1	The shared setting	1
1.2	The two criteria, and why they are two	2
1.3	What each part contributes	2
1.4	Relation to reports already in the collection	3
1.5	What is and is not claimed as new	4
2	The coefficient category	5
2.1	One set-sized workspace	5
2.2	Two standing conventions that differ between the two parts	5
2.3	Comparison of names across the two parts	6
2.4	Surcomplex evaluation on a halo	7
2.5	Ordinary paths, not fine-surreal paths	7
3	Support calculus without any convergence assumption	7
4	The ordinary analytic inputs	9
4.1	A fixed-disk Laurent splitting	9
4.2	Isolated singular parts: the Part I packaging	9
4.3	The period map and a linear section: the Part II packaging	10
5	Local positive factorizations at a puncture	11
5.1	The normalized polar factorization	11
5.2	Where the obstruction is manufactured	13
5.3	The positive Frobenius form	13
	Part I. Multiplicative gluing: the polar-support criterion	14
6	The sharp global matrix Cousin criterion	14
6.1	Why the converse is sharp	15
6.2	The space of solutions	16
7	Invariance, and the scalar case that is recovered	16
8	A rank-three obstruction invisible in the raw poles	17
8.1	The closed factorization in the Heisenberg group	17
8.2	The obstructed datum	18
8.3	Adding a singular term repairs the problem	18
8.4	Why a projected matrix logarithm is not a repair	19

9	Obstructions at arbitrary nilpotent depth	19
9.1	A chain identity	19
9.2	The exponent design	20
10	Integral vector bundles and local invisibility	21
10.1	From positive matrices to actual vector bundles	21
10.2	Every bounded restriction is trivial	22
10.3	The localization boundary	22
11	Positive rigidity on Stein manifolds	23
11.1	An arbitrary-cover theorem with a support hypothesis	23
11.2	Deformations of a nontrivial ordinary bundle	23
11.3	A cyclic versus noncyclic dichotomy	24
12	The two well-ordered-support criteria are not the same criterion	25
Part II. Inverse monodromy: the meridian-support criterion		26
13	Connections, fundamental matrices, and holonomy	26
13.1	The explicit iterated-integral expression	27
14	The exact realization theorem	27
14.1	An intrinsic version of the support condition	29
15	Framed gauge classification and removable gauges	29
16	Finite-puncture inversion and determinant one	31
17	Sharp global obstructions and calibrating examples	32
17.1	Finite solvability does not imply global solvability	32
17.2	The union condition must not be strengthened to point-finiteness	32
17.3	Admissible supports need not be locally finite in an interval	33
18	A noncommuting calculation across two valuation ranks	33
18.1	The targets and the first-order guess	33
18.2	The representation is genuinely nonabelian	34
18.3	Specialization to a genuinely higher-rank Hahn group	35
Part III. Beyond the localization boundary: essential-singularity bundles over $\mathcal{H}_\Gamma$ and $\text{Mer}((t^\Gamma))$		35
19	What survives monomial localization	35
19.1	The question inherited from Part I	35

19.2	An explicit model, and the results at a glance	36
19.3	What is new here, what is inherited, and what overlaps with Part I	37
20	The meromorphic coefficient sheaf	38
20.1	Three coefficient categories on the ordinary plane	38
20.2	Leading terms and units	39
20.3	Bundles and base change	40
20.4	The surcomplex reading	40
20.5	Names in this part	40
21	An essential singularity protects a descending support	41
21.1	The fixed cover and the local data	41
21.2	A cohomological formulation without a cohomology assumption	42
22	Nilpotent bundles and localization-resistant nontriviality	43
22.1	Finite exponentials and the gluing convention	43
22.2	Every bounded restriction is trivial	44
22.3	Exact sequences and the standard flag	44
22.4	All global sections, and nontriviality after localization	44
22.5	Which further localizations are covered	45
23	Every morphism and the intrinsic flag	46
23.1	The internal Hom is another nilpotent bundle	46
23.2	A representation category on the ordinary plane	47
23.3	Recovering the flag intrinsically	48
24	Exact classification of exponent profiles	48
24.1	Necessity: the rank-two calculation	49
24.2	Sufficiency: explicit finite gauges	49
24.3	Examples that separate the categories	50
25	Value groups: a localization-stable dichotomy and continuum many classes	51
25.1	Which groups admit a descending positive profile	51
25.2	The localization-stable rigidity dichotomy	51
25.3	Continuum many localized classes	52
26	Tensor products and duality	53
27	Poles versus essential singularities	54
27.1	Pole data are removable in the meromorphic category	54
27.2	Why this does not contradict ordinary analytic bundle theorems	55
27.3	Why determinant and finite tests miss the obstruction	56

27.4 Other essential singularities	56
Part IV. Effectivity, verification, and boundaries	57
28 How to compute the invariants, and what computation cannot decide	57
28.1 A support-aware procedure for Criterion P	57
28.2 Finite truncations	57
28.3 What the Criterion M recursion actually depends on	57
28.4 Why an infinite linear period section cannot be continuous	58
28.5 Part III: a complete criterion that is not semidecidable	59
29 The accompanying exact and numerical checks	60
30 Exactly what is not claimed	62
31 Questions exposed by the results	67
32 Conclusion	70
Part V. Pole localization and omnific interpolation	72
33 The printed examples after monomial localization: question, source and conventions	72
33.1 What this part answers	72
33.2 Source and status	72
33.3 Conventions, and the three gluing conventions	73
34 Explicit Laurent frames for pole-only unipotent cocycles	74
34.1 An ordinary principal-part construction	74
34.2 The rank-two theorem	75
34.3 Every fixed nilpotent direction	77
35 The printed examples become trivial over $\mathcal{H}_\Gamma$	78
35.1 The hidden rank-three example	78
35.2 The deep examples	79
35.3 What this settles and what it does not	79
36 Line bundles and extensions that localization does not kill	80
36.1 Line bundles	80
36.2 Nonsplit extensions with a free middle bundle	81
37 An omnific-valued cardinal-sine family and a global Bézout obstruction	82
37.1 The family	82

37.2 A proper, locally unit, nonprincipal ideal	83
37.3 Ultrafilter maximal ideals	84
37.4 Taylor prolongation to finite halos	85
38 Zeros of the cardinal-sine family on finite halos	86
38.1 Scope of Part V	87
A A proof and support audit	88
B A dependency map	91
C Provenance, reproduction, and the merge record	92
C.1 Sources of this report	92
C.2 Repository provenance	95
C.3 Building this report	96
C.4 Rerunning the delivered checks	96
References	97

1 Two questions, one category

1.1 The shared setting

Let Γ be a set-sized totally ordered abelian group and let t^Γ be its monomial group. A coefficient of our objects is an ordinary holomorphic function on a *single fixed* ordinary complex domain, and an object is a Hahn series in t with well-ordered support whose coefficients all live on that same domain. The matrices we glue and the matrices we realize as monodromy both lie in

$$I_r + \text{Mat}_r(\text{positive-support part}),$$

so they are congruent to the identity modulo positive exponents. When Γ is an ordered additive subgroup of the surreal numbers $\mathbf{No}$, the Conway normal form substitution $t^\gamma = \omega^{-\gamma}$ interprets everything inside $\mathbf{SC} = \mathbf{No}[i]$. That interpretation is a reading of a Hahn model; it is never an assertion that a fine-topological surcomplex manifold or a fine-continuous surcomplex path is involved.

Two problems are asked in this one category.

Part I: multiplicative gluing (a matrix Cousin problem). Let X be a connected open Riemann surface, let $A \subset X$ be closed and discrete, let (D_a, u_a) be pairwise disjoint coordinate disks, and put $U_0 = X \setminus A$, $W_a = D_a \setminus \{a\}$. Given, *separately for each a* , a positive matrix G_a on W_a , when are there positive matrices B_0 on U_0 and B_a on D_a with

$$G_a = B_0 B_a^{-1} \quad \text{on } W_a? \tag{1.1}$$

Each local equation is harmless. The difficulty is that the entries of the *one global matrix* B_0 must have a legitimate, that is well ordered, Hahn support.

Part II: inverse monodromy (a Riemann–Hilbert problem). Let $D \subset \mathbb{C}$ be closed and discrete, $U = \mathbb{C} \setminus D$, and let $\rho : \pi_1(U, p) \rightarrow I_r + \text{Mat}_r(\mathfrak{m}_\Gamma)$ be a near-identity representation. When is ρ the monodromy of a connection $dY = \Omega Y$ whose coefficients are ordinary holomorphic matrix one-forms on all of U and whose Hahn support is a single well-ordered positive set?

These are different questions sharing a coefficient category and an ordinary Mittag–Leffler input applied one exponent at a time. Part I first normalizes each transition matrix; Part II corrects the lower-weight monodromy prediction by ordinary periods. Mixed products matter in both constructions, but the support obstruction in Part II can already occur for an abelian representation. In both problems a certain *union of Hahn supports must be well ordered*. The objects in those unions must be distinguished.

1.2 The two criteria, and why they are two

Criterion P (Part I, [theorem 6.2](#)). Gluing (1.1) is solvable if and only if

$$\mathcal{E}(G) = \bigcup_{a \in A} \text{supp}(\text{Pol}(G_a) - I_r) \quad \text{is well ordered,}$$

where $\text{Pol}(G_a)$ is the normalized polar factor produced by an *independent local factorization* of each transition matrix $G_a = \text{Pol}(G_a) \text{Reg}(G_a)$. This is a criterion on the polar factors, not on the transition matrices. Well-ordering of the full raw union $\bigcup_a \text{supp}(G_a - I_r)$ is *sufficient but not necessary*. Well-ordering of the raw *singular-coefficient* support is neither necessary nor sufficient ([section 8](#)).

Criterion M (Part II, [theorem 14.2](#)). A near-identity representation ρ is realizable if and only if

$$E_\rho = \bigcup_{d \in D} \text{supp}(\rho(\ell_d) - I_r) \quad \text{is well ordered,}$$

the union being over the matrices themselves, for a compatible system of based meridians ℓ_d freely generating $\pi_1(U, p)$. No factorization is applied first, and none may be: this is a criterion on the monodromy matrices, and it is basis- and basepoint-independent ([proposition 14.4](#)).

A reader who carries one “well-ordered support criterion” away from this report will test the wrong object in one of the two problems.

The distinction is not pedantic. [Section 8](#) contains a rank-three family over $\Gamma = \mathbb{Q}$ whose raw singular support is the single exponent $\{1\}$ and which nevertheless cannot be glued, because the normalized polar factors are forced to carry the descending set $\{1 + 1/n\}$; and it contains the opposite example, in which *adding* a term that introduces exactly that descending set among the raw poles *removes* the obstruction. So in Part I the raw singular support is neither sufficient nor necessary, and the factorization must be performed first. In Part II, by contrast, well-ordering of the raw monodromy support is the invariant condition, and applying a factorization to $\rho(\ell_d)$ before testing would be a mistake of the opposite kind.

1.3 What each part contributes

Part I owns the machinery the two parts share: the integral common-domain Hahn sheaf on an ordinary base, the noncommutative support calculus, the normalized local polar factorization with right-gauge invariance, invariance under ordered value-group extension, the recovery of the known rank-one scalar test, and the Stein branch. Its own new content is the exact criterion $\mathcal{E}(G)$, the rank-three cross-term obstruction with its repair and its logarithm diagnostic, the arbitrary-depth central hierarchy, the resulting determinant-one integral vector bundles that are trivial on every bounded plane domain, the explicit localization boundary, and the cyclic-versus-noncyclic rigidity dichotomy.

Part II reads as the connection-and-monodromy application of that machinery, with its own new content: the three-way equivalence of [theorem 14.2](#) including *necessity*, which comes from a uniform support bound on holonomy valid for every group element; the fixed complex-linear period section that selects a unique normalized realizer; the framed gauge classification and the entire extension of logarithmic comparison gauges; the positive Frobenius form in which no spectral resonance operator is ever inverted; the SL_r refinement; three calibrating examples that pin the criterion from both

sides; the explicit first nonabelian correction with its $\mathbb{Q} + \mathbb{Q}\omega$ specialization; and the proof that an unrestricted infinite-puncture linear period right inverse cannot also be continuous.

Part III, built from a third manuscript, leaves the integral category on purpose. Part I’s bundles are proved nontrivial over $\mathcal{I}_\Gamma$ only, and [section 31](#) asked whether its examples survive monomial localization. Part III does not answer that question for those examples. It answers the broader existence question with a different family: descending positive exponents placed in front of ordinary *essential* singularities give determinant-one unipotent bundles on the plane that remain indecomposable and not stably trivial over $\mathcal{H}_\Gamma$ and over the coefficientwise meromorphic sheaf $\mathcal{M}_\Gamma = \text{Mer}((t^\Gamma))$, while being integral-trivial on every bounded domain. Its own content is the zero-annihilator obstruction behind this, the complete calculation of global sections and morphisms, the exact classification of the family (eventual equality of exponent profiles over $\mathcal{I}_\Gamma$, eventual equality up to one translation over $\mathcal{H}_\Gamma$ and $\mathcal{M}_\Gamma$), the extension of the cyclic dichotomy to $\mathcal{H}_\Gamma$ and $\mathcal{M}_\Gamma$, continuum many classes, a tensor rule, and the failure of semidecidability of isomorphism ([theorem 28.4](#)). It also records that Part I’s pole-data examples, and the global-divisor report’s plane Picard classes, become trivial over $\mathcal{M}_\Gamma$ ([corollary 27.2](#)).

Part V, built from a fourth manuscript, answers the printed question for Part I’s examples, negatively. An explicit determinant-one Laurent frame, placing interpolation coefficients at the admissible negative exponents $-\alpha_n$, trivializes every descending-scale cocycle $\exp(t^{\alpha_n} p_n N)$ with finite-order poles and one fixed nilpotent N after one Hahn monomial is inverted; an integral reduction brings the hidden and deep examples to that form, so all of them are trivial over $\mathcal{H}_\Gamma$ ([theorem 35.1](#) and [corollary 35.2](#)). In rank one the opposite holds, as the global-divisor report already shows. Part V also constructs nonsplit self-extensions with free middle bundle, an omnific-valued cardinal-sine family, a proper locally unit ideal whose quotient has projective dimension two, ultrafilter maximal ideals, and the classification of the zeros of the sine family on finite halos. To keep the numbering of Parts I–IV unchanged, Part V is placed after Part IV; its checks, limitations and questions are recorded in [sections 29](#) to [31](#).

1.4 Relation to reports already in the collection

This report is a continuation of the repository `VladimirReshetnikov/Surreal`, reviewed at commit

39f2be6667ade51bca2b45daa47e289d69c09764,

inspected September 21, 2026. Two companion reports are leaned on and are *cited rather than reproved*.

Global Support Obstructions: Moving Divisors, Mittag–Leffler, and a Picard Dichotomy, at `docs/surcomplex/global-divisors/article.tex` [[2](#)], proves the scalar support criteria and the Picard-group dichotomy for this workspace, and its subsection “Further precise directions” asks for exactly the present work: “the finite-rank vector-bundle theory: matrix-valued positive transition functions no longer reduce to an abelian logarithm identity, so a noncommutative support recursion must replace the rank-one argument”, and the same package over general ordinary open Riemann surfaces. Part I answers both. Its rank-one specialization *recovers* that report’s exact Cousin criterion, its Theorem 13.1, stated there in terms of the singular support $\Sigma(b)$; we cite that theorem as the scalar case rather than reprove it ([proposition 7.4](#)). Likewise the Picard-group vanishing dichotomy of that report — vanishing exactly when $\Gamma = \{0\}$ or $\Gamma = \mathbb{Z}\delta$ — is the rank-one shadow of our Stein dichotomy ([theorem 11.4](#)), driven by the same universal well-ordered set $\{\delta, 2\delta, \dots\}$.

Differential Algebra and Differential Equations over the Surreal and Surcomplex Numbers, at `docs/surcomplex/differential-equations/article.tex` [[3](#)], contains the *forward* map that

Part II inverts: its theorem “Monodromy valued in $\mathrm{GL}_d(K_\Gamma)$ ” constructs M_ℓ in $\mathrm{GL}_d(K_\Gamma)$ from a coherent connection, records that the coefficient of M_ℓ at exponent 0 is the ordinary monodromy matrix and that all other exponents lie in $S^* \setminus \{0\}$, and then states the categorical boundary we adopt verbatim rather than re-argue: “No fine-continuous surcomplex path is asserted: the loop is an ordinary complex path, continuation is coefficientwise, and the result is a matrix of Hahn constants.” [Theorem 14.2](#) is the converse of that map for near-identity representations, together with the exact condition on which ones arise.

Part III was written against a later commit, 0097304c7ae9d5de46d2ea342e2494f8fc126303, at which this report and the global-divisor report already had their present text. It inherits the global-divisor report’s scalar Picard dichotomy as the rank-one antecedent of its localization-stable dichotomy ([theorem 25.1](#)), and the merge records one consequence for that report: the explicit plane Picard classes there, nontrivial over $\mathcal{I}_\Gamma$ and $\mathcal{H}_\Gamma$, become trivial over the coefficientwise meromorphic sheaf $\mathcal{M}_\Gamma$ ([corollary 27.2](#)). This is not a contradiction, since that report’s dichotomy concerns $\mathcal{I}_\Gamma$ and $\mathcal{H}_\Gamma$ only. The sheaf $\mathcal{M}_\Gamma$ is not one of the four coefficient constructions tabulated in the collection’s index; the global-divisor report uses the same recipe over a compact base, and the analysis report [4] uses its rings for a coefficientwise residue ([section 20](#)).

Part V was written against commit 0865f043aec113c14c69ef45006bbc7546a4e75a, at which this report had exactly the text of Parts I–IV. Its source inspected this report only. The merge records its overlaps with three further reports, all cited rather than reprinted: the global-divisor report’s Picard reduction, line-bundle obstruction, Cousin criterion and interpolation criterion; the analytic-geometry report’s fixed-divisor Bézout, corona and ultrafilter results on the unit disk [5], of which Part V’s are plane versions; and the trigonometry report’s zero sets of the surcomplex sine [6]. The entire-functions report’s Bézout theorem [7] concerns a different ring and does not conflict with Part V ([section 33.2](#)).

1.5 What is and is not claimed as new

The Hahn support calculus is classical [9, 10]. The analytic interpretation of restricted ordinary functions over surreal workspaces is classical [12], and the broader landscape is surveyed in [13]. Algebraic Birkhoff factorization in complete filtered noncommutative algebras, including its recursive form and the necessity of commutator corrections, is established [15, 16], as are the ordinary Laurent, Mittag–Leffler, Behnke–Stein and Cartan inputs [17, 18, 19], iterated path integrals [24], and the classical regular-singular and Riemann–Hilbert background [25, 26]. We reprove only what we need with explicit arbitrary-ordered-group support control, and we claim no new general factorization principle and no new ordinary analytic theorem.

We do not classify bundles or connections beyond the stated questions. Part I decides triviality on a puncture-and-disk cover for arbitrary finite rank and proves a sufficient theorem for arbitrary Stein covers; Part II inverts holonomy for near-identity representations on a punctured plane; Part III classifies one explicit family of bundles and nothing beyond it; Part V trivializes one family of pole-only cocycles after localization and classifies nothing. The boundaries in [section 30](#) are part of the theorem statements, not qualifications added afterwards.

2 The coefficient category

2.1 One set-sized workspace

For a complex algebra R , the Hahn algebra $R((t^\Gamma))$ consists of formal sums $F = \sum_{\gamma \in S} f_\gamma t^\gamma$ with $f_\gamma \in R$ whose nonzero support S is well ordered in the given order; addition is coefficientwise and multiplication is convolution. For matrices the support is the union of the supports of the finitely many entries, and an exponent occurs exactly when the corresponding coefficient matrix is not identically zero. The Hahn field of constants is

$$K_\Gamma = \mathbb{C}((t^\Gamma)), \quad \mathfrak{m}_\Gamma = \{a \in K_\Gamma : v(a) > 0\},$$

where v of a nonzero series is its least exponent and $v(0) = +\infty$. Divisibility of Γ and algebraic closedness of K_Γ are never used: we extract no roots and take no algebraic closure.

When $\Gamma \subseteq \mathbf{No}$ is an ordered additive subgroup, Conway normal forms give the interpretation

$$\mathbb{C}((t^\Gamma)) \longrightarrow \mathbf{No}[i] = \mathbf{SC}, \quad \sum c_\gamma t^\gamma \longmapsto \sum c_\gamma \omega^{-\gamma}, \quad (2.1)$$

with real and imaginary parts interpreted separately [12, 10, 14]. All supports here are sets; no sum indexed by a proper class, no proper-class fundamental group and no proper-class coefficient ring occurs anywhere.

All ordinary manifolds and Riemann surfaces carry their usual Hausdorff, second-countable topology. For a connected ordinary complex domain V set

$$\mathcal{H}_\Gamma(V) = \mathcal{O}(V)((t^\Gamma)), \quad (2.2)$$

$$\mathcal{I}_\Gamma(V) = \{F \in \mathcal{H}_\Gamma(V) : \text{supp } F \subseteq \Gamma_{\geq 0}\}, \quad (2.3)$$

$$\mathcal{J}_\Gamma(V) = \{F \in \mathcal{H}_\Gamma(V) : \text{supp } F \subseteq \Gamma_{> 0}\}. \quad (2.4)$$

Every coefficient must be holomorphic on the *same fixed domain* V ; we never substitute a family of convergent germs with unrelated radii, and $\mathcal{H}_\Gamma(V)$ is in general *not* the algebraic tensor product $K_\Gamma \otimes_{\mathbb{C}} \mathcal{O}(V)$, since a legitimate family may involve infinitely many linearly independent coefficient functions. For disconnected open sets, sections are specified component by component. Writing $\Omega_{\text{hol}}^1(V)$ for ordinary holomorphic one-forms, the same notation with one-form coefficients denotes one-form families. There is a reduction map

$$\text{red} : \mathcal{I}_\Gamma \longrightarrow \mathcal{O}, \quad \sum_{\gamma \geq 0} f_\gamma t^\gamma \longmapsto f_0, \quad \ker \text{red} = \mathcal{J}_\Gamma, \quad (2.5)$$

and the positive matrix group is

$$\mathcal{G}_{r,\Gamma}(V) = I_r + \text{Mat}_r(\mathcal{J}_\Gamma(V)). \quad (2.6)$$

Its elements are automatically two-sided invertible by lemma 3.4. Throughout, “positive” refers to *Hahn exponents only*: never to positivity of a Hermitian matrix, never to positivity of entries, and never to positive-definiteness. Elements of $I_r + \text{Mat}_r(\mathfrak{m}_\Gamma)$ need not be algebraically unipotent.

2.2 Two standing conventions that differ between the two parts

The two parts of this report were developed independently and do not make identical structural assumptions. Both assumptions are correct as stated and neither is weakened here.

Part I *proves and uses* the sheaf property of these assignments on the ordinary base. The point is a support point. A nonzero ordinary holomorphic coefficient on a connected domain cannot vanish on a nonempty open subset, so restriction between nonempty connected ordinary domains preserves support. If compatible local sections are given on a cover of a connected domain, the identity theorem along chains of intersecting patches shows that their coefficient functions glue and that the global support equals the support seen on any one nonempty connected patch; that support is well ordered. For disconnected opens one takes the product over connected components. In particular, if a section on $X \setminus A$ extends separately across each point of a discrete A , it extends *with its existing global support*: no new exponent can appear only at an isolated ordinary point. This observation is used whenever a matrix quotient is extended across a puncture. Part III follows Part I’s convention: it proves and uses the same sheaf property, for $\mathcal{H}_\Gamma$, $\mathcal{I}_\Gamma$, $\mathcal{J}_\Gamma$ and for the meromorphic sheaf $\mathcal{M}_\Gamma$ (lemma 20.2).

Part II deliberately does *not* assume that the global-support assignment is a sheaf for arbitrary covers, and none of its proofs uses that. All its gluing is done directly, by descending families already supported in one specified monoid, and its only analytic input is the ordinary Mittag–Leffler theorem applied one exponent at a time. Consequently no application of a Cartan-type theorem to an infinite-dimensional Hahn coefficient sheaf is hidden in Part II, and no companion claim about Hahn Noetherianity or a Hahn Nullstellensatz is relied upon anywhere in this report.

2.3 Comparison of names across the two parts

The following table fixes one vocabulary. Where the two source manuscripts used different names for the same object, the merged name is given first; where they used the same word for genuinely different objects, the objects are separated. The renamings made for Part III are listed separately, in section 20.5.

Merged name	Part I usage	Part II usage
$\mathcal{H}_\Gamma(V), \mathcal{I}_\Gamma(V), \mathcal{J}_\Gamma(V)$	full, integral and positive common-domain sheaves; sheaf property proved on the ordinary base	$\mathcal{J}_\Gamma(V)$ is the positive part of the common-domain family; no sheaf property assumed
$\mathcal{G}_{r,\Gamma}(V) = I_r + \text{Mat}_r(\mathcal{J}_\Gamma(V))$	the positive matrix group being split	the group of positive gauges; framed gauges have value I_r at the base point, and ρ takes values in the constant subgroup
$\mathfrak{m}_\Gamma$	not used; positivity is imposed sheaf-wise	positive-valuation elements of K_Γ ; ρ takes values in $I_r + \text{Mat}_r(\mathfrak{m}_\Gamma)$
Well-ordered union of supports	taken over normalized <i>polar factors</i> $\text{Pol}(G_a) - I_r$: well-ordering of $\mathcal{E}(G)$ is invariant	taken over the <i>monodromy matrices</i> $\rho(\ell_d) - I_r$: well-ordering of E_ρ is invariant
Local positive factorization	$G = \text{Pol}(G) \text{Reg}(G)$ on a punctured disk, normalized Birkhoff type	$Y = H \exp(R \text{Log } z)$ on a punctured disk, positive Frobenius type
$\Sigma(\cdot)$	the rank-one logarithmic singular support, used only for the scalar comparison	not used
Strong summability	implicit in Hahn summability of finite expressions	a named condition on set-indexed families; distinguished sharply from valuation-topology convergence

Remark 2.1 (Resonance does not occur in this report). No part of this report imposes, or needs, any arithmetic or small-divisor hypothesis, and the word *resonance* appears here only as an absence: in [theorem 5.4](#) no spectral resonance operator is inverted, because positivity forces the exponent-zero part of the residue to vanish. Other reports in this collection use “resonance” in several mutually inequivalent senses; none of them is in force here, and no theorem of this report may be combined with one of those hypotheses by analogy.

Remark 2.2 (Which exponential and which logarithm). The matrix $\exp$ and $\log$ in this report are formal one-variable substitutions from [lemma 3.4](#), computed coefficientwise on a fixed ordinary domain. They are mutually inverse maps $\exp : \text{Mat}_r(\mathcal{J}_\Gamma(V)) \rightarrow \mathcal{G}_{r,\Gamma}(V)$ and $\log : \mathcal{G}_{r,\Gamma}(V) \rightarrow \text{Mat}_r(\mathcal{J}_\Gamma(V))$. For commuting arguments the usual addition identities hold; they are not imposed on noncommuting arguments. Ordinary scalar logarithms such as $\log 2$ and the branch $\text{Log } z$ retain their ordinary complex meanings. No matrix exponential here is the Ehrlich–Kaplan surcomplex exponential, or any radial or quaternionic extension of it, whose homomorphism properties are separate questions belonging to separate reports. The scalar identity $\text{Pol}(G) = \exp(\pi_- \log G)$ of [\(5.5\)](#) is used only in rank one, and [proposition 8.4](#) shows what happens if one tries to use it in rank three.

2.4 Surcomplex evaluation on a halo

At an ordinary point $z_0 \in V$ and an $\varepsilon \in \mathfrak{m}_\Gamma$, a positive family is evaluated by the ordinary Taylor series of its coefficients:

$$F^\#(z_0 + \varepsilon) = \sum_\gamma \sum_{k \geq 0} t^\gamma \frac{F_\gamma^{(k)}(z_0)}{k!} \varepsilon^k, \quad (2.7)$$

the exponent-zero term being allowed separately in the integral case. If S is the Hahn support and $T = \text{supp } \varepsilon$, the resulting support lies in $S + \langle T \rangle_\mathbb{N}$, and each fixed exponent receives only finitely many contributions by [lemma 3.2](#). So our matrices do give matrices on finite surcomplex thickenings of ordinary charts, entry by entry. This is offered only to explain why the category deserves to be called surcomplex. The ordinary complex base is not being replaced by a fine-topological surcomplex manifold, and no monodromy or gluing statement below is obtained by transport along a fine-continuous surcomplex loop.

2.5 Ordinary paths, not fine-surreal paths

For a piecewise smooth ordinary path c in an ordinary domain, integrals of coefficient one-forms and their iterated integrals are ordinary complex integrals; their values are then assembled as Hahn coefficients. Paths, the universal cover, iterated integrals and Mittag–Leffler are all ordinary complex. Only the assembly of coefficients is Hahn. We assert no convergence of partial Hahn sums in the fine topology of **SC**, and [example 17.3](#) shows that such partial sums can fail to converge even in the valuation topology. This is the same domain separation stated in the repository’s differential-equations report [\[3\]](#), whose wording we adopt rather than re-argue: the loop is an ordinary complex path, continuation is coefficientwise, and the result is a matrix of Hahn constants.

3 Support calculus without any convergence assumption

Definition 3.1 (Strong summability). A set-indexed family of Hahn series, or of matrices of Hahn series, is *strongly summable* if the union of its supports is well ordered and each exponent occurs in

the support of only finitely many members. Its sum is then the finite coefficientwise sum at each exponent. This is not valuation-topology convergence, which is never used below.

Lemma 3.2 (Neumann support calculus; classical input). *Let $E \subseteq \Gamma_{>0}$ be well ordered and let $M = \langle E \rangle_{\mathbb{N}} = \{0\} \cup \{e_1 + \cdots + e_n : n \geq 1, e_j \in E\}$. Then M is well ordered, and for each $\gamma \in \Gamma$ only finitely many finite words in E , across all word lengths, have sum γ . Consequently, if $S \subseteq \Gamma$ is well ordered then $S + M$ is well ordered, a fixed element has only finitely many decompositions with one summand in each of two well-ordered sets, and every fixed coefficient of an expression formed by adjoining finitely many positive-support factors receives only finitely many contributions.*

This is the positive-support form of Neumann’s lemma [9], with the maximally complete field background in [10]. We use it as an established combinatorial input and do not rename it. It is strictly stronger than the statement that all exponents are positive, and the finite-word clause across all word lengths is exactly what powers noncommutative Dyson expansions and geometric inverses below. Both source manuscripts of this report state and use it independently, in the same form.

Remark 3.3 (No hidden Archimedean or cofinality assumption). A least positive input exponent δ need not have the property that $n\delta$ eventually exceeds a prescribed γ . In $\mathbb{Z}^2$ with lexicographic order, $n(0, 1) < (1, 0)$ for every ordinary integer n . In $\Gamma = \mathbb{Q} + \mathbb{Q}\omega$ with its induced surreal order, $n < \omega$ for every ordinary positive integer n , so repeatedly increasing a valuation by 1 never reaches ω ; nevertheless $\sum_{n \geq 0} t^n$ is a valid Hahn sum and the support calculus works equally well at the exponents ω and $\omega + 1$, which is where the correction of [proposition 18.1](#) will sit. No proof below infers coefficient finiteness from that false cofinality. Every recursion is a well-founded recursion on a well-ordered support set.

Lemma 3.4 (Positive inversion and formal functions). *Let V be an ordinary domain and $A \in \text{Mat}_r(\mathcal{J}_\Gamma(V))$, with $\text{supp } A \subseteq E$ for a positive well-ordered E . Then*

$$(I_r + A)^{-1} = \sum_{k \geq 0} (-A)^k, \quad \exp A = \sum_{k \geq 0} \frac{A^k}{k!}, \quad \log(I_r + A) = \sum_{k \geq 1} \frac{(-1)^{k+1}}{k} A^k$$

are strongly summable coefficientwise on V , with nonconstant supports in $\langle E \rangle_{\mathbb{N}} \setminus \{0\}$. The inverse is two-sided, and $\exp$ and $\log$ are mutually inverse as one-variable formal substitutions. They do not make matrix multiplication additive. The same statements hold with coefficients in any associative complex algebra for which the finitely many operations in each displayed coefficient are defined.

Proof. Expand an entry of A^k : its terms are indexed by a word of k positive exponents together with a path through r matrix indices. At a fixed total exponent only finitely many exponent words occur by [lemma 3.2](#), and each word admits only finitely many matrix-index paths. Hence every displayed coefficient sum is finite and its value is holomorphic on the original V , and all supports lie in $\langle E \rangle_{\mathbb{N}}$. Ordinary formal identities in one variable, and the geometric-series cancellation, may therefore be checked coefficientwise, because at a fixed exponent all sufficiently long words fail to contribute. Noncommuting independent arguments do not satisfy the scalar exponential addition law. $\square$

Lemma 3.5 (Finite decomposition inside the generated monoid). *Put $S = \langle E \rangle_{\mathbb{N}} \setminus \{0\}$ for a positive well-ordered E . Each $\gamma \in \Gamma$ has only finitely many ordered decompositions into members of S , and only finitely many members of S can occur as a part in such a decomposition of a fixed γ .*

Proof. Choose an E -word representing each member of S . Flattening a word in S of total weight γ gives an E -word of that weight, and there are finitely many such by [lemma 3.2](#). A fixed finite word

has only finitely many partitions into nonempty consecutive blocks, and every original S -word is recovered from one of these finitely many possibilities. Taking all parts of those words gives the second assertion. $\square$

[Lemma 3.5](#) is what makes the effectivity statement [proposition 28.1](#) possible; it is not needed for [lemma 3.4](#).

4 The ordinary analytic inputs

Everything analytic in this report is an ordinary complex theorem applied at *one Hahn exponent at a time*. That is the reason the criteria exist at all: the ordinary problems are always solvable, and the obstruction is that their outputs across exponents must assemble into one legitimate Hahn section. The two parts use two different packagings of the ordinary input, and both are kept, because each theorem below uses its own.

4.1 A fixed-disk Laurent splitting

Fix $D = \{|u| < \rho\}$ and $D^* = D \setminus \{0\}$. A function $f \in \mathcal{O}(D^*)$ has the Laurent decomposition

$$f = f_- + f_+, \quad f_- = \sum_{k \geq 1} a_{-k} u^{-k}, \quad f_+ = \sum_{k \geq 0} a_k u^k. \quad (4.1)$$

Write $\mathcal{P}$ for the algebra of all such negative parts and π_-, π_+ for the projections, acting entrywise on matrices.

Lemma 4.1 (The ordinary splitting algebras). *There is a direct-sum decomposition of complex vector spaces $\mathcal{O}(D^*) = \mathcal{P} \oplus \mathcal{O}(D)$. Both summands are closed under multiplication, although $\mathcal{P}$ has no multiplicative identity; their intersection is zero; and the decomposition does not shrink D .*

Proof. For every $s \in (0, \rho)$, Cauchy’s formula bounds the negative Laurent coefficients by $|a_{-k}| \leq M_s s^k$, so the series in $1/u$ converges for $|u| > s$ with s arbitrarily small, and vanishes at infinity. Conversely a function holomorphic on $\mathbb{C} \setminus \{0\}$ and vanishing at infinity has only negative Laurent powers. Products retain these properties. A function in the intersection extends across zero and across infinity, hence is constant on the sphere with value zero at infinity. The positive part converges on the original disk of radius ρ . $\square$

The negative part may have an essential singularity at zero; no bound on pole orders is imposed. The explicit obstruction families below use only simple poles.

4.2 Isolated singular parts: the Part I packaging

Lemma 4.2 (Isolated singular parts on the plane). *Let $A \subset \mathbb{C}$ be closed and discrete and let p_a be a prescribed negative Laurent part in $z - a$ at each $a \in A$. There exists $f \in \mathcal{O}(\mathbb{C} \setminus A)$ such that $f - p_a$ extends holomorphically through each disk D_a . One may additionally prescribe $f(z_*)$ at one fixed $z_* \notin A$.*

Proof. The finite case is a finite sum. Otherwise enumerate $A = \{a_n\}$ with $|a_n| \rightarrow \infty$ and choose radii $R_n \rightarrow \infty$ with $R_n < |a_n|$ for all large n . Each $p_{a_n}(z - a_n)$ is holomorphic on $|z| < |a_n|$; choose

a Taylor polynomial q_n at zero with $\sup_{|z| \leq R_n} |p_{a_n}(z - a_n) - q_n(z)| < 2^{-n}$. After separating finitely many early indices, $f(z) = \sum_n (p_{a_n}(z - a_n) - q_n(z))$ converges normally on every compact subset of $\mathbb{C} \setminus A$. Near a fixed a_m all terms except the m th form a normally convergent holomorphic series, so the required negative part is p_{a_m} , and the differences extend over the full D_{a_m} by uniqueness. Adding a constant fixes $f(z_*)$. $\square$

The proof allows essential isolated singularities, because it approximates the entire negative part away from its centre, and uses no bound on pole orders. The same correcting-polynomial mechanism appears in the scalar Cousin step of the repository’s global-divisor report [2].

Lemma 4.3 (Open-surface version). *The conclusion of lemma 4.2 holds on any connected open Riemann surface X , with A closed and discrete, pairwise disjoint coordinate disks D_a , and negative Laurent parts in the chosen coordinates.*

Proof. An open Riemann surface is Stein by the Behnke–Stein theorem [17]. The data p_a define an ordinary additive holomorphic cocycle on the cover $X \setminus A$, D_a . Cartan’s theorem B gives $H^1(X, \mathcal{O}) = 0$ and hence a splitting of that cocycle; in the chosen sign convention the central component is the desired f . A trivial additive torsor has a section expressed in the original charts, so no acyclicity assumption on this particular cover is needed. A constant again prescribes one value. See [18, 19] for the ordinary coherent-sheaf input. $\square$

On the plane, lemma 4.2 makes the analytic existence step of theorem 6.2 self-contained. The general open-surface statement uses exactly the two established ordinary theorems named in its proof, and not an unproved theorem B for a Hahn coefficient sheaf.

4.3 The period map and a linear section: the Part II packaging

Let $D \subset \mathbb{C}$ be closed and discrete. Every compact set meets D in a finite set, so D is at most countable, and $U = \mathbb{C} \setminus D$ is connected.

Lemma 4.4 (Free based meridians). *There is a collection of based positively oriented meridians $(\ell_d)_{d \in D}$ forming a free basis of $\pi_1(U, p)$; every group element is a finite word in these meridians and their inverses.*

Proof. Exhaust $\mathbb{C}$ by nested disks centred at p with boundaries missing D . Each disk contains finitely many punctures and has free fundamental group generated by based meridians. When the disk is enlarged, the old meridians can be retained and meridians for the new punctures added: cutting the annular region along arcs to the newly included punctures, or applying van Kampen, exhibits the old group as a free factor. Choose these bases compatibly. A loop and any specified homotopy have compact image, hence occur in a sufficiently large disk, so the fundamental group of the union is free on the union of the compatible bases. For $D = \emptyset$ the group is trivial. $\square$

We fix such a basis throughout Part II. The numerical matrices assigned to based meridians depend on the chosen tails and on the base point, and that dependence is not suppressed. So that $dY = \Omega Y$ yields a homomorphism for matrix multiplication, $\alpha\beta$ means traversal of β first and then α . All meridians are positively oriented.

Let $\mathcal{L}(D)$ be the complex vector space of meromorphic one-forms on $\mathbb{C}$ with at most simple poles at the points of D and no other finite poles. There is *no condition at infinity*. Define

$$P : \mathcal{L}(D) \longrightarrow \mathbb{C}^D, \quad (P\alpha)_d = \int_{\ell_d} \alpha = 2\pi i \operatorname{Res}_d \alpha. \quad (4.2)$$

The target is the full *product* of complex vector spaces, not the direct sum. That is the whole reason no summability across punctures is needed in [theorem 14.2](#).

Lemma 4.5 (A linear period right inverse). *The map P is surjective, and there is a complex-linear $\sigma : \mathbb{C}^D \rightarrow \mathcal{L}(D)$ with $P\sigma = \operatorname{id}$. For finite D one may take*

$$\sigma((v_d)_{d \in D}) = \frac{1}{2\pi i} \sum_{d \in D} v_d \frac{dz}{z - d}. \quad (4.3)$$

For $D = \emptyset$ the right inverse has zero image.

Proof. Prescribing the principal part $v_d dz/(2\pi i(z - d))$ at each d and applying the ordinary Mittag–Leffler theorem proves surjectivity; the elementary mechanism is the correcting-polynomial argument of [lemma 4.2](#), applied to the one-form data. That construction need not be linear in the input, because the chosen polynomial degrees may depend on it. To obtain a linear section, split the surjection of complex vector spaces: choose a basis of $\mathbb{C}^D$, choose one preimage of each basis element, and extend linearly. This uses the axiom of choice for vector spaces. For finite D the displayed finite sum is an explicit linear section. $\square$

Remark 4.6 (What is and is not required of the section). For infinite D we assert neither continuity nor a norm estimate nor an effective algorithm for σ ; [proposition 28.3](#) shows that continuity is in fact impossible for the natural topologies. In particular $\sum_d v_d/(z - d)$ cannot simply be written without a convergence correction. The theorem needs only an ordinary complex-linear splitting. The section is chosen once, independently of the Hahn exponent, and is applied *entrywise* to matrix data; that last point is what makes the traceless refinement [theorem 16.3](#) true.

Both packagings are the ordinary Mittag–Leffler theorem, and neither is derived from the other here. [Lemma 4.2](#) gives, for each exponent separately, one function with prescribed negative parts of unbounded order; [lemma 4.5](#) gives a single *linear* choice of simple-pole one-form, valid simultaneously for all exponents, at the cost of the axiom of choice and of continuity. Part I needs the first because its prescribed local data are arbitrary negative Laurent matrices; Part II needs the second because uniqueness of its normalized realizer is exactly injectivity of P on $\operatorname{im} \sigma$.

5 Local positive factorizations at a puncture

Two local normal forms are used, one per part. They are different normalizations of different objects and neither is a consequence of the other.

5.1 The normalized polar factorization

For the chosen disk and coordinate put

$$\mathcal{P}_{r,\Gamma} = I_r + \operatorname{Mat}_r(\mathcal{P}((t^\Gamma))_{>0}),$$

meaning that the nonconstant coefficients lie in $\mathcal{P}$ and have positive well ordered support. This is a group: nonconstant coefficients of a product or of a positive inverse are finite sums of products of negative Laurent parts, and each such product still vanishes at auxiliary infinity.

Theorem 5.1 (Local normalized factorization). *Let $G \in \mathcal{G}_{r,\Gamma}(D^*)$ and $S = \text{supp}(G - I_r)$. There exist unique matrices*

$$P \in \mathcal{P}_{r,\Gamma}, \quad H \in \mathcal{G}_{r,\Gamma}(D), \quad G = PH, \quad (5.1)$$

and both nonconstant supports lie in $M = \langle S \rangle_{\mathbb{N}} \setminus \{0\}$. We write $P = \text{Pol}(G)$ and $H = \text{Reg}(G)$. Every coefficient of H is holomorphic on the original disk, not on an exponent-dependent smaller disk.

Proof. The case $S = \emptyset$ is immediate. Otherwise write $G = I_r + \sum_{\gamma > 0} g_{\gamma} t^{\gamma}$, $P = I_r + \sum_{\gamma \in M} p_{\gamma} t^{\gamma}$, $H = I_r + \sum_{\gamma \in M} h_{\gamma} t^{\gamma}$, with $g_{\gamma} = 0$ at exponents not originally present, and use well-founded recursion on M . Once all smaller coefficients are chosen, define

$$r_{\gamma} = g_{\gamma} - \sum_{\substack{\eta + \theta = \gamma \\ \eta, \theta > 0}} p_{\eta} h_{\theta}, \quad (5.2)$$

$$p_{\gamma} = \pi_{-}(r_{\gamma}), \quad h_{\gamma} = \pi_{+}(r_{\gamma}). \quad (5.3)$$

Every summand involves strictly smaller exponents and the sum is finite by lemma 3.2, so each step produces a negative Laurent matrix and a matrix holomorphic on D . The series have supports in M and satisfy $PH = G$ coefficientwise; their inverses exist by lemma 3.4.

For uniqueness, suppose $P_1 H_1 = P_2 H_2$, so $P_2^{-1} P_1 = H_2 H_1^{-1}$. The nonconstant coefficients on the left lie in $\mathcal{P}$ and those on the right are holomorphic on D . After placing the finitely many input supports in a common positive generated monoid, coefficientwise comparison and lemma 4.1 force both sides to equal I_r . $\square$

Proposition 5.2 (Right-gauge invariance). *If $Q \in \mathcal{G}_{r,\Gamma}(D)$ then*

$$\text{Pol}(GQ) = \text{Pol}(G). \quad (5.4)$$

The analogous statement for multiplication on the left is generally false.

Proof. The factorization $GQ = \text{Pol}(G)(\text{Reg}(G)Q)$ has the defining properties of theorem 5.1, and uniqueness gives the equality. For failure on the left, choose $\delta > 0$, $s = t^{\delta}$, $G = I_3 + (s/u)E_{23}$ and $Q = I_3 + sE_{12}$. Direct multiplication gives

$$QG = (I_3 + (s/u)E_{23} + (s^2/u)E_{13})Q.$$

Here E_{ij} are matrix units. The first factor is polar and Q is regular, so it is $\text{Pol}(QG)$, which differs from $\text{Pol}(G) = G$. This exhibits the multiplication order explicitly. $\square$

Corollary 5.3 (Unitriangular factors). *If G is upper unitriangular then so are $\text{Pol}(G)$ and $\text{Reg}(G)$. If a fixed pattern of strictly upper-triangular entries is closed under matrix multiplication, the factors preserve that pattern too.*

Proof. Both Laurent projections preserve zero entries, and every nonlinear term in (5.2) is a product of previously constructed matrices in the specified multiplicatively closed pattern. Induction finishes the proof. $\square$

5.2 Where the obstruction is manufactured

For a single positive parameter, writing $G = I_r + tg_1 + t^2g_2 + t^3g_3 + \cdots$, the recursion begins

$$\begin{aligned} p_1 &= \pi_- g_1, & h_1 &= \pi_+ g_1, \\ p_2 &= \pi_- (g_2 - p_1 h_1), & h_2 &= \pi_+ (g_2 - p_1 h_1), \\ p_3 &= \pi_- (g_3 - p_1 h_2 - p_2 h_1). \end{aligned}$$

The second-order correction already contains the product of a polar part and a regular part, and it can be nonzero even when $g_2 = 0$. With different positive exponents at different punctures, the exponents of such products can form a descending sequence absent from the raw singular data. That single mechanism is the whole content of [theorem 8.2](#) and [theorem 9.2](#).

In rank one the commuting logarithmic parts give the closed form

$$\text{Pol}(G) = \exp(\pi_- \log G), \quad \text{Reg}(G) = \exp(\pi_+ \log G), \quad (5.5)$$

with the projections acting coefficientwise and the scalar exponential addition law verifying the factorization. The same argument applies to a matrix input if its two projected logarithmic parts commute. For general matrix inputs, replacing [theorem 5.1](#) by (5.5) is invalid; the algebraic Birkhoff literature describes the associated Baker–Campbell–Hausdorff corrections in a filtered setting [[15](#), [16](#)], and our direct recursion avoids expanding that formula. [Proposition 8.4](#) exhibits the exact numerical discrepancy in rank three.

5.3 The positive Frobenius form

The next theorem is the local normal form used in Part II. It explains why residues still control *local* monodromy while being unreadable directly from the prescribed *based* monodromy matrices. It is a support-controlled version of the elementary nonresonant regular-singular construction, not a new claim about classical Frobenius theory.

Theorem 5.4 (A common-disk positive Frobenius factorization). *Let Δ be an ordinary disk about 0 and suppose a positive Hahn connection on $\Delta \setminus \{0\}$ has the form*

$$\Omega = R \frac{dz}{z} + B(z) dz, \quad (5.6)$$

with $R \in \text{Mat}_r(\mathfrak{m}_\Gamma)$ and $B \in \text{Mat}_r(\mathcal{J}_\Gamma(\Delta))$ sharing one positive well ordered support E . Then there is a unique $H \in I_r + \text{Mat}_r(\mathcal{J}_\Gamma(\Delta))$ with $H(0) = I_r$ such that, on a branch of the logarithm,

$$Y(z) = H(z) \exp(R \text{Log } z) \quad (5.7)$$

solves $dY = \Omega Y$. The supports of $H - I_r$ and $H^{-1} - I_r$ lie in $\langle E \rangle_{\mathbb{N}} \setminus \{0\}$, and the local monodromy based at $z_0 \in \Delta \setminus \{0\}$ is

$$H(z_0) \exp(2\pi i R) H(z_0)^{-1}. \quad (5.8)$$

Proof. The required equation is $H' = [R, H]/z + BH$ with $H(0) = I_r$. Construct its coefficients recursively on $\langle E \rangle_{\mathbb{N}} \setminus \{0\}$: at weight γ one must solve

$$H'_\gamma = B_\gamma + \sum_{\substack{\alpha+\beta=\gamma \\ \alpha, \beta > 0}} \left(B_\alpha H_\beta + \frac{[R_\alpha, H_\beta]}{z} \right), \quad H_\gamma(0) = 0,$$

a finite sum. The previously constructed H_β vanish at 0, so H_β/z is holomorphic on the entire disk; the right-hand side is therefore holomorphic on that same disk, and integration from 0 defines a unique holomorphic H_γ vanishing there. The resulting family has the claimed support and its inverse is given by [lemma 3.4](#). A least-difference argument proves uniqueness.

Because R has positive support, $\exp(R \operatorname{Log} z)$ is a strongly summable family of ordinary holomorphic functions on any logarithm branch, and coefficientwise differentiation gives $(\exp(R \operatorname{Log} z))' = R \exp(R \operatorname{Log} z)/z$; the equation for H then verifies (5.7). Continuing once positively around 0 increases $\operatorname{Log} z$ by $2\pi i$ and multiplies the fundamental matrix on the right by $\exp(2\pi i R)$. Normalizing at z_0 conjugates that factor by $H(z_0) \exp(R \operatorname{Log} z_0)$, and the exponential factor commutes with $\exp(2\pi i R)$ and cancels, giving (5.8). $\square$

There is no division by a spectral resonance operator in this proof: the exponent-zero coefficient of the residue is zero, so the equation at each new exponent is an ordinary primitive problem. Positivity is doing real work here (compare [remark 2.1](#)).

If a meridian is based at a distant point p , its tail contributes a further transport conjugation to (5.8), and with several punctures these conjugations interact. Choosing $R_d = (2\pi i)^{-1} \log M_d$ separately at each puncture therefore need not solve the based inverse problem. The explicit failure appears as a mixed commutator term in [proposition 18.1](#).

Part I. Multiplicative gluing: the polar-support criterion

6 The sharp global matrix Cousin criterion

Let X be a connected open Riemann surface, $A \subset X$ closed and discrete, and choose pairwise disjoint coordinate disks (D_a, u_a) with $u_a(a) = 0$. Put $U_0 = X \setminus A$ and $W_a = D_a \setminus \{a\}$; the punctured surface U_0 is connected. For each a , *independently*, let $G_a \in \mathcal{G}_{r,\Gamma}(W_a)$. The disks do not meet one another, so this is a complete positive matrix cocycle on the displayed cover, with the inverse transition on the reversed overlap. There need not be one well-ordered set containing the supports of all the G_a .

Definition 6.1 (Normalized polar support). Factor $G_a = P_a H_a$ by [theorem 5.1](#) and define

$$\mathcal{E}(G) = \bigcup_{a \in A} \operatorname{supp}(P_a - I_r) \subseteq \Gamma_{>0}. \quad (6.1)$$

The datum is *polar-admissible* when $\mathcal{E}(G)$ is well ordered. No sum over a is intended in (6.1): it is a set-theoretic union of supports, taken across all matrix entries.

Theorem 6.2 (Exact nonabelian Cousin criterion: Criterion P). *The following are equivalent.*

- (i) *There exist $B_0 \in \mathcal{G}_{r,\Gamma}(U_0)$ and $B_a \in \mathcal{G}_{r,\Gamma}(D_a)$ with $G_a = B_0 B_a^{-1}$ on every W_a .*
- (ii) *The set $\mathcal{E}(G)$ is well ordered.*

When these hold, let $M = \langle \mathcal{E}(G) \rangle_{\mathbb{N}}$. One can choose

$$\operatorname{supp}(B_0 - I_r) \subseteq M \setminus \{0\}, \quad (6.2)$$

and in the local factorizations $B_0 = P_a Q_a$ the matrices $Q_a - I_r$ can also be supported in $M \setminus \{0\}$. The disk trivializers are $B_a = H_a^{-1} Q_a$ and hence have individually legitimate supports. A value $B_0(z_*) = I_r$ may be imposed at any chosen $z_* \in U_0$.

Proof. Necessity. Suppose $G_a = B_0 B_a^{-1}$. By right-gauge invariance (5.4),

$$P_a = \text{Pol}(G_a) = \text{Pol}(B_0|_{W_a}).$$

Let $S = \text{supp}(B_0 - I_r)$. The local support estimate in theorem 5.1 is independent of the puncture and of the Laurent coefficients of the input, so $\text{supp}(P_a - I_r) \subseteq \langle S \rangle_{\mathbb{N}} \setminus \{0\}$ for every a . That one generated monoid is well ordered by lemma 3.2, and $\mathcal{E}(G)$ is a subset of it.

Sufficiency. Assume $E = \mathcal{E}(G)$ is well ordered and put $M = \langle E \rangle_{\mathbb{N}}$. We construct a global $F \in \mathcal{G}_{r,\Gamma}(U_0)$ and local $Q_a \in \mathcal{G}_{r,\Gamma}(D_a)$ with

$$F = P_a Q_a \tag{6.3}$$

and every nonconstant coefficient supported in M . Write $P_a = I_r + \sum_{\gamma > 0} p_{a,\gamma} t^\gamma$, $F = I_r + \sum_{\gamma > 0} f_\gamma t^\gamma$, $Q_a = I_r + \sum_{\gamma > 0} q_{a,\gamma} t^\gamma$. At a positive $\gamma \in M$, with all smaller coefficients constructed, define on W_a

$$R_{a,\gamma} = p_{a,\gamma} + \sum_{\substack{\eta + \theta = \gamma \\ \eta, \theta > 0}} p_{a,\eta} q_{a,\theta}, \tag{6.4}$$

a finite sum. Apply lemma 4.3 entry by entry to the prescribed ordinary negative Laurent matrices $\pi_- R_{a,\gamma}$. It supplies a single matrix $f_\gamma \in \text{Mat}_r(\mathcal{O}(U_0))$ such that $f_\gamma - R_{a,\gamma}$ extends holomorphically across D_a for every a ; set

$$q_{a,\gamma} = f_\gamma - R_{a,\gamma}. \tag{6.5}$$

The prescribed zero value at z_* can be imposed on every f_γ .

Recursion on the well-ordered set $M \setminus \{0\}$ constructs all coefficients. The scalar analytic step changes coefficient functions but introduces no new Hahn exponents, and every coefficient of every Q_a is holomorphic on its original disk. The resulting Hahn matrices satisfy (6.3) coefficientwise and are invertible by lemma 3.4. Finally $G_a = P_a H_a$ and $F = P_a Q_a$ give $G_a = F(H_a^{-1} Q_a)^{-1}$; take $B_0 = F$ and $B_a = H_a^{-1} Q_a$. The support of $H_a - I_r$ lies in the monoid generated by the original support of $G_a - I_r$ and the support of $Q_a - I_r$ lies in M ; a finite union of two well-ordered positive sets generates a legitimate monoid, so B_a is a well-defined section on D_a . There is no requirement that the disk supports have a globally well-ordered union. $\square$

6.1 Why the converse is sharp

An infinite puncture set does not by itself prevent a solution: at each fixed exponent the ordinary analytic problem is always solvable, and the obstruction is not a divergent ordinary sum, since the ordinary coefficient construction uses normal convergence independent of the formal exponents. The failure occurs exactly when the *necessary normalized polar information* cannot fit into a single well-ordered support.

Write $E_{\text{raw}} = \bigcup_a \text{supp}(G_a - I_r)$. If this set is well ordered, theorem 5.1 puts every $P_a - I_r$ in $\langle E_{\text{raw}} \rangle_{\mathbb{N}} \setminus \{0\}$, so Criterion P gives a splitting. Thus well-ordering of the full raw union is a sufficient condition. It is not necessary: G_a may itself be holomorphic on D_a , in which case $P_a = I_r$ and the solution $B_0 = I_r$, $B_a = G_a^{-1}$ exists even if those raw supports have a descending union. In contrast, the raw singular-coefficient union $\bigcup_a \text{supp}(\pi_-(G_a - I_r))$ fails both directions: discarding raw holomorphic coefficients can lose polar information created by their products with retained terms, and retained singular terms can cancel in the normalization. Section 8 gives both examples.

6.2 The space of solutions

Proposition 6.3 (Right torsor of splittings). *For a polar-admissible datum, all splittings are obtained from any one by*

$$(B_0, B_a) \mapsto (B_0 T, B_a T|_{D_a}), \quad T \in \mathcal{G}_{r,\Gamma}(X),$$

and the action is free and transitive. Under the normalization $B_0(z_*) = I_r$, use the subgroup with $T(z_*) = I_r$.

Proof. Right multiplication by a common T preserves $B_0 B_a^{-1}$. Conversely two splittings give $B_0^{-1} B'_0 = B_a^{-1} B'_a$ on W_a ; the central quotient extends through every disk by that equality, and the sheaf observation of [section 2.2](#) yields a global positive section T with one legitimate support on connected X . It is the unique element relating the two splittings. $\square$

This describes all solutions once a splitting exists. It is not a classification of the nonsplit objects; see [section 30](#).

7 Invariance, and the scalar case that is recovered

Corollary 7.1 (Removal of arbitrarily supported regular data). *For independently chosen $R_a \in \mathcal{G}_{r,\Gamma}(D_a)$, the families (G_a) and $(G_a R_a)$ have the same normalized polar factors and the same solvability status. No well-ordering condition on $\bigcup_a \text{supp}(R_a - I_r)$ is needed.*

Proof. Immediate from [proposition 5.2](#) and [theorem 6.2](#). $\square$

This is stronger than invariance under finitely many transformations: the infinitely many regular parts are genuinely irrelevant, but only *after* they have been removed in the correct multiplication order.

Corollary 7.2 (Coordinate independence of admissibility). *Changing the local coordinates, or shrinking the disks, can change individual normalized polar coefficients but cannot change whether their global support union is well ordered.*

Proof. Existence of the splitting in [theorem 6.2](#) is invariant under these changes. Shrinking a disk cannot turn a previously nonextendible germ into an extendible one: a local splitting on smaller disks yields extension of the same central matrix, hence a splitting on the original disks by analytic continuation of $G_a^{-1} B_0$. Alternatively, re-expressing a previously constructed polar factor in the new coordinate and applying [theorem 5.1](#) puts every new support in the monoid generated by the old support union, and the inverse coordinate change gives the converse. $\square$

Corollary 7.3 (Absoluteness under ordered extension). *Let $\Gamma \hookrightarrow \Delta$ be an order-preserving embedding of set-sized ordered abelian groups. A datum defined over Γ splits in the positive matrix category over Δ if and only if it splits over Γ .*

Proof. The recursive local polar factor computed in Γ is also a factor in Δ and remains the unique normalized factor, so its support union is the image of $\mathcal{E}(G)$. An order embedding preserves and reflects well-ordering of a subset. Apply [theorem 6.2](#) in both groups. $\square$

In particular the obstructions below cannot be removed by allowing more surreal exponents in the same integral positive-matrix problem. This is not a claim about passing to another analytic category, and in particular not a claim about admitting negative-valuation gauges; see [section 10](#) and item 2 of [section 30](#). For a different family of bundles, Part III proves the localized statement ([corollary 22.8](#)).

Proposition 7.4 (Recovery of the rank-one singular-support test). *Let $r = 1$, put $L_a = \log G_a$, and define the logarithmic singular support*

$$\Sigma(G) = \bigcup_a \text{supp}(\pi_- L_a).$$

Then

$$\mathcal{E}(G) \text{ well ordered} \iff \Sigma(G) \text{ well ordered.} \quad (7.1)$$

Proof. By (5.5), $\log P_a = \pi_- L_a$ is supported in the positive monoid generated by $\mathcal{E}(G)$, which gives the forward direction; conversely exponentiation puts every $P_a - I_1$ inside the monoid generated by $\Sigma(G)$. Apply [lemma 3.2](#). $\square$

Combined with [theorem 6.2](#), (7.1) says that in rank one our criterion is exactly the exact Cousin criterion already proved in the repository’s global-divisor report — its Theorem 13.1, stated there in terms of the singular support of the additive data on a puncture-and-disk cover [2]. We cite that theorem as the case we recover; it is not reproved here, and no priority for the scalar statement is claimed. What is new is that the correct matrix replacement for $\pi_- \log$ is the normalized polar factor and not the Laurent projection of a matrix logarithm: the scalar formula is not valid for general matrix inputs because logarithms need not turn products into sums ([remark 2.2](#)), and [proposition 8.4](#) measures the error.

8 A rank-three obstruction invisible in the raw poles

This section contains the example that forces the distinction of [section 12](#). Read it before generalizing either criterion.

8.1 The closed factorization in the Heisenberg group

Proposition 8.1 (Closed Heisenberg factorization). *Let*

$$G = \begin{pmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{pmatrix}, \quad a, b, c \in \mathcal{J}_\Gamma(D^*),$$

and write $a = a_- + a_+$, $c = c_- + c_+$ by coefficientwise Laurent projection. Then

$$\text{Pol}(G) = \begin{pmatrix} 1 & a_- & \pi_-(b - a_- c_+) \\ 0 & 1 & c_- \\ 0 & 0 & 1 \end{pmatrix}, \quad \text{Reg}(G) = \begin{pmatrix} 1 & a_+ & \pi_+(b - a_- c_+) \\ 0 & 1 & c_+ \\ 0 & 0 & 1 \end{pmatrix}. \quad (8.1)$$

Proof. All products are legitimate, being finite expressions in finitely many Hahn series. Direct multiplication verifies $\text{Pol}(G) \text{Reg}(G) = G$, the left factor is polar and the right factor is holomorphic on the full disk, so uniqueness in [theorem 5.1](#) identifies them as the canonical factors. $\square$

The subtraction of a_-c_+ is the essential term: a product of a *polar* coefficient of one entry with a *holomorphic* coefficient of another. Its exponent is a sum of two input exponents that need not appear in the input.

8.2 The obstructed datum

Take $X = \mathbb{C}$, $A = \{n : n \geq 2\}$, $D_n = \{|z - n| < 1/4\}$, $\Gamma = \mathbb{Q}$, and $u_n = z - n$. Consider

$$G_n = I_3 + \frac{t}{u_n} E_{12} + t^{1/n} E_{23}, \quad (8.2)$$

with E_{ij} the matrix units. Each G_n is a finite Hahn expression, has determinant exactly one, and reduces to I_3 . By (8.1),

$$P_n = I_3 + \frac{t}{u_n} E_{12} - \frac{t^{1+1/n}}{u_n} E_{13}, \quad H_n = I_3 + t^{1/n} E_{23}, \quad (8.3)$$

and the product $P_n H_n$ has zero $(1, 3)$ entry because its two contributions cancel.

Theorem 8.2 (Hidden rank-three obstruction). *The matrices (8.2) admit no splitting by general positive 3×3 matrices. Their abelianized upper-unitriangular cocycle does split. Their raw nonremovable coefficient support is the singleton $\{1\}$, and their determinant cocycle is identically one.*

Proof. The normalized polar support contains $\{1 + 1/n : n \geq 2\}$, a strictly decreasing infinite sequence, hence is not well ordered, and theorem 6.2 rules out a splitting even in the full positive matrix group.

The abelianization of the upper-unitriangular 3×3 group remembers the $(1, 2)$ and $(2, 3)$ entries and adds them componentwise. Let $\phi \in \mathcal{O}(U_0)$ have principal part $1/(z - n)$ at every $n \geq 2$ and no singularities outside A ; such ϕ exists by lemma 4.2, one explicit choice being

$$\phi(z) = \sum_{n=2}^{\infty} \left(\frac{1}{z - n} + \frac{1}{n} \right), \quad (8.4)$$

whose n th summand is $O_K(n^{-2})$ on each compact set away from A . The $(1, 2)$ component splits using the central function $t\phi$ and the disk functions $t(\phi - 1/u_n)$; the $(2, 3)$ component splits using central function 0 and disk functions $-t^{1/n}$. Finally the only original singular coefficient is E_{12}/u_n at exponent 1, and unitriangular matrices have determinant one. $\square$

The example is stronger than a determinant obstruction, and stronger even than the statement that a noncommutative cocycle has nontrivial abelianization: the abelianized problem *is* solvable, and the obstruction appears only when its solutions are required to fit together in the original multiplication law. The obstruction also lives in a matrix entry that is identically zero in the input.

8.3 Adding a singular term repairs the problem

Example 8.3 (The repair, which worsens the raw support). Define

$$\tilde{G}_n = I_3 + \frac{t}{u_n} E_{12} + t^{1/n} E_{23} + \frac{t^{1+1/n}}{u_n} E_{13}. \quad (8.5)$$

Then

$$\tilde{G}_n = \left(I_3 + \frac{t}{u_n} E_{12}\right)(I_3 + t^{1/n} E_{23}),$$

so $\text{Pol}(\tilde{G}_n) = I_3 + tE_{12}/u_n$ and $\mathcal{E}(\tilde{G}) = \{1\}$, and a global splitting exists. Explicitly, with ϕ as in (8.4), put $F = I_3 + t\phi E_{12}$,

$$Q_n = I_3 + t\left(\phi - \frac{1}{u_n}\right)E_{12}, \quad \tilde{H}_n = I_3 + t^{1/n}E_{23};$$

then $F = \text{Pol}(\tilde{G}_n)Q_n$ and $\tilde{G}_n = F(\tilde{H}_n^{-1}Q_n)^{-1}$, all disk factors being holomorphic on D_n . Yet the raw singular support of (8.5) contains the descending set $\{1 + 1/n\}$.

So (8.2) has raw singular support $\{1\}$ and is obstructed, while (8.5) has a descending raw singular support and is solvable. Raw singular support is therefore neither a sufficient nor a necessary test, in the strongest possible sense: the two directions fail on data that differ by one term.

8.4 Why a projected matrix logarithm is not a repair

Proposition 8.4 (Logarithm diagnostic). *The matrix logarithm of (8.2) is finite, namely*

$$\log G_n = \frac{t}{u_n} E_{12} + t^{1/n} E_{23} - \frac{1}{2} \frac{t^{1+1/n}}{u_n} E_{13},$$

so exponentiating its negative Laurent part alone gives the $(1, 3)$ coefficient $-\frac{1}{2}t^{1+1/n}/u_n$, whereas the correct polar coefficient in (8.3) is $-t^{1+1/n}/u_n$.

Proof. $G_n - I_3$ is nilpotent of order three, so the series of lemma 3.4 terminates: with $a = t/u_n$ and $c = t^{1/n}$ one has $(G_n - I_3)^2 = ac E_{13}$ and $(G_n - I_3)^3 = 0$, giving $\log G_n = (G_n - I_3) - \frac{1}{2}ac E_{13}$, which is the display. Comparison with (8.3) gives the factor $\frac{1}{2}$. The symbolic check is recorded in the companion code (section 29). $\square$

Thus even the true matrix logarithm is not a substitute for the factorization: one still needs the noncommutative correction, and the global criterion is built from the actual normalized factor rather than from an *ad hoc* linearization. Note also that the discrepancy is only a factor, not a change of exponent, so a reader who used the logarithm would find the same descending set here — and example 8.3 shows that agreement is an accident of this example, not a method.

9 Obstructions at arbitrary nilpotent depth

9.1 A chain identity

Lemma 9.1 (Chain identity). *Let $r \geq 3$, let a be a negative Laurent Hahn function with positive support, and let $c_2, \dots, c_{r-1}$ be positive Hahn constants. Put*

$$C = \sum_{j=2}^{r-1} c_j E_{j,j+1}, \quad H = I_r + C, \quad G = I_r + C + a E_{12}.$$

Since $C^{r-1} = 0$,

$$P = GH^{-1} = I_r + aE_{12}(I_r + C)^{-1} = I_r + \sum_{k=2}^r (-1)^{k-2} a \left(\prod_{j=2}^{k-1} c_j \right) E_{1k}, \quad (9.1)$$

the empty product for $k = 2$ being 1, and (9.1) is the normalized polar factorization $P = \text{Pol}(G)$, $H = \text{Reg}(G)$.

Proof. The finite geometric inverse of $I_r + C$ gives the displayed signs. Every nonconstant coefficient of P is a negative Laurent part, because each is a constant multiple of a , while H is holomorphic on the disk. Uniqueness in [theorem 5.1](#) identifies the factors. $\square$

9.2 The exponent design

For $r \geq 4$ and $n \geq 2$ set

$$a_n = \frac{t}{z-n}, \quad c_{2,n} = t^{1-1/n}, \quad c_{j,n} = t \quad (3 \leq j \leq r-2), \quad c_{r-1,n} = t^{2/n}, \quad (9.2)$$

the middle range being empty for $r = 4$. Let C_n and G_n be the matrices of [lemma 9.1](#) built from these data. Every G_n is finite, upper unitriangular, and has exactly one row singular matrix coefficient, namely $E_{12}/(z-n)$ at exponent 1. For $3 \leq k \leq r-1$ the exponent in $(P_n)_{1k}$ is

$$\lambda_{k,n} = k - 1 - \frac{1}{n}, \quad (9.3)$$

whereas the exponent in the final entry is

$$\lambda_{r,n} = r - 2 + \frac{1}{n}. \quad (9.4)$$

For each fixed $k < r$ the exponents (9.3) increase with n and their set has order type ω ; there are finitely many such sets, so their union with $\{1\}$ is well ordered. The final exponents (9.4) strictly decrease.

Theorem 9.2 (Arbitrarily deep central support obstructions). *For each integer $r \geq 3$ there is a positive upper-unitriangular cocycle on the puncture-and-disk cover of $\mathbb{C}$ with $A = \{2, 3, \dots\}$, over $\Gamma = \mathbb{Q}$, such that:*

- (i) *its original singular coefficient support is $\{1\}$ and its determinant is one;*
- (ii) *its image in the quotient of the upper-unitriangular group by $Z_r = \{I_r + cE_{1r} : c \in \mathcal{J}_\Gamma\}$ is a coboundary;*
- (iii) *it is not a coboundary in the full group of positive $r \times r$ matrices;*
- (iv) *it remains nonsplit after every ordered value-group extension.*

For $r = 3$ use (8.2); for $r \geq 4$ use (9.2) with [lemma 9.1](#).

Proof. Property (i) was checked directly above. [Lemma 9.1](#) and (9.4) show that the normalized polar support contains a strictly descending infinite sequence, so [theorem 6.2](#) proves (iii) and [corollary 7.3](#) proves (iv).

For (ii), let P'_n be obtained from P_n by deleting its $(1, r)$ entry. Their global support union is well ordered by (9.3), the case $r = 3$ leaving only exponent 1. Every $P'_n - I_r$ belongs to the row-one algebra

$$V = \text{span}\{E_{12}, \dots, E_{1,r-1}\}, \quad V^2 = 0,$$

so the gluing problem for the P'_n is additive and splits by the scalar construction with this one well-ordered support union: solve the ordinary principal-part problem for each coefficient of each allowed row entry, using [lemma 4.2](#). This gives F' on U_0 and Q'_n on D_n , valued in $I_r + V$, with $F' = P'_n Q'_n$. In the quotient by Z_r one has $\bar{P}_n = \bar{P}'_n$, and since $G_n = P_n H_n$ with H_n holomorphic,

$$\bar{G}_n = \bar{F}' (\bar{H}_n^{-1} \bar{Q}'_n)^{-1}$$

provides the required coboundary. All factors have individually legitimate positive supports. No matrix representative for the quotient is silently assumed: the bars denote the ordinary group quotient by the central subgroup. $\square$

Here Z_r is the central subgroup with $c \in \mathcal{J}_\Gamma$. Over $\Gamma = \mathbb{Q}$ one has $\mathcal{J}_\Gamma^m = \mathcal{J}_\Gamma$ for every positive integer m , since a nonzero section of least exponent $\delta > 0$ factors into $m - 1$ monomials $t^{\delta/m}$ and one remaining positive section. The elementary commutator identity $[I + aE_{ij}, I + bE_{jk}] = I + abE_{ik}$ then shows that Z_r is the final nonzero term of the lower central series of the positive upper-unitriangular group, and every further lower-central quotient factors through the quotient in (ii). So all those lower-depth tests pass, and the full obstruction first appears at depth $r - 1$, for arbitrarily large finite r .

Remark 9.3 (What the depth assertion means). The theorem does not say that a scalar equation cannot detect the final obstruction once a noncommutative normal form has been found — the final coefficient is of course scalar. It says that discarding the last central level *before* normalizing loses an obstruction created by the multiplication law. The new information sits in a genuinely higher noncommutative compatibility condition.

10 Integral vector bundles and local invisibility

10.1 From positive matrices to actual vector bundles

The transition matrices G_a define a locally free rank- r sheaf $\mathcal{E}_G$ over $\mathcal{J}_\Gamma$ on the ordinary surface X . Its ordinary reduction is canonically the trivial rank- r holomorphic bundle, because every transition reduces to I_r , and a positive splitting is a trivialization compatible with that reduced framing.

Proposition 10.1 (For triviality, the reduced framing costs nothing). *For a bundle defined by positive transition matrices the following are equivalent, on an arbitrary ordinary complex base and not only on a star cover:*

- (i) $\mathcal{E}_G$ is trivial as an $\mathcal{J}_\Gamma$ -module;
- (ii) its transition cocycle splits by positive matrices.

Proof. A positive splitting gives a trivialization. Conversely, suppose $A_i \in \mathrm{GL}_r(\mathcal{J}_\Gamma(U_i))$ satisfy $G_{ij} = A_i A_j^{-1}$ and reduce modulo $\mathcal{J}_\Gamma$. Since $\mathrm{red} G_{ij} = I_r$, the ordinary invertible matrices $\mathrm{red} A_i$ agree on overlaps and glue to $A \in \mathrm{GL}_r(\mathcal{O}(X))$. Right multiplication by the ordinary matrix A^{-1} produces $B_i = A_i A^{-1}$ with reduction I_r , hence $B_i \in \mathcal{G}_{r,\Gamma}(U_i)$ and $G_{ij} = B_i B_j^{-1}$. $\square$

Corollary 10.2 (Nontrivial determinant-one integral bundles). *The examples of [theorems 8.2](#) and [9.2](#) define nontrivial locally free $\mathcal{J}_\Gamma$ -modules even after forgetting the upper-triangular flag and the chosen reduced framing. Their ordinary reductions and determinant line bundles are trivial, the transition determinants being exactly one and not merely one modulo positive exponents.*

Proof. A general integral trivialization would give a positive matrix splitting by [proposition 10.1](#), contradicting [theorem 6.2](#). $\square$

10.2 Every bounded restriction is trivial

Corollary 10.3 (Exhaustion-invisible nontriviality). *Each integral bundle in [corollary 10.2](#) restricts trivially to every bounded ordinary domain in $\mathbb{C}$. In particular it is trivial on every disk of an increasing exhaustion of the plane, but not on the plane itself.*

Proof. A bounded domain meets only finitely many of the fixed disks D_n , so on the restricted cover there are only finitely many nonidentity transition matrices, each with legitimate positive support, and their union is well ordered. Every ordinary plane domain is Stein, so [theorem 11.1](#) applies. Equivalently, apply the finite-puncture construction directly and absorb transitions on pieces containing no puncture into holomorphic local gauges. $\square$

The failure is compatibility of infinitely many local trivializations, not existence on any one bounded domain. Consequently a numerical experiment involving finitely many punctures cannot establish global triviality: it will necessarily miss these examples. This is an explicit limit on what any implementation can decide, and it is recorded again as item 4 of [section 30](#).

10.3 The localization boundary

Let $\mathcal{H}_\Gamma = \mathcal{O}((t^\Gamma))$ be the sheaf in which arbitrary monomial shifts are allowed; the natural map $\mathcal{I}_\Gamma \rightarrow \mathcal{H}_\Gamma$ extends scalars from $\mathcal{E}_G$ to a rank- r $\mathcal{H}_\Gamma$ -module. We have *not* proved that the resulting module is nontrivial, and we decline the claim rather than gesture at it. The reduction argument of [proposition 10.1](#) cannot be repeated over $\mathcal{H}_\Gamma$: there is no reduction ring homomorphism $\mathcal{H}_\Gamma \rightarrow \mathcal{O}$ killing positive monomials, because those monomials are units in $\mathcal{H}_\Gamma$.

Accordingly the precise conclusions of Part I are nonexistence of a positive matrix splitting with common-domain Hahn entries and nontriviality of the corresponding *integral* locally free module. Whether negative-valuation changes of frame can trivialize particular higher-rank examples is a separate question. The scalar report proves more in rank one by a special leading-unit normalization [2]; no matrix analogue of that argument is assumed here.

Part III, added from a third source, sharpens this boundary without crossing it for these examples. It constructs *different* bundles, with ordinary essential rather than polar singularities in their transitions, that remain nontrivial over $\mathcal{H}_\Gamma$ and even over the coefficientwise meromorphic sheaf $\mathcal{M}_\Gamma = \text{Mer}((t^\Gamma))$ ([corollary 22.5](#)), and it shows that the examples of [corollary 10.2](#) become trivial over $\mathcal{M}_\Gamma$ ([corollary 27.2](#)). Part III leaves open whether they remain nontrivial over $\mathcal{H}_\Gamma$.

Part V, added from a fourth source, settles this: they do not. The hidden example becomes trivial over $\mathcal{I}_\mathbb{Q}[t^{-1}]$ by explicit determinant-one frames of valuation at least $-3/4$ ([theorem 35.1](#)), and the deep examples in the same way ([corollary 35.2](#)). The claim declined above was therefore rightly declined: these integral bundles are nontrivial over $\mathcal{I}_\Gamma$ and trivial over $\mathcal{H}_\Gamma$.

This distinction matters for the surcomplex interpretation. Integral models describe holomorphic matrices with a defined ordinary reduction and infinitesimal corrections. They are not the entire unrestricted matrix geometry over $\mathbf{SC}$, and our theorems are strong within the former category without conflating the two.

11 Positive rigidity on Stein manifolds

11.1 An arbitrary-cover theorem with a support hypothesis

Let X be a connected Stein manifold and $(U_i)_{i \in I}$ a set-indexed open cover; on disconnected intersections, support statements refer to all connected components. Suppose

$$G_{ij} \in \mathcal{G}_{r,\Gamma}(U_i \cap U_j), \quad G_{ij}G_{jk} = G_{ik}, \quad G_{ii} = I_r.$$

Call the cocycle *globally support-admissible* if

$$E = \bigcup_{i,j} \text{supp}(G_{ij} - I_r) \quad (11.1)$$

is well ordered. This is a sufficient condition *on a presentation*, not an invariant necessary condition: regular local gauges may destroy it without destroying triviality. It must not be conflated with the invariant $\mathcal{E}(G)$ of [definition 6.1](#), nor with E_ρ of [section 14](#).

Theorem 11.1 (Support-controlled Stein splitting). *A globally support-admissible positive matrix cocycle on a Stein manifold splits on its original cover: $G_{ij} = B_i B_j^{-1}$ with $B_i \in \mathcal{G}_{r,\Gamma}(U_i)$. All supports $\text{supp}(B_i - I_r)$ can be contained in the one monoid $\langle E \rangle_{\mathbb{N}} \setminus \{0\}$.*

Proof. Set $M = \langle E \rangle_{\mathbb{N}}$ and solve $G_{ij}B_j = B_i$ coefficientwise with $B_i = I_r + \sum_{\gamma \in M \setminus \{0\}} b_{i,\gamma} t^\gamma$. Assume the equations hold at all exponents below γ ; the coefficient equation at γ is

$$b_{i,\gamma} - b_{j,\gamma} = g_{ij,\gamma} + \sum_{\substack{\eta+\theta=\gamma \\ \eta,\theta>0}} g_{ij,\eta} b_{j,\theta} =: R_{ij,\gamma}, \quad (11.2)$$

a finite sum using only earlier coefficients. The cocycle identity $G_{ij}G_{jk} = G_{ik}$ together with the equations already solved below γ implies $R_{ij,\gamma} + R_{jk,\gamma} = R_{ik,\gamma}$ on triple overlaps; expand $G_{ij}(G_{jk}B_k - B_j) + (G_{ij}B_j - B_i) = G_{ik}B_k - B_i$ at exponent γ and note that all positive multipliers of a lower defect vanish by induction.

Thus $R_{ij,\gamma}$ is an ordinary additive cocycle with values in $\mathcal{O}^{r^2}$. Cartan's theorem B gives $H^1(X, \mathcal{O}^{r^2}) = 0$ [[18](#), [19](#)], so it has a coboundary $b_{i,\gamma}$ on the original cover; this is again the torsor interpretation of first cohomology and assumes nothing about coherence of the Hahn sheaf. Recursion on $M \setminus \{0\}$ gives the required matrices with a common support bound, and positive inversion proves they are invertible. No new ordinary domains and no new Hahn exponents are introduced. $\square$

11.2 Deformations of a nontrivial ordinary bundle

The principle is not limited to a trivial ordinary reduction. Let $\mathcal{V}$ be an ordinary holomorphic rank- r bundle on X with transition matrices g_{ij}^0 , and suppose Hahn transitions G_{ij} have reduction g_{ij}^0 and are invertible over $\mathcal{A}_\Gamma$. The relative transitions $G_{ij}(g_{ij}^0)^{-1} - I_r$ then have positive support on overlaps.

Theorem 11.2 (Support-controlled deformation rigidity). *If the union of these relative positive supports is well ordered and X is Stein, there exist positive local matrices B_i with*

$$G_{ij} = B_i g_{ij}^0 B_j^{-1}, \quad (11.3)$$

and the nonconstant supports of all B_i lie in the monoid generated by the given support union. Thus the integral Hahn deformation is isomorphic, by an isomorphism reducing to the identity, to the scalar extension of its ordinary bundle.

Proof. Proceed as in [theorem 11.1](#), now solving $G_{ij}B_j = B_i g_{ij}^0$. At a fixed positive exponent, multiply the residual equation on the right by $(g_{ij}^0)^{-1}$; the unknown coefficients appear as $b_{i,\gamma} - g_{ij}^0 b_{j,\gamma} (g_{ij}^0)^{-1}$. The residual is an additive cocycle for the ordinary endomorphism sheaf $\mathcal{E}nd(\mathcal{V})$, the cocycle identity following from $G_{ij}G_{jk} = G_{ik}$ and the lower solved equations exactly as before. That sheaf is locally free, hence coherent, and Cartan’s theorem B gives $H^1(X, \mathcal{E}nd(\mathcal{V})) = 0$. Solve the ordinary coefficient cocycle and continue the support-controlled recursion. All multiplications by g_{ij}^0 and its inverse occur at exponent zero and do not enlarge the exponent set. $\square$

This is a formal deformation-rigidity argument with a precise support certificate. It is not a new proof of the ordinary Oka–Grauert principle, and it is not an unconditional theorem about every Hahn-valued deformation on a noncompact base. Note that the cohomological input is acyclicity of $\mathcal{E}nd(\mathcal{V})$, not of $\mathcal{O}^{r^2}$: the theorem is about deformations of a genuinely nontrivial ordinary bundle.

11.3 A cyclic versus noncyclic dichotomy

Lemma 11.3 (Ordered-group alternative). *For a nonzero ordered abelian group Γ , the positive cone contains no infinite strictly decreasing sequence if and only if $\Gamma = \mathbb{Z}\delta$ for a least positive element δ .*

Proof. The cyclic case is immediate. Conversely, if there is no least positive element, choose a smaller positive element repeatedly. If there is a least positive δ but $\Gamma \neq \mathbb{Z}\delta$, choose a positive $\gamma \notin \mathbb{Z}\delta$; then $\gamma - n\delta > 0$ for every integer $n \geq 0$, since the first nonpositive difference would either leave an element strictly between 0 and δ or force $\gamma \in \mathbb{Z}\delta$. The sequence $\gamma, \gamma - \delta, \dots$ is strictly decreasing and positive. $\square$

Theorem 11.4 (Universal positive rigidity and its failure). *Fix a rank $r \geq 1$. Every positive rank- r cocycle on every Stein manifold is a coboundary if and only if*

$$\Gamma = \{0\} \quad \text{or} \quad \Gamma = \mathbb{Z}\delta \text{ for some } \delta > 0.$$

For every noncyclic nonzero Γ and every $r \geq 2$ there is a nonsplit determinant-one positive rank- r cocycle already on the puncture-and-disk cover of $\mathbb{C}$.

Proof. In the cyclic case every transition support is a subset of the universal well ordered positive set $\{\delta, 2\delta, \dots\}$, so its global union is automatically well ordered and [theorem 11.1](#) applies on any cover; the zero group gives only the identity cocycle.

For the converse choose a strictly decreasing positive sequence γ_n by [lemma 11.3](#) and set

$$g_n = \exp\left(\frac{t^{\gamma_n}}{z - n}\right).$$

This scalar matrix is its own normalized polar factor, every nonconstant coefficient being a multiple of a negative power of $z - n$, and its support union contains the descending set $\{\gamma_n\}$, so it is nonsplit. In rank $r \geq 2$ use

$$G_n = \text{diag}(g_n, g_n^{-1}, 1, \dots, 1). \tag{11.4}$$

Every nonconstant coefficient of this matrix is polar, so $P_n = G_n$; the same descending support prevents splitting in the full positive matrix group, while $\det G_n = 1$. For rank one the scalar example suffices. [Proposition 10.1](#) converts these examples into nontrivial integral vector bundles. $\square$

The rank-one part agrees with the Picard-group vanishing dichotomy of the repository’s global-divisor report [2], which vanishes exactly for $\Gamma = \{0\}$ or $\Gamma = \mathbb{Z}\delta$; that statement is the rank-one shadow of [theorem 11.4](#), driven by the same universal set $\{\delta, 2\delta, \dots\}$, and we cite rather than reprove it. The higher-rank conclusion here is stated for integral positive deformations, and the diagonal family (11.4) by itself supplies no nonabelian information: that is what [theorem 8.2](#) and [theorem 9.2](#) are for. Part III extends the dichotomy to $\mathcal{H}_\Gamma$ and to the meromorphic sheaf $\mathcal{M}_\Gamma$ ([theorem 25.1](#)); the diagonal family cannot serve there, since it becomes trivial over $\mathcal{M}_\Gamma$ ([corollary 27.2](#)).

12 The two well-ordered-support criteria are not the same criterion

Both headline theorems of this report read “the union of the supports is well ordered”. The unions concern different objects: full raw transition support gives only a sufficient gluing test, while raw meridian support gives the exact monodromy test.

	Criterion P (theorem 6.2)	Criterion M (theorem 14.2)
Problem	multiplicative gluing $G_a = B_0 B_a^{-1}$ on a puncture-and-disk cover	realization of a prescribed monodromy by a global positive connection
Tested object	$\text{Pol}(G_a) - I_r$, the normalized polar factor of an <i>independent local factorization</i> of each transition matrix	$\rho(\ell_d) - I_r$, the prescribed monodromy matrices themselves, for a compatible based-meridian basis
Tested union	$\mathcal{E}(G) = \bigcup_a \text{supp}(\text{Pol}(G_a) - I_r)$	$E_\rho = \bigcup_d \text{supp}(\rho(\ell_d) - I_r)$
Raw data	well-ordering of full raw transition support is sufficient, not necessary; raw singular support fails both directions (theorem 8.2 and example 8.3)	well-ordering of raw meridian support is the invariant condition, independent of basis and base point by proposition 14.4
Must one factor first?	Yes. Skipping the factorization tests an object that is provably inequivalent	No. The monodromy matrices are already the right object; a local factorization of $\rho(\ell_d)$ is not part of the criterion
Invariance	under arbitrary local holomorphic right gauges (corollary 7.1), coordinate change (corollary 7.2) and ordered value-group extension (corollary 7.3)	under change of free meridian basis and base point (proposition 14.4), one fixed $\text{GL}_r(K_\Gamma)$ conjugation and ordered value-group enlargement (proposition 14.5)

The trap is concrete, and it has already been printed. In [theorem 8.2](#) the raw singular support is the single exponent $\{1\}$ and no splitting exists; in [example 8.3](#), *adding* a term whose exponents form the descending set $\{1 + 1/n\}$ among the raw poles *removes* the obstruction. So for Criterion P the raw singular support is not the invariant, and the factorization must be performed first. Nothing of the kind happens for Criterion M: there the union is over the prescribed matrices, [proposition 14.4](#) shows the condition does not depend on the presentation at all, and [example 17.1](#) obstructs precisely because the prescribed matrices themselves carry $\{1/n\}$.

Two further consequences of the difference are worth stating so that neither criterion is quoted for the other’s conclusion. First, the objects tested live in different places: $\text{Pol}(G_a)$ is a matrix of negative Laurent tails in a local coordinate at a , whereas $\rho(\ell_d)$ is a matrix of Hahn constants. Second, the two criteria are not two instances of one general theorem proved here; they are two theorems with two proofs, sharing [lemmas 3.2](#) and [3.4](#) and an ordinary Mittag–Leffler input. This report does not claim a common generalization, and [section 31](#) lists the obvious attempt as open.

Part II. Inverse monodromy: the meridian-support criterion

13 Connections, fundamental matrices, and holonomy

A positive connection on the trivial rank- r bundle over an ordinary domain U means

$$dY = \Omega Y, \quad \Omega = \sum_{\gamma \in S} t^\gamma \Omega_\gamma, \quad S \subseteq \Gamma_{>0} \text{ well ordered}, \quad (13.1)$$

where each Ω_γ is an ordinary holomorphic matrix one-form on *all* of U . In complex dimension one such connections are automatically flat: both the exterior derivative of a holomorphic one-form and the wedge of two of them vanish. As always, “positive” concerns exponents only.

Let $\tilde{U} \rightarrow U$ be the ordinary universal covering with a chosen lift $\tilde{p}$ of the base point p ; pullbacks of coefficients are understood.

Theorem 13.1 (Fundamental matrix with a support certificate). *For a connection (13.1) there is a unique positive fundamental matrix*

$$Y = I_r + \sum_{\gamma > 0} t^\gamma Y_\gamma \text{ on } \tilde{U}, \quad dY = \Omega Y, \quad Y(\tilde{p}) = I_r.$$

Every coefficient is holomorphic on all of $\tilde{U}$ and

$$\text{supp}(Y - I_r) \cup \text{supp}(Y^{-1} - I_r) \subseteq \langle S \rangle_{\mathbb{N}} \setminus \{0\}.$$

Analytic continuation determines a representation $\text{Hol}_\Omega : \pi_1(U, p) \rightarrow I_r + \text{Mat}_r(\mathfrak{m}_\Gamma)$ with the same uniform support bound for every group element.

Proof. Write $T = \langle S \rangle_{\mathbb{N}} \setminus \{0\}$ and define the coefficients by well-founded recursion on T , missing coefficients of Ω being zero. At weight γ set

$$dY_\gamma = \Omega_\gamma + \sum_{\substack{\alpha + \beta = \gamma \\ \alpha, \beta > 0}} \Omega_\alpha Y_\beta, \quad Y_\gamma(\tilde{p}) = 0. \quad (13.2)$$

Only finitely many pairs contribute by [lemma 3.2](#), and $\beta < \gamma$. The right-hand side is a holomorphic matrix one-form on the simply connected Riemann surface $\tilde{U}$, so its entries have unique holomorphic primitives vanishing at $\tilde{p}$; this constructs Y_γ on the whole cover. The family has support in the well-ordered set T and satisfies the equation coefficientwise, and [lemma 3.4](#) gives its inverse with the same monoid support.

For uniqueness even against a solution with a different positive well-ordered support, take the least exponent at which two solutions differ, in the well-ordered union of their supports: at that exponent the difference has zero derivative and zero value at $\tilde{p}$, hence is zero, a contradiction.

A continued fundamental matrix differs from the original branch on a connected overlap by a constant matrix on the right, since $d(Y^{-1}\hat{Y}) = 0$ follows coefficientwise from the equation. Continuing around a based loop gives the normalized monodromy matrix, whose coefficients are values of coefficients of Y ; therefore all its positive exponents lie in T , independently of the loop. Uniqueness of continuation gives the group law with the convention of [lemma 4.4](#), and homotopy invariance follows from the universal-cover construction. $\square$

This theorem is the precise form of the *forward* map that the repository’s differential-equations report constructs from a coherent connection, recording that the exponent-zero coefficient is the ordinary monodromy matrix and that all other exponents lie in the generated monoid [3]. Here the exponent-zero coefficient is I_r by positivity. Theorem 14.2 is the converse.

13.1 The explicit iterated-integral expression

For a piecewise smooth path $c : [0, 1] \rightarrow U$ write $c^*\Omega_\gamma = A_\gamma(s) ds$. Its transport is

$$T_\Omega(c) = I_r + \sum_{n \geq 1} \sum_{\gamma_1, \dots, \gamma_n \in S} t^{\gamma_1 + \dots + \gamma_n} \int_{0 < s_1 < \dots < s_n < 1} A_{\gamma_n}(s_n) \cdots A_{\gamma_1}(s_1) ds_1 \cdots ds_n. \quad (13.3)$$

This is the usual ordered-integral solution, a classical mechanism related to iterated path integrals [24]; the name “Dyson expansion” is used for it purely as classical mathematics, with no physical reading attached. Its meaning here is strict: the outer family is strongly summable by the finite-word clause of lemma 3.2, and each individual coefficient is a *finite* sum of ordinary iterated integrals. The expression agrees with theorem 13.1, since termwise differentiation gives (13.2); for a loop it gives Hol_Ω .

Lemma 13.2 (Triangular change of monodromy). *Suppose two positive connections Ω and Ψ agree at every exponent strictly below γ . Then their transports along any fixed path agree below γ , and for a based loop ℓ ,*

$$[t^\gamma](\text{Hol}_\Psi(\ell) - \text{Hol}_\Omega(\ell)) = \int_\ell (\Psi_\gamma - \Omega_\gamma). \quad (13.4)$$

Terms strictly above γ cannot affect coefficients at or below γ .

Proof. Compare the word expansions (13.3) over the well-ordered union of the two positive supports. A differing coefficient of weight at least γ cannot occur in a word of total weight less than γ ; at weight exactly γ it can occur only as the single letter of the word, every additional letter having strictly positive weight. The length-one term is the displayed ordinary period. All comparisons are finite at each exponent. $\square$

The crucial feature is that the linearized map at *every* successive weight is the same ordinary period map P of (4.2). Noncommutativity appears in the already determined lower-weight terms; it does not change the operator needed to solve the new correction equation.

14 The exact realization theorem

Definition 14.1 (Uniform support admissibility and σ -normalization). For $\rho : \pi_1(U, p) \rightarrow I_r + \text{Mat}_r(\mathfrak{m}_\Gamma)$ with $U = \mathbb{C} \setminus D$ and $M_d = \rho(\ell_d)$, put

$$E_\rho = \bigcup_{d \in D} \text{supp}(M_d - I_r), \quad (14.1)$$

the union being over all matrix entries and all based meridians of the fixed free basis of lemma 4.4. The representation is *uniformly support-admissible* when E_ρ is well ordered. For a fixed period section σ of lemma 4.5, a connection is σ -normalized if every entry of every coefficient lies in $\text{im } \sigma \subseteq \mathcal{L}(D)$.

Theorem 14.2 (Support-sensitive Riemann–Hilbert realization: Criterion M). *Let $D \subset \mathbb{C}$ be closed and discrete, $p \in U = \mathbb{C} \setminus D$, with free based meridians fixed. For $\rho : \pi_1(U, p) \rightarrow I_r + \text{Mat}_r(\mathfrak{m}_\Gamma)$ the following are equivalent:*

- (i) E_ρ is well ordered;
- (ii) ρ is the monodromy of a positive globally common-domain Hahn connection on U ;
- (iii) ρ is the monodromy of such a connection whose coefficients are meromorphic on $\mathbb{C}$ with at most simple poles in D and no other finite poles.

If these hold then for every fixed complex-linear period section σ there is exactly one σ -normalized realizer Ω^σ , and

$$\text{supp } \Omega^\sigma \subseteq \langle E_\rho \rangle_{\mathbb{N}} \setminus \{0\}. \quad (14.2)$$

Uniqueness is among all positive globally supported σ -normalized connections, not only those whose support was assumed in advance to lie in the right-hand side of (14.2).

Proof. Necessity. If $\rho = \text{Hol}_\Omega$ with $\text{supp } \Omega = S$ then [theorem 13.1](#) gives $\text{supp}(\rho(g) - I_r) \subseteq \langle S \rangle_{\mathbb{N}} \setminus \{0\}$ for every group element g , so E_ρ is a subset of one well-ordered set. This proves (ii) $\Rightarrow$ (i), and (iii) $\Rightarrow$ (ii) is immediate.

Construction. Assume (i), put $T = \langle E_\rho \rangle_{\mathbb{N}} \setminus \{0\}$, and write $M_d = I_r + \sum_{\gamma \in T} t^\gamma m_{d,\gamma}$, missing coefficients being zero. We construct $\Omega_\gamma \in \text{Mat}_r(\mathcal{L}(D))$ for $\gamma \in T$ by recursion. Suppose all lower coefficients have been constructed and form the legitimate positive connection $\Omega^{<\gamma} = \sum_{\delta \in T, \delta < \gamma} t^\delta \Omega_\delta$, whose support is a well-ordered subset of T ; the initial segment need not be finite. Define the lower-order prediction and the remaining defect by

$$h_{d,\gamma}^{<\gamma} = [t^\gamma] \text{Hol}_{\Omega^{<\gamma}}(\ell_d), \quad b_{d,\gamma} = m_{d,\gamma} - h_{d,\gamma}^{<\gamma}. \quad (14.3)$$

The prediction exists by [theorem 13.1](#); equivalently it is the finite sum of all iterated-integral words of total weight γ formed from previously constructed coefficients. Apply the period section entrywise and set

$$\Omega_\gamma = \sigma((b_{d,\gamma})_{d \in D}). \quad (14.4)$$

The datum at one exponent is an arbitrary matrix-valued element of the product $\mathbb{C}^D$, and [lemma 4.5](#) applies even when infinitely many of its entries are nonzero; this is why no summability across punctures is required.

The coefficient Ω_γ is meromorphic on $\mathbb{C}$ with at most simple poles in D . It corrects exactly the defect at weight γ , since $\int_{\ell_d} \Omega_\gamma = b_{d,\gamma}$ and [lemma 13.2](#) applies, and it leaves all lower weights unchanged. Transfinite recursion on the well-ordered set T therefore constructs $\Omega^\sigma = \sum_{\gamma \in T} t^\gamma \Omega_\gamma$, all of whose coefficients are holomorphic on the same domain U and whose support is a subset of T . No limiting assertion at a limit ordinal is involved: a coefficient is simply assigned to every member of a well-ordered set.

To verify the monodromy, fix $\gamma \in T$. Coefficients at weights above γ cannot affect it by [lemma 13.2](#), and the recursion has already set it equal to $m_{d,\gamma}$. The monodromy support of the final connection lies in $\langle T \rangle_{\mathbb{N}} \setminus \{0\} = T$, so no exponent outside T appears. Thus every meridian has its prescribed monodromy, and the meridians freely generate the fundamental group by [lemma 4.4](#), proving $\text{Hol}_{\Omega^\sigma} = \rho$ and (iii).

Uniqueness. Let Ω and Ψ be two positive σ -normalized realizers. If they differ, the union of their supports is well ordered and there is a least differing exponent γ . Their holonomies are equal, so [lemma 13.2](#) gives $P(\Psi_\gamma - \Omega_\gamma) = 0$. But $\Psi_\gamma - \Omega_\gamma$ lies entrywise in $\text{im } \sigma$, and $P\sigma = \text{id}$ says that P is injective there. Hence the difference is zero, a contradiction. $\square$

Remark 14.3 (No hidden convergence over the puncture set). The construction is not a sum of one independently built differential equation per puncture. At a fixed Hahn exponent it first collapses *all* puncture data into one ordinary meromorphic one-form using the period section, and only then are these ordinary forms assembled in Hahn exponents. This move is indispensable when infinitely many punctures use the same exponent; see [example 17.2](#).

14.1 An intrinsic version of the support condition

Proposition 14.4 (Independence of the meridian basis). *For a near-identity representation ρ put $E_{\text{all}}(\rho) = \bigcup_{g \in \pi_1(U, p)} \text{supp}(\rho(g) - I_r)$. Then E_ρ is well ordered if and only if $E_{\text{all}}(\rho)$ is well ordered. Hence admissibility does not depend on the chosen free meridians or, after a base-point identification, on the base point.*

Proof. One implication follows from $E_\rho \subseteq E_{\text{all}}(\rho)$. For the other, assume E_ρ well ordered. Each group element is a finite word in the M_d and their inverses; [lemma 3.4](#) expands the inverses in the common monoid $\langle E_\rho \rangle_{\mathbb{N}}$, and multiplication preserves that monoid, so $E_{\text{all}}(\rho) \subseteq \langle E_\rho \rangle_{\mathbb{N}} \setminus \{0\}$. Changing based meridians, or identifying fundamental groups at different base points, changes the presentation of the same representation and not this condition on all its values. $\square$

The support set E_ρ itself can change with the meridian basis; its *well-ordering* is the invariant proved here. Contrast [section 12](#): Criterion P is *not* a condition on the raw transition data, and the analogue of [proposition 14.4](#) for $\bigcup_a \text{supp}(G_a - I_r)$ is false.

Proposition 14.5 (Fixed conjugation and enlargement cannot repair support). *For a family (M_d) over K_Γ , well-ordering of $\bigcup_d \text{supp}(M_d - I_r)$ is invariant under conjugation by any one fixed $C \in \text{GL}_r(K_\Gamma)$, and is unchanged when the family is embedded by an order-preserving value-group embedding into a larger Hahn workspace. An arbitrary conjugation need not preserve near-identity positivity; the assertion about well-ordered supports does not require that property.*

Proof. Let A and B be the supports of the finitely many entries of C and C^{-1} ; they are well ordered. If E is the union of the supports of $M_d - I_r$ then $\bigcup_d \text{supp}(C(M_d - I_r)C^{-1}) \subseteq A + E + B$, which is well ordered by the last clause of [lemma 3.2](#) when E is. The converse follows by conjugating with C^{-1} . An order embedding preserves and reflects the ordering of a subset, hence preserves and reflects its being well ordered. $\square$

[Proposition 14.5](#) for Criterion M and [corollary 7.3](#) for Criterion P are two independent proofs, for two different objects, of the same moral: enlarging the surreal workspace never repairs these obstructions.

15 Framed gauge classification and removable gauges

Definition 15.1 (Positive framed gauge). A positive gauge on U is a matrix $G \in \mathcal{G}_{r, \Gamma}(U)$. It is *framed* at p when $G(p) = I_r$ as an equality of Hahn matrices. Its inverse is a positive globally supported holomorphic matrix by [lemma 3.4](#), and it transforms a connection by

$$\Psi = dG G^{-1} + G \Omega G^{-1}. \quad (15.1)$$

Theorem 15.2 (Framed classification with controlled support). *Two positive globally Hahn-supported connections Ω and Ψ on U have the same based monodromy if and only if there is a positive framed gauge G satisfying (15.1). Such a gauge is unique, and if $E = \text{supp } \Omega \cup \text{supp } \Psi$ then*

$$\text{supp}(G - I_r) \cup \text{supp}(G^{-1} - I_r) \subseteq \langle E \rangle_{\mathbb{N}} \setminus \{0\}. \quad (15.2)$$

Consequently holonomy is a bijection between framed positive gauge classes and uniformly support-admissible near-identity representations.

Proof. Let Y_Ω, Y_Ψ be the normalized fundamental matrices on $\tilde{U}$. If the based monodromies agree, form $\tilde{G} = Y_\Psi Y_\Omega^{-1}$. Upon continuation around any loop both fundamental matrices acquire the same right factor M , so the product becomes $Y_\Psi M(Y_\Omega M)^{-1} = Y_\Psi Y_\Omega^{-1}$. Each ordinary coefficient descends to U , and the whole family already has support in $\langle E \rangle_{\mathbb{N}}$ by [theorem 13.1](#) and [lemma 3.4](#). Thus the descended matrix is a globally supported positive gauge G with $G(p) = I_r$, and differentiation gives $dG = \Psi G - G\Omega$, equivalent to (15.1).

Conversely, if a framed gauge exists then GY_Ω solves the Ψ equation with initial value I_r , so uniqueness in [theorem 13.1](#) gives $GY_\Omega = Y_\Psi$. Since G is single valued on U the monodromy matrices agree, and the equality also forces $G = Y_\Psi Y_\Omega^{-1}$, proving uniqueness and the support bound. The bijection now follows from [theorem 14.2](#). $\square$

The logarithmic subclass has a stronger rigidity property: although the comparison gauge is initially defined only on the punctured plane, no coefficient of it actually needs a puncture.

Theorem 15.3 (Entire extension of a logarithmic comparison gauge). *Suppose Ω and Ψ are as in [theorem 15.2](#), have the same based monodromy, and every coefficient of both lies in $\text{Mat}_r(\mathcal{L}(D))$. Then the unique framed gauge and its inverse extend coefficientwise holomorphically to all of $\mathbb{C}$; in particular $G, G^{-1} \in \mathcal{G}_{r,\Gamma}(\mathbb{C})$, with the same support bound (15.2).*

Proof. Write $G = I_r + \sum_{\gamma > 0} t^\gamma G_\gamma$. The gauge equation at weight γ reads

$$dG_\gamma = \Psi_\gamma - \Omega_\gamma + \sum_{\substack{\alpha+\beta=\gamma \\ \alpha, \beta > 0}} (\Psi_\alpha G_\beta - G_\beta \Omega_\alpha). \quad (15.3)$$

Induct on the positive support monoid in (15.2). If every lower G_β is entire, the right-hand side is an ordinary meromorphic one-form on $\mathbb{C}$ with at most simple poles in D , the sum being finite. Fix $d \in D$ and a disk containing no other point of D . The coefficient G_γ is already single valued and holomorphic on the punctured disk, so the integral of its derivative around a small circle vanishes; hence the right-hand side of (15.3) has residue zero at d , and having at most a simple pole it is holomorphic there. Its primitive on the full disk differs from G_γ by a constant on the punctured disk, so G_γ extends holomorphically across d . Doing this for all d extends G_γ to an entire function, with no uniform bound on the sizes of coefficient functions required, and the ordinary domain after extension is the same $\mathbb{C}$ for every coefficient. Finally the geometric inverse of a positive entire family has entire coefficients and the same monoid support. $\square$

Corollary 15.4 (A global normal slice for logarithmic connections). *For a fixed σ , every positive logarithmic connection has exactly one σ -normalized representative under framed positive entire gauges, namely the connection constructed from its monodromy by (14.4).*

Proof. Its monodromy is admissible by [theorem 13.1](#); [theorems 14.2](#) and [15.3](#) give existence, and uniqueness of the normalized connection and of the framed gauge has already been proved. $\square$

Example 15.5 (A nonlogarithmic connection with trivial monodromy). On $\mathbb{C}^*$ consider the scalar connection $\Omega = -t dz/z^2$ with base point $p = 1$. Its normalized solution is $Y = \exp(t(1/z - 1))$, a positive globally supported holomorphic family on $\mathbb{C}^*$, single valued, so the monodromy is trivial. The unique framed gauge from Ω to the zero connection is $G = Y^{-1} = \exp(-t(1/z - 1))$, whose coefficient at t has a pole at 0. Therefore [theorem 15.3](#) cannot be extended to connections with arbitrary higher-order coefficient poles.

16 Finite-puncture inversion and determinant one

Corollary 16.1 (Unique positive residue matrices for finite data). *Let $D = \{a_1, \dots, a_m\}$ be finite, fix based meridians, and let $M_j \in I_r + \text{Mat}_r(\mathfrak{m}_\Gamma)$ be arbitrary. There are unique residue matrices $R_j \in \text{Mat}_r(\mathfrak{m}_\Gamma)$ for which*

$$\Omega = \sum_{j=1}^m R_j \frac{dz}{z - a_j} \quad (16.1)$$

has based meridian monodromies M_j . If $E = \bigcup_j \text{supp}(M_j - I_r)$ then $\text{supp } R_j \subseteq \langle E \rangle_{\mathbb{N}} \setminus \{0\}$. The form has at most a simple pole at infinity, with residue $-\sum_j R_j$ there.

Proof. A finite union of well-ordered sets in a linear order is well ordered, so E is admissible. Use the explicit section (4.3) in [theorem 14.2](#): being normalized for that section is precisely the form (16.1), and coefficientwise uniqueness gives uniqueness of all the R_j . In the coordinate $w = 1/z$ one has $dz/(z - a_j) = -dw/(w(1 - a_j w))$, proving the assertion at infinity. $\square$

This is a near-identity Hahn version of a classical Fuchsian inverse problem, not a realization result for arbitrary complex monodromy on a globally trivial bundle. The residue matrices must have positive support, a regular singularity at infinity is permitted, no assertion is made that the connection is holomorphic at infinity, and uniqueness is not claimed among arbitrary residue matrices with the same monodromy: ordinary integer shifts of exponents, which already destroy such unrestricted uniqueness, are excluded by the positive-support hypothesis.

Lemma 16.2 (Determinants and positive connections). *For the normalized fundamental matrix Y of a positive connection, $d(\det Y) = (\text{tr } \Omega) \det Y$. Consequently a traceless positive connection has determinant-one transport along every path.*

Proof. The determinant is a finite polynomial in the entries of Y , so the adjugate identity and the Leibniz rule apply coefficientwise, and Y is invertible by [theorem 13.1](#). Jacobi's formula reduces to the displayed identity. If $\text{tr } \Omega = 0$ the determinant is constant along the normalized solution and equals 1 at the starting point. All identities involve finite coefficientwise algebra or already justified support products. $\square$

Theorem 16.3 (Traceless normalized realization). *Suppose an admissible representation in [theorem 14.2](#) takes values in $\text{SL}_r(K_\Gamma)$. For an entrywise complex-linear period section its unique normalized realizer satisfies $\text{tr } \Omega^\sigma = 0$. Conversely every traceless positive connection has monodromy in $\text{SL}_r(K_\Gamma)$, and the unique framed gauge between two traceless connections with the same monodromy has determinant one. These statements retain all the support and logarithmic-extension conclusions above.*

Proof. Use the recursion (14.4) and assume every lower coefficient Ω_δ traceless. Then $\Omega^{<\gamma}$ is traceless, so its monodromy N_d has determinant one by lemma 16.2; by construction N_d and the target M_d agree below γ , and both have constant term I_r . In the determinant polynomial a difference at weight γ can contribute to that weight only with all other factors of weight zero, so

$$[t^\gamma](\det M_d - \det N_d) = \text{tr}(m_{d,\gamma} - h_{d,\gamma}^{<\gamma}).$$

The left side is zero, hence the defect vector in (14.3) is traceless at every puncture. Since the same linear section is used for each matrix entry, $\text{tr } \Omega_\gamma = \sigma((\text{tr } b_{d,\gamma})_{d \in D}) = 0$, proving the induction. The converse is lemma 16.2. For a gauge satisfying $dG = \Psi G - G\Omega$, Jacobi’s formula gives $d \det G = (\text{tr } \Psi - \text{tr } \Omega) \det G = 0$, and its value at p is 1. $\square$

17 Sharp global obstructions and calibrating examples

17.1 Finite solvability does not imply global solvability

Example 17.1 (A countable special-linear obstruction). Let $D = \{1, 2, 3, \dots\}$, $\Gamma = \mathbb{Q}$, and

$$N = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad M_n = I_2 + t^{1/n} N.$$

Because the meridians form a free basis, this prescription defines a representation $\rho : \pi_1(\mathbb{C} \setminus D, p) \rightarrow \text{SL}_2(K_\Gamma)$. Each generator matrix has one positive exponent, is invertible, and is an entirely legitimate Hahn matrix, and every finite initial collection of the prescribed monodromies is realizable on the corresponding finitely punctured plane by corollary 16.1. Nevertheless no positive globally Hahn-supported holomorphic connection on $\mathbb{C} \setminus D$ has all the prescribed monodromies.

Proof. The determinant is 1 because $N^2 = 0$ and N is strictly upper triangular. A finite set of the exponents $1/n$ is well ordered, but the full union $E_\rho = \{1, 1/2, 1/3, \dots\}$ is a strictly decreasing sequence with no least member, hence not well ordered. Nonexistence follows from theorem 14.2, even without any requirement of simple poles, and proposition 14.5 shows that a fixed change of matrix basis or an enlargement of the value group cannot repair it. $\square$

One-puncture realizers can even be written down:

$$\Omega_n = \frac{t^{1/n}}{2\pi i} N \frac{dz}{z - n}, \quad \exp\left(\int_{\ell_n} \Omega_n\right) = I_2 + t^{1/n} N.$$

The failure is not an inability to solve a differential equation at any one puncture; it is the impossibility of placing the entire prescription in one global support set. This is a genuine local-to-global obstruction in the chosen coefficient category, not a claim that no ordinary local system exists: the representation itself already defines an algebraic local system over K_Γ .

17.2 The union condition must not be strengthened to point-finiteness

Example 17.2 (Infinitely many punctures at one exponent). Let $D = \mathbb{Z}$, $U = \mathbb{C} \setminus \mathbb{Z}$, $\Gamma = \mathbb{Q}$, and use the same nilpotent N . Then

$$\Omega = \frac{t}{2i} N \cot(\pi z) dz \tag{17.1}$$

has monodromy $I_2 + tN$ at every positive based meridian. The family $(M_n - I_2)_{n \in \mathbb{Z}}$ is *not* strongly summable as a family indexed by punctures, because exponent 1 appears infinitely many times; it is nevertheless globally realizable.

Proof. The ordinary form $\cot(\pi z) dz$ has residue $1/\pi$ at every integer, so each meridian period of (17.1) is tN . Every product of two coefficient matrices vanishes, both being multiples of N with $N^2 = 0$, so the Dyson expansion (13.3) terminates after its length-one term and gives $I_2 + tN$. The connection has the singleton support $\{1\}$ and is holomorphic on one common domain U . Nothing in the definition asks that its periods over *different* loops form a summable family. $\square$

Together, examples 17.1 and 17.2 identify the precise strength of Criterion M. Requiring only that individual matrices be legitimate Hahn matrices is too weak; requiring a strongly summable family over punctures is too strong; a well-ordered *union* of supports is exactly right.

17.3 Admissible supports need not be locally finite in an interval

Example 17.3 (Infinitely many exponents below one bound). On $D = \{1, 2, \dots\}$ prescribe $M_n = I_2 + t^{2-1/n}N$. The support $E = \{2 - 1/n : n \geq 1\}$ is increasing of order type ω , hence well ordered, although it has infinitely many elements below 2. These data are realizable: for a fixed period section put $\alpha_n = \sigma(e_n)$ and

$$\Omega = N \sum_{n \geq 1} t^{2-1/n} \alpha_n,$$

where $e_n \in \mathbb{C}^D$ is 1 at n and 0 elsewhere. The connection has support in E , and its holonomy is exactly the prescribed family because $N^2 = 0$.

The displayed sum is a Hahn family, not an assertion that its ordinary partial sums converge in the valuation topology. In $\mathbb{Q}$ the tail valuations $2 - 1/n$ stay below 2, so those partial sums do not even approach the full family in that topology. The example also shows why our theorems supply no procedure enumerating all coefficients below a fixed bound in finitely many steps.

18 A noncommuting calculation across two valuation ranks

18.1 The targets and the first-order guess

Take $U = \mathbb{C} \setminus \{0, 1\}$ and base point $b = 1/2$, with the usual positive based meridians: a real tail to a small circle about the puncture, a positive circuit, and the reversed tail. Define

$$N_+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad N_- = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad H = [N_+, N_-] = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

so $N_+^2 = N_-^2 = 0$. Initially use two independent formal bookkeeping variables u, v of positive degree and prescribe

$$M_0 = I_2 + uN_+, \quad M_1 = I_2 + vN_-. \quad (18.1)$$

At the end set $u = t^\alpha$ and $v = t^\beta$ for positive, rationally independent $\alpha, \beta \in \Gamma$; the choice $\alpha = 1$, $\beta = \omega$ is the rank-two surcomplex example. Put

$$\eta_0 = \frac{dz}{2\pi iz}, \quad \eta_1 = \frac{dz}{2\pi i(z-1)}, \quad C = \frac{\log 2}{2\pi i}.$$

The obvious first-order guess is

$$\Omega^{(1)} = uN_+\eta_0 + vN_-\eta_1. \quad (18.2)$$

Its first-order periods are correct. Its mixed-order monodromy is not.

Proposition 18.1 (The first nonabelian correction). *For the connection (18.2) the coefficient of uv in the based monodromy is CH at the meridian of 0 and $-CH$ at the meridian of 1. Consequently the unique normalized realizer has expansion*

$$\Omega = uN_+\eta_0 + vN_-\eta_1 - uvCH(\eta_0 - \eta_1) + O_{\deg \geq 3}, \quad (18.3)$$

and its coefficients of u^2 and v^2 are zero. The remainder notation refers only to total degree in u, v , not to a claimed valuation bound.

Proof. Along a chosen meridian let $L_j(s) = \int_0^s \eta_j$ be the accumulated integral from the base point. The first-order transport is $I_2 + uN_+L_0 + vN_-L_1$, so the mixed term in the coefficient equation is

$$N_+N_- \int_\ell L_1\eta_0 + N_-N_+ \int_\ell L_0\eta_1. \quad (18.4)$$

For the meridian of 0, L_1 is the single-valued ordinary function $L_1(z) = (\log(1-z) - \log(1-b))/2\pi i$ on a neighbourhood containing the tail and the disk about 0 but not 1; it satisfies $L_1(b) = 0$ and $L_1(0) = C$. Integrating $d(L_0L_1)$ along the full loop gives $\int_\ell L_0\eta_1 = -\int_\ell L_1\eta_0$, because the product vanishes at both endpoints, L_0 ending at 1 but L_1 ending at 0. The residue theorem gives $\int_\ell L_1\eta_0 = L_1(0) = C$, contributions of the outgoing and incoming tails cancelling. Thus (18.4) equals $C(N_+N_- - N_-N_+) = CH$.

For the meridian of 1 use instead $L_0(z) = (\log z - \log b)/2\pi i$, which is single valued near that tail and disk with $L_0(b) = 0$ and $L_0(1) = C$; now $\int_\ell L_0\eta_1 = C$ and $\int_\ell L_1\eta_0 = -C$, so the mixed coefficient is $-CH$. The prescribed matrices (18.1) have zero uv coefficient, so the ordinary period section for $\{0, 1\}$ assigns the correcting one-form $-CH\eta_0 + CH\eta_1$ at that degree, which is (18.3). Pure second-order terms in the uncorrected transport vanish because $N_+^2 = N_-^2 = 0$, so their defects and normalized correction coefficients are zero. $\square$

In residue form the same calculation reads

$$R_0 = \frac{u}{2\pi i} N_+ - \frac{uv \log 2}{(2\pi i)^2} H + O_{\deg \geq 3}, \quad (18.5)$$

$$R_1 = \frac{v}{2\pi i} N_- + \frac{uv \log 2}{(2\pi i)^2} H + O_{\deg \geq 3}. \quad (18.6)$$

The mixed corrections cancel in the residue at infinity but are essential at the two finite poles. All displayed coefficient matrices are traceless, in agreement with [theorem 16.3](#).

18.2 The representation is genuinely nonabelian

The exact group commutator of the prescribed matrices is

$$M_0M_1M_0^{-1}M_1^{-1} = \begin{pmatrix} 1 + uv + u^2v^2 & -u^2v \\ uv^2 & 1 - uv \end{pmatrix}, \quad (18.7)$$

with determinant 1 and leading mixed term $uvH \neq 0$. So the realization theorem here is not an entrywise construction of independent scalar equations: the extra commutator in (18.3) is exactly

the interaction that a scalar period prescription would miss. This is the Part II counterpart of the cross term $-a_-c_+$ of [proposition 8.1](#): both constructions require mixed products absent from the abelianized equations. Here the mixed term is a solvable correction; in Part I such products can also create a globally inadmissible polar-support union.

18.3 Specialization to a genuinely higher-rank Hahn group

Set $\Gamma = \mathbb{Q} + \mathbb{Q}\omega$, $u = t$ and $v = t^\omega$. The monoid generated by 1 and ω is $\{a + b\omega : a, b \in \mathbb{N} \cup \{0\}\}$, ordered first by b and then by a , and it is well ordered. The correction in [\(18.3\)](#) occurs at exponent $1 + \omega$. No finite number of increments by 1 reaches even ω , but the well-founded recursion handles ω and $1 + \omega$ directly, with no change in the proof or in its support certificate; compare [remark 3.3](#).

The notation $O_{\deg \geq 3}$ is especially important here. A term t^3 has total degree three and yet smaller valuation than t^ω , while $t^{3\omega}$ has much larger valuation. A finite bivariate degree truncation is therefore *not* an approximation uniformly valid below a single Hahn valuation threshold. The accompanying computation checks a finite coefficient identity, not a truncation of all coefficients below $1 + \omega$.

Part III. Beyond the localization boundary: essential-singularity bundles over $\mathcal{H}_\Gamma$ and $\text{Mer}((t^\Gamma))$

19 What survives monomial localization

19.1 The question inherited from Part I

Part I proves nontriviality of its bundles over the integral sheaf $\mathcal{I}_\Gamma$ only. [Section 10.3](#) explains why its argument stops there: positive monomials are units in $\mathcal{H}_\Gamma$, there is no reduction homomorphism $\mathcal{H}_\Gamma \rightarrow \mathcal{O}$, and the framing normalization of [proposition 10.1](#) cannot be repeated. Item 2 of [section 30](#) declines the stronger claim, and [section 31](#) asks two questions: whether the hidden and deep examples of [theorems 8.2](#) and [9.2](#) remain nontrivial after monomial localization, and how isomorphism classes, not merely triviality, could be classified.

This part is built from a third, independently written manuscript, called *the localization manuscript* below ([appendix C](#)). It answers two *broad*er questions and leaves the printed one open.

1. *Existence*. There are finite-rank bundles on the ordinary plane, with trivial ordinary reduction and trivial determinant, that remain nontrivial — indeed indecomposable and not stably trivial — after every Hahn monomial is inverted, and even after holomorphic coefficients are replaced by meromorphic ones. Every bounded restriction of these bundles is already trivial over $\mathcal{I}_\Gamma$.
2. *Classification*. Within the resulting family the isomorphism classes are determined exactly, over $\mathcal{I}_\Gamma$, over $\mathcal{H}_\Gamma$, and over the coefficientwise meromorphic sheaf $\mathcal{M}_\Gamma = \text{Mer}((t^\Gamma))$ of [definition 20.1](#).

The examples are *not* those of Part I. Part I’s hidden and deep examples use ordinary poles; the new ones put descending exponents in front of ordinary *essential* singularities. The localization manuscript states that it does not settle the fate of Part I’s pole-only examples, and nothing in this part settles it. What this part does add about those examples is narrow: they become trivial

over $\mathcal{M}_\Gamma$ (corollary 27.2), so no argument that survives meromorphic extension can prove them nontrivial over $\mathcal{H}_\Gamma$. This part leaves their status over $\mathcal{H}_\Gamma$ open; Part V, added later from a fourth source, settles it negatively (theorem 35.1 and corollary 35.2).

The mechanism fits in one sentence. A nonzero meromorphic coefficient cannot cancel an essential singularity, so a gluing equation that attaches an essential singularity to each member of a strictly decreasing sequence of exponents forces that sequence into the support of one global Hahn section, which is impossible. Neither a determinant, nor a positive normalization, nor a bound on the valuation of a hypothetical gauge enters the argument; this is why it works after localization, where Part I's integral argument does not.

19.2 An explicit model, and the results at a glance

Let $\Gamma = \mathbb{Q}$, let $A = \{1, 2, 3, \dots\}$, and put

$$c_n(z) = t^{1/n} \exp\left(\frac{1}{z-n}\right). \quad (19.1)$$

Use the cover of $\mathbb{C}$ by $U_0 = \mathbb{C} \setminus A$ and pairwise disjoint small disks D_n centred at n . If J_r denotes the nilpotent Jordan matrix with ones immediately above the diagonal, glue rank- r free modules by

$$G_n = \exp(c_n J_r) = I_r + c_n J_r + \frac{c_n^2 J_r^2}{2!} + \dots + \frac{c_n^{r-1} J_r^{r-1}}{(r-1)!}. \quad (19.2)$$

The exponential is a finite expression. Each G_n and G_n^{-1} is an integral Hahn matrix, even though the union $\{1/n : n \geq 1\}$ of the first nonconstant exponents is not well ordered. Write $\mathcal{B}$ for any one of the integral holomorphic, full holomorphic, or full coefficientwise meromorphic Hahn sheaves $\mathcal{I}_\Gamma$, $\mathcal{H}_\Gamma$, $\mathcal{M}_\Gamma$ of definition 20.1, and $A_{\mathcal{B}} = \mathcal{B}(\mathbb{C})$. For the resulting bundle $\mathcal{N}_{\alpha,r}^{\mathcal{B}}$ we prove

$$H^0(\mathbb{C}, \mathcal{N}_{\alpha,r}^{\mathcal{B}}) = A_{\mathcal{B}} e_1, \quad \text{End}_{\mathcal{B}}(\mathcal{N}_{\alpha,r}^{\mathcal{B}}) \cong A_{\mathcal{B}}[T]/(T^r). \quad (19.3)$$

For $r \geq 2$ the bundle is indecomposable, is not stably trivial, and has no coefficientwise meromorphic frame even over $\mathcal{M}_\Gamma$. Yet every bounded restriction is already trivial over the integral sheaf. This is not a rank-one Picard-group phenomenon placed on a diagonal, as in (11.4): the determinant is trivial, all successive flag quotients are trivial lines, and the bundle itself is indecomposable.

The profile $1/n$ in (19.1) can be replaced by any strictly decreasing sequence $\alpha_1 > \alpha_2 > \dots > 0$ in any set-sized ordered abelian group.

Theorem 19.1 (Summary of Part III). *Let $\alpha = (\alpha_n)$ be such a profile and let $r \geq 2$.*

- (i) *The bundle $\mathcal{N}_{\alpha,r}$ has trivial ordinary reduction and trivial determinant. Every bounded restriction is trivial over $\mathcal{I}_\Gamma$, but the global bundle is indecomposable and not stably trivial over $\mathcal{I}_\Gamma$, $\mathcal{H}_\Gamma$ or $\mathcal{M}_\Gamma$.*
- (ii) *For nilpotent complex matrices N and M the corresponding bundles satisfy*

$$\text{Hom}_{\mathcal{B}}(\mathcal{N}_\alpha(N), \mathcal{N}_\alpha(M)) = A_{\mathcal{B}} \otimes_{\mathbb{C}} \{B : MB = BN\}.$$

Thus the construction realizes the scalar-extended category of finite nilpotent representations, including its direct sums, duals and tensors.

- (iii) *For another strictly decreasing positive profile $\beta = (\beta_n)$,*

$$\mathcal{N}_{\alpha,r}^{\mathcal{I}_\Gamma} \cong \mathcal{N}_{\beta,r}^{\mathcal{I}_\Gamma} \iff \alpha_n = \beta_n \text{ for all sufficiently large } n,$$

whereas for $\mathcal{B} = \mathcal{H}_\Gamma$ or $\mathcal{B} = \mathcal{M}_\Gamma$,

$$\mathcal{N}_{\alpha,r}^{\mathcal{B}} \cong \mathcal{N}_{\beta,r}^{\mathcal{B}} \iff \beta_n - \alpha_n = \delta \text{ eventually, for one } \delta \in \Gamma.$$

All isomorphisms here cover the identity of the ordinary base $\mathbb{C}$.

- (iv) Every noncyclic nonzero ordered abelian group admits such profiles and, for each $r \geq 2$, continuum many pairwise nonisomorphic bundles of this kind even over $\mathcal{M}_\Gamma$. Ordered group enlargement cannot identify two previously distinct classes in this family.

The proof is assembled from [theorems 22.3](#) and [23.3](#) and [corollary 22.5](#) for (i), [theorem 23.1](#) and [corollary 23.4](#) for (ii), [theorem 24.1](#) for (iii), and [lemma 11.3](#), [theorem 25.2](#) and [corollary 24.2](#) for (iv). The effective version, [theorem 28.4](#) in Part IV, shows that neither isomorphism nor nonisomorphism is semidecidable for a simple class of computably presented rational profiles. The distinction is between a complete mathematical criterion and an algorithm for unrestricted infinite input.

19.3 What is new here, what is inherited, and what overlaps with Part I

The localization manuscript inspected the repository at commit

0097304c7ae9d5de46d2ea342e2494f8fc126303.

At that commit this report and the global-divisor report [\[2\]](#) were textually identical to the versions into which this part is merged, so its quotations of both are current. By its own assessment, the scalar full-Hahn Picard dichotomy of the global-divisor report and the integral cyclic rigidity theorem of Part I ([theorem 11.4](#)) are antecedents, not new results. Its proposed new contributions are the zero-annihilator essential-singularity obstruction, its meromorphic-resistant finite-rank realization, the complete morphism calculation in that realization, and the tail-translation classification; the localization-stable dichotomy and the computability consequences follow from those.

There are classical precedents. The Hahn support calculus is classical [\[9\]](#). Atiyah’s bundles on an ordinary elliptic curve are a familiar indecomposable unipotent family with related section and tensor patterns [\[28\]](#). Here the base is instead the ordinary plane, the coefficient sheaf is different, and the profile and localization invariants are essential. The Jordan-matrix tensor identity of [lemma 26.2](#) is classical linear algebra, not a new result; a proof is included so that the application does not depend on a comparison with elliptic curves. Müller–Strohmaier’s Hahn-meromorphic function theory [\[31\]](#) uses generalized series with a convergence condition near a distinguished point on a logarithmic covering; the common-domain, coefficientwise-meromorphic sheaf used here is defined differently, and the word “meromorphic” must not be used to identify the two categories (compare item [21](#) of [section 30](#)). Within the repository material and primary sources it inspected, the localization manuscript located no statement matching the combined construction and exact classification. That is a bounded novelty assessment, not a proof of historical priority.

The following overlaps with Part I and with the global-divisor report are recorded by the merge; the localization manuscript does not state them.

- Over $\mathcal{J}_\Gamma$, nontriviality (though not stable nontriviality, indecomposability or classification) of the new bundles is already an instance of Criterion P; see [remark 22.2](#).
- The ordered-group lemma used in [section 25](#) is [lemma 11.3](#); the global-divisor report proves it again as its `global:lem:descending`, and the localization manuscript proves it a third time. It is cited here, not reprinted.

- The equivalence (i) $\Leftrightarrow$ (ii) of [theorem 25.1](#) is [theorem 11.4](#) on the plane.
- [Theorem 22.3](#) parallels [corollary 10.3](#); its explicit gauges are kept because the sufficiency half of [theorem 24.1](#) reuses them.
- For $\mathcal{H}_\Gamma$ and $\mathcal{I}_\Gamma$, [lemma 20.3](#) is the unit criterion of the global-divisor report (its `global:prop:units`).

What is new relative to the collection is everything over $\mathcal{H}_\Gamma$ and $\mathcal{M}_\Gamma$: the zero-annihilator class, the global sections, the morphisms, the classification, the tensor rule, the continuum count and the undecidability statement.

20 The meromorphic coefficient sheaf

20.1 Three coefficient categories on the ordinary plane

Throughout this part Γ is a set-sized totally ordered abelian group. No rank, divisibility, Archimedean, or countable-cofinality assumption is imposed (compare [remark 3.3](#)).

Definition 20.1 (Common-domain Hahn sections). For a nonempty connected ordinary open set $U \subseteq \mathbb{C}$, let $\mathcal{H}_\Gamma(U)$, $\mathcal{I}_\Gamma(U)$ and $\mathcal{J}_\Gamma(U)$ be as in (2.2)–(2.4), and put

$$\mathcal{M}_\Gamma(U) = \text{Mer}(U)((t^\Gamma)). \quad (20.1)$$

An element of $\mathcal{M}_\Gamma(U)$ is a formal sum $f = \sum_{\gamma \in S} f_\gamma t^\gamma$ with a single well-ordered support S and every coefficient f_γ an ordinary meromorphic function on the same U . Zero coefficients are omitted from the support. For a disconnected open set use the product over its connected components, with no common support requirement between different components.

The field $\text{Mer}(U)$ consists of ordinary meromorphic functions: poles are allowed, essential singularities at points of U are not. On a punctured disk $\exp(1/z)$ is holomorphic and hence meromorphic there; it does not extend as a meromorphic function across the missing centre.

The distinction between common-domain and coefficientwise-germ constructions is load-bearing. A section on U has all its coefficients on that same U ; the domain may not shrink as the exponent varies, and a global section cannot be assembled from a family whose total support is not well ordered.

Where these sheaves sit in the collection. $\mathcal{H}_\Gamma$ is the row $\mathcal{O}(D)((t^\Gamma))$ of the four-ring table in the collection’s index [1], all coefficients holomorphic on one fixed ordinary domain, sheafified over the plane. It is the same sheaf as in Parts I and II and in the global-divisor report [2]; its stalk at an ordinary point is the common-domain germ ring of that table, and it is neither the radius-free ring $\mathbb{C}\{z\}((t^\Gamma))$ nor the formal ring $\mathbb{C}[[z]]((t^\Gamma))$. The sheaf $\mathcal{M}_\Gamma$ is *not* one of the four tabulated constructions: its coefficients are meromorphic, not holomorphic, on the specified domain, and its sections on a connected domain form a field. It is not new to the collection, however. The global-divisor report defines the same recipe over a compact ordinary Riemann surface, under the same name $\mathcal{M}_\Gamma$ (its `global:def:compactsheaf`), and uses it for coherent meromorphic sections and Cartier data there; and on a connected domain the ring $\mathcal{M}_\Gamma(U)$ is the class $\mathcal{M}(U)$ of Hahn series with meromorphic coefficients on which the analysis report [4] defines its coefficientwise cluster residue (its `d:def:clusterres`). What this part adds is its use on the plane as a coefficient sheaf for vector bundles.

Lemma 20.2 (Sheaf property and preservation of support). *The assignments $\mathcal{H}_\Gamma$, $\mathcal{I}_\Gamma$, $\mathcal{J}_\Gamma$ and $\mathcal{M}_\Gamma$ are sheaves on the ordinary complex plane, of commutative rings except for $\mathcal{J}_\Gamma$, which is a sheaf of rings without unit. If $V \subseteq U$ are nonempty connected open sets, restriction preserves the support of every section.*

Proof. For $\mathcal{H}_\Gamma$, $\mathcal{I}_\Gamma$, $\mathcal{J}_\Gamma$ this is the observation of [section 2.2](#). For $\mathcal{M}_\Gamma$ the argument is the same: a nonzero meromorphic function on a connected domain cannot vanish identically on a nonempty open subdomain, so every nonzero coefficient survives restriction and no support exponent is lost. For gluing, first refine an open cover to connected sets. On two overlapping sets, equal restrictions have equal supports. On a connected union, any two members can be joined by a finite chain of intersecting members, since otherwise the unions of the two components of the intersection graph would disconnect the union. All local supports are consequently one and the same well-ordered set. At each exponent the ordinary coefficient functions glue, producing a section with support contained in that set; uniqueness is coefficientwise. The positive and nonnegative conditions are preserved. Disconnected unions are handled component by component. $\square$

20.2 Leading terms and units

We use the Neumann support calculus of [lemma 3.2](#); it justifies convolution and the positive-support geometric inverse, and these are algebraic support statements, not convergence of partial sums. For nonzero f write

$$v(f) = \min \operatorname{supp} f, \quad \operatorname{lc}(f) = f_{v(f)}.$$

On a connected domain the coefficient rings $\mathcal{O}(U)$ and $\operatorname{Mer}(U)$ are domains, so

$$v(fg) = v(f) + v(g), \quad \operatorname{lc}(fg) = \operatorname{lc}(f) \operatorname{lc}(g). \quad (20.2)$$

The three global section rings $\mathcal{I}_\Gamma(\mathbb{C})$, $\mathcal{H}_\Gamma(\mathbb{C})$ and $\mathcal{M}_\Gamma(\mathbb{C})$ are therefore integral domains, and $\mathcal{M}_\Gamma(\mathbb{C}) = \operatorname{Mer}(\mathbb{C})((t^\Gamma))$ is a field.

Lemma 20.3 (Scalar units). *On a connected domain U , a nonzero $f \in \mathcal{H}_\Gamma(U)$ is a unit exactly when its leading coefficient is a nowhere-zero holomorphic function. Every nonzero $f \in \mathcal{M}_\Gamma(U)$ is a unit. A section of $\mathcal{I}_\Gamma(U)$ is a unit exactly when $v(f) = 0$ and its coefficient at zero is nowhere zero.*

Proof. If $fg = 1$, (20.2) makes the leading coefficient of f a unit of the ordinary coefficient ring. Conversely write $f = t^\gamma a(1 + h)$ with $\operatorname{supp} h \subseteq \Gamma_{>0}$ and a the invertible leading coefficient. The inverse is $t^{-\gamma} a^{-1} \sum_{k \geq 0} (-h)^k$, a legitimate section on the original domain by the positive-support Hahn lemma ([lemma 3.4](#)). For $\mathcal{M}_\Gamma$, every nonzero meromorphic a has a meromorphic inverse. In the integral ring both the unit and its inverse have nonnegative valuation, forcing $\gamma = 0$; the converse follows by the same formula. $\square$

For $\mathcal{H}_\Gamma$ and $\mathcal{I}_\Gamma$ this is the global-divisor report's exact unit criterion, and the global-divisor report records the same invertibility for its compact $\mathcal{M}_\Gamma$. Every full holomorphic Hahn section becomes integral after multiplication by a suitable monomial, so $\mathcal{H}_\Gamma$ is the monomial localization of $\mathcal{I}_\Gamma$. Only $\mathcal{I}_\Gamma$ carries the ordinary reduction map (2.5), with kernel $\mathcal{J}_\Gamma$; there is no such homomorphism on $\mathcal{H}_\Gamma$, since $t^\gamma t^{-\gamma} = 1$.

20.3 Bundles and base change

For $\mathcal{B}$ one of $\mathcal{I}_\Gamma$, $\mathcal{H}_\Gamma$, $\mathcal{M}_\Gamma$, a rank- r $\mathcal{B}$ -bundle is a sheaf of $\mathcal{B}$ -modules locally isomorphic to $\mathcal{B}^r$ on the ordinary base. This is a definition in a ringed-space category; no assumption that the stalks are local rings is needed. Put

$$A_{\mathcal{I}_\Gamma} = \mathcal{I}_\Gamma(\mathbb{C}), \quad A_{\mathcal{H}_\Gamma} = \mathcal{H}_\Gamma(\mathbb{C}), \quad A_{\mathcal{M}_\Gamma} = \mathcal{M}_\Gamma(\mathbb{C}).$$

When a statement holds in all three cases, $\mathcal{B}$ denotes the chosen sheaf and $A_{\mathcal{B}}$ its global section ring. Tensor extension along $\mathcal{I}_\Gamma \rightarrow \mathcal{H}_\Gamma \rightarrow \mathcal{M}_\Gamma$ takes a bundle given by transition matrices to the bundle given by the same matrices over the larger sheaf. No claim that global sections commute with tensor extension is assumed: global sections are computed independently in each category.

20.4 The surcomplex reading

When Γ is an ordered additive subgroup of $\mathbf{No}$, the substitution $t^\gamma = \omega^{-\gamma}$ of (2.1) identifies, at each ordinary point where the coefficients are defined, a complex Hahn series with a Conway normal-form expression in $\mathbf{SC} = \mathbf{No}[i]$; the support is a set and Hahn addition and multiplication agree with the normal-form operations [27]. The domain of every coefficient is still an *ordinary* complex domain. In particular $\exp(1/(z-n))$ is an ordinary complex function, not a proposed value of an exponential at an infinite imaginary surreal. For $\Gamma = \mathbb{Q}$, the model (19.1) reads $c_n(z) = \omega^{-1/n} \exp(1/(z-n))$. The theorems of this part thus describe global geometry with surcomplex-valued Hahn coefficients. They are not assertions about a sheaf on the whole proper class $\mathbf{No}[i]$ with its fine topology, and this separation also avoids any use of set-indexed fine-topological limits.

20.5 Names in this part

The localization manuscript was written independently, and several of its symbols collide with symbols of Parts I and II. They are renamed here, in the new material only; the conventions below hold throughout Part III and in the passages of Part IV that refer to it.

This report	Localization manuscript	Reason
$\mathcal{M}_\Gamma = \text{Mer}((t^\Gamma))$	$\mathcal{P}_\Gamma$	$\mathcal{P}$ is Part I's algebra of negative Laurent parts (lemma 4.1) and $\mathcal{P}_{r,\Gamma}$ its polar group; $\mathcal{M}_\Gamma$ is the global-divisor report's name for the same recipe
$\mathcal{N}_\alpha(N), \mathcal{N}_{\alpha,r}$	$\mathcal{E}_\alpha(N), \mathcal{E}_{\alpha,r}$	$\mathcal{E}(G)$ is Part I's polar support union and $\mathcal{E}_G$ its integral bundles
$A = \{1, 2, \dots\}, D_n, W_n = D_n \setminus \{n\}$	$D, U_n, U_n \setminus \{n\}$	Part I's names for punctures, disks and overlaps
Λ_δ	D_δ	the diagonal monomial matrix of (24.5)
$\mathbf{Nil}_{A_{\mathcal{B}}}$	$\mathcal{N}_{A_{\mathcal{B}}}$	category of nilpotent representations, section 23.2
v, lc	v, lc	v is Part I's valuation symbol

Four further points are conventions of meaning. The profile letters α_n, β_n play the role of the γ_n in theorem 11.4 and in the global-divisor report. The function $h_n = \exp(1/(z-n))$ has an *ordinary* essential singularity at an ordinary point; it is the same device as the analysis report's family $te^{1/Z}$, and it is *not* a surreal-scale essential singularity, which the global-divisor report disclaims and the

analysis report limits to one specified function. The only property of h_n used is that its singularity at n is not a pole or a removable point ([lemma 21.1](#)); no theorem of Picard type is used, and “Picard” in this report always means the Picard group of line bundles. The letter A alone always denotes the puncture set; the global section rings $A_{\mathcal{I}_\Gamma}$, $A_{\mathcal{H}_\Gamma}$, $A_{\mathcal{M}_\Gamma}$, $A_{\mathcal{B}}$ always carry their subscript. Finally, the gluing convention of this part differs from Part I’s; see [section 22.1](#).

21 An essential singularity protects a descending support

21.1 The fixed cover and the local data

Throughout this part set

$$A = \{1, 2, 3, \dots\}, \quad U_0 = \mathbb{C} \setminus A, \quad D_n = \{z : |z - n| < 1/4\}, \quad W_n = D_n \setminus \{n\} \quad (n \geq 1).$$

The disks are pairwise disjoint, and the only nonempty overlaps between unequal members of the cover are $U_0 \cap D_n = W_n$. This is a puncture-and-disk cover of $X = \mathbb{C}$ in the sense of [section 6](#). Choose a strictly decreasing positive profile

$$\alpha_1 > \alpha_2 > \dots > 0, \quad \alpha_n \in \Gamma, \quad (21.1)$$

and define

$$h_n(z) = \exp\left(\frac{1}{z - n}\right), \quad c_n = t^{\alpha_n} h_n. \quad (21.2)$$

Each c_n is a positive Hahn section on W_n ; since h_n is holomorphic on $\mathbb{C} \setminus \{n\}$, it is even a positive section on $\mathbb{C} \setminus \{n\}$. The family (c_n) need not be, and is not asserted to be, a summable Hahn family over all n .

Lemma 21.1 (Essential protection). *Let g be a nonzero meromorphic germ at n . The germ $h_n g$ on the punctured neighbourhood of n does not extend meromorphically across n .*

Proof. If $h_n g$ extended meromorphically, division by the nonzero meromorphic germ g would extend h_n meromorphically. But its Laurent expansion

$$h_n(z) = \sum_{k=0}^{\infty} \frac{(z - n)^{-k}}{k!}$$

has infinitely many nonzero negative coefficients, so the singularity is essential. The expansion here is the ordinary complex Laurent expansion, not a Hahn sum over Γ . $\square$

Zeros of g at n do not invalidate the argument: a nonzero meromorphic germ can have a zero or a pole only of finite order. This is exactly what makes the construction survive the passage from holomorphic to meromorphic coefficients.

Theorem 21.2 (Zero-annihilator descent obstruction). *Let $\mathcal{B} \in \{\mathcal{I}_\Gamma, \mathcal{H}_\Gamma, \mathcal{M}_\Gamma\}$ and let $f \in A_{\mathcal{B}}$ be nonzero. There do not exist $b_0 \in \mathcal{B}(U_0)$ and $b_n \in \mathcal{B}(D_n)$ satisfying*

$$b_n = b_0 + c_n f \quad \text{on } W_n, \quad n \geq 1. \quad (21.3)$$

The same assertion holds after every order-preserving group embedding $\Gamma \hookrightarrow \Gamma'$.

Proof. Put $\beta = v(f)$ and $g = f_\beta$. The function g is a nonzero global holomorphic or meromorphic coefficient, and its germ at every n is nonzero, since otherwise the identity theorem would make it zero everywhere.

At the exponent $\beta + \alpha_n$ the product $c_n f$ has exactly the coefficient $h_n g$, because c_n has a single Hahn exponent and β is the least exponent of f . By [lemma 21.1](#) that coefficient is not meromorphic at n . The coefficient of b_n at the same exponent is meromorphic at n (holomorphic, in the two holomorphic categories). Therefore [\(21.3\)](#) forces the coefficient of b_0 at $\beta + \alpha_n$ to be nonzero, and consequently

$$\{\beta + \alpha_n : n \geq 1\} \subseteq \text{supp } b_0.$$

The set on the left is strictly decreasing and cannot be a subset of a well-ordered support. This contradiction proves the assertion.

An order embedding preserves the strict inequalities in [\(21.1\)](#). A hypothetical solution in the enlarged group still has a least exponent β , now possibly outside Γ , and the same translated decreasing set gives the same contradiction. $\square$

Remark 21.3 (The missing normalization is unnecessary). For $\mathcal{H}_\Gamma$ and $\mathcal{M}_\Gamma$ the number β in the proof can be negative, and there is no bound on the negative valuation of a prospective gauge. Translation by any β preserves strict decrease. Thus the proof does not smuggle the integral reduction normalization of [proposition 10.1](#) into the localized category.

21.2 A cohomological formulation without a cohomology assumption

For the displayed cover form the additive quotient

$$Q_{\mathcal{B}} = \frac{\prod_{n \geq 1} \mathcal{B}(W_n)}{\{(b_n - b_0)_n : b_0 \in \mathcal{B}(U_0), b_n \in \mathcal{B}(D_n)\}}.$$

It is an $A_{\mathcal{B}}$ -module. Since there are no nonempty triple overlaps of distinct members, these are precisely the additive Čech cocycles and coboundaries on this cover.

Corollary 21.4 (A free cyclic obstruction submodule). *The map*

$$A_{\mathcal{B}} \longrightarrow Q_{\mathcal{B}}, \quad f \longmapsto f[(c_n)_n]$$

is injective. In particular the obstruction class has zero annihilator over $A_{\mathcal{B}}$; it is not merely a nonzero torsion class that a global scalar might remove.

Proof. Its kernel is exactly the set of f for which [\(21.3\)](#) has a solution. Apply [theorem 21.2](#). $\square$

No identification of $Q_{\mathcal{B}}$ with derived sheaf cohomology is needed in any proof below, and none is claimed. It is notation for the exact obstruction that actually occurs in the gluing equations.

22 Nilpotent bundles and localization-resistant nontriviality

22.1 Finite exponentials and the gluing convention

Let V be a finite-dimensional complex vector space and let $N \in \text{End}_{\mathbb{C}}(V)$ be nilpotent, say $N^m = 0$. Define

$$G_n(N) = \exp(c_n N) = \sum_{k=0}^{m-1} \frac{c_n^k N^k}{k!}. \quad (22.1)$$

Its inverse is $\exp(-c_n N)$. After an ordinary constant change of basis, N is strictly upper triangular, so $G_n(N)$ has determinant one. These statements hold over each of our coefficient sheaves. The finite support of $G_n(N) - I$ is contained in $\{\alpha_n, 2\alpha_n, \dots, (m-1)\alpha_n\}$.

Definition 22.1 (The nilpotent bundle). The bundle $\mathcal{N}_{\alpha}(N)^{\mathcal{B}}$ is defined by the free modules $\mathcal{B}|_{U_0} \otimes_{\mathbb{C}} V$ and $\mathcal{B}|_{D_n} \otimes_{\mathbb{C}} V$ and the transition convention

$$s_n = G_n(N)s_0 \quad \text{on } W_n. \quad (22.2)$$

For the standard $r \times r$ Jordan block J_r write $\mathcal{N}_{\alpha,r}^{\mathcal{B}} = \mathcal{N}_{\alpha}(J_r)^{\mathcal{B}}$. Superscripts are omitted when the coefficient category is fixed or the statement holds in all three categories.

There is no triple-overlap condition to check beyond inverse transitions, so these data are a genuine bundle, not merely a formal list of local matrices. The bundle is already defined over $\mathcal{I}_{\Gamma}$, its ordinary reduction is the trivial bundle $\mathcal{O} \otimes_{\mathbb{C}} V$, and its determinant is canonically trivial.

The convention, once. In (22.2) a section is a family of coordinate vectors s_0, s_n related by $G_n(N)$, and a trivialization is a family of invertible matrices with $B_n = G_n(N)B_0$ on W_n , whose columns are a global frame. Part I writes its gluing problem as $G_a = B_0 B_a^{-1}$ (theorem 6.2). The two conventions differ by inverting the transitions: $B_n = G_n(N)B_0$ is the same as $G_n(N)^{-1} = B_0 B_n^{-1}$, so $\mathcal{N}_{\alpha}(N)^{\mathcal{I}_{\Gamma}}$ is Part I's bundle $\mathcal{E}_{G'}$ for $G'_n = G_n(N)^{-1} = \exp(-c_n N)$. Since $\exp(-c_n N)$ is of the same form with h_n replaced by $-h_n$, no statement below depends on the choice.

Remark 22.2 (Over $\mathcal{I}_{\Gamma}$ this is an instance of Criterion P). This remark is a merge observation, not a statement of the localization manuscript. Let $N \neq 0$. In Part I's convention the cocycle is $G'_n = \exp(-c_n N) \in \mathcal{G}_{r,\Gamma}(W_n)$. The Laurent decomposition of h_n at n is $h_n = (h_n - 1) + 1$, where $h_n - 1 = \sum_{k \geq 1} (z - n)^{-k} / k!$ is a negative Laurent part in the sense of lemma 4.1 and 1 is holomorphic on D_n . Scalar multiples of the one matrix N commute, so

$$\exp(-c_n N) = \exp(-t^{\alpha_n} (h_n - 1)N) \exp(-t^{\alpha_n} N).$$

The first factor lies in the polar group $\mathcal{P}_{r,\Gamma}$ of section 5, its nonconstant coefficients being finite sums of products of negative Laurent parts, and the second lies in $\mathcal{G}_{r,\Gamma}(D_n)$. By uniqueness in theorem 5.1, $\text{Pol}(G'_n) = \exp(-t^{\alpha_n} (h_n - 1)N)$, whose coefficient at t^{α_n} is $-(h_n - 1)N \neq 0$. Hence $\mathcal{E}(G') \supseteq \{\alpha_n : n \geq 1\}$ is not well ordered, theorem 6.2 excludes a positive splitting, and proposition 10.1 shows that $\mathcal{N}_{\alpha}(N)^{\mathcal{I}_{\Gamma}}$ is nontrivial. The same computation with h_n replaced by a pole datum such as $1/(z - n)$ also gives integral nontriviality; within Part III the two kinds of data are separated only after passage to $\mathcal{M}_{\Gamma}$ (proposition 27.1). Part V separates them over $\mathcal{H}_{\Gamma}$: with finite-order poles in place of h_n the bundle becomes trivial after one monomial localization (theorem 34.6). What Part I does not give, and this part does, is nontriviality over $\mathcal{H}_{\Gamma}$ and $\mathcal{M}_{\Gamma}$, stable nontriviality, indecomposability, and the classification.

22.2 Every bounded restriction is trivial

Theorem 22.3 (Finite-puncture trivialization). *If an ordinary open set $Y \subseteq \mathbb{C}$ meets only finitely many points of A , then $\mathcal{N}_\alpha(N)|_Y$ is trivial over $\mathcal{I}_\Gamma$. In particular every bounded restriction is integral-trivial.*

Proof. Let $F = A \cap Y$, a finite set, and on $Y \setminus F$ set

$$p = \sum_{n \in F} c_n.$$

This is a finite Hahn sum whose ordinary coefficient functions are holomorphic away from F . Choose the matrices

$$B_0 = \exp(-pN), \quad B_n = \exp((c_n - p)N) \quad (n \in F). \quad (22.3)$$

The function $c_n - p$ is a finite sum of $-c_j$ with $j \neq n$, hence is holomorphic on the disk D_n . Each matrix and its inverse is an integral Hahn matrix. Since all factors are polynomials in the same matrix N , they commute and

$$B_n = \exp(c_n N) \exp(-pN) = G_n(N) B_0.$$

Their columns therefore define a global frame on the induced cover of Y . Disks whose centres lie outside Y are unnecessary, because $Y \setminus F$ already covers their intersection with Y . $\square$

For bounded Y the integral statement is also the argument of [corollary 10.3](#): finitely many transition supports have a well-ordered union and [theorem 11.1](#) applies. The explicit gauges (22.3) are recorded because the sufficiency proof of [theorem 24.1](#) reuses them. The finite formula cannot simply be sent to an infinite limit: the first exponents α_n strictly decrease, so the putative infinite $p = \sum_n c_n$ is not a Hahn section. The theorem gives local trivializations, not one compatible exhaustion-limit frame.

22.3 Exact sequences and the standard flag

The standard coordinate subspaces $0 \subset \mathbb{C}e_1 \subset \mathbb{C}e_1 + \mathbb{C}e_2 \subset \cdots \subset \mathbb{C}^r$ are invariant under J_r . They yield a flag of subbundles

$$0 = F_0 \subset F_1 \subset \cdots \subset F_r = \mathcal{N}_{\alpha,r}, \quad F_k \cong \mathcal{N}_{\alpha,k}, \quad F_k/F_{k-1} \cong \mathcal{B}, \quad (22.4)$$

and in particular an exact sequence

$$0 \longrightarrow \mathcal{B} \longrightarrow \mathcal{N}_{\alpha,r} \longrightarrow \mathcal{N}_{\alpha,r-1} \longrightarrow 0. \quad (22.5)$$

These sequences are locally split, being sequences of free modules in each chart. They need not split globally. The global obstruction is computed next, and it will also show that the flag is intrinsic rather than an arbitrary choice of coordinates.

22.4 All global sections, and nontriviality after localization

Theorem 22.4 (All global sections). *For $\mathcal{B} \in \{\mathcal{I}_\Gamma, \mathcal{H}_\Gamma, \mathcal{M}_\Gamma\}$ and every $r \geq 1$,*

$$H^0(\mathbb{C}, \mathcal{N}_{\alpha,r}^{\mathcal{B}}) = A_{\mathcal{B}} e_1. \quad (22.6)$$

More generally, for every finite-dimensional complex nilpotent N ,

$$H^0(\mathbb{C}, \mathcal{N}_\alpha(N)^{\mathcal{B}}) = A_{\mathcal{B}} \otimes_{\mathbb{C}} \ker N. \quad (22.7)$$

The equalities identify a section with its common coordinate vector in the fixed kernel of N .

Proof. First take $N = J_r$ and write a section in local coordinates as $s_i = (s_{i,1}, \dots, s_{i,r})^T$. The bottom row of G_n is the last row of the identity, so $s_{n,r} = s_{0,r}$ on overlaps, and the sheaf property glues these entries to a global $f_r \in A_{\mathcal{B}}$.

For $r \geq 2$ the next row of the transition equation is

$$s_{n,r-1} = s_{0,r-1} + c_n f_r.$$

If $f_r \neq 0$, this contradicts [theorem 21.2](#); therefore $f_r = 0$. The entries in row $r-1$ now agree on overlaps and define a global f_{r-1} . The equation one row higher has reduced to $s_{n,r-2} = s_{0,r-2} + c_n f_{r-1}$, because all terms involving f_r have vanished, and the same argument forces $f_{r-1} = 0$. Continuing upward kills every coordinate except the first, which glues to an arbitrary element of $A_{\mathcal{B}}$.

Conversely $J_r e_1 = 0$ implies $G_n e_1 = e_1$, so every common vector $f e_1$ with $f \in A_{\mathcal{B}}$ defines a section. This proves (22.6), including $r = 1$. A constant complex change of basis expresses any nilpotent N as a finite direct sum of Jordan blocks; the construction respects that change of basis and finite direct sums. Applying the proved assertion to each block, the resulting invariant lines are precisely $\ker N$. $\square$

The proof never assumes that a global section has nonzero ordinary reduction. Its first nonzero coefficient can occur at any exponent allowed in the chosen coefficient category, which is why the argument remains valid after localization.

Corollary 22.5 (No stable trivialization). *If $r \geq 2$, then for every $m \geq 0$*

$$\mathcal{N}_{\alpha,r}^{\mathcal{B}} \oplus \mathcal{B}^m \not\cong \mathcal{B}^{r+m}.$$

In particular $\mathcal{N}_{\alpha,r}^{\mathcal{B}}$ is nontrivial in all three coefficient categories.

Proof. An isomorphism of bundles would induce an isomorphism of global $A_{\mathcal{B}}$ -modules. By [theorem 22.4](#) these are $A_{\mathcal{B}}^{m+1}$ and $A_{\mathcal{B}}^{m+r}$. The ring $A_{\mathcal{B}}$ is a domain, so tensoring with its fraction field would equate vector spaces of different finite dimensions. $\square$

Corollary 22.6 (No coefficientwise meromorphic frame). *For $r \geq 2$, the space of global $\mathcal{M}_\Gamma$ -sections of $\mathcal{N}_{\alpha,r}^{\mathcal{M}_\Gamma}$ has dimension one over the field $A_{\mathcal{M}_\Gamma} = \text{Mer}(\mathbb{C})((t^\Gamma))$. The bundle admits no coefficientwise meromorphic frame, and not even two globally independent coefficientwise meromorphic sections.*

Proof. This is [theorem 22.4](#) for $\mathcal{M}_\Gamma$: any two sections are multiples of e_1 over the indicated field. $\square$

22.5 Which further localizations are covered

A nonzero $f \in \mathcal{H}_\Gamma(U)$ on a connected U has an inverse in $\mathcal{M}_\Gamma(U)$: its leading holomorphic coefficient may have zeros, but its reciprocal is meromorphic, and the positive-support geometric inverse still exists. Therefore the fraction field of $\mathcal{H}_\Gamma(U)$ embeds in $\mathcal{M}_\Gamma(U)$. At the sheaf level the local fraction sheaf of $\mathcal{H}_\Gamma$ embeds in $\mathcal{M}_\Gamma$ as well: represent a nonzero germ by a section on a connected neighbourhood and apply the same inversion.

Corollary 22.7 (Intermediate coefficient categories). *Let $\mathcal{C}$ be a sheaf of rings with compatible inclusions $\mathcal{H}_\Gamma \subseteq \mathcal{C} \subseteq \mathcal{M}_\Gamma$. No bundle $\mathcal{N}_{\alpha,r}$ with $r \geq 2$ becomes stably trivial over $\mathcal{C}$. This includes every such localization obtained by inverting nonzero holomorphic Hahn sections.*

Proof. A stable trivialization over $\mathcal{C}$ would remain a stable trivialization after tensor extension to $\mathcal{M}_\Gamma$, contrary to [corollary 22.5](#). $\square$

The statement is deliberately about one specified inclusion of sheaves. It does not identify $\mathcal{M}_\Gamma$ with every possible definition of a surreal or Hahn meromorphic function, and it does not concern a coefficient category that allows essential singularities at every point of the ordinary base.

Corollary 22.8 (Absolute resistance under ordered enlargement). *If $\Gamma \hookrightarrow \Gamma'$ is an order-preserving group embedding, the base-changed bundle $\mathcal{N}_{\alpha,r}$ with $r \geq 2$ remains nontrivial and not stably trivial over each of the three Γ' -coefficient sheaves. Its global section formula is still (22.6), with the enlarged global ring.*

Proof. The image profile is still strictly decreasing and positive. Apply [theorems 21.2](#) and [22.4](#) and [corollary 22.5](#) in Γ' . $\square$

Thus enlarging a set-sized surreal workspace cannot repair this obstruction. The claim follows from the order of an explicit sequence, not from an assumption that the original coefficient group was maximal. This is the localized counterpart of [corollary 7.3](#), which is a statement about positive integral splittings only.

23 Every morphism and the intrinsic flag

23.1 The internal Hom is another nilpotent bundle

Theorem 23.1 (Complete morphism formula). *Let $N \in \text{End}_\mathbb{C}(V)$ and $M \in \text{End}_\mathbb{C}(V')$ be nilpotent, and use the same profile α for both bundles. Then*

$$\begin{aligned} & \text{Hom}_{\mathcal{B}}(\mathcal{N}_\alpha(N)^{\mathcal{B}}, \mathcal{N}_\alpha(M)^{\mathcal{B}}) \\ &= \{B \in A_{\mathcal{B}} \otimes_{\mathbb{C}} \text{Hom}_{\mathbb{C}}(V, V') : MB = BN\} \\ &= A_{\mathcal{B}} \otimes_{\mathbb{C}} \text{Hom}_{\mathbb{C}[T]}((V, N), (V', M)). \end{aligned} \tag{23.1}$$

Every such morphism is represented by the same global Hahn matrix in all charts, and there are no other morphisms.

Proof. In local coordinates a bundle morphism is a family of matrices B_i with compatibility

$$B_n = \exp(c_n M) B_0 \exp(-c_n N). \tag{23.2}$$

On the finite-dimensional complex space $\text{Hom}_{\mathbb{C}}(V, V')$ set

$$L_M(B) = MB, \quad R_N(B) = BN, \quad T = L_M - R_N.$$

The operators L_M and R_N commute. If $M^a = N^b = 0$, the binomial formula gives $T^{a+b-1} = 0$, since each term contains L_M^a or R_N^b . Their finite exponentials consequently satisfy

$$\exp(c_n T)(B) = \exp(c_n M) B \exp(-c_n N).$$

Thus the internal Hom bundle is exactly $\mathcal{N}_\alpha(T)$, and its global sections are $A_{\mathcal{B}} \otimes_{\mathbb{C}} \ker T$ by [theorem 22.4](#). The condition $T(B) = 0$ is $MB = BN$, proving the formula. A matrix in this kernel is fixed by all transitions, which explains the common global representative. Since the defining linear equations have complex coefficients and finite dimension, extending their solution space from $\mathbb{C}$ to the complex algebra $A_{\mathcal{B}}$ commutes with taking the kernel. $\square$

Corollary 23.2 (Hom spaces, endomorphisms and automorphisms). *For $r, s \geq 1$ the module $\text{Hom}_{\mathcal{B}}(\mathcal{N}_{\alpha,r}, \mathcal{N}_{\alpha,s})$ is free of rank $\min(r, s)$ over $A_{\mathcal{B}}$. Moreover*

$$\text{End}_{\mathcal{B}}(\mathcal{N}_{\alpha,r}) \cong A_{\mathcal{B}}[T]/(T^r), \quad (23.3)$$

$$\text{Aut}_{\mathcal{B}}(\mathcal{N}_{\alpha,r}) = \left\{ \sum_{k=0}^{r-1} a_k J_r^k : a_k \in A_{\mathcal{B}}, a_0 \in A_{\mathcal{B}}^\times \right\}. \quad (23.4)$$

Proof. An intertwiner B from J_r to J_s is determined by $x = Be_r$, since e_r is cyclic and $Be_{r-k} = J_s^k x$ for $0 \leq k < r$. The sole constraint is $J_s^r x = 0$, whose solution module is free of rank $\min(r, s)$. For $r = s$, the image of e_r can be written uniquely as $\sum_{k=0}^{r-1} a_k J_r^k e_r$, so every commuting matrix is the corresponding polynomial in J_r ; this proves (23.3). A polynomial in J_r is upper triangular with diagonal a_0 , and it is invertible exactly when a_0 is a unit; then its inverse is a finite geometric polynomial in its nilpotent nonconstant part. $\square$

Theorem 23.3 (Indecomposability). *Every $\mathcal{N}_{\alpha,r}^{\mathcal{B}}$ is indecomposable, including after all the coefficient and ordered-group extensions above.*

Proof. A nontrivial direct-sum decomposition would provide an idempotent in the endomorphism ring different from zero and one. In $A_{\mathcal{B}}[T]/(T^r)$, reduction modulo T maps an idempotent to an idempotent of the domain $A_{\mathcal{B}}$, hence to zero or one. An idempotent in the nilpotent ideal (T) is zero: if $e^2 = e$ and $e^r = 0$, then $e = e^r = 0$. If the reduction is one, apply this to $1 - e$. There are no nontrivial idempotents. The same reasoning applies over the enlarged coefficient rings, which are still domains. $\square$

This distinguishes the construction from a mere nontrivial extension class whose middle bundle might nevertheless be decomposable: the full endomorphism calculation controls the middle bundle itself.

23.2 A representation category on the ordinary plane

Let $\mathbf{Nil}_{A_{\mathcal{B}}}$ be the category whose objects are finite-dimensional complex vector spaces equipped with a nilpotent endomorphism, and whose morphism modules are

$$\text{Hom}_{\mathbf{Nil}_{A_{\mathcal{B}}}}((V, N), (V', M)) = A_{\mathcal{B}} \otimes_{\mathbb{C}} \text{Hom}_{\mathbb{C}[T]}((V, N), (V', M)).$$

Composition is the scalar extension of ordinary matrix composition.

Corollary 23.4 (Fully faithful realization). *The construction $N \mapsto \mathcal{N}_\alpha(N)^{\mathcal{B}}$ is an $A_{\mathcal{B}}$ -linear fully faithful functor from $\mathbf{Nil}_{A_{\mathcal{B}}}$ to $\mathcal{B}$ -bundles on $\mathbb{C}$. It preserves finite direct sums and duals, and it is compatible with*

$$(V, N) \otimes (V', M) = (V \otimes V', N \otimes I + I \otimes M).$$

A short exact sequence of finite-dimensional complex nilpotent representations is sent to a short exact sequence of bundles.

Proof. Full faithfulness is [theorem 23.1](#). Direct sums are immediate from the block-diagonal exponential. The dual transition is $\exp(-c_n N^T)$, and the tensor transition is

$$\exp(c_n N) \otimes \exp(c_n M) = \exp(c_n (N \otimes I + I \otimes M)),$$

by the finite binomial formula for commuting nilpotent matrices. An exact sequence of the original complex vector spaces is split as a sequence of vector spaces, so it stays exact after tensoring with each local coefficient ring. The representation maps commute with transitions, and the local exact sequences glue. $\square$

The scalar extension in the definition of $\mathbf{Nil}_{A_{\mathcal{B}}}$ matters: the functor from the original $\mathbb{C}$ -linear category alone would not be full, because nonconstant global Hahn scalars produce additional morphisms.

Corollary 23.5 (Classification within one profile). *For a fixed profile, $\mathcal{N}_{\alpha}(N)$ and $\mathcal{N}_{\alpha}(M)$ are isomorphic if and only if N and M have the same Jordan block sizes, with multiplicities. Hence every object in this image is a finite direct sum of the indecomposables $\mathcal{N}_{\alpha,r}$, and the multiset of summands is unique.*

Proof. Identical Jordan types give an ordinary constant conjugacy and hence an isomorphism. Conversely an isomorphism is an invertible global intertwiner by [theorem 23.1](#). Extend the coefficient domain $A_{\mathcal{B}}$ to its fraction field. The matrices become similar there, so the ranks of all powers agree; those ranks determine the nilpotent Jordan partition and have the same values before and after this field extension. $\square$

No Krull–Schmidt theorem for the entire category of Hahn bundles is invoked; the classification is for the fully specified image of this functor.

23.3 Recovering the flag intrinsically

Proposition 23.6 (The canonical global-section flag). *For $\mathcal{N}_{\alpha,r}$, the subsheaf generated by all global sections is precisely F_1 in [\(22.4\)](#). Repeating this operation in the quotient and taking inverse images recovers each F_k . Thus every isomorphism between two Jordan-profile bundles of the same rank preserves the indicated flags, even if the two profiles differ.*

Proof. By [theorem 22.4](#) all global sections lie in $A_{\mathcal{B}}e_1$, and the global section e_1 generates $\mathcal{B}e_1 = F_1$ as a sheaf. The quotient is the Jordan bundle of rank $r - 1$ for the same profile, so the same reasoning identifies its first flag step, and induction recovers all steps. The construction uses only global sections, quotient sheaves and inverse images, and is therefore preserved by an isomorphism. $\square$

An arbitrary localized gauge need not preserve a flag given in advance. The proposition is what permits triangular form in the next proof: the flag is recovered from the bundle, not imposed as extra structure on its isomorphisms.

24 Exact classification of exponent profiles

Keep the same punctures, disks and functions h_n . Let $\alpha = (\alpha_n)$ and $\beta = (\beta_n)$ be two strictly decreasing positive profiles. All bundle isomorphisms in this section are over the identity map of $\mathbb{C}$; base reparametrizations and permutations of punctures are not included.

Theorem 24.1 (Integral equality versus localized translation). *For every fixed $r \geq 2$,*

$$\mathcal{N}_{\alpha,r}^{\mathcal{J}_\Gamma} \cong \mathcal{N}_{\beta,r}^{\mathcal{J}_\Gamma} \iff \beta_n = \alpha_n \text{ eventually,} \quad (24.1)$$

$$\mathcal{N}_{\alpha,r}^{\mathcal{H}_\Gamma} \cong \mathcal{N}_{\beta,r}^{\mathcal{H}_\Gamma} \iff \beta_n - \alpha_n = \delta \text{ eventually, for some } \delta \in \Gamma, \quad (24.2)$$

$$\mathcal{N}_{\alpha,r}^{\mathcal{M}_\Gamma} \cong \mathcal{N}_{\beta,r}^{\mathcal{M}_\Gamma} \iff \beta_n - \alpha_n = \delta \text{ eventually, for some } \delta \in \Gamma. \quad (24.3)$$

Whenever the relevant condition holds, an isomorphism is obtained from one constant diagonal monomial matrix and finite-exponential gauges supported at finitely many exceptional punctures.

24.1 Necessity: the rank-two calculation

Proof of necessity. First take $r = 2$. The intrinsic flag of [proposition 23.6](#) forces every local isomorphism matrix to be upper triangular. Write its central representative as

$$B_0 = \begin{pmatrix} a & b_0 \\ 0 & d \end{pmatrix}.$$

The diagonal entries are unchanged by the unipotent transition equation, so they glue to global scalars a, d ; the inverse matrices glue their inverses, hence $a, d \in A_{\mathcal{D}}^\times$. Let b_n denote the upper-right entry of the matrix on D_n . With $c_{\alpha,n} = t^{\alpha_n} h_n$ and $c_{\beta,n} = t^{\beta_n} h_n$, the compatibility $B_n = \exp(c_{\beta,n} J_2) B_0 \exp(-c_{\alpha,n} J_2)$, which is the two-profile form of [\(23.2\)](#), gives by direct multiplication

$$b_n = b_0 + d c_{\beta,n} - a c_{\alpha,n}. \quad (24.4)$$

Divide by the global unit d and put

$$q = a/d \in A_{\mathcal{D}}^\times, \quad \gamma = v(q), \quad q_\gamma = \text{lc}(q).$$

The coefficient of $q c_{\alpha,n}$ at $\lambda_n = \gamma + \alpha_n$ is $q_\gamma h_n$. If $\lambda_n \neq \beta_n$, the term $c_{\beta,n}$ has zero coefficient at λ_n . As b_n/d is meromorphic at n , [lemma 21.1](#) forces

$$\lambda_n \in \text{supp}(b_0/d) \quad \text{for every } n \text{ with } \beta_n \neq \gamma + \alpha_n.$$

If infinitely many such exceptional indices existed, their λ_n would form a strictly decreasing subset of a well-ordered support. Thus only finitely many are exceptional, and

$$\beta_n - \alpha_n = \gamma \quad \text{eventually.}$$

In the integral category a global unit has valuation zero by [lemma 20.3](#), so $\gamma = 0$. In the full holomorphic and meromorphic categories any $\gamma \in \Gamma$ is allowed.

For $r > 2$, [proposition 23.6](#) makes an isomorphism restrict to an isomorphism between the intrinsic rank-two subbundles F_2 . These are the rank-two bundles for the two profiles, and the proved necessity applies to that restriction. This reduction does not assume that the original isomorphism was a polynomial in J_r . $\square$

24.2 Sufficiency: explicit finite gauges

Proof of sufficiency. Suppose $\beta_n = \alpha_n + \delta$ except at a finite set F of indices. In $\mathcal{H}_\Gamma$ or $\mathcal{M}_\Gamma$ the invertible constant diagonal matrix

$$\Lambda_\delta = \text{diag}(t^{(r-1)\delta}, t^{(r-2)\delta}, \dots, t^\delta, 1) \quad (24.5)$$

satisfies $\Lambda_\delta J_r \Lambda_\delta^{-1} = t^\delta J_r$. It therefore identifies the α bundle with the bundle whose transition matrices are $\exp(t^{\alpha_n + \delta} h_n J_r)$. The shifted exponents at the finitely many exceptional indices need not be positive; finite exponentials remain valid full Hahn matrices regardless.

Put

$$d_n = (t^{\beta_n} - t^{\alpha_n + \delta}) h_n, \quad p = \sum_{n \in F} d_n.$$

Then $d_n = 0$ outside F . On U_0 take $B_0 = \exp(-p J_r)$, and on D_n take $B_n = \exp((d_n - p) J_r)$. The function $d_n - p$ is holomorphic on D_n , because the only possible singularity there, from the n th term, has cancelled. Since these are polynomials in J_r ,

$$B_n = \exp(t^{\beta_n} h_n J_r) B_0 \exp(-t^{\alpha_n + \delta} h_n J_r).$$

Thus they give the desired isomorphism from the shifted bundle to the β bundle. Compose with Λ_δ .

For the integral statement, eventual equality means $\delta = 0$ and no diagonal localization is needed. Every exponent in the finite differences d_n is positive, so all the gauges just constructed and their inverses lie in $\mathcal{S}_\Gamma$. This proves sufficiency in all three categories. $\square$

Corollary 24.2 (Group enlargement creates no new profile identifications). *An order-preserving embedding $\Gamma \hookrightarrow \Gamma'$ neither identifies nor separates isomorphism classes of this profile family in any of the three coefficient categories.*

Proof. Existing isomorphisms extend. Conversely, the classification over Γ' forces eventual equality or an eventual translation δ' . In the latter case choose one index in the eventual tail; then $\delta' = \beta_n - \alpha_n$ lies in the image of Γ , so the classification over Γ already gives an isomorphism. $\square$

Corollary 24.3 (No additional identifications from meromorphic coefficients). *For this family, extension from $\mathcal{H}_\Gamma$ to $\mathcal{M}_\Gamma$ reflects isomorphism. The same exact classification holds in every intermediate sheaf $\mathcal{H}_\Gamma \subseteq \mathcal{C} \subseteq \mathcal{M}_\Gamma$.*

Proof. An isomorphism over $\mathcal{C}$ extends to $\mathcal{M}_\Gamma$, where necessity is known. The necessary eventual-translation condition already gives an isomorphism over $\mathcal{H}_\Gamma$, hence over $\mathcal{C}$. $\square$

24.3 Examples that separate the categories

Example 24.4 (Localization forgets a constant shift). Over $\Gamma = \mathbb{Q}$ take $\alpha_n = 1 + 1/n$ and $\beta_n = 2 + 1/n$. The corresponding bundles are not integral-isomorphic, but they are full-Hahn-isomorphic by Λ_1 . Both remain indecomposable and nontrivial over $\mathcal{M}_\Gamma$. Thus localization has nontrivial fibres on integral isomorphism classes without making the individual bundles trivial.

Example 24.5 (A nonconstant perturbation survives localization). Take $\alpha_n = 1/n$ and $\beta_n = 1/n + 1/(10n(n+1))$. Both profiles decrease strictly to zero, and their difference is never zero and not eventually constant. The associated rank- r bundles are nonisomorphic even over $\mathcal{M}_\Gamma$, for every $r \geq 2$.

Example 24.6 (Arbitrary finite changes are invisible). Modify finitely many values of a decreasing profile while retaining strict positive decrease. The resulting bundle is already integral-isomorphic to the original one, by the finite gauge of the sufficiency proof; there is no need to solve an infinite interpolation problem.

These examples exhibit the invariant explicitly. It is not the set of exponents, their infimum, a total support cardinality, or the determinant. It is the indexed eventual profile, modulo one global translation in the localized categories.

25 Value groups: a localization-stable dichotomy and continuum many classes

25.1 Which groups admit a descending positive profile

By [lemma 11.3](#), a set-sized ordered abelian group has no infinite strictly decreasing sequence in $\Gamma_{>0}$ exactly when $\Gamma = \{0\}$ or $\Gamma = \mathbb{Z}\delta$ for a least positive element δ . That lemma is printed once, in Part I; the global-divisor report and the localization manuscript each prove it again, by essentially the same argument, and the localization manuscript notes that the recursive choice of a decreasing sequence uses only ordinary countable choice. Having a least positive element is not enough to prevent the obstruction: a non-Archimedean ordered group may have a least positive element and still contain elements larger than every integer multiple of it ([example 25.4](#)).

25.2 The localization-stable rigidity dichotomy

Theorem 25.1 (Universal rigidity versus resistant examples). *For a set-sized ordered abelian group Γ the following are equivalent:*

- (i) $\Gamma = \{0\}$ or Γ is ordered cyclic;
- (ii) every finite-rank $\mathcal{S}_\Gamma$ -bundle on $\mathbb{C}$ with trivial ordinary reduction is already trivial over $\mathcal{S}_\Gamma$;
- (iii) every such bundle becomes trivial over $\mathcal{H}_\Gamma$;
- (iv) every such bundle becomes trivial over $\mathcal{M}_\Gamma$.

When these conditions fail, for every $r \geq 2$ there is a determinant-one rank- r counterexample that is indecomposable and not stably trivial over $\mathcal{M}_\Gamma$, but integral-trivial on every bounded domain.

Proof. (i) $\Rightarrow$ (ii). For $\Gamma = \{0\}$ the integral sheaf is $\mathcal{O}$ and the assertion is the hypothesis. Let $\Gamma = \mathbb{Z}\delta$ with $\delta > 0$. Choose local frames of the bundle on an open cover (U_i) of $\mathbb{C}$ with transition matrices $G_{ij} \in \mathrm{GL}_r(\mathcal{S}_\Gamma(U_i \cap U_j))$. Their reductions are transition matrices of the trivial ordinary bundle, so $\mathrm{red} G_{ij} = g_i g_j^{-1}$ with $g_i \in \mathrm{GL}_r(\mathcal{O}(U_i))$, and after the ordinary changes of frame g_i the transitions $g_i^{-1} G_{ij} g_j$ reduce to I_r , that is, they form a positive cocycle. Its supports lie in the universal well-ordered set $\{\delta, 2\delta, \dots\}$, so it is globally support-admissible, and [theorem 11.1](#) splits it on the same cover: this is the cyclic half of [theorem 11.4](#). The localization manuscript proves this implication directly by the q -adic form of the same coefficientwise recursion: with $q = t^\delta$ one has $\mathcal{S}_\Gamma(U) = \mathcal{O}(U)[[q]]$ on connected domains; on a fixed locally finite disk cover, frames congruent to the identity modulo q^m are corrected by matrices $I + q^m B_i$ solving an ordinary additive matrix cocycle, which is a coboundary by $H^1(\mathbb{C}, \mathcal{O}) = 0$ (equivalently, the ordinary additive Cousin theorem); the coefficient at each q^k is unchanged by all steps with $m > k$, the final frames are defined by these stabilized coefficients, and no analytic convergence is used. The two routes use the same analytic input, and the recursion is not printed twice. Neither is claimed as a new Stein theorem.

(ii) $\Rightarrow$ (iii) $\Rightarrow$ (iv) follow by tensor extension. If (i) fails, [lemma 11.3](#) gives a strictly decreasing positive profile. The bundles of [definition 22.1](#) have trivial reduction, but [corollary 22.6](#) refutes (iv) already in rank two. The stronger claims follow from [theorems 22.3](#) and [23.3](#) and [corollary 22.5](#). $\square$

As the localization manuscript says, the new part of this theorem is that the noncyclic side continues to fail under the negative-valuation and meromorphic extensions, with determinant-one, indecomposable, not stably trivial witnesses. Failure of (iii) is already visible in rank one, for noncyclic $\Gamma \subseteq \mathbf{No}$, from the global-divisor report’s Picard dichotomy. But (iv) cannot be refuted by any of the earlier pole-data examples — neither by the scalar Picard classes of the global-divisor report nor by (11.4) and the examples of Part I — because all of them become trivial over $\mathcal{M}_\Gamma$ (corollary 27.2). The essential-singularity bundles are what make the equivalence with (iv) true.

25.3 Continuum many localized classes

Theorem 25.2 (Cardinality lower bound in every noncyclic group). *If Γ is nonzero and noncyclic, then for every $r \geq 2$ there exist $2^{\aleph_0}$ pairwise nonisomorphic rank- r bundles of this construction over $\mathcal{M}_\Gamma$, and hence over $\mathcal{H}_\Gamma$ and $\mathcal{J}_\Gamma$. All have trivial determinant and reduction, and all their bounded restrictions are trivial.*

Proof. Choose a strictly decreasing positive sequence $(\theta_k)_{k \geq 1}$. For each binary sequence $\varepsilon \in \{0, 1\}^{\mathbb{N}_{>0}}$ define $\alpha_n^\varepsilon = \theta_{2n+\varepsilon_n}$. These profiles are strictly decreasing, since $2n + \varepsilon_n < 2(n+1) + \varepsilon_{n+1}$, and they are distinct for distinct binary sequences.

Fix one profile. If a second is eventually a constant translation of it, the translation belongs to the countable difference set $\{\theta_j - \theta_k : j, k \geq 1\}$. For one fixed translation and one fixed starting index of the eventual tail, each value of the second tail is uniquely determined, since the θ_k are distinct, and there are only finitely many choices for the preceding binary prefix. Thus each translation-equivalence class among these profiles is countable.

There are $2^{\aleph_0}$ binary sequences. A partition of a set of that cardinality into countable classes has $2^{\aleph_0}$ classes, by standard cardinal arithmetic with choice. By theorem 24.1, distinct classes give pairwise nonisomorphic $\mathcal{M}_\Gamma$ -bundles. The other properties have already been proved. $\square$

Example 25.3 (A rational binary family). Over $\Gamma = \mathbb{Q}$ a transparent family is

$$\alpha_n^\varepsilon = \frac{1}{n} + \frac{\varepsilon_n}{10n(n+1)}, \quad \varepsilon_n \in \{0, 1\}. \quad (25.1)$$

For every choice of bits

$$\alpha_n^\varepsilon - \alpha_{n+1}^\varepsilon \geq \frac{1}{n(n+1)} - \frac{1}{10(n+1)(n+2)} > 0.$$

All these profiles tend to zero as ordinary rational sequences, so an eventual constant translation between two of them must be zero. Hence, in each of the three coefficient categories,

$$\mathcal{N}_{\alpha^\varepsilon, r} \cong \mathcal{N}_{\alpha^\eta, r} \iff \varepsilon_n = \eta_n \text{ eventually.} \quad (25.2)$$

The ordinary limit is used only to rule out a nonzero constant rational translation. It is not a convergence assertion about Hahn sections or surcomplex-valued functions.

Example 25.4 (A genuinely non-Archimedean profile). In the ordered group $\Gamma = \mathbb{Z}\omega + \mathbb{Z} \subseteq \mathbf{No}$, ordered lexicographically by the coefficient of ω , take $\alpha_n = \omega - n$. Every exponent is positive, and they strictly decrease while remaining larger than every ordinary integer. The least positive group element is 1, but the group is not cyclic. The same bundle theorems apply without replacing this order by any real-valued valuation. The global-divisor report uses the same profile for its scalar Picard classes.

26 Tensor products and duality

The morphism theorem turns ordinary nilpotent linear algebra into explicit bundle structure. This section records the resulting tensor rule, with its classical algebraic proof separated from the realization.

Theorem 26.1 (Tensor decomposition). *For a fixed profile and all $r, s \geq 1$,*

$$\mathcal{N}_{\alpha,r} \otimes_{\mathcal{B}} \mathcal{N}_{\alpha,s} \cong \bigoplus_{j=0}^{\min(r,s)-1} \mathcal{N}_{\alpha,r+s-1-2j}. \quad (26.1)$$

Moreover $(\mathcal{N}_{\alpha,r})^\vee \cong \mathcal{N}_{\alpha,r}$.

Proof. By [corollary 23.4](#) the tensor bundle is attached to the nilpotent matrix $J_r \otimes I + I \otimes J_s$. Its Jordan block sizes are $r + s - 1, r + s - 3, \dots, |r - s| + 1$, by the finite-dimensional identity [lemma 26.2](#) below, and a constant change of basis gives (26.1). The dual is attached to $-J_r^T$, which has the same single Jordan block as J_r and is conjugate to it by an invertible complex matrix. $\square$

For example $\mathcal{N}_{\alpha,2} \otimes \mathcal{N}_{\alpha,2} \cong \mathcal{N}_{\alpha,3} \oplus \mathcal{N}_{\alpha,1}$ and $\mathcal{N}_{\alpha,2} \otimes \mathcal{N}_{\alpha,3} \cong \mathcal{N}_{\alpha,4} \oplus \mathcal{N}_{\alpha,2}$. Consequently a tensor product of two indecomposable objects need not be indecomposable. The rank of its global section module is the number of summands, $\min(r, s)$, in agreement with self-duality and the Hom formula.

Lemma 26.2 (The Jordan tensor rule). *Over a field of characteristic zero, the matrix $J_r \otimes I_s + I_r \otimes J_s$ has Jordan block sizes $r + s - 1 - 2j$ for $0 \leq j < \min(r, s)$.*

Proof. It is enough to work over $\mathbb{C}$, since the Jordan chains constructed below use rational coefficients and the resulting rank identities are characteristic-zero identities. Put $a = r - 1$ and $b = s - 1$. Let $P_{a,b}$ be the space of polynomials separately homogeneous of degrees a and b in the pairs (X_1, Y_1) and (X_2, Y_2) ; its dimension is $(a + 1)(b + 1)$. Define

$$\begin{aligned} E &= X_1 \partial_{Y_1} + X_2 \partial_{Y_2}, \\ F &= Y_1 \partial_{X_1} + Y_2 \partial_{X_2}, \\ H &= X_1 \partial_{X_1} - Y_1 \partial_{Y_1} + X_2 \partial_{X_2} - Y_2 \partial_{Y_2}. \end{aligned}$$

Direct calculation gives $[E, F] = H$, $[H, E] = 2E$ and $[H, F] = -2F$. On the degree- a polynomials in one pair of variables, the first summand of E has one Jordan block of size $a + 1$: it sends $X^{a-k} Y^k$ to $k X^{a-k+1} Y^{k-1}$, and rescaling that basis gives J_{a+1} . Thus the operator E on $P_{a,b}$ represents the tensor-sum matrix of the statement, up to a constant basis change.

For $0 \leq j \leq \min(a, b)$ set

$$w_j = (X_1 Y_2 - Y_1 X_2)^j X_1^{a-j} X_2^{b-j}, \quad m_j = a + b - 2j.$$

The determinant factor is killed by both E and F and has H -weight zero, so $E w_j = 0$ and $H w_j = m_j w_j$. Commuting E past successive factors F shows inductively that

$$E F^k w_j = k(m_j - k + 1) F^{k-1} w_j. \quad (26.2)$$

Since the X -degree of w_j remaining after the determinant factor is m_j , one has $F^{m_j+1} w_j = 0$. Formula (26.2) implies that none of $F^k w_j$ for $0 \leq k \leq m_j$ is zero: apply E^k and obtain a nonzero

multiple of w_j . These vectors have distinct H -weights and are linearly independent; their span W_j is invariant and carries a single E -Jordan block of size $m_j + 1$.

To see that the different W_j form a direct sum without invoking a semisimplicity theorem, use the operator $C = H^2 + 2H + 4FE$. The three commutator relations give $[C, E] = [C, F] = [C, H] = 0$ by direct expansion. Since $EW_j = 0$, one has $Cw_j = m_j(m_j + 2)w_j$, so C is that scalar on W_j . These scalars are distinct as j varies, and eigenspaces for distinct scalars are independent, so the sum of the W_j is direct. Finally, assuming $a \leq b$ only to evaluate the sum, $\sum_{j=0}^a (m_j + 1) = \sum_{j=0}^a (a + b - 2j + 1) = (a + 1)(b + 1)$. The independent subspaces fill $P_{a,b}$, and their block sizes are exactly $r + s - 1 - 2j$. $\square$

Atiyah’s ordinary elliptic-curve theory is a notable precedent for unipotent indecomposables with this tensor pattern [28]. What is proved here is a realization of the same finite nilpotent algebra in a localized Hahn category on the ordinary plane, together with the profile classification. The tensor pattern by itself is not evidence of novelty.

27 Poles versus essential singularities

27.1 Pole data are removable in the meromorphic category

Proposition 27.1 (Pole data are removable in the meromorphic category). *Replace h_n in (21.2) by any function p_n that is meromorphic on the entire disk D_n , and keep any positive profile, without a global well-ordering assumption on that profile. Then the bundle with transitions $\exp(t^{\alpha_n} p_n N)$ is trivial over $\mathcal{M}_\Gamma$. More generally, any star-cover transition matrix G_n for which both G_n and G_n^{-1} extend as $\mathcal{M}_\Gamma$ -matrices to D_n defines a trivial $\mathcal{M}_\Gamma$ -bundle.*

Proof. Set $B_0 = I$ on U_0 and $B_n = G_n$ on each D_n . These matrices are invertible over the stated local sheaves and satisfy $B_n = G_n B_0$; their columns are a global frame. For the nilpotent matrices of the first assertion, G_n and its inverse are finite sums of powers of the meromorphic coefficient p_n , so they do extend over D_n . $\square$

There is no requirement that the matrices on different disks have one common support: they are different local coordinate descriptions of a bundle frame, not restrictions of one scalar Hahn function. The common-support constraint enters only when a particular coordinate is shown to glue as a scalar, as in the proof of [theorem 22.4](#).

This also explains why essential protection is a substantial hypothesis. With $p_n = (z - n)^{-1}$, multiplication by a global holomorphic function vanishing at all n can cancel the coefficient poles, and the zero-annihilator proof fails. In the essential case no nonzero meromorphic coefficient can perform that cancellation.

The proof of [proposition 27.1](#) uses nothing about the cover beyond the disjointness of the disks, so it applies verbatim to every puncture-and-disk cover of the plane, in either gluing convention (in Part I’s convention take $B_0 = I$ and $B_a = G_a^{-1}$). The localization manuscript observes that Part I’s hidden and deep examples use pole data, and says that their meromorphic extension is not a model for its resistance mechanism *when* their transitions and inverses extend as in [proposition 27.1](#). The following merge observation checks that they do, and adds the global-divisor report’s plane Picard classes, for which the same check applies.

Corollary 27.2 (Pole-data bundles die over $\mathcal{M}_\Gamma$). *After extension of scalars to $\mathcal{M}_\Gamma$ the following bundles on $\mathbb{C}$ become trivial.*

- (a) The integral bundles of [corollary 10.2](#): the hidden rank-three cocycle (8.2) and the deep cocycles of [theorem 9.2](#).
- (b) The scalar cocycles $\exp(t^{\gamma_n}/(z-n))$ and the determinant-one diagonal cocycles (11.4) in the proof of [theorem 11.4](#).
- (c) In the global-divisor report, every line bundle $\mathcal{L}(a)$ glued by $\text{Exp}_H(a_j/(z-j))$ with a sequence of positive Hahn constants (its `global:eq:transitions` and `global:thm:obstruction`), in particular every explicit plane Picard class $\text{Exp}_H(a_n t^{\gamma_n}/(z-n))$, $a_n \in \mathbb{C}$, of its Picard dichotomy (`global:eq:explicitbundles`, `global:thm:dichotomy`); and the line bundles attached to root-cluster data in its `global:prop:divPic`.

Proof. In each case the transitions live on a puncture-and-disk cover of $\mathbb{C}$, and it suffices, by [proposition 27.1](#) and the remark preceding this corollary, that each transition and its inverse extend across the centre with meromorphic coefficients.

(a) The hidden matrices $I_3 + tE_{12}/(z-n) + t^{1/n}E_{23}$ and the deep matrices $I_r + C_n + a_n E_{12}$ of [lemma 9.1](#), with $a_n = t/(z-n)$ and Hahn-constant C_n , are finite unitriangular matrices whose entries are Hahn constants or constant multiples of $t/(z-n)$. Their inverses are finite geometric sums of the nilpotent part, whose entries are polynomials in those entries. All coefficients are rational functions with poles only at n .

(b) $\exp(\pm t^{\gamma_n}/(z-n)) = \sum_{k \geq 0} (\pm 1)^k t^{k\gamma_n} / (k!(z-n)^k)$ has, at each exponent, a single pole of finite order at n ; the diagonal matrices are built from these.

(c) For a positive Hahn constant a_j , the coefficient of $\text{Exp}_H(\pm a_j/(z-j)) = \sum_k (\pm a_j)^k / (k!(z-j)^k)$ at a fixed exponent receives contributions from only finitely many k ([lemma 3.2](#)), each a complex constant times $(z-j)^{-k}$; so every coefficient is rational with a pole only at j . The explicit Picard classes are the special case of the Hahn constants $a_n t^{\gamma_n}$ with $a_n \in \mathbb{C}$. For root-cluster data the transition is $P_c(z-c)/(z-c)^{m_c} = 1 + \sum_{i < m_c} a_{c,i} (z-c)^{i-m_c}$ with positive Hahn constants $a_{c,i}$, a nonzero element of $\mathcal{M}_\Gamma$ on the disk with meromorphic coefficients; it is a unit of $\mathcal{M}_\Gamma$ there by [lemma 20.3](#). $\square$

What this does and does not say. First, it contradicts nothing. The global-divisor report’s Picard dichotomy is a statement about $\mathcal{I}_\Gamma$ and $\mathcal{H}_\Gamma$ only, and there the classes in (c) are nontrivial for every noncyclic $\Gamma \subseteq \mathbf{No}$, that report’s standing hypothesis. Together with (c) this shows that extension of scalars from $\mathcal{H}_\Gamma$ to $\mathcal{M}_\Gamma$ sends nonzero classes of the plane Picard group to the trivial bundle whenever $\Gamma \subseteq \mathbf{No}$ is noncyclic: in rank one, pole-data classes survive monomial localization and die under meromorphic extension. Second, the corollary concerns the displayed families. It does not assert that every class of $\text{Pic}(\mathbb{C}, \mathcal{H}_\Gamma)$ dies over $\mathcal{M}_\Gamma$, and no statement about the Picard group of $\mathcal{M}_\Gamma$, or about rank-one $\mathcal{M}_\Gamma$ -bundles in general, is made anywhere in this report. Third, for Part I’s examples it settles nothing over $\mathcal{H}_\Gamma$: a proof of their nontriviality would have had to use the holomorphy of gauge coefficients at the punctures, which the argument of [theorem 21.2](#) never uses. Part V answers it by exactly such frames: the examples do not survive monomial localization ([theorem 35.1](#) and [corollary 35.2](#)). Over a compact base the global-divisor report identifies exactly which Hahn line bundles become trivial over its $\mathcal{M}_\Gamma$ (those with vanishing valuation class, by its `global:thm:meromorphic`); no plane analogue is claimed.

27.2 Why this does not contradict ordinary analytic bundle theorems

Ordinary holomorphic bundle theory over the plane has very strong triviality results, within the setting of the Oka–Grauert principle [29]. Those theorems concern finite-rank locally free

modules over $\mathcal{O}$, not sheaves built from infinitely many ordinary coefficient functions with a joint Hahn-support constraint. Our ordinary reductions are trivial. The obstruction is not an ordinary holomorphic obstruction remaining in the reduction; it is the impossibility of fitting infinitely many forced exponents into one well-ordered support. Applying a theorem for coherent $\mathcal{O}$ -modules directly to $\mathcal{H}_\Gamma$ would change the coefficient category and discard exactly the constraint used in [theorem 21.2](#).

Similarly, $\mathcal{M}_\Gamma$ is not a constant field sheaf merely because its sections on each connected domain form a field. Restriction maps are injective but in general not surjective: an essential singularity at an omitted point prevents extension. A global frame requires compatibility across all these restriction maps together with admissible supports. The one-dimensional global section formula over the field $A_{\mathcal{M}_\Gamma}$ is therefore not paradoxical.

27.3 Why determinant and finite tests miss the obstruction

The transition determinants are identically one, and their ordinary reductions are identically the identity. On every finite-puncture subproblem there is the explicit integral frame [\(22.3\)](#). None of these tests detects the global nontriviality. For any positive integer K , the restrictions over a bounded region containing the first K punctures are indistinguishable, at the level of bundle isomorphism, from a trivial bundle. A proof of global nontriviality must use infinite information; here it uses a quantified statement about the whole decreasing profile, not a numerical experiment on a long prefix. This is the localized counterpart of item 4 of [section 30](#).

27.4 Other essential singularities

Proposition 27.3 (Replacing the ordinary exponential). *The global section, morphism and indecomposability theorems, and the necessity direction of the profile classification, remain valid if each h_n is replaced by a holomorphic function on W_n with an essential singularity at n , provided the same h_n is used for both profiles being compared. If each h_n additionally extends holomorphically to $\mathbb{C} \setminus \{n\}$, then the full profile classification and the explicit finite-puncture trivialization theorem remain valid as well.*

Proof. Essential protection follows by the same quotient argument for every nonmeromorphic germ. No other property of h_n is used in [theorems 21.2, 22.4 and 23.1](#) or in the necessity part of [theorem 24.1](#). The sufficiency gauges require the finite sums of the h_n to be holomorphic away from their indicated centres; the stated global extension hypothesis guarantees exactly this property, and it also proves bounded triviality as before. $\square$

The assertion about classification in the proposition includes its sufficiency under the stated global extension hypothesis. Without that extra hypothesis, the necessity argument and the section and morphism theorems still hold on the fixed cover, but a finite global correction needs a separate additive Cousin construction. We do not silently assume such an extension of arbitrary locally specified functions.

Part IV. Effectivity, verification, and boundaries

28 How to compute the invariants, and what computation cannot decide

28.1 A support-aware procedure for Criterion P

For a presented star-cover datum the mathematical procedure is:

1. at each puncture compute the normalized factor by (5.2)–(5.3), or exploit a finite matrix identity such as (8.1) or (9.1);
2. extract the union of the supports of the normalized *polar* coefficients — not the supports of the raw entries, and not just the leading valuations;
3. supply a proof of well-ordering, or a descending-sequence obstruction, for that union. In the admissible case use (6.4)–(6.5) to construct a splitting with a monoid support certificate.

These steps give an exact criterion, not a decision algorithm for an arbitrary black-box infinite family of functions. In general an input description must support a proof of the required infinite order property: checking many exponents or many punctures does not settle it, and [corollary 10.3](#) shows that a finite-puncture computation provably cannot.

28.2 Finite truncations

For finitely generated positive rational exponent monoids every bounded exponent interval is finite, so a sparse dictionary keyed by exact rational exponents suffices for a finite-cutoff implementation of the local recursion, each dictionary value being a matrix of ordinary Laurent polynomials, with π_- keeping the negative powers of u and π_+ the nonnegative ones. For a general higher-rank ordered group a bounded interval can be infinite and no such finite-cutoff assumption is justified: an implementation must either work with a certified finite dependency set or restrict its representation class. This is the computational counterpart of [remark 3.3](#), and [example 17.3](#) is a case where even the admissible answer cannot be enumerated below a fixed bound in finitely many steps.

28.3 What the Criterion M recursion actually depends on

Proposition 28.1 (Finite exponent dependence). *Fix an admissible support E , a period section σ , a target exponent $\gamma \in \langle E \rangle_{\mathbb{N}} \setminus \{0\}$, and the meridian paths. The coefficient Ω_{γ}^{σ} is determined by finitely many exponent-indexed vectors of target coefficients $(m_{d,\delta})_{d \in D}$, $\delta \in E$, and finitely many exponent patterns of ordinary products, iterated integrals, and applications of σ . The set of punctures inside one such vector may still be infinite.*

Proof. Every lower coefficient entering the defect at γ has a weight occurring as a part in a decomposition of γ into members of $\langle E \rangle_{\mathbb{N}}$, and there are finitely many such decompositions and parts by [lemma 3.5](#). Recursively replacing a constructed coefficient by its own defining defect only refines one of these finite exponent-word decompositions; equivalently all expressions are organized by finite rooted trees whose leaves are E -letters summing to γ , and there are finitely many possible leaf words by [lemma 3.2](#) and finitely many tree shapes and partitions for each fixed finite number of

leaves. Applying σ may use an entire element of $\mathbb{C}^D$, so this argument says nothing about finite dependence on puncture coordinates. $\square$

A relative algorithm is therefore explicit: represent the support well enough to list the finitely many words relevant to a requested exponent, compute the lower-weight iterated-integral prediction, subtract it from the target coefficient vector, and apply the chosen ordinary period section. This does not imply that the steps are computable for every set-theoretically specified Hahn series, nor that a non-effective support can be enumerated, nor that an infinite product vector can be supplied to a finite program. For finite puncture sets the period section is explicit and the remaining tasks are ordinary symbolic or numerical integrations.

Proposition 28.2 (Compatibility with enlargement of the workspace). *Let $\iota : \Gamma \hookrightarrow \Gamma'$ be an order-preserving group embedding, with the same complex coefficient data, meridians and period section. For an admissible representation, its normalized realizer in the larger Hahn workspace is obtained from the original realizer by replacing t^γ with $t^{\iota(\gamma)}$, and the same compatibility holds for normalized fundamental matrices and framed comparison gauges.*

Proof. Order embeddings preserve Hahn summability and addition of exponents, and the finite coefficient formulas for products, iterated integrals and ordinary primitives are unchanged, so the transfinite recursion on $\langle E \rangle_{\mathbb{N}}$ maps to the same recursion on $\langle \iota(E) \rangle_{\mathbb{N}}$ with no extra coefficients outside that monoid. Alternatively the transported family satisfies all defining equations and normalizations, so uniqueness in [theorems 13.1, 14.2 and 15.2](#) identifies it with the larger-workspace construction. $\square$

28.4 Why an infinite linear period section cannot be continuous

The choice-dependent section of [lemma 4.5](#) is not merely missing an estimate that could be supplied later. Give $\mathbb{C}^D$ its product topology and give holomorphic one-forms on U the topology of uniform convergence of the coefficient functions on compact subsets, restricted to $\mathcal{L}(D)$. The following is an auxiliary observation, not a claim of novelty for the general functional-analytic phenomenon.

Proposition 28.3 (No continuous linear period right inverse). *If D is infinite there is no continuous complex-linear right inverse $\sigma : \mathbb{C}^D \rightarrow \mathcal{L}(D)$ to P for these topologies. No such continuous linear right inverse exists even if its values are allowed to be arbitrary holomorphic one-forms on U with the specified meridian periods.*

Proof. Choose a closed ordinary disk $K \subset U$ with nonempty interior, write a one-form on U as $f(z) dz$ and put $q_K(f dz) = \sup_{z \in K} |f(z)|$. If σ were continuous and linear, the continuous seminorm $q_K \circ \sigma$ on the product $\mathbb{C}^D$ would be bounded by a constant times a maximum of finitely many coordinate absolute values, so there would be a finite $F \subset D$ and $C_0 > 0$ with

$$q_K(\sigma(v)) \leq C_0 \max_{d \in F} |v_d| \quad (v \in \mathbb{C}^D);$$

this standard product-topology bound also follows by taking a basic finite-coordinate neighbourhood on which the seminorm is less than one and scaling. Choose $d \in D \setminus F$ and take the coordinate vector e_d . The bound implies $q_K(\sigma(e_d)) = 0$, so the coefficient function of $\sigma(e_d)$ vanishes on a disk and the ordinary identity theorem on the connected domain U makes it identically zero; its period around ℓ_d is then zero, contradicting $(P\sigma(e_d))_d = 1$. $\square$

So the algebraic splitting used in the infinite-puncture normal form cannot have product-topology continuity. This does not contradict [theorem 14.2](#), whose coefficient category imposes no continuity, and it does not rule out nonlinear choices with other regularity properties, or useful linear choices on smaller weighted sequence spaces; those are different problems. For finite D the explicit section [\(4.3\)](#) is continuous and the obstruction disappears.

28.5 Part III: a complete criterion that is not semidecidable

[Theorem 24.1](#) is a complete mathematical criterion for isomorphism in its family. It is not an algorithm, and for unrestricted infinite input no algorithm exists.

The input model. An *effective profile* is a program that, on input $n \geq 1$, outputs an exact positive rational α_n , with the promise that the program halts on every input and that $\alpha_n > \alpha_{n+1}$. Fix the ordinary punctures, disks, essential functions h_n of [section 21](#), and a rank $r \geq 2$ once and for all. The only input is thus the exponent profile; there is no hidden oracle for ordinary analytic continuation or for coefficient functions. A semidecision procedure for isomorphism halts with a positive answer exactly on isomorphic pairs of valid inputs and may run forever on nonisomorphic pairs; a semidecision procedure for nonisomorphism reverses the two cases. Correctness is required only on inputs satisfying the promise.

Theorem 28.4 (Neither side is semidecidable). *For the rational binary family [\(25.1\)](#), neither isomorphism nor nonisomorphism of the corresponding rank- r bundles is semidecidable, in any of the three coefficient categories $\mathcal{I}_\Gamma$, $\mathcal{H}_\Gamma$, $\mathcal{M}_\Gamma$. In particular there is no decision algorithm for isomorphism on this promised computable input class.*

Proof. Fix the zero-bit profile $\alpha_n^0 = 1/n$. By [\(25.2\)](#), the bundle for a bit sequence ε is isomorphic to this fixed bundle exactly when $\varepsilon_n = 0$ eventually. For a Turing machine e on a fixed input, define a uniformly computable bit sequence

$$u_n(e) = \begin{cases} 1, & e \text{ halts within its first } n \text{ steps,} \\ 0, & \text{otherwise.} \end{cases}$$

Computing one bit requires only a bounded simulation, so the resulting profile is always a total computable valid input. If e never halts, $u(e)$ is all zero and the two bundles are isomorphic; if e halts, $u(e)$ is eventually one and they are not. A semidecision procedure for isomorphism would therefore semidecide the set of machines that never halt. For the other direction put $v_n(e) = 1 - u_n(e)$: if e halts, this sequence is eventually zero and gives an isomorphic bundle, and if it never halts, it is all one and gives a nonisomorphic bundle. A semidecision procedure for nonisomorphism would again semidecide nonhalting.

The halting set is semidecidable by simulation, whereas its complement is not: if both were semidecidable, dovetailing the two procedures would decide halting, contradicting the classical diagonal argument [\[30\]](#). Both proposed semidecision procedures are therefore impossible. Every profile produced in the reductions satisfies the promise, by the inequality in [example 25.3](#); no undecidable validity check is hidden in the input encoding. $\square$

This is a reduction to a classical undecidability phenomenon, not a new theorem about Turing machines. Its content here is that the exact geometric isomorphism relation already carries that obstruction, in a family with rational exponents and fixed elementary coefficient functions. For

restricted finite or eventually periodic presentations the tail criterion can of course be evaluated by ordinary finite methods. The theorem is the Part III counterpart of item 4 of [section 30](#): there, finite-puncture computation cannot detect triviality; here, even a complete criterion for isomorphism is not semidecidable on a computable input class.

29 The accompanying exact and numerical checks

Four independent check suites were delivered with the four source manuscripts and are preserved unchanged in `code/` and `data/` of this report directory. No suite verifies a theorem; each audits finite identities, and each prints its own scope disclaimer. The facts recorded in this section were confirmed by an independent run of the code, described in [appendix C](#).

Part I: exact symbolic algebra. The script `08-polar-support-matrix-cousin-verify.py` uses Python with SymPy and exact rational arithmetic, with no floating-point comparison in any checked identity. It verifies: the rank-three factors and the missing cross term, including $PH = G$, $\det = 1$, nilpotence of $G - I_3$, the repaired product of [example 8.3](#), the matrix logarithm of [proposition 8.4](#) with its $(1, 3)$ coefficient $-xy/2u$ and the exact discrepancy $xy/2u$ between the naive exponentiated-logarithm factor and the true polar factor; the chain inverse, chain factorization, row-one square-zero property and determinant for ranks 3 through 10 with symbolic c_j , together with the closed exponent formulas (9.3)–(9.4) and their monotonicity in n for ranks 4 through 12 and $n = 2, \dots, 25$; a genuinely noncommuting 2×2 series over the monoid generated by $\{1/2, 2/3\}$ truncated at cutoff 3 (sixteen exponents including zero), on which it runs the actual π_-/π_+ recursion (5.2)–(5.3) and checks $PH = G$ coefficientwise, that polar outputs are negative Laurent and regular outputs holomorphic, holomorphic right-gauge invariance $\text{Pol}(GQ) = \text{Pol}(G)$ with $\text{Reg}(GQ) = \text{Reg}(G)Q$, and a two-sided inverse; and an exact four-puncture splitting with extension of every local frame. The recorded run reports **2,899 exact checks passed** under Python 3.13.5 and SymPy 1.14.0; the independent rerun reproduced the same four PASS lines and the identical count of 2,899 under Python 3.14.4 and SymPy 1.14.0, the version banner being the only differing output line.

Part II: exact algebra plus a coefficient-ODE audit. The script `09-support-monodromy-realization-verify-examples.py` has two components. SymPy checks eight exact algebraic assertions: $N_+^2 = N_-^2 = 0$, $[N_+, N_-] = H$, both target determinants equal one, the exact group commutator (18.7) and its determinant, and tracelessness of the correction. A second component integrates the degree-graded coefficient ODE system over the bivariate degrees $(0, 0), (1, 0), (0, 1), (2, 0), (1, 1), (0, 2)$ with `scipy solve_ivp DOP853` at `rtol` $2 \cdot 10^{-12}$ and `atol` $2 \cdot 10^{-13}$, along explicit meridian paths: from $1/2$ to $1/4$, once positively around the radius- $1/4$ circle about 0, and back; and from $1/2$ to $3/4$, around the radius- $1/4$ circle about 1 from angle π to 3π , and back. It does *not* substitute floating-point surrogates for u or v . The uncorrected mixed term is compared with $\pm CH$ and the corrected transport with the target through degree two. Recorded maximum entrywise errors are $1.17 \cdot 10^{-13}$ and $1.47 \cdot 10^{-13}$ respectively at both meridians; the assertions use a looser absolute tolerance of $2 \cdot 10^{-9}$ so as not to depend on platform-specific last digits. The recorded environment is Python 3.13.5 on Linux with NumPy 2.3.5, SciPy 1.17.0 and SymPy 1.14.0.

Part III: exact finite algebra and exponent samples. The script `10-localization-bundles-verify.py` checks the finite algebra used in Part III with exact SymPy arithmetic and exact

Python fractions. Its dependency pin is `data/10-localization-bundles-requirements.txt` (SymPy 1.14.0). The recorded run used maximum rank six under Python 3.13.5 and SymPy 1.14.0 and is stored twice, byte-identically, as `data/10-localization-bundles-verification.json` and `data/10-localization-bundles-verification.txt`.

Check family	Count
Nilpotent exponential addition, inverse and determinant	18
Monomial diagonal conjugation	12
Finite-puncture gauge identities	6
Centralizer basis identities	21
Intertwiner-space dimensions for ranks one through six	36
Internal-Hom finite exponential identities	9
Tensor-sum Jordan rank profiles	252
The upper-triangular rank-two gauge equation (24.4)	1
Exact rational binary-profile inequalities (25.1)	8,000
Lexicographic non-Archimedean profile samples	4,000
Total exact checks	12,355

All recorded checks passed. The rational inequalities use exact fractions for $n \leq 2000$ and all four bit patterns of consecutive indices; the non-Archimedean samples use exact lexicographic pairs of rationals, a model of $\mathbb{Q}\omega + \mathbb{Q}$, for the profile $\omega - n$ and one translate of it, not floating-point stand-ins for ω . The matrix tests use symbolic identities or exact ranks, not approximate eigenvalues. The script writes nothing unless an output path is given with `--output`, so running it in place does not overwrite the recorded evidence. An independent rerun on a copy, under Python 3.14.4 with SymPy 1.14.0, reproduced the recorded JSON exactly except for its `python_version` field, with the same count of 12,355.

Part V: exact finite algebra. The script `11-pole-localization-verify.py` is a deterministic SymPy program with exact arithmetic; its dependency pin is `data/11-pole-localization-requirements.txt` (SymPy 1.14.0). It checks the finite matrix identities of the rank-two and rank-three constructions (the frame product, determinants and inverses, the extension row and column, the difference of local lifts, the composition of the resolution maps, the polar factor (8.3), the reduction (35.1) and the frame equation (33.1) for (35.3)); symmetric-power identities in degrees one through six; finite-node pole cancellation with simple and multiple poles; the cardinal-sine values and derivatives, including the leading root balance of theorem 38.2; and finite support sanity checks, including the balanced budget 3/4.

Check family	Count
Matrix identities	17
Symmetric-power representation	36
Pole cancellation	48
Cardinal-sine formulas	106
Finite support sanity	8
Total exact checks	215

The recorded run, `data/11-pole-localization-verification_results.json`, passed all 215 under Python 3.13.5 and SymPy 1.14.0. An independent rerun on a copy, under Python 3.14.4 with SymPy 1.14.0, exited zero and reproduced the recorded JSON except for its `python_version` field.

By default the script writes `verification_results.json` beside itself; in this layout an in-place run would create that file under `code/` without touching the record under `data/`, but reruns should use a copy and `--output`, as in [appendix C](#). The suite does not check [corollary 35.2](#); the merge checked its matrix identities symbolically for $3 \leq r \leq 8$, with symbolic row-one entries and symbolic f and $\mathcal{S}$, in a check that is not shipped with this report.

A warning about rerunning the Part II script. That script *writes its own evidence*: it emits `verification_results.json` beside itself, via `Path(__file__).with_name`. In the original archive layout this could overwrite the recorded output beside the script. In the maintained layout it creates or overwrites an output under `code/`, while the separately named record under `data/` is unchanged. Reruns should use a copy outside the report directory, as in [appendix C](#); the delivered `09-support-monodromy-realization-verification-results.json` in `data/` is the original record.

What the checks do not establish. The Part I checks audit algebraic signs, multiplication order and finite coefficient identities; they do not prove the support lemma, Cartan’s theorem B, any infinite descending obstruction, an analytic existence theorem, or novelty. The Part II numerical results are observed floating-point errors, not rigorous enclosures, and test only a finite bivariate degree-two identity; the exact checks are algebraic identities in two formal variables. Neither suite tests an arbitrary Hahn field, justifies a transfinite recursion, proves any support set well ordered, verifies anything in a proof assistant, or establishes that the results are unpublished. The Part III checks likewise cannot establish the well-ordering contradiction for all indices, prove the identity theorem for meromorphic functions, certify the sheaf construction, the essential-singularity theorem, descent or the classification, verify the undecidability statement, or establish historical novelty; they are a reproducible algebraic audit of finite steps. The Part V checks do not prove that an infinite support is well ordered or that a forbidden descending support cannot occur, and they do not check the existence part of Weierstrass’s theorem, sheaf cohomology or the transfinite implicit recursion; no numerical experiment with a large real parameter is used as evidence for a surreal identity. No implication in this report depends on the outcome of any script: the general proofs are the arguments in the body.

30 Exactly what is not claimed

The following list merges the honest limitations stated by all four source manuscripts. Every one of them survives this merge, and none is weakened. Items [1](#) through [11](#) come from Part I’s scope, items [12](#) through [21](#) from Part II’s, items [23](#) through [28](#) from Part III’s, and items [30](#) through [38](#) from Part V’s; item [39](#) records what the merge added to Part V. The pole-localization manuscript’s generic limitations are added to items [1](#), [9](#), [10](#) and [11](#). The localization manuscript’s generic limitations (refereeing, priority, credits, code, formalization, and the Müller–Strohmaier distinction) are added to items [1](#), [8](#), [9](#), [10](#), [11](#) and [21](#) rather than repeated, and items [2](#) and [6](#) are re-scoped to Part III’s results. Item [29](#) is a caution added by the merge. The boundary note at the end records the one place where the parts make different standing assumptions.

1. Nothing here is refereed or proof-assistant verified. All three source manuscripts are self-described AI-assisted research drafts and say so on their title pages and in their READMEs; the localization manuscript’s provenance record (`data/10-localization-bundles-PROVENANCE.json`) also lists peer review and proof-assistant verification as false. The pole-localization

manuscript describes itself as an unrefereed AI-assisted research manuscript whose proofs have not been formalized in Lean or independently reviewed, and its source audit claims no repository build, axiom audit, Wolfram check or automated prover check of the infinite arguments; its layout record `data/11-pole-localization-build_report.json` lists formal proof-assistant verification and independent peer review as false.

2. Nontriviality in Part I is proved *only* over the integral sheaf $\mathcal{I}_\Gamma$. We explicitly do not claim that the examples survive after all monomials are inverted, that is over $\mathcal{H}_\Gamma$, because positive monomials become units there, no reduction homomorphism $\mathcal{H}_\Gamma \rightarrow \mathcal{O}$ exists, and the framing-normalization argument of [proposition 10.1](#) cannot be repeated. This was listed as an open question, not as a believed consequence, and Part V has since answered it negatively: the hidden and deep examples are trivial over $\mathcal{H}_\Gamma$ ([theorem 35.1](#) and [corollary 35.2](#)), so the declined claim is in fact false. Part III does not change this for Part I’s examples: it proves nontriviality over $\mathcal{H}_\Gamma$ and over $\mathcal{M}_\Gamma$ for a different, essential-singularity family ([corollary 22.5](#)), and it shows that Part I’s examples become trivial over $\mathcal{M}_\Gamma$ ([corollary 27.2](#)), which decides nothing over $\mathcal{H}_\Gamma$. In particular neither criterion of this report is restated over a coefficient field in which negative-valuation changes of frame are admitted.
3. The exact criterion of [theorem 6.2](#) is only for the puncture-and-disk (star) cover, where each overlap has a distinguished isolated Laurent singularity. For general covers only the *sufficient* globally support-admissible theorem [theorem 11.1](#) is available, and its hypothesis ([11.1](#)) is a condition on a *presentation*, not an invariant necessary condition: regular local gauges can destroy it without destroying triviality.
4. Neither criterion is a decision algorithm for a black-box infinite family. An input description must support a proof of the infinite order property; checking many exponents or many punctures does not settle it, and [corollary 10.3](#) shows that finite-puncture computation provably cannot detect the Part I examples.
5. Finite-cutoff implementation is justified only for finitely generated positive rational exponent monoids. For a general higher-rank ordered group a bounded exponent interval can be infinite.
6. Part I decides triviality rather than classifying all bundle isomorphism classes. [Proposition 6.3](#) describes all solutions once a splitting exists, not the nonsplit objects. Part II does classify framed positive gauge classes by their admissible monodromy representations ([theorem 15.2](#)). Part III classifies one explicit family and nothing beyond it; see item [26](#).
7. The sharp star-cover invariant does not generalize to positive-dimensional analytic singular loci: there is no one-variable Laurent projection there, and an ideal-theoretic or cohomological replacement may have nontrivial relations.
8. Credited as established mathematics and not new: the Neumann support calculus; the ordinary Laurent, Mittag–Leffler, Cartan B and Behnke–Stein inputs; and algebraic noncommutative Birkhoff factorization in complete filtered algebras. The local factorization is reproved only for explicit arbitrary-ordered-group support control. The citation of [\[15\]](#) is to the *algebra* of recursive Birkhoff factorization and to the necessity of commutator corrections; nothing in this report imports, endorses or reinterprets any application of that algebra, and no assertion here is physical, empirical or applied in any way. Likewise “positive” never means Hermitian positivity, and the normalized polar factorization is *not* the Hilbert-space polar decomposition. For Part III: the Jordan tensor identity ([lemma 26.2](#)) is classical linear algebra, Atiyah’s elliptic-curve bundles [\[28\]](#) are a precedent for its pattern, and the tensor pattern is not evidence of novelty; the undecidability statement is a reduction to the classical halting theorem [\[30\]](#); the cyclic integral direction of [theorem 25.1](#) is recalled and proved, not claimed as a new Stein theorem; and the global-divisor report’s scalar Picard dichotomy and Part I’s integral cyclic rigidity are antecedents, not Part III results.

9. The literature checks were targeted and are explicitly not certifications of priority. Broad searches that returned irrelevant material were not treated as evidence of absence, a repository code-search response marked incomplete was not used as evidence of absence, and no named published conjecture is claimed solved. The localization manuscript’s novelty assessment is likewise bounded by what it inspected — for this report, its README and two line ranges of its article; for the global-divisor report, a truncated README — and is not a proof of historical priority. The pole-localization manuscript’s targeted search (Hahn holomorphic coefficient rings, monomial localization of bundles, Picard groups, omnific normal forms) found no matching explicit pole-localization theorem; it is not exhaustive across generalized-series, Cousin-problem or ringed-space terminology, and the manuscript claims a proved argument as presented, not a certified breakthrough or a resolved named conjecture.
10. The companion code checks exact finite algebra, finitely many exponent samples, and one finite numerical example only; see the final paragraph of [section 29](#). The Part V suite’s 215 exact checks do not verify infinite supports, Weierstrass existence, sheaf cohomology or the transfinite implicit recursion.
11. Formal verification is future work. The local support recursion, the chain identity and the necessity directions appear well suited to formalization with Hahn series and finite matrix algebra, but the global sufficiency proofs additionally need a formal ordinary Mittag–Leffler theorem or a suitable sheaf-cohomology interface. For Part III the localization manuscript proposes a formalization boundary: first an abstract kernel with restriction maps, distinguished local extensions, and a predicate saying that multiplication of h_n by a nonzero global leading coefficient cannot extend at n , to be instantiated later with holomorphic and meromorphic germs; then three milestones — the descending-support contradiction and its preservation under ordered group embeddings; finite nilpotent exponentials, the Jordan induction and $B \mapsto MB - BN$; and bundle descent with the intrinsic global-section flag, so that arbitrary isomorphisms, not only flag-preserving gauges, satisfy the classification. Omitting the last milestone would weaken the classification to a framed statement that is not what is proved. For Part V the pole-localization manuscript proposes four layers: the support reversal $\alpha_n \mapsto -\alpha_n$, translated-support unions and the fact that ordinary evaluation creates no exponent; the determinant-one rank-two construction from the five cancellation hypotheses of [lemma 34.1](#), as pure ring algebra; the ordinary finite-pole cancellation lemma with Weierstrass existence as a declared analytic dependency; and the cocycle equivalences, the nonsplitting argument and the implicit recursion. The finite rank-three identity is a small first target, but a check of that identity alone would not verify the localization theorem. No Lean implementation is included or claimed, and the repository’s build status is not evidence that any of these theorems has been checked. No script is a substitute for such a formalization.
12. Positivity of exponents is essential throughout Part II. There is no treatment of nontrivial exponent-zero monodromy, of arbitrary ordinary residues with resonances, or of arbitrary algebraic groups.
13. [Theorem 14.2](#) is not an unrestricted classical Riemann–Hilbert theorem and not a Riemann–Hilbert correspondence on a general surcomplex domain; likewise [theorem 6.2](#) is not the unrestricted classical Cousin theorem over $\mathcal{H}_\Gamma$. The base, the universal cover and the path category are *ordinary* complex by design, not provisional notation for a fine-topological construction.
14. There is no fine-topological theory of surcomplex loops here. Convergence of Hahn partial sums in the fine topology is never asserted, and [example 17.3](#) shows that partial sums can fail to converge even in the valuation topology. The surcomplex reading is an interpretation of a Hahn model via $t^\gamma = \omega^{-\gamma}$ and does not assert an ambient surcomplex manifold.

15. The infinite-puncture period section is obtained by a vector-space splitting using the axiom of choice. It is not continuous — [proposition 28.3](#) proves that impossible. No norm estimate or effective procedure is supplied; only an ordinary complex-linear splitting is claimed, with no computability assertion about a specified representation of the infinite input data.
16. For infinitely many punctures, no regularity or growth condition at infinity is imposed on realizers. This is deliberate, and the freedom to use arbitrary ordinary entire correction terms at each exponent is important to the existence proof.
17. Part II does not assume that the global-support assignment is a sheaf for arbitrary covers, uses no Cartan theorem and no sheaf cohomology for a Hahn coefficient sheaf, and relies on no companion claim about Hahn Noetherianity or a Hahn Nullstellensatz.
18. Deformations of a *nontrivial ordinary connection* are open. The linearized operator there becomes a twisted period map with values in an adjoint local system rather than the scalar period map P , and the triangular proof of [lemma 13.2](#) supplies no right inverse for it. This is not to be confused with [theorem 11.2](#), which deforms a nontrivial ordinary *vector bundle* on a Stein base and does have its cohomological input available.
19. Higher-dimensional bases are open for Part II: flatness is automatic only for holomorphic one-forms on a curve, a proposed correction must satisfy the nonlinear flatness equation as well as the period conditions, and the result cannot be inferred by changing z to a tuple.
20. The Part II numerical checks are observed floating-point errors, not rigorous enclosures, and test only a finite bivariate degree-two identity.
21. The Hahn-meromorphic category of Müller–Strohmaier [31] is explicitly *not* identified with ours: there the parameter is tied to an expansion near zero on a logarithmic cover, whereas here t is an independent Hahn parameter and z the ordinary analytic variable. The localization manuscript refuses the same identification for the meromorphic sheaf $\mathcal{M}_\Gamma$ of Part III.
22. No theorem of either part is claimed to be an instance of a common generalization proved here. The two criteria are two theorems with two proofs ([section 12](#)); they share only [lemmas 3.2](#) and [3.4](#) and an ordinary Mittag–Leffler input.
23. The localization-resistant bundles of Part III form one explicit essential-singularity family. Nothing in Part III settles whether Part I’s pole-only hidden and deep examples survive monomial localization; Part V settles it negatively ([theorem 35.1](#) and [corollary 35.2](#)). The family shows that allowing all negative exponents, all ordinary meromorphic coefficients, or all exponents of a larger ordered workspace need not remove the obstruction. It does not say that these extensions never remove other obstructions: [example 24.4](#) and [proposition 27.1](#) show that they can identify classes or trivialize other families.
24. The theorems of Part III concern a specified category: the ordinary plane as base, a set-sized exponent group, coefficient functions on a common ordinary domain, and finite-dimensional complex nilpotent matrices. These restrictions are part of the statements, not technicalities to be removed without proof. $\mathcal{M}_\Gamma$ is one specified inclusion of sheaves; it is not identified with every notion of surreal or Hahn meromorphic function, and it is not a category allowing essential singularities at every point of the base. Nothing is claimed about arbitrary fine-topological bundles on the surcomplex class, or about a sheaf on the proper class $\mathbf{No}[i]$ with its fine topology; the functions $\exp(1/(z - n))$ are ordinary complex functions. Ordinary Oka–Grauert triviality theorems do not apply, since the coefficient ring is changed, and nothing here contradicts them. The existence of a decreasing profile in a noncyclic group uses ordinary countable choice, and the continuum count of [theorem 25.2](#) uses cardinal arithmetic with choice.
25. The matrix exponentials $\exp(c_n N)$ and all gauges of Part III are finite polynomials, because N is nilpotent; they are not limits, and no convergence or infinite matrix-exponential assertion occurs. The local family (c_n) is not a global Hahn sum, the finite-puncture trivializations of

- theorem 22.3 are not a compatible exhaustion-limit frame, and the additive quotient $Q_{\mathcal{B}}$ of corollary 21.4 is not identified with derived sheaf cohomology. No norm on the surcomplex class, no real-valued valuation for an arbitrary-rank group, and no limit of bounded-domain trivializations is used.
26. The classification of Part III is for the profile family and for the image of the nilpotent-representation functor only. It compares bundles with the same fixed punctures and the same fixed functions h_n , by isomorphisms covering the identity of the base; reparametrizations of the base and permutations of punctures are excluded. No Krull–Schmidt theorem for the category of all Hahn bundles is invoked. There is no classification of all finite-rank Hahn bundles, no assertion that every unipotent bundle is one of these, no calculation of the whole nonabelian first cohomology set, no general theorem that global sections commute with scalar extension, no computation of extension groups between these bundles, no description of their connections, and no claim about differential equations driven by the Berarducci–Mantova derivation.
 27. Theorem 28.4 is a reduction to the classical halting problem, not a new theorem about Turing machines. It concerns the promised computable input class of rational binary profiles, with fixed punctures, fixed h_n and fixed rank; for restricted finite or eventually periodic presentations the tail criterion is decidable by ordinary finite methods. It separates a complete mathematical criterion from an algorithm for unrestricted infinite input.
 28. In proposition 27.3 the sufficiency half of the classification and the finite-puncture trivialization need each h_n to extend holomorphically to $\mathbb{C} \setminus \{n\}$. Without that hypothesis only the necessity direction and the section, morphism and indecomposability theorems are claimed, and no extension of arbitrary locally specified functions is assumed.
 29. Corollary 27.2 is about the displayed pole-data families only. It does not assert that every class of $\text{Pic}(\mathbb{C}, \mathcal{H}_{\Gamma})$ becomes trivial over $\mathcal{M}_{\Gamma}$, it makes no statement about the Picard group of $\mathcal{M}_{\Gamma}$ or about rank-one $\mathcal{M}_{\Gamma}$ -bundles in general, and it decides nothing about Part I’s examples over $\mathcal{H}_{\Gamma}$ (Part V decides them).
 30. Part V’s localization theorems concern descending-scale cocycles $\exp(t^{\alpha_n} p_n N)$ with finite-order poles and one fixed nilpotent N , the hidden example, and (by the merge) the deep examples. They settle the printed repository question, not every pole-only matrix cocycle: there is no classification of cocycles with varying or noncommuting nilpotent directions, no exact pole-only criterion after localization, and nothing about essential singularities, for which the construction is unavailable; Part III’s results are unaffected. The frames have holomorphic coefficients; meromorphic coefficients are never used as a substitute for holomorphic ones.
 31. The bounds $-3/4$, $-(2r-3)/4$, $-\alpha_1$ and $-(\nu-1)\alpha_1$ are proved upper bounds on the negative budget, not claims of optimality. Corollary 34.5 is an infimum statement: the infimum is 0 and is not attained, and no least positive budget is asserted. Inverting one monomial is distinguished from full Laurent localization.
 32. The Picard decomposition of proposition 36.1 is elementary unit-group algebra, printed in the global-divisor report, not a new cohomology theorem; $H^1(\mathbb{C}, \mathcal{J}_{\Gamma})$ is not computed completely. The persistence of pole-data line classes over $\mathcal{H}_{\Gamma}$ is the global-divisor report’s for simple poles.
 33. There is no classification of all self-extensions or of all vector bundles. Ext^1 in theorem 36.4 is taken in sheaves of $\mathcal{H}_{\Gamma}$ -modules, not in modules over $A_{\mathcal{H}_{\Gamma}}$, and the free-middle classes are not claimed to form a subspace outside the displayed family.
 34. Omnific values are claimed only at ordinary real arguments: $\mathcal{S}$ is not claimed to give an omnific-valued map at every surreal argument, or a function continuous for any topology on $\mathbf{No}$. Theorem 37.1(iii) concerns simultaneous common-domain interpolation, not field inversion.
 35. Taylor prolongation is defined and used only at finite surcomplex arguments; no value, zero or zero theorem at infinite arguments is asserted, and $\mathcal{S}$ is not a strongly entire series. Theorem 38.2

- is complete on finite halos only. [Theorem 37.3](#) uses “Bézout” in its literal equation sense; it is not a classical corona theorem for bounded functions and says nothing about the ring of strongly entire series.
36. The ideal statements (properness, non-Bézout, regular sequence, projective dimensions, Tor, non-flatness, ultrafilter maximal ideals) concern the global ring $A_{\mathcal{H}_\Delta}$, not sheaves of modules. The ultrafilter ideals are not claimed to exhaust the maximal ideals above $\mathfrak{a}$, and no description of the maximal spectrum is claimed.
 37. Credited to the literature and not new: the Weierstrass, Mittag–Leffler and discrete-interpolation theorems (cited by the pole-localization manuscript to course notes [21], and by the merge to [22]); the Hahn support calculus [10]; symmetric powers of SL_2 and their determinant; the implicit root lemma, which is not a new Hensel theorem; the interpolation criterion [proposition 37.2](#), a standard-style consequence (`global:cor:interp`) and not a priority claim. No Oka principle is transferred to the Hahn sheaf; [23] is cited for the ordinary context only. [11] is used for normal-form conventions only, with no factorization or Gonshor-type claim.
 38. The pole-localization manuscript audited this report only, on a branch that was changing during its inspection; it fixed the version at commit 0865f04 (an earlier tree response at b895e86 was not used for provenance). It consulted the Wikipedia overview of surreal numbers for orientation only, not to certify conventions or novelty. Its novelty statements are bounded by that audit; the overlaps with the global-divisor and analytic-geometry reports were found by the merge.
 39. [Corollary 35.2](#), the consequences (a)–(d) in [section 35](#), the credits to other reports in Part V, and the question on the diagonal examples in [section 31](#) are added by the merge, not stated by the pole-localization manuscript. The merge’s symbolic check of [corollary 35.2](#) is not shipped, and the corollary rests on its written proof. The diagonal examples (11.4) are not decided over $\mathcal{H}_\Gamma$.

Boundary note on the sheaf convention. As recorded in [section 2.2](#), Part I proves and uses the sheaf property of the integral common-domain assignment on the ordinary base, while Part II deliberately assumes nothing of the kind. Both are correct as stated, the difference is one of standing hypotheses rather than of fact, and no theorem of Part II is strengthened here by importing Part I’s convention. Part III follows Part I’s convention, for $\mathcal{M}_\Gamma$ as well ([lemma 20.2](#)).

31 Questions exposed by the results

Monomial localization (answered for the printed examples; re-scoped). Does $\mathcal{E}_G \otimes_{\mathcal{I}_\Gamma} \mathcal{H}_\Gamma$ remain nontrivial for the explicit hidden and deep examples? A proof would need control of general Hahn matrix gauges, not just positive ones, and the ordinary reduction normalization is unavailable after localization. Parts I–III left this question open. Part V answers it: *no*. Both families become trivial over $\mathcal{I}_\mathbb{Q}[t^{-1}]$, by determinant-one frames with valuation bounds $-3/4$ and $-(2r-3)/4$ ([theorem 35.1](#), and [corollary 35.2](#), added by the merge), and so does every descending-scale cocycle with finite-order poles and one fixed nilpotent direction ([theorem 34.6](#)). The re-scoped remainder is stated in the next three paragraphs. What Part III, from the localization manuscript, had settled around the question is the following. (a) The broader existence question has a positive answer: there are determinant-one bundles with trivial ordinary reduction that remain nontrivial, indecomposable and not stably trivial over $\mathcal{H}_\Gamma$, and even over $\mathcal{M}_\Gamma = \mathrm{Mer}((t^\Gamma))$ ([corollary 22.5](#) and [theorem 23.3](#)); they use essential singularities, not the pole data of the printed examples. (b) The printed examples themselves, and the diagonal counterexamples of [theorem 11.4](#), become trivial

over $\mathcal{M}_\Gamma$ ([corollary 27.2](#)); so a proof of their nontriviality over $\mathcal{H}_\Gamma$ must use the holomorphy of the gauge coefficients at the punctures, which the essential-singularity argument never uses. (c) In rank one the gap between $\mathcal{H}_\Gamma$ and $\mathcal{M}_\Gamma$ is real: for noncyclic $\Gamma \subseteq \mathbf{No}$ the global-divisor report’s pole-data line bundles are nontrivial over $\mathcal{H}_\Gamma$ and trivial over $\mathcal{M}_\Gamma$. As the localization manuscript puts it, a complete answer for the printed examples must account for holomorphic zeros, possible negative-valuation gauges, and the lack of an ordinary reduction map after localization. Part V’s answer uses exactly these: an entire function vanishing at the punctures and a negative-valuation frame.

Varying nilpotent directions (Part V). For finite-pole data p_n and a descending positive profile, when does $\exp(t^{\alpha_n} p_n N_n)$ with puncture-dependent nilpotent N_n become trivial over $\mathcal{H}_\Gamma$? Does simultaneous upper triangularity suffice, and what obstruction appears when the directions run through infinitely many flags? A simultaneous flag permits induction on rank, but the induction must control the supports of the extension corrections, and the failure of global Bézout in [theorem 37.3](#) warns that a naive extension-by-extension argument may not suffice.

An exact pole-only criterion after localization (Part V). Is there a necessary and sufficient invariant for Laurent triviality of a pole-only positive cocycle, computable from its normalized polar factors, and can it be split into finite-dimensional ordinary pole data and a well-ordering condition stable under Laurent gauges? Criterion P is not such an invariant: the hidden example fails it and is Laurent-trivial. A replacement must see the interpolation cancellation of [theorem 34.3](#) while still detecting essential or genuinely noncommutative obstructions; a uniform nilpotency class with finitely many ordinary directions is a natural first case.

The diagonal determinant-one examples over $\mathcal{H}_\Gamma$ (raised by the merge). The cocycles ([11.4](#)) are nontrivial over $\mathcal{I}_\Gamma$ and trivial over $\mathcal{M}_\Gamma$ ([corollary 27.2](#)), and their two line-bundle summands stay nontrivial over $\mathcal{H}_\Gamma$ ([corollary 36.2](#)). Is their determinant-one direct sum trivial over $\mathcal{H}_\Gamma$? Part V’s construction applies to unipotent cocycles only, and neither the pole-localization manuscript nor this merge decides this case.

The exact denominator budget (Part V). For a Laurent-trivial integral cocycle, determine the set of $b \geq 0$ for which frames and inverse frames can be chosen with valuation at least $-b$: its lower cut, and whether the infimum is attained. What is it for the hidden example, where $3/4$ is an upper bound? For the profiles of [corollary 34.5](#) the infimum is 0 and not attained. At higher rank of Γ the answer is an ordered cut rather than a real number, and inverting one monomial should be distinguished from full localization.

Arbitrary-cover normal forms. The polar support is exact on the puncture-and-disk cover because every overlap has a distinguished isolated Laurent singularity. An equally explicit invariant for general covers would require a substitute for that canonical local negative factor; [theorem 11.1](#) does not provide one.

Classification beyond triviality (partly answered). Deciding whether a positive star-cover cocycle represents the trivial integral bundle is not classifying all isomorphism classes, which would require understanding the action of global and local gauges on *pairs* of polar families. Part III answers this for one explicit family only: for the essential-singularity profile bundles, integral

isomorphism is eventual equality of profiles and localized isomorphism is eventual equality up to one translation ([theorem 24.1](#)), and within one profile the isomorphism type is the Jordan type ([corollary 23.5](#)). The general question remains open, including the classification of Part I’s examples over $\mathcal{I}_T$; over $\mathcal{H}_T$ they are trivial (Part V).

The full extension category. The morphisms between the nilpotent bundles of Part III are known exactly, but the extension groups between them are not computed. Do extensions inside the common-domain Hahn category admit a useful decomposition into ordinary analytic data, support-tail data and finite nilpotent representation data? The zero-annihilator class of [corollary 21.4](#) supplies an explicit submodule of an additive obstruction group, not a classification of that group.

Several independent descending profiles. One can use different profiles at different super-diagonal entries. The resulting transitions need not be exponentials of one fixed nilpotent matrix. Do their morphism equations realize representations of several noncommuting operators? No such realization, and no claim of a wild classification problem, is proved here.

Other ordinary bases and growth restrictions. An open Riemann surface may support analogous escaping puncture families, whereas a compact surface cannot contain an infinite closed discrete subset. Behaviour under coefficient growth bounds at infinity may also change, because the finite and infinite correction spaces would be different. Any extension of Part III should state both the ordinary analytic domain and its allowed coefficient class before asserting a Hahn gluing theorem.

A common generalization of the two criteria. Both criteria say that a union of supports must be well ordered, over different objects ([section 12](#)). Is there a single statement, in some category of positive Hahn objects on a punctured base with a distinguished local normalization, of which both are instances? We do not know, and nothing in this report should be read as evidence that the naive common statement — taking the union over raw local data — is true: [theorem 8.2](#) and [example 8.3](#) refute it for Criterion P.

Deformations of a nontrivial ordinary connection. Fix an ordinary connection Ω_0 and a representation reducing to its monodromy rather than to the identity. The first correction operator is then a twisted period map into the adjoint local system. A precise extension should specify when that map has a right inverse within a chosen pole class, and should track local resonances.

Controlled growth and effective infinite data. For a prescribed weight or growth condition on residues at D , identify spaces on which a useful continuous or computable period section exists. The exact question is to characterize the residue sequence spaces admitted by the chosen coefficient growth class, and then to prove that nonlinear monodromy corrections stay inside them; arbitrary algebraic splittings need not preserve any such structure.

Higher-dimensional bases. For Stein manifolds of dimension greater than one the support-controlled rigidity theorems still apply, but the sharp star-cover invariant does not generalize, and for connections the nonlinear flatness equation enters. Both directions are open.

Free-middle extension classes (Part V). Characterize the classes in $\text{Ext}_{\mathcal{H}_\Gamma}^1(\mathcal{H}_\Gamma, \mathcal{H}_\Gamma)$ whose middle bundle is free. Is that set closed under Baer sum outside the family of [theorem 36.4](#), and how is it read from global columns that admit local but no global unimodular completion? The image of $\mathbb{C}^{\mathbb{N}}/\mathbb{C}^{(\mathbb{N})}$ there is a subspace of free-middle classes, but the whole set need not be a subspace.

Maximal ideals beyond finite-halo evaluation (Part V). Describe the maximal ideals of $A_{\mathcal{H}_\Gamma}$ above $(\sin \pi z, \mathcal{S})$. Are they parametrized by support cuts, by ultrafilters on the nodes, or by both, and which parameters survive ordered extension of the value group? Do the ideals of [theorem 37.5](#) exhaust them? A first test is the quotient by f , whose realized value sequences are those with well-ordered combined support ([proposition 37.2](#); compare `analytic:thm:fixed-quotient`).

A coefficient enlargement with controlled reciprocals (Part V). Is there a useful enlargement of the common-domain families that contains the reciprocal data $t^{1/n}$ of [theorem 37.1](#) without invalidating ordinary evaluation, ring operations and Taylor prolongation? Any proposal must state how products and coefficient sums are defined; a sheaf of locally varying support bounds or a sequence-valued target solves a different problem, and the difference should be explicit.

Mixed poles and essential singularities (Part V). For $\exp(t^{\alpha_n} p_n N)$ in which some p_n have essential singularities, find a sharp condition for Laurent triviality, perhaps through a finite-rank measure of the principal-part space interpolating between [theorem 34.6](#) and Part III. The failure of one particular frame is not a proof that every frame fails; finitely many essential punctures, and singular germs obeying fixed finite-dimensional relations, are natural intermediate cases.

Other ordinary bases for pole localization (Part V). Extend [theorem 34.3](#) from the plane to open Riemann surfaces, tracking the divisor class of the function f and the constant-sheaf summand of [\(36.1\)](#). A compact base has no infinite closed discrete puncture set, so there the plane statement cannot simply be repeated (compare the paragraph on other ordinary bases above).

Omnific-valued coefficient families (Part V). Classify tractable subclasses of $\mathcal{O}(\mathbb{C})((t^{\mathbb{Q}}))$ whose values at all ordinary real arguments are omnific integers, and study their finite-halo zero sets. Which infinitesimal displacements can be realized with a simple interpolation formula? The normal-form condition constrains exponents and the constant coefficient, while the zeros depend on the leading ordinary coefficient after a monomial normalization; a theory should relate the two without conflating ordinary real arguments with all surreal ones.

Formal verification. See item 11 of [section 30](#).

32 Conclusion

In the positive integral Hahn category over a set-sized ordered value group, both the multiplicative gluing problem and the inverse monodromy problem have exact answers, and in both the answer is a well-ordering condition on a union of Hahn supports. They are not the same condition.

For gluing, the correct obstruction is not the support of the input entries, nor the support of their raw principal parts, nor a determinant. One must first remove the holomorphic factors in the correct noncommutative order; the resulting normalized polar support is then an exact necessary and

sufficient criterion on a puncture-and-disk cover. The explicit examples show why that normalization is consequential: a family with one singular input exponent can force a descending support at a hidden matrix entry; adding a cross term can repair the family while worsening its raw singular support; and the same phenomenon can be postponed to the final central layer of upper-unitriangular groups of arbitrarily large finite rank, yielding nontrivial determinant-one integral Hahn bundles with trivial ordinary reduction that are trivial on every bounded plane domain. At the opposite boundary, one globally well-ordered support certificate restores Stein rigidity in every finite rank; cyclic value groups supply such a certificate automatically and noncyclic ones do not.

For monodromy, the correct obstruction is failure of well-ordering of the raw support union of the prescribed matrices. This condition is an invariant of the representation. Realizers can be taken coefficientwise logarithmic, a fixed linear period section selects a unique normalized one, framed positive gauges classify connections with equal monodromy, and within the logarithmic class the comparison gauge extends across every puncture. The order of the value group may have several ranks, the singularity set may be infinite, and the monodromy may be noncommutative; none of this requires a false topological limiting argument.

Past the integral boundary, an essential singularity does what a pole cannot. Placed behind a strictly decreasing sequence of positive exponents, it forces a descending support that no nonzero meromorphic multiplier can cancel. The resulting determinant-one bundles exist in every finite rank, are indecomposable and not stably trivial even over coefficientwise meromorphic Hahn coefficients, and are nevertheless trivial over every bounded ordinary region. Their isomorphism invariant is exactly the eventual exponent profile in the integral category, and that profile modulo one eventual translation after localization; it is independent of valuation rank and survives every ordered exponent enlargement. Pole data, by contrast, die over the meromorphic sheaf, so this argument cannot reach Part I's own examples; Part V shows that they already die over $\mathcal{H}_\Gamma$. Here finite-dimensional linear algebra and ordinary local analytic solvability both behave perfectly, while the single global well-ordered support required by a Hahn section still does not exist.

What Parts I and II separate cleanly is ordinary analytic solvability, noncommutative compatibility, and generalized-series admissibility. The analytic work is done coefficientwise on the ordinary base; the Hahn support calculus is exactly what makes those ordinary constructions coexist in one surcomplex workspace; and the noncommutative correction — the cross term $-a_-c_+$ in Part I, the mixed commutator $-uvCH(\eta_0 - \eta_1)$ in Part II — is what the abelianization cannot see.

Part V shows how thin the boundary between the integral and the localized category is for pole data. A single entire function vanishing at the punctures, with interpolation coefficients placed at the admissible exponents $-\alpha_n$, turns a forbidden descending positive support into an admissible increasing negative one, and one monomial localization then kills every descending-scale pole cocycle with a fixed nilpotent direction, Part I's hidden and deep examples included. The same global support information does not disappear: it persists in rank one, in the filtration of a free rank-two bundle, in a proper locally unit ideal of projective dimension two, and in the infinitesimally displaced zeros of an omnific-valued sine family.

Part V. Pole localization and omnific interpolation

33 The printed examples after monomial localization: question, source and conventions

33.1 What this part answers

Part I proves its hidden and deep examples nontrivial over the integral sheaf $\mathcal{I}_\Gamma$ only, and until this part was added the report asked whether they stay nontrivial after monomial localization, over $\mathcal{H}_\Gamma$ (section 10.3, item 2 of section 30, and the first question of section 31). Part III answered a broader existence question with a different, essential-singularity family and left the printed examples open. This part, built from a fourth manuscript, answers the printed question, and the answer is *negative*:

- the hidden rank-three example (8.2) of theorem 8.2 becomes trivial after the single monomial t is inverted, by explicit determinant-one frames whose entries and inverse entries all have valuation at least $-3/4$ (theorem 35.1);
- the deep examples of theorem 9.2, $r \geq 4$, become trivial in the same way, with valuation bound $-(2r - 3)/4$ (corollary 35.2). This case is not treated in the source manuscript; it is added by the merge, with a complete proof;
- more generally, every descending-scale cocycle $\exp(t^{\alpha_n} p_n N)$ with finite-order poles p_n and one fixed nilpotent matrix N is integrally nontrivial and becomes trivial after inverting one Hahn monomial (theorems 34.3 and 34.6).

So every bundle of corollary 10.2 is nontrivial over $\mathcal{I}_\Gamma$ and trivial over $\mathcal{H}_\Gamma$. Nothing printed in this report is retracted: no earlier part claimed nontriviality over $\mathcal{H}_\Gamma$, and item 2 declined that claim explicitly.

In rank one the answer is the opposite, and this is not new: pole-data line bundles stay nontrivial over $\mathcal{H}_\Gamma$ (corollary 36.2), by the Picard reduction and the Cousin criterion of the global-divisor report. The same interpolation construction then yields self-extensions of the trivial line that stay nonsplit although their middle bundle is free (theorems 36.3 and 36.4); an explicit cardinal-sine family with omnific-integer values at every ordinary real argument (theorem 37.1); a proper, nonprincipal, locally unit ideal of the global ring whose quotient has projective dimension two (theorem 37.3 and corollary 37.4); ultrafilter maximal ideals above it (theorem 37.5); and the classification of the zeros of the sine family on finite surcomplex halos (theorem 38.2).

What stays open is recorded in section 31: pole cocycles whose nilpotent directions vary with the puncture, an exact pole-only criterion after localization, the optimal negative budget, and the determinant-one diagonal examples (11.4), to which neither the source manuscript nor this merge applies.

33.2 Source and status

This part is built from *Pole Localization and Omnific Interpolation: Explicit bundle trivializations, persistent line classes, and a surcomplex Bézout obstruction*, dated September 23, 2026, called *the pole-localization manuscript* below (appendix C). It inspected this report at commit

0865f043aec113c14c69ef45006bbc7546a4e75a,

file blob c2dc2f9, which is exactly the text of Parts I–IV into which this part was merged; its quotations of this report are therefore current. The manuscript audited this report only. The comparisons with the global-divisor, analytic-geometry, trigonometry and entire-functions reports below were made by the merge.

Its own assessment of novelty is bounded: it proposes as new the explicit pole-localization construction, its resolution of (8.2), and the package of localization, extension, Bézout, omnific-interpolation and zero-geometry consequences, and it credits Hahn multiplication, positive inversion, the Weierstrass and Mittag–Leffler theorems, the description of Picard groups by unit cocycles and the symmetric powers of SL_2 as classical [10, 21, 22, 23]. The merge found further overlaps, which are printed once and credited:

- The Picard decomposition (proposition 36.1) and the persistence of pole-data line classes (corollary 36.2) are the global-divisor report’s `global:prop:explog`, `global:prop:Picreduction`, `global:thm:piccompact`, `global:thm:obstruction` and `global:thm:dichotomy` [2]; the discrete interpolation criterion (proposition 37.2) is its `global:cor:interp` exactly; the nonsplitting argument of theorem 36.3 is an instance of its `global:thm:cousincriterion`.
- The Bézout, corona and ultrafilter statements of section 37 are plane versions of the analytic-geometry report’s unit-disk results `analytic:thm:fixed-quotient`, `analytic:thm:fixed-bézout`, `analytic:thm:fixed-corona` and `analytic:thm:fixed-two-ideals(2)` [5], which use the same cardinal-function mechanism with the sine $\sin(\pi/(1-z))$ on the unit disk.
- The integral nontriviality statements are instances of Criterion P (theorem 6.2); the pole-only cocycles of theorem 34.6 are the Part III bundles with h_n replaced by pole data, whose integral nontriviality remark 22.2 already records.

What is new relative to the collection is the Laurent frame construction and everything it gives over $\mathcal{H}_\Gamma$ for pole data (sections 34 and 35), the free-middle nonsplit extensions, the projective-dimension computation, the omnific-valued sine family and the classification of its zeros on finite halos. None of this has received the maintained mathematical review recorded for Parts I and II.

33.3 Conventions, and the three gluing conventions

Throughout this part Γ is a set-sized totally ordered abelian group, with no divisibility or Archimedean hypothesis unless stated, and the sheaves $\mathcal{H}_\Gamma$, $\mathcal{I}_\Gamma$, $\mathcal{J}_\Gamma$ are those of (2.2)–(2.4). “Integral” means nonnegative Hahn valuation, as everywhere in this report, not membership in the omnific integers **Oz**. Let $A = \{a_n : n \geq 1\} \subset \mathbb{C}$ be closed, discrete and infinite, choose pairwise disjoint disks D_n about a_n , and put $U_0 = \mathbb{C} \setminus A$ and $W_n = D_n \setminus \{a_n\}$: a puncture-and-disk cover in the sense of section 6. For another ordered group Δ write $\mathcal{H}_\Delta = \mathcal{O}((t^\Delta))$ and $A_{\mathcal{H}_\Delta} = \mathcal{O}(\mathbb{C})((t^\Delta))$; as in Part III the letter A alone is the puncture set, and $A_{\mathcal{H}_\Gamma} = \mathcal{H}_\Gamma(\mathbb{C})$.

The gluing convention, once. This part writes a trivialization of transitions G_n on W_n as matrices B_0 over U_0 and B_n over D_n with

$$B_n = B_0 G_n \quad \text{on } W_n. \quad (33.1)$$

This is a third convention. Part I writes $G_a = B_0 B_a^{-1}$ (theorem 6.2); Part III writes $B_n = G_n B_0$ (section 22.1). If (B_0, B_n) satisfies (33.1), then (B_0^{-1}, B_n^{-1}) solves Part I’s equation $G_n = B_0^{-1} (B_n^{-1})^{-1}$ for the *same* cocycle, so triviality statements for a given cocycle mean the same thing in Parts I and V; the frames of this part are the inverses of Part I’s splitting matrices, and an integral frame in this part is a Part I splitting after the normalization of proposition 10.1. Part III’s equation

$B_n = G_n B_0$ is (33.1) for the cocycle G_n^{-1} with frames B_i^{-1} . For the unipotent pole cocycles of this part, G_n^{-1} has the same form with p_n replaced by $-p_n$, so no statement depends on the choice.

Names. The pole-localization manuscript was written independently and several of its symbols collide with symbols already in use here. They are renamed in this part only:

This report	Manuscript	Reason
$A = \{a_n\}, D_n, W_n$	D, U_n, W_n	Part I's names for punctures and disks
$A_{\mathcal{H}_\Gamma}, A_{\mathcal{H}_\Delta}$	A_Γ, A_Δ	Part III's rule: A alone is the puncture set
$\mathcal{S}$	S	S is used for supports throughout
κ_n (and the cardinal sines)	F_n, ℓ_n	F is Part I's central matrix, ℓ_d Part II's meridians
π_-	PP_{a_n}	$\mathcal{P}$ is Part I's algebra of negative Laurent parts; the principal part is π_- of lemma 4.1
F, Q_n, H_n of example 8.3	L^{-1}, C_n, H_n	the manuscript's gauges are Part I's, and $R_n = H_n^{-1} Q_n$
Ξ_n	Q_n	Q_n is already the disk gauge of example 8.3
$\mathfrak{a} = (f, \mathcal{S}), \bar{A} = A_{\mathcal{H}_\Delta}/\mathfrak{a}$	I, Q	I is the identity matrix
$\mathcal{K}, k_\gamma, \mu$	H, h_γ, d	$\mathcal{H}_\Gamma, h_n$ and $d \in D$ are in use
$\theta_m = 1 - 1/m, \hat{\mathcal{S}} = t\mathcal{S}$	β_m, T	β_n is Part III's second profile

The functions p_n with a pole at a_n keep their name; it is the same use as in proposition 27.1. The profile α_n is Part III's. The frames T_0, T_n keep the manuscript's names; they are unrelated to the gauge T of proposition 6.3. The function ϕ is (8.4), and $t^q = \omega^{-q}$ is the reading (2.1). Where the manuscript counts pole orders it writes m_n ; its δ in corollary 34.4 is a local exponent, not the generator of a cyclic group.

34 Explicit Laurent frames for pole-only unipotent cocycles

34.1 An ordinary principal-part construction

The following lemma is ordinary complex analysis; it is the only analytic preparation needed for the localization theorem.

Lemma 34.1 (Pole cancellation by an entire coefficient). *Suppose $p_n \in \mathcal{O}(W_n)$ has a pole of exact finite order $m_n \geq 1$ at a_n . There are an entire function f and entire functions κ_n such that*

- (i) *the zeros of f are exactly the a_n , with multiplicities m_n ;*
- (ii) *$f p_n$ extends holomorphically across a_n ;*
- (iii) *$p_n \kappa_j$ extends holomorphically across a_n whenever $j \neq n$;*
- (iv) *$1/f + p_n \kappa_n$ extends holomorphically across a_n ;*
- (v) *$\kappa_n(a_n) \neq 0$ and $\kappa_j(a_n) = 0$ for $j \neq n$.*

One may take

$$q_n = \pi_- \left(-\frac{1}{f^2 p_n} \right) \text{ at } a_n, \quad \kappa_n = f q_n, \quad (34.1)$$

where the negative Laurent part q_n is a rational function of $z - a_n$.

Proof. The ordinary Weierstrass theorem supplies f with the prescribed divisor [22, 21]. Near a_n write $u = z - a_n$, $f = u^{m_n}h(u)$ and $p_n = u^{-m_n}k(u)$ with $h(0)k(0) \neq 0$. The germ $-1/(f^2p_n)$ has a pole of order exactly m_n ; subtracting its principal part q_n leaves a holomorphic germ r_n , so $-1/(f^2p_n) = q_n + r_n$. The rational function q_n has no pole away from a_n and pole order at most m_n there, so $\kappa_n = fq_n$ is entire, and

$$\frac{1}{f} + p_n\kappa_n = -fp_nr_n \quad (34.2)$$

is holomorphic at a_n . For $j \neq n$ the rational function q_j is holomorphic at a_n , so κ_j vanishes there to order at least m_n , which gives (iii) and the second half of (v); (ii) holds since $fp_n = hk$. Finally the leading term of q_n gives $\kappa_n(a_n) = -1/(h(0)k(0)) \neq 0$. Only finite-order ordinary poles occur; no infinite Hahn sum is exchanged with an ordinary one. $\square$

Example 34.2 (Simple poles). If $p_n = c_n/(z - a_n)$ with $c_n \neq 0$ and the zeros of f are simple, then

$$\kappa_n(z) = -\frac{f(z)}{c_nf'(a_n)^2(z - a_n)}, \quad \kappa_n(a_n) = -\frac{1}{c_nf'(a_n)}. \quad (34.3)$$

The singularity at a_n is removable, and (34.2) cancels one ordinary residue; the coefficient functions for different punctures will sit at different Hahn exponents.

No bound on the pole orders m_n is needed, but each singularity must have finite order: an entire function cannot cancel an essential singularity by a finite-order zero, and this is exactly where the construction stops applying. The essential-singularity bundles of Part III lie beyond it (lemma 21.1).

34.2 The rank-two theorem

Fix a strictly decreasing profile

$$\alpha_1 > \alpha_2 > \cdots > 0 \quad \text{in } \Gamma, \quad (34.4)$$

take f, κ_n from lemma 34.1, and put

$$\mathcal{S} = \sum_{j \geq 1} t^{-\alpha_j} \kappa_j \in A_{\mathcal{H}_\Gamma}, \quad T_0 = \begin{pmatrix} f & 0 \\ \mathcal{S} & f^{-1} \end{pmatrix} \text{ on } U_0. \quad (34.5)$$

The support of $\mathcal{S}$ is the strictly increasing, hence well-ordered, sequence $(-\alpha_j)$, and f has no zero on U_0 .

Theorem 34.3 (Pole-only unipotent bundles become trivial after one monomial localization). *The transitions*

$$G_n = \begin{pmatrix} 1 & t^{\alpha_n} p_n \\ 0 & 1 \end{pmatrix} \quad (34.6)$$

define a nontrivial rank-two $\mathcal{S}_\Gamma$ -bundle whose extension to $\mathcal{H}_\Gamma$ is trivial. An explicit determinant-one trivialization in the convention (33.1) is T_0 and

$$T_n = T_0 G_n = \begin{pmatrix} f & t^{\alpha_n} f p_n \\ \mathcal{S} & f^{-1} + t^{\alpha_n} p_n \mathcal{S} \end{pmatrix} \in \mathrm{SL}_2(\mathcal{H}_\Gamma(D_n)). \quad (34.7)$$

Every entry of T_0, T_n and their inverses has valuation at least $-\alpha_1$, so the bundle is already trivial over $\mathcal{S}_\Gamma[(t^{\alpha_1})^{-1}]$.

Proof. First $\det T_0 = 1$, and $T_0^{-1} = \begin{pmatrix} f^{-1} & 0 \\ -\mathcal{S} & f \end{pmatrix}$ is a Laurent matrix on U_0 . In (34.7) the first column is entire coefficientwise and fp_n extends across a_n . The lower-right entry is

$$f^{-1} + p_n \kappa_n + \sum_{j \neq n} t^{\alpha_n - \alpha_j} p_n \kappa_j. \quad (34.8)$$

The first two terms combine into a function holomorphic on D_n by lemma 34.1(iv), and each other coefficient is holomorphic on D_n by (iii). For fixed n the exponents $\alpha_n - \alpha_j$ increase strictly with j , so their union with $\{0\}$ is well ordered and each exponent carries one coefficient: (34.8) is a section of $\mathcal{H}_\Gamma(D_n)$, not a formal expression requiring a completion. The identity $\det T_n = 1$ holds on W_n and hence coefficientwise on D_n , so the adjugate is the inverse, with the same support bounds. No displayed exponent lies below $-\alpha_1$; multiplying by t^{α_1} makes every entry integral.

Integral nontriviality, a direct route. Suppose integral frames B_0, B_n satisfy (33.1). Their reductions modulo $\mathcal{J}_\Gamma$ agree on overlaps, since G_n reduces to I_2 , and glue to $C \in \mathrm{GL}_2(\mathcal{O}(\mathbb{C}))$; replacing every B_i by $C^{-1}B_i$ makes them positive. At exponent α_n the $(1, 2)$ entry of $B_n = B_0 G_n$ is $(B_n)_{12, \alpha_n} = (B_0)_{12, \alpha_n} + p_n$, only the exponent-zero coefficient I_2 of B_0 meeting $t^{\alpha_n} p_n E_{12}$. As p_n has a pole, $(B_0)_{12, \alpha_n} \neq 0$ for every n , and $\mathrm{supp}(B_0 - I_2)$ contains the descending set $\{\alpha_n\}$, which is impossible. This is also an instance of Criterion P: $\mathrm{Pol}(G_n) = I_2 + t^{\alpha_n} \pi_-(p_n) E_{12}$, so $\mathcal{E}(G) \supseteq \{\alpha_n\}$ is not well ordered. $\square$

What the construction changes. A positive frame would have to place pole-cancelling data at the forbidden exponents α_n . The Laurent frame instead places the interpolation coefficients κ_n at the admissible exponents $-\alpha_n$; multiplication by the transition at the n th puncture moves that one term to exponent zero, where it cancels $1/f$, while all other coefficients already vanish to sufficient order there. The determinant is exactly one, so the adjugate supplies the inverse inside the required ring; invertibility of ordinary values alone would not.

Corollary 34.4 (Balanced denominator bound). *For $\delta \in \Gamma$ with $0 \leq \delta \leq \alpha_1$, the frames $\mathrm{diag}(t^{-\delta}, t^\delta)T_i$ and their inverses have valuation at least $-\max\{\delta, \alpha_1 - \delta\}$. If $\alpha_1/2 \in \Gamma$ this bound is $-\alpha_1/2$.*

Proof. The nonzero entries of the new T_0 have valuations $-\delta$, $\delta - \alpha_1$ and δ , and its inverse has the same bound. The disk frames differ by right multiplication by G_n , integral with integral inverse. The determinant stays one. $\square$

Corollary 34.5 (Arbitrarily small negative budgets). *Let $\Gamma = \mathbb{Q}$ and $\alpha_n \rightarrow 0$ in the ordinary sense. For every rational $\varepsilon > 0$ the bundle of theorem 34.3 admits determinant-one Laurent frames whose entries and inverse entries have valuation at least $-\varepsilon$; it admits no integral frame.*

Proof. Choose N with $\alpha_{N+1} < \varepsilon$. On U_0 put $g_0 = I_2 + \sum_{n \leq N} t^{\alpha_n} \pi_-(p_n) E_{12}$, the principal parts being rational functions with poles only at a_n ; put $g_n = G_n^{-1} g_0$ for $n \leq N$ and $g_n = g_0$ for $n > N$. All these matrices commute, and each g_n is holomorphic on D_n : for $n \leq N$ the coefficient at α_n is $\pi_-(p_n) - p_n$, and every other principal part is holomorphic near a_n . The integral gauge $B_i \mapsto B_i g_i$ turns the cocycle into I_2 at the first N punctures and leaves G_n unchanged for $n > N$. Apply theorem 34.3 to the tail, with f vanishing only at a_n , $n > N$; its frames extend across the first N punctures, where the cocycle is now the identity, and have valuation at least $-\alpha_{N+1} > -\varepsilon$. Composing with the integral gauge does not lower this bound. Integral nontriviality is theorem 34.3. $\square$

This is an infimum statement: in this family the infimum of admissible negative budgets is 0 and is not attained. It is not a claim that a least positive budget exists.

34.3 Every fixed nilpotent direction

Theorem 34.6 (Fixed-nilpotent pole localization). *Let $N \in \text{Mat}_r(\mathbb{C})$ be nonzero and nilpotent with nilpotency index ν , and keep the hypotheses on p_n and α_n . The transitions*

$$G_n = \exp(t^{\alpha_n} p_n N) = \sum_{j=0}^{\nu-1} \frac{t^{j\alpha_n} p_n^j N^j}{j!} \quad (34.9)$$

define an integrally nontrivial bundle that admits determinant-one Laurent frames, with frames and inverses of valuation at least $-(\nu-1)\alpha_1$, and at least $-(\nu-1)\alpha_1/2$ when $\alpha_1/2 \in \Gamma$. One monomial localization suffices.

Proof. Integral nontriviality. Normalize an assumed integral frame to have exponent-zero coefficient I_r as in [theorem 34.3](#). The coefficient at α_n of $B_n = B_0 G_n$ is $(B_0)_{\alpha_n} + p_n N$; the higher terms of the exponential have exponent at least $2\alpha_n$ and meet no negative support of B_0 . A nonzero entry of N forces $\alpha_n \in \text{supp}(B_0 - I_r)$ for every n . (With h_n replaced by p_n this is the computation of [remark 22.2](#).)

Existence. First let N be one Jordan block of size $d+1$. Apply the d th symmetric power of the standard representation of SL_2 , acting on homogeneous polynomials of degree d in two variables, to the frames of [theorem 34.3](#). The matrix $\exp(aE_{12})$ acts by the substitution $Y \mapsto Y + aX$, which is $\exp(aX\partial_Y)$, and $X\partial_Y(X^{d-k}Y^k) = kX^{d-k+1}Y^{k-1}$; so $X\partial_Y$ is one nilpotent Jordan block with nonzero weights $1, \dots, d$, and a constant diagonal change of basis conjugates it to the standard block. Both sides of $\text{Sym}^d(\exp(aE_{12})) = \exp(aX\partial_Y)$ are polynomial in a , so the identity holds for Hahn-series values of a . Every entry of $\text{Sym}^d M$ is a homogeneous polynomial of degree d in the entries of M , so the represented frames are Hahn matrices with holomorphic coefficients on the same domains, and valuation bounds are multiplied by d ([lemma 3.2](#)). The represented inverse is the representation of the inverse, and $\det \text{Sym}^d M = (\det M)^{d(d+1)/2} = 1$ for $M \in \text{SL}_2$: the identity holds for diagonal matrices by multiplying the weights, hence for diagonalizable ones, hence for all, since both sides are polynomials and diagonalizable matrices are Zariski dense. Since Sym^d is a homomorphism, $\text{Sym}^d T_n = \text{Sym}^d T_0 \text{Sym}^d G_n$.

For general $N = C(\bigoplus_k J_k)C^{-1}$ in constant Jordan form, take the block sum of the frames just built, identity frames on zero blocks, and conjugate by C ; then $B_0 G_n = B_n$ blockwise. Constant conjugation preserves valuation bounds and determinants. The largest d is $\nu-1$; starting from the frames of [corollary 34.4](#) gives the improved bound. $\square$

The hypothesis that one N serves every puncture is used. The argument does not triangularize puncture-dependent nilpotent matrices simultaneously and does not solve a general noncommutative Cousin problem after localization; it gives an exact family, not a classification of Laurent-trivial bundles.

35 The printed examples become trivial over $\mathcal{H}_\Gamma$

35.1 The hidden rank-three example

Take the data of [section 8](#): $A = \{2, 3, \dots\}$, $u_n = z - n$, $\Gamma = \mathbb{Q}$, the cocycle (8.2), the function ϕ of (8.4), and the matrices

$$F = I_3 + t\phi E_{12}, \quad Q_n = I_3 + t\left(\phi - \frac{1}{u_n}\right)E_{12}, \quad H_n = I_3 + t^{1/n}E_{23}$$

of [example 8.3](#) and (8.3). They are positive, F on U_0 and Q_n, H_n on D_n , with determinant one.

Theorem 35.1 (The hidden obstruction does not survive monomial localization). *The transitions (8.2) are integrally nontrivial but admit determinant-one Laurent frames whose entries and inverse entries all have valuation at least $-3/4$. In particular the bundle $\mathcal{E}_G$ of [corollary 10.2](#) is already trivial over $\mathcal{I}_\mathbb{Q}[t^{-1}]$, and hence over $\mathcal{H}_\Gamma$.*

Proof. Put $\Xi_n = I_3 - t^{1+1/n}u_n^{-1}E_{13}$. By (8.3), $G_n = P_n H_n$, and $P_n = \Xi_n (I_3 + tu_n^{-1}E_{12})$ because $E_{13}E_{12} = 0$. By [example 8.3](#), $I_3 + tu_n^{-1}E_{12} = FQ_n^{-1}$. The matrix Ξ_n commutes with every upper unitriangular matrix, so

$$G_n = F \Xi_n (H_n^{-1}Q_n)^{-1}, \quad \text{equivalently} \quad F^{-1}G_n H_n^{-1}Q_n = \Xi_n. \quad (35.1)$$

This integral equivalence turns the example into a rank-two pole cocycle on coordinates 1 and 3, coordinate 2 being untouched. Apply [theorem 34.3](#) to it with $\alpha_n = 1 + 1/n$ and $p_n = -1/u_n$, $n \geq 2$: choose f entire with simple zeros exactly at $2, 3, \dots$ (for instance $f(z) = 1/\Gamma_E(2 - z)$, with Γ_E Euler's gamma function), so that by (34.3)

$$\kappa_n(z) = \frac{f(z)}{f'(n)^2(z - n)}, \quad \mathcal{S} = \sum_{n \geq 2} t^{-(1+1/n)} \kappa_n,$$

and let

$$T_0 = \begin{pmatrix} f & 0 & 0 \\ 0 & 1 & 0 \\ \mathcal{S} & 0 & f^{-1} \end{pmatrix}, \quad T_n = T_0 \Xi_n. \quad (35.2)$$

These are determinant-one Laurent frames for Ξ_n . Consequently

$$B_0 = T_0 F^{-1}, \quad B_n = T_n Q_n^{-1} H_n \quad (35.3)$$

satisfy $B_0 G_n = T_0 \Xi_n Q_n^{-1} H_n = B_n$, exactly in the convention (33.1); in Part I's convention $G_n = B_0^{-1}(B_n^{-1})^{-1}$.

For the bound replace each T_i by $\text{diag}(t^{-3/4}, 1, t^{3/4})T_i$. The largest exponent of the profile is $3/2$, at $n = 2$, so [corollary 34.4](#) gives valuation at least $-3/4$ for these matrices and their inverses, and multiplication by the integral matrices $F^{\pm 1}$, $Q_n^{\pm 1}$, $H_n^{\pm 1}$ cannot lower it. Multiplying every entry by t makes it integral, so all frames and inverse frames lie over $\mathcal{I}_\mathbb{Q}[t^{-1}]$. Finally Ξ_n is integrally nontrivial by [theorem 34.3](#) applied to its (1,3) entry, and (35.1) is integral in both directions, which reproves the integral half of [theorem 8.2](#) without Criterion P. $\square$

The number $3/4$ is a proved upper bound on the negative budget, not a claim of optimality. The frames lie in the intended ring $\mathcal{H}_\Gamma$, not in the coefficientwise meromorphic sheaf $\mathcal{M}_\Gamma$, over which triviality was already known ([corollary 27.2](#)).

35.2 The deep examples

The pole-localization manuscript treats only the hidden example. The deep examples reduce in the same way. The following corollary is added by the merge.

Corollary 35.2 (The deep obstructions do not survive monomial localization; merge). *For every $r \geq 3$ the cocycle $G_n = I_r + C_n + a_n E_{12}$ of [theorem 9.2](#) (for $r = 3$ the cocycle (8.2)) is integrally equivalent to the rank-two cocycle*

$$\Xi_n = I_r + (-1)^r \frac{t^{r-2+1/n}}{z-n} E_{1r}, \quad n \geq 2, \quad (35.4)$$

acting on coordinates 1 and r . It admits determinant-one Laurent frames whose entries and inverse entries have valuation at least $-(2r-3)/4$, so the integral bundle of [corollary 10.2](#) is trivial over $\mathcal{I}_{\mathbb{Q}}[t^{-1}]$ and hence over $\mathcal{H}_{\Gamma}$.

Proof. By [lemma 9.1](#), $G_n = P_n H_n$ with $H_n = I_r + C_n$, and the $(1, r)$ entry of P_n is

$$(-1)^{r-2} a_n \prod_{j=2}^{r-1} c_{j,n} = (-1)^r \frac{t}{z-n} t^{1-1/n} t^{r-4} t^{2/n} = (-1)^r \frac{t^{r-2+1/n}}{z-n}$$

for $r \geq 4$ by (9.2), in agreement with (9.4); for $r = 3$ it is $-t^{1+1/n}/u_n$ by (8.3). Let P'_n be P_n with this entry deleted, as in the proof of [theorem 9.2](#). Both $P_n - I_r$ and $P'_n - I_r$ are supported in row one, whose products vanish, so $P_n = \Xi_n P'_n$ with Ξ_n as in (35.4). The proof of [theorem 9.2\(ii\)](#) supplies positive F' on U_0 and Q'_n on D_n , valued in $I_r + V$, with $F' = P'_n Q'_n$ (for $r = 3$, $F' = F$ and $Q'_n = Q_n$ of [example 8.3](#)). The matrix Ξ_n commutes with every upper unitriangular matrix, because $E_{1r} M = E_{1r} = M E_{1r}$ for such M . Hence

$$G_n = \Xi_n F' Q_n'^{-1} H_n = F' \Xi_n (H_n^{-1} Q_n')^{-1},$$

an integral equivalence with Ξ_n . The cocycle Ξ_n is the rank-two cocycle of [theorem 34.3](#) on coordinates 1, r with the strictly decreasing positive profile $\alpha_n = r - 2 + 1/n$ and the simple poles $p_n = (-1)^r/(z-n)$, $n \geq 2$. With f as in [theorem 35.1](#), $\kappa_n = (-1)^{r+1} f/(f'(n)^2(z-n))$ by (34.3), and $\mathcal{S} = \sum_{n \geq 2} t^{-\alpha_n} \kappa_n$, let T_0 be I_r with the entries f at $(1, 1)$, $\mathcal{S}$ at $(r, 1)$ and f^{-1} at (r, r) , and $T_n = T_0 \Xi_n$. Then

$$B_0 = T_0 F'^{-1}, \quad B_n = T_n Q_n'^{-1} H_n$$

are determinant-one Laurent frames with $B_0 G_n = B_n$. The largest exponent of the profile is $r - 3/2$, at $n = 2$; [corollary 34.4](#) with $\delta = (2r-3)/4$, applied on coordinates 1, r , gives the bound $-(2r-3)/4$, which the integral factors $F'^{\pm 1}$, $Q_n'^{\pm 1}$, $H_n^{\pm 1}$ do not lower. For $r = 3$ this is [theorem 35.1](#). $\square$

The identities used in this proof were checked symbolically by the merge for $3 \leq r \leq 8$, with symbolic row-one entries of F' and symbolic f and $\mathcal{S}$ ([section 29](#)); the proof above does not depend on that check.

35.3 What this settles and what it does not

[Theorem 35.1](#) and [corollary 35.2](#) settle the printed question: every explicit example of [corollary 10.2](#) is nontrivial over $\mathcal{I}_{\Gamma}$ and trivial over $\mathcal{H}_{\Gamma}$, already after inverting t . Four consequences for the rest of the report are recorded here.

- (a) *Criterion P does not survive localization as a criterion.* The hidden example fails the integral polar-support test and is Laurent-trivial. An exact criterion after localization would have to recognize the interpolation cancellation of [theorem 34.3](#); none is known ([section 31](#)).
- (b) *Poles and essential singularities are separated over $\mathcal{H}_\Gamma$.* [Remark 22.2](#) observed that pole and essential data are separated only after passage to $\mathcal{M}_\Gamma$. With one fixed nilpotent N , the bundles $\exp(t^{\alpha_n} p_n N)$ with finite-order poles p_n are $\mathcal{H}_\Gamma$ -trivial ([theorem 34.6](#)), whereas those with $h_n = \exp(1/(z - n))$ are not even stably trivial over $\mathcal{H}_\Gamma$ ([corollary 22.5](#)). The dividing line is exactly the finite order in [lemma 34.1](#).
- (c) *Consistency.* Triviality over $\mathcal{H}_\Gamma$ implies the triviality over $\mathcal{M}_\Gamma$ of [corollary 27.2\(a\)](#), by a different mechanism. Bounded triviality ([corollary 10.3](#)) and the finite-puncture blindness of item 4 of [section 30](#) are unaffected.
- (d) *Not settled.* The determinant-one diagonal cocycles ([11.4](#)) are not unipotent; their summands are the rank-one classes of [corollary 36.2](#), which persist over $\mathcal{H}_\Gamma$, and whether their direct sum becomes trivial over $\mathcal{H}_\Gamma$ is not decided here. Pole cocycles with puncture-dependent nilpotent directions are not treated either.

36 Line bundles and extensions that localization does not kill

36.1 Line bundles

Write $\underline{\Gamma}$ for the sheaf of locally constant Γ -valued functions on an ordinary domain $X \subseteq \mathbb{C}$. Picard groups below classify locally free rank-one modules over the specified sheaf of rings (the group Pic_Γ of the global-divisor report; compare its `global:rem:piclf`).

Proposition 36.1 (Picard decomposition; global-divisor report). *There are natural decompositions of abelian sheaves $\mathcal{H}_\Gamma^\times \simeq \underline{\Gamma} \times \mathcal{I}_\Gamma^\times$ and $\mathcal{J}_\Gamma^\times \simeq \mathcal{O}_X^\times \times \mathcal{J}_\Gamma$, and hence*

$$\text{Pic}(\mathcal{H}_\Gamma) \simeq H^1(X, \underline{\Gamma}) \oplus \text{Pic}(\mathcal{J}_\Gamma), \quad (36.1)$$

$$\text{Pic}(\mathcal{J}_\Gamma) \simeq \text{Pic}(\mathcal{O}_X) \oplus H^1(X, \mathcal{J}_\Gamma). \quad (36.2)$$

Extension of scalars $\text{Pic}(\mathcal{J}_\Gamma) \rightarrow \text{Pic}(\mathcal{H}_\Gamma)$ is the inclusion of the second summand of (36.1), hence injective. For $X = \mathbb{C}$ it is an isomorphism, and both groups are $H^1(\mathbb{C}, \mathcal{J}_\Gamma)$.

Proof. The sheaf decompositions are `global:eq:unitsplit` and `global:eq:Iunits` of `global:prop:explog` [2], proved there on every ordinary open set: valuation splits off by the unit criterion (`global:prop:units`; [lemma 20.3](#)) with section $\beta \mapsto t^\beta$, and the positive logarithm identifies the principal units with $\mathcal{J}_\Gamma$. Taking H^1 of these direct-product decompositions, and using that locally free rank-one modules are classified by unit cocycles, gives (36.1) and (36.2). For $X = \mathbb{C}$ the first summand and $\text{Pic}(\mathcal{O}_\mathbb{C})$ vanish, which is `global:prop:Picreduction` and its display `global:eq:PicH1`. $\square$

The global-divisor report proves this on the plane and, with the valuation summand, over a compact base (`global:thm:piccompact`); the pole-localization manuscript states it on an arbitrary ordinary domain, which the same argument covers. It is elementary unit-group algebra, not a new cohomology theorem, and the manuscript says so. On the puncture-and-disk cover the injectivity is also visible directly: a Laurent splitting $b_n = b_0 g_n$ of a cocycle $g_n \in 1 + \mathcal{J}_\Gamma(W_n)$ has one common valuation β and leading coefficients that glue to a nowhere-zero entire h , and dividing by $t^\beta h$ gives a positive splitting. This is the normalization in the proof of `global:thm:obstruction`.

Corollary 36.2 (Descending pole line bundles stay nontrivial over $\mathcal{H}_\Gamma$). *Under the hypotheses of lemma 34.1 and (34.4), the scalar transitions $g_n = \exp(t^{\alpha_n} p_n)$ define a line bundle that is nontrivial and not stably trivial over $\mathcal{H}_\Gamma$, and*

$$(c_n) \mapsto [\exp(c_n t^{\alpha_n} p_n)] \quad (36.3)$$

induces an injective group homomorphism $\mathbb{C}^\mathbb{N}/\mathbb{C}^{(\mathbb{N})} \hookrightarrow \text{Pic}(\mathcal{H}_\Gamma)$, where $\mathbb{C}^{(\mathbb{N})}$ denotes the finitely supported sequences.

Proof. By the normalization just described a Laurent splitting would give a positive one, and taking logarithms would split the additive cocycle $c_n t^{\alpha_n} p_n$ in $\mathcal{J}_\Gamma$. Its singular support is $\{\alpha_n : c_n \neq 0\}$, so by `global:thm:cousincriterion` it splits exactly when finitely many c_n are nonzero; for finitely many, the finite sum of the principal parts at their exponents is an explicit splitting. Exponentials of commuting cocycles multiply as their logarithms add. Stable triviality of a line bundle implies triviality by taking determinants. $\square$

For simple poles this is `global:thm:obstruction` with `global:eq:injPic` and, over subgroups of **No**, `global:thm:dichotomy`; the manuscript’s contribution is only the finite-order generalization and the contrast with theorem 34.3: in rank one monomial localization kills nothing, in rank two it kills the descending pole cocycles. It agrees with the observation after corollary 27.2 that pole-data line classes survive monomial localization.

36.2 Nonsplit extensions with a free middle bundle

For f and $\mathcal{S}$ as in (34.5) put

$$v = \begin{pmatrix} f \\ \mathcal{S} \end{pmatrix}, \quad w = \begin{pmatrix} -\mathcal{S} & f \end{pmatrix}, \quad wv = 0. \quad (36.4)$$

The frames T_i of theorem 34.3 have first column v , and their second columns s_i satisfy $ws_i = 1$.

Theorem 36.3 (A free middle bundle does not split this extension). *The sequence of sheaves of $\mathcal{H}_\Gamma$ -modules*

$$0 \longrightarrow \mathcal{H}_\Gamma \xrightarrow{a \mapsto av} \mathcal{H}_\Gamma^2 \xrightarrow{(x,y) \mapsto -\mathcal{S}x + fy} \mathcal{H}_\Gamma \longrightarrow 0 \quad (36.5)$$

is exact but not split. Equivalently, there are no $x, y \in A_{\mathcal{H}_\Gamma}$ with $-\mathcal{S}x + fy = 1$. Its extension class is represented, up to the sign convention, by the additive cocycle $t^{\alpha_n} p_n$.

Proof. In the coordinates given by the local determinant-one completions T_i of v , the sequence is the inclusion of the first coordinate followed by the projection to the second, so it is exact. From $T_n = T_0 G_n$ one gets $s_n - s_0 = t^{\alpha_n} p_n v$ on W_n . A splitting would give a global s with $ws = 1$; on each chart $s - s_i = b_i v$ with $b_i \in \mathcal{H}_\Gamma(U_i)$ (here U_i is U_0 or D_n), and on overlaps $t^{\alpha_n} p_n = b_0 - b_n$. At exponent α_n the coefficient of b_0 must have a nonremovable pole at a_n , since b_n is holomorphic there, so $\text{supp } b_0 \supseteq \{\alpha_n\}$, which is impossible. No positivity of the b_i is assumed: negative support cannot remove the forced descending positive support. This is the necessity half of `global:thm:cousincriterion` for the full sheaf $\mathcal{H}_\Gamma$. $\square$

Theorem 36.4 (Free-middle self-extensions). *The cocycles $c_n t^{\alpha_n} p_n$ induce an injective linear map*

$$\mathbb{C}^\mathbb{N}/\mathbb{C}^{(\mathbb{N})} \hookrightarrow \text{Ext}_{\mathcal{H}_\Gamma}^1(\mathcal{H}_\Gamma, \mathcal{H}_\Gamma), \quad (36.6)$$

Ext^1 being taken in sheaves of $\mathcal{H}_\Gamma$ -modules, not in modules over $A_{\mathcal{H}_\Gamma}$. Every class in the image has a representative whose middle sheaf is isomorphic to $\mathcal{H}_\Gamma^2$.

Proof. Gluing by upper unitriangular matrices realizes an additive cocycle as an extension; addition of cocycles is Baer sum and scalar multiplication acts on the upper-right entry. The argument of [theorem 36.3](#) proves nontriviality whenever infinitely many c_n are nonzero, and finite support gives a coboundary by the finite principal-part construction, so the kernel is $\mathbb{C}^{(\mathbb{N})}$. A splitting found on a refinement would still give local coefficients b_i on the original charts and the same contradiction. For an infinite active set $\{n : c_n \neq 0\}$ choose an entire function with zeros exactly at the active a_n , with the original orders, and apply [theorem 34.3](#) to the pole functions $c_n p_n$ on that subsequence; the frames extend across inactive punctures, where the cocycle is the identity, so the middle bundle is free. For finite support the extension splits and the middle bundle is free again. $\square$

The injectivity alone is the additive statement of the global-divisor report (`global:cor:H1injection` for $\mathcal{J}_\Gamma$, `global:prop:fullH1` for $\mathcal{H}_\Gamma$); what is new is that every class in this image has a *free* middle bundle. There is no conflict with projectivity of free modules over $A_{\mathcal{H}_\Gamma}$: the epimorphism in (36.5) is a sheaf epimorphism whose map on global sections is not surjective. Each chart solves $-\mathcal{S}x + fy = 1$, but no single well-ordered support carries a global solution. Nor does an isomorphism of middle bundles preserve the chosen subline and quotient: the Laurent trivialization moves the subline from a constant coordinate to the column v , and the missing global complementary lift is the surviving class. One cocycle is thus nontrivial as an integral bundle, trivial as a Laurent bundle, and nonsplit as a Laurent extension; forgetting the filtration loses information, and a line bundle has no filtration to forget.

37 An omnific-valued cardinal-sine family and a global Bézout obstruction

37.1 The family

Enumerate the integers by

$$a_1 = 0, \quad a_{2k} = k, \quad a_{2k+1} = -k \quad (k \geq 1), \quad (37.1)$$

and put

$$f(z) = \sin(\pi z), \quad \kappa_n(z) = \frac{\sin(\pi(z - a_n))}{\pi(z - a_n)} = \frac{f(z)}{f'(a_n)(z - a_n)}, \quad \mathcal{S} = \sum_{n \geq 1} t^{-1/n} \kappa_n. \quad (37.2)$$

The singularities of κ_n are removable, $\kappa_n(a_j) = \delta_{nj}$, and $\mathcal{S}(a_n) = t^{-1/n}$. Since $-1, -1/2, -1/3, \dots$ increases, $\mathcal{S} \in \mathcal{O}(\mathbb{C})((t^{\mathbb{Q}}))$. With

$$p_n(z) = -\frac{1}{f'(a_n)(z - a_n)}, \quad \alpha_n = \frac{1}{n}, \quad (37.3)$$

(34.3) returns exactly these κ_n , so every frame, localization and extension statement of [sections 34](#) and [36](#) applies to this pair $f, \mathcal{S}$. The analytic-geometry report uses the same device on the unit disk, with $h(z) = \sin(\pi/(1 - z))$ and cardinal functions e_n (`analytic:lem:fixed-cardinal`) [5].

Read $t^q = \omega^{-q}$ as in (2.1). A real Hahn family evaluated at an ordinary real point is then a surreal normal form, and complex coefficients are read by real and imaginary parts in **SC**. A surreal

number is an omnific integer exactly when its normal form has no term of negative exponent and an ordinary integer as coefficient of ω^0 [11].

Theorem 37.1 (Omnific interpolation with no Hahn-holomorphic reciprocal data). *The family (37.2) has the following properties.*

- (i) *For every ordinary $x \in \mathbb{R}$ the value*

$$\mathcal{S}(x) = \sum_{n \geq 1} \kappa_n(x) \omega^{1/n}$$

is a nonzero omnific integer with zero constant coefficient.

- (ii) $\mathcal{S}(a_n) = \omega^{1/n}$ for every n .
 (iii) *For every order-preserving extension $\mathbb{Q} \hookrightarrow \Delta$ there is no $B \in A_{\mathcal{H}_\Delta}$ with $B(a_n) = t^{1/n}$ for all n ; no common-domain Hahn family interpolates the reciprocal values $1/\mathcal{S}(a_n)$.*

Proof. For real x every $\kappa_n(x)$ is real, the only exponents are the decreasing $1/n > 0$, and the constant coefficient is zero, so $\mathcal{S}(x) \in \mathbf{Oz}$. If $x \notin \mathbb{Z}$ then $\kappa_1(x) = \sin(\pi x)/(\pi x) \neq 0$ is the leading coefficient; at an integer exactly one monomial survives. This proves (i) and (ii). For (iii), ordinary evaluation can delete exponents but not create them, so $B(a_n) = t^{1/n}$ forces $1/n \in \text{supp } B$ for all n , a strictly decreasing sequence in Δ . $\square$

The values in (i) are evaluations of one specified Hahn-holomorphic family at ordinary real arguments; they are not claimed to define a function continuous for any topology on $\mathbf{No}$, and the reciprocals $\omega^{-1/n}$ exist individually in $\mathbf{No}$: (iii) concerns their simultaneous common-domain interpolation, not field inversion. The argument of (iii) is the evaluation-support argument of `analytic:thm:fixed-corona` and the necessity half of `global:cor:interp`.

Proposition 37.2 (Discrete interpolation: the support test). *Let $\{a_n\} \subset \mathbb{C}$ be closed and discrete and $b_n \in K_\Gamma$. There is $F \in A_{\mathcal{H}_\Gamma}$ with $F(a_n) = b_n$ for all n if and only if $\bigcup_n \text{supp } b_n$ is well ordered.*

Proof. This is `global:cor:interp` of the global-divisor report [2], whose proof interpolates the complex coefficients at each exponent of the well-ordered union by an ordinary entire function; it is not reprinted. The unit-disk counterpart is the surjectivity in `analytic:thm:fixed-quotient`. $\square$

For (37.2) the values $t^{-1/n}$ pass this test and the reciprocal values $t^{1/n}$ fail it. The pole-localization manuscript proved the proposition independently and presented it as a standard-style consequence, not as a priority claim.

37.2 A proper, locally unit, nonprincipal ideal

Theorem 37.3 (A support-controlled Bézout obstruction). *Let f and $\mathcal{S}$ be as in (37.2). For every ordered extension $\mathbb{Q} \hookrightarrow \Delta$ the ideal*

$$\mathfrak{a} = (f, \mathcal{S}) \subset A_{\mathcal{H}_\Delta} \tag{37.4}$$

is proper, although the associated ideal sheaf is the unit sheaf on the ordinary plane. After Taylor prolongation f and $\mathcal{S}$ have no common zero at any finite surcomplex argument. Every common divisor of f and $\mathcal{S}$ is a unit, $\mathfrak{a}$ is not principal, and $A_{\mathcal{H}_\Delta}$ is not a Bézout domain.

Proof of properness and local generation. If $fX + SY = 1$ in $A_{\mathcal{H}_\Delta}$, evaluation at a_n gives $Y(a_n) = t^{1/n}$, contrary to [theorem 37.1\(iii\)](#). On U_0 , f has the ordinary inverse $1/f$; on D_n the determinant-one frame T_n built from [\(37.3\)](#) completes $(f, \mathcal{S})^\top$ to a basis, and its second column solves $-\mathcal{S}x + fy = 1$ there. The same matrices remain valid over every ordered extension. If h divides f and $\mathcal{S}$, local generation of the unit ideal makes h locally invertible; the local inverses are unique and glue by the sheaf property ([lemma 20.2](#)), since $\mathcal{H}_\Delta(\mathbb{C}) = A_{\mathcal{H}_\Delta}$, so h is a unit. A principal $\mathfrak{a}$ would therefore be generated by a unit, contrary to properness. The statement about finite surcomplex zeros is proved in [section 37.4](#). $\square$

Consequently $A_{\mathcal{H}_\Delta}$ has maximal ideals containing $\mathfrak{a}$ that are not ordinary evaluation ideals: at each a_n , where f vanishes, $\mathcal{S}(a_n) \neq 0$. The word Bézout is used in its literal equation sense. The theorem does not concern the classical bounded-function corona problem, and it is about the common-domain ring $\mathcal{O}(\mathbb{C})((t^\Delta))$, not the different ring of strongly entire series, which *is* Bézout when Γ has an order unit (`ent:thm:bezout` of the entire-functions report [\[7\]](#)); there is no conflict. The unit-disk version, with an exact Bézout criterion modulo a fixed simple divisor and norm estimates, is `analytic:thm:fixed-bezout` and `analytic:thm:fixed-corona`; the plane version here adds the explicit sine, the omnific values and the following homological computation.

Corollary 37.4 (Regular sequence and projective dimensions). *For every ordered extension $\mathbb{Q} \hookrightarrow \Delta$ put $A = A_{\mathcal{H}_\Delta}$, $\mathfrak{a} = (f, \mathcal{S})$ and $\bar{A} = A/\mathfrak{a}$. Then $f, \mathcal{S}$ is a regular sequence in A , and*

$$0 \longrightarrow A \xrightarrow{a \mapsto (-Sa, fa)} A^2 \xrightarrow{(x, y) \mapsto fx + Sy} A \longrightarrow \bar{A} \longrightarrow 0 \quad (37.5)$$

is an exact free resolution. Hence $\mathrm{pd}_A \bar{A} = 2$, $\mathrm{pd}_A \mathfrak{a} = 1$, $\mathrm{Tor}_2^A(\bar{A}, \bar{A}) \simeq \bar{A} \neq 0$, and $\mathfrak{a}$ is not flat. These are statements about modules over the global ring A , not about sheaves.

Proof. The ring A is a domain, so f is a nonzerodivisor. For $h \in A$,

$$h \in (f) \iff h(a_n) = 0 \text{ for every } n, \quad (37.6)$$

since then every coefficient of h vanishes at every integer and is divisible by $\sin(\pi z)$, whose zeros are simple and exactly the integers, with no new exponents; this is the plane counterpart of the kernel statement of `analytic:thm:fixed-quotient`. If $Sh \in (f)$ then $t^{-1/n}h(a_n) = 0$ in the field K_Γ for every n , so $h \in (f)$: multiplication by $\mathcal{S}$ is injective on $A/(f)$, and with $\mathfrak{a} \neq A$ this is regularity. If $fx + Sy = 0$ then $y = fa$ for some a , and cancelling f gives $x = -Sa$; the first map is injective, and the rest is the definition of $\mathfrak{a}$ and $\bar{A}$. Tensoring [\(37.5\)](#) with $\bar{A}$ kills both differentials, whose entries lie in $\mathfrak{a}$, so $\mathrm{Tor}_2^A(\bar{A}, \bar{A}) = \bar{A} \neq 0$ and $\mathrm{pd} \bar{A} = 2$. The truncation $0 \rightarrow A \rightarrow A^2 \rightarrow \mathfrak{a} \rightarrow 0$ gives $\mathrm{Tor}_1^A(\mathfrak{a}, \bar{A}) \simeq \bar{A} \neq 0$, so $\mathfrak{a}$ is not flat and has projective dimension exactly one. $\square$

The analytic-geometry report proves non-Noetherianity and the failure of the Nullstellensatz for its fixed-domain ring; it does not compute projective dimensions, and the merge found no such computation elsewhere in the collection.

37.3 Ultrafilter maximal ideals

For an ultrafilter $\mathcal{U}$ on $\mathbb{N}$ let $\mathbb{C}_{\mathcal{U}} = \mathbb{C}^{\mathbb{N}}/\mathcal{U}$ be the ultrapower field: sequences are identified when they agree on a member of $\mathcal{U}$.

Theorem 37.5 (Ultrafilter residue fields). *For every ordered extension $\mathbb{Q} \hookrightarrow \Delta$ and every nonprincipal ultrafilter $\mathcal{U}$ on $\mathbb{N}$,*

$$\Theta_{\mathcal{U}} : A_{\mathcal{H}_{\Delta}} \rightarrow \mathbb{C}_{\mathcal{U}}((t^{\Delta})), \quad \sum_{\gamma} F_{\gamma} t^{\gamma} \mapsto \sum_{\gamma} [(F_{\gamma}(a_n))_{n \geq 1}]_{\mathcal{U}} t^{\gamma}, \quad (37.7)$$

is a surjective ring homomorphism whose kernel is a maximal ideal containing $\mathfrak{a}$, with residue field exactly $\mathbb{C}_{\mathcal{U}}((t^{\Delta}))$. Different nonprincipal ultrafilters give different maximal ideals.

Proof. The output support is contained in the input support; at a fixed exponent Hahn multiplication involves finitely many pairs, and finite sums and products commute with the ultrapower quotient. For surjectivity, choose representing sequences for the coefficients of a target series and interpolate each by an ordinary entire function on the nodes; the functions assemble with the target’s support. The target is a field, so the kernel is maximal. The function f vanishes at every node, and the coefficient κ_k of $\mathcal{S}$ at $-1/k$ has node values δ_{nk} , which is zero in every nonprincipal ultrapower; so $\Theta_{\mathcal{U}}(f) = \Theta_{\mathcal{U}}(\mathcal{S}) = 0$. If $\mathcal{U} \neq \mathcal{V}$, pick $E \in \mathcal{U} \setminus \mathcal{V}$ and an entire g equal to 1 at a_n for $n \in E$ and 0 otherwise; then $\Theta_{\mathcal{U}}(g) = 1$ and $\Theta_{\mathcal{V}}(g) = 0$. $\square$

This is the plane counterpart of `analytic:thm:fixed-two-ideals(2)`, where coefficientwise reduction gives $\mathcal{O}(D)((t^{\mathbb{R}})) \rightarrow k_{\mathcal{U}}((t^{\mathbb{R}}))$ on the unit disk. The calculation is coefficientwise; it is not a limit of surcomplex points, and it does not classify the maximal ideals above $\mathfrak{a}$.

37.4 Taylor prolongation to finite halos

For a set-sized ordered group Δ containing the relevant exponents, $c \in \mathbb{C}$, $\varepsilon \in \mathbb{C}((t^{\Delta}))$ with $v(\varepsilon) > 0$, and $F = \sum t^{\gamma} F_{\gamma}$ with coefficients holomorphic near c , put

$$F^{\sharp}(c + \varepsilon) = \sum_{\gamma} \sum_{k \geq 0} t^{\gamma} \frac{F_{\gamma}^{(k)}(c)}{k!} \varepsilon^k. \quad (37.8)$$

This is the halo evaluation (2.7) extended from positive and integral families to arbitrary sections of $\mathcal{H}_{\Gamma}$: the support lies in $\text{supp } F + \langle \text{supp } \varepsilon \rangle_{\mathbb{N}}$, which is well ordered with finite contributions at each exponent by lemma 3.2 even when $\text{supp } F$ has negative elements, and ordinary Taylor identities make (37.8) a ring homomorphism. A finite surcomplex number is $c + \varepsilon$ with ordinary standard part c and infinitesimal ε ; its support, together with the exponents of the functions involved, lies in a set-sized ordered subgroup of $\mathbf{No}$, so (37.8) interprets statements at all finite surcomplex arguments without a class-indexed sum. Nothing is evaluated at infinite arguments.

Completion of the proof of theorem 37.3. If $c \notin \mathbb{Z}$, the constant coefficient of $f^{\sharp}(c + \varepsilon)$ is $\sin(\pi c) \neq 0$. If $c = a_n \in \mathbb{Z}$, the zero is simple and $f^{\sharp}(a_n + \varepsilon) = \varepsilon(f'(a_n) + \text{positive-valuation terms})$, the bracket being a unit, so $f^{\sharp}$ vanishes on that halo only at $\varepsilon = 0$, where $\mathcal{S}^{\sharp}(a_n) = t^{-1/n} \neq 0$. Every step survives enlargement of the value group. $\square$

The zero set of $f^{\sharp}$ on finite arguments is thus $\mathbb{Z}$, in agreement with the trigonometry report, whose canonical sine has zeros exactly $\pi\mathbb{Z}$ on its strip of finite imaginary part and is the Taylor lift of the ordinary sine at finite surcomplex arguments (`trigonometry:thm:striptrig`); at infinite arguments that report’s global sine vanishes exactly on $\pi\mathbf{Oz}$ (`trigonometry:cor:globalzeros`) [6], a region where this part evaluates nothing. Although $\mathcal{S}$ has no zero at an ordinary real point, $\mathcal{S}^{\sharp}$ does have zeros on finite halos; they are classified next, and none is a zero of $f^{\sharp}$.

38 Zeros of the cardinal-sine family on finite halos

Put

$$\widehat{\mathcal{S}} = t\mathcal{S} = \kappa_1 + \sum_{m \geq 2} t^{\theta_m} \kappa_m, \quad \theta_m = 1 - \frac{1}{m}. \quad (38.1)$$

The set $\{\theta_m : m \geq 2\} = \{1/2, 2/3, 3/4, \dots\}$ is positive and strictly increasing, so $\widehat{\mathcal{S}}$ has ordinary leading coefficient κ_1 and a positive-support remainder; multiplication by the monomial t does not change zeros.

Lemma 38.1 (A support-controlled implicit root). *Let $\mathcal{K}(Z) = k_0(Z) + \sum_{\gamma \in E} t^\gamma k_\gamma(Z)$ with $E \subseteq \Gamma_{>0}$ well ordered and all coefficients holomorphic near c , and suppose $k_0(c) = 0$ and $k'_0(c) = \mu \neq 0$. Then $\mathcal{K}^\sharp(c + \varepsilon) = 0$ has exactly one infinitesimal solution ε , zero allowed; it has support in $\langle E \rangle_{\mathbb{N}} \setminus \{0\}$, it is 0 when all $k_\gamma(c) = 0$, it is unique also among infinitesimals of every ordered value-group extension, and it is real when c and all Taylor coefficients are real.*

Proof. Write the equation as $\mu\varepsilon + \sum_{k \geq 2} a_k \varepsilon^k + \sum_{\gamma \in E} t^\gamma \sum_{k \geq 0} b_{\gamma k} \varepsilon^k = 0$ and seek $\varepsilon = \sum_{\eta \in M \setminus \{0\}} u_\eta t^\eta$ with $M = \langle E \rangle_{\mathbb{N}}$. By lemma 3.2 M is well ordered and the coefficient at a fixed η is a finite sum, in which the unknown u_η occurs only as μu_η : every other term has at least two positive factors or a positive γ , so it involves only coefficients at smaller exponents. Well-founded recursion determines each u_η uniquely; with no forcing term the recursion gives zero. For uniqueness among arbitrary infinitesimals x, y of an extension, factoring differences of powers gives $\mathcal{K}^\sharp(c + x) - \mathcal{K}^\sharp(c + y) = (x - y)(\mu + R)$ with $v(R) > 0$, legitimately by the support lemma, and $\mu + R$ is a unit. Real data give a real recursion. $\square$

This is the positive-support version of the simple-root implicit function construction, written as a recursion because sequential valuation convergence is not available at arbitrary rank; it is not a new Hensel theorem.

Theorem 38.2 (Exact finite-halo zero classification). *The zeros of $\mathcal{S}^\sharp$ at finite surcomplex arguments are exactly one real surreal point $z_n = a_n + \varepsilon_n$ for each $n \geq 2$. There is no zero in the halo of 0 and none whose standard part is not a nonzero integer. The displacement is*

$$\varepsilon_n = (-1)^{a_n+1} a_n \omega^{-(1-1/n)} + r_n, \quad v(r_n) > 1 - \frac{1}{n}, \quad (38.2)$$

valuation being read through $t^q = \omega^{-q}$ and r_n possibly zero. All these roots lie in $\mathbb{R}((t^{\mathbb{Q}}))$, and ordered value-group extension produces no further roots on finite halos.

Proof. The exponent-zero coefficient of $\widehat{\mathcal{S}}^\sharp(c + \varepsilon)$ is $\kappa_1(c) = \sin(\pi c)/(\pi c)$, with value 1 at 0, so a zero needs c to be a nonzero integer; these are the zeros of κ_1 and are simple. For $c = a_n$, $n \geq 2$,

$$\widehat{\mathcal{S}}(a_n) = t^{\theta_n}, \quad \mu_n = \kappa'_1(a_n) = \frac{(-1)^{a_n}}{a_n} \neq 0. \quad (38.3)$$

Lemma 38.1 with $E = \{\theta_m\}$ gives a unique real infinitesimal root supported in $\langle \{\theta_m\} \rangle_{\mathbb{N}} \subset \mathbb{Q}$. Below θ_n there is no forcing term, so by induction all coefficients vanish there; at θ_n the equation reads $\mu_n u_{\theta_n} + 1 = 0$, whence $u_{\theta_n} = -\mu_n^{-1} = (-1)^{a_n+1} a_n$, which is (38.2). Uniqueness in the lemma excludes other roots in the halo even over a larger group, and the leading-coefficient argument excludes every other standard part. $\square$

With (37.1), the first three roots are

$$z_2 = 1 + \omega^{-1/2} + \cdots, \quad z_3 = -1 - \omega^{-2/3} + \cdots, \quad z_4 = 2 - 2\omega^{-3/4} + \cdots,$$

the omitted terms having valuation greater than $1/2$, $2/3$, $3/4$. These are exact leading-term descriptions, not numerical approximations with a chosen large real ω . At the nodes themselves $\mathcal{S}$ is an infinite nonzero omnific integer.

A limitation on the word “entire”. Every coefficient of $\mathcal{S}$ is an ordinary entire function. That does not make $\mathcal{S}$ one power series strongly evaluable at every point of $\mathbb{C}((t^{\mathbb{Q}}))$, still less of $\mathbf{SC}$; it belongs to the common-domain ring $\mathcal{O}(\mathbb{C})((t^{\mathbb{Q}}))$ with the finite-halo prolongation (37.8), and theorem 38.2 is complete on that domain only. The strongly entire series of the entire-functions report form a different ring.

Where the classical theorems enter. Ordinary Weierstrass, Mittag–Leffler and interpolation theorems construct individual coefficient functions; they do not by themselves assemble outputs into one well-ordered Hahn support. Ordinary Oka and Cartan theorems belong to their own finite-dimensional analytic categories [23] and are not invoked for $\mathcal{H}_{\Gamma}$ because its coefficients are holomorphic; (37.4) shows concretely that a locally surjective map of free Hahn sheaves need not be surjective on global sections. Ordinary estimates are used only for the normal convergence of (8.4) and inside classical inputs; no numerical scale parameter tends to zero, and no inadmissible infinite Hahn family is exchanged with an ordinary sum.

38.1 Scope of Part V

Question	Proved in Part V	Not asserted
Pole-only localization	every fixed-nilpotent descending-scale family; the hidden example; the deep examples (merge)	arbitrary varying or noncommuting pole matrices; the diagonal examples (11.4)
Coefficient categories	holomorphic common-domain families; one monomial localization suffices	meromorphic coefficients as a substitute for holomorphic ones
Line bundles	integral Picard injection; plane isomorphism; persistent classes (global-divisor report)	a complete calculation of $H^1(\mathbb{C}, \mathcal{I}_{\Gamma})$
Extensions	an injected sequence quotient with free middle representatives	a classification of all self-extensions or all vector bundles
Omnific values	$\mathcal{S}(x) \in \mathbf{Oz}$ at every ordinary real x	an omnific-valued map at every surreal argument
Zeros and ideals	exact finite-halo zeros of $\mathcal{S}$; the pair $(f, \mathcal{S})$ has none; $\text{pd} = 2$; ultrafilter maximal ideals	a zero theorem at infinite arguments; a classical corona theorem; all maximal ideals above $\mathfrak{a}$
Verification	written proofs; 215 finite symbolic checks	peer review, Lean formalization, certified priority

The corresponding limitations are items 30–39 of section 30, and the questions left open are in section 31.

A A proof and support audit

The table records every place where an infinite formal expression occurs, together with the set that controls it and the ordinary analytic fact required. A support in the middle column is a *containing* support; cancellation may make the actual support smaller.

Part I constructions.

Construction	Support control	Ordinary analytic requirement
Positive matrix inverse, exp, log	Positive monoid generated by the input support; finitely many exponent words and finitely many matrix paths at each coefficient (lemmas 3.2 and 3.4)	Coefficients remain on the original domain.
Local polar factorization	$\langle \text{supp}(G - I_r) \rangle_{\mathbb{N}}$; well-founded triangular recursion	Laurent splitting on one fixed punctured disk, with no shrinking (lemma 4.1).
Necessity of Criterion P	All $P_a - I_r$ lie in the one monoid generated by $\text{supp}(B_0 - I_r)$	Right-gauge invariance and uniqueness of the local factorization.
Sufficiency of Criterion P	$\langle \mathcal{E}(G) \rangle_{\mathbb{N}}$, for the central matrix and for the Q_a	One ordinary isolated-singularity interpolation per matrix coefficient and exponent (lemmas 4.2 and 4.3).
Disk trivializers	Generated monoid of the global polar support together with that disk's own original support	Products of matrices holomorphic on that disk.
Right torsor of splittings	Finite union of the two solution supports, then positive monoid closure	Identity-theorem extension across each puncture.
Stein splitting	One monoid generated by all transition supports	Ordinary $H^1(X, \mathcal{O}^{r^2}) = 0$.
Deformation rigidity	The same positive monoid; ordinary transitions are exponent-zero coefficients	Ordinary $H^1(X, \mathcal{E}nd(\mathcal{V})) = 0$.
Rank-three obstruction	Forced descending set $\{1 + 1/n\}$	Only simple poles are used.
Depth- $(r - 1)$ obstruction	Lower partial-sum supports well ordered; final support $\{r - 2 + 1/n\}$ descends	Only simple poles and holomorphic constants are used.
Integral nontriviality	A hypothetical integral frame normalizes to a positive frame	Reduction of invertible matrices and gluing of the reduced frame.

Part II constructions.

Construction	Support control	Ordinary analytic requirement
Fundamental matrix on the cover	$\langle \text{supp } \Omega \rangle_{\mathbb{N}}$, for Y and for Y^{-1} ; the same bound for <i>every</i> holonomy matrix	Holomorphic primitives on a simply connected Riemann surface.
Dyson transport	Strong summability by the finite-word clause; each coefficient a finite sum of iterated integrals	Ordinary iterated path integrals along an ordinary path.
Necessity of Criterion M	$E_{\rho} \subseteq \langle \text{supp } \Omega \rangle_{\mathbb{N}} \setminus \{0\}$, uniformly in the group element	None beyond the fundamental-matrix theorem.
Sufficiency of Criterion M	$\langle E_{\rho} \rangle_{\mathbb{N}}$; transfinite recursion with no limit-ordinal passage	Ordinary Mittag–Leffler through the period map P , plus one fixed linear section σ (axiom of choice for infinite D).
Uniqueness of the normalized realizer	Least differing exponent in a well-ordered union	Injectivity of P on $\text{im } \sigma$.
Framed gauge and its inverse	$\langle \text{supp } \Omega \cup \text{supp } \Psi \rangle_{\mathbb{N}}$	Single-valuedness on U ; descent of coefficients from the cover.
Entire extension of a logarithmic gauge	The same monoid, unchanged by the extension	A simple pole with zero residue is removable.
Positive Frobenius form	$\langle E \rangle_{\mathbb{N}}$ for H and H^{-1}	Primitives on a disk; $H_{\beta}(0) = 0$ makes H_{β}/z holomorphic.
Traceless refinement	No change of support	Finite polynomial determinant identity; one entrywise linear section.
Finite exponent dependence	Finitely many decompositions and parts inside $\langle E \rangle_{\mathbb{N}}$ (lemma 3.5)	Finitely many rooted trees and ordinary operations per exponent.

Part III constructions.

Construction	Support control	Ordinary analytic requirement
A section $\sum f_\gamma t^\gamma$ of $\mathcal{H}_\Gamma$, $\mathcal{I}_\Gamma$ or $\mathcal{M}_\Gamma$	One well-ordered support and one ordinary coefficient domain, by definition	Identity theorem for holomorphic and meromorphic functions (lemma 20.2).
Products; scalar inverse near one	Hahn support addition; the positive-support Neumann lemma (lemmas 3.2 and 3.4)	None beyond the fixed domain.
$\exp(c_n N)$ and every matrix gauge	Finite polynomials, N being nilpotent; supports in $\{k\alpha_n\}$ or finite sums of such	None; no convergence and no infinite matrix exponential.
The local family $(c_n)_n$	Not a global Hahn sum; each member is a separate overlap section	h_n is holomorphic on $\mathbb{C} \setminus \{n\}$.
Zero-annihilator obstruction and classification necessity	An infinite decreasing set $\{\beta + \alpha_n\}$ or $\{\gamma + \alpha_n\}$ forced into one putatively well-ordered support; no summation is performed	$h_n g$ is not meromorphic at n for a nonzero meromorphic germ g (lemma 21.1).
Finite-puncture and sufficiency gauges	Finite sums of c_n or d_n	Finite sums of the h_j are holomorphic away from their centres.
Cyclic coefficientwise frame corrections	For each degree, later corrections leave its coefficient fixed; all supports in $\mathbb{N}\delta$, domains fixed	Ordinary $H^1(\mathbb{C}, \mathcal{O}) = 0$ on a fixed cover.
Triviality of pole data over $\mathcal{M}_\Gamma$	None needed: $B_0 = I$, $B_n = G_n$	Meromorphic extension of $G_n^{\pm 1}$ over D_n .

The Part III proofs never use a norm on the full surcomplex class, a real-valued valuation for an arbitrary-rank group, or a limit of bounded-domain trivializations.

Part V constructions.


Construction	Support control	Ordinary analytic requirement
Interpolation series $\mathcal{S} = \sum t^{-\alpha_j} \kappa_j$	The increasing sequence $\{-\alpha_j\}$; one coefficient per exponent	Weierstrass products and finite principal parts (lemma 34.1).
Disk frames $T_n = T_0 G_n$	$\{0\} \cup \{\alpha_n - \alpha_j\}$, increasing in j	Cancellation (iii)–(iv) of lemma 34.1 ; the determinant is one.
Symmetric powers	Degree- d polynomials in the entries; bounds multiplied by d	None.
Hidden and deep reductions	Finite unitriangular products; F' , Q'_n from the proof of theorem 9.2	lemma 4.2 through that proof.
Persistent line and extension classes	A forced descending set $\{\alpha_n\}$ in one global support	Nonremovability of a pole; the Cousin criterion of the global-divisor report.
Interpolation, ideal, ultrafilter maps	Value supports contained in input supports; no exponent created by evaluation	Ordinary interpolation on a discrete set; division by $\sin(\pi z)$.
Taylor prolongation and implicit root	$\text{supp } F + \langle \text{supp } \varepsilon \rangle_{\mathbb{N}}$; well-founded recursion on $\langle E \rangle_{\mathbb{N}}$	Taylor expansion at an ordinary point.

The indispensable hypotheses are these. The value group is a set and is totally ordered. Supports

are well ordered, not merely bounded below. Matrix rank is finite. The puncture set is ordinary closed and discrete. Coefficient functions share their stated ordinary domains. In Part I the gluing matrices reduce to the identity, or to a specified ordinary bundle in [theorem 11.2](#); in Part II the representation is near-identity with positive support; in Part III the transitions are finite exponentials of one constant nilpotent matrix on a fixed puncture-and-disk cover of the plane, the punctures and the functions h_n are fixed, and isomorphisms cover the identity of the base; in Part V the poles have finite order, the nilpotent direction is fixed, and prolongation is used only on finite halos. None of these hypotheses is replaced anywhere by a fine-topological convergence assumption.


B A dependency map

The Part I chain is



The arbitrary-cover Stein theorem is a separate positive branch whose only cohomological input is ordinary coherent-sheaf acyclicity. It is not used to justify the infinite star-cover criterion on the plane, whose interpolation step has its own direct proof; it is used for the statement about all bounded restrictions and for the universal cyclic-group result.

The Part II chain is



The gauge classification uses only the fundamental matrices; its entire extension uses only single-valuedness and removability of a simple pole with zero residue. The traceless refinement adds a finite polynomial determinant identity. The counterexamples use the necessity half of [theorem 14.2](#) and explicit support orderings. No implication in either chain depends on the outcome of a verification script.

The two chains meet only at [lemmas 3.2](#) and [3.4](#) and at the shared observation that the ordinary analytic input is applied one exponent at a time. Neither criterion is derived from the other.

The Part III chain is

an essential singularity cannot be cancelled by a meromorphic germ
 $\Downarrow$
 a nonzero global coefficient forces a decreasing Hahn support
 $\Downarrow$
 the global sections of a Jordan bundle form one invariant line
 $\Downarrow$
 all morphisms are constant-in-chart nilpotent intertwiners
 $\Downarrow$
 indecomposability and an intrinsic flag
 $\Downarrow$
 the rank-two gauge equation gives exact tail translation.

Here “constant-in-chart” means the same global Hahn matrix in every local frame, not constant as an ordinary function of z ; its coefficients can be arbitrary global holomorphic or meromorphic functions. The only general analytic input for the resistant examples is ordinary uniqueness of holomorphic and meromorphic continuation and the existence of the displayed essential singularities; the cyclic rigidity direction separately uses the ordinary additive Cousin theorem, through [theorem 11.1](#). The tensor and undecidability consequences add finite linear algebra and the classical halting theorem, respectively. The Part III chain uses [lemmas 3.2](#) and [3.4](#) and [lemma 11.3](#); it does not use Criterion P or Criterion M, and [remark 22.2](#) is an independent cross-check over $\mathcal{S}_\Gamma$ only.

The Part V chain is

Weierstrass and finite principal parts ([lemma 34.1](#))
 $\Downarrow$ with the Neumann support lemma
 determinant-one rank-two Laurent frames
 $\Downarrow$ symmetric powers; integral reductions of Part I
 fixed-nilpotent, hidden and deep examples trivial over $\mathcal{H}_\Gamma$
 $\Downarrow$ with the sine as the entire function
 free-middle extensions, the proper ideal, $\text{pd} = 2$, ultrafilter ideals
 $\Downarrow$ with Taylor prolongation and a well-founded recursion
 finite-halo zeros of the sine family.

The rank-one persistence and the interpolation criterion are cited from the global-divisor report and are not links of this chain. The reduction of the hidden and deep examples uses the factorizations of [section 8](#) and the proof of [theorem 9.2](#), but not Criterion P.

C Provenance, reproduction, and the merge record

C.1 Sources of this report

This report merges four independently written manuscripts. The first two are dated September 21, 2026, the third September 22, 2026, the fourth September 23, 2026; each was delivered with

its own code and recorded output, preserved unchanged in `code/` and `data/` of this directory. The manuscripts themselves are not shipped with this report.

- *Nonabelian Support Obstructions in Surcomplex Analysis: A Sharp Matrix Cousin Criterion and Arbitrarily Deep Unipotent Examples*, from the archive `surcomplex_nonabelian_support`; source file `08-polar-support-matrix-cousin.tex`, delivered with its README. It contributes Part I entire, the shared coefficient category and support calculus, the Laurent and isolated-singular-part inputs, the normalized polar factorization, the exact Criterion P with its invariance corollaries and torsor statement, the rank-one recovery, the rank-three obstruction with its repair and logarithm diagnostic, the chain identity and depth hierarchy, the integral vector-bundle statements with the localization boundary, and the Stein splitting, deformation rigidity and cyclic dichotomy.
- *Global Hahn Support and the Surcomplex Riemann–Hilbert Problem: Noncommutative monodromy, countably many singularities, and an exact realization obstruction*, from the archive `surcomplex_riemann_hilbert`; source file `09-support-monodromy-realization.tex`, delivered with its README and its `BUILD_AND_PROVENANCE.json`. It contributes Part II entire: strong summability and the finite-decomposition lemma, the free-meridian lemma, the period map with its linear section, the fundamental-matrix and triangular lemmas, Criterion M with its intrinsic and conjugation-invariance forms, the framed gauge classification and entire extension, the positive Frobenius form, the finite Fuchsian specialization and the SL_r refinement, the three calibrating examples, the mixed commutator correction with its $\mathbb{Q} + \mathbb{Q}\omega$ specialization, the finite-dependence and base-change propositions, and the impossibility of a continuous linear period right inverse.
- *Localization-resistant bundles in Hahn-coherent surcomplex analysis: Essential singularities, nilpotent representations, and exact isomorphism classification*, from the archive `surreal_localization_bundles`; despite the archive name its objects are surcomplex, complex-valued Hahn coefficients read in `No[i]`. Called *the localization manuscript* in the text, it is a 28-page article delivered with a README, a provenance record, a SHA-256 manifest, a verification script with its requirement pin, and three data files, placed here as `code/10-localization-bundles-verify.py` and `data/10-localization-bundles-*`. It contributes Part III entire: the meromorphic sheaf and its unit and sheaf lemmas, the zero-annihilator obstruction, the nilpotent bundles with their sections, stable nontriviality, bounded triviality and resistance to group enlargement, the morphism formula, indecomposability, the fully faithful realization and the intrinsic flag, the profile classification with its corollaries and examples, the localization-stable dichotomy, the continuum count, the tensor rule, the pole proposition and the essential-singularity generalization. It also contributes the undecidability theorem of [section 28.5](#), the Part III check suite of [section 29](#), the Part III rows of [appendices A](#) and [B](#), items [23–28](#) of [section 30](#) with its additions to the generic items there, and three questions of [section 31](#) (the full extension category, several independent descending profiles, and other ordinary bases).
- *Pole Localization and Omnific Interpolation: Explicit bundle trivializations, persistent line classes, and a surcomplex Bézout obstruction*, from the archive `pole_localization_research`. Called *the pole-localization manuscript* in the text, it is a 27-page article delivered with a README, a source audit, a SymPy verification script with its recorded results and requirement pin, a layout record and a SHA-256 manifest. The script, its record, the pin and the layout record are placed here as `code/11-pole-localization-verify.py` and `data/11-pole-localization-*`, and the source audit as `11-pole-localization-SOURCE_AUDIT.md` at the top of this directory. It contributes Part V except [corollary 35.2](#) and the merge observations listed

below: the pole-cancellation lemma, the rank-two, balanced, small-budget and fixed-nilpotent theorems, the trivialization of the hidden example, the Picard and line-bundle statements (credited to the global-divisor report), the free-middle extensions, the omnific sine family, the interpolation criterion (credited), the Bézout obstruction, the resolution, the ultrafilter ideals, the prolongation, the implicit root lemma and the zero classification. It also contributes the Part V check suite of [section 29](#), items 30–38 of [section 30](#) with its additions to the generic items there, its formalization route in item 11, and nine questions of [section 31](#).

Where the two manuscripts proved the same fact — the Neumann support calculus as an input, and the positive inversion, exponential and logarithm lemma — the statement is printed once, in [lemmas 3.2](#) and [3.4](#), and both members verified it independently in the same form. Where they supplied genuinely different routes to a needed ingredient, both routes are kept and marked: the ordinary Mittag–Leffler input appears twice, as the unbounded-order isolated-singular-part construction [lemmas 4.2](#) and [4.3](#) and as the linear period section [lemma 4.5](#), for the reasons given at the end of [section 4](#); and the impossibility of repair by enlarging the value group is proved twice, for two different objects, in [corollary 7.3](#) and [proposition 14.5](#). The local factorizations of [theorem 5.1](#) and [theorem 5.4](#) are likewise two different normalizations of two different objects, and neither is derived from the other here.

For Part III the merge prints once what the localization manuscript shares with the rest of the collection: the Neumann calculus and positive inverse ([lemmas 3.2](#) and [3.4](#)); the ordered-group lemma, which is [lemma 11.3](#) and also the global-divisor report’s `global:lem:descending`, the localization manuscript’s proof being a third copy of the same argument; the sheaf property, extended to $\mathcal{M}_\Gamma$ in [lemma 20.2](#); and the unit lemma, whose holomorphic cases are the global-divisor report’s unit criterion. Two genuinely different routes are kept and marked. Bounded integral triviality is proved by explicit finite gauges in [theorem 22.3](#) and by the Stein theorem in [corollary 10.3](#). Integral nontriviality of the new bundles follows both from the zero-annihilator argument and from Criterion P ([remark 22.2](#), a merge observation). The cyclic direction of [theorem 25.1](#) is printed through [theorem 11.1](#); the localization manuscript’s q -adic version of the same recursion is described in that proof and not printed a second time.

The merge made these choices. Part III is placed after Part II and before the effectivity part, which is renumbered as Part IV. The undecidability theorem is moved into the effectivity section, the manuscript’s check table into [section 29](#), its table of infinite expressions into [appendix A](#), its dependency chain into [appendix B](#), and its scope statements and open questions into [sections 30](#) and [31](#). Its summary theorem is kept as [theorem 19.1](#). Colliding symbols are renamed in the new material only, as listed in [section 20.5](#); in particular the meromorphic sheaf is written $\mathcal{M}_\Gamma$, the global-divisor report’s name for the same recipe. The merge adds four observations that the localization manuscript does not state: [remark 22.2](#), [corollary 27.2](#) with the discussion after it, the relation of $\mathcal{M}_\Gamma$ to the other reports in [section 20](#), and item 29 of [section 30](#).

For Part V the merge made these choices. Part V is placed after Part IV, not before it, so that every section, statement and item number of Parts I–IV is unchanged; its checks, limitations and questions are added to [sections 29](#) to [31](#), and its rows to [appendices A](#) and [B](#). Colliding symbols are renamed in the new material only ([section 33.3](#)), and its gluing convention is reconciled there with those of Parts I and III. Results the collection already had are printed once: the Picard decomposition and the persistence of pole-data line classes ([proposition 36.1](#) and [corollary 36.2](#)) are reduced to citations of the global-divisor report with the manuscript’s finite-order generalization kept, and the interpolation criterion ([proposition 37.2](#)) is stated with a citation to `global:cor:interp` and not reproved. The Bézout, corona and ultrafilter statements are kept with their proofs, because they are plane versions, not copies, of the analytic-geometry report’s unit-disk results, which are

credited at each. The manuscript’s direct proof of integral nontriviality is kept as a second route beside Criterion P. The manuscript’s citation of course notes for the classical analytic inputs is kept, and a textbook reference is added. The merge adds [corollary 35.2](#) with its proof, the consequences (a)–(d) of [section 35](#), the explicit choice $1/\Gamma_E(2-z)$ in [theorem 35.1](#), the credits to the global-divisor, analytic-geometry, trigonometry and entire-functions reports, the question on the diagonal examples, and item 39; and it updates every statement of Parts I–IV that called the monomial-localization question open. No earlier statement is retracted.

C.2 Repository provenance

Repository: <https://github.com/VladimirReshetnikov/Surreal>, reviewed at commit 39f2be6667ade51bca2b45daa47e289d69c09764, inspected September 21, 2026. No repository files were changed by either source manuscript and nothing was pushed by them. The report catalogue [1] supplies the report map and the distinctions between coefficient rings, which both members call essential. The global-divisor report [2] is the source of the motivating open direction and of the scalar results we recover rather than reprove; the differential-equations report [3] supplies the forward monodromy map and the ordinary-paths wording. The analysis, analytic-geometry and contour packages [4, 5, 8] are credited by Part II as antecedents for scalar global support obstructions and for the coefficientwise differential and integration viewpoint; they are not contributions claimed here, and no draft ring theorem or Hahn sheaf-cohomology theorem from them is used in any proof.

Neither source manuscript conducted an exhaustive audit of every source manuscript in the repository, and one repository code-search response was marked incomplete; its empty result is not evidence of absence. The comparison above is with the documented contents actually reviewed.

The localization manuscript reviewed the same repository at commit

0097304c7ae9d5de46d2ea342e2494f8fc126303,

inspected September 22, 2026, and records that it changed nothing there. By its provenance record it inspected lines 1–220 of the documentation index, this report’s README, lines 1900–2150 and 2880–3070 of this report’s article as it then stood, and a truncated README of the global-divisor report. Between that commit and the tree into which Part III was merged (commit e9a9650), the article texts of this report, of the global-divisor report and of the analysis report did not change; the documentation index did change, but its four-ring table was present throughout. The comparisons with the global-divisor and analysis reports made in Part III were made by the merge against that tree.

The pole-localization manuscript reviewed the repository at commit

0865f043aec113c14c69ef45006bbc7546a4e75a,

inspected September 23, 2026. By its source audit it read the repository README, the documentation inventory, selected manifest entries, and lines 1–210, 510–800 and 1160–1420 of this report’s article, whose blob c2dc2f9 it records; that blob is exactly the text of Parts I–IV into which Part V was merged. It audited no other report and no Lean declaration. The global-divisor, analytic-geometry, trigonometry and entire-functions reports cited in Part V did not change between that commit and the tree into which Part V was merged; the comparisons with them were made by the merge.

C.3 Building this report

The source is standalone, with an internal bibliography, no external bibliography file, no graphics and no separately distributed fonts. A standard TeX Live installation builds it with

```
latexmk -pdf -interaction=nonstopmode article.tex
```

Alternatively, run `pdflatex` with `-interaction=nonstopmode -halt-on-error` three times to settle the cross-references and contents.

C.4 Rerunning the delivered checks

The four scripts are preserved under `code/` with their original contents and their recorded outputs under `data/`. The Part I script needs Python 3.10 or later with SymPy; the Part II script needs NumPy, SciPy and SymPy. The Part III script needs Python 3.9 or later and SymPy, pinned at 1.14.0 in `data/10-localization-bundles-requirements.txt`. From this report directory, run copies in a temporary directory:

```
check_dir=$(mktemp -d)
cp code/*-verify*.py "$check_dir/"
cd "$check_dir"
python 08-polar-support-matrix-cousin-verify.py
python 09-support-monodromy-realization-verify-examples.py
python 10-localization-bundles-verify.py --max-rank 6 \
    --output fresh-verification.json
python 11-pole-localization-verify.py \
    --output fresh-pole-verification.json
```

The Part V script needs Python 3.9 or later and SymPy, pinned at 1.14.0 in `data/11-pole-localization-requirements.txt`; without `--output` it writes `verification_results.json` beside itself. The Part III script writes only to the path given with `--output`; the recorded files are therefore preserved by an in-place run, and the command above writes a fresh file for comparison. It makes no network calls.

Two operational cautions about the Part II suite, both verified directly rather than taken from a README.

First, the Part II script writes `verification_results.json` beside itself. An in-place run in the maintained layout writes under `code/`; in the original archive layout it could overwrite the historical record. Running a temporary copy keeps generated evidence separate from both the script and the delivered record.

Second, the Part II README as delivered lists a file `SHA256SUMS.txt` among the archive contents. That file is *not* present in the delivered directory and is not present here, so no checksum verification of the original archive is possible from what was shipped.

An independent rerun of both suites, on copies and not in the extraction tree, found: the Part I suite exits zero and prints its four PASS lines and `SUCCESS: 2899 exact checks passed`, identical in count to the recorded run, under Python 3.14.4 with SymPy 1.14.0 against the recorded Python 3.13.5 with SymPy 1.14.0, the version banner being the only differing line; and the Part II suite exits zero with all eight exact checks and all four numerical error checks passing. The recorded Part II build metadata claims zero warnings, zero overfull or underfull boxes, zero undefined references, a

25-page document, and a Poppler render review of all pages, and records `formal_verification: false, priority_certified: false, peer_reviewed: false`. Those three flags remain false for this merged report.

For Part III: the localization manuscript’s SHA-256 manifest, which is not shipped, verified all nine of its entries at merge time, and the six files placed here from that archive match it byte for byte. An independent rerun of its script on a copy, under Python 3.14.4 with SymPy 1.14.0, exited zero and reproduced the recorded JSON exactly except for the Python version field, with 12,355 exact checks passed. The file `data/10-localization-bundles-build_qa.json` describes the build of that manuscript’s own 28-page PDF, which is not shipped; it is layout quality assurance, not theorem verification, and says nothing about this merged document. The provenance record `data/10-localization-bundles-PROVENANCE.json` lists 12,355 finite exact checks with result PASS, peer review false, proof-assistant verification false, and no redistribution of external source files.

For Part V: the pole-localization manuscript’s SHA-256 manifest, which is not shipped, verified all eight of its entries at placement, and the five files placed here from that archive are byte-identical to the delivered ones. An independent rerun of its script on a copy, under Python 3.14.4 with SymPy 1.14.0, exited zero with all 215 exact checks passed and reproduced the recorded JSON except for the Python version field. The file `data/11-pole-localization-build_report.json` describes the build of that manuscript’s own 27-page PDF, which is not shipped; it is layout quality assurance, not theorem verification.

References

- [1] V. Reshetnikov, *Surreal repository: report catalogue*, `docs/README.md`, commit `39f2be6667ade51bca2b45daa47e289d69c09764`, inspected September 21, 2026.
<https://github.com/VladimirReshetnikov/Surreal/blob/39f2be6667ade51bca2b45daa47e289d69c09764/docs/README.md>.
- [2] *Global Support Obstructions: Moving Divisors, Mittag–Leffler, and a Picard Dichotomy*, AI-assisted merged research draft in V. Reshetnikov’s *Surreal* repository, `docs/surcomplex/global-divisors/`. Pinned commit as above; see especially the exact Cousin criterion (Theorem 13.1), the Picard-group vanishing dichotomy, and the subsection “Further precise directions”. Part III refers to its text at commit `0097304c7ae9d5de46d2ea342e2494f8fc126303`, identical to the current tree, where its compact-base part is present; its labels are quoted there verbatim.
<https://github.com/VladimirReshetnikov/Surreal/blob/39f2be6667ade51bca2b45daa47e289d69c09764/docs/surcomplex/global-divisors/article.tex>.
- [3] *Differential Algebra and Differential Equations over the Surreal and Surcomplex Numbers*, AI-assisted merged research draft in V. Reshetnikov’s *Surreal* repository, `docs/surcomplex/differential-equations/`. Pinned commit as above; see especially the theorem “Monodromy valued in $GL_d(K_\Gamma)$ ” and the ordinary-path remark following it.
<https://github.com/VladimirReshetnikov/Surreal/blob/39f2be6667ade51bca2b45daa47e289d69c09764/docs/surcomplex/differential-equations/article.tex>.
- [4] *Surcomplex analysis: holomorphic functions on $\mathbf{No}[i]$* , `docs/surcomplex/analysis/README.md`, same repository and commit. AI-assisted, unrefereed research documentation. The class $\mathcal{M}(U)$ cited in Part III is defined in its `article.tex` (label `d:def:clusterres`) at that commit and in the current tree.

- [5] *Infinitesimal analytic geometry over the surcomplex numbers: radius-free germs, a Nullstellensatz, and finite maps*, docs/surcomplex/analytic-geometry/README.md, same repository and commit. AI-assisted, unrefereed research documentation.
- [6] *Surcomplex trigonometry*, docs/surcomplex/trigonometry/, same repository. AI-assisted, unrefereed research documentation. Part V cites its labels `trigonometry:thm:striptrig` and `trigonometry:cor:globalzeros` as they stand in the tree into which Part V was merged.
- [7] *Entire functions at arbitrary rank*, docs/surcomplex/entire-functions-at-arbitrary-rank/, same repository. AI-assisted, unrefereed research documentation. Part V cites its label `ent:thm:bezout`.
- [8] *Contours and Stokes: Jordan separation, Cauchy theory, and Stokes calculus on the surcomplex plane*, docs/surcomplex/contours-and-stokes/README.md, same repository and commit. AI-assisted, unrefereed research documentation.
- [9] B. H. Neumann, On ordered division rings, *Transactions of the American Mathematical Society* **66** (1949), 202–252. <https://www.jstor.org/stable/1990552>.
- [10] B. Poonen, *Maximally complete fields*, L’Enseignement Mathématique (2) **39** (1993), 87–106. DOI: <https://doi.org/10.5169/seals-60414>.
- [11] S. L’Innocente and V. Mantova, A factorisation theory for generalised power series and omnific integers, *Advances in Mathematics* **442** (2024), 109513; arXiv:1710.07304. Used in Part V only for the normal-form description of omnific integers. <https://arxiv.org/abs/1710.07304>.
- [12] L. van den Dries and P. Ehrlich, Fields of surreal numbers and exponentiation, *Fundamenta Mathematicae* **167** (2001), 173–188. <https://doi.org/10.4064/fm167-2-3>.
- [13] V. Mantova and M. Matusinski, *Surreal numbers with derivation, Hardy fields and transseries: a survey*, arXiv:1608.03413. <https://arxiv.org/abs/1608.03413>.
- [14] O. Bournez and Q. Guilmant, *Surreal fields stable under exponential and logarithmic functions*, preprint (2022), arXiv:2201.08199. Cited for normal-form and set-sized-surreal-field background only. <https://arxiv.org/abs/2201.08199>.
- [15] K. Ebrahimi-Fard, L. Guo, and D. Kreimer, Spitzer’s identity and the algebraic Birkhoff decomposition in pQFT, *Journal of Physics A: Mathematical and General* **37** (2004), 11037–11052. Cited solely as the closest algebraic antecedent for recursive Birkhoff factorization in complete filtered noncommutative algebras and for the necessity of commutator corrections; no application of that algebra is imported here. <https://arxiv.org/abs/hep-th/0407082>.
- [16] K. Ebrahimi-Fard, D. Manchon, and F. Patras, *A noncommutative Bohnenblust–Spitzer identity for Rota–Baxter algebras solves Bogoliubov’s recursion*, arXiv:0705.1265. <https://arxiv.org/abs/0705.1265>.
- [17] H. Behnke and K. Stein, Entwicklung analytischer Funktionen auf Riemannschen Flächen, *Mathematische Annalen* **120** (1947/1949), 430–461. <https://doi.org/10.1007/BF01447838>.
- [18] H. Cartan, Variétés analytiques complexes et cohomologie, in *Colloque sur les fonctions de plusieurs variables*, Brussels, 1953, 41–55. Classical source for coherent-sheaf acyclicity on Stein manifolds.
- [19] M. M. Schmitt, *Kohomologie mit Schranken und Fortsetzung holomorpher Funktionen durch lineare stetige Operatoren*, doctoral thesis, arXiv:math/0202263 (2002). Includes a proof of Cartan’s theorem B for coherent subsheaves of finite free holomorphic sheaves and coefficientwise coboundary constructions. <https://arxiv.org/abs/math/0202263>.
- [20] O. Wone, *The Birkhoff–Grothendieck theorem*, arXiv:2304.08037, version 2. Historical and mathematical exposition of bundle splitting and Birkhoff factorization; its usual single-loop setting differs from the infinitely many independently supported gluing data treated here. <https://arxiv.org/abs/2304.08037>.

- [21] M. Greenfield, *Constructing holomorphic functions*, Rutgers University, Math 503 course notes, modified December 1, 2007. Cited by the pole-localization manuscript for the ordinary Weierstrass, Mittag–Leffler and discrete-interpolation constructions.
https://sites.math.rutgers.edu/~greenfie/mill_courses/math503a/construction.html.
- [22] W. Rudin, *Real and Complex Analysis*, 3rd ed., McGraw–Hill, 1987, Chapters 13 and 15 (Mittag–Leffler and Weierstrass theorems). Added by the merge as a textbook reference for the same classical inputs; cited at chapter level, not re-verified against a particular theorem number.
- [23] F. Forstnerič and J. Prezelj, Oka’s principle for holomorphic fiber bundles with sprays, *Mathematische Annalen* **317** (2000), 117–154. Cited in Part V for the ordinary analytic context only; none of its results is transferred to the Hahn sheaf.
- [24] K.-T. Chen, *Iterated path integrals*, Bulletin of the American Mathematical Society **83** (1977), 831–879. DOI: <https://doi.org/10.1090/S0002-9904-1977-14320-6>.
- [25] P. Deligne, *Équations différentielles à points singuliers réguliers*, Lecture Notes in Mathematics, vol. 163, Springer, 1970. A classical reference point, not a theorem reproved here in its original generality. DOI: <https://doi.org/10.1007/BFb0061194>.
- [26] Y. Ilyashenko and S. Yakovenko, *Lectures on Analytic Differential Equations*, Graduate Studies in Mathematics, vol. 86, American Mathematical Society, 2008. DOI: <https://doi.org/10.1090/gsm/086>.
- [27] P. Ehrlich, The absolute arithmetic continuum and the unification of all numbers great and small, *Bulletin of Symbolic Logic* **18** (2012), 1–45. DOI: <https://doi.org/10.2178/bsl/1327328438>.
- [28] M. F. Atiyah, Vector bundles over an elliptic curve, *Proceedings of the London Mathematical Society* (3) **7** (1957), 414–452. Cited in Part III as a precedent for indecomposable unipotent bundles with a related tensor pattern, on a different base. DOI: <https://doi.org/10.1112/plms/s3-7.1.414>.
- [29] H. Grauert, Analytische Faserungen über holomorph-vollständigen Räumen, *Mathematische Annalen* **135** (1958), 263–273. DOI: <https://doi.org/10.1007/BF01351803>.
- [30] A. M. Turing, On computable numbers, with an application to the Entscheidungsproblem, *Proceedings of the London Mathematical Society* (2) **42** (1937), 230–265; received and read in 1936. DOI: <https://doi.org/10.1112/plms/s2-42.1.230>.
- [31] J. Müller and A. Strohmaier, *The theory of Hahn-meromorphic functions, a holomorphic Fredholm theorem, and its applications*, Analysis & PDE **7** (2014), no. 3, 745–770. Their analytic category is explicitly not identified with ours; see item 21 of section 30. DOI: <https://doi.org/10.2140/apde.2014.7.745>.