

A countable genetic cut across the finite–infinite gap

Counterexamples concerning uniqueness of primitives
and the sup property of entire genetic functions

Research note prepared for Vladimir Reshetnikov

20 September 2026

Abstract

Put $q(t) = t/(1 + t^2)$ and form the countable Conway cut

$$T(x) = \{ q(n - x) : n \in \mathbb{N} \mid \}, \quad B(x) = 1 - T(x).$$

We prove that B is the indicator of the positive infinite surreal numbers. The option functions are globally defined rational functions, so this is an admissible genetic construction under the set-sized option rules in Rubinstein–Salzedo–Swaminathan and in Berenbeim’s later formulation; the latter’s global cofinality requirement is checked directly. Yet B is locally constant everywhere and nonconstant globally. Consequently x and $x - B(x)$ are genetic primitives of 1 that agree on every real argument but do not differ by a constant. The second is an increasing bijection of **No**. A bounded sublevel class of B has no surreal supremum, giving a second counterexample. We extend the construction to arbitrary nonempty sets of surreals without a largest element, construct ordinal-indexed independent families in the kernel of pointwise differentiation, and explain precisely why real-variable uniqueness arguments do not apply. The accompanying standard-library Python program performs exact finite checks, not a formal verification of the infinitary proofs.

Status and provenance. The targets are Conjecture 53 of [1] and Chapter 8, Conjecture 3 of [2]. The piecewise primitive itself already appears in [1]; the contribution here is the explicit genetic formula and the deductions from it. The arguments below give a negative answer under the published option conventions. No later resolution was located in the searches recorded with this note, but those searches do not establish priority or exclude unpublished work. The proof has not been independently refereed or checked in a proof assistant.

Contents

1	The published questions and the intended scope	2
2	Cuts, infinitude, and the derivative in use	3
3	A countable cut that detects positive infinity	4
4	Why the construction is genetic	5
5	Nonunique primitives, even with the same real values	6
6	Failure of the sup property	8
7	Every set-generated endpoint-free cut gives a variant	8
8	Order automorphisms and an ordinal-indexed kernel	10
9	Why familiar uniqueness arguments do not apply	11
10	An ordered-field analogue and the hyperreal distinction	13
11	Finite truncations and exact computational checks	13
12	Conclusions and remaining questions	15
A	Proof dependency and objection audit	15
B	Source locations and status of the search	16

Reading guide. Sections 2–5 contain the complete central argument. Section 4 is essential: a nonconstant zero-derivative function alone would not answer the genetic-function conjecture. Section 6 treats the second target. The remaining sections give extensions, failure analyses, and a precise account of what the computational checks establish.

1 The published questions and the intended scope

A primitive of a function f is a function F with $F' = f$. The question studied here asks whether restricting both f and its primitives to *genetic* surreal functions restores uniqueness up to a constant. This is the substance of Rubinstein-Salzedo and Swaminathan’s Conjecture 53 [1, p. 37]. Berenbeim’s later Conjecture 3 instead asks whether every entire genetic function has the *sup property* [2, pp. 225–226].

The term *genetic* refers to specified constructions of functions using Conway cuts, previously available functions, and recursion on simpler arguments. It does not mean merely “definable,” and it does not by itself mean analytic, continuous, or finitely presented. The distinction matters: a

direct piecewise definition is not, without further work, a certificate of membership in a genetic class.

Our main construction requires only a particularly small part of the published machinery: rational functions with no poles, a countable fixed family of options, a single nonrecursive cut, and ordinary arithmetic on the resulting functions. We state the operational requirements explicitly in Section 4 instead of assuming that every reasonable piecewise function is genetic.

There are two separate issues of validation. The mathematical issue is whether the displayed construction has the stated properties under the specified rules. That is addressed by proofs. The historical issue is whether this particular construction or an equivalent argument is already known. The searches supporting this note locate the published conjectures and their relevant definitions; they do not settle historical priority.

2 Cuts, infinitude, and the derivative in use

2.1 The small amount of surreal theory needed

The surreal numbers form a proper-class ordered field. If L and R are *sets* of surreal numbers and every element of L is less than every element of R , then $\{L \mid R\}$ is the simplest surreal strictly between them. Empty option sets are allowed. We use the standard simplicity rule, in particular $0 = \{\mid\}$ and $1 = \{0 \mid\}$; these conventions are recalled in [1, Section 2.1].

The braces in $\{L \mid R\}$ do *not* denote an ordinary real supremum. For example, a set of negative left options may approach 0 in various senses, but if there are no right options, 0 itself is already the simplest number above all of them. Conversely, one nonnegative left option excludes 0 even if that option is an infinitesimal.

All algebra below is *surreal field algebra*. In particular, $\omega - 1$ is meaningful in **No** and is not an ordinal predecessor of ω . Whenever ordinal exponents are used later, ω^α denotes the usual embedded ordinal monomial. Neither noncommutative ordinal addition nor ordinal subtraction is being used in the rational formulas.

Definition 2.1. Throughout, $\mathbb{N} = \{0, 1, 2, \dots\}$ denotes the ordinary, finite natural numbers. Define

$$\mathcal{U} = \{x \in \mathbf{No} : (\forall n \in \mathbb{N}) x > n\}, \quad (2.1)$$

$$\mathcal{D} = \{x \in \mathbf{No} : (\exists n \in \mathbb{N}) x \leq n\}, \quad (2.2)$$

$$\mathcal{O} = \{x \in \mathbf{No} : (\exists n \in \mathbb{N}) |x| \leq n\}. \quad (2.3)$$

Thus $\mathcal{U}$ is the class of positive infinite numbers, $\mathcal{O}$ is the class of finite numbers, and $\mathcal{D} = \mathbf{No} \setminus \mathcal{U}$ contains both the finite numbers and all negative infinite numbers.

There is no surreal number sitting at the boundary between $\mathcal{D}$ and $\mathcal{U}$. The left class has no greatest element: from $x \leq n$ we get $n + 1 > x$ with $n + 1 \in \mathcal{D}$. The right class has no least element: if $u \in \mathcal{U}$, then $0 < u/2 < u$ and $u/2 \in \mathcal{U}$, since $u > 2n$ for every natural n . We shall refer to this boundary as the *finite–infinite gap*. The word “finite” in this name is not a claim that every member of $\mathcal{D}$ is finite.

A useful stronger separation is

$$d \in \mathcal{D}, u \in \mathcal{U} \implies u - d \in \mathcal{U}. \quad (2.4)$$

Indeed, choose n with $d \leq n$. For any $m \in \mathbb{N}$ we have $u > m + n$, so $u - d \geq u - n > m$.

2.2 Local differentiation, not a derivation of field elements

For a function on a neighborhood of x , use

$$F'(x) = \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h}. \quad (2.5)$$

An order-field ε - δ limit quantifies over all positive surreal ε and δ , not only positive real ones. This is the pointwise derivative at issue in [1, Section 5]. Our difference quotients will be exactly constant on a punctured neighborhood, so there is no subtle convergence claim to establish.

This operator acts on *functions*. It should not be confused with a chosen derivation $\partial : \mathbf{No} \rightarrow \mathbf{No}$ on the surreal field itself. The assertion $F'(x) = 1$ concerns the variation of F with its argument; it is not an assertion about the value of a field derivation at the number $F(x)$. No result about a distinguished field derivation is being challenged.

3 A countable cut that detects positive infinity

Lemma 3.1 (A bounded sign-preserving rational function). *In every ordered field, the function*

$$q(t) = \frac{t}{1+t^2} \quad (3.1)$$

is everywhere defined, has the same sign as t , and satisfies $-1 < q(t) < 1$. In fact, $|q(t)| \leq \frac{1}{2}$.

Proof. The denominator is positive. For the strict bounds, the identities

$$1+t^2-t = (t-\frac{1}{2})^2 + \frac{3}{4} > 0, \quad 1+t^2+t = (t+\frac{1}{2})^2 + \frac{3}{4} > 0$$

give $q(t) < 1$ and $q(t) > -1$. The sharper bound follows from $(|t| - 1)^2 \geq 0$, equivalently $2|t| \leq 1+t^2$. $\square$

Lemma 3.2 (A binary cut test). *Let L be a set of surreal numbers, all strictly less than 1. If every member of L is negative, then $\{L \mid\} = 0$. If L contains a nonnegative member, then $\{L \mid\} = 1$.*

Proof. In the first case 0 lies above every left option and is the simplest surreal of all. In the second case 0 is excluded, whereas 1 lies above all left options. The only other number born on the day when 1 is born is -1 , and it is excluded by the nonnegative option. Thus 1 is the simplest admissible number. The first case also covers $L = \emptyset$. $\square$

Definition 3.3. For every $x \in \mathbf{No}$, set

$$T(x) = \left\{ \frac{n-x}{1+(n-x)^2} : n \in \mathbb{N} \mid \right\}, \quad B(x) = 1 - T(x). \quad (3.2)$$

For each fixed x , the left side of this cut is a countable set of surreal numbers. There are no right options, so the order condition is vacuous. The definition therefore produces a surreal number at every input. The natural-number quantifier is part of the option indexing; it is not a limit operation and it is not replaced by an ordinal-length summation.

Theorem 3.4 (Exact evaluation). *The functions in definition 3.3 satisfy*

$$T(x) = \begin{cases} 1, & x \in \mathcal{D}, \\ 0, & x \in \mathcal{U}, \end{cases} \quad B(x) = \begin{cases} 0, & x \in \mathcal{D}, \\ 1, & x \in \mathcal{U}. \end{cases} \quad (3.3)$$

Proof. Every left option lies in $[-\frac{1}{2}, \frac{1}{2}]$, by lemma 3.1. If $x \in \mathcal{D}$, choose $n \in \mathbb{N}$ with $x \leq n$. Then $n - x \geq 0$, so the corresponding left option is nonnegative. The second case of lemma 3.2 gives $T(x) = 1$.

If $x \in \mathcal{U}$, then $n - x < 0$ for every $n \in \mathbb{N}$. All left options are negative, so the first case gives $T(x) = 0$. Subtracting from 1 gives the formula for B . $\square$

Example 3.5. The distinction remains exact for infinitesimals and infinite numbers:

x	$-\omega$	0	$1/\omega$	1	$1 + 1/\omega$	$\omega/2$	$\omega - 1$	ω^2
$T(x)$	1	1	1	1	1	0	0	0
$B(x)$	0	0	0	0	0	1	1	1

For example, the $n = 2$ option is positive at $x = 1 + 1/\omega$, while every finite- n option is negative at $x = \omega/2$. No approximation of a sign is being made.

4 Why the construction is genetic

4.1 The admissibility certificate

We isolate the exact constructor being used. Begin with the rational function q , take a fixed set of affine translates of it, and use their values as left options in a Conway cut. There is no reference to the function currently being defined, nor to a left or right option of the input. Such a formula is a *nonrecursive instance* of a genetic constructor, not an additional operation being postulated here.

In the original construction the admissible option expressions include $c_1 h(c_2 x + c_3) + c_4$, where h is already genetic and the c_i may be surreal constants. The chosen option families are set-sized. Arithmetic closure is also available [1, pp. 9–10]. For our n th option use

$$h = q, \quad c_1 = 1, \quad c_2 = -1, \quad c_3 = n, \quad c_4 = 0.$$

Thus $q(n - x)$ is an allowed expression before T is introduced. Its membership does not depend on knowing whether x is infinite.

The original paper explicitly accepts q and the one-option cut $\{q(x) \mid\}$ as genetic [1, p. 11]. This matters because it establishes that an option expression depending on x but on no options of x is admissible. We are using the same type of expression with a countable option set instead of a singleton.

Proposition 4.1. *T , B , id , and $\text{id} - B$ are genetic under the constructor just specified. The option formula for T is independent of the representation chosen for x .*

Proof. The set of option expressions

$$\mathcal{L} = \{x \mapsto q(n - x) : n \in \mathbb{N}\}, \quad \mathcal{R} = \emptyset$$

is countable and consists of allowed previously constructed expressions. For any x , all options have defined surreal values. No left–right conflict is possible. The genetic adjunction therefore defines T . Polynomial and arithmetic closure gives $B = 1 - T$, id , and $\text{id} - B$.

Changing a presentation $x = \{L_x \mid R_x\}$ changes none of the numerical expressions $q(n - x)$. Hence it changes neither option set nor cut value. This is uniformity with respect to the presentation of x . $\square$

4.2 The stronger cofinality check

Berenbeim’s Definition 55 uses a positive-denominator localization and an additional global cofinality condition [2, pp. 126–128]. For a left option expression $\ell(v, w; x)$, the relevant requirement is that $\ell(y, z; x) < T(x)$ whenever $y < x < z$; a dual inequality applies to right options. Here

$$\ell_n(v, w; x) = q(n - x)$$

ignores both v and w . If $x \in \mathcal{D}$, then $\ell_n(y, z; x) \leq \frac{1}{2} < 1 = T(x)$. If $x \in \mathcal{U}$, then $\ell_n(y, z; x) < 0 = T(x)$. The right-option condition is vacuous.

The denominator $1 + (n - x)^2$ is strictly positive on the entire domain, so it is also compatible with the localization. Thus the certificate checks the extra condition directly, rather than deducing it from an informal assertion that the function “looks uniform.” The thesis also explicitly treats the one-option rational step construction as entire and uniform [2, pp. 133–134].

Remark 4.2 (What “entire” means here). In this setting “entire” concerns a genetic function defined at every surreal argument, without a pole or an excluded input. It is not the complex-analytic notion of a globally holomorphic function. Formula (3.2) is entire in the former sense.

4.3 A notational issue that must not be hidden

Genetic formulas often display unions over x^L and x^R even when a particular option term uses neither variable. Such a term remains present when one or both input option sets are empty. This convention is necessary already for the accepted one-option step function at $x = 0$. Reading a schematic union as a literal Cartesian-product index that deletes every independent term when $R_x = \emptyset$ would not reproduce that example. Fornasiero makes the convention about unused option variables explicit [3, Example 2.5].

Our proof therefore uses the usual option-expression semantics exhibited by the sources themselves. It does not exploit undefined terms, hidden conditional inclusion, class-sized options, or a choice of an unusual presentation of x . A materially narrower notion, for example one allowing only finitely many option expressions, would be a different question; this note does not settle it.

5 Nonunique primitives, even with the same real values

Lemma 5.1 (Invariance under finite translation). *If $c \in \mathcal{O}$, then*

$$x \in \mathcal{U} \iff x + c \in \mathcal{U}.$$

Consequently $B(x + c) = B(x)$ and $T(x + c) = T(x)$.

Proof. Choose $k \in \mathbb{N}$ with $|c| \leq k$. If $x \in \mathcal{U}$, then for every $n \in \mathbb{N}$, $x > n + k$, and therefore $x + c \geq x - k > n$. Thus $x + c \in \mathcal{U}$. The reverse implication follows by applying the same argument to $-c$. $\square$

Theorem 5.2 (The kernel is not just the constants). *B and T are locally constant at every surreal argument. More precisely,*

$$|h| < 1 \implies B(x + h) = B(x) \text{ for every } x \in \mathbf{No}. \quad (5.1)$$

Hence $B'(x) = T'(x) = 0$ for all x , although neither function is constant. All their higher pointwise derivatives exist and vanish.

Proof. Every h with $|h| < 1$ is finite, so lemma 5.1 proves (5.1). For $0 < |h| < 1$ the difference quotient for B is exactly 0. Given any surreal $\varepsilon > 0$, choosing $\delta = 1$ makes its distance from 0 less than ε . The same applies to T .

But $B(0) = 0$ and $B(\omega) = 1$, so B is not constant, and neither is $T = 1 - B$. The first derivative is the zero function; repeated differentiation therefore gives zero at every higher order. $\square$

The radius 1 in (5.1) is independent of x and of the requested accuracy. The apparent paradox is that no finite chain of steps of size less than 1 can carry a finite number to a positive infinite number. A real proof that connects two points by finitely many such steps uses the Archimedean property, which is unavailable here.

Theorem 5.3 (Counterexample to genetic primitive uniqueness). *Define, on all of $\mathbf{No}$,*

$$F_0(x) = x, \quad F_1(x) = x - B(x) = x - 1 + T(x). \quad (5.2)$$

Then F_0 and F_1 are genetic, $F'_0 = F'_1 = 1$ everywhere, and $F_0(r) = F_1(r) = r$ for every real r . Nevertheless $F_1 - F_0$ is not constant. The same failure occurs on the proper bounded open interval $(0, \omega)$.

Proof. Geneticity is proposition 4.1. For $0 < |h| < 1$,

$$\frac{F_1(x + h) - F_1(x)}{h} = \frac{h - B(x + h) + B(x)}{h} = 1.$$

Thus its derivative is 1, and the identity function has the same derivative. Every real number belongs to $\mathcal{D}$, so $B(r) = 0$ and the real values agree. The difference is $-B$, which takes both 0 and -1 .

On $(0, \omega)$ use the two points 1 and $\omega/2$. Both belong to the interval, and the differences there are 0 and -1 , respectively. For differentiation within this interval, shrink the radius at x to $\min\{1, x/2, (\omega - x)/2\} > 0$. This leaves the same quotient calculation valid on a neighborhood contained in the domain. $\square$

The piecewise description of F_1 is precisely

$$F_1(x) = \begin{cases} x, & x \in \mathcal{D}, \\ x - 1, & x \in \mathcal{U}. \end{cases}$$

This is the function already displayed in equation (7) immediately before the conjecture in [1, p. 37]. The present answer does not claim to have discovered this piecewise primitive. It supplies the genetic representation that makes it a counterexample to the restriction proposed there.

Corollary 5.4 (Failure of normalized global uniqueness). *The global initial-value problem $F' = 1$, $F(0) = 0$ has at least two genetic solutions. Even prescribing $F(r) = r$ simultaneously for every $r \in \mathbb{R}$ does not select one of them. Likewise $G' = 0$, $G(0) = 0$ has the distinct genetic solutions $G = 0$ and $G = B$.*

Proof. Apply theorems 5.2 and 5.3. No assertion of failure of local uniqueness on a sufficiently small neighborhood of 0 is intended: the displayed solutions agree on every finite argument. $\square$

6 Failure of the sup property

For a single-variable function f , the sup property requires that for $a < b$ and c in the field, the infima and suprema of both classes

$$\{x : a < x < b, f(x) \leq c\}, \quad \{x : a < x < b, f(x) \geq c\}$$

are field elements or the two end gaps. This is Fornasiero's Definition 3.2 [3], used in the later conjecture. It is enough to exhibit one nonempty bounded sublevel class with no surreal supremum.

Theorem 6.1 (Counterexample to the entire-genetic sup conjecture). *The entire genetic function B fails the sup property. In particular,*

$$E = \{x \in \mathbf{No} : 0 < x < \omega, B(x) \leq \tfrac{1}{2}\} \quad (6.1)$$

has no supremum in $\mathbf{No} \cup \{\mathbf{Off}, \mathbf{On}\}$.

Proof. By theorem 3.4, E is exactly the class of positive finite surreals. The condition $x < \omega$ is automatic for every such number. The class contains every positive integer and is bounded above by ω .

If u is any surreal upper bound of E , then $u \geq n + 1 > n$ for every $n \in \mathbb{N}$, since $n + 1 \in E$. Hence $u \in \mathcal{U}$. Conversely, every $u \in \mathcal{U}$ is an upper bound of E : for $e \in E$, choose $n \in \mathbb{N}$ with $e \leq n < u$. Thus the class of surreal upper bounds of E is exactly $\mathcal{U}$.

This class has no least member, because $u/2 \in \mathcal{U}$ and $u/2 < u$ for every $u \in \mathcal{U}$. There is therefore no surreal least upper bound. The bottom end gap cannot be an upper bound of a nonempty positive class, and the top end gap cannot be least because ω is already an upper bound. This proves the claim. $\square$

For completeness, the complementary superlevel class in the same interval,

$$E^+ = \{x : 0 < x < \omega, B(x) \geq \tfrac{1}{2}\} = \mathcal{U} \cap (0, \omega),$$

has no surreal infimum. Every $d \in \mathcal{D}$ is a lower bound. If a lower bound u belonged to $\mathcal{U}$, then either $u < \omega$, in which case $u/2 \in E^+$ lies below it, or $u \geq \omega$, in which case $\omega/2 \in E^+$ lies below it. Thus all lower bounds lie in $\mathcal{D}$, and $\mathcal{D}$ has no greatest member.

Notice the precise obstruction: boundedness supplies many upper bounds, not a *least* upper bound. Being defined everywhere and having no rational poles does not remove an unrealized Dedekind boundary of a sublevel class.

7 Every set-generated endpoint-free cut gives a variant

The use of $\mathbb{N}$ is not essential for the evaluation mechanism. What matters for differentiability is whether the lower indexing set has a last element.

Definition 7.1. For a nonempty set $A \subset \mathbf{No}$, define

$$T_A(x) = \{q(a - x) : a \in A\}, \quad B_A(x) = 1 - T_A(x), \quad (7.1)$$

$$\mathcal{U}_A = \{x : (\forall a \in A) a < x\}, \quad \mathcal{D}_A = \{x : (\exists a \in A) x \leq a\}. \quad (7.2)$$

The set A is fixed before the function is defined.

Theorem 7.2 (General gap-indicator construction). *For every nonempty set $A \subset \mathbf{No}$, T_A and B_A are entire uniform genetic functions under the same option rules, with $T_A = \mathbf{1}_{\mathcal{D}_A}$ and $B_A = \mathbf{1}_{\mathcal{U}_A}$. Both values occur. If A has no largest element, there is one surreal $\delta_A > 0$ such that*

$$|h| < \delta_A \implies B_A(x + h) = B_A(x), \quad T_A(x + h) = T_A(x) \quad (x \in \mathbf{No}). \quad (7.3)$$

In particular, both functions are locally constant everywhere and have zero derivative everywhere. If A has a largest element m , they are discontinuous at m and are not differentiable there.

Proof. The option expressions are a set of affine translates of q with surreal constants. They ignore all options of x , and their denominators are positive. The construction and cofinality checks of Section 4 apply with a in place of n . The sign argument of theorem 3.4 proves the stated values. Any $a \in A$ lies in $\mathcal{D}_A$, while $\{A \mid \}$ lies in $\mathcal{U}_A$, so the function is nonconstant.

Suppose first that A has no largest element. Form the set of all its positive differences and one separating cut:

$$\Delta_A = \{b - a : a, b \in A, a < b\}, \quad \delta_A = \{0 \mid \Delta_A\}.$$

Replacement makes Δ_A a set, every member is positive, and the cut therefore gives $0 < \delta_A < d$ for every $d \in \Delta_A$. The radius is chosen once from A , independently of the point x .

If $x \in \mathcal{D}_A$, choose $a \in A$ with $x \leq a$ and then $b \in A$ with $b > a$. For $|h| < \delta_A$,

$$x + h < x + \delta_A \leq a + \delta_A < b,$$

so $x + h \in \mathcal{D}_A$. If $x \in \mathcal{U}_A$, fix any $a \in A$ and choose $b \in A$ with $b > a$. Since $x > b$ and $\delta_A < b - a$, the same radius gives

$$x + h > x - \delta_A > b - \delta_A > a.$$

This holds for every $a \in A$, so $x + h \in \mathcal{U}_A$. No simultaneous selection of the witnesses b is needed: each inequality is proved for an arbitrary fixed a . The two indicator formulas now give (7.3), and every punctured difference quotient is exactly zero on this radius.

Suppose instead that $m = \max A$. Then $B_A(x) = 0$ for $x \leq m$ and $B_A(x) = 1$ for $x > m$. In every neighborhood of m there are points of both types, so continuity fails with tolerance $\frac{1}{2}$. For a direct derivative argument, if a derivative L at m existed, choose $h > 0$ smaller than both its proposed tolerance-1 radius and $1/(2(|L| + 1))$. The quotient equals $1/h > 2(|L| + 1)$ and hence cannot be within 1 of L . Thus no derivative exists at m . $\square$

A finite nonempty A has a maximum, so it cannot produce an everywhere zero-derivative nonconstant function by this particular formula. A countable set without a maximum already can. This is a minimality statement about the *displayed schema*, not about all possible finite genetic definitions.

Proposition 7.3 (Cofinality is the invariant). *Suppose $A, C \subset \mathbf{No}$ are nonempty sets and each is weakly cofinal in the other: every member of either set is at most some member of the other. Then $T_A = T_C$ and $B_A = B_C$.*

Proof. A number is strictly greater than all of A if and only if it is strictly greater than all of C . Indeed, an $a \in A$ can be bounded by $c \in C$, and the same argument works in reverse. Thus $\mathcal{U}_A = \mathcal{U}_C$, and the explicit indicator formulas apply. $\square$

The indicator depends only on the cut encoded by A , not on an enumeration or on the magnitudes of the rational options. The displayed choice of δ_A is sufficient, but not intrinsic to the indicator or necessarily largest: cofinal sets can give the same indicator with different such radii. No real supremum of A is used. The uniform radius comes from a cut with a *set* of right options; this argument does not supply one radius simultaneously for a proper class of indexing sets.

8 Order automorphisms and an ordinal-indexed kernel

8.1 A group of normalized primitives

For $c \in \mathbf{No}$, let

$$\Phi_c(x) = x + cB(x). \quad (8.1)$$

Every Φ_c is genetic and has derivative 1, whatever the size of c . Finite c give a stronger conclusion.

Theorem 8.1. Φ_c is an increasing bijection $\mathbf{No} \rightarrow \mathbf{No}$ exactly when c is finite. For finite c, d ,

$$\Phi_c \circ \Phi_d = \Phi_{c+d}, \quad \Phi_c^{-1} = \Phi_{-c}. \quad (8.2)$$

Proof. Let c be finite. Translation by c preserves both $\mathcal{U}$ and $\mathcal{D}$ by lemma 5.1. On $\mathcal{D}$, Φ_c is the identity; on $\mathcal{U}$, it is translation by c , a bijection of that class. It is strictly increasing on each class. If $d \in \mathcal{D}$ and $u \in \mathcal{U}$, then $u - d$ is positive infinite by (2.4), so $(u - d) + c > 0$. Thus $\Phi_c(u) = u + c > d = \Phi_c(d)$, establishing strict increase across the gap.

Since $B(\Phi_d(x)) = B(x)$ for finite d , substitution gives

$$\Phi_c(\Phi_d(x)) = x + dB(x) + cB(\Phi_d(x)) = x + (c + d)B(x).$$

This proves (8.2), including the inverse formula.

Conversely, if c is positive infinite, then c is not in the range of Φ_c : points in $\mathcal{D}$ remain in $\mathcal{D}$, while for $x \in \mathcal{U}$, $x + c > c$. Thus the map is not onto. If c is negative infinite, then $-c \in \mathcal{U}$ and $\Phi_c(-c) = 0 = \Phi_c(0)$, so the map is not injective. These exhaust the nonfinite cases. $\square$

Taking $c = -1$ recovers F_1 . Thus the central counterexample is not merely an arbitrary discontinuous rearrangement of values: it is an order automorphism with exactly the same derivative as the identity.

8.2 Many independent zero-derivative functions

For each ordinal α , put

$$s_\alpha = \omega^\alpha, \quad A_\alpha = \{ns_\alpha : n \in \mathbb{N}\}, \quad B_\alpha = B_{A_\alpha}.$$

Here ns_α is field multiplication. Because $s_\alpha > 0$, A_α has no maximum, so every B_α is a nonconstant entire genetic function with zero pointwise derivative.

Theorem 8.2 (Ordinal-indexed finite linear independence). *If $\alpha_1 < \dots < \alpha_k$ are ordinals and $c_1, \dots, c_k \in \mathbf{No}$, then*

$$\sum_{i=1}^k c_i B_{\alpha_i} = 0 \implies c_1 = \dots = c_k = 0.$$

Consequently there are independent families of these kernel elements indexed by sets of arbitrarily large ordinal order type.

Proof. Evaluate at $x_j = \omega^{\alpha_j+1}$. If $i \leq j$, then $x_j > n\omega^{\alpha_i}$ for every natural n , so $B_{\alpha_i}(x_j) = 1$. For $i = j$ this follows from $\omega^{\alpha_j+1} = \omega^{\alpha_j}\omega$; for earlier i the scale is no larger. If $i > j$, ordinal order gives $\alpha_i \geq \alpha_j + 1$, hence $x_j \leq \omega^{\alpha_i} \in A_{\alpha_i}$, so $B_{\alpha_i}(x_j) = 0$.

Thus the evaluation matrix has entries

$$B_{\alpha_i}(x_j) = \begin{cases} 1, & i \leq j, \\ 0, & i > j. \end{cases}$$

The first evaluation gives $c_1 = 0$, the second gives $c_1 + c_2 = 0$, and successive subtraction gives every coefficient zero. $\square$

This is a statement about finite linear relations, so no summability convention for a proper-class family is needed. Each finite linear combination remains genetic and locally constant: at a fixed point, intersect finitely many neighborhoods on which its summands are constant. Adding such a combination to x gives further primitives of 1. We do not assign a set-valued dimension to a proper-class function space.

9 Why familiar uniqueness arguments do not apply

9.1 The missing point in a mean-value argument

Over the real numbers, zero derivative on an interval implies constancy through the mean value theorem. That theorem is not an algebraic consequence of the ordered-field axioms. Our example makes the failing implication explicit: B has derivative 0 everywhere but $B(\omega) - B(0) = 1$.

There is no discontinuity at a hidden surreal “infinity point.” No such point exists. Every actual surreal has a neighborhood entirely on one side of the gap. In particular, testing differentiability at ω , $\omega/2$, $\omega - 1$, or an arbitrary positive infinite number cannot locate a jump. The jump belongs to a Dedekind boundary not realized by a field element.

The failure is visible on the bounded interval $[0, \omega]$. Boundedness in a non-Archimedean field is not compactness or Dedekind completeness. A supremum proof of Rolle’s theorem cannot simply be imported from real analysis: the class whose endpoint one needs may have a gap as its supremum, as in theorem 6.1.

9.2 Local openness versus the set-union topology

Two openness conventions occur here and must be named separately. The local order-field convention declares a class open when each point has an interval contained in it; it is the convention relevant to the ε - δ derivative in (2.5). The original article instead uses a global openness convention built from *set-sized* unions of intervals whose endpoints are surreal numbers or the two end gaps [1, Definition 7]. This is weaker than allowing every locally open proper class to count as open. The distinction can be seen directly here. It is not a distinction between two choices of small numerical tolerance: it is a restriction on the allowed global unions.

Both $\mathcal{D}$ and $\mathcal{U}$ are locally open. Also

$$\mathcal{D} = \bigcup_{n \in \mathbb{N}} (-\infty_{\text{end}}, n)$$

is a set-sized union of allowed intervals. But $\mathcal{U}$ is not such a union. To prove this, suppose $\mathcal{U} = \bigcup_{i \in I} (a_i, b_i)$ with I a set. Every nonempty interval in the union must have its lower endpoint $a_i \in \mathcal{U}$: an endpoint in $\mathcal{D}$ would leave some points of $\mathcal{D}$ in the interval, because $\mathcal{D}$ has no greatest element. A bottom-end endpoint would have the same problem. Now

$$y = \{ \mathbb{N} \mid a_i : i \in I \}$$

is positive infinite but below every a_i , so belongs to none of the intervals, a contradiction. Hence $B^{-1}((\frac{1}{2}, \frac{3}{2})) = \mathcal{U}$ is not open in this set-union sense, despite pointwise local continuity.

However, F_1 is an order automorphism by theorem 8.1, so it is a homeomorphism for this topology: preimages of basic intervals are intervals with the corresponding inverse-image endpoints, and preimages preserve set-sized unions. The same is true of its inverse. Thus demanding that the primitive itself be continuous in this global sense does not remove the example. The original paper already describes the piecewise F_1 as strongly continuous; this note uses the directly proved order property rather than relying on that terminology.

There is no valid shortcut from continuity of F_1 and id to global continuity of their difference in this particular framework. Indeed, that difference is $-B$, for which the preceding inverse-image obstruction applies. The familiar closure of continuous real-valued functions under arithmetic must not be assumed without checking compatibility with the chosen proper-class topology.

9.3 Why Fornasiero's preservation theorem is not contradicted

Fornasiero proves preservation of the sup property under specified uniform recursive constructions with an additional *tameness* hypothesis [3, Theorem 3]. In his Definition 3.1, a nonconstant tame function must, on each side of a gap and sufficiently near it, stay strictly above or strictly below any specified comparison value.

Our B fails that condition. At the finite–infinite gap, choose comparison value 0. On the entire left side $B = 0$; no left neighborhood has $B > 0$ everywhere or $B < 0$ everywhere. The global constant-function exception does not apply because B is nonconstant. Thus a published theorem about tame functions cannot be used to infer the sup property for B . Uniformity alone and tameness are different restrictions.

9.4 A correct narrow uniqueness statement

Proposition 9.1 (Rational primitives remain unique). *Let K be an ordered field of characteristic zero. If two rational functions without poles on a nonempty open interval have equal pointwise derivatives there, then their difference is constant on that interval. The same argument applies to rational surreal functions.*

Proof. Write their difference as P/Q in lowest terms over K , with Q nonzero on the interval. Differentiating algebraically gives $(P'Q - PQ')/Q^2$. Since it vanishes at every point of the interval and the interval contains infinitely many elements, the numerator polynomial is identically zero.

If $P = 0$ we are done. Otherwise, $P'Q = PQ'$ and $\gcd(P, Q) = 1$ imply $P \mid P'$. In characteristic zero a nonconstant P has $\deg P' < \deg P$, so this is impossible unless P is constant. Then the identity forces $Q' = 0$, making Q constant as well. All operations involve only finitely many surreal coefficients, so the same polynomial argument is legitimate over the proper-class field. $\square$

Thus the countable *cut* is doing real work. It converts signs of rational functions into an infinitary Boolean condition; the resulting function is not itself a rational function. A repair based on a smaller

function class must control this operation rather than only the regularity of individual rational option functions.

10 An ordered-field analogue and the hyperreal distinction

Proposition 10.1. *Let K be any non-Archimedean ordered field. Define*

$$b_K(x) = \begin{cases} 1, & x > n \text{ for all ordinary } n \in \mathbb{N}, \\ 0, & \text{otherwise.} \end{cases}$$

As an unrestricted function $K \rightarrow K$, it is nonconstant, invariant under finite translations, and has order-field derivative zero everywhere.

Proof. Non-Archimedeaness supplies a positive element greater than every natural number, whereas $b_K(0) = 0$. The integer-bound proof of lemma 5.1 uses only ordered-field arithmetic and applies in K . In particular, $b_K(x + h) = b_K(x)$ for $|h| < 1$, and every relevant punctured difference quotient is exactly zero. $\square$

The exceptional feature in the surreal problem is therefore not the mere existence of such a function. It is the ability to certify it by the published *genetic syntax*. The Conway cut provides that certificate.

In a hyperreal setting with the usual internal/external distinction, b_K must be treated as an external function, not as the transferred extension of an ordinary real function. Here is a direct diagnostic, assuming the standard transfer framework. Were b_K internal, the class of hyperintegers H with $b_K(H) = 1$ would be an internal nonempty subset of the hypernaturals. The transferred least-element principle would give it a least member H . But H is greater than every ordinary natural, and so is $H - 1$, contradicting minimality. This argument invokes the least-element principle only for an internal set, which is precisely where the contradiction enters.

The ordered-field derivative calculation is valid independently of such extra structure. It neither refutes transferred theorems for internal objects nor identifies genetic functions with internal hyperreal functions. These are different restrictions on which functions are admitted.

11 Finite truncations and exact computational checks

11.1 What finite option sets do

Let

$$T_N(x) = \{ q(n - x) : 0 \leq n \leq N \mid \}, \quad B_N(x) = 1 - T_N(x).$$

The maximum of the option-indexing set is N , so the binary cut lemma gives

$$T_N(x) = \begin{cases} 1, & x \leq N, \\ 0, & x > N. \end{cases}$$

Each B_N is an ordinary step at the actual surreal point N , where it is not differentiable. In contrast, the countably indexed B has no exceptional point.

For every fixed surreal x , the sequence $T_N(x)$ is eventually exactly $T(x)$. If $x \in \mathcal{D}$, choose a natural upper bound n for x ; all $N \geq n$ give $T_N(x) = 1$. If $x \in \mathcal{U}$, every N gives $T_N(x) = 0$. This eventual stabilization is stronger than a numerical convergence claim at the chosen input, but it is *not uniform* in x : at $x = N + 1$, $T_N(x) = 0$ while $T(x) = 1$.

In particular, one must not replace a countable option family by a large finite list and then assert that the resulting function has zero derivative everywhere. The exact construction is infinitary, even though its sign argument is short.

11.2 The supplied verifier

The file `code/verify.py` uses Python's standard library and exact rational arithmetic. It implements the ordered rational-function field $\mathbb{Q}(t)$ with t positive infinite. For a nonzero rational function $p(t)/r(t)$ with positively normalized denominator, its sign is determined by the leading coefficient of p , and it is positive infinite exactly when that sign is positive and $\deg p > \deg r$. This is a symbolic classification, not a test against finitely many large integers.

The program separately enumerates a finite early-day dyadic surreal tree to find the simplest number above selected finite sets of rational left options. This checks the finite-cut implementation without hard-coding the output of each example. It does not claim to implement arbitrary surreal birthdays or countable cuts.

The archived run completed **7,248 exact checks**, all passing, with a fixed random seed 20260920 and 84 rational-function inputs. These are assertion counts, not counts of independently proved theorems. The groups are summarized below; the machine-readable details are in `results/verification.json`.

Check group	Assertions
Rational-field identities	335
Sign and rational bounds	336
Polynomial square certificates	168
Explicit natural upper-bound witnesses	134
Selected infinite-option signs	85
Inverse maps	168
Punctured-neighborhood bounds	6
Local constancy	504
Primitive difference quotients	1008
Indicator difference quotients	504
Preservation of order	3486
Finite-translation group law	180
Finite Conway-cut searches	270
Triangular scale-evaluation matrix	64
Total	7248

These computations can expose arithmetic, sign, boundary-case, and implementation mistakes. They do not verify membership in a published genetic class, settle a quantifier over all surreal numbers, show that a set-sized cut exists, or establish novelty. Those parts of the argument are explicitly mathematical or bibliographic, and remain separate from the test report.

11.3 Reproduction

From the archive directory, run

```
python code/verify.py
latexmk -pdf -interaction=nonstopmode -halt-on-error article.tex
latexmk -pdf -interaction=nonstopmode -halt-on-error short_proof.tex
```

No network access or third-party Python packages are required for the verification. The TeX documents use standard TeX Live packages; a build script for PowerShell and a POSIX shell are supplied. External papers are referenced, not redistributed.

12 Conclusions and remaining questions

The central algebraic observation is that the cut

$$\{ q(a - x) : a \in A \mid \}$$

turns a set-indexed family of sign tests into an existential assertion: it equals 1 exactly when $x \leq a$ for at least one $a \in A$. For an indexing set with no largest element, this assertion changes truth value across an unrealized gap, not at any surreal point. A function can therefore be locally constant at every input while being globally nonconstant, and the broad genetic constructors admit it.

Under the definitions checked in this note, the outcomes are exact. Genetic primitives need not be unique up to a constant; even real normalization and an increasing-bijection requirement do not restore uniqueness. Entire genetic functions need not have the sup property. The pointwise differentiation kernel contains the ordinal-indexed independent family of [theorem 8.2](#). These are consequences of the explicit cut, not extrapolations from computation.

Several narrower research questions remain worth separating from the resolved targets. What restrictions on option families preserve the sup property? Which recursive subclasses exclude gap indicators while still containing the desired transcendental functions? Can one characterize an analytically useful class in which pointwise differentiation has only constant kernel? Restricting to finitely many rational option expressions removes the present schema, but no assertion that this restriction alone settles those broader questions is made here.

The appropriate next external check is focused: verify the admissibility certificate against the intended genetic-function conventions, and search for prior publication of this countable-cut observation. The core proof is short enough for such a check to be independent of the implementation and of every later generalization in this article.

A Proof dependency and objection audit

Potential objection	Explicit resolution
The left options may form a proper class.	They are indexed by $\mathbb{N}$, or by a specified set A . For each input, replacement gives a set of values.
The cut may fail to define a number.	Its right option set is empty. Moreover every left option lies strictly below 1, so even an explicit separating candidate is available.
The formula may hide a conditional option rule.	Every option $q(n - x)$ is always included and always defined. The finite/infinite split is derived from its sign, not used in the genetic definition.

Potential objection	Explicit resolution
The denominator may vanish at a nonreal input.	$1 + t^2 > 0$ in every ordered field, including No .
Only a nongenetic piecewise function has been constructed.	The nonrecursive option terms fit the published affine-template constructor, and the fixed option family is countable.
The construction may fail uniformity.	The expressions ignore the option variables of the argument. All representation choices leave exactly the same numerical cut.
The later cofinality condition may exclude it.	For every $y < x < z$, each left option is still $q(n - x) < T(x)$; right inequalities are vacuous.
A jump may occur at ω .	The finite–infinite gap is not ω . Every number in $(\omega - 1, \omega + 1)$ is positive infinite; B is constant there.
The derivative may use only real increments.	The quotient identity holds for every surreal h with $0 < h < 1$, including arbitrary infinitesimals.
A bounded interval might restore uniqueness.	$(0, \omega)$ already contains the witnesses 1 and $\omega/2$.
The primitive might lack monotonicity.	$F_1 = \Phi_{-1}$ is an increasing bijection, with inverse Φ_1 .
The sup property might allow the end gaps.	The witness class is nonempty and bounded by the surreal ω . Neither end gap can be its least upper bound.
A known theorem about tame functions may contradict the result.	B is not tame in Fornasiero’s strict gap-comparison sense.
The finite computations might be presented as a formal proof.	Their scope is limited to exact finite shadows; the proper-class and countable-cut claims are proved in the text, not verified by the program.
The primitive itself may already be known.	It is: the source prints its piecewise form. Only the genetic realization and the deductions are presented as the contribution here.

B Source locations and status of the search

The primary-text locations checked for this note are as follows. Page numbers are printed page numbers, not PDF-viewer indices.

Source	Locations used
[1]	Section 2.1 (cut and simplicity conventions); pp. 9–11 (genetic constructor, closure, rational unit step); pp. 27–28 (function limits and derivatives); p. 37, equation (7) and Conjecture 53.
[2]	Definition 55, pp. 126–128 (adjunction and global cofinality); pp. 133–134 (entire functions and rational step); Chapter 8, pp. 225–226, Conjecture 3.
[3]	Example 2.5 (unused option variables); Definitions 3.1–3.2 and Theorem 3 (tameness and sup property).

Searches on 20 September 2026 included exact-conjecture queries and combinations of “genetic,” “surreal,” “primitive,” “counterexample,” and “sup property,” together with the authors’ names. Relevant results located the original article, its arXiv version, the dissertation, and Fornasiero’s preprint. Some broader queries returned mostly irrelevant material, so absence from these results

is weak evidence about priority. No located result supplied a later resolution of the two targets. The source manifest records this limitation rather than labelling an unsuccessful search as an exhaustive literature review.

The theorem statements and proofs in this note were developed from the explicit formulas displayed here. Their evaluation does not depend on a claim that the questions remained universally recognized as open on the access date. If earlier work contains the same observation, the mathematical counterexamples stand but the novelty claim must be revised.

References

- [1] Simon Rubinstein-Salzedo and Ashvin Swaminathan, *Analysis on Surreal Numbers*, Journal of Logic & Analysis **6:5** (2014), 1–39. doi:10.4115/jla.2014.6.5. Published PDF; arXiv:1307.7392.
- [2] Alexander Michael Berenbeim, *Foundations for the Analysis of Surreal-Valued Genetic Functions*, Ph.D. dissertation, University of Illinois at Chicago, 2022. UIC repository PDF, version 1, file 36188862.
- [3] Antongiulio Fornasiero, *Recursive definitions on surreal numbers*, arXiv:math/0612234, version 1, 9 December 2006. <https://arxiv.org/abs/math/0612234>.