

Term counts in the derivatives of power towers

One framework and three sequences:

OEIS A293239 (x^x), A290268 (x^{x^2}), A281434 (x^{x^x})

Research notes prepared for Vladimir Reshetnikov

September 2026

Abstract

For a function f , let $a(n)$ be the number of distinct monomials in the fully collected n th derivative $f^{(n)}$. This article treats three instances of that question at once: $f = x^x$ (OEIS A293239), $f = x^{x^2}$ (A290268) and $f = x^{x^x}$ (A281434).

Part 1 proves, once, the apparatus the three share: a canonical differential-polynomial form $f^{(n)} = f \cdot P_n$ or $f \cdot x^{-n} P_n$ with a first-order recurrence for P_n over $\mathbb{Z}$, integrality of its coefficients, and the envelope bound that turns an upper estimate into a cancellation count. It also proves the linear independence of logarithmic monomials that makes “the number of terms” an invariant of f rather than of a differentiation algorithm, and the resulting identity $a(n) = |\text{supp } P_n|$ — for two of the three frames; x^{x^x} falls outside that statement and supplies its own, as Part 1 is careful to say. Two normalizations are stated, because neither one covers all three parts in the frames those parts use; Remark 1.3 is careful about what that does and does not mean.

The three remaining parts are the three original investigations, unabridged. They reach different answers by different routes. **None of the three headline questions is fully settled here, and this article does not obscure that.** Two of the three entries print a conjectured closed form and neither is proved; the third prints no closed form, and there the growth *order* is settled while the constant is not. For x^x the term count is reduced exactly to the zeros of the first-kind Lehmer–Comtet triangle A008296, giving $a(n) = 1 + \binom{n+1}{2} - Z(n)$ and $a(n) = \Theta(n^2)$, but the principal closed formula needs a nonvanishing statement that is isolated and left open; separately, the recurrence printed in the entry with the range $n > 6$ is *disproved* at $n = 8$. Later additions at the end of that part (Section 2.11) prove the nonvanishing statement on further infinite families of offsets of the triangle — every offset $\mu(p-1)$ for primes $p > \mu$ under explicit conditions, and every prime p for $\mu = 2, 4$, every odd one for $\mu = 3, 5$ — and give exact p -adic valuations on divisibility progressions; the zero classification itself stays open. [Added 1 October 2026, batch 72A: the sibling report **power-tower-exponent-supports** proves $a(n) \sim n^2/2$, that is $Z(n) = o(n^2)$, without an effective threshold.] For x^{x^2} the conjectured expression is proved to be an upper bound, every predicted cancellation is explained, and $a(n) = \Theta(n^2)$ is established, but the conjectured leading constant is not; a later addition to that part (Section 3.14) proves the missing nonvanishing statement at logarithmic deficits three and four, leaving deficits five and above open; a second later addition (Section 3.27) re-proves deficit three by another route; a third (Section 3.35) proves deficits five and six, so that deficits seven and above are now open, reduces every fixed deficit to a finite rectangle of singly exponential size, and derives an explicit lower bound $3n^2/8 + (n/2) \log n - (n/2) \log \log n - 5n/4$ for $n \geq e^{10}$. [Added 1 October 2026, batch 73: a fourth addition (Section 3.42) proves deficits seven and eight, deficit seven by two independent certificates, so that deficits nine and above are now open.] [Added 1 October 2026, batch 72A: the same sibling report proves $a(n) \sim n^2/2$ here too, so the leading constant is $\frac{1}{2}$; the proof is ineffective, and $3/8$ remains the leading constant of every explicit lower bound.] For x^{x^x} two explicit polynomial bounds give $a(n) = \Theta(n^3)$ unconditionally, while the sharper equivalent $a(n) \sim \frac{2}{3}n^3$ is not proved — and there, unlike the other two, the obstruction is not a nonvanishing statement at all but a density one,

because additional zeros provably exist. Part 5 sets the three side by side, says what the shared method does and does not buy, and records the one piece of structure they share beyond it: every cancellation any of them can prove is a parity zero of the same falling factorial.

All computations reported here are exact. Every modular certificate is confirmed by exact integer arithmetic before a coefficient is declared to vanish, so no claim rests on a probabilistic zero test; but every computation certifies a finite range and none of them is an all- n proof.

Contents

How this article is arranged	5
1 The common framework	6
1.1 Setting	6
1.2 The canonical form and its recurrence	7
1.3 Independence, and why the count is well posed	8
1.4 Envelopes, upper bounds and exact computation	9
1.5 Notation across the parts	10
2 x^x: OEIS A293239	12
2.1 The question and the status of the answer	12
2.2 A canonical polynomial for each derivative	13
2.3 The Lehmer–Comtet coefficient triangle	14
2.4 The exact zero-count reduction	16
2.5 Diagonal polynomials and exact nonvanishing	18
2.6 The first three columns for all row indices	20
2.7 Unconditional quadratic growth	21
2.8 The rational function, quasipolynomial, and recurrence	22
2.9 Reproducible verification	24
2.10 What remains to finish the requested proof	25
2.11 Lehmer–Comtet zeros on composite offsets: three later additions	26
2.12 Complete nonvanishing on prime-multiple offsets (manuscript 12)	27
2.13 Stirling residues and finite candidate lists (manuscript 09)	34
2.14 Uniform p -adic valuations (manuscript 17)	39
2.15 Lehmer–Comtet additions: provenance, files and formal status	43
3 x^{x^2}: OEIS A290268	46
3.1 The problem and the status of the result	47
3.2 A canonical polynomial expansion	48
3.3 Exact formulas for individual coefficients	49
3.4 A transformation proving positivity	50
3.5 The two forced families of zeros	51

3.6	Counting the forced zeros	52
3.7	Nonvanishing on the first two deficit diagonals	53
3.8	Generating functions and equivalent formulas for the upper bound	55
3.9	Exact finite verification and certificates	57
3.10	The precise missing step	58
3.11	Conclusion	59
3.12	Reproducing the computations	60
3.13	Files included in the archive	60
3.14	Depth certificates: deficits three and four	61
3.15	Depth coordinates: exact coefficient and polynomial models	64
3.16	Depth coordinates: bulk positivity and symmetry holes	67
3.17	Depth coordinates: the branch-cut integral and its folding	68
3.18	Depth coordinates: monotone comparisons	70
3.19	Depth coordinates: an explicit finite reduction at every depth	71
3.20	Depth coordinates: completing depths one through four	73
3.21	Depth coordinates: the integer certificates and independent checks	74
3.22	Depth coordinates: an all-depth bound on the number of zeros	76
3.23	Depth coordinates: consequences for the term count	77
3.24	Depth coordinates: further research questions	79
3.25	Depth coordinates: conclusion, verifier and source audit	81
3.26	Depth certificates: provenance, files and formal status	83
3.27	A second route to depth three	85
3.28	Second route to depth three: the coefficient model and a falling-factorial transform	89
3.29	Second route to depth three: the beta-sine transform and centred moments	90
3.30	Second route to depth three: the harmonic phase transition	92
3.31	Second route to depth three: the complete tail and the full classification	93
3.32	Second route to depth three: further exact consequences	96
3.33	Second route to depth three: certificate, formalization blueprint and research questions	97
3.34	Second route to depth three: provenance, files and formal status	102
3.35	Depths five and six, and count bounds	103
3.36	Depths five and six: the theorem and the known regions	104
3.37	Depths five and six: the depth-five moment certificates	107
3.38	Depths five and six: the depth-six completion	111
3.39	Depths five and six: exponential cutoffs at every depth	112
3.40	Pointwise and summatory count bounds (manuscript 68)	114
3.41	Depths five and six and count bounds: provenance, files and formal status	120
3.42	Depths seven and eight	122
3.43	Depths seven and eight: the model and the known regions	126

3.44	Depths seven and eight: the largest-root tangent	129
3.45	Depths seven and eight: single-exponential cutoffs, a weaker form of Theorem 3.81	131
3.46	Depths seven and eight: three moment anchors at every depth	133
3.47	Depths seven and eight: the theorems and manuscript 37's certificates	135
3.48	Depth seven: a second, independent certificate (manuscript 46)	137
3.49	Depths seven and eight: term counts	140
3.50	Depths seven and eight: research questions	143
3.51	Depths seven and eight: status, provenance, files and formal status	146
4	x^{x^x}: OEIS A281434	151
4.1	The problem and the scope of the results	151
4.2	A canonical differential-polynomial representation	152
4.3	The support envelope and the cubic upper bound	153
4.4	An independent exact formula for every coefficient	154
4.5	A rigorous cubic lower bound	157
4.6	Exact cancellation families	159
4.7	Algorithms and complexity	160
4.8	Computational validation and results	162
4.9	What is and is not established asymptotically	163
4.10	A compact exact implementation	164
4.11	Reproducing the supplied computations	164
5	The three answers side by side	166
5.1	What was proved	166
5.2	Why the orders differ	167
5.3	What the shared framework buys, and what it does not	167
5.4	What the three do share, beyond the framework	168
5.5	What would settle each one	168

How this article is arranged

Three separate reports were written on three OEIS sequences that ask the same question about three different functions. They are collected here as one article because they share a construction, and kept as four distinct parts because the questions they answer are different: three functions, three recurrences, three answers. They do share one thing beyond the construction, and Section 5.4 sets it out — all three reduce to integer zeros of derivatives and differences of the same falling factorial — but no argument in any part carries over to another. Nothing that was proved in any of the three has been dropped, weakened or strengthened in the merge.

- **Part 1** is new to this article. It states and proves the machinery that the three reports had each established separately, and is explicit about where that generality stops: the canonical form, the recurrence and the envelope bound cover all three, the independence statement covers two of the three. It settles none of the three counting problems and is not intended to: the content of the subject is in the nonvanishing questions, and those do not transfer between the three functions.
- **Parts 2, 3 and 4** are the three original articles. Each opens with a status box saying what it does and does not prove, and a note on how it instantiates Part 1. Each part retains the setup section it was written with, so it can still be read on its own; those sections are now instances of Part 1 rather than independent developments, and are flagged as such where they begin.
- **Part 5** compares the three.
- **Sections 2.11–2.15**, at the end of Part 2, were added on 1 October 2026 from three later manuscripts on the Lehmer–Comtet triangle. They prove nonvanishing on further infinite families of offsets and exact p -adic valuations, and leave Conjecture 2.9 open. They follow the part they extend, so that no number elsewhere in the article moved, and they rename their sources’ letters by Table 2.A.
- **Sections 3.14–3.26**, at the end of Part 3, were added on 29 September 2026 from a later manuscript on A290268, which closes that part’s missing step at logarithmic deficits three and four. They follow the part they extend, so that no number elsewhere in the article moved, and they use their own coordinates (Table 3).
- **Sections 3.27–3.34**, after those, were added on 30 September 2026 from a manuscript on A290268 that gives a second proof at logarithmic deficit three, through a harmonic sign law; they use the same depth coordinates, with the further renamings of Table 5.
- **Sections 3.35–3.41**, after those, were added on 1 October 2026 from two manuscripts on A290268: one proves logarithmic deficits five and six and gives singly exponential cutoffs at every deficit, the other derives explicit pointwise and summatory bounds on the term count. They use the same depth coordinates, with the renamings of Table 3.A.
- [Added 1 October 2026, batch 73: **Sections 3.42–3.51**, after those, were added from two further manuscripts on A290268. They classify logarithmic deficits seven (twice, by independent certificates) and eight, so that the open deficits are now nine and above, and give an all-deficit three-anchor reduction and an eight-deficit count bound. One of them was written before the batch-72A addition and re-proves parts of it; those parts are printed as second certificates or remarks. They use the same depth coordinates, with the renamings of Table 3.B.]

- [Added 1 October 2026, batch 72A: two sibling reports, at the paths `../power-tower-exponent-supports` and `../a290268-unbounded-deficits` relative to this article's directory, build on it without being part of it. The first proves $a(n) \sim n^2/2$ for x^x and for x^{x^2} , ineffectively, as the cases $a = 1, 2$ of a theorem on x^{x^a} ; the second studies the coefficients of x^{x^2} when the logarithmic deficit grows with n . Dated notes of this date point to them where they bear on the text.]

The parts keep the notation of their sources, which is not uniform across them. Section 1.5 sets out what a reader crossing between parts needs. Two things are worth carrying in advance: **j is the exponent of $\log x$ in Parts 2 and 3 but the exponent of x^{-1} in Part 4**, where ℓ is the logarithm exponent; and **z is x^2 in Part 3, $\log x$ in Part 4, and a formal generating-function variable in Part 2**. Also $a(n)$ means a different sequence in each of the three parts.

Each part ends with a description of the code and data that accompanied it. Those descriptions have been updated to the merged package, in which every script lives under `code/` and every generated table under `data/`, and the article is `article.tex` and `article.pdf`. The three groups of scripts do not collide and each was run from the merged layout before this article was built.

These are AI-assisted research notes. None of the three has been refereed or machine-checked.

1 The common framework

Each of the three parts that follow studies a different function, obtains a different recurrence and arrives at a different answer. What they share is the apparatus in this part: a canonical polynomial form for the n th derivative, a uniqueness statement that makes “the number of terms” well defined, a first-order recurrence on that form, and the reduction of the counting problem to a nonvanishing problem. Each of the three source reports established this apparatus for its own function. Here it is proved once.

Nothing in this part decides any of the three counting problems. It is deliberately cheap: every statement below is an exercise, and the work in this subject is entirely in the nonvanishing questions that Parts 2, 3 and 4 take up.

1.1 Setting

Throughout, functions are real-valued and smooth on $(0, \infty)$, and f is positive there, so $\log f$ is defined and f'/f makes sense.

Definition 1.1 (Differential frame). A *differential frame* is a tuple $w = (w_1, \dots, w_m)$ of smooth functions on $(0, \infty)$ together with the polynomial ring $R = \mathbb{Z}[w_1, \dots, w_m]$, where the w_i are treated as independent indeterminates. We write $\text{supp } P$ for the set of exponent vectors of the monomials occurring in $P \in R$ with a nonzero coefficient, and $|\text{supp } P|$ for its cardinality.

The distinction between R as an abstract polynomial ring and the functions the w_i denote is the point of Section 1.3: a priori a polynomial identity in R is a stronger statement than the corresponding identity of functions, and it is the converse that has to be proved.

1.2 The canonical form and its recurrence

Theorem 1.2 (Canonical differential-polynomial form). *Let w be a differential frame with ring $R = \mathbb{Z}[w_1, \dots, w_m]$, and let $D: R \rightarrow R$ be the $\mathbb{Z}$ -derivation determined by $Dw_i = \delta_i$ for some $\delta_1, \dots, \delta_m \in R$.*

(a) (Unnormalized form.) *Suppose*

$$w'_i = \delta_i(w) \quad (1 \leq i \leq m), \quad \frac{f'}{f} = M(w) \text{ for some } M \in R.$$

Then for every $n \geq 0$,

$$f^{(n)} = f \cdot P_n(w), \quad P_0 = 1, \quad P_{n+1} = M P_n + D P_n, \quad (1.1)$$

and $P_n \in R$; in particular every coefficient of P_n is an integer.

(b) (Form normalized by x^{-n} .) *Suppose instead*

$$w'_i = \frac{1}{x} \delta_i(w) \quad (1 \leq i \leq m), \quad x \frac{f'}{f} = M(w) \text{ for some } M \in R.$$

Then for every $n \geq 0$,

$$f^{(n)} = f \cdot x^{-n} P_n(w), \quad P_0 = 1, \quad P_{n+1} = (M - n) P_n + D P_n, \quad (1.2)$$

and again $P_n \in R$.

Proof. For (a), induct on n . The case $n = 0$ is $P_0 = 1$. Assuming $f^{(n)} = f P_n(w)$ and differentiating,

$$f^{(n+1)} = f' P_n(w) + f \sum_{i=1}^m w'_i \frac{\partial P_n}{\partial w_i}(w) = f \left(M(w) P_n(w) + \sum_{i=1}^m \delta_i(w) \frac{\partial P_n}{\partial w_i}(w) \right),$$

which is $f(M P_n + D P_n)(w)$. Since $M, \delta_i \in R$ and $P_0 = 1 \in R$, induction also gives $P_n \in R$, and R consists of polynomials with integer coefficients.

For (b), the case $n = 0$ reads $f = f x^0 \cdot 1$. Assume $f^{(n)} = f x^{-n} P_n(w)$ and differentiate:

$$f^{(n+1)} = f' x^{-n} P_n - n f x^{-n-1} P_n + f x^{-n} \sum_i w'_i \frac{\partial P_n}{\partial w_i}.$$

Multiplying by x^{n+1}/f and using $x f'/f = M$ and $x w'_i = \delta_i$ gives

$$\frac{x^{n+1} f^{(n+1)}}{f} = (M - n) P_n + \sum_i \delta_i \frac{\partial P_n}{\partial w_i} = ((M - n) P_n + D P_n)(w),$$

which is the assertion for $n + 1$. Integrality follows as before. $\square$

Remark 1.3 (Why both clauses are kept). Neither clause covers all three parts *in the frames those parts use*, and since the parts are reproduced here unchanged, both clauses are needed.

Part 3 uses the frame $z = x^2$, $L = \log x$, for which $z' = 2x \notin \mathbb{Z}[z, L]$: clause (a) does not apply, and clause (b) applies verbatim. Conversely Part 4 uses $u = x^x$, $v = x^{-1}$, $z = \log x$, for which $xu' = u(z + 1)/v \notin \mathbb{Z}[u, v, z]$: clause (b) does not apply, while clause (a) does.

It would overstate the situation to say that no single clause *could* have covered all three. Clause (b) in fact does, once the frames are changed so that x appears in place of x^{-1} :

$$x^x : (x, \log x), \quad x^{x^2} : (x^2, \log x), \quad x^{x^x} : (x^x, x, \log x),$$

each of which satisfies $xw'_i \in R$ and $xf'/f \in R$, and each of which gives the same term counts. For x^{x^x} this is legitimate only because the exponent of x^{-1} in that part's envelope never exceeds n , so multiplying by x^n really does clear it. What one cannot do is change the frames and keep the parts: each part's transition table, envelope, coefficient indices and data files are written in its own variables, and rewriting some two and a half thousand lines of dense manipulation to gain a cosmetic uniformity in this section would be a poor trade. So the parts keep their frames and this theorem keeps two clauses.

One genuine obstruction is worth recording separately, since it is about independence rather than convenience. In Part 3 one might try to make clause (a) apply by adjoining $y = x^{-1}$ to the frame $(x^2, \log x)$. The derivatives then are polynomial, but $zy^2 = 1$ identically, so the monomials of the enlarged frame are no longer independent as functions and $|\text{supp } P_n|$ stops counting anything at all. That failure is not a matter of taste.

Each of the three parts instantiates Theorem 1.2 in one line; the resulting recurrences are collected in Table 1.

Part	f	frame and clause	recurrence for P_{n+1}
2	x^x	$t = x^{-1}$, $L = \log x$; (a)	$(1 + L)P_n + t \partial_L P_n - t^2 \partial_t P_n$
3	x^{x^2}	$z = x^2$, $L = \log x$; (b)	$(z(2L + 1) - n)P_n + 2z \partial_z P_n + \partial_L P_n$
4	x^{x^x}	$u = x^x$, $v = x^{-1}$, $z = \log x$; (a)	$u(z^2 + z + v)P_n + u(z + 1)\partial_u P_n - v^2 \partial_v P_n + v \partial_z P_n$

Table 1: The three instances. In each case the displayed recurrence is exactly the one derived independently in the corresponding part (Propositions 2.1, 3.3 and 4.2). Note that z denotes x^2 in Part 3 and $\log x$ in Part 4; see Section 1.5.

1.3 Independence, and why the count is well posed

Counting “the terms of the n th derivative” is meaningless until one fixes what a term is, because a symbolic differentiator can be made to produce almost any number of summands by factoring differently. The following classical statement is what makes the count an invariant of the function rather than of an algorithm. Parts 2 and 3 each prove the special case they need by this argument. Part 4's frame is not of this form, for the reason given after Corollary 1.5, and it proves its own independence statement by a related but genuinely different asymptotic argument.

Theorem 1.4 (Independence of logarithmic monomials). *Let $(\lambda_1, j_1), \dots, (\lambda_N, j_N)$ be distinct pairs with $\lambda_k \in \mathbb{R}$ and $j_k \in \mathbb{Z}_{\geq 0}$. Then the functions $x^{\lambda_k}(\log x)^{j_k}$ are linearly independent over $\mathbb{R}$ on every interval (a, ∞) with $a \geq 0$.*

Proof. Suppose $\sum_k c_k x^{\lambda_k}(\log x)^{j_k} = 0$ on (a, ∞) with not all c_k zero. Substitute $x = e^s$ and group by exponent: the identity becomes

$$\sum_{\lambda} e^{\lambda s} p_{\lambda}(s) = 0 \quad (s > \log \max(a, 1)), \quad (1.3)$$

where λ runs over the distinct values among the λ_k and $p_{\lambda}(s) = \sum_{k: \lambda_k = \lambda} c_k s^{j_k}$ is a polynomial. Because the pairs are distinct, p_{λ} is not the zero polynomial for at least one λ . Let Λ be the

largest such λ . Dividing (1.3) by $e^{\Lambda s}$ gives

$$p_{\Lambda}(s) = - \sum_{\lambda < \Lambda} e^{(\lambda - \Lambda)s} p_{\lambda}(s) \xrightarrow{s \rightarrow +\infty} 0,$$

since each exponent $\lambda - \Lambda$ is strictly negative and an exponential decay beats a polynomial. No nonzero polynomial tends to 0 at $+\infty$, so $p_{\Lambda} = 0$, a contradiction. $\square$

Corollary 1.5 (The count is well defined). *Let f and w be as in either clause of Theorem 1.2, and suppose that the monomials w^{α} of the frame correspond to pairwise distinct pairs $(\lambda(\alpha), j(\alpha))$ in the sense that $w^{\alpha} = x^{\lambda(\alpha)}(\log x)^{j(\alpha)}$. Then the representation of $f^{(n)}$ in (1.1), respectively (1.2), is unique, and the number of distinct monomials in the fully collected n th derivative is*

$$a(n) = |\text{supp } P_n|. \quad (1.4)$$

Proof. Two representations would differ by a finite vanishing linear combination of the functions w^{α} , hence of pairwise distinct $x^{\lambda}(\log x)^j$, which Theorem 1.4 forbids. Uniqueness turns (1.4) into a definition that does not refer to any particular way of performing the differentiation. $\square$

The hypothesis of Corollary 1.5 is verified separately in each part, since it depends on the frame: for x^x the monomials $t^k L^j$ are $x^{-k}(\log x)^j$; for x^{x^2} the monomials $z^k L^j$ carry the extra factor x^{-n} and give $x^{2k-n}(\log x)^j$; for x^{x^x} the variable $u = x^x$ is itself not of the form x^{λ} with constant λ , and Part 4 accordingly proves its independence statement by a related but genuinely different asymptotic argument, which is retained there in full.

1.4 Envelopes, upper bounds and exact computation

Proposition 1.6 (Envelope bound). *Write $P_{n+1} = \Phi_n(P_n)$ for either recurrence of Theorem 1.2, so that $\Phi_n(P) = MP + DP$ in clause (a) and $\Phi_n(P) = (M - n)P + DP$ in clause (b); note that in the second case the operator depends on n , although $\text{supp}((M - n)P) \subseteq \text{supp}(MP) \cup \text{supp } P$ for every n , so the support behaviour does not. Suppose $\mathcal{E}_0, \mathcal{E}_1, \dots$ are finite sets of exponent vectors with $\text{supp } P_0 \subseteq \mathcal{E}_0$ and with the property that Φ_n maps any polynomial supported in $\mathcal{E}_n$ to a polynomial supported in $\mathcal{E}_{n+1}$. Then $\text{supp } P_n \subseteq \mathcal{E}_n$ for every n ; and if moreover the count is well defined in the sense of Corollary 1.5, or by the part's own substitute for it, then*

$$a(n) \leq |\mathcal{E}_n|.$$

Proof. Immediate induction for the support statement. Only the direction matters: Φ_n sends each monomial to a bounded set of monomials, so a cell outside $\mathcal{E}_{n+1}$ receives no contribution at all. Cancellation between contributions can remove a cell of $\mathcal{E}_{n+1}$ from the support, but it can never create one outside. The bound on $a(n)$ is then (1.4). $\square$

The deficit

$$\Delta(n) = |\mathcal{E}_n| - a(n) \geq 0 \quad (1.5)$$

is therefore exactly the number of cells inside the envelope whose coefficient vanishes. This is the shape every one of the three problems takes: an envelope that is easy to count, and a cancellation count that is not. The three parts differ precisely in how much they can say about Δ .

Remark 1.7 (The parts count deficits differently). A reader moving between the parts must not compare their deficit numbers directly. Part 4 writes $\Delta(n)$ for *all* vanishing cells of its envelope, including the ones it proves must vanish; that quantity is itself a published sequence, OEIS

A352697. Part 2 writes $E(n)$ for only the zeros *outside* the families it has already proved to vanish, so E is the residual defect and is conjectured to be identically zero, which Δ certainly is not. Part 3 likewise subtracts its proved reflection zeros before counting. Both conventions are useful and each part keeps its own; the generic Δ of (1.5) is the first kind.

Proposition 1.8 (Exact and modular computation). *With notation as above, $P_0, \dots, P_N$ can be computed in $O(\sum_{n \leq N} |\mathcal{E}_n|)$ integer ring operations, using only integer arithmetic; the implied constant is the frame’s transition count $|\text{supp } M| + \sum_i |\text{supp } \delta_i|$. Moreover, for any modulus $q \geq 2$, a coefficient of P_n whose residue modulo q is nonzero is nonzero.*

Proof. Each recurrence applies at most $|\text{supp } M| + \sum_i |\text{supp } \delta_i|$ monomial-to-monomial transitions to each cell of the support — a constant depending on the frame and not on n — so one step costs $O(|\mathcal{E}_n|)$ operations; the coefficients are integers by Theorem 1.2. Reduction modulo q is a ring homomorphism, so it commutes with the recurrence; a nonzero image forces a nonzero preimage. $\square$

Proposition 1.8 is the reason the computations reported in the three parts are certifications and not samples: a nonzero residue settles nonvanishing outright. The converse direction is the weak one — a zero residue on its own certifies nothing — and the three parts close it in three different ways, which is worth stating precisely rather than in one sentence:

- Part 2 computes the triangle in exact integer arithmetic outright, so no residue has to be resolved; its modular lemma is used for a different purpose, to rule out integer roots of a fixed polynomial over a whole diagonal.
- Part 3 runs two moduli and checks that the two lists of vanishing residues are disjoint, so at least one residue of every coefficient is nonzero. Its exact integer run stops at $n = 200$; the certification to 3000 is modular.
- Part 4 re-evaluates every vanishing residue with an independent exact coefficient formula, and never accepts a zero residue as evidence.

All three are deterministic. No statement anywhere in this article rests on a probabilistic zero test. None of them, however, is an all- n proof: each certifies a finite range, and each part says so where it reports the range.

1.5 Notation across the parts

The three parts below retain the notation of the reports they come from. That notation is not consistent between them, and rewriting it would have meant altering some two and a half thousand lines of dense manipulation with no way to check the result; declaring the clash is safer than silently repairing it.

The table that matters is not a dictionary of letters — nearly every letter is reused somewhere as a summation index, a residue or a generating-function variable, and no short table can be both complete and correct about that. What a reader does need, before crossing from one part to another, is the canonical monomial each part counts, and in particular which of its indices is the exponent of $\log x$.

	Part 2: x^x	Part 3: x^{x^2}	Part 4: x^{x^x}
frame	$t = x^{-1}, L = \log x$	$z = x^2, L = \log x$	$u = x^x, v = x^{-1}, z = \log x$
monomial	$x^{x-k}(\log x)^j$	$x^{x^2+2k-n}(\log x)^j$	$x^{x^x+kx-j}(\log x)^\ell$
log x exponent	j	j	ℓ
power of x^{-1}	k	— (absorbed in x^{-n})	j
normalization	$f^{(n)} = x^x P_n$	$f^{(n)} = f x^{-n} P_n$	$f^{(n)} = f P_n$

Two warnings follow from that table, and they are the two places where a reader who is skimming will go wrong.

1. **j changes role.** In Parts 2 and 3 the index j is the exponent of $\log x$. In Part 4 it is the exponent of x^{-1} , and the exponent of $\log x$ is ℓ . Every envelope, every bound and every cancellation family in Part 4 is indexed in the second convention.
2. **z has three meanings.** It is x^2 in Part 3 and $\log x$ in Part 4 — a direct contradiction — and in Part 2 it is neither: there it is the formal variable in which the Lehmer–Comtet numbers are defined, through $\Phi(z) = (1+z)\log(1+z)$ and $b(m, r) = \frac{m!}{r!} [z^m] \Phi(z)^r$.

A third warning concerns only the later Sections 3.14–3.26 at the end of Part 3: they use the depth coordinates of their source and of the Lean development, in which k **is the exponent of $\log x$** and j the exponent of x — the reverse of the rest of that part — and in which d is the logarithmic deficit. Table 3 gives the full dictionary with the false readings. The second later addition, Sections 3.27–3.34, uses the same coordinates and renames its source’s letters by Table 5; so does the third, Sections 3.35–3.41, by Table 3.A, where in particular the cutoffs $\hat{Q}_d, \hat{K}_d$ are *not* the constants Q_d, K_d of the first addition. [Added 1 October 2026, batch 73: so does the fourth, Sections 3.42–3.51, by Table 3.B, where the checked constants $\check{K}_d, \check{Q}_d$ of one source are neither K_d, Q_d nor $\hat{K}_d, \hat{Q}_d$, and ϱ_d is the largest root, not the bulk count $\beta(N)$.]

A fourth warning concerns the later Sections 2.11–2.15 at the end of Part 2. Their sources call the offset of the Lehmer–Comtet triangle r — which is Part 2’s *column* — and call a multiplier m — which is Part 2’s *row*. They are printed with the offset d of Section 2.5 and the multiplier μ ; Table 2.A gives the full dictionary with the false readings.

Beyond these, the letters u, v, t, q, U, Z, E and Δ all occur in more than one part with unrelated meanings, as frame variables in one place and as summation indices, residues, primes, proposed closed forms or generating-function variables in another. Each is introduced where it is used, and within any one paragraph the meaning is fixed; but nothing in this article guarantees that a letter carries the same meaning two parts later, and the reader should not assume it. Finally, $a(n)$ denotes a different sequence in each of the three parts — its own — and the deficit conventions differ as described in Remark 1.7.

2 x^x : OEIS A293239

Framework. This part instantiates Theorem 1.2(a) with the frame $t = x^{-1}$, $L = \log x$; Proposition 2.1 below is that instance, derived here independently. Lemma 2.2 is the special case of Theorem 1.4 this part needs. Both are retained in full so that the part can be read on its own. Throughout this part $a(n)$ denotes the term count for x^x .

Abstract of the original report. *The sequence A293239 counts the nonzero terms in the fully expanded n th derivative of x^x . Its proposed generating function is rational, and its proposed eventual formula is a quadratic quasipolynomial of period four. We reduce the counting problem exactly to the zero set of the first-kind Lehmer–Comtet numbers. All cancellations predicted by the conjecture are proved: a symmetry family and one additional zero. The converse assertion, excluding every further zero, is not established here; consequently this report does not claim a complete proof of the principal closed formula. The unconditional results include an exact defect formula, upper and lower bounds proving quadratic-order growth, nonvanishing on every prime-offset diagonal with an exact prime-adic valuation, and complete zero classifications for the first three columns and offsets one through sixteen. Independently implemented exact computations confirm the formula through $n = 3000$. A separate conjectured recurrence in the OEIS entry is false with its printed range: it already fails at $n = 8$. For the proposed sequence its correct eventual starting point is $n = 14$. The accompanying code and certificates distinguish finite verification from statements proved for infinitely many indices.*

2.1 The question and the status of the answer

For $x > 0$, put $f(x) = x^x = \exp(x \log x)$. Let $a(n)$ be the number of nonzero summands after $f^{(n)}(x)$ is expanded into the monomials

$$x^{x-k}(\log x)^j, \quad k, j \in \mathbb{Z}_{\geq 0}, \quad (2.1)$$

and coefficients of identical monomials are collected. In particular, an unexpanded expression such as $x^x(1 + \log x)^n$ is not one term for this purpose. This is the convention in OEIS A293239, introduced by Vladimir Reshetnikov in 2017 [1].

The entry attributes the rational generating function and closed formula to Colin Barker and the proposed asymptotic equivalent to Václav Kotešovec. To separate a proved object from the proposed answer, define

$$q(0) = 1, \quad q(n) = 1 + \frac{n(n+1)}{2} - \left\lfloor \frac{n-1}{4} \right\rfloor - \mathbf{1}_{\{n \geq 8\}} \quad (n \geq 1). \quad (2.2)$$

The principal conjecture is $a(n) = q(n)$. For $n \geq 8$, this says simply

$$a(n) \stackrel{?}{=} \frac{n(n+1)}{2} - \left\lfloor \frac{n-1}{4} \right\rfloor. \quad (2.3)$$

The proposed generating function is

$$Q(z) := \sum_{n \geq 0} q(n)z^n = \frac{1 + z^2 + z^3 + z^6 - z^8 + z^9 + z^{12} - z^{13}}{(1-z)^2(1-z^4)}. \quad (2.4)$$

The rational function in (2.4) is indeed the generating function of the explicitly defined sequence q . What remains to be proved is its identification with the derivative count a .

2.1.1 What is proved

The central unconditional identity will be

$$a(n) = q(n) - E(n), \quad (2.5)$$

where $E(n)$ counts a precisely defined set of additional zero coefficients. In particular, $E(n)$ is a nonnegative, nondecreasing integer. Thus the proposed answer is an upper bound, not a formula that can be exceeded by a previously overlooked cancellation.

We prove, for all $n \geq 0$,

$$n + 1 + \left\lfloor \frac{n^2}{4} \right\rfloor \leq a(n) \leq q(n). \quad (2.6)$$

Consequently $a(n) = \Theta(n^2)$. This does not establish the sharper claim $a(n) \sim n^2/2$; that claim is equivalent to $E(n) = o(n^2)$. [Added 1 October 2026, batch 72A: the sharper claim is proved, without an effective threshold, in the sibling report [power-tower-exponent-supports](#) (its Corollary 12.1, the case $a = 1$ of its Theorem 3.1); the bounds (2.6) and (2.34) remain the best explicit ones.] We also prove infinite nonvanishing results that substantially restrict where additional zeros could lie. The remaining gap is stated explicitly in Conjecture 2.9.

2.1.2 A correction that is unconditional

The OEIS entry also proposes

$$a(n) = 2a(n-1) - a(n-2) + a(n-4) - 2a(n-5) + a(n-6) \quad (n > 6). \quad (2.7)$$

This range is incorrect. Exact low-order differentiation gives

$$a(2) = 4, \quad a(3) = 7, \quad a(4) = 11, \quad a(6) = 21, \quad a(7) = 28, \quad a(8) = 35,$$

whereas the right side of (2.7) at $n = 8$ is

$$2 \cdot 28 - 21 + 11 - 2 \cdot 7 + 4 = 36 \neq 35.$$

For q , the recurrence holds for every $n \geq 14$, and it fails at $n = 13$. This correction does not disprove the proposed generating function or closed formula. Section 2.8 explains the distinction.

2.2 A canonical polynomial for each derivative

Set $t = x^{-1}$ and $L = \log x$. There is a polynomial $P_n \in \mathbb{Z}[t, L]$ such that

$$f^{(n)}(x) = x^x P_n(x^{-1}, \log x). \quad (2.8)$$

Differentiating this expression and using $t' = -t^2$, $L' = t$, and $(x^x)' = x^x(1 + L)$ gives the following exact recurrence.

Proposition 2.1. *The derivative polynomials satisfy*

$$P_0 = 1, \quad P_{n+1} = (1 + L)P_n + t \frac{\partial P_n}{\partial L} - t^2 \frac{\partial P_n}{\partial t}. \quad (2.9)$$

In particular, they have integer coefficients and can be computed using integer arithmetic only.

For example,

$$\begin{aligned} P_1 &= 1 + L, \\ P_2 &= 1 + 2L + L^2 + t, \\ P_3 &= 1 + 3L + 3L^2 + L^3 + 3t + 3tL - t^2. \end{aligned}$$

Thus $a(3) = 7$, consistently with the example in the entry.

Lemma 2.2. *The monomials in (2.1) are linearly independent over $\mathbb{R}$ when only finitely many are used. Consequently the collected expansion is unique, and $a(n) = |\text{supp } P_n|$.*

Proof. Divide a putative relation by the positive function x^x and set $x = e^s$. The result is $\sum_{k=0}^K e^{-ks} p_k(s) = 0$, with polynomial p_k . If some p_k is nonzero, let k_0 be the smallest such index. Multiplication by $e^{k_0 s}$ shows that $p_{k_0}(s)$ tends to zero as $s \rightarrow +\infty$, because all remaining summands decay exponentially times a polynomial. No nonzero polynomial has that property. This is a contradiction. $\square$

The independence statement matters: the variables t and L are related by $t = e^{-L}$ after substitution, but this relation causes no additional finite polynomial identities.

2.3 The Lehmer–Comtet coefficient triangle

2.3.1 Definition and generating function

Put

$$\Phi(z) = (1+z) \log(1+z) = z + \sum_{d \geq 1} \frac{(-1)^{d-1}}{d(d+1)} z^{d+1}. \quad (2.10)$$

Define

$$b(m, r) = \frac{m!}{r!} [z^m] \Phi(z)^r \quad (m, r \geq 0). \quad (2.11)$$

Here $b(0, 0) = 1$, $b(m, 0) = 0$ for $m > 0$, and $b(m, r) = 0$ when $r > m$. These are the first-kind Lehmer–Comtet numbers, tabulated in OEIS A008296 [4]. The general derivative problem and these numbers belong to the classical literature of Comtet, Lehmer, and Gould [7, 8, 9]. A recent treatment of their identities is Jeong’s preprint [10]. We derive the identities needed here rather than assuming a nonvanishing conclusion from those formulas.

Theorem 2.3 (Exact coefficient formula). *For every $n \geq 0$,*

$$f^{(n)}(x) = x^x \sum_{m=0}^n \binom{n}{m} (\log x)^{n-m} \sum_{r=0}^m b(m, r) x^{r-m}. \quad (2.12)$$

Equivalently,

$$[t^k][L^j] P_n(t, L) = \binom{n}{j} b(n-j, n-j-k), \quad (2.13)$$

with out-of-range entries interpreted as zero.

Proof. Taylor expansion at a fixed $x > 0$ gives

$$\sum_{n \geq 0} \frac{f^{(n)}(x)}{x^x} \frac{u^n}{n!} = \frac{(x+u)^{x+u}}{x^x} = \exp(u \log x) \exp(x \Phi(u/x)).$$

Using (2.11), expand the second exponential:

$$\begin{aligned}\exp(x\Phi(u/x)) &= \sum_{r \geq 0} \frac{x^r}{r!} \Phi(u/x)^r \\ &= \sum_{m \geq 0} \frac{u^m}{m!} \sum_{r=0}^m b(m, r) x^{r-m}.\end{aligned}$$

Multiplication by $\exp(u \log x)$ and extraction of the coefficient of u^n proves (2.12). All manipulations can also be performed formally, since only finitely many terms affect any prescribed coefficient. $\square$

2.3.2 A shifted falling-factorial formula

Let

$$F_m(y) = y^m = \prod_{h=0}^{m-1} (y - h), \quad F_0(y) = 1.$$

Proposition 2.4. For $0 \leq r \leq m$,

$$b(m, r) = \frac{F_m^{(r)}(r)}{r!} = [u^r] \prod_{h=0}^{m-1} (u + r - h). \quad (2.14)$$

In particular, all $b(m, r)$ are integers.

Proof. The binomial-series identity is $(1+z)^y = \sum_{m \geq 0} F_m(y) z^m / m!$. Differentiate r times in y and set $y = r$. The left side becomes $(1+z)^r \log(1+z)^r = \Phi(z)^r$. Comparing coefficients gives the first formula. The second is the polynomial Taylor formula at $y = r$. $\square$

Writing $F_m(y) = \sum_{\ell=0}^m s(m, \ell) y^\ell$ in terms of the signed Stirling numbers of the first kind also gives

$$b(m, r) = \sum_{\ell=r}^m \binom{\ell}{r} r^{\ell-r} s(m, \ell). \quad (2.15)$$

This is another exact formula, but its alternating summands do not by themselves resolve whether the result can vanish.

2.3.3 An integer recurrence

Proposition 2.5. For $m \geq 2$ and $1 \leq r \leq m$,

$$b(m, r) = (r - m + 1)b(m - 1, r) + b(m - 1, r - 1) + (m - 1)b(m - 2, r - 1). \quad (2.16)$$

Together with rows $b(0, 0) = 1$ and $(b(1, 0), b(1, 1)) = (0, 1)$, this determines the entire triangle.

Proof. Let $B_r(z) = \Phi(z)^r / r!$. Since $(1+z)\Phi'(z) = 1 + z + \Phi(z)$, we have

$$(1+z)B'_r(z) = (1+z)B_{r-1}(z) + rB_r(z).$$

Compare coefficients of $z^{m-1} / (m-1)!$. The result is

$$b(m, r) + (m - 1)b(m - 1, r) = b(m - 1, r - 1) + (m - 1)b(m - 2, r - 1) + rb(m - 1, r),$$

which rearranges to (2.16). $\square$

For reference, the first rows, omitting column zero, are

$m \backslash r$	1	2	3	4	5	6	7	8	9
1	1								
2	1	1							
3	-1	3	1						
4	2	-1	6	1					
5	-6	0	5	10	1				
6	24	4	-15	25	15	1			
7	-120	-28	49	-35	70	21	1		
8	720	188	-196	49	0	154	28	1	
9	-5040	-1368	944	0	-231	252	294	36	1

2.4 The exact zero-count reduction

Definition 2.6. For $n \geq 0$, let

$$Z(n) = \#\{(m, r) : 1 \leq r \leq m \leq n, b(m, r) = 0\}.$$

Column zero is not included: its zeros for $m > 0$ are forced by the definition and never represent potential nonzero derivative terms.

Theorem 2.7. For every $n \geq 0$,

$$a(n) = 1 + \frac{n(n+1)}{2} - Z(n). \quad (2.17)$$

Proof. In (2.12), the pair (m, r) gives the monomial

$$x^{x-(m-r)}(\log x)^{n-m}$$

with coefficient $\binom{n}{m}b(m, r)$. The map $(m, r) \mapsto (m-r, n-m)$ is injective for fixed n : the second coordinate recovers m , and the first then recovers r . Thus different pairs do not cancel each other. The factor $\binom{n}{m}$ is positive. There is one nonzero entry at $(0, 0)$, and m potentially nonzero entries $1 \leq r \leq m$ in every row $m \geq 1$. Removing exactly the zero entries proves the formula. $\square$

This argument isolates the main difficulty. Once the triangle is known, term counting is easy; proving that its list of zeros is complete is not a consequence of merely writing down the triangle's recurrence.

2.4.1 Cancellations that are proved

Proposition 2.8 (Symmetry zeros). For every integer $h \geq 1$,

$$b(4h+1, 2h) = 0. \quad (2.18)$$

Proof. Center the product in (2.14) at $r = 2h$:

$$F_{4h+1}(2h+u) = u \prod_{s=1}^{2h} (u^2 - s^2).$$

This polynomial is odd. Its coefficient of the even power u^{2h} is zero. $\square$

This family is not claimed as new: it appears as Proposition 3.10(iii) in Jeong [10], with attribution there to Theorem 7 of Lehmer [8]. Its elementary proof is included to make the reduction self-contained.

There is one further zero required by the proposed formula:

$$b(8, 5) = 0. \quad (2.19)$$

It follows immediately from the factorization proved in Section 2.5:

$$b(m, m-3) = \frac{m(m-1)(m-2)(m-3)(m-5)(m-8)}{48}.$$

The root $m = 5$ is the first symmetry zero; $m = 8$ is genuinely additional.

2.4.2 The precise missing assertion

Conjecture 2.9 (Complete zero classification). For integers $1 \leq r \leq m$,

$$b(m, r) = 0 \iff (m, r) = (8, 5) \text{ or } (m, r) = (4h + 1, 2h) \text{ for some } h \geq 1. \quad (2.20)$$

The right-to-left implication is proved above. The left-to-right implication is the unproved step in this report.

Let

$$\mathcal{Z}_0 = \{(8, 5)\} \cup \{(4h + 1, 2h) : h \geq 1\}$$

be the proved set of zeros, and define

$$e_m = \#\{r : 1 \leq r \leq m, b(m, r) = 0, (m, r) \notin \mathcal{Z}_0\},$$

$$E(n) = \sum_{m=1}^n e_m.$$

Thus E counts only zeros not in the proved list.

Theorem 2.10 (Exact defect formula). For every $n \geq 0$,

$$a(n) = q(n) - E(n).$$

The following are equivalent: Conjecture 2.9; $E(n) = 0$ for all n ; $a(n) = q(n)$ for all n ; and the proposed eventual formula (2.3) for every $n \geq 8$.

Proof. For $n \geq 1$, the number of symmetry zeros with first coordinate at most n is $\lfloor (n-1)/4 \rfloor$. The zero $(8, 5)$ contributes $\mathbf{1}_{\{n \geq 8\}}$. Substituting

$$Z(n) = \left\lfloor \frac{n-1}{4} \right\rfloor + \mathbf{1}_{\{n \geq 8\}} + E(n)$$

into (2.17) proves the identity for $n \geq 1$; $n = 0$ is immediate. The first three equivalences follow from the definitions. The eventual formula also forces $E(n) = 0$ for every $n \geq 8$, and nonnegativity and monotonicity then force $E(n) = 0$ for smaller n as well. $\square$

A hypothetical new zero at row M would decrease every count $a(n)$ for $n \geq M$ relative to $q(n)$. It would not merely alter a single isolated term of the sequence. This persistent-deficit property is a useful safeguard when testing proposed proofs or numerical data.

2.5 Diagonal polynomials and exact nonvanishing

2.5.1 A polynomial for each fixed offset

Define

$$H(z) = \frac{\Phi(z)}{z} = 1 + \frac{z}{2} - \frac{z^2}{6} + \frac{z^3}{12} - \frac{z^4}{20} + \cdots, \quad T_d(Y) = [z^d]H(z)^Y. \quad (2.21)$$

For a fixed integer $d \geq 0$, T_d is a polynomial over $\mathbb{Q}$ in Y .

Lemma 2.11. *For $m \geq d$,*

$$b(m, m-d) = m^d T_d(m-d). \quad (2.22)$$

For $d \geq 1$, $T_d(Y)$ has degree d , leading coefficient $1/(2^d d!)$, and a factor Y . Consequently $T_d(Y)/Y$ is a nonzero polynomial of degree $d-1$.

Proof. Since $\Phi(z)^{m-d} = z^{m-d} H(z)^{m-d}$, definition (2.11) gives (2.22). Now assume $d \geq 1$ and write $h(z) = H(z) - 1$. Then

$$T_d(Y) = \sum_{\ell=1}^d \binom{Y}{\ell} [z^d] h(z)^\ell. \quad (2.23)$$

Every summand is divisible by Y . The only term of degree d in Y comes from $\ell = d$, and $[z^d]h(z)^d = 2^{-d}$. This proves the remaining assertions. $\square$

The first three cases are

$$b(m, m-1) = \frac{m(m-1)}{2}, \quad (2.24)$$

$$b(m, m-2) = \frac{m(m-1)(m-2)(3m-13)}{24}, \quad (2.25)$$

$$b(m, m-3) = \frac{m(m-1)(m-2)(m-3)(m-5)(m-8)}{48}. \quad (2.26)$$

For instance,

$$T_3(Y) = \frac{Y}{12} - \frac{Y(Y-1)}{12} + \frac{Y(Y-1)(Y-2)}{48} = \frac{Y(Y-2)(Y-5)}{48}.$$

This proves both zeros on the third offset and excludes all others on that entire diagonal, not just up to a numerical cutoff.

2.5.2 Prime offsets: an infinite family of nonvanishing results

For a nonzero rational number v , write $v_p(v)$ for its exponent of the prime p in its prime factorization. A rational number is called p -integral when its denominator in lowest terms is not divisible by p .

Theorem 2.12 (Prime-offset nonvanishing). *Let p be any prime and let $m \geq p$. Then*

$$b(m, m-p+1) \neq 0, \quad v_p(b(m, m-p+1)) = v_p\left(\left\lfloor \frac{m}{p} \right\rfloor\right). \quad (2.27)$$

In particular, an additional zero cannot occur on any offset $d = m - r$ for which $d+1$ is prime.

Proof. Put $d = p - 1$ and $h_j = [z^j](H(z) - 1)$. For $1 \leq j \leq p - 2$, the number $h_j = (-1)^{j-1}/(j(j+1))$ is p -integral. Moreover,

$$ph_{p-1} \equiv 1 \pmod{p}.$$

For odd p , this follows from $ph_{p-1} = -1/(p-1)$; for $p = 2$, it is $2h_1 = 1$.

Divide (2.23) by Y . Its $\ell = 1$ summand is h_{p-1} . For every $2 \leq \ell \leq p - 1$, a composition of $p - 1$ into ℓ positive parts has all parts at most $p - 2$. Therefore every coefficient contributing to $[z^{p-1}]h(z)^\ell$ is p -integral. Also,

$$\frac{1}{Y} \binom{Y}{\ell} = \frac{(Y-1) \cdots (Y-\ell+1)}{\ell!}$$

has p -integral coefficients, because $\ell!$ is not divisible by p . It follows, as a polynomial congruence over the p -integral rationals, that

$$p \frac{T_{p-1}(Y)}{Y} \equiv 1 \pmod{p}. \quad (2.28)$$

For every positive integer Y , the left side is therefore nonzero and has p -adic valuation zero. Hence $v_p(T_{p-1}(Y)) = v_p(Y) - 1$.

Set $Y = m - p + 1 > 0$ and use (2.22). This yields

$$v_p(b(m, m - p + 1)) = v_p(m^p) - 1.$$

Among the p consecutive positive integers $m - p + 1, \dots, m$, exactly one is divisible by p : it is $p \lfloor m/p \rfloor$. Its valuation is $1 + v_p(\lfloor m/p \rfloor)$, proving (2.27). $\square$

This proof explains, for example, why the residual polynomial $3m - 13$ in (2.25) is a nonzero constant modulo 3, and why analogous residual polynomials for offsets 4, 6, 10, 12, 16 are nonzero constants modulo 5, 7, 11, 13, 17, respectively.

2.5.3 Finite certificates that cover infinitely many row indices

For a fixed $d \geq 1$, put

$$S_d(X) = T_d(X - d).$$

The factor $X - d$ is forced by Lemma 2.11. When $d \geq 3$ is odd, the symmetry zero also forces $X - (2d - 1)$; for $d = 3$ there is in addition the factor $X - 8$. Remove these factors from S_d , clear rational denominators, divide out the greatest common divisor of the integer coefficients, and normalize the leading coefficient to be positive. Call the resulting polynomial $R_d(X)$. Thus, with a positive rational constant c_d ,

$$S_d(X) = c_d(X - d) \begin{cases} (X - 5)(X - 8)R_d(X), & d = 3, \\ (X - (2d - 1))R_d(X), & d \geq 5 \text{ odd}, \\ R_d(X), & \text{otherwise.} \end{cases} \quad (2.29)$$

Lemma 2.13 (Modular certificate). *If $R \in \mathbb{Z}[X]$ satisfies $R(t) \not\equiv 0 \pmod{p}$ for every $t \in \{0, 1, \dots, p - 1\}$, then R has no integer root.*

Proof. An integer root, reduced modulo p , would be one of those residues and would give a zero value modulo p . $\square$

Theorem 2.14 (Computer-assisted diagonal classification). *Conjecture 2.9 holds for every pair $1 \leq r \leq m$ with $m - r \leq 16$.*

Proof. For offset zero, $b(m, m) = 1$. For $1 \leq d \leq 16$, compute S_d exactly from (2.23) and factor it as in (2.29). The supplied `data/diagonal_certificates.json` records the full rational coefficients of S_d , the scalar c_d , all removed linear factors, the full integer coefficients of R_d , and every value of $R_d(t)$ modulo the certifying prime. All of those values are nonzero. The following summary identifies the certificates:

d	$\deg R_d$	p	roots mod p	d	$\deg R_d$	p	roots mod p
1	0	2	0	9	7	7	0
2	1	3	0	10	9	11	0
3	0	2	0	11	9	13	0
4	3	5	0	12	11	13	0
5	3	3	0	13	11	11	0
6	5	7	0	14	13	23	0
7	5	5	0	15	13	19	0
8	7	11	0	16	15	17	0

The standard-library program `code/verify_diagonals.py` reconstructs every polynomial using rational arithmetic and checks the factorization and each residue. The modular lemma excludes every integer root of every R_d . For $m > d$, the removed factor $m - d$ cannot vanish; the remaining removed factors give exactly the stated symmetry and exceptional zeros. $\square$

These certificates are not extrapolations from finitely many values of m : each modular argument excludes integer roots for *all* m on its fixed diagonal. What is finite is the set of offsets treated. For a small example not requiring the machine-readable file,

$$\begin{aligned} R_4(X) &= 15X^3 - 390X^2 + 3245X - 8638 \equiv 2 \pmod{5}, \\ R_5(X) &= 3X^3 - 103X^2 + 1118X - 3776 \equiv 2X^2 + 2X + 1 \pmod{3}. \end{aligned}$$

The second reduced polynomial takes the values 1, 2, 1 at the three residues modulo 3. Hence there are no additional zeros on offset 4, and the only zero on offset 5 is the symmetry zero $m = 9$.

2.6 The first three columns for all row indices

The diagonal results keep $m - r$ fixed. A complementary calculation keeps r fixed and proves nonvanishing in a different infinite region.

Proposition 2.15. *Column $r = 1$ has no zero. Column $r = 2$ has exactly the zero $m = 5$. Column $r = 3$ has no zero.*

Proof. For $r = 1$, (2.14) gives

$$b(1, 1) = 1, \quad b(m, 1) = (-1)^m(m-2)! \quad (m \geq 2).$$

For $r = 2$, put $t = m - 3 \geq 0$. The shifted product is

$$F_m(2+u) = u(u+1)(u+2) \prod_{j=1}^t (u-j).$$

The coefficient of u^2 is

$$b(m, 2) = (-1)^m(m-3)!(2H_{m-3} - 3), \quad m \geq 3, \quad (2.30)$$

where $H_t = \sum_{j=1}^t 1/j$ and $H_0 = 0$. The harmonic numbers strictly increase, and $H_2 = 3/2$, so the only zero is $m-3 = 2$. The remaining boundary entry is $b(2, 2) = 1$.

For $r = 3$, the boundary entry is $b(3, 3) = 1$. Put $t = m - 4 \geq 0$ and define

$$H_t^{(2)} = \sum_{j=1}^t \frac{1}{j^2}, \quad g(t) = 6 - 11H_t + 3(H_t^2 - H_t^{(2)}).$$

Extracting the coefficient of u^3 from $u(u+1)(u+2)(u+3) \prod_{j=1}^t (u-j)$ gives

$$b(m, 3) = (-1)^t t! g(t). \quad (2.31)$$

The elementary identity

$$g(t+1) - g(t) = \frac{6H_t - 11}{t+1} \quad (2.32)$$

controls its entire sign pattern. We have $g(0) = 6$, $g(1) = -5$, and $H_3 = 11/6$. Thus g decreases through $t = 3$, has $g(3) = g(4) = -49/6$, and is strictly increasing thereafter. Finally, exact rational evaluation gives

$$g(19) = -\frac{4571462267}{97772875200} < 0, \quad g(20) = \frac{537826687}{1150269120} > 0.$$

It follows that $g(t) < 0$ for $1 \leq t \leq 19$, whereas $g(t) > 0$ for $t \geq 20$. There is no integer zero. $\square$

The last argument illustrates a distinction from a numerical search: exact values at two adjacent integers, together with a proved monotonicity statement outside a finite initial segment, settle the entire column.

2.7 Unconditional quadratic growth

Theorem 2.16. *For every integer $n \geq 0$,*

$$n + 1 + \left\lfloor \frac{n^2}{4} \right\rfloor \leq a(n) \leq q(n). \quad (2.33)$$

In particular,

$$\frac{1}{4} \leq \liminf_{n \rightarrow \infty} \frac{a(n)}{n^2} \leq \limsup_{n \rightarrow \infty} \frac{a(n)}{n^2} \leq \frac{1}{2}, \quad a(n) = \Theta(n^2).$$

Proof. The upper bound is Theorem 2.10. For the lower bound, count nonzero entries along diagonals of the triangle in Theorem 2.7. The main diagonal, including $(0, 0)$, contributes $n + 1$ entries.

Fix $1 \leq d \leq n - 1$. The diagonal $m - r = d$ has $n - d$ relevant positions, $m = d + 1, \dots, n$. By Lemma 2.11, its zero positions are roots of the nonzero polynomial $T_d(m - d)/(m - d)$ of degree $d - 1$. There are therefore at most $d - 1$ such positions, and at least

$$\max(0, n - d - (d - 1)) = \max(0, n - 2d + 1)$$

nonzero entries on the diagonal. Summing over all $d \geq 1$ gives

$$a(n) \geq n + 1 + \sum_{d \geq 1} \max(0, n - 2d + 1) = n + 1 + \left\lfloor \frac{n^2}{4} \right\rfloor.$$

The last equality follows by summing the odd numbers $1, 3, \dots, 2s - 1$ when $n = 2s$, and the even numbers $2, 4, \dots, 2s$ when $n = 2s + 1$. The asymptotic bounds follow immediately. $\square$

Corollary 2.17 (A stronger explicit lower bound). *For every $n \geq 0$,*

$$a(n) \geq n + 1 + \left\lfloor \frac{n^2}{4} \right\rfloor + \sum_{\substack{p \leq n \\ p \text{ prime}}} \min(p - 2, n - p + 1). \quad (2.34)$$

Proof. On offset $d = p - 1$, Theorem 2.12 guarantees all $n - d = n - p + 1$ entries, rather than just $\max(0, n - 2p + 3)$. The improvement is $\min(p - 2, n - p + 1)$. These diagonals are disjoint, so their improvements add. $\square$

The prime-enhanced bound is useful without invoking any theorem about the density of primes. It still does not close the factor-of-two gap in the leading quadratic coefficient. In terms of the exact defect,

$$a(n) \sim \frac{n^2}{2} \iff E(n) = o(n^2). \quad (2.35)$$

Thus the proposed asymptotic equivalent is potentially a weaker target than the complete zero classification: it would allow a sufficiently sparse set of additional zeros. [Added 1 October 2026, batch 72A: the weaker target is now proved: the sibling report **power-tower-exponent-supports** shows $E(n) = o(n^2)$, hence $a(n) \sim n^2/2$ (its Corollary 12.1), without an effective threshold. Conjecture 2.9, and with it the closed formula, remains open.]

2.8 The rational function, quasipolynomial, and recurrence

2.8.1 Deriving the proposed generating function

There is one potential term at $(0, 0)$ and m potential terms in row m . The cumulative potential count therefore has generating function

$$\sum_{n \geq 0} \left(1 + \frac{n(n+1)}{2}\right) z^n = \frac{1}{1-z} + \frac{z}{(1-z)^3}.$$

The symmetry zeros are born at rows 5, 9, 13, $\dots$, and each remains absent from every later derivative. Their cumulative generating function is $z^5/((1-z)(1-z^4))$. The exceptional zero is born at row 8 and contributes $z^8/(1-z)$. Subtracting these two *proved* families gives

$$\begin{aligned} Q(z) &= \frac{1}{1-z} + \frac{z}{(1-z)^3} - \frac{z^5}{(1-z)(1-z^4)} - \frac{z^8}{1-z} \\ &= \frac{1 + z^2 + z^3 + z^6 - z^8 + z^9 + z^{12} - z^{13}}{(1-z)^2(1-z^4)}. \end{aligned} \quad (2.36)$$

This explains every correction term in the proposed rational function.

Theorem 2.18 (Exact generating-function correction). *With $A(z) = \sum_{n \geq 0} a(n)z^n$ and $\mathcal{E}(z) = \sum_{m \geq 1} e_m z^m$,*

$$A(z) = Q(z) - \frac{\mathcal{E}(z)}{1-z}. \quad (2.37)$$

Consequently the proposed rational function equals $A(z)$ if and only if Conjecture 2.9 is true.

Proof. Sum $a(n) = q(n) - \sum_{m \leq n} e_m$ as a formal power series. Each e_m contributes $e_m z^m / (1 - z)$ to the cumulative sum. $\square$

2.8.2 The real form of the complex expression

For $n \geq 8$, division into the four residue classes gives

$$q(n) = \frac{n^2}{2} + \frac{n}{4} + \begin{cases} 1, & n \equiv 0 \pmod{4}, \\ \frac{1}{4}, & n \equiv 1 \pmod{4}, \\ \frac{1}{2}, & n \equiv 2 \pmod{4}, \\ \frac{3}{4}, & n \equiv 3 \pmod{4}. \end{cases} \quad (2.38)$$

The same periodic correction is

$$\frac{5 + (-1)^n + (1 - i)(-i)^n + (1 + i)i^n}{8}, \quad i^2 = -1.$$

Substitution yields exactly the complex-valued-looking expression printed in OEIS, whose total value is real and integral. This is an algebraic identity for q , independent of the unresolved equality $a = q$.

For all $n \geq 0$, another convenient form is

$$q(n) = \frac{n(n+1)}{2} - \left\lfloor \frac{n-1}{4} \right\rfloor + \mathbf{1}_{\{1 \leq n \leq 7\}}.$$

Here $\lfloor -1/4 \rfloor = -1$ correctly gives $q(0) = 1$. The first terms are

$$1, 2, 4, 7, 11, 15, 21, 28, 35, 43, 53, 64, 76, 88, 102, 117, 133, \dots$$

2.8.3 Why the recurrence starts at fourteen

Set

$$\begin{aligned} D(z) &= (1 - z)^2(1 - z^4) = 1 - 2z + z^2 - z^4 + 2z^5 - z^6, \\ N(z) &= 1 + z^2 + z^3 + z^6 - z^8 + z^9 + z^{12} - z^{13}. \end{aligned}$$

The identity $D(z)Q(z) = N(z)$ implies, for $n \geq 6$,

$$q(n) - 2q(n-1) + q(n-2) - q(n-4) + 2q(n-5) - q(n-6) = [z^n]N(z). \quad (2.39)$$

The right side vanishes for every $n \geq 14$, but at $n = 13$ it equals -1 . Thus the least starting index from which the homogeneous recurrence holds for *every* subsequent index of q is 14. For $7 \leq n \leq 13$, its only nonzero residuals are

n	8	9	12	13
residual	-1	1	1	-1

This proves the range correction asserted at the start of this part. If the original derivative sequence satisfies the corrected recurrence for all $n \geq 14$, its directly verified initial values $a(0), \dots, a(13)$ force $a = q$ by induction. Consequently the corrected recurrence for a is itself equivalent to the main conjecture, once those initial values are supplied; it is not an independent proved consequence for arbitrary orders.

The second forward differences $q(n+2) - 2q(n+1) + q(n)$ begin

$$1, 1, 1, 0, 2, 1, 0, 1, \underbrace{2, 1, 1, 0, 2, 1, 1, 0, \dots}_{n \geq 8}$$

Again, this is proved for q ; transferring the entire pattern to a requires the missing nonvanishing statement.

2.9 Reproducible verification

2.9.1 What was actually checked

All reported checks use exact arithmetic. No floating-point threshold was used to decide that a coefficient was zero.

The C++ implementation with GMP computes the recurrence (2.16) through row 3000. It tests all 4,501,500 entries with $1 \leq r \leq m \leq 3000$. Exactly 750 are zero: 749 symmetry zeros and (8, 5). There are no other zeros in that range, and therefore

$$a(n) = q(n) \quad (0 \leq n \leq 3000), \quad a(3000) = 4,500,751. \quad (2.40)$$

These are finite computational results, not an induction proving all n .

A separate Python standard-library implementation computes the triangle through row 1500 and agrees with the C++ term counts at all 1501 indices $0, \dots, 1500$. Further independent checks include the complete polynomials from the differential recurrence (2.9) through order 60, all 496 shifted-product coefficients (2.14) with $0 \leq r \leq m \leq 30$, and 11,865 instances of the prime-adic valuation formula. All 54 values displayed directly in the OEIS entry agree with the exact computation. The separately linked OEIS table through 500 was not used as an independent data source for this report.

The sixteen diagonal certificates were reconstructed and checked with a third, rational-polynomial computation. In addition to showing no unexpected zeros in a finite triangular region, these certificates prove there are no unexpected zeros on those fixed diagonals for any row index.

2.9.2 Algorithm and complexity

Only two preceding rows are required for (2.16). A schematic implementation is

```
previousprevious = [1]
previous = [0, 1]
count = 2
for m = 2, ..., N:
    current[0] = 0
    for r = 1, ..., m:
        current[r] = (r-m+1)*previous[r]
            + previous[r-1] + (m-1)*previousprevious[r-1]
    count += number of nonzero current[r] for 1 <= r <= m
    rotate the three rows
```

Indices beyond the end of a row mean zero. The supplied implementations handle these boundaries explicitly.

The computation uses $O(N^2)$ arithmetic operations and stores $O(N)$ integers. These are not bit-complexity claims: the integers grow with N . From the product formula,

$$|b(m, r)| \leq \prod_{h=0}^{m-1} (1 + |r - h|) \leq (m + 1)^m,$$

so a coefficient has $O(m \log m)$ bits. The rolling-row implementation thus has an $O(N^2 \log N)$ -bit storage bound. A crude schoolbook arithmetic bound for the running time is $O(N^3 \log^2 N)$ bit operations.

2.9.3 Files and commands

The archive contains `article.tex`, the compiled `article.pdf`, a build script, both triangle verifiers, the rational certificate verifier, the complete count and zero-position tables through 3000, and the exact verification reports. From the extracted directory, run

```
python3 code/verify.py -max-n 1500 -data-dir data
python3 code/verify_diagonals.py -output-dir data -check-only
```

The C++ implementation requires a C++17 compiler and GMP development files:

```
c++ -O3 -std=c++17 code/verify_gmp.cpp -lgmpxx -lgmp -o verify_gmp
./verify_gmp 3000 data
```

The Python programs use only the standard library. The diagonal checker constructs the coefficients afresh rather than trusting the saved residue tables. Its proof obligations are elementary rational polynomial identities and finite modular evaluations; it does not require a general-purpose computer algebra system.

2.10 What remains to finish the requested proof

The derivation of the candidate rational function is complete, but the identification of that function with the derivative count is conditional. The exact outstanding statement is

$$[u^r] \prod_{h=0}^{m-1} (u + r - h) \neq 0$$

for all $1 \leq r \leq m$ outside the explicitly proved zero set $\mathcal{Z}_0$. The preceding results settle this for $r \leq 3$, for $m - r \leq 16$, and whenever $m - r + 1$ is prime, in addition to all rows through 3000 by exact computation. They do not cover every remaining pair.

The diagonal-polynomial formulation gives a particularly precise route to a complete proof. For every offset d , it would suffice to show that the residual polynomial R_d in (2.29) has no integer root $m > d$. Irreducibility would be stronger than necessary. A uniform family of modular nonvanishing certificates would also suffice, but the fact that such certificates work for $d \leq 16$ does not establish that they work for all d .

Update (1 October 2026, batch 72A). The later Subsections 2.11–2.15 add infinitely many settled offsets to the list above: every offset $d = \mu(p - 1)$ with $p > \mu$ prime under the conditions of Theorem 2.19, in particular for all $p \geq \max\{4\mu, 1 + 3^{4\mu-7}\}$; all offsets $2(p - 1)$ and $4(p - 1)$, and $3(p - 1)$ and $5(p - 1)$ for odd p (Corollaries 2.20, 2.21, 2.22 and 2.35). On those offsets the only zeros are the structural ones in column zero. The offset $(p - 1)^2$ is reduced to the single candidate $m = 2p^2 - p - 1$ (that is, $Y = p^2 + p - 2$; Corollary 2.36) and the offset $p^2 - 1$ to

the columns $1 \leq r \leq p-1$ for $p \geq 5$ (Theorem 2.38); exact p -adic valuations exclude zeros on divisibility progressions of the column (Theorems 2.39 and 2.40). On the route of the previous paragraph, Theorem 2.28 shows that at the offsets $p^2 - 1$ no sieve modulo a fixed power p^e of that same prime can work by itself (it says nothing about other moduli), and the positivity cone of Corollary 2.25 confines every zero on offset d to the rows $m < 5d$. Conjecture 2.9 is still open.

For the asymptotic conjecture alone, an exclusion of every additional zero is unnecessary: by (2.35), it suffices to bound their cumulative number by $o(n^2)$. The present root-counting argument proves quadratic order but not this sharper estimate. [Added 1 October 2026, batch 72A: the sharper estimate is proved in the sibling report **power-tower-exponent-supports**, by a contour argument rather than root counting; its Section 12 prints manuscript 31's proof at $a = 1$, a second route to the case $a = 1$ of its Theorem 3.1. It gives no rate and no threshold, so Theorem 2.16 and Corollary 2.17 remain the explicit bounds.]

Conclusion. The printed recurrence with range $n > 6$ is disproved, and its correct range for the proposed sequence is identified. The principal closed formula is supported by an exact structural reduction, several infinite nonvanishing theorems, and extensive reproducible verification. A complete all-order proof of that formula is *not* supplied: the missing converse in Conjecture 2.9 is left explicit rather than being replaced by an inference from numerical agreement.

2.11 Lehmer–Comtet zeros on composite offsets: three later additions

What this addition is. Subsections 2.11–2.15 were written into this part on 1 October 2026 from three later manuscripts on the first-kind Lehmer–Comtet triangle, all dated 1 October 2026 and all pinned to this report's text at commit `ca81647a9`:

- manuscript 12, *Complete Nonvanishing on Prime-Multiple Offsets* (Subsection 2.12);
- manuscript 09, *Stirling Residues and Finite Candidate Lists* (Subsection 2.13), written after manuscript 12 and building on it;
- manuscript 17, *Uniform p -adic Nonvanishing* (Subsection 2.14).

The numbers are those of batch 72A; provenance is in Subsection 2.15. The three manuscripts attack Conjecture 2.9 along the route of Subsection 2.5 — one fixed offset $d = m - r$ at a time — and extend Theorem 2.12 from the offsets $d = p - 1$ to composite offsets. **Conjecture 2.9 remains open, and with it the principal closed formula $a(n) = q(n)$; each of the three manuscripts says so.**

What is new relative to this part.

- Every offset $d = \mu(p - 1)$ with $p > \mu$ prime is zero-free under two explicit conditions, which hold for every prime $p \geq \max\{4\mu, 1 + 3^{4\mu-7}\}$ (Theorem 2.19). The offsets $2(p - 1)$ and $4(p - 1)$ are zero-free for every prime p , and $3(p - 1)$ and $5(p - 1)$ for every odd prime (Corollaries 2.20, 2.22, 2.21 and 2.35).
- The universal positivity cone: $T_d(Y) > 0$ whenever $Y \geq 4d$ (Corollary 2.25), so every zero on offset d lies in a row $m < 5d$.
- An exact Stirling residue for the small columns of the offset $\mu(p - 1)$, the p -adic displacement of the roots, and a complete list of candidate zeros for every such offset (Theorems 2.30, 2.32 and 2.33).
- The offset $(p - 1)^2$ is reduced to the single candidate $Y = p^2 + p - 2$ (Corollary 2.36), and the offset $p^2 - 1$ to $1 \leq Y \leq p - 1$ for $p \geq 5$ (Theorem 2.38). It is also explained why a sieve with a fixed modulus cannot by itself do better at $p^2 - 1$ (Theorem 2.28).
- Exact p -adic valuations on divisibility progressions of Y : for the offsets $(p - 1)\mu$, including $\mu \geq p$, and for the odd offsets $(p - 1)\mu + 1$ (Theorems 2.39 and 2.40).

What is not new. Some statements re-derive results printed above in this part, or in a sibling manuscript. Each is printed once and the earlier result is credited:

- Theorem 2.39 at $\mu = 1$ is the valuation $v_p(T_{p-1}(Y)) = v_p(Y) - 1$ in the proof of Theorem 2.12, proved by a different method (a minimal-valuation multinomial count after Davis). Manuscript 17 says so. It is a second route to that case.
- The congruence $pT_{p-1}(Y)/Y \equiv 1$ that manuscript 12 recalls in proving Corollary 2.20 is (2.28).
- The offset-eight certificate displayed in the same proof is the row $d = 8$, $p = 11$ of Theorem 2.14. The manuscript’s polynomial $F(Y)$ is $R_8(Y + 8)$ of (2.29), with the same scalar $c_8 = 1/1393459200$; the intake checked this against `data/diagonal_certificates.json`.
- The cases $j = 1, 2$ of Lemma 2.27 re-derive the columns $r = 1, 2$ of Proposition 2.15.
- The derivative identity (2.50) is Proposition 2.4 combined with (2.22).
- Manuscript 09 re-proves three lemmas of manuscript 12: the positivity cone, the ordinary congruence and the prime-square congruence. They are printed once, as manuscript 12’s. Manuscript 09’s versions are recorded in Remarks 2.29, 2.31 and 2.37.
- Manuscript 09’s candidate list also gives a second route, not stated there, to Corollary 2.21 for $p \geq 5$ and to Corollary 2.22 for $p \geq 7$, $p \neq 11$ (Remark 2.34).
- For $p > \mu$, Theorem 2.39 follows from the congruence in Lemma 2.23 (Remark 2.42). Manuscript 17 is new for $\mu \geq p$ and on the shifted offsets.

2.11.1 Notation of the additions

The three manuscripts write the offset polynomial of Subsection 2.5 as $T_r(Y)$, with r the offset. In this part r is the *column* of $b(m, r)$, and the offset is $d = m - r$; the manuscripts also call a multiplier m , which is this part’s *row*. The additions are therefore printed with the letters of Table 2.A. The first three rows are where a reader will go wrong. No normalization was changed.

Three further conventions. The falling factorial $(Y)_{\underline{\ell}}$ of the manuscripts (17 writes $(Y)_{\ell}$) is $Y^{\underline{\ell}}$ here, as in the rest of the article. The manuscripts’ “factor defect r ” and “log degree k ” of a derivative monomial are this part’s exponents d and j in $x^{x-d}(\log x)^j$. And all three cite this report as “Part I” of the pinned source: they mean Part 2, Section 2 of this article. Their bibliography entry for that source is replaced below by references into this part, and manuscript 09’s entry for its “preceding companion” by references to Subsection 2.12.

The relation between the two Stirling conventions is $x^{\bar{\mu}} = x(x+1) \cdots (x+\mu-1) = \sum_j c(\mu, j)x^j$ and $F_{\mu}(y) = y^{\underline{\mu}} = \sum_{\ell} s(\mu, \ell)y^{\ell}$, so that $c(\mu, j) = (-1)^{\mu-j}s(\mu, j)$. The two uses do not overlap: (2.15) is an exact formula for $b(m, r)$, while $c(\mu, j)$ below is a residue modulo p .

2.12 Complete nonvanishing on prime-multiple offsets (manuscript 12)

Abstract of manuscript 12. *We prove that every positive-column entry on offset $2(p-1)$ of the Lehmer–Comtet triangle is nonzero, for every prime p , and likewise on offset $4(p-1)$ for every prime and $3(p-1)$ for every odd prime. More generally, for every fixed positive multiplier μ , the entire offset $\mu(p-1)$ is zero-free for all sufficiently large primes p , with an explicit harmonic-number condition and a fully integer sufficient threshold. A denominator argument confines every possible positive integer root to a small exponent or a residue class beyond p^2 . A classical derivative-root barrier proves the universal positivity cone $Y \geq 4d$ and excludes the latter region; an elementary harmonic-product estimate excludes the former. We also explain a limitation of a fixed finite-modulus sieve at prime-square offsets. The complete integer-zero conjecture remains open.*

Table 2.A: Letters of Subsections 2.11–2.15 against manuscripts 12, 09 and 17 and against the rest of Part 2. No normalization was changed.

Here	Manuscripts	Meaning	Tempting false reading
d	r (all three); n (in 12's and 09's prime-square proofs, 17's proofs)	offset $m - Y$; $T_d(Y) = [z^d]H(z)^Y$ as in (2.21)	false: the column r of $b(m, r)$
μ	m	multiplier: $d = \mu(p - 1)$, or $d = \mu(p - 1) + 1$	false: the row m of $b(m, r)$
m	M	row: $m = Y + d$, or $m = \mu(p - 1) + j$ in 09	false: the multiplier
Y	Y	column, $b(Y + d, Y)$; the r of $b(m, r)$	—
j	j (12, 09)	a small column value $Y = j < \mu$	the exponent j of $\log x$ in (2.1)
$H(z), h_j$	$\mathcal{H}(z)$ (12, 09), $H(z)$ (17); h_j , h_n	$\Phi(z)/z$ and its coefficients, (2.21)	the harmonic H_t
H_n	H_n	harmonic number $\sum_{i \leq n} 1/i$, the H_t of (2.30)	the series $H(z)$
$F_m(x)$	$P_M(x)$	falling factorial $\prod_{h=0}^{m-1} (x - h)$	the canonical polynomial P_n
$\Psi_P(\xi)$	$\Phi_P(b)$ (12), $\Phi(b)$ (09)	$P'(\xi)/P(\xi)$ at a real point ξ above the roots	$\Phi(z) = (1 + z) \log(1 + z)$; the coefficient $b(m, r)$
$c(\mu, j)$	$c(m, j)$ (09)	unsigned Stirling number of the first kind, $c(\mu, j) = (-1)^{\mu-j} s(\mu, j)$	the signed $s(m, \ell)$ of (2.15); the scalar c_d of (2.29)
ω	U	total multiplicity (12) or total weight (12, § 2.12.5) of bad indices	the bound $U(n)$ of Part 3
$\mathcal{V}$	D	total denominator valuation of a term	$D(z)$ of Subsection 2.8
Λ, ν	L, q (12)	$H_{d-1} - H_j$, and $j - 1$	$L = \log x$; the proposed count $q(n)$
$\Theta, \gamma_s, \beta_t, \mathcal{R}, \chi$	Q, c_s, b_t, E, x (12)	local objects of the proof of Lemma 2.27	$Q(z), c_d, b(m, r), E(n), x$
σ	c (12), s (09)	the ratio Y/d	—
w_i	x_i (12)	local variables in the proof of Lemma 2.24	the variable x of x^x
κ	k (12)	order of a derivative in Lemma 2.24	the exponent k of x^{-1}
λ, η, g_0	L, s, n_0 (09)	$\sum \ell_i, \sum \varepsilon_i$, the good index	$L = \log x$; $s(m, \ell)$; the order n
$\mathbf{e}_j$	E_j (17)	unit vector at position j	the defect $E(n)$
n, j	N, k (12, 09, 17)	derivative order, exponent of $\log x$	—

Editorial note. Printed in the letters of Table 2.A: the manuscript's multiplier m is μ and its offset r is d .

2.12.1 The coefficients and the main theorem

Write

$$H(z) = \frac{(1+z)\log(1+z)}{z} = 1 + \sum_{j \geq 1} h_j z^j, \quad h_j = \frac{(-1)^{j-1}}{j(j+1)}, \quad T_d(Y) = [z^d]H(z)^Y, \quad (2.41)$$

which is (2.21). For every fixed $d \geq 1$, $T_d \in \mathbb{Q}[Y]$ has degree d and the factor Y (Lemma 2.11). The Lehmer–Comtet coefficients are

$$b(m, Y) = \frac{m!}{Y!} [z^m]((1+z)\log(1+z))^Y = \frac{m!}{Y!} T_{m-Y}(Y), \quad 1 \leq Y \leq m, \quad (2.42)$$

which is (2.22). In particular, $T_d(Y) \neq 0$ for all positive integers Y is exactly nonvanishing of the entire offset $m - Y = d$. These coefficients enter the derivative expansion (2.12). Throughout, $H_n = \sum_{i=1}^n 1/i$, with $H_0 = 0$, denotes a harmonic number.

Theorem 2.19. *Let $\mu \geq 2$ be an integer, let $p > \mu$ be prime, and set $d = \mu(p-1)$. If*

$$p^2 + 1 \geq 4d, \quad (2.43)$$

$$H_{d-1} - H_{\mu-1} \geq 4(\mu-2), \quad (2.44)$$

then $T_d(Y) \neq 0$ for every positive integer Y .

For each fixed μ , both conditions hold for all sufficiently large p . The explicit integer condition

$$p \geq \max\{4\mu, 1 + 3^{4\mu-7}\} \quad (2.45)$$

is sufficient.

Corollary 2.20. *For every prime p and every positive integer Y ,*

$$T_{2(p-1)}(Y) \neq 0.$$

Equivalently, $b(m, m-2p+2) \neq 0$ for every $m \geq 2p-1$.

Corollary 2.21. *For every odd prime p and every positive integer Y ,*

$$T_{3(p-1)}(Y) \neq 0.$$

The exclusion of $p=2$ is necessary: $T_3(2) = T_3(5) = 0$.

Corollary 2.22. *For every prime p and every positive integer Y ,*

$$T_{4(p-1)}(Y) \neq 0.$$

For derivatives of x^x , the monomial $x^{x-d}(\log x)^j$ of $f^{(n)}$ — offset d , logarithmic degree j — has coefficient $\binom{n}{m} b(m, Y)$ with $m = n - j$ and $Y = n - d - j$ (Theorem 2.3). On every covered offset, the only physical cancellation is therefore the structural boundary $Y = 0$, or $j = n - d$; all cells with $Y > 0$ are nonzero. The covered offsets are even, so this conclusion does not remove the known odd-offset reflection family.

The inspected repository proves the prime-offset case $d = p - 1$ (Theorem 2.12) and completely checks offsets through 16 (Theorem 2.14). Corollaries 2.20–2.22 and Theorem 2.19 extend those source-relative conclusions to complete infinite families, including offsets whose successor is composite. No claim of exhaustive literature priority is intended.

2.12.2 Arithmetic localization to two or more digits

Let $\mathbb{Z}_{(p)}$ be the rationals with denominator coprime to p . Polynomial congruences in this subsection are over that ring.

Lemma 2.23. *Suppose $p > \mu \geq 2$ and $d = \mu(p-1)$. Then $G(Y) = p^\mu T_d(Y)$ and $G(Y)/Y$ lie in $\mathbb{Z}_{(p)}[Y]$, and*

$$G(Y) \equiv \frac{Y^\mu}{\mu!} \pmod{p}. \quad (2.46)$$

For $1 \leq j \leq \mu-1$,

$$v_p(G(j)) \geq \mu - j + 1. \quad (2.47)$$

Every positive integer zero of T_d therefore satisfies

$$Y \equiv j \pmod{p^{\mu-j+1}} \quad \text{for some } 1 \leq j \leq \mu-1. \quad (2.48)$$

In particular, it either lies in $\{1, \dots, \mu-1\}$ or is at least $p^2 + 1$.

Proof. The falling-factorial expansion is

$$T_d(Y) = \sum_{\sum j i_j = d} Y^\ell \prod_{j \geq 1} \frac{h_j^{i_j}}{i_j!}, \quad \ell = \sum i_j. \quad (2.49)$$

Since $d < p^2 - 1$, each h_j occurring here has valuation 0 or -1 . Call an index bad when its valuation is -1 , and let ω be the total multiplicity of bad indices. Every bad index is at least $p-1$, so $\omega \leq \mu$. Equality forces the unique partition $i_{p-1} = \mu$; its total denominator valuation is μ , because $\mu!$ is a unit.

If $\omega < \mu$, every bad-index multiplicity is less than p . The total multiplicity of good indices is at most $(\mu - \omega)(p-1)$. Since all multiplicities are below p^2 , their factorial valuations sum to at most

$$\left\lfloor \frac{(\mu - \omega)(p-1)}{p} \right\rfloor = \mu - \omega - 1.$$

Thus every nondistinguished term in (2.49) has denominator valuation at most $\mu-1$. Only $i_{p-1} = \mu$ survives after multiplication by p^μ and reduction. Since $ph_{p-1} \equiv 1 \pmod{p}$, this proves (2.46). Every falling factorial has a factor Y , so division by that exact factor also preserves p -integrality.

To prove (2.47), expand H^j as a product of j series. A term could have denominator valuation j only if all j chosen indices were positive and bad. Each such index is 0 or -1 modulo p . Their sum cannot equal $d \equiv -\mu \pmod{p}$, since $j < \mu < p$. Hence the denominator valuation is at most $j-1$.

The reduction of G/Y has simple roots precisely $1, \dots, \mu-1$ modulo p . An integer zero must first lie in one of these classes. At its representative j , $G'(j)$ is a unit. In the Taylor expansion at j , if $1 \leq v_p(Y-j) < \mu-j+1$, the linear term has strictly smaller valuation than the constant term and all higher terms. It cannot cancel. This proves (2.48) and the asserted localization. $\square$

The denominator comparison uses the classical multinomial method also appearing in Davis's analysis of powers of $\log(1+z)/z$ [15]; here the additional factor $1+z$ produces the denominators $j(j+1)$.

2.12.3 A derivative-root barrier excludes large exponents

We reproduce the needed classical barrier estimate. A primary source is Ravichandran's Theorem 1.4, attributed there to Marcus [16].

Lemma 2.24. *Let P have only real roots, let $\xi > \max \text{root } P$, and put $\Psi_P(\xi) = P'(\xi)/P(\xi)$. For $0 \leq \kappa < \deg P$,*

$$\max \text{root } P^{(\kappa)} < \xi - \frac{\kappa}{\Psi_P(\xi)}.$$

Proof. Put $\delta = 1/\Psi_P(\xi)$. For degree at least two, let $u_i = \delta/(\xi - \lambda_i)$, where the λ_i are the roots with multiplicity. Then $0 < u_i < 1$ and $\sum u_i = 1$, so $\xi - \delta$ remains above every root of P . With

$$w_i = \frac{u_i}{1 - u_i}, \quad A = \sum w_i, \quad B = \sum w_i^2,$$

logarithmic differentiation gives

$$\Psi_{P'}(\xi - \delta) = \delta^{-1}(A - B/A).$$

Regard the u_i as probability weights. Then

$$\mathbb{E}[w] = A - 1, \quad \mathbb{E}[1 + w] = A, \quad \mathbb{E}[w(1 + w)] = B,$$

so $B - A(A - 1) = \text{Var}(w) \geq 0$. Hence $\Psi_{P'}(\xi - \delta) \leq 1/\delta$. A smaller displacement gives the same upper bound by monotonicity of the logarithmic derivative above the roots. Iterate using this fixed δ ; every shifted point stays strictly above the corresponding derivative's roots. $\square$

Corollary 2.25 (A universal positivity cone). *For positive integers Y, d , if $Y \geq 4d$, then $T_d(Y) > 0$. Consequently every positive integer zero satisfies $Y < 4d$.*

Proof. Set $m = Y + d$ and $F_m(x) = \prod_{h=0}^{m-1} (x - h)$. Differentiating $(1+z)^x = \sum_{m \geq 0} F_m(x) z^m / m!$ exactly Y times in x and setting $x = Y$ gives

$$T_d(Y) = \frac{F_m^{(Y)}(Y)}{m!}. \quad (2.50)$$

Choose $\xi = m + d = Y + 2d$. Its barrier is

$$\Psi_{F_m}(\xi) = H_{Y+2d} - H_d < \int_d^{Y+2d} \frac{dt}{t} = \log \frac{Y+2d}{d}.$$

Write $\sigma = Y/d \geq 4$. The function $\sigma - 2 \log(\sigma + 2)$ increases for $\sigma \geq 4$ and is positive there, since $\log 6 < 2$. Thus $Y > 2d \Psi_{F_m}(\xi)$. Lemma 2.24 yields $\max \text{root } F_m^{(Y)} < \xi - Y/\Psi_{F_m}(\xi) < Y$. The derivative has positive leading coefficient, so its value at Y is strictly positive. $\square$

Editorial note. Identity (2.50) is Proposition 2.4 combined with (2.22), derived there by the same differentiation of the binomial series.

Condition (2.43) therefore excludes every possible zero with $Y \geq p^2 + 1$. This uses real-rootedness only of the falling-factorial polynomial F_m , not of T_d as a polynomial in Y .

Remark 2.26 (Editorial: the cone and the asymptotic sign boundary). In the ratio $\rho = m/Y$ of row to column, the cone of Corollary 2.25 is $\rho \leq 5/4$, and it holds on every offset, bounded ones included. Manuscript 31 of the same batch, placed in the new sibling report **power-tower-exponent-supports**, proves positivity of $b(m, Y)$ eventually on compact ranges of ρ inside $(1, \rho_*)$, $\rho_* \approx 1.29843$, without an effective threshold. Neither statement implies the other. The cone is valid but not sharp: in an exact scan of all rows $m \leq 200$ at the intake, no entry in the cone is nonpositive, and the smallest ratio m/Y of a nonpositive entry is $196/148 \approx 1.324$.

2.12.4 Small exponents and completion of the proof

Lemma 2.27. *Let $d, j \geq 1$, put $\nu = j - 1$ and $\Lambda = H_{d-1} - H_j$. If $j = 1$, then $T_d(j) \neq 0$. If $\nu \geq 1$ and $\Lambda \geq 4\nu$, then $T_d(j) \neq 0$, with sign $(-1)^{d+j}$.*

Proof. The case $j = 1$ follows from $T_d(1) = h_d \neq 0$. Otherwise

$$F_{d+j}(j+u) = (-1)^{d-1} j!(d-1)! u \Theta(u), \quad \Theta(u) = \prod_{h=1}^j (1+u/h) \prod_{h=1}^{d-1} (1-u/h).$$

Its logarithm is $-\Lambda u + \sum_{s \geq 2} \gamma_s u^s$, with $|\gamma_s| \leq 2$. Let $\beta_t = [u^t] \exp(2u^2/(1-u))$. Coefficientwise majorization gives

$$[u^\nu] \Theta(u) = \frac{(-\Lambda)^\nu}{\nu!} + \mathcal{R}, \quad \frac{|\mathcal{R}|}{\Lambda^\nu/\nu!} \leq \sum_{t=2}^{\nu} \beta_t \left(\frac{\nu}{\Lambda} \right)^t.$$

For $\chi = \nu/\Lambda \leq 1/4$, the right side is at most $\exp(2\chi^2/(1-\chi)) - 1 \leq e^{1/6} - 1 < 1$. Thus $[u^\nu] \Theta$ has sign $(-1)^\nu$. The derivative identity (2.50) proves the assertion. $\square$

Editorial note. At $j = 1$ and $j = 2$ the lemma re-derives, on every offset, the column formulas of Proposition 2.15: $T_d(1) = h_d$ is $b(m, 1) = (-1)^m(m-2)!$, and the sign of $[u^1] \Theta = -(H_{d-1} - H_2)$ is that of $2H_{m-3} - 3$ in (2.30). For larger j it has no counterpart in this part: every fixed column $Y = j$ has $T_d(j) \neq 0$ as soon as $H_{d-1} - H_j \geq 4(j-1)$, so each column has only finitely many possible zeros, below an explicit offset (an observation of the intake, not stated in the manuscript).

Proof of Theorem 2.19. Arithmetic localization leaves only $1 \leq Y \leq \mu - 1$ and $Y \geq p^2 + 1$. The latter is excluded by (2.43) and Corollary 2.25. For $1 \leq j \leq \mu - 1$, condition (2.44) gives

$$H_{d-1} - H_j \geq 4(\mu - 2) \geq 4(j - 1),$$

so Lemma 2.27 excludes all remaining exponents.

For the explicit threshold, $p \geq 4\mu$ gives $p^2 + 1 > 4\mu(p-1)$, proving (2.43). Also

$$H_{d-1} - H_{\mu-1} > \log d - 1 - \log(\mu - 1) > \log(p - 1) - 1.$$

If $p - 1 \geq 3^{4\mu-7}$, then $\log 3 > 1$ makes this greater than $4(\mu - 2)$. Thus (2.44) holds as well. $\square$

Proof of Corollary 2.20. Take $\mu = 2$. Condition (2.44) holds for every $p > 2$, and $p^2 + 1 \geq 8(p-1)$ for every $p \geq 7$. Theorem 2.19 handles those primes.

The cases $p = 2, 3$ have offsets 2, 4, already prime offsets for the primes 3, 5. For completeness, that known theorem follows from $pT_{p-1}(Y)/Y \equiv 1 \pmod{p}$: only the one-part contribution h_{p-1} survives after division by Y and multiplication by p . (Editorial: this is (2.28) in the proof of Theorem 2.12.)

For $p = 5$, a primitive numerator of $T_8(Y)/Y$ is

$$\begin{aligned} F(Y) = & 135Y^7 - 8820Y^6 + 229950Y^5 - 3075240Y^4 \\ & + 22538215Y^3 - 89214420Y^2 + 172998164Y - 122821584. \end{aligned}$$

Its values modulo 11 at $Y = 0, \dots, 10$ are 5, 9, 6, 2, 2, 5, 5, 4, 10, 4, 3, all nonzero. This restates one of the source's small-offset certificates solely to close the uniform all-prime assertion. Finally $Y = 1$ is always explicitly nonzero by (2.41). $\square$

Editorial note. The source certificate restated here is the row $d = 8$ of Theorem 2.14: $F(Y) = R_8(Y + 8)$, with R_8 the primitive residual of (2.29) recorded in `data/diagonal_certificates.json`, and $T_8(Y) = c_8 Y R_8(Y + 8)$ with $c_8 = 1/1393459200$. The residues listed above are those of R_8 at $X = 8, \dots, 18$, the same eleven classes as the residues recorded for $X = 0, \dots, 10$. The intake checked the identity $F(Y) = R_8(Y + 8)$ coefficient by coefficient.

Proof of Corollary 2.21. Take $\mu = 3$. If $p \geq 11$, then $p^2 + 1 \geq 12(p - 1) = 4d$. The small sector in Lemma 2.23 contains only $j = 1, 2$. For $j = 2$, the coefficient in Lemma 2.27 is exactly $-(H_{d-1} - H_2)$, which is nonzero for $d > 3$. Thus the stronger condition (2.44) is unnecessary here. The remaining odd primes 3, 5, 7 give offsets 6, 12, 18, whose successors 7, 13, 19 are prime. The known prime-offset theorem covers these cases. $\square$

Proof of Corollary 2.22. Take $\mu = 4$. If $p \geq 17$, then $p^2 + 1 \geq 16(p - 1) = 4d$ and $d \geq 64$. The cases $j = 1, 2$ are as above. For $j = 3$, the relevant coefficient is

$$[u^2]\Theta(u) = \Lambda^2/2 + \gamma_2, \quad \Lambda = H_{d-1} - H_3, \quad |\gamma_2| < 2.$$

Here $\Lambda \geq H_{63} - H_3 > \int_4^{64} dt/t = \log 16 > 2$, so this coefficient is strictly positive. Arithmetic localization and the positivity cone exclude all other Y .

The remaining primes 2, 3, 5, 7, 11, 13 give offsets 4, 8, 16, 24, 40, 48. They are covered respectively by the prime-offset theorem at 5, the displayed offset-eight certificate, the prime-offset theorem at 17, Corollary 2.20 at 13, the prime-offset theorem at 41, and Corollary 2.21 at 17. This completes the proof. $\square$

2.12.5 Why prime-square offsets need additional input

The following obstruction concerns a fixed finite-modulus sieve, not every possible p -adic argument.

Theorem 2.28. *For every prime p , put $d = p^2 - 1$. Then*

$$J_p(Y) = p^{p+2} \frac{T_d(Y)}{Y} \in \mathbb{Z}_{(p)}[Y], \quad J_p(Y) \equiv -(Y^p - Y) \pmod{p}.$$

Consequently J_p has exactly p simple roots in $\mathbb{Z}_p$, one in each residue class modulo p .

Proof. In (2.49), let $v_j = v_p(j(j+1))$, $\omega = \sum i_j v_j$ and $\mathcal{V} = \omega + \sum v_p(i_j!)$. We use

$$j \geq (p-1)v_j, \quad j \geq 1 + (p-2)v_j,$$

with equality in the first only at $j = p - 1$. For $v_j > 0$, these follow from $j \geq p^{v_j} - 1$; the case $v_j = 0$ is immediate. Set $\mu = p + 1$. Then $\omega \leq \mu$, and equality forces $i_{p-1} = \mu$, whose denominator valuation is $p + 2$.

For every other partition put $\delta = \mu - \omega \geq 1$. The second inequality gives $\ell \leq p + 1 + (p - 2)\delta$. Since each multiplicity is below p^2 ,

$$\sum v_p(i_j!) \leq \lfloor \ell/p \rfloor \leq \delta + 1 - \left\lceil \frac{2\delta - 1}{p} \right\rceil \leq \delta.$$

Thus $\mathcal{V} \leq p + 1$ for all nondistinguished terms. Only the distinguished term survives after scaling by p^{p+2} . Wilson's congruence gives its coefficient -1 modulo p (also for $p = 2$). Dividing by the exact factor Y leaves $\prod_{j=1}^p (Y - j) = Y^p - Y$ modulo p .

The derivative of the reduced polynomial is 1. Each residue class is a root modulo p and has exactly one lift at every higher power, since the linearized congruence determines the next digit uniquely. These compatible residues are exactly the asserted simple roots in $\mathbb{Z}_p$. $\square$

Thus every fixed modulus p^e retains positive integer candidates in all p compatible classes. This does not exclude a more refined argument showing that the particular p -adic roots are not ordinary integers. For odd p , the offset is even and has no known reflection factor to remove. The complete prime-power-offset classification remains open.

Editorial note. Theorem 2.38 below (manuscript 09) is such a refined argument for $p \geq 5$: it locates the p -adic roots to three digits and, with the cone, reduces the offset $p^2 - 1$ to $1 \leq Y \leq p - 1$. The classification of that offset stays open.

2.12.6 Verification and source scope

The accompanying standard-library checker (`code/05-prime-multiple-verify.py`) constructs T_d exactly through degree 72, verifies the coefficient congruences and 64 stronger lift inequalities, and compares 3,132 coefficients against an independent integer-triangle recurrence. It checks 14,000 instances of the three complete prime families, 15,880 positivity-cone instances, 29,456 localization instances through triangle row 400, the prime-square congruence for $p = 2, 3, 5, 7$, the full offset-eight identity and its modular certificate, and the main conditions for $(\mu, p) = (2, 11), (3, 83), (3, 89), (4, 3001)$. A second exact convolution replay (`code/05-prime-multiple-replay_convolution.py`) independently checks 1,320 double-family instances, 3,000 localization instances, 56 lifts and 1,176 derivative-bridge identities; finite-difference interpolation also recovers the offset-eight certificate. These finite replays support reproducibility; the preceding proofs establish the infinite statements.

The relevant default-branch source was rechecked on 1 October 2026 and still had blob 7344f3aa4dc9344fb8648326e74e0d4bcc33eada, matching the pinned source (this report at commit ca81647a9). It already contains the prime-offset theorem and the finite offset-16 classification. The global exact zero list remains unsettled. No real-rootedness assertion about all T_d is used or proved here.

2.13 Stirling residues and finite candidate lists (manuscript 09)

Abstract of manuscript 09. *We identify a scaled block of the Lehmer–Comtet triangle modulo a prime with an ordinary unsigned Stirling row. The formula determines the first nontrivial digits of the associated p -adic roots. Combined with a derivative-root bound, it confines every positive integer zero on offset $\mu(p - 1)$, for $p > \mu$, to an explicit set of small exponents and at most one larger integer. It proves complete nonvanishing on offset $5(p - 1)$ for every odd prime. Two different square families have particularly short remaining lists: offset $(p - 1)^2$ has at most one candidate for odd primes, while offset $p^2 - 1$ has candidates only in $1, \dots, p - 1$ for $p \geq 5$. These are necessary conditions, not assertions that the listed candidates are zeros. The global exact zero classification remains open.*

Editorial note. Printed in the letters of Table 2.A. The manuscript’s “preceding companion” is manuscript 12, printed in Subsection 2.12; manuscript 09 was delivered with an unchanged copy of it, which is not shipped a second time.

2.13.1 Coefficients, scope, and the root bound

The manuscript defines $H(z)$, h_j , $T_d(Y)$ and $b(Y + d, Y)$ as in (2.41) and (2.42), and notes that T_d has the structural factor Y . Write $c(\mu, j)$ for the unsigned Stirling number of the first kind, defined by

$$X(X + 1) \cdots (X + \mu - 1) = \sum_{j=1}^{\mu} c(\mu, j) X^j.$$

All congruences of rational numbers or polynomials are over $\mathbb{Z}_{(p)}$, the rationals with denominator coprime to p .

Remark 2.29 (Manuscript 09’s restatement of the cone). The manuscript opens with the positivity bound of Corollary 2.25 — for integers $d, Y \geq 1$, $Y \geq 4d$ implies $T_d(Y) > 0$ — taken from manuscript 12 “with its brief derivation included”. That derivation is the proof of Corollary 2.25 in condensed form: it applies the classical barrier [16, Theorem 1.4] to $F_m^{(Y)}$, $m = Y + d$, with $\Psi(\xi) = \sum_{h=0}^{m-1} 1/(\xi - h)$ at $\xi = Y + 2d$, bounds $\Psi(\xi) = H_{Y+2d} - H_d < \log((Y + 2d)/d)$, and uses $\sigma > 2 \log(\sigma + 2)$ for $\sigma = Y/d \geq 4$. It is the same argument, quoting the barrier rather than proving it, and is not printed a second time.

This note concerns positive Y . In derivatives of x^x , the offset d and log degree j of a monomial $x^{x-d}(\log x)^j$ satisfy $Y = n - d - j$; the structural line $Y = 0$, or $j = n - d$, remains a cancellation line. The inspected source gives the prime-offset theorem and fixed-offset certificates through 16 (Theorems 2.12 and 2.14). The preceding companion (Subsection 2.12) proves complete families with multipliers 2, 3, 4 and a general sufficiently-large-prime theorem. Here the additional input is an exact Stirling residue and its root displacement. The partition method belongs to the classical p -adic Bernoulli/Stirling framework [15, 17]; the results below are proved directly, without an exhaustive priority claim.

2.13.2 The Stirling residue

Theorem 2.30. *Let $p > \mu \geq 2$ be prime and integer, respectively, and let $1 \leq j < \mu$. For $d = \mu(p - 1)$,*

$$p^{j-1}T_d(j) \equiv -\frac{(j!)^2(\mu - j - 1)!}{\mu!(\mu - 1)!}c(\mu, j) \pmod{p}. \quad (2.51)$$

The left side is p -integral. If $p \nmid c(\mu, j)$, then $v_p(T_d(j)) = 1 - j$, in particular $T_d(j) \neq 0$. Equivalently, the triangle entries satisfy

$$\frac{b(\mu(p - 1) + j, j)}{p^{\mu-j}} \equiv (-1)^j \frac{j!}{\mu!}c(\mu, j) \pmod{p}. \quad (2.52)$$

Proof. Since $d < p^2 - 1$, each nonconstant coefficient h_i with $i \leq d$ has valuation 0 or -1 . Call an index bad in the latter case. In an ordered-product expansion of H^j , all j indices cannot be bad: their residues are 0 or -1 modulo p , whereas their sum is $d \equiv -\mu$, and $j < \mu < p$. Hence $p^{j-1}T_d(j)$ is integral at p .

Only terms with exactly $j - 1$ bad indices survive. The remaining index is positive and good; a constant there would again make the residue sum impossible. Choose its position in j ways, and write the bad indices as

$$\ell_i p - \varepsilon_i, \quad \ell_i \geq 1, \quad \varepsilon_i \in \{0, 1\}, \quad \lambda = \sum \ell_i, \quad \eta = \sum \varepsilon_i.$$

The good index is $g_0 = (\mu - \lambda)p - \mu + \eta$, positive exactly when $\lambda \leq \mu - 1$. Both g_0 and $g_0 + 1$ are units, since $\eta \leq j - 1 \leq \mu - 2$. Because p is odd,

$$ph_{\ell_i p - \varepsilon_i} \equiv \frac{(-1)^{\ell_i+1}}{\ell_i}, \quad h_{g_0} \equiv \frac{(-1)^{\eta-\lambda-1}}{(\mu - \eta)(\mu - \eta - 1)} \pmod{p}.$$

The product sign is $(-1)^{\eta+j-2}$, independently of λ . Therefore the surviving residue is

$$j(-1)^{j-2}A_{\mu,j}B_{\mu,j}, \quad (2.53)$$

where

$$A_{\mu,j} = \sum_{\substack{\ell_1, \dots, \ell_{j-1} \geq 1 \\ \ell_1 + \dots + \ell_{j-1} \leq \mu-1}} \frac{1}{\ell_1 \cdots \ell_{j-1}},$$

$$B_{\mu,j} = \sum_{\eta=0}^{j-1} \frac{(-1)^\eta \binom{j-1}{\eta}}{(\mu-\eta)(\mu-\eta-1)}.$$

For $j = 1$, the empty tuple gives $A_{\mu,1} = 1$. The standard logarithmic generating function for unsigned Stirling numbers gives

$$A_{\mu,j} = [t^{\mu-1}] \frac{(-\log(1-t))^{j-1}}{1-t} = \frac{(j-1)!}{(\mu-1)!} c(\mu, j).$$

A finite binomial sum and a beta integral give

$$B_{\mu,j} = \int_0^1 x^{\mu-2} (1-x)(1-x^{-1})^{j-1} dx$$

$$= (-1)^{j-1} \frac{(\mu-j-1)! j!}{\mu!}.$$

The integral converges because $j < \mu$. Substitution into (2.53) proves (2.51), including the sign.

Finally put $m = \mu(p-1) + j$. Then $v_p(m!) = \mu - 1$. Splitting into p -blocks and applying Wilson's congruence gives

$$\frac{m!}{p^{\mu-1}} \equiv (-1)^{j+1} \frac{(\mu-1)!}{(\mu-j-1)!} \pmod{p}.$$

Combining this with (2.42) proves (2.52). $\square$

2.13.3 Root displacements and the complete candidate list

Remark 2.31 (Manuscript 09's ordinary congruence). The manuscript's Lemma 3.1 is the congruence half of Lemma 2.23: for $p > \mu \geq 2$ and $d = \mu(p-1)$, $G(Y) = p^\mu T_d(Y) \in \mathbb{Z}_{(p)}[Y]$, $G(Y) \equiv Y^\mu/\mu! \pmod{p}$, and G/Y is also p -integral. Its proof, which the manuscript calls “the next elementary denominator observation”, is a condensed version of the first two paragraphs of the proof of Lemma 2.23 (at most μ bad factors, equality only at $i_{p-1} = \mu$, factorial valuations of the other terms at most $\mu - \omega - 1$) and is not printed a second time.

Theorem 2.32. *For every $j = 1, \dots, \mu-1$, the polynomial G/Y has one simple root $\rho_j \in \mathbb{Z}_p$ in class j , and*

$$\rho_j \equiv j + (-1)^{\mu-1-j} \frac{j! c(\mu, j)}{(\mu-1)!} p^{\mu-j+1} \pmod{p^{\mu-j+2}}. \quad (2.54)$$

There are no other roots in $\mathbb{Z}_p$.

Proof. Congruence (2.46) gives precisely the simple residue roots $1, \dots, \mu-1$, each with a unique lift at every successive power of p . Its derivative before division by Y is

$$G'(j) \equiv \frac{j! (-1)^{\mu-1-j} (\mu-1-j)!}{\mu!} \pmod{p}.$$

Theorem 2.30 determines $G(j)/p^{\mu-j+1}$ modulo p . Taylor expansion at j therefore gives (2.54). The quadratic remainder has valuation at least $2(\mu-j+1) \geq \mu-j+2$. The formula remains valid if $p \mid c(\mu, j)$; it then asserts an additional zero digit. $\square$

Theorem 2.33 (Finite candidate list). *Let $p > \mu \geq 2$, $d = \mu(p-1)$, and let $C_p \in \{1, \dots, p-1\}$ be the least nonnegative residue of $\binom{\mu}{2}$ modulo p . Every positive integer zero of T_d belongs to*

$$\{j : 1 \leq j < \mu, p \mid c(\mu, j)\} \cup \{\mu - 1 + C_p p^2\}, \quad (2.55)$$

where the last candidate may be omitted when it is at least $4d$.

Proof. Corollary 2.25 confines zeros to $0 < Y < 4d$. Since $\mu \leq p-1$ and $p \geq 3$,

$$p^3 > 4(p-1)^2 \geq 4\mu(p-1) = 4d.$$

For $j \leq \mu-2$, formula (2.54) fixes the root at j modulo at least p^3 . Its only possible positive integer representative below $4d$ is j itself, excluded by Theorem 2.30 unless $p \mid c(\mu, j)$.

For $j = \mu-1$, the displacement simplifies to

$$\rho_{\mu-1} \equiv \mu - 1 + \binom{\mu}{2} p^2 \pmod{p^3}.$$

Here $p \nmid \binom{\mu}{2}$, since $p > \mu$. There is at most one representative in $0 < Y < 4d$, namely the last candidate in (2.55). The small representative $\mu-1$ is not a root, again by Theorem 2.30. $\square$

Thus a sufficient complete-offset criterion is: no entry $c(\mu, j)$, $1 \leq j < \mu$, is divisible by p , and $\mu-1 + C_p p^2 \geq 4\mu(p-1)$. The candidates in (2.55) are necessary possibilities, not a claimed zero list.

Remark 2.34 (Editorial: a second route to Corollaries 2.21 and 2.22). Manuscript 09 does not say so, but its criterion re-proves most of manuscript 12's small-multiplier corollaries without the harmonic condition (2.44). For $\mu = 3$, the row is $c(3, 1) = 2$, $c(3, 2) = 3$, prime to every $p \geq 5$, and $C_p = 3$; the candidate $2 + 3p^2$ exceeds $12(p-1)$ for every p , so Theorem 2.33 gives Corollary 2.21 for every $p \geq 5$. For $\mu = 4$, the row is 6, 11, 6 and $C_p = 6 \bmod p$: at $p = 5$ the candidate $3 + 25 = 28$ lies below $4d = 64$ and survives; at $p = 11$ the entry $c(4, 2) = 11$ leaves the candidate $Y = 2$; for $p = 7$ and every $p \geq 13$ the list is empty, so the criterion gives Corollary 2.22 for $p \geq 7$, $p \neq 11$. For $\mu = 2$ the candidate $1 + p^2$ is at least $8(p-1)$ exactly when $p \geq 7$, which is condition (2.43). The exceptional primes are covered, in manuscript 12, by the prime-offset theorem and the offset-eight certificate.

2.13.4 Two consequences of the Stirling row

Corollary 2.35. *For every odd prime p and every positive integer Y ,*

$$T_{5(p-1)}(Y) \neq 0.$$

Proof. The unsigned Stirling row for $\mu = 5$, omitting its last entry 1, is 24, 50, 35, 10. Its only prime divisor above 5 is 7. For every prime $p \geq 11$, the small set in (2.55) is empty, $C_p = 10$, and $4 + 10p^2 > 20(p-1)$. Theorem 2.33 applies. The primes 3, 5, 7 give offsets 10, 20, 30, covered respectively by the prime-offset theorem at 11, the double family at 11 (Corollary 2.20), and the prime-offset theorem at 31. The prime 2 is correctly excluded: offset 5 has its known reflection zero at $Y = 4$. $\square$

Editorial note. The reflection zero at offset 5, $Y = 4$ is $b(9, 4) = 0$, the case $h = 2$ of Proposition 2.8.

Corollary 2.36. *For every odd prime p ,*

$$T_{(p-1)^2}(Y) = 0, \quad Y \in \mathbb{Z}_{>0} \implies Y = p^2 + p - 2.$$

No assertion is made that this one remaining value vanishes.

Proof. Take $\mu = p - 1$. Over $\mathbb{F}_p$,

$$\sum_{j=1}^{p-1} c(p-1, j) X^j = X(X+1) \cdots (X+p-2) = \frac{X^p - X}{X - 1} = X + \cdots + X^{p-1}.$$

Thus every relevant Stirling entry is a unit and $C_p = \binom{p-1}{2} \bmod p = 1$. Theorem 2.33 leaves only $p - 2 + p^2$. $\square$

2.13.5 A different square family: offset $p^2 - 1$

This subsection concerns $p^2 - 1$, not the offset $(p - 1)^2$ in Corollary 2.36.

Remark 2.37 (Manuscript 09's prime-square lemma). The manuscript's Lemma 5.1 is the congruence of Theorem 2.28: for any prime p and $d = p^2 - 1$, $J(Y) = p^{p+2} T_d(Y)/Y \in \mathbb{Z}_{(p)}[Y]$ and $J(Y) \equiv -(Y^p - Y) \pmod{p}$; its J is J_p . The manuscript says so ("This is the prime-square denominator lemma from" the companion) and repeats the estimates of the proof of Theorem 2.28 — the two inequalities for j in terms of $v_j = v_p(j(j+1))$, the bound $\omega \leq p + 1$ with equality only at $i_{p-1} = p + 1$, and $\sum v_p(i_j!) \leq \delta$ for the other terms — which are not printed a second time.

Theorem 2.38. *For every prime $p \geq 5$,*

$$T_{p^2-1}(Y) = 0, \quad Y \in \mathbb{Z}_{>0} \implies 1 \leq Y \leq p - 1.$$

Proof. Put $d = p^2 - 1$. Theorem 2.28 gives one simple p -adic root in each residue class. For $1 \leq a \leq p - 1$, expand H^a directly. Every index at most $d = p^2 - 1$ has denominator valuation at most one, except d itself, whose valuation is two. If that index occurs, all other chosen indices are zero. Hence

$$v_p(T_d(a)) \geq -\max(a, 2), \quad v_p(J_p(a)) \geq p + 2 - \max(a, 2) \geq 3.$$

The corresponding root is therefore a modulo p^3 .

To determine the zero-residue branch, expand H^p . Every degree- d term has a multinomial coefficient divisible by p : a pure p th-power term would require $pi = d$, which is impossible. A term containing index d has valuation -1 , greater than $1 - p$. Otherwise all bad indices have valuation -1 . Terms with at most $p - 1$ bad factors have valuation at least $2 - p$.

If all p factors are bad, their minimum total degree is $p(p - 1)$, leaving increment $p - 1$. A bad index greater than p is at least $2p - 1$, already requiring increment p . Thus the unique all-bad pattern has one index $p - 1$ and $p - 1$ indices p . Its contribution is $ph_{p-1}h_p^{p-1}$, so

$$p^{p-1} T_d(p) \equiv 1 \pmod{p}, \quad J_p(p) \equiv p^2 \pmod{p^3}.$$

Since $J'_p(p) \equiv 1$, the zero-residue branch equals $p - p^2$ modulo p^3 .

By Corollary 2.25, every positive integer zero has $Y < 4(p^2 - 1)$. For $p \geq 5$, both p^3 and $p^3 - p^2 + p$ exceed this bound. The only possible representatives of the identified branches in that interval are $1, \dots, p - 1$. $\square$

The remaining small-column assertions are not proved here. In particular, the existence of p -adic roots is fully compatible with the absence of ordinary integer roots; the archimedean bound is essential to both square-family reductions.

2.13.6 Verification and remaining scope

The standard-library checker (`code/04-stirling-residues-verify_stirling.py`) independently expands rational powers of H and verifies 134 instances each of the Stirling residue, the scaled triangle congruence, and the derivative/root-displacement formula, covering multipliers through 8 and coefficient degrees through 144. Separate exact computations lift the prime-square roots for $p = 3, 5, 7, 11$ and check the single-candidate values for $(p-1)^2$ at $p = 3, 5, 7, 11, 13, 17, 19$. Those finitely many values are nonzero; they do not establish a general nonvanishing theorem for the remaining candidate. A separate integer falling-product replay verifies 162 instances each of the two residues, derivatives and first displacements, all prime-square branches through $p = 19$, and the seven stated single candidates. It uses finite p -steps for derivatives and passes with Python assertions disabled.

The repository comparison is relative to the pinned source, whose relevant default-branch blob was unchanged at the check on 1 October 2026. Our bounded literature review found the classical partition, higher-order Bernoulli, and Stirling valuation framework [15, 17]; the formulas in this note are proved directly for the Lehmer–Comtet specialization. No worldwide priority claim, complete global zero list, or formal proof-assistant verification is asserted.

2.14 Uniform p -adic valuations (manuscript 17)

Abstract of manuscript 17. *We give exact valuations for two infinite families of coefficients of $((1+z)\log(1+z)/z)^Y$. The first extends the inspected source’s prime-offset theorem to offsets $(p-1)\mu$ when $\mu \leq p^{v_p(Y)}$. A second treats offsets $(p-1)\mu + 1$ for odd p under an explicit nonresonance condition. The proof adapts Davis’s minimal-valuation multinomial method directly to the different coefficient denominators $j(j+1)$, including a lower bound that handles competitors with many factors. The resulting criteria exclude new infinite families of possible Lehmer–Comtet zeros; the complete zero classification remains open.*

Editorial note. Printed in the letters of Table 2.A: the manuscript’s m is μ , its offset r (and n in its proofs) is d .

2.14.1 The coefficient and its derivative interpretation

The manuscript writes $H(z)$, h_j and $T_d(Y)$ as in (2.41). Throughout this subsection Y is a *positive integer*; the structural case $Y = 0$ is excluded. We write v_p for the valuation on nonzero rational numbers, so negative values are allowed.

The first-kind Lehmer–Comtet numbers satisfy $b(Y+d, Y) = \frac{(Y+d)!}{Y!} T_d(Y)$, which is (2.42). They determine the expanded derivative coefficients of x^x : at total derivative order $n \geq m = Y + d$, the monomial

$$x^{x-d}(\log x)^{n-m}$$

has coefficient $\binom{n}{m} b(m, Y)$. Thus nonvanishing of a fixed $b(m, Y)$ proves nonvanishing on its entire persistent derivative line. These identities are proved in the inspected source (Theorem 2.3 and Lemma 2.11) and belong to the classical Lehmer–Comtet framework.

Theorem 2.39 (Principal progression). *Let p be prime, $\mu \geq 1$ and $e = v_p(Y)$. If $\mu \leq p^e$, then*

$$v_p(T_{(p-1)\mu}(Y)) = e - v_p(\mu) - \mu. \quad (2.56)$$

In particular the coefficient is nonzero.

Theorem 2.40 (Shifted progression). *Let p be an odd prime, $\mu \geq 1$ and $e = v_p(Y)$. If*

$$\mu < p^e, \quad p \nmid (p-1)\mu + 1, \quad (2.57)$$

then

$$v_p(T_{(p-1)\mu+1}(Y)) = e - \mu. \quad (2.58)$$

The second condition in (2.57) is equivalently $\mu \not\equiv 1 \pmod{p}$.

Davis [15, Theorem 1.1] proves the valuation analogous to (2.56) for powers of $\log(1+z)/z$. Multiplication by $(1+z)^Y$ cannot simply be ignored. We instead prove both claims directly from (2.41), adapting the same minimal-multinomial mechanism. The $\mu = 1$ case of Theorem 2.39 is exactly the source's prime-offset valuation.

Editorial note. That valuation is $v_p(T_{p-1}(Y)) = v_p(Y) - 1$, in the proof of Theorem 2.12. The proof below is a second route to it: the one in this part reduces $pT_{p-1}(Y)/Y$ modulo p ((2.28)), the one below compares the valuations of all terms of the multinomial expansion.

2.14.2 Two elementary valuation bounds

For a multiplicity vector $I = (i_1, \dots, i_d)$ with

$$\sum_j j i_j = d, \quad \ell = \sum_j i_j,$$

the corresponding term in the multinomial expansion of $T_d(Y)$ is

$$C_I = \frac{Y^\ell}{\prod_j i_j!} \prod_j h_j^{i_j}, \quad Y^\ell = Y(Y-1) \cdots (Y-\ell+1). \quad (2.59)$$

Terms with $\ell > Y$ are zero and may be omitted.

Lemma 2.41. *If $1 \leq \ell \leq Y$ and $\ell < p^{e+1}$, where $e = v_p(Y)$, then*

$$v_p(Y^\ell) \geq e + v_p((\ell-1)!). \quad (2.60)$$

There is equality when $\ell \leq p^e$.

Proof. For $1 \leq s < \ell < p^{e+1}$, one has $v_p(s) \leq e$. If $v_p(s) < e$, then $v_p(Y-s) = v_p(s)$. If $v_p(s) = e$, both Y and s are divisible by p^e , giving $v_p(Y-s) \geq v_p(s)$. Sum these inequalities over s and include the factor Y . When $\ell \leq p^e$, every $s < \ell$ has valuation below e , so all comparisons are equalities. $\square$

In particular, we do *not* assert the equality when a competing term has more than p^e factors; such competitors can occur for $p > 2$.

The integer identity

$$\frac{d(\ell-1)!}{\prod_j i_j!} = \sum_{j: i_j > 0} j \frac{(\ell-1)!}{(i_j-1)! \prod_{h \neq j} i_h!} \in \mathbb{Z}_{>0} \quad (2.61)$$

gives

$$v_p((\ell-1)!) - \sum_j v_p(i_j!) \geq -v_p(d). \quad (2.62)$$

Every summand on the right of (2.61) is an integer multiple of a multinomial coefficient. Combining (2.59), Lemma 2.41 and (2.62), whenever $\ell < p^{e+1}$ we obtain

$$v_p(C_I) \geq e - v_p(d) - \sum_j i_j v_p(j(j+1)). \quad (2.63)$$

Because consecutive integers are coprime, $v = v_p(j(j+1)) \geq 1$ forces $j \geq p^v - 1 \geq v(p-1)$. Equality occurs only for $v = 1$ and $j = p-1$. Thus

$$v_p(j(j+1)) \leq \frac{j}{p-1}, \quad \text{with equality exactly at } j = p-1. \quad (2.64)$$

The case $v = 0$ is strictly below $j/(p-1)$ since $j \geq 1$.

2.14.3 The principal progression

Proof of Theorem 2.39. Set $d = (p-1)\mu$. Every nonzero term has

$$\ell \leq d \leq (p-1)p^e < p^{e+1},$$

so (2.63) applies. Inequality (2.64) gives

$$\sum_j i_j v_p(j(j+1)) \leq \mu,$$

with equality only for $I = \mu \mathbf{e}_{p-1}$, where $\mathbf{e}_j$ has a one at position j . Every other vector has this integer sum at most $\mu-1$. Since $v_p(d) = v_p(\mu)$, its term has valuation at least

$$e - v_p(\mu) - \mu + 1.$$

The distinguished term is $\binom{Y}{\mu} h_{p-1}^\mu$. It is nonzero because $\mu \leq p^e \leq Y$. The equality part of Lemma 2.41 gives

$$v_p \binom{Y}{\mu} = e - v_p(\mu), \quad v_p(h_{p-1}) = -1.$$

Thus its valuation is $e - v_p(\mu) - \mu$, uniquely smaller than all competitors. A sum with a unique least valuation has that valuation, proving (2.56). $\square$

Remark 2.42 (Editorial: manuscripts 12 and 17 on a common range). For $p > \mu \geq 2$ the hypothesis $\mu \leq p^e$ of Theorem 2.39 says $p \mid Y$. On that range the theorem also follows from the proof of Lemma 2.23: G/Y is p -integral and congruent to $(Y-1) \cdots (Y-\mu+1)/\mu!$, which at $Y \equiv 0$ is the unit $(-1)^{\mu-1}/\mu$; so $v_p(G(Y)) = v_p(Y)$ and $v_p(T_d(Y)) = e - \mu$, which is (2.56) since $v_p(\mu) = 0$. Manuscript 12 states only that the zeros lie in the classes $1 \leq j \leq \mu-1$, not the valuation. Conversely, its complete-offset theorems are not implied by manuscript 17, which needs $p \mid Y$ there. Manuscript 17 is new relative to manuscript 12 and to this part for $\mu \geq p$ (for instance every offset at $p = 2$ or 3) and on the shifted, odd offsets of Theorem 2.40. The three manuscripts share the “bad index” multinomial count after Davis; (2.63) is its cleanest general form.

2.14.4 One step above the progression

Proof of Theorem 2.40. Set $d = (p-1)\mu + 1$. The strict bound $\mu < p^e$ again gives $d < p^{e+1}$, and $p \nmid d$ removes the term $v_p(d)$ in (2.63). The denominator-valuation sum is an integer at most μ . If it is at most $\mu-1$, the corresponding term has valuation at least $e - \mu + 1$.

Otherwise its value is μ , and hence

$$\sum_j i_j (j - (p-1)v_p(j(j+1))) = 1.$$

For odd p , the only indices whose parenthesized value is zero or one are $j = p - 1$ (value zero), and $j = 1, p$ (value one). Indeed, valuation at least two would require

$$j \geq p^v - 1 > v(p - 1) + 1 \quad (v \geq 2, p \geq 3).$$

Therefore only the two vectors

$$\mu \mathbf{e}_{p-1} + \mathbf{e}_1, \quad (\mu - 1) \mathbf{e}_{p-1} + \mathbf{e}_p$$

remain. Put $A = h_{p-1}$, $B = h_p$. Their combined contribution is exactly

$$\frac{Y^\mu}{(\mu - 1)!} A^{\mu-1} \left(B + \frac{Y - \mu}{2\mu} A \right). \quad (2.65)$$

The entire prefactor outside the brackets has valuation $e - \mu + 1$. Since $\mu < p^e$, one has $v_p(\mu) < e$, and so $(Y - \mu)/\mu \equiv -1 \pmod{p}$. Moreover (2.41) gives $pA \equiv pB \equiv 1 \pmod{p}$. Multiplying the bracket in (2.65) by p gives

$$1 - \frac{1}{2} = \frac{1}{2} \not\equiv 0 \pmod{p}.$$

Thus the bracket has valuation -1 , and the pair has valuation $e - \mu$, strictly below every other term. This proves (2.58). $\square$

The conditions have real content. Allowing $\mu = p^e$ in the shifted theorem would include the symmetry zeros: for example $p = 3, \mu = 3, Y = 6$ gives $T_7(6) = 0$. Dropping $p \nmid d$ can also invalidate the valuation formula: $p = 3, \mu = 4, Y = 9$ gives

$$T_9(9) = \frac{2021}{268800}, \quad v_3(T_9(9)) = -1,$$

whereas $e - \mu = -2$. This is a failure of the unrestricted valuation formula, not a new coefficient zero.

Editorial note. $T_7(6) = 0$ is the symmetry zero $b(13, 6) = 0$, the case $h = 3$ of Proposition 2.8.

2.14.5 New infinite regions and the remaining obstruction

By (2.42), either theorem gives the valuation of $b(Y + d, Y)$ by adding $v_p((Y + d)!) - v_p(Y!)$. Here are four infinite divisibility progressions beyond the source's completed offsets $d \leq 16$; each listed $d + 1$ is composite.

d	p	condition on Y	$v_p(T_d(Y))$
17	2	$32 \mid Y$	$v_2(Y) - 17$
17	3	$9 \mid Y$	$v_3(Y) - 8$
20	3	$27 \mid Y$	$v_3(Y) - 10$
21	5	$25 \mid Y$	$v_5(Y) - 5$

For their first positive Y , the respective valuations of $b(Y + d, Y)$ are 3, 0, 1, 1. The supplied exact output records the corresponding integers.

Every hypothetical additional integer zero must avoid all divisibility progressions covered by the two theorems. In particular, taking $p = 2$ in Theorem 2.39 gives the necessary restriction

$$T_d(Y) = 0 \implies 2^{v_2(Y)} < d.$$

The criteria do not exclude every pair (d, Y) , and they do not prove the conjectured complete zero list consisting of the symmetry family and $b(8, 5)$.

2.14.6 Verification and positioning

An integer-triangle verifier (`code/06-padic-valuations-verify_padic.py`) checks both valuation formulas through row $m = 360$, for the eleven primes through 31: 7,749 principal and 2,299 shifted instances pass. A separate direct rational power-series recurrence checks 3,073 principal and 437 shifted instances for $1 \leq Y \leq 120$, $d \leq 60$. A third checker, using direct integer-scaled convolution, passes 2,421 principal and 463 shifted cases, including 1,352 principal boundary cases $\mu = p^e$, and 5,500 denominator checks. These independent finite checks supplement the infinite proof and are included with their exact outputs.

The source comparison is with the pinned ProveIt manuscript, this report at commit `ca81647a9`; its current default-branch file was checked again on 1 October 2026 and still contains the prime-offset theorem and fixed-offset certificates through 16. Davis’s paper [15] supplies the classical valuation method for a related log series. The contribution here is its direct adaptation to (2.41), the two-term shifted calculation, and the resulting exact nonvanishing families for this coefficient array. No exhaustive priority claim is made. A complete integer-root classification, and hence the exact A293239 term-count formula, remain open. The proof is not a proof-assistant formalization or externally refereed publication.

2.15 Lehmer–Comtet additions: provenance, files and formal status

2.15.1 Provenance

Subsections 2.11–2.15 were built from **three** manuscripts of batch 72A, which arrived in commit `1512ef835` and were placed in this report by commit `d292c6765`, as its additions 05, 04 and 06. All three are dated 1 October 2026, carry the line “Research note prepared for Vladimir Reshetnikov with OpenAI”, and are pinned to ProveIt commit `ca81647a9f679e96d49ffdbc54b0c0ab13d10a36`, whose text of this article (blob `7344f3aa4dc9`) they cite.

- Manuscript 12 (addition 05), *Complete Nonvanishing on Prime-Multiple Offsets: Lehmer–Comtet coefficients and derivatives of x^x* , 6-page PDF, archive `ProveIt_Lehmer_Comtet_Prime_Multiple_Nonvanishing.zip`. Its six sections form Subsection 2.12. It contributes Theorem 2.19, its three corollaries, the localization, barrier, cone and small-exponent lemmas, and Theorem 2.28.
- Manuscript 09 (addition 04), *Stirling Residues and Finite Candidate Lists: Arithmetic restrictions on Lehmer–Comtet zeros*, 6-page PDF, archive `ProveIt_Lehmer_Comtet_Stirling_Residues_and_Candidate_Lists.zip`. Its six sections form Subsection 2.13. It contributes the Stirling residue, the root displacements, the candidate list, Corollaries 2.35 and 2.36, and Theorem 2.38.
- Manuscript 17 (addition 06), *Uniform p -adic Nonvanishing: Two valuation families for Lehmer–Comtet coefficients*, 4-page PDF, archive `ProveIt_Lehmer_Comtet_Uniform_Padic_Nonvanishing.zip`. Its six sections form Subsection 2.14. It contributes Theorems 2.39 and 2.40 and their valuation lemmas.

Every theorem, lemma and corollary of the three manuscripts is printed, with its proof and with every non-claim and limitation the manuscripts state; none was weakened or strengthened. The one exception to printing in full is recorded in the second item below. The editorial choices were these.

- The additions are placed as subsections at the end of Part 2, so that no existing section, theorem or equation number moves; the order is 12, 09, 17, because 09 builds on 12.

- Three lemmas that manuscript 09 re-proves from manuscript 12 are printed once, as manuscript 12's; 09's statements and its condensed proofs are recorded in Remarks 2.29, 2.31 and 2.37. The three manuscripts' identical definitions of H , T_d and $b(Y + d, Y)$ are printed once, as (2.41) and (2.42).
- Results of this part that the manuscripts re-derive are credited where they occur (the list in Subsection 2.11); the restated offset-eight certificate is kept, with a note identifying it as $R_8(Y + 8)$.
- The letters are changed as in Table 2.A, with no change of normalization. The manuscripts' citations of "ProveIt, Part I" are re-targeted to this part, manuscript 09's citations of its companion to Subsection 2.12, and their file names to the shipped names. The references of Davis, Ravichandran and Adelberg–Lengyel are merged into the bibliography.
- Remarks 2.26, 2.34 and 2.42, the notation subsection and the editorial notes are additions of the intake, not of the manuscripts.

Not shipped: the three manuscripts' sources, PDFs and README.md files; manuscript 09's embedded copy of manuscript 12 (`prime_multiple_companion_source.zip`, byte-identical to archive 12, and `prime_multiple_companion.pdf`, byte-identical to 12's PDF); and the three copies of the generic TeX wrapper `build_local.sh`. The archives survive in the history of commit 1512ef835. The READMEs add no mathematical claim beyond the manuscripts; those of manuscripts 12 and 17 say that the proofs received "independent review", which this report cannot confirm.

2.15.2 Files

The manuscripts' code and data are shipped byte-identical to the delivery, with the prefixes 04-stirling-residues- (09), 05-prime-multiple- (12) and 06-padic-valuations- (17). All programs use only the Python standard library.

Shipped path (delivery name)	Purpose
<code>code/05-prime-multiple-verify.py</code> (<code>verify.py</code>)	Exact checks of Subsection 2.12; writes <code>checks.json</code> beside itself.
<code>data/05-prime-multiple-checks.json</code> (<code>checks.json</code>)	Its recorded output.
<code>code/05-prime-multiple-replay_convolution.py</code> (<code>replay_convolution.py</code>)	Independent convolution replay; writes a <code>.json</code> of its own name beside itself.
<code>data/05-prime-multiple-replay_convolution.json</code> (<code>replay_convolution.json</code>)	Its recorded output.
<code>code/04-stirling-residues-verify_stirling.py</code> (<code>verify_stirling.py</code>)	Stirling residues, triangle congruences, root displacements; writes <code>stirling_checks.json</code> beside itself.
<code>data/04-stirling-residues-stirling_checks.json</code> (<code>stirling_checks.json</code>)	Its recorded output.
<code>code/04-stirling-residues-verify_prime_square_lifts.py</code> (<code>verify_prime_square_lifts.py</code>)	Roots modulo p^3 at offset $p^2 - 1$, $p = 3, 5, 7, 11$; writes <code>prime_square_lifts.json</code> beside itself.

Shipped path (delivery name)	Purpose
data/04-stirling-residues-prime_square_lifts.json (prime_square_lifts.json)	Its recorded output.
code/04-stirling-residues-verify_square_candidates.py (verify_square_candidates.py)	Exact values at the one $(p-1)^2$ candidate, seven primes; writes <code>square_candidates.json</code> beside itself.
data/04-stirling-residues-square_candidates.json (square_candidates.json)	Its recorded output.
code/04-stirling-residues-replay_falling_products.py (replay_falling_products.py)	Independent integer falling-product replay (run with <code>-0</code>); writes a <code>.json</code> of its own name beside itself.
data/04-stirling-residues-replay_falling_products.json (replay_falling_products.json)	Its recorded output.
code/06-padic-valuations-verify_padic.py (verify_padic.py)	Integer-triangle check of both valuation families through row 360; writes <code>padic_checks.json</code> beside itself.
data/06-padic-valuations-padic_checks.json (padic_checks.json)	Its recorded output.
code/06-padic-valuations-check_rational_powers.py (check_rational_powers.py)	Direct rational power recurrence; prints its JSON to standard output and writes nothing.
data/06-padic-valuations-rational_power_checks.json (rational_power_checks.json)	Its recorded output.
code/06-padic-valuations-replay_padic_powers.py (replay_padic_powers.py)	Integer-scaled convolution replay with boundary and denominator checks; writes a <code>.json</code> of its own name beside itself.
data/06-padic-valuations-replay_padic_powers.json (replay_padic_powers.json)	Its recorded output.

The JSON files keep the manuscripts' letters: their `r` is the offset d and their `m` the multiplier μ , and the field `primitive_r8` of the convolution replay holds the coefficients of $F(Y) = R_8(Y+8)$, not of $R_8(X)$. At the intake every program was run under its shipped name on a copy of this directory and exited with status 0; every output equalled the shipped file, up to line endings (a Windows Python may write CRLF). The programs write beside themselves, never into `data/`; the README gives the commands.

2.15.3 Formal status

Nothing in this addition is formalized, and its placement in this report confers no formal status. No Lean or Rocq file in the repository mentions the Lehmer–Comtet numbers, OEIS A008296 or OEIS A293239. The manuscripts claim no formal verification, no referee and no exhaustive priority.

3 x^{x^2} : OEIS A290268

Framework. This part instantiates Theorem 1.2(b) — the clause normalized by x^{-n} — with the frame $z = x^2$, $L = \log x$; Proposition 3.3 below is that instance, derived here independently, and the $-n$ in its recurrence is the normalizer’s contribution. Lemma 3.2 is the special case of Theorem 1.4 this part needs.

Notation. In this part z **denotes** x^2 , not $\log x$; compare Part 4, where z is $\log x$. See Section 1.5. Throughout this part $a(n)$ denotes the term count for x^{x^2} .

Later addition (29 September 2026). Sections 3.14–3.26, at the end of this part, add a later manuscript that closes the missing nonvanishing step at logarithmic deficits $m = 3, 4$ and improves the lower bound $\Lambda(n)$ below. They use depth coordinates in which the roles of j and k differ from this part’s (Table 3). The text of the original report is unchanged; dated notes like this one point forward.

Second later addition (30 September 2026). Sections 3.27–3.34, after the first addition, print a further manuscript whose main theorem is the case $m = 3$ of that addition’s depth theorem, proved by a second route: a harmonic sign law and positivity propagation in the logarithm exponent. The open region is unchanged.

Third later addition (1 October 2026). Sections 3.35–3.41, after the second, print two further manuscripts. The first closes the missing step at deficits $m = 5, 6$ (Theorem 3.71), so the open region is now $m \geq 7$, $n \geq 2k + 2$, and replaces the tower-type cutoffs of the first addition by $\lceil (2m + 1)e^m \rceil$ (Theorem 3.81). The second improves the lower bound to $3n^2/8 + (n/2) \log n - (n/2) \log \log n - 5n/4$ for $n \geq e^{10}$ (Theorem 3.83) and bounds the summatory count (Theorem 3.85); the leading constant of the explicit lower bound is still $3/8$.

Fourth later addition (1 October 2026, batch 73). Sections 3.42–3.51, after the third, print two further manuscripts. They close the missing step at deficits $m = 7$ (Theorem 3.99, by two independent certificates) and $m = 8$ (Theorem 3.100), so the open region is now $m \geq 9$, $n \geq 2k + 2$. They also give a practical reduction at every deficit from three exact moment anchors (Theorem 3.97) and the eight-deficit count bound of Corollary 3.106; the leading constant of the explicit lower bound is still $3/8$. One of them re-proves deficits five and six and, more weakly, the cutoffs and the $n \log n$ bound of the third addition; those parts are printed as a second certificate and as remarks.

[Added 1 October 2026, batch 72A: two sibling reports bear on this part without being part of it. **power-tower-exponent-supports** proves $a(n) = n^2/2 + o(n^2)$ (its Corollary 13.1), without an effective threshold, which answers Question 6 of Subsection 3.24; **a290268-unbounded-deficits** studies the coefficients when the deficit grows with n (phase laws, an Airy law at the fold, cosine laws, arithmetic families). Neither proves Conjecture 3.19.]

Abstract of the original report. Let $a(n)$ be the number of nonzero monomials in the fully collected n th derivative of x^{x^2} . The conjecture in OEIS A290268 predicts

$$a(n) = U(n) := \frac{n^2}{2} + n + 1 - (n \bmod 2) \left(\frac{1}{2} + \left\lfloor \frac{n+1}{8} \right\rfloor \right).$$

This article does not establish the equality for all n . It proves the unconditional bound $a(n) \leq U(n)$ by identifying every cancellation needed to explain the proposed formula. It also proves positivity throughout the upper half of the coefficient triangle, classifies all zeros on the first two logarithmic-deficit diagonals, and gives explicit quadratic lower bounds. An exact generating-function identity and a finite-difference formula reduce the full conjecture to a precise nonvanishing assertion. Integer arithmetic verifies the support through $n = 200$; two complete modular computations, combined with the proved zero identities, certify $a(n) = U(n)$ for every $0 \leq n \leq 3000$. All source code, finite certificates, and reproduction instructions accompany the article. The remaining nonvanishing step is stated explicitly rather than assumed.

3.1 The problem and the status of the result

Put

$$f(x) = x^{x^2} = \exp(x^2 \log x), \quad x > 0.$$

The sequence A290268 counts the terms in the fully expanded derivative $f^{(n)}(x)$, after identical monomials have been collected. The OEIS entry was contributed by Vladimir Reshetnikov in October 2017; its formula section labels both the generating function and the explicit expression as conjectural. Peter Luschny's equivalent floor-function form is explicitly conditional on that conjecture [2].

For clarity, throughout this article the letter U denotes the *proposed* answer, not an already proved formula for a :

$$U(n) = \frac{n^2}{2} + n + 1 - (n \bmod 2) \left(\frac{1}{2} + \left\lfloor \frac{n+1}{8} \right\rfloor \right). \quad (3.1)$$

Equivalently,

$$U(2r) = 2r^2 + 2r + 1, \quad U(2r+1) = 2(r+1)^2 - \left\lfloor \frac{r+1}{4} \right\rfloor. \quad (3.2)$$

The two central issues are different. One must explain why certain coefficients vanish, and one must also exclude every other cancellation. The first task is completed here. The second is completed in substantial regions and in a large finite range, but not in full generality.

Theorem 3.1 (Unconditional bounds). *For every integer $n \geq 0$,*

$$\Lambda(n) \leq a(n) \leq U(n), \quad \Lambda(n) = \begin{cases} (3n^2 + 10n + 8)/8, & n \text{ even}, \\ (3n^2 + 12n + 1)/8, & n \text{ odd}. \end{cases} \quad (3.3)$$

In particular, $a(n) = \Theta(n^2)$. The proposed leading constant $1/2$ is not established by these bounds.

The upper bound is proved in Section 3.6; the lower bound is proved in Section 3.4. Section 3.9 gives a reproducible finite certification of the stronger equality through 3000. The exact outstanding assertion appears in Section 3.10.

Update (29 September 2026). The lower bound $\Lambda(n)$ is improved by Corollary 3.45 to $(3n^2 + 26n - 72)/8$ for even $n \geq 8$ and $(3n^2 + 28n - 111)/8$ for odd $n \geq 17$ (Remark 3.46), a linear-term gain of $2n - 10$, respectively $2n - 14$; the leading constant $3/8$ is unchanged. Theorem 3.1 stands as stated.

Update (1 October 2026, batch 72A). Corollary 3.78 adds the classified depths five and six, a further linear gain, and Theorem 3.83 gives $a(n) \geq 3n^2/8 + (n/2) \log n - (n/2) \log \log n - 5n/4$ for every $n \geq e^{10}$; the leading constant of every explicit lower bound is still $3/8$. Theorem 3.85 bounds the cumulative count, $\sum_{n \leq X} a(n) \geq 11X^3/72 + 5X^2/6 - O(X)$. [Added 1 October 2026, batch 73: Corollary 3.106 adds depths seven and eight: $a(n) \geq (3n^2 + 42n - 280)/8$ for even $n \geq 16$ and $(3n^2 + 44n - 351)/8$ for odd $n \geq 33$, a further gain of $n - 15$, respectively $n - 17$; the leading constant of every explicit lower bound is still $3/8$.]

[Added 1 October 2026, batch 72A: the proposed leading constant $1/2$ is now established in the limit: the sibling report `power-tower-exponent-supports` proves $\lim a(n)/n^2 = 1/2$ (its Corollary 13.1, the case $a = 2$ of its Theorem 3.1), without an effective threshold. Theorem 3.1 stands as stated, and as explicit bounds $\Lambda(n)$ and the improvements above, all with leading constant $3/8$, are still the best known.]

3.2 A canonical polynomial expansion

Set $z = x^2$ and $L = \log x$. Define polynomials $P_n \in \mathbb{Z}[z, L]$ by

$$f^{(n)}(x) = f(x)x^{-n}P_n(x^2, \log x), \quad P_0 = 1. \quad (3.4)$$

Write

$$P_n(z, L) = \sum_{k,j} c_{n,k,j} z^k L^j. \quad (3.5)$$

Thus $c_{n,k,j}$ is the coefficient of $x^{x^2+2k-n}(\log x)^j$ in the derivative. Counting terms means counting nonzero $c_{n,k,j}$, rather than counting the summands generated by a particular symbolic differentiation algorithm.

Lemma 3.2 (Uniqueness of the collected expansion). *A finite identity $\sum_{k,j} b_{k,j} x^{2k} (\log x)^j = 0$ for all $x > 0$ forces every $b_{k,j}$ to vanish. Consequently, the expansion (3.4)–(3.5) is canonical.*

Proof. Substitute $x = e^L$ and group the identity as $\sum_k e^{2kL} B_k(L) = 0$, with polynomial B_k . If K is the largest index with $B_K \neq 0$, division by e^{2KL} shows that $B_K(L)$ tends to zero as $L \rightarrow +\infty$: all remaining terms are polynomials multiplied by decaying exponentials. A nonzero polynomial cannot tend to zero at $+\infty$. This contradiction eliminates K , and then all indices. $\square$

The normalized derivative obeys a particularly small recurrence.

Proposition 3.3 (Polynomial and coefficient recurrences). *For every $n \geq 0$,*

$$P_{n+1} = (z(2L+1) - n)P_n + 2z \frac{\partial P_n}{\partial z} + \frac{\partial P_n}{\partial L}. \quad (3.6)$$

With nonexistent coefficients interpreted as zero,

$$\boxed{c_{n+1,k,j} = (2k - n)c_{n,k,j} + (j+1)c_{n,k,j+1} + c_{n,k-1,j} + 2c_{n,k-1,j-1}.} \quad (3.7)$$

For $n \geq 1$, the only possible nonzero indices satisfy

$$1 \leq k \leq n, \quad 0 \leq j \leq k. \quad (3.8)$$

Proof. We have $f'/f = x(2L+1)$, $dz/dx = 2z/x$, and $dL/dx = 1/x$. Differentiate (3.4) and factor out $f(x)x^{-n-1}$ to obtain (3.6). Taking the coefficient of $z^k L^j$ gives (3.7). Starting from $P_0 = 1$, the first derivative is $P_1 = z(2L+1)$. The four operations in (3.6) preserve $k \geq 1$ and $j \leq k$, and increase k by at most one. Induction proves (3.8) and integrality. $\square$

For example,

$$\begin{aligned} P_1 &= z(2L+1), \\ P_2 &= z(2L+3) + z^2(2L+1)^2, \\ P_3 &= 2z + z^2(12L^2 + 24L + 9) + z^3(2L+1)^3. \end{aligned}$$

These have respectively 2, 5, and 8 nonzero coefficients, in agreement with the definition. Already P_3 illustrates a degree loss: its z -coefficient has no L -term. Later cancellations are less obvious. For instance,

$$[z^2]P_7(z, L) = 28, \quad [z^2]P_9(z, L) = -864L. \quad (3.9)$$

The missing L -term in the first polynomial and the missing constant in the second will follow from the same reflection principle.

3.3 Exact formulas for individual coefficients

3.3.1 An exponential generating function

Define two formal power series

$$A(t) = t(2+t) = (1+t)^2 - 1, \quad B(t) = (1+t)^2 \log(1+t). \quad (3.10)$$

Formal coefficient extraction is denoted by $[t^n]$. All logarithms in formal series are expanded at the origin.

Proposition 3.4 (Generating function). *The complete coefficient array is described by*

$$\sum_{n \geq 0} P_n(z, L) \frac{t^n}{n!} = \exp(z(A(t)L + B(t))). \quad (3.11)$$

In particular, if $m = k - j \geq 0$, then

$$c_{n,k,j} = \frac{n!}{j!m!} [t^n] A(t)^j B(t)^m. \quad (3.12)$$

Proof. Taylor expansion at the point x gives

$$\frac{f(x(1+t))}{f(x)} = \sum_{n \geq 0} \frac{x^n f^{(n)}(x)}{f(x)} \frac{t^n}{n!}.$$

The logarithm of the ratio on the left is

$$x^2(((1+t)^2 - 1) \log x + (1+t)^2 \log(1+t)).$$

This proves (3.11) after substitution of z, L . Alternatively, its coefficients satisfy (3.6), so the identity also holds with z, L algebraically independent. Expanding separately the two exponentials $\exp(zA(t)L)$ and $\exp(zB(t))$ gives (3.12). $\square$

Because both A and B start with a nonzero term of degree one, their product in (3.12) starts at degree k . This independently explains $k \leq n$. The difficulty is that B does not have only positive coefficients:

$$B(t) = t + \frac{3}{2}t^2 + \frac{1}{3}t^3 - \frac{1}{12}t^4 + \frac{1}{30}t^5 - \dots$$

Consequently, the coefficient formula is not itself a proof of nonvanishing.

3.3.2 Falling factorials and finite differences

Let

$$F_n(u) = u^{\underline{n}} = u(u-1) \cdots (u-n+1), \quad F_0(u) = 1,$$

and let $\Delta_2 G(u) = G(u+2) - G(u)$. Set $Q_{n,j}(u) = \Delta_2^j F_n(u)$.

Proposition 3.5 (Finite-difference formula). *For $m = k - j \geq 0$,*

$$c_{n,k,j} = \frac{1}{j!m!} \left. \frac{d^m}{du^m} Q_{n,j}(u) \right|_{u=2m} \quad (3.13)$$

$$= \frac{1}{j!} [v^m] \sum_{h=0}^j (-1)^{j-h} \binom{j}{h} F_n(2m+2h+v). \quad (3.14)$$

Proof. The generalized binomial theorem gives $n![t^n](1+t)^u = F_n(u)$. Differentiating m times in u introduces $\log^m(1+t)$. Taking j forward differences of step two introduces $((1+t)^2 - 1)^j = A(t)^j$. Evaluation at $u = 2m$ introduces $(1+t)^{2m}$, which combines with the logarithm to give $B(t)^m$. Equation (3.12) now proves (3.13); Taylor's formula proves (3.14). $\square$

Derivatives of falling factorials also occur in the theory of shifted Stirling numbers and elementary symmetric harmonic sums; Poon gives a broader account of these connections [11]. No general nonvanishing theorem from that literature is being assumed here. Formula (3.14) is also an independent way to compute a coefficient using integer polynomial products.

3.4 A transformation proving positivity

A fractional-linear change of variable turns a large region of the coefficient array into a manifestly positive expression. Define

$$H(u) = \frac{\operatorname{atanh} u}{u} = \sum_{r \geq 0} \frac{u^{2r}}{2r+1}, \quad H(0) = 1. \quad (3.15)$$

The value at zero is understood by its formal power series.

Proposition 3.6 (Transformed coefficient formula). *For $m = k - j \geq 0$ and $n \geq k$,*

$$c_{n,k,j} = \frac{n!}{j!m!} 2^{2j+m-n} [u^{n-k}](1-u)^{n-2k-1}(1+u)^{2m} H(u)^m. \quad (3.16)$$

Negative integer powers of $1-u$ have their usual formal expansions.

Proof. In the residue expression for $[t^n]A(t)^j B(t)^m$, substitute $t = 2u/(1-u)$. Then

$$1+t = \frac{1+u}{1-u}, \quad A(t) = \frac{4u}{(1-u)^2}, \quad B(t) = \frac{2(1+u)^2}{(1-u)^2} \operatorname{atanh} u, \quad dt = \frac{2 du}{(1-u)^2}.$$

Using $[t^n]G(t) = \operatorname{Res}_{t=0} G(t)t^{-n-1} dt$, the constant factor becomes 2^{2j+m-n} . The remaining power of $1-u$ is $n-2k-1$. The factors $u^j(\operatorname{atanh} u)^m$ contribute $u^k H(u)^m$, so the resulting residue extracts the coefficient of u^{n-k} . Multiplying by $n!/(j!m!)$ proves the formula. The formal change-of-variable rule applies because $2u/(1-u)$ has zero constant term and nonzero linear coefficient. $\square$

Theorem 3.7 (A nonvanishing region). *For $1 \leq k \leq n$ and $0 \leq j \leq k$,*

$$\begin{aligned} n \leq 2k &\implies c_{n,k,j} > 0, \\ n = 2k+1, j < k &\implies c_{n,k,j} > 0. \end{aligned} \quad (3.17)$$

Proof. If $n \leq 2k$, the factor $(1-u)^{n-2k-1}$ is $(1-u)^{-q}$ with $q \geq 1$. Every coefficient of this factor is strictly positive. The other two factors in (3.16) have nonnegative coefficients and constant term one. Their product therefore has a strictly positive coefficient in every nonnegative degree.

If $n = 2k+1$, that factor disappears. The condition $j < k$ means $m \geq 1$. The series $H(u)^m$ has a strictly positive coefficient in every even degree. The polynomial $(1+u)^{2m}$ has positive constant and linear coefficients. An even target degree can be supplied entirely by $H(u)^m$; an odd target degree can use the linear term of $(1+u)^{2m}$ and an even degree from $H(u)^m$. Thus the target coefficient of degree $n-k$ is strictly positive in both cases. The prefactor in (3.16) is positive. $\square$

Corollary 3.8 (Quadratic lower bounds). *The lower bound in Theorem 3.1 holds.*

Proof. For $n = 2r \geq 2$, Theorem 3.7 supplies all coefficients with $r \leq k \leq 2r$, giving

$$a(2r) \geq \sum_{k=r}^{2r} (k+1) = \frac{(r+1)(3r+2)}{2}.$$

For $n = 2r + 1$ with $r \geq 1$, it supplies all coefficients with $r + 1 \leq k \leq 2r + 1$ and the r coefficients $0 \leq j < r$ at $k = r$. Hence

$$a(2r+1) \geq r + \sum_{k=r+1}^{2r+1} (k+1) = \frac{3(r+1)(r+2)}{2} - 1.$$

These are precisely (3.3). The cases $n = 0, 1$ follow directly from $P_0 = 1$ and $P_1 = z(2L+1)$. $\square$

3.5 The two forced families of zeros

3.5.1 Degree-forced zeros on the top diagonal

Setting $j = k$, and therefore $m = 0$, in (3.12) yields an explicit coefficient.

Proposition 3.9 (Top logarithmic degree). *For $k \geq 1$ and $k \leq n \leq 2k$,*

$$c_{n,k,k} = \frac{n!}{k!} \binom{k}{n-k} 2^{2k-n} > 0. \quad (3.18)$$

For $n > 2k$, one has $c_{n,k,k} = 0$.

Proof. The relevant series is the polynomial $A(t)^k = t^k(2+t)^k$, of degree $2k$. Its coefficient of t^n is $\binom{k}{n-k} 2^{2k-n}$ in the indicated range and zero above it. $\square$

For a fixed $n \geq 1$, this removes exactly $\lfloor (n-1)/2 \rfloor$ cells from the possible triangle.

3.5.2 Reflection-forced cancellations

The less obvious zeros follow from symmetry about the midpoint of the roots of a falling factorial.

Lemma 3.10 (Reflection of a finite difference). *For all $n, j \geq 0$,*

$$Q_{n,j}(n-2j-1-u) = (-1)^{n-j} Q_{n,j}(u). \quad (3.19)$$

Proof. First, $F_n(n-1-u) = (-1)^n F_n(u)$, by reversing the order of the factors. Expand the left side of (3.19):

$$\sum_{h=0}^j (-1)^{j-h} \binom{j}{h} F_n(n-2j-1-u+2h).$$

Reflection turns its last factor into $(-1)^n F_n(u+2(j-h))$. Reindexing by $\ell = j-h$ leaves the factor $(-1)^{n-j}$ times the usual expansion of $\Delta_2^j F_n(u)$. $\square$

Theorem 3.11 (Reflection zeros). *Suppose $1 \leq k \leq n$ and $0 \leq j < k$. Then*

$$k \text{ even}, \quad n = 4k - 2j + 1 \quad \implies \quad c_{n,k,j} = 0. \quad (3.20)$$

Proof. Put $m = k - j$. The reflection center in Lemma 3.10 is $(n - 2j - 1)/2$. Under the displayed condition this equals $2m$, precisely the evaluation point in (3.13). The parity of $Q_{n,j}(2m + v)$ as a polynomial in v is $n - j$. Its coefficient of v^m therefore vanishes whenever $n - j - m = n - k$ is odd. The relation $n = 4k - 2j + 1$ makes n odd, so $n - k$ is odd when k is even. Equation (3.14) completes the proof. $\square$

There is also a direct proof using (3.16). The relation in (3.20) gives $n - 2k - 1 = 2m$, so the series whose coefficient is extracted becomes

$$(1 - u^2)^{2m} H(u)^m. \quad (3.21)$$

This series is even. When k is even, $n - k$ is odd, and its coefficient of that degree is zero. This second proof makes the cancellation visible without evaluating any large integer coefficient.

Remark 3.12 (What symmetry does not say). The theorem is an implication, not an equivalence. An even power series has zero odd coefficients, but it can also have zero even coefficients. Similarly, away from the reflection center, a derivative of a polynomial can vanish for reasons unrelated to symmetry. Excluding such additional zeros is a separate requirement of the full conjecture.

3.6 Counting the forced zeros

For $n \geq 1$, the triangle (3.8) has

$$T(n) = \sum_{k=1}^n (k+1) = \frac{n(n+3)}{2} \quad (3.22)$$

cells. Let Z_n^{top} be the top-diagonal cells removed by Proposition 3.9, and let Z_n^{ref} be the cells in Theorem 3.11. These sets are disjoint, since the second requires $j < k$.

Lemma 3.13 (Number of reflection zeros). *For even n , Z_n^{ref} is empty. For odd n ,*

$$|Z_n^{\text{ref}}| = \left\lfloor \frac{n+1}{8} \right\rfloor. \quad (3.23)$$

Proof. Reflection zeros have odd n . Write $n = 2r + 1$ and $k = 2\ell$. The reflection relation determines $j = 2k - r = 4\ell - r$. The inequalities $0 \leq j < k$ become

$$\frac{r}{4} \leq \ell < \frac{r}{2}.$$

For $r \geq 1$, these automatically give $\ell \geq 1$ whenever the interval contains an integer. Its number of integers is

$$\left\lceil \frac{r}{2} \right\rceil - \left\lceil \frac{r}{4} \right\rceil = \left\lfloor \frac{r+1}{4} \right\rfloor,$$

as is checked in the four residue classes of r modulo four. For $r = 0$ both sides are zero. Since $(r+1)/4 = (n+1)/8$, the result follows. $\square$

Theorem 3.14 (The conjectured expression is an upper bound). *For every $n \geq 0$,*

$$a(n) \leq U(n). \quad (3.24)$$

Equality holds for a given n if and only if the coefficient triangle has no zeros other than $Z_n^{\text{top}} \cup Z_n^{\text{ref}}$.

Proof. For even $n = 2r \geq 2$, subtraction of the top-diagonal zeros from (3.22) gives

$$2r^2 + 3r - (r - 1) = 2r^2 + 2r + 1.$$

For odd $n = 2r + 1$, subtraction of both disjoint families gives

$$2r^2 + 5r + 2 - r - \left\lfloor \frac{r+1}{4} \right\rfloor = 2(r+1)^2 - \left\lfloor \frac{r+1}{4} \right\rfloor.$$

These are (3.2). Additional zeros could only decrease the number of nonzero coefficients, proving the inequality and the stated equivalence. Finally $a(0) = U(0) = 1$. $\square$

For example, at $n = 7$ there are 35 potential cells, three top-diagonal zeros, and the reflection zero $(k, j) = (2, 1)$, leaving 31. At $n = 9$, there are 54 potential cells, four top-diagonal zeros, and the reflection zero $(2, 0)$, leaving 49. The exact expansions in (3.9) display those two different cancellations. The floor term in the proposed formula is thus explained by a proved symmetry, not merely by fitting initial values.

3.7 Nonvanishing on the first two deficit diagonals

Call $m = k - j$ the *logarithmic deficit*. This section gives a complete zero classification for $m = 1$ and $m = 2$. Together with Theorem 3.7, it narrows the unproved portion of the problem.

3.7.1 A beta-integral representation

Suppose $n > 2k$, $m = k - j \geq 1$, and put

$$d = n - 2k - 1 \geq 0. \quad (3.25)$$

For sufficiently small complex ε , define

$$I_{m,j,d}(\varepsilon) = \int_0^1 t^{2m+\varepsilon} (1-t)^{d-\varepsilon} (2t-1)^j dt. \quad (3.26)$$

The integral and all derivatives required below converge uniformly on compact subsets of $|\varepsilon| < 1/2$.

Lemma 3.15 (Integral formula). *Under these assumptions,*

$$Q_{n,j}(2m + \varepsilon) = (-1)^{n-1} n! \frac{\sin(\pi\varepsilon)}{\pi} I_{m,j,d}(\varepsilon), \quad (3.27)$$

and consequently

$$c_{n,k,j} = \frac{(-1)^{n-1} n!}{j!} [\varepsilon^m] \left(\frac{\sin(\pi\varepsilon)}{\pi} I_{m,j,d}(\varepsilon) \right). \quad (3.28)$$

Proof. Euler's reflection formula and beta integral [13, 14] give, initially away from integer α ,

$$\begin{aligned} F_n(\alpha) &= (-1)^{n-1} \frac{\sin(\pi\alpha)}{\pi} \Gamma(\alpha+1) \Gamma(n-\alpha) \\ &= (-1)^{n-1} n! \frac{\sin(\pi\alpha)}{\pi} \int_0^1 t^\alpha (1-t)^{n-\alpha-1} dt. \end{aligned}$$

In the j th difference put $\alpha = 2m + 2h + \varepsilon$. The sine factor is independent of the integer h . The finite sum inside the integral is

$$\sum_{h=0}^j (-1)^{j-h} \binom{j}{h} \left(\frac{t}{1-t} \right)^{2h} = \frac{(2t-1)^j}{(1-t)^{2j}}.$$

The remaining exponent of $1-t$ is $d-\varepsilon$, proving (3.27). All terms are analytic near $\varepsilon = 0$, so the identity extends to zero as well. Taking the coefficient of ε^m and using (3.14) gives (3.28). $\square$

3.7.2 Deficit one

Theorem 3.16 (Complete signs for $m = 1$). *Suppose $m = k - j = 1$ and $n > 2k$, and define d by (3.25). Then*

$$\operatorname{sgn} c_{n,k,j} = (-1)^{n-1} \begin{cases} 1, & j \text{ even}, \\ \operatorname{sgn}(2-d), & j \text{ odd}. \end{cases} \quad (3.29)$$

In particular the only zeros in this range have j odd and $d = 2$.

Proof. The coefficient of ε in (3.28) is

$$c_{n,k,j} = \frac{(-1)^{n-1} n!}{j!} \int_0^1 t^2 (1-t)^d (2t-1)^j dt. \quad (3.30)$$

For even j , the integrand is nonnegative and positive except at finitely many points, so the integral is strictly positive. For odd j , pair t with $1-t$ and integrate over $1/2 < t < 1$. The paired integrand is

$$(2t-1)^j (t^2(1-t)^d - t^d(1-t)^2).$$

The ratio of the two positive terms in parentheses is $(t/(1-t))^{2-d}$. Since $t/(1-t) > 1$, their difference is identically zero when $d = 2$ and otherwise has the strict sign of $2-d$ throughout the open interval. This proves the sign formula. $\square$

These zeros are exactly the previously identified reflection zeros: j odd means $k = j + 1$ is even, and $d = 2$ means $n = 2k + 3 = 4k - 2j + 1$.

3.7.3 Deficit two

Theorem 3.17 (Complete signs for $m = 2$). *Suppose $m = k - j = 2$ and $n > 2k$. Then*

$$\operatorname{sgn} c_{n,k,j} = (-1)^{n-1} \begin{cases} 1, & j \text{ odd}, \\ \operatorname{sgn}(4-d), & j \text{ even}. \end{cases} \quad (3.31)$$

The only zeros in this range have j even and $d = 4$.

Proof. The coefficient of ε^2 in (3.28) is

$$c_{n,k,j} = \frac{(-1)^{n-1} n!}{j!} \int_0^1 t^4 (1-t)^d (2t-1)^j \log \frac{t}{1-t} dt. \quad (3.32)$$

For odd j , the factors $(2t-1)^j$ and $\log(t/(1-t))$ have the same sign, so the integral is strictly positive. For even j , their product is antisymmetric under $t \mapsto 1-t$. Pairing the integral as before leaves, on $1/2 < t < 1$,

$$(2t-1)^j \log \frac{t}{1-t} (t^4(1-t)^d - t^d(1-t)^4).$$

The first two factors are positive there. The bracket has the strict sign of $4 - d$, or is identically zero when $d = 4$. This proves the claim. $\square$

Again the zero condition is exactly reflection: $k = j + 2$ is even and $n = 2k + 5 = 4k - 2j + 1$. For $n \leq 2k$, positivity has already been proved, and the borderline $n = 2k + 1$ is included in both sign arguments. Thus these two theorems, together with Proposition 3.9, classify all zeros for $k - j \leq 2$ throughout the possible triangle.

3.7.4 Why the same sign proof does not settle higher deficits

At $m = 3$, formula (3.28) gives the integral of

$$t^6(1-t)^d(2t-1)^j \left(\frac{1}{2} \log^2 \frac{t}{1-t} - \frac{\pi^2}{6} \right) \quad (3.33)$$

multiplied by $(-1)^{n-1}n!/j!$. The last factor changes sign inside $(0, 1)$. Thus positivity of the weight, or pairing t with $1 - t$, no longer excludes cancellation. This is a concrete obstruction to simply extending the proofs above. It is not a counterexample to the conjecture.

Update (29 September 2026). At $m = 3$ and $m = 4$ the obstruction has since been overcome by a different route: the kernel is written for every deficit as $R_m/m!$ (Remark 3.31), its sign change is controlled by monotone likelihood-ratio comparisons, and a finite certificate finishes the proof (Theorem 3.23). The obstruction as stated remains for $m \geq 5$.

Update (30 September 2026). At $m = 3$ it has been overcome a second time (Theorem 3.59): factoring the root of the falling factorial at the evaluation point leaves a harmonic quantity whose sign changes exactly once in the tail (Lemma 3.54), and a propagation recurrence in the logarithm exponent finishes the proof with a different finite certificate.

Update (1 October 2026, batch 72A). At $m = 5$ and $m = 6$, where the residual polynomial Π_m is quadratic and no longer monotone, it has been overcome by exact lower bounds on the mean of τ^2 and two larger finite certificates (Theorem 3.71). The obstruction as stated remains for $m \geq 7$. [Added 1 October 2026, batch 73: at $m = 7$ and $m = 8$, where Π_m is cubic, it has been overcome in the same way (Theorems 3.99 and 3.100); it remains for $m \geq 9$.]

3.8 Generating functions and equivalent formulas for the upper bound

All identities in this section are proved identities for $U(n)$. They become identities for $a(n)$ only after the remaining nonvanishing assertion is supplied, or when restricted to a certified finite range.

3.8.1 Derivation of the rational generating function

By (3.2), the even and odd subsequences have generating functions

$$E(y) = \sum_{r \geq 0} U(2r)y^r = \frac{(1+y)^2}{(1-y)^3}, \quad (3.34)$$

$$O(y) = \sum_{r \geq 0} U(2r+1)y^r = \frac{2(1+y)}{(1-y)^3} - \frac{y^3}{(1-y)(1-y^4)}. \quad (3.35)$$

For the last term, write $\lfloor (r+1)/4 \rfloor = \sum_{h \geq 1} \mathbf{1}_{r \geq 4h-1}$ and sum each geometric tail. Therefore

$$\begin{aligned} G(t) &:= \sum_{n \geq 0} U(n)t^n = E(t^2) + tO(t^2) \\ &= \frac{1+t^2}{(1-t)^3(1+t)} - \frac{t^7}{(1-t^2)(1-t^8)} \\ &= \frac{N(t)}{(1-t)(1-t^2)(1-t^8)}, \end{aligned} \quad (3.36)$$

where

$$N(t) = 1 + t + 2t^2 + 2t^3 + 2t^4 + 2t^5 + 2t^6 + t^7 + 2t^8 + t^9. \quad (3.37)$$

Indeed, $N(t) = (1+t^2)(1+t+\cdots+t^7) - t^7(1-t)$. This is exactly the rational function proposed in the OEIS entry [2]. The derivation shows why period eight appears: it counts the even values of k on the reflection line, in addition to the parity split of the ordinary triangle.

3.8.2 The eight constituent polynomials

Writing $n = 8q + s$ with $0 \leq s \leq 7$, (3.1) becomes the following degree-two quasipolynomial.

s	$U(8q+s)$
0	$32q^2 + 8q + 1$
1	$32q^2 + 15q + 2$
2	$32q^2 + 24q + 5$
3	$32q^2 + 31q + 8$
4	$32q^2 + 40q + 13$
5	$32q^2 + 47q + 18$
6	$32q^2 + 56q + 25$
7	$32q^2 + 63q + 31$

The trigonometric expression from the OEIS entry is another way to write the same eight polynomials:

$$U(n) = \frac{8n^2 + 15n + 14 + (n+2)(-1)^n + (2 - 4\sqrt{2}\sin(\pi n/4))\sin(\pi n/2)}{16}. \quad (3.38)$$

To verify the equivalence without an approximation, substitute $n = 8q + s$. The sine factors take their exact values at multiples of $\pi/4$ and $\pi/2$. Substitution in each of the eight residue classes yields the polynomials in the table. In particular, no irrational value remains.

3.8.3 Linear recurrences

Multiplication of (3.36) by its denominator yields

$$\begin{aligned} U(n) &= U(n-1) + U(n-2) - U(n-3) + U(n-8) \\ &\quad - U(n-9) - U(n-10) + U(n-11), \quad n \geq 11. \end{aligned} \quad (3.39)$$

Multiplying the denominator by $1+t$ gives the convenient order-twelve form recorded in the OEIS entry:

$$U(n) = 2U(n-2) - U(n-4) + U(n-8) - 2U(n-10) + U(n-12), \quad n \geq 12. \quad (3.40)$$

The twelve initial values are

$$1, 2, 5, 8, 13, 18, 25, 31, 41, 49, 61, 71.$$

Using these recurrences to *generate* the claimed values of $a(n)$ would be circular in a proof of the conjecture. The computations in the next section instead use the independently derived coefficient recurrence (3.7).

3.9 Exact finite verification and certificates

3.9.1 Integer arithmetic and independent formula checks

The accompanying Python program applies (3.7) with arbitrary-precision integers. Through $n = 200$, it checks not only the term count but the entire zero pattern: a coefficient is zero if and only if it belongs to one of the two proved zero families. This examines 1,373,500 cells with $1 \leq k \leq n \leq 200$, $0 \leq j \leq k$.

For $0 \leq n \leq 16$, the program independently recomputes 969 triples (n, k, j) , including the $k = 0$ boundary, using each of (3.12), (3.14), and (3.16). Fractions are exact rational numbers. All three formulas agree with the integer recurrence. The positivity and deficit-one and -deficit-two sign theorems are also checked throughout the integer range. These checks test the derivations and their implementation; the proofs in the preceding sections do not depend on them.

3.9.2 Why modular arithmetic gives a rigorous finite certificate

A nonzero residue modulo any integer $M \geq 2$ proves that the original integer is nonzero. A zero residue, in contrast, might mean only that M divides a nonzero integer. We use two moduli,

$$M_1 = 1,000,000,007, \quad M_2 = 1,000,000,009. \quad (3.41)$$

Their primality is not needed for this argument: only exact reduction of an integer recurrence is used, with no modular division.

For each modulus, a C++ program computes every row through $n = 3000$ and records every zero residue outside the two mathematically proved zero families. There are three such residues modulo M_1 and four modulo M_2 . The two lists are disjoint. Their union and the corresponding residues are as follows.

n	k	j	$c_{n,k,j} \bmod M_1$	$c_{n,k,j} \bmod M_2$
1471	1046	521	847578491	0
1701	610	505	0	439228697
1854	1713	549	0	366891231
2336	1689	145	0	782191592
2543	1785	663	405694396	0
2807	2566	502	607368440	0
2842	2673	2118	474540977	0

Every entry in this table is therefore a nonzero integer coefficient. All other cells outside the proved zero families have a nonzero residue in both runs. Each modulus run checks 4,509,002,501 potential cells, including the single cell at $n = 0$.

Proposition 3.18 (Finite computational certification). *The accompanying exact computations, together with Proposition 3.9 and Theorem 3.11, certify*

$$a(n) = U(n) \quad \text{for every integer } 0 \leq n \leq 3000. \quad (3.42)$$

Certificate argument. The two analytic results prove that every predicted zero is an actual integer zero. For any other coefficient in the finite range, membership in neither, one, or both modular zero lists exhausts the possibilities. The intersection of the lists is empty. Thus at least one of its two residues is nonzero, proving that the integer coefficient is nonzero. Theorem 3.14 now gives the equality for every row in the range. $\square$

This is a deterministic finite verification, not a probabilistic claim based on the size of the moduli. It is also not a formal proof-assistant certificate: it relies on the supplied ordinary programs and their execution, which can be independently reproduced.

3.9.3 Implementation details and limitations

The C++ program stores two coefficient rows in arrays of unsigned 32-bit integers. An update is evaluated in signed 64-bit arithmetic before reduction. Its explicit input bounds guarantee that these intermediates cannot overflow. It never approximates logarithms or transcendental values. The arrays have $O(N^2)$ entries and the computation uses $O(N^3)$ fixed-modulus arithmetic operations through order N .

The Python implementation uses a different update organization: it scatters each existing coefficient into its four successors, whereas the C++ implementation gathers four contributions into each destination. The small-range generating-function and finite-difference calculations provide further independent comparisons. The archive retains both raw modular zero lists, row-by-row support statistics, explicit nonzero witnesses, the combined certificate, and the exact small-range counts.

No finite range, regardless of its size, proves the nonvanishing assertion for all n . In particular, the certificate supplies neither an induction step beyond 3000 nor a theorem bounding where a first additional zero could occur.

3.10 The precise missing step

The coefficient problem can now be stated independently of the original transcendental function.

Conjecture 3.19 (Complete support classification). Let $n \geq 1$, $1 \leq k \leq n$, $0 \leq j \leq k$, and $m = k - j$. Then

$$\frac{1}{j!m!} \frac{d^m}{du^m} \Delta_2^j u^n \Big|_{u=2m} = 0 \quad (3.43)$$

if and only if at least one of the following holds:

$$(j = k \text{ and } n > 2k), \quad (k \text{ even and } n = 4k - 2j + 1). \quad (3.44)$$

The implication from (3.44) to (3.43) is proved above. The converse remains unproved in the following precise region:

$$m = k - j \geq 3, \quad n \geq 2k + 2, \quad (3.45)$$

away from the reflection zeros. All other regions have been handled by the preceding analytic theorems.

Update (29 September 2026). For $m = 3$ and $m = 4$ the converse is now proved, for every n : Theorem 3.23 shows that on those two deficits the only zeros are the reflection zeros. The region still unproved is therefore

$$m = k - j \geq 5, \quad n \geq 2k + 2,$$

away from the reflection zeros; the full conjecture remains open. At every fixed $m \geq 3$ the region is also reduced to finitely many cells by Theorem 3.22, but with bounds too large to compute.

Update (1 October 2026, batch 72A). For $m = 5$ and $m = 6$ the converse is now proved as well (Theorem 3.71). The region still unproved is

$$m = k - j \geq 7, \quad n \geq 2k + 2,$$

away from the reflection zeros, and the full conjecture remains open. [Added 1 October 2026, batch 73: $m = 7$ and $m = 8$ are proved as well (Theorems 3.99 and 3.100); the region still unproved is $m \geq 9$, $n \geq 2k + 2$.] The finite reduction at every fixed $m \geq 3$ now has cutoffs $\lceil (2m+1)e^m \rceil$ and twice that (Theorem 3.81), a rectangle of $2\lceil (2m+1)e^m \rceil^2$ cells — still large: about $5.4 \cdot 10^8$ cells at $m = 7$.

Theorem 3.20 (Exact reduction of the OEIS conjecture). *The equality $a(n) = U(n)$ for every $n \geq 0$ is equivalent to Conjecture 3.19. It is also equivalent to proving the nonvanishing assertion only in the remaining region (3.45), excluding the reflection-zero family.*

Proof. Formula (3.13) identifies the expression in (3.43) with the actual derivative coefficient. Theorem 3.14 counts exactly the cells left after removal of the two proved, disjoint zero families. Hence any additional zero makes the corresponding inequality strict, and no additional zeros makes it an equality. Theorem 3.7, Proposition 3.9, and Theorems 3.16–3.17 eliminate the complementary regions. $\square$

A tempting shortcut would assert that a derivative of a falling factorial cannot vanish at an integer other than its reflection center. That assertion is false. Direct differentiation gives

$$\frac{d^5}{du^5} u^{\underline{8}} = 3360(u-5)(u-2)(2u-7). \quad (3.46)$$

It vanishes at $u = 2$ and $u = 5$, as well as at the reflection center $u = 7/2$. This is *not* a counterexample to Conjecture 3.19, because the derivative order, evaluation point, and finite-difference order in that conjecture are linked in a special way. It does show why those constraints must be used in a genuine proof of the missing implication.

The transformed formula provides an alternative target. With $d = n - 2k - 1 \geq 1$, the outstanding task is to prove that

$$[u^{k+d+1}](1-u)^d(1+u)^{2m} \left(\frac{\operatorname{atanh} u}{u} \right)^m \quad (3.47)$$

is nonzero for $m \geq 3$, except when $d = 2m$ and k is even. The reflection case is transparent because the series then becomes even; the remaining finite-difference cancellations are the essential issue. Equation (3.47) is an equivalent residual lemma, not an additional theorem claimed by this article. (Since the update above it is needed only for $m \geq 5$; since 1 October 2026, Theorem 3.71, only for $m \geq 7$; since batch 73, Theorems 3.99 and 3.100, only for $m \geq 9$.)

3.11 Conclusion

The proposed formula has a rigorous structural explanation. The obvious degree loss removes the top logarithmic coefficients with $n > 2k$. A separate reflection symmetry removes exactly $\lfloor (n+1)/8 \rfloor$ further coefficients for odd n . Their total produces the OEIS floor expression and its rational generating function as an unconditional upper bound.

A fractional-linear transformation proves that a large region of the coefficient triangle is strictly positive, and beta-integral arguments exclude additional zeros on the first two deficit diagonals. These results prove quadratic growth and reduce the full formula to the explicit residual nonvanishing problem (3.47). Exact finite computations settle every derivative order through 3000. The global equality is not proved here: its remaining obligation is the nonvanishing statement in (3.45), not any of the counting or generating-function manipulations. (Update, 29 September 2026: that statement is now proved at deficits $m = 3, 4$, Theorem 3.23; it remains open for $m \geq 5$. Update, 1 October 2026: also at $m = 5, 6$, Theorem 3.71; it remains open for $m \geq 7$. Update, 1 October 2026, batch 73: also at $m = 7, 8$, Theorems 3.99 and 3.100; it remains open for $m \geq 9$.)

3.12 Reproducing the computations

The archive needs Python 3.10 or later and a C++17 compiler for the computations. Python uses only its standard library. The PDF can be rebuilt with a conventional LaTeX installation containing the packages listed in the source preamble.

Run the following commands from the archive directory. The Makefile wraps the same operations.

```
python3 code/verify_exact.py --max-n 200 --formula-n 16
mkdir -p build
g++ -O3 -std=c++17 -Wall -Wextra \
    code/verify_modular.cpp -o build/verify_modular
build/verify_modular 3000 1000000007 data/p1000000007 \
    data/watch_coordinates.csv
build/verify_modular 3000 1000000009 data/p1000000009 \
    data/watch_coordinates.csv
python3 code/combine_certificates.py
pdflatex -interaction=nonstopmode -halt-on-error \
    A290268_partial_results.tex
pdflatex -interaction=nonstopmode -halt-on-error \
    A290268_partial_results.tex
```

The watch file requests explicit residues at the seven exceptional coordinates found in the modular runs. It does not restrict the search: the C++ program checks every coefficient in every row regardless of the watch file. The combiner checks that the complete exceptional zero lists are disjoint and that all listed witnesses are consistent with them.

For a smaller test, the optional watch file can be omitted. For example, `build/verify_modular 100 1000000007 data/test` performs the full modular recurrence through 100. A watch file containing indices beyond the chosen bound is rejected, rather than silently ignored.

3.13 Files included in the archive

File or directory	Purpose
<code>article.tex</code>	Complete source of the merged article; this part is its Part 3.
<code>article.pdf</code>	The compiled merged article.
<code>README.md</code> , <code>Makefile</code>	Status, dependency notes, and reproduction commands.

File or directory	Purpose
<code>code/verify_exact.py</code>	Arbitrary-precision recurrence and three independent coefficient-formula checks.
<code>code/verify_modular.cpp</code>	Complete modular coefficient search with explicit overflow bounds.
<code>code/combine_certificates.py</code>	Verification and combination of the two finite modular certificates.
<code>data/exact_counts.csv</code>	Exact term counts through 200.
<code>data/certified_counts.csv</code>	Certified term counts through 3000.
<code>data/p1000000007_*</code> and <code>data/p1000000009_*</code>	Complete exceptional zero lists, row statistics, witness residues, run logs, and summaries.
<code>data/modular_witnesses.csv</code>	The seven coordinates at which one residue vanishes, with the nonzero residue in the other modulus.
<code>data/*summary.json</code>	Machine-readable verification scope and results.
<code>data/sample_coefficients.json</code>	Fully collected low-order coefficient arrays and cancellation examples.

3.14 Depth certificates: deficits three and four

What this addition is. Subsections 3.14–3.26 were written into this part on 29 September 2026 from a later manuscript, *From Infinite Nonvanishing to Finite Certificates: Bulk positivity, fixed-depth finiteness, and a four-depth solution for OEIS A290268* (batch 42, manuscript 06; provenance in Section 3.26). It closes the region $m = k - j \in \{3, 4\}$ of the precise missing step (3.45): at logarithmic deficits three and four, for every n , the only vanishing coefficients are the reflection zeros of Theorem 3.11. The unproved region is now $m \geq 5$, $n \geq 2k + 2$. The full A290268 conjecture is *not* proved. (*Update, 1 October 2026*: deficits five and six are closed in Subsection 3.35ff., Theorem 3.71; the unproved region is $m \geq 7$, $n \geq 2k + 2$. *Update, batch 73*: deficits seven and eight are closed in Subsection 3.42ff.; the unproved region is $m \geq 9$, $n \geq 2k + 2$.)

Written without this part. The manuscript was prepared against the repository notes `0eis/A290268/README.md` and the conditional Lean reduction, and never read this report. Several of its results therefore re-derive results printed above. They are printed here once more because their proofs are different, and each is marked as a second route with the earlier result credited:

- its all-depth bulk positivity (Theorem 3.21) is Theorem 3.7 restricted to $m \geq 1$;
- its symmetry holes (Theorem 3.29) are the reflection zeros of Theorem 3.11;
- its depths one and two are Theorems 3.16 and 3.17;
- its upper bound (3.84) is Theorem 3.14;
- its exact depth bridge (Proposition 3.24) is Proposition 3.4 re-indexed, and its branch-cut formula (Proposition 3.30) is Lemma 3.15 with its ε -coefficient evaluated in closed form (Remark 3.31).

Its finite rectangles reach only derivative orders $N \leq 359$, inside the range $n \leq 3000$ that Proposition 3.18 already certifies. What is new is the reduction that makes finitely many cells suffice for *every* n ; see Remark 3.42.

Notation. These subsections use the *depth coordinates* of the manuscript, which are also the coordinates of the Lean development. In them k is the **exponent of** $\log x$ — the index called j everywhere else in this part — and d is the logarithmic deficit m , not the d of (3.25). Table 3 is binding for everything from here to the end of the part.

Abstract of manuscript 06. *OEIS A290268 counts the terms in the fully expanded derivatives of x^{x^2} . Its conjectured quasipolynomial reduces to a nonvanishing problem for $\gamma(k, d, M) = [w^M](2 + w)^k((1 + w)^2 \log(1 + w))^d$. The ProveIt investigation identifies symmetry-forced zeros and leaves the higher-depth nonvanishing problem open. We prove positivity throughout the whole bulk $d \leq M \leq k + 2d + 1$, uniformly in every depth $d \geq 1$. We then prove an effective finiteness theorem: at each fixed depth, every possible additional zero is confined to an explicitly bounded finite rectangle in two integer parameters. The proof combines a centered Meixner–Pollaczek factorization, a branch-cut integral, and monotone likelihood-ratio comparisons on a half interval. At depths three and four the bounds sharpen to rectangles of 494 and 15,540 cells. Two independent integer recurrences, supplemented by modular nonzero witnesses, verify these rectangles exactly. Consequently, for $1 \leq d \leq 4$ and $M \geq d$, the only zeros are $M = k + 4d + 1$ with $k + d$ even. This is an infinite-parameter result, not a finite derivative-order experiment. We also bound the number of real zeros of an interpolating Mellin transform by at most d , derive an unconditional term-count lower bound, and formulate concrete next research targets. The full A290268 formula is not proved here: the remaining nonvanishing problem lies at depths $d \geq 5$. The proofs are conventional and computer-assisted where indicated; no Lean kernel certification or independent refereeing is claimed.*

Editorial note. “The ProveIt investigation” is the note `Oeis/A290268/README.md`. The bulk positivity claimed in the abstract was already Theorem 3.7 of this part; the manuscript’s proof of it is a second route.

3.14.1 Notation: depth coordinates

The manuscript writes the N th derivative in the lattice of monomials $x^{x^2+j}(\log x)^k$. That is the lattice of the Lean development (`coeff n j k`, Section 3.26.3) and of the shipped data files, whose columns are named `d`, `k`, `q`. The subsections below keep those letters rather than translate some four hundred lines of indices, and the price is the table below. Its first four rows are the places where a reader coming from the rest of this part will go wrong.

Table 3: Depth coordinates (Subsections 3.14–3.26) against the rest of Part 3. Throughout, $c_{n,k,j}$ is the coefficient of (3.5).

Here	Meaning	Elsewhere in Part 3	Tempting false reading
k	exponent of $\log x$	j	false: the power of $z = x^2$, the k of $c_{n,k,j}$
j	exponent of x in $x^{x^2+j}(\log x)^k$	$2k - n$	false: the exponent of $\log x$
d	depth, $d = (N + j)/2 - k$	$m = k - j$	false: $d = n - 2k - 1$ of (3.25)
q	$M - k - 2d - 1 = N - 2k - 2d - 1$	$d = n - 2k - 1$ of (3.25)	false: the exponent in $(1 - u)^{-q}$ in the proof of Theorem 3.7
N	derivative order	n	—
M	coefficient index $N - k$	$n - j$	the moduli M_1, M_2 of (3.41)
$c_N(j, k)$	coefficient of $x^{x^2+j}(\log x)^k$	$c_{N,(N+j)/2,k}$	—
$\varphi(w)$	$(1 + w)^2 \log(1 + w)$	$B(w)$ of (3.10)	—
$\gamma(k, d, M)$	$[w^M](2 + w)^k \varphi(w)^d$	$[t^{M+k}]A(t)^k B(t)^d$	—
$H_{d,k,q}$	the integer (3.49)	—	$H(u) = \operatorname{atanh}(u)/u$ of (3.15), which is unrelated
r	$M - k$	—	r in $n = 2r$, $n = 2r + 1$
K_d, Q_d	constants (3.74)	—	$A(t)$ of (3.10), $Q_{n,j}$ of
A_d, C_d, E_d			Section 3.3

Two index translations follow from the table and are used repeatedly:

$$c_N(j, k) = c_{N, k+d, k}, \quad N = 2k + 2d + 1 + q \iff n = 2(j_{\text{III}} + m) + 1 + d_{\text{III}},$$

where the subscript III marks the letters of the rest of this part. In particular the tail line $q = 0$ is the borderline $n = 2k + 1$ of Theorem 3.7, and the region $q \geq 1$ is the region $n \geq 2k + 2$ of (3.45).

Renamed symbols. Where the manuscript's letter names a different object of this part, the object is renamed here, with no change of normalization: the manuscript's residual polynomial $P_d(z)$ is $\Pi_d(\zeta)$ (this part's P_n is the canonical polynomial); its centered polynomial $Q_k^{(r)}$ is $\mathcal{Q}_k^{(r)}$; its $A_{M,k}(s)$ is $\mathcal{A}_{M,k}(s)$; its base moment $B_d(r)$ is $\mathcal{B}_d(r)$ and its bulk count $B(N)$ is $\beta(N)$; its half-interval logarithm $L = \log(t/(1-t))$ is τ (this part's L is $\log x$); its hole indicator $h_d(N)$ is $\chi_d(N)$ (the manuscript also used h_d for a second object, kept below as $h_d(\tau)$); its $m = \lfloor (d-1)/2 \rfloor$ is ν_d and the m of its count formulas is ρ (this part's m is the deficit); its Mellin transform $F_{d,k}$ is $\mathcal{F}_{d,k}$ (this part's F_n is the falling factorial); its generating variable z is ξ ; the auxiliary combination $h(t)$ in the proof of Lemma 3.43 is $\psi(t)$; and product indices written j in the manuscript are ℓ . The manuscript's falling factorial $(s)_r$ is printed s^r , as in the rest of this article. The letters s, t, u, w, y are local to the paragraph in which they are introduced; as in the manuscript, y is a centered variable in Subsection 3.15 and $1-t$ in Subsection 3.17, and s is both a polynomial variable and $2t-1$.

3.14.2 Main results of the addition

Put

$$\varphi(w) = (1+w)^2 \log(1+w), \quad \gamma(k, d, M) = [w^M](2+w)^k \varphi(w)^d. \quad (3.48)$$

The depth is d , the logarithmic degree is k , and M is a coefficient index. All three are nonnegative integers.

Theorem 3.21 (Bulk positivity; second route to Theorem 3.7). *For every $d \geq 1$ and $k \geq 0$,*

$$d \leq M \leq k + 2d + 1 \implies \gamma(k, d, M) > 0.$$

In the coordinates of this part the hypothesis reads $m \geq 1, k \leq n \leq 2k + 1$, so the theorem is exactly the case $m \geq 1$ of Theorem 3.7; with (3.57) below, which is Proposition 3.9, it is all of it. The manuscript's proof in Subsection 3.16 goes through a centered factorization instead of the transformation (3.16).

For the remaining region it is useful to put

$$q = M - k - 2d - 1 \geq 0, \quad H_{d,k,q} = \frac{(k + 2d + 1 + q)!}{d!} \gamma(k, d, k + 2d + 1 + q). \quad (3.49)$$

The number $H_{d,k,q}$ is an integer. The $q = 0$ line overlaps the already positive boundary of the bulk.

Theorem 3.22 (Effective fixed-depth finiteness). *For each $d \geq 3$ there are explicit positive integers K_d, Q_d , given in Subsection 3.19, such that whenever $k \geq K_d$ or $q \geq Q_d$,*

$$H_{d,k,q} = 0 \iff q = 2d \text{ and } k + d \equiv 0 \pmod{2}.$$

Thus the complete fixed-depth nonvanishing conjecture is equivalent to a finite, effectively specified collection of integer nonzero tests.

Theorem 3.23 (Complete first-four-depth classification). *For $1 \leq d \leq 4$, $k \geq 0$, and $M \geq d$,*

$$\gamma(k, d, M) = 0 \iff M = k + 4d + 1 \text{ and } k + d \equiv 0 \pmod{2}. \quad (3.50)$$

For $d = 1, 2$ the proof is entirely analytic. For $d = 3, 4$ it uses finite exact-integer certificates after a proved infinite-to-finite reduction.

In the coordinates of this part: for $1 \leq m \leq 4$ and every $n \geq k$, the coefficient $c_{n,k,j}$ vanishes if and only if k is even and $n = 4k - 2j + 1$. For $m = 1, 2$ this is Theorems 3.7, 3.16 and 3.17; for $m = 3, 4$ it is new, and it removes $m = 3, 4$ from the region (3.45).

The practical exceptional rectangles for this theorem are

$$\begin{aligned} d = 3 : \quad & 0 \leq k \leq 12, \quad 0 \leq q \leq 37, \\ d = 4 : \quad & 0 \leq k \leq 83, \quad 0 \leq q \leq 184. \end{aligned} \quad (3.51)$$

They contain 48 symmetry zeros and 15,986 nonzero entries in total. Subsection 3.21 explains every finite obligation and its verification. Subsection 3.22 gives a separate all-depth bound on the number of zeros for each fixed k .

3.14.3 What is and is not new here

The manuscript's own assessment, relative to the repository notes it read: the support reduction, the target quasipolynomial, the reflection idea, and the first two depths are prior repository material [18]. Meixner–Pollaczek polynomials are classical [20]; no new orthogonal-polynomial family is proposed. The contribution is their use to prove all-depth bulk positivity, the effective uniform fixed-depth reduction, and its rigorous completion at depths three and four. The real-zero bound and term-count lower bound are additional consequences.

Relative to this part the assessment is narrower. All-depth bulk positivity is Theorem 3.7, and the reflection zeros, the deficit-one and deficit-two classification and the upper bound are all proved above; the manuscript's proofs of them are second routes. New relative to this part are: the centered factorization (Proposition 3.26); the closed form of the branch-cut kernel for every depth and its folding (Subsection 3.17); the monotone comparisons and the effective finite reduction at every fixed depth $d \geq 3$ (Theorem 3.22); the completion at deficits three and four (Theorem 3.23); the zero-count bound (Theorem 3.44); and the lower bound of Corollary 3.45, which improves $\Lambda(n)$ of Theorem 3.1.

The manuscript's literature and repository checks support it as a continuation beyond the inspected A290268 notes, not an assertion of absolute historical priority. Related results on generalized Bernoulli polynomials, including integer interlacing in the order-equals-degree family, appear in Štampach's work [21]. Those results concern a different parameter regime and do not by themselves settle the coefficient nonvanishing considered here. The addition is a research draft with explicit proof and computation boundaries, not an announcement that the entire OEIS conjecture has been solved.

3.15 Depth coordinates: exact coefficient and polynomial models

3.15.1 The lattice recurrence and the depth reduction

For $x > 0$, write

$$\frac{d^N}{dx^N} x^{x^2} = x^{x^2} \sum_{j \in \mathbb{Z}} \sum_{k \geq 0} c_N(j, k) x^j (\log x)^k. \quad (3.52)$$

Only finitely many coefficients are nonzero, and all are integers. The sequence in question is $a(N) = \#\{(j, k) : c_N(j, k) \neq 0\}$. The OEIS entry, proposed by Reshetnikov in 2017 and subsequently given alternative forms by Luschny, records the conjecture

$$a(2\rho) = 2\rho^2 + 2\rho + 1, \quad a(2\rho - 1) = 2\rho^2 - \lfloor \rho/4 \rfloor \quad (\rho \geq 1), \quad (3.53)$$

which is (3.2) with $\rho = r + 1$ in the odd case. The equivalent rational generating function remained labeled conjectural in the entry accessed for the manuscript on 29 September 2026 [2]. The precise coefficient model is used rather than the output conventions of any computer algebra system.

The functions $x^j(\log x)^k$ appearing in a finite expansion are linearly independent; this is Lemma 3.2, and the manuscript's argument is the same one. After setting $x = e^t$, a putative relation becomes a finite sum of exponentials $\exp(jt)$ times polynomials in t . Divide by the largest exponential and let $t \rightarrow +\infty$; the corresponding polynomial must vanish. Repeat. Thus the coefficient support is an intrinsic object within this normal form.

The repository notes already reduce the support problem to a three-parameter coefficient and prove or discuss several initial strata [18]. The Lean module `Main.lean` explicitly consumes the support characterization as a hypothesis in its support-to-count theorem [19]; that conditional theorem is not treated as an existing proof of the conjecture (Section 3.26.3).

Differentiating a basis term in (3.52) yields

$$\begin{aligned} c_{N+1}(j, k) = & 2c_N(j-1, k-1) + c_N(j-1, k) \\ & + (j+1)c_N(j+1, k) + (k+1)c_N(j+1, k+1), \end{aligned} \quad (3.54)$$

with $c_0(0, 0) = 1$ and all other initial coefficients zero. Negative logarithmic indices are interpreted as zero. This is the recurrence (3.7) re-indexed, and it is verbatim the definition of `coeff` in the Lean development. In particular,

$$|j| \leq N, \quad j \equiv N \pmod{2}, \quad 0 \leq k \leq (N+j)/2$$

are necessary for a nonzero coefficient.

Proposition 3.24 (Exact depth bridge). *For an integer pair (j, k) in this support window, set $d = (N+j)/2 - k$ and $M = N - k$. Then*

$$c_N(j, k) = \frac{N!}{k!d!} \gamma(k, d, M). \quad (3.55)$$

If $d \geq 1$ and $j \leq -1$, so that $q = -j - 1 \geq 0$, this becomes the integer identity

$$c_N(j, k) = \binom{N}{k} H_{d,k,q}, \quad N = 2k + 2d + 1 + q. \quad (3.56)$$

Proof. Taylor expansion gives

$$\sum_{N \geq 0} \frac{\xi^N}{N!} x^{-x^2} \frac{d^N}{dx^N} x^{x^2} = \exp((2x\xi + \xi^2) \log x) \exp((x + \xi)^2 \log(1 + \xi/x)).$$

Set $w = \xi/x$. The contribution of logarithmic degree k in the first exponential and degree d in the second is

$$\frac{x^{2k+2d}}{k!d!} w^k (2+w)^k \varphi(w)^d (\log x)^k.$$

Extracting ξ^N introduces x^{-N} , so $j = 2k + 2d - N$ and the remaining coefficient index is $N - k = M$. This proves (3.55). Substitution of (3.49) gives (3.56). $\square$

Since $A(t)^k B(t)^d = t^k (2+t)^k \varphi(t)^d$, the bridge (3.55) is (3.12) re-indexed, and the proof above is essentially the Taylor-expansion proof of Proposition 3.4.

For completeness,

$$\gamma(k, 0, M) = \begin{cases} 2^{k-M} \binom{k}{M}, & 0 \leq M \leq k, \\ 0, & M > k, \end{cases} \quad (3.57)$$

which is Proposition 3.9. Also $\gamma(k, d, M) = 0$ for $M < d$, since φ has order one at the origin.

3.15.2 A parameter derivative

Define

$$\mathcal{A}_{M,k}(s) = [w^M](2+w)^k(1+w)^s.$$

This is a polynomial in s . Differentiation with respect to s gives

$$\gamma(k, d, M) = \left. \frac{\partial^d}{\partial s^d} \mathcal{A}_{M,k}(s) \right|_{s=2d}. \quad (3.58)$$

Thus the question is about Taylor coefficients of explicit polynomials, not the cancellation of numerical transcendental values. (This is the depth-coordinate form of the finite-difference formula (3.13).)

Definition 3.25 (Centered polynomial). For an integer $r \geq 0$, define $\mathcal{Q}_k^{(r)}(y)$ by

$$\sum_{k \geq 0} \frac{\mathcal{Q}_k^{(r)}(y)}{k!} \xi^k = (1 - \xi^2)^{-(r+1)/2} \exp(2y \operatorname{atanh} \xi). \quad (3.59)$$

Proposition 3.26 (Centered factorization). If $M \geq k$ and $r = M - k$, then

$$M! \mathcal{A}_{M,k}(s) = s^r \mathcal{Q}_k^{(r)}\left(s - \frac{r-1}{2}\right). \quad (3.60)$$

The polynomial $\mathcal{Q}_k^{(r)}$ has degree k , leading coefficient 2^k , nonnegative coefficients, and parity k . It satisfies

$$\mathcal{Q}_0^{(r)} = 1, \quad \mathcal{Q}_1^{(r)} = 2y, \quad \mathcal{Q}_{k+1}^{(r)} = 2y \mathcal{Q}_k^{(r)} + k(k+r) \mathcal{Q}_{k-1}^{(r)} \quad (k \geq 1). \quad (3.61)$$

Proof. Expanding $(2+w)^k$ and factoring the common falling factorial gives

$$M! \mathcal{A}_{M,k}(s) = s^r \sum_{\ell=0}^k \binom{k}{\ell} 2^\ell \frac{(r+k)!}{(r+\ell)!} (s-r)^\ell.$$

The sum divided by $k!$ is

$$[t^k](1+t)^{r+k}(1+2t)^{s-r}.$$

Under the formal residue substitution $t = \xi/(1-\xi)$, this coefficient is

$$[\xi^k](1+\xi)^{s-r}(1-\xi)^{-s-1}.$$

With $y = s - (r-1)/2$, the last generating function is exactly (3.59). This proves the factorization. Both $(1-\xi^2)^{-(r+1)/2}$ and $\operatorname{atanh} \xi = \xi + \xi^3/3 + \dots$ have nonnegative coefficients. Expansion of the exponential proves coefficient nonnegativity and parity, and its top y -degree term gives $2^k y^k$. Finally, if G is the generating function in (3.59), then

$$(1-\xi^2)G'(\xi) = (2y + (r+1)\xi)G(\xi).$$

Coefficient comparison gives (3.61). □

In the standard normalization of the classical Meixner–Pollaczek polynomials $P_k^{(\lambda)}(x; \phi)$ (classical notation, unrelated to the P_n of this part), the generating function in [20, Eq. 18.23.7] gives

$$\mathcal{Q}_k^{(r)}(y) = k! i^k P_k^{((r+1)/2)}(-iy; \pi/2). \quad (3.62)$$

All uses of this family in the proof follow directly from Proposition 3.26, so no orthogonality theorem is an unstated dependency.

Corollary 3.27 (Mixed real and imaginary zeros). *The zeros of $\mathcal{Q}_k^{(r)}$ are simple and purely imaginary. More precisely, for positive numbers $\eta_1, \dots, \eta_{\lfloor k/2 \rfloor}$,*

$$\mathcal{Q}_k^{(r)}(y) = 2^k y^{k \bmod 2} \prod_{\ell=1}^{\lfloor k/2 \rfloor} (y^2 + \eta_\ell^2).$$

Thus (3.60) separates the consecutive real roots of the falling factorial from a centered imaginary-root factor.

Proof. The monic polynomials $p_k(x) = 2^{-k} i^{-k} \mathcal{Q}_k^{(r)}(ix)$ satisfy

$$p_{k+1}(x) = x p_k(x) - \frac{k(k+r)}{4} p_{k-1}(x).$$

They are the characteristic polynomials of real symmetric tridiagonal matrices with zero diagonal and positive off-diagonal entries $\sqrt{\ell(\ell+r)}/2$. The eigenvalues are real. They are simple because an eigenvector is determined by its first coordinate through the tridiagonal recurrence, and a zero first coordinate would force the entire vector to vanish. The zero diagonal and alternating sign conjugation make the spectrum symmetric about zero. Transform back to $\mathcal{Q}_k^{(r)}$. $\square$

3.16 Depth coordinates: bulk positivity and symmetry holes

Both results of this subsection were proved earlier in this part, by other means: the bulk is Theorem 3.7 and the holes are Theorem 3.11. The manuscript's proofs are printed as second routes.

Second proof of Theorem 3.7 for $m \geq 1$ (manuscript 06): proof of Theorem 3.21. First suppose $M \geq k$ and put $r = M - k$. The stated range implies $r \leq 2d + 1$. In (3.60), set $s = 2d + u$. Every factor of

$$(2d + u)^x = \prod_{\ell=0}^{r-1} (u + 2d - \ell)$$

has nonnegative constant term. At most one has constant term zero. If no factor vanishes at $u = 0$, all coefficients through degree r are positive; if one vanishes, all coefficients of degrees 1 through r are positive. The empty product is one.

The argument of $\mathcal{Q}_k^{(r)}$ is $u + y_0$ with $y_0 = 2d - (r - 1)/2 \geq d > 0$. Its coefficients through degree k are all positive: the leading term $2^k(u + y_0)^k$ already has that property, and all remaining contributions are nonnegative. Since $1 \leq d \leq r + k = M$, the coefficient of u^d in the product is strictly positive. Equation (3.58) proves the claim in this case.

To handle $M < k$, induct on k for the entire claimed range, using

$$\gamma(k, d, M) = 2\gamma(k - 1, d, M) + \gamma(k - 1, d, M - 1). \quad (3.63)$$

The $k = 0$ case has already been proved. When $M < k$ and $M \geq d$, the first term on the right is positive by induction. The second is positive when $M - 1 \geq d$ and is zero otherwise. Both indices lie in the preceding bulk range. This completes the induction. $\square$

Corollary 3.28 (Positive derivative columns). *Within the structural support window, every coefficient with $j \geq 0$ is positive. In the $j = -1$ column every coefficient with $d \geq 1$ is positive, while the $d = 0$ coefficient is zero.*

Proof. For $d \geq 1$, the condition $M \leq k + 2d + 1$ is exactly $j \geq -1$ by Proposition 3.24. For $d = 0$, use (3.57); its nonzero condition $M \leq k$ is exactly $j \geq 0$. $\square$

In the coordinates of this part, $j \geq 0$ is $n \leq 2k$ and $j = -1$ is $n = 2k + 1$, so the corollary restates Theorem 3.7 together with Proposition 3.9.

Theorem 3.29 (All-depth symmetry holes; second route to Theorem 3.11). *For $d \geq 1$ and $k \geq 0$,*

$$k + d \equiv 0 \pmod{2} \implies \gamma(k, d, k + 4d + 1) = 0.$$

Proof. Here $r = 4d + 1$ and the center $(r - 1)/2$ is $2d$. With $s = 2d + u$,

$$s^r = u \prod_{\ell=1}^{2d} (u^2 - \ell^2)$$

is odd, whereas $\mathcal{Q}_k^{(r)}(u)$ has parity k . The product has parity $k + 1$. Its u^d coefficient therefore vanishes if $k + d$ is even. Apply (3.58). $\square$

This proves the hole direction without relying on the analytic integral below. The polynomial proof also distinguishes the symmetry statement from the separate assertion that there are no other zeros. Translated, $M = k + 4d + 1$ is $q = 2d$, that is $n = 2k + 2m + 1 = 4k - 2j + 1$ in this part's letters, and $k + d$ even is “ k even” there; so the hole family is exactly the reflection family of Theorem 3.11, and the proof above is the odd-falling-factorial parity argument that Section 5.4 identifies as common to all three parts.

3.17 Depth coordinates: the branch-cut integral and its folding

3.17.1 An exact integral valid in the negative region

Let

$$R_d(v) = \frac{\operatorname{Im}(v + i\pi)^d}{\pi}. \quad (3.64)$$

It is a real polynomial of degree $d - 1$ with leading coefficient d .

Proposition 3.30 (Branch-cut formula). *For $d \geq 1$, $k \geq 0$, and $M = k + 2d + 1 + q$ with integer $q \geq 0$,*

$$\gamma(k, d, M) = (-1)^{M+1} \int_0^1 t^q (1-t)^{2d} (2t-1)^k R_d\left(\log \frac{1-t}{t}\right) dt. \quad (3.65)$$

The integral is absolutely convergent.

Proof. Apply Cauchy's coefficient formula to $(2+w)^k(1+w)^{2d}\log^d(1+w)$, using the principal logarithm and deforming the circle to the cut $(-\infty, -1]$. The small circle about -1 gives zero in the limit because its contribution is bounded by a constant times $\varepsilon^{2d+1}(1+|\log \varepsilon|^d)$. The large circle vanishes because $M > k + 2d$, with bound $O(R^{k+2d-M}\log^d R)$.

At $w = -x$, $x > 1$, the upper and lower logarithms are $\log(x-1) \pm i\pi$. Their jump divided by $2\pi i$ is $R_d(\log(x-1))$. The factor w^{-M-1} contributes $(-1)^{M+1}$. Consequently the coefficient is

$$(-1)^{M+1} \int_1^\infty \frac{(2-x)^k(x-1)^{2d}}{x^{M+1}} R_d(\log(x-1)) dx.$$

The substitution $t = 1/x$ gives (3.65). At $t = 0$ and $t = 1$ the integrand is bounded by a constant times a nonnegative integral power of the distance to the endpoint, multiplied by a fixed power of its logarithm. This is integrable. $\square$

Remark 3.31 (Relation to Lemma 3.15; editorial). Formula (3.65) is the ε -expansion of Lemma 3.15 carried out for every deficit. Indeed $\frac{\sin(\pi\varepsilon)}{\pi} \left(\frac{t}{1-t}\right)^\varepsilon = \text{Im exp}((\sigma + i\pi)\varepsilon)/\pi$ with $\sigma = \log \frac{t}{1-t}$, whose coefficient of ε^m is $R_m(\sigma)/m!$. Hence (3.28) reads

$$c_{n,k,j} = \frac{(-1)^{n-1}n!}{j!m!} \int_0^1 t^{2m}(1-t)^d(2t-1)^j R_m\left(\log \frac{t}{1-t}\right) dt$$

in the letters of this part, and the substitution $t \mapsto 1-t$ together with the dictionary of Table 3 turns it into (3.56) with (3.65). For $m = 1, 2, 3$ the kernel $R_m/m!$ is 1 , σ and $\frac{1}{2}\sigma^2 - \frac{\pi^2}{6}$: the integrands (3.30), (3.32) and (3.33). What is new below is the treatment of the sign change that Section 3.7 identified as the obstruction at $m = 3$.

3.17.2 The residual polynomial

For $d \geq 1$, put

$$\nu_d = \left\lfloor \frac{d-1}{2} \right\rfloor, \quad h_d(\tau) = \begin{cases} 1, & d \text{ odd,} \\ \tau, & d \text{ even,} \end{cases}$$

and define the monic polynomial

$$\Pi_d(\zeta) = \prod_{\ell=1}^{\nu_d} \left(\zeta - \pi^2 \cot^2 \frac{\ell\pi}{d} \right). \quad (3.66)$$

The empty product is one. Factoring the imaginary part in (3.64) gives

$$R_d(-\tau) = (-1)^{d-1} d h_d(\tau) \Pi_d(\tau^2) \quad (\tau > 0). \quad (3.67)$$

Indeed, the roots of R_d are the distinct numbers $\pi \cot(\ell\pi/d)$ for $1 \leq \ell \leq d-1$; pairing opposite roots proves the identity, including its leading coefficient.

The initial cases are

$$\begin{aligned} \Pi_1(\zeta) &= \Pi_2(\zeta) = 1, \\ \Pi_3(\zeta) &= \zeta - \pi^2/3, \quad \Pi_4(\zeta) = \zeta - \pi^2, \\ \Pi_5(\zeta) &= \zeta^2 - 2\pi^2\zeta + \pi^4/5, \\ \Pi_6(\zeta) &= \zeta^2 - \frac{10}{3}\pi^2\zeta + \pi^4. \end{aligned} \quad (3.68)$$

Thus the first two depths have no residual sign change. At depths three and four the residual polynomial is increasing. At higher depths that last property fails and cannot be silently reused.

3.17.3 Positive measures on the half interval

For $1/2 < t < 1$, abbreviate

$$y = 1 - t, \quad s = 2t - 1, \quad \tau = \log(t/y), \quad p = 2d, \quad \delta = q - p, \quad b_d(t) = (ty)^p h_d(\tau).$$

Reflect the part $0 < t < 1/2$ in (3.65). Apart from the sign and positive factor $(-1)^{d-1}d$, the result is

$$\int_{1/2}^1 b_d(t) s^k \Pi_d(\tau^2) (t^\delta + (-1)^{k+d-1} y^\delta) dt. \quad (3.69)$$

When $k + d$ is odd, the bracket is the positive sum $S_\delta(t) = t^\delta + y^\delta$. When $k + d$ is even, it is the difference $D_\delta(t) = t^\delta - y^\delta$; it is identically zero for $\delta = 0$ and otherwise has constant sign $\operatorname{sgn} \delta$.

For a sum case, define a probability measure by normalizing $b_d s^k S_\delta dt$. For a nonzero difference case, normalize $b_d s^k |D_\delta| dt$. Write $\mathcal{E}_{d,k,q}$ for expectation with respect to the relevant measure. The following sign rule is exact:

$$\operatorname{sgn} H_{d,k,q} = \begin{cases} (-1)^q \operatorname{sgn} \mathcal{E}_{d,k,q}[\Pi_d(\tau^2)], & k + d \text{ odd}, \\ (-1)^{q+1} \operatorname{sgn}(q - 2d) \operatorname{sgn} \mathcal{E}_{d,k,q}[\Pi_d(\tau^2)], & k + d \text{ even, } q \neq 2d. \end{cases} \quad (3.70)$$

At a symmetry hole $H_{d,k,2d} = 0$ by Theorem 3.29. All the measures above have finite logarithmic moments of every fixed order.

3.18 Depth coordinates: monotone comparisons

Lemma 3.32 (Covariance comparison). *Let μ be a probability measure on an interval, and let f, g be nondecreasing functions with the needed integrability, $g > 0$. Then*

$$\frac{\int f g d\mu}{\int g d\mu} \geq \int f d\mu.$$

In particular, multiplying a positive density by a nondecreasing function can only increase expectations of nondecreasing functions.

Proof. The covariance equals

$$\frac{1}{2} \iint (f(u) - f(v))(g(u) - g(v)) d\mu(u) d\mu(v) \geq 0.$$

Truncation proves the same conclusion for integrable unbounded functions. $\square$

Lemma 3.33 (Radial likelihood ratios). *On $(1/2, 1)$ the following assertions hold.*

- (i) *For every integer δ , S_δ is nondecreasing. The cases $\delta = 0, 1$ are constant.*
- (ii) *For every nonzero integer δ , $|D_\delta|/s$ is positive and nondecreasing. The positive cases $\delta = 1, 2$ are constant.*
- (iii) *If $\delta \geq 1$ is an integer, both $S_{\delta+1}/S_\delta$ and $D_{\delta+1}/D_\delta$ are nondecreasing.*

Proof. Write $v = \frac{1}{2} \log(t/y) > 0$, so $t = e^v/(2 \cosh v)$ and $y = e^{-v}/(2 \cosh v)$. Direct calculation gives

$$\begin{aligned} \frac{S_{\delta+1}}{S_\delta} &= \frac{1}{2} (1 + \tanh v \tanh(\delta v)), \\ \frac{D_{\delta+1}}{D_\delta} &= \frac{1}{2} (1 + \tanh v \coth(\delta v)). \end{aligned} \quad (3.71)$$

The first expression is nondecreasing for $\delta \geq 0$. For $\delta \geq 1$,

$$\frac{d}{dv} \log(\tanh v \coth(\delta v)) = \frac{2}{\sinh(2v)} - \frac{2\delta}{\sinh(2\delta v)} \geq 0,$$

using $\sinh(\delta x) \geq \delta \sinh x$ for $\delta \geq 1$, $x > 0$. This proves (iii). Products of these ratios, starting at $S_1 = 1$ and $D_1 = s$, prove (i) and (ii) for positive δ . For a negative integer $\delta = -a$, use

$$S_{-a} = (ty)^{-a} S_a, \quad |D_{-a}| = (ty)^{-a} D_a.$$

The extra factor is nondecreasing because ty decreases on the half interval. The stated constant cases are immediate. $\square$

For real $r \geq 0$, let $\mu_{d,r}$ denote the probability measure obtained by normalizing $b_d(t)s^r dt$, and set

$$\mathcal{B}_d(r) = \mathbb{E}_{\mu_{d,r}}[\tau^2]. \quad (3.72)$$

Proposition 3.34 (Uniform comparison with a symmetric base). *For every integer $q \geq 0$ outside a symmetry hole,*

$$\begin{aligned} \mathcal{E}_{d,k,q}[\tau^2] &\geq \mathcal{B}_d(k), & k+d \text{ odd}, \\ \mathcal{E}_{d,k,q}[\tau^2] &\geq \mathcal{B}_d(k+1), & k+d \text{ even}. \end{aligned} \quad (3.73)$$

The function $\mathcal{B}_d(r)$ is nondecreasing in r . Within either parity class of k , the expectations of every increasing function of τ^2 increase when k increases by two. They also increase when q increases by one in the region $q \geq 2d+1$.

Proof. For a sum measure, its ratio to the base density with exponent k is S_δ , which is nondecreasing by Lemma 3.33. For a difference measure, its ratio to the base with exponent $k+1$ is $|D_\delta|/s$, again nondecreasing. Apply Lemma 3.32. Increasing r multiplies a base density by a positive power of s . Increasing k by two multiplies either type of density by s^2 . The assertion about q follows from the adjacent ratios in (3.71), with $\delta \geq 1$. $\square$

Remark 3.35 (The role of integer parameters). The global base comparison uses integer δ . For noninteger $0 < \delta < 1$, the sum S_δ has different monotonicity. No conclusion about that interval is being smuggled into the integer coefficient problem. The real-zero theorem in Subsection 3.22 uses a separate argument and does not depend on the integer comparison.

3.19 Depth coordinates: an explicit finite reduction at every depth

This subsection proves Theorem 3.22 with deliberately crude but fully explicit bounds. Their purpose is decidability of a fixed-depth stratum, not efficient computation. The next subsection replaces them by small bounds at depths three and four.

3.19.1 A monotone lower bound for every residual polynomial

For $d \geq 3$, put

$$\begin{aligned} \nu_d &= \lfloor (d-1)/2 \rfloor, & C_d &= [3(d^2+1)]^{\nu_d} + d^2 + 1, \\ A_d &= 4C_d + 2, & E_d &= (2d+1)A_d + 6d + 2, \\ K_d &= 2^{A_d}(E_d + A_d + 2), \\ Q_d &= 2d + 2^{A_d+1}(E_d + A_d + 6). \end{aligned} \quad (3.74)$$

These are positive integers constructed by elementary integer operations and exponentiation.

Lemma 3.36 (Affine minorant). *For all $\zeta \geq 0$,*

$$\Pi_d(\zeta) \geq \frac{1}{2}\zeta - C_d. \quad (3.75)$$

Proof. Every root β of Π_d satisfies $0 < \beta < d^2$, because $\pi \cot(\pi/d) < d$. For $\zeta \geq 2(d^2 + 1)$, each factor $\zeta - \beta$ is at least $\zeta/2$, and $\nu_d \geq 1$ implies $\Pi_d(\zeta) \geq (\zeta/2)^{\nu_d} \geq \zeta/2$. For $0 \leq \zeta \leq 2(d^2 + 1)$, the absolute value of the product is at most $[3(d^2 + 1)]^{\nu_d}$. Meanwhile $\zeta/2 - C_d$ is at most the negative of that quantity. This proves the inequality on both ranges. $\square$

3.19.2 Explicit concentration of the base measure

Write $A = A_d$, $E = E_d$, and choose rational points

$$t_0 = 1 - 2^{-A}, \quad t_1 = 1 - 2^{-A-1}, \quad t_2 = 1 - 2^{-A-2}.$$

On $t \geq t_0$,

$$\tau^2 \geq \log^2(2^A - 1) > \frac{(A-1)^2}{4} > 4C_d, \quad (3.76)$$

where $\log 2 > 1/2$ and $A = 4C_d + 2$ were used. Below t_0 , the base factor b_d has integral at most A : use $(t(1-t))^{2d} \leq 1$ and, when d is even, $\tau < A$. On $[t_1, t_2]$, the interval length is 2^{-A-2} , $t(1-t) \geq 2^{-A-3}$, and $h_d(\tau) \geq 1$. Therefore

$$\int_{t_1}^{t_2} b_d(t) dt \geq 2^{-E}. \quad (3.77)$$

Lemma 3.37 (Large base exponent). *If $r \geq K_d$, then $\mathcal{B}_d(r) > 2C_d$.*

Proof. Let $s_0 = 2t_0 - 1 = 1 - 2^{1-A}$ and $s_1 = 2t_1 - 1 = 1 - 2^{-A}$. The ratio of the unnormalized mass below t_0 to the mass on $[t_1, t_2]$ is at most

$$A2^E(s_0/s_1)^r \leq A2^E \exp(-2^{-A}r) < 1.$$

For the last inequality, substitute $r \geq 2^A(E + A + 2)$ and use $\log A < A$, $\log 2 < 1$. Thus the normalized mass above t_0 exceeds one half. Equation (3.76) gives $\mathcal{B}_d(r) > 2C_d$. $\square$

3.19.3 Explicit concentration in the tail parameter

Lemma 3.38 (Large coefficient tail). *If $q \geq Q_d$, then in both parity types and for every $k \geq 0$,*

$$\mathcal{E}_{d,k,q}[\tau^2] > 2C_d.$$

Proof. Within each parity type it suffices, by Proposition 3.34, to use its smallest possible k , either zero or one. Put $\delta = q - 2d \geq 1$. On the part below t_0 , the bracket S_δ or D_δ is at most $2t_0^\delta$, so the unnormalized mass is at most $2At_0^\delta$. On $[t_1, t_2]$, the factor s^k is at least $1/2$. Both brackets are at least $t_1^\delta/2$: for the difference use $D_\delta = t^\delta(1 - (y/t)^\delta)$ and $y/t < 1/2$ there. The interval mass is consequently at least $2^{-E}t_1^\delta/4$. The ratio of the bad mass to this interval mass is at most

$$8A2^E(t_0/t_1)^\delta \leq 8A2^E \exp(-2^{-A-1}\delta) < 1,$$

by $\delta \geq 2^{A+1}(E + A + 6)$. Once again, more than half of the probability lies above t_0 , and (3.76) finishes the proof. $\square$

Proof of Theorem 3.22. If $k \geq K_d$, the base comparison and Lemma 3.37 imply $\mathcal{E}_{d,k,q}[\tau^2] > 2C_d$ for every nonhole $q \geq 0$. If $q \geq Q_d$, Lemma 3.38 gives the same conclusion for every k . In either case, Lemma 3.36 gives

$$\mathcal{E}_{d,k,q}[\Pi_d(\tau^2)] \geq \frac{1}{2}\mathcal{E}_{d,k,q}[\tau^2] - C_d > 0.$$

The sign rule (3.70) proves nonvanishing outside the stated symmetry holes. The holes themselves were proved independently. Combined with bulk positivity and the structural zeros, this confines every possible extra zero to $0 \leq k < K_d$, $0 \leq q < Q_d$. $\square$

Corollary 3.39 (A terminating fixed-depth decision algorithm). *Given $d \geq 1$, there is an algorithm that decides whether (3.50) holds for every pair of integer parameters (k, M) at that depth. For $d \geq 3$, it computes the finite rectangle specified by (3.74) using integer polynomial recurrences. If the answer is negative, it returns an exact counterexample.*

Remark 3.40 (The bound depends on the depth). The bound is uniform in both k and q , but depends on d . It is not a finite verification of the full conjecture over all depths. Nor do the enormous universal constants justify a practical exhaustive computation. Their value is the proved reduction; efficient certificates need sharper analysis, as the next subsection demonstrates.

3.20 Depth coordinates: completing depths one through four

3.20.1 The first two depths

For $d = 1, 2$, the residual polynomial Π_d is identically one. Every positive-weight expectation in (3.70) is strictly positive. Together with bulk positivity and the symmetry theorem, this proves (3.50) at the first two depths. In particular, no higher Peano-kernel argument is needed for depth two. (This is a second route to the zero classification of Theorems 3.16 and 3.17, which prove more: the complete sign pattern. The manuscript's folding (3.69) is the pairing $t \leftrightarrow 1 - t$ used in their proofs.)

3.20.2 A small-certificate criterion for depths three and four

At these depths, $\Pi_d(\tau^2)$ is an increasing affine function of τ^2 . The comparisons in Proposition 3.34 apply to its expectation itself, not merely to a lower bound.

Proposition 3.41 (Three signs and a finite rectangle). *Let $d \in \{3, 4\}$. Choose $R \geq 2$ of parity $d - 1$ and an integer $Q \geq 2d + 1$. Put $k_s = (d - 1) \bmod 2$ and $k_a = d \bmod 2$. Suppose*

$$H_{d,R,2d} > 0, \quad (-1)^Q H_{d,k_s,Q} > 0, \quad (-1)^{Q+1} H_{d,k_a,Q} > 0. \quad (3.78)$$

Then every nonhole entry with $k \geq R - 1$ or $q \geq Q$ is nonzero. Consequently the full depth is settled by testing $0 \leq k \leq R - 2$, $0 \leq q \leq Q - 1$.

Proof. Since $R + d$ is odd and $2d$ is even, the first sign in (3.78) says exactly that the base expectation $\mathbb{E}_{\mu_{d,R}} \Pi_d(\tau^2)$ is positive. In a sum case with $k \geq R - 1$, parity forces $k \geq R$. In a difference case the comparison base has exponent $k + 1 \geq R$. Thus all nonhole entries in this k -range have positive reduced expectation.

The remaining two signs are exactly positive reduced expectations for the smallest sum and difference parity classes at $q = Q$; here $Q > 2d$. Increasing k by two and increasing q by one preserves that positivity by Proposition 3.34. This covers every $q \geq Q$. $\square$

For $d = 3$, take $(R, Q) = (14, 38)$. For $d = 4$, take $(R, Q) = (85, 185)$. The six seed signs and the two rectangles are verified in the next subsection. This proves Theorem 3.23.

3.20.3 What the result says about individual coefficients

At depth three the only holes have

$$q = 6, \quad k \text{ odd}; \quad j = -7, \quad N = 2k + 13.$$

At depth four they have

$$q = 8, \quad k \text{ even}; \quad j = -9, \quad N = 2k + 17.$$

(In this part's letters: $m = 3$, j odd, $n = 2j + 13$; and $m = 4$, j even, $n = 2j + 17$ — in both cases $k = j + m$ even and $n = 4k - 2j + 1$.) All other entries at those depths are nonzero, regardless of derivative order. Above the practical k or q thresholds their signs are the explicit signs in (3.70) with the reduced expectation replaced by a positive number.

Nonzero does not mean positive. For example, in the $d = 3, k = 0$ row the reduced expectation is positive at $q = 0$, negative for $1 \leq q \leq 37$, and positive for all $q \geq 38$. These sign changes explain why a naive one-sign induction can fail without producing an integer zero. The exact finite sign data are shipped as `data/02-depth-certificates-sign_thresholds.csv`. *Update (30 September 2026)*: the sign of the $d = 3, k = 0$ row is that of the harmonic quantity E_q , so this observation is now proved (Lemma 3.54); the second route of Subsection 3.31 reproduces every $d = 3$ row of that file (Remark 3.61).

3.21 Depth coordinates: the integer certificates and independent checks

3.21.1 First recurrence: fixed tail column

Set $r = 2d + 1 + q$ and work in $\mathbb{Z}[u]/(u^{d+1})$. From (3.60),

$$H_{d,k,q} = [u^d](2d + u)^r \mathcal{Q}_k^{(r)}\left(u + \frac{2d - q}{2}\right). \quad (3.79)$$

Although the displayed center may be half-integral, the recurrence only uses the integer linear polynomial $2u + 2d - q$. If $V_k(u)$ denotes the product on the right, truncated after degree d , then

$$\begin{aligned} V_0(u) &= (2d + u)^r \pmod{u^{d+1}}, \\ V_1(u) &= (2u + 2d - q)V_0(u) \pmod{u^{d+1}}, \\ V_{k+1}(u) &= (2u + 2d - q)V_k(u) + k(k + r)V_{k-1}(u) \pmod{u^{d+1}}. \end{aligned} \quad (3.80)$$

The required integer is the coefficient of u^d in V_k . Multiplication by each linear factor of V_0 and every recurrence step are implemented directly on lists of integers. Induction proves that truncation cannot affect any coefficient being tested.

3.21.2 Second recurrence: Pascal propagation

Equation (3.63) gives the independent array recurrence

$$H_{d,k+1,q} = 2H_{d,k,q+1} + (k + 2d + 2 + q)H_{d,k,q}, \quad (3.81)$$

with

$$H_{d,0,q} = [u^d](2d + u)^{2d+1+q}.$$

To compute $0 \leq k \leq K$, $0 \leq q \leq Q$, initialize the $k = 0$ row through $q = Q + K$ and shorten it at each step. This recurrence comes from multiplying by $(2 + w)$, rather than from the centered-polynomial three-term recurrence. Every entry of both finite rectangles was recomputed by both methods and compared as an exact integer.

3.21.3 Finite obligations and results

Depth	k maximum	q maximum	Cells	Symmetry zeros	Nonzero
3	12	37	494	6	488
4	83	184	15,540	42	15,498
Total			16,034	48	15,986

No additional zero occurs in either rectangle. Every nonzero integer also has a nonzero residue modulo at least one of 1009 and 1013. The first modulus suffices for every nonzero depth-three entry; nineteen depth-four entries require the second modulus. Primality is not needed for this particular witness rule: a nonzero residue modulo any integer greater than one already proves the original integer is nonzero.

The necessary seed signs are

d	k	q	$\text{sgn } H_{d,k,q}$	Role
3	14	6	+	central base
3	0	38	+	sum tail
3	1	38	−	difference tail
4	85	8	+	central base
4	1	185	−	sum tail
4	0	185	+	difference tail

For example,

$$H_{3,14,6} = 9707604961443840000 > 0, \quad H_{3,12,6} = -14670288691200000 < 0.$$

The exact larger seed integers, rather than floating-point approximations, are stored in `data/02-depth-certificates-verification.json`. The preceding central values at $k = 12$ and $k = 83$ are negative, so the chosen central exponents are the first positive ones in their relevant parity classes. This observation is not needed for validity of the certificates.

Remark 3.42 (The rectangles lie inside the range certified in this part; editorial). By $N = 2k + 2d + 1 + q$, the depth-three rectangle reaches derivative order $N \leq 2 \cdot 12 + 7 + 37 = 68$ and the depth-four rectangle $N \leq 2 \cdot 83 + 9 + 184 = 359$; the six seeds sit at $N = 41, 45, 47$ ($d = 3$) and $N = 187, 194, 196$ ($d = 4$). All of these lie inside the range $n \leq 3000$ of Proposition 3.18, whose modular certificate already shows that every rectangle cell outside the reflection family is nonzero. The nonvanishing half of the manuscript's finite computation is therefore a second, independent confirmation of cells this part had already certified; what Proposition 3.18 does not supply is the six seed *signs* (3.78), which the reduction needs and which the manuscript computes. The contribution of the addition is the proved reduction (Proposition 3.41) that makes a finite set of cells decide every n , not the finite range itself.

3.21.4 Additional cross-check against differentiation

The shipped program independently generates the original lattice using (3.54) through $N = 80$. It checks:

Exact test	Number of cells
Positive bulk cells	69,701
Identity $c_N(j, k) = \binom{N}{k} H_{d,k,-j-1}$	21,320
First-four-depth zero classification in that lattice	5,780

These are consistency checks on implementation and index translation. They are not the justification for the infinite theorem; that justification is Proposition 3.41 together with the finite obligations just listed.

3.21.5 Reproduction and trust boundary

From the report directory, on a copy (see below), run

```
python3 code/02-depth-certificates-minimal_verify.py
python3 code/02-depth-certificates-verify.py
```

The first prints PASS; the second prints a JSON report with "status": "PASS" and writes nothing. The manuscript's regeneration command, `verify.py -write-data`, writes the unprefixed files `finite_certificate.csv`, `sign_thresholds.csv` and `verification.json` into `data/` beside the shipped prefixed copies, so it should be run only on a copy of the report directory. The code uses only Python's standard library and arbitrary-precision integers. It uses no floating point, symbolic algebra library, random search, external service, or precomputed presumed support formula. Failures raise explicit exceptions, even if Python assertions are disabled. The CSV file supplies the sign and both modular residues for every rectangle cell.

The finite checks were run for the manuscript and returned PASS; they were rerun on a copy during the intake and passed again, with byte-identical CSV files. Ordinary program execution is still not a proof-assistant kernel check. A formal version would need the polynomial-model bridge, correctness of one recurrence, and a proved modular checker; the analytic lemmas would also need formalization. No such formalization is claimed. The two exact recurrences reduce the risk of implementation or indexing error, but do not replace independent mathematical review.

3.22 Depth coordinates: an all-depth bound on the number of zeros

The branch-cut formula also controls how often cancellation can occur, without any fixed-depth classification assumption. For fixed $d \geq 1$ and $k \geq 0$, define for real $q > -1$

$$\mathcal{F}_{d,k}(q) = \int_0^1 t^q (1-t)^{2d} (2t-1)^k R_d \left(\log \frac{1-t}{t} \right) dt. \quad (3.82)$$

For integer $q \geq 0$, its zeros are precisely those of $H_{d,k,q}$.

Lemma 3.43 (Mellin zero bound from sign changes). *Let W be a nonzero continuous real weight on $(0, 1)$ with finitely many zeros and exactly J sign changes. Assume that $t^q W(t)$ is integrable for every $q > -1$. Then $F(q) = \int_0^1 t^q W(t) dt$ has at most J distinct real zeros in $(-1, \infty)$. Here a sign change refers to a crossing of sign, not an even-multiplicity zero of the weight.*

Proof. Suppose $q_0 < \dots < q_J$ are zeros. There is a nonzero linear combination $\psi(t) = \sum_{i=0}^J a_i t^{q_i}$ vanishing at the J sign-change points of W : solve the corresponding homogeneous system of J linear equations. A nonzero linear combination of $J+1$ distinct real powers has at most J positive zeros, counting multiplicities. This follows inductively by dividing by the smallest power, differentiating, and applying Rolle's theorem. It also establishes linear independence of the powers.

Hence the prescribed zeros of ψ are simple and there are no others. After changing the overall sign, ψW is nonnegative throughout the interval and positive on a set of positive measure.

Therefore

$$0 < \int_0^1 \psi(t)W(t) dt = \sum_{i=0}^J a_i F(q_i) = 0,$$

a contradiction. Continuity and the finite zero set ensure strict positivity of ψW on an interval, as needed above. $\square$

Theorem 3.44 (At most d cancellations in a fixed row). *The number of distinct real zeros of $\mathcal{F}_{d,k}$ on $(-1, \infty)$ is at most*

$$J(d, k) = \begin{cases} d-1, & k \text{ even}, \\ d, & k \text{ odd and } d \text{ odd}, \\ d-2, & k \text{ odd and } d \text{ even}. \end{cases} \quad (3.83)$$

In particular, for fixed k, d there are at most d integer zeros in the entire negative-region tail. If a symmetry zero is present, it consumes one of these permitted zeros.

Proof. The polynomial R_d has $d-1$ distinct real roots. The logarithmic argument in (3.82) is strictly decreasing from $+\infty$ to $-\infty$, so these initially give $d-1$ sign changes. If k is even, $(2t-1)^k$ adds no sign change. If k is odd and d is odd, $R_d(0) \neq 0$, and the factor adds one sign change at $t = 1/2$. If k is odd and d is even, R_d has a simple zero at zero; its sign change at $t = 1/2$ is canceled by the odd power of $2t-1$, leaving $d-2$. The remaining factors are positive. Apply Lemma 3.43. $\square$

The theorem is a zero-count statement, not an assertion that all zeros are simple or that the upper bound is always attained. It also does not rule out an integer zero by locating nearby real zeros; the finite certificates are what supply that additional information at depths three and four. In this part's letters, it bounds by $J(m, j)$ the number of n at which a fixed (j, m) diagonal can vanish with $n \geq 2k+1$.

3.23 Depth coordinates: consequences for the term count

3.23.1 An unconditional upper bound (second route to Theorem 3.14)

The triangular support window and (3.57) imply that, in negative columns, the top $d=0$ term is absent. Removing those structural zeros, the number of remaining cells is $2\rho^2 + 2\rho + 1$ when $N = 2\rho$ and $2\rho^2$ when $N = 2\rho - 1$. For even N no symmetry hole is possible. For odd $N = 2\rho - 1$, a hole has

$$k = \rho - 2d - 1 \geq 0, \quad d \equiv \rho - 1 \pmod{2}.$$

Counting these depths gives $\lfloor \rho/4 \rfloor$ holes (or check the four residue classes of ρ). Thus, independently of unknown nonvanishing,

$$a(2\rho) \leq 2\rho^2 + 2\rho + 1, \quad a(2\rho - 1) \leq 2\rho^2 - \lfloor \rho/4 \rfloor. \quad (3.84)$$

The conjecture is exactly equality in these bounds. This is Theorem 3.14 and its equality clause, with the hole count $\lfloor \rho/4 \rfloor = \lfloor (n+1)/8 \rfloor$ of Lemma 3.13 counted by depth instead of by k .

3.23.2 A certified lower bound from the proved regions

Let $\beta(N)$ count the proved positive bulk cells, including $d = 0$ where it is nonzero. Direct summation gives

$$\beta(2\rho) = \frac{(\rho+1)(3\rho+2)}{2}, \quad \beta(2\rho-1) = \frac{3\rho^2+3\rho-2}{2} \quad (\rho \geq 1). \quad (3.85)$$

These are exactly the two cases of $\Lambda(N)$ in (3.3): the bulk is the region of Theorem 3.7, and Corollary 3.8 counts the same cells. For a fixed depth $1 \leq d \leq 4$, the cells outside this bulk have $q \geq 1$. The identity $N = 2k + 2d + 1 + q$ shows that their number before holes is $\max(0, \lfloor N/2 \rfloor - d)$. Define

$$\chi_d(N) = \mathbf{1}\{N \geq 4d + 1 \text{ and } N \equiv 2d + 1 \pmod{4}\}. \quad (3.86)$$

This is exactly the number, zero or one, of symmetry holes at that depth and derivative order.

Corollary 3.45 (Explicit unconditional lower bound). *For all $N \geq 0$,*

$$a(N) \geq \beta(N) + \sum_{d=1}^4 \max(0, \lfloor N/2 \rfloor - d) - \sum_{d=1}^4 \chi_d(N). \quad (3.87)$$

In particular, the lower bound has leading term $3N^2/8$, whereas the conjectured upper bound has leading term $N^2/2$.

Proof. Count the positive bulk and the disjoint first-four-depth cells with $q \geq 1$. By Theorem 3.23, only the indicated holes must be removed. The counts give (3.87). $\square$

For example, for even $N = 2\rho$ with $\rho \geq 4$, the lower bound simplifies to $(3\rho^2 + 13\rho - 18)/2$. It agrees with the conjectured value at $N = 8, 10$ and is 84 at $N = 12$, compared with the conjectured upper value 85. The unresolved cell there has depth five. These examples illustrate the precise boundary of the argument; they are not evidence that depth five fails. No asymptotic equivalence $a(N) \sim N^2/2$ is deduced here. (*Update, 1 October 2026:* depth five is now classified, Theorem 3.71, and the cell at $N = 12$ is nonzero; Corollary 3.78 adds depths five and six to this bound, and Theorem 3.83 lets the number of depths grow with N .) [Added 1 October 2026, batch 72A: the equivalence $a(N) \sim N^2/2$ is proved, ineffectively, in the sibling report **power-tower-exponent-supports** (its Corollary 13.1); the leading term $3N^2/8$ of (3.87) is still that of every explicit lower bound.]

Remark 3.46 (How much this improves Λ ; editorial). Since $\beta = \Lambda$, the gain of (3.87) over Theorem 3.1 is the sum of the last two terms. For even $N \geq 8$ no hole occurs and every maximum is attained by $N/2 - d \geq 0$; for odd $N \geq 17$ each maximum is $(N-1)/2 - d > 0$ and exactly two indicators equal one (at $d = 1, 3$ if $N \equiv 3$, at $d = 2, 4$ if $N \equiv 1 \pmod{4}$). Hence

$$a(N) \geq \begin{cases} \Lambda(N) + 2N - 10 = \frac{3N^2 + 26N - 72}{8}, & N \text{ even, } N \geq 8, \\ \Lambda(N) + 2N - 14 = \frac{3N^2 + 28N - 111}{8}, & N \text{ odd, } N \geq 17. \end{cases}$$

The leading constant $3/8$ is unchanged; the linear term improves. [Added 1 October 2026, batch 72A: unchanged for explicit bounds, including Theorem 3.83; in the limit the constant is $\frac{1}{2}$, by the sibling report **power-tower-exponent-supports**, ineffectively.] Below these thresholds the gain is read off (3.87) directly; the bound equals $U(N)$ for every $N \leq 11$. Of the even gain $2N - 10$, the part $N - 3$ from depths one and two already followed from Theorems 3.16–3.17,

although Part 3 did not add it to Λ ; the part $N - 7$ from depths three and four is new. At odd $N \geq 17$ the split is $(N - 5) + (N - 9)$. The two closed forms were checked against a direct count of the proved cells for every $N < 400$. $\Lambda(n)$ remains the lower bound stated in Theorem 3.1; the certified range $n \leq 3000$ of Proposition 3.18 makes both bounds irrelevant there, since $a(n) = U(n)$ is known for those n .

3.24 Depth coordinates: further research questions

3.24.1 Depths five and six: a stronger moment criterion

The explicit quadratics in (3.68) suggest a practical next step. Since $\mathbb{E}[Z^2] \geq (\mathbb{E}Z)^2$, where $Z = \tau^2$, sufficient conditions for positive reduced expectation are

$$\begin{aligned} d = 5 : \quad \mathbb{E}Z &> \pi^2(1 + 2/\sqrt{5}), \\ d = 6 : \quad \mathbb{E}Z &> 3\pi^2. \end{aligned} \tag{3.88}$$

These follow by applying the variance inequality to the two quadratic polynomials and placing the mean above their largest root. Unlike the crude affine minorant, these criteria retain the exact quadratic structure. Their usefulness is that $\mathbb{E}Z$ is monotone under all the proved likelihood-ratio comparisons even though Π_5 and Π_6 are not monotone on the entire positive axis.

Question 1. Determine certified practical k and q cutoffs from (3.88), and settle the resulting finite rectangles at depths five and six. A successful computation must be accompanied by the mean-moment bounds; an unbounded inference from a numerical scan would not suffice. (Editorial: the range $n \leq 3000$ of Proposition 3.18 already certifies nonvanishing of every depth-five and depth-six cell with $n \leq 3000$; what is missing is the reduction, and the seed signs it will require.)

Update (1 October 2026, batch 72A): answered. Manuscript 70 carries out exactly this programme (Subsections 3.37 and 3.38). The cutoffs are $(R, Q) = (410, 839)$ at depth five and $(1571, 3048)$ at depth six. They are certified by exact lower bounds on the mean of τ^2 (Lemmas 3.75 and 3.79) — moment seeds rather than sign seeds, since Π_5 and Π_6 are not monotone. The rectangles of 343,151 and 4,785,360 cells are settled by two recurrences each, and Theorem 3.71 classifies both depths.

3.24.2 Sharpening the fixed-depth reduction

Question 2. How small can uniform fixed-depth cutoffs be made as functions of d ? The constants in (3.74) are intentionally wasteful. The largest residual root is $\pi^2 \cot^2(\pi/d)$, of order d^2 , so endpoint concentration on a logarithmic scale is a natural quantity to analyze. This observation does not yet prove an optimal cutoff growth rate.

Update (1 October 2026, batch 72A): answered, not optimally. Theorem 3.81 takes exactly this route — the largest residual root, below d^2 , and beta logarithmic moments — and gives the cutoffs $\widehat{Q}_d = \lceil (2d + 1)e^d \rceil$, $\widehat{K}_d = 2\widehat{Q}_d$, a rectangle of $O(d^2 e^{2d})$ cells in place of the tower-type constants (3.74). No optimality is claimed; the practical cutoffs at $d = 3, \dots, 6$ are far smaller.

Question 3. Can lower bounds involving the first few moments of τ^2 , or positivity in an orthogonal-polynomial basis adapted to $\mu_{d,r}$, replace the affine minorant for all depths? The target is a small finite certificate rather than an enormous merely terminating algorithm. (*Update, 30 September 2026:* at $d = 3$ the second route of Subsections 3.30–3.31 gives an explained cutoff, the exact sign change of E_q between $q = 37$ and $q = 38$, and a different small certificate; it bears on Questions 2 and 3 without answering them for general d .) (*Update,*

1 October 2026: answered in part. Proposition 3.80 replaces the affine minorant at every depth by the sharp distribution-free first-moment criterion $\mathbb{E}\tau^2 > \varrho_d$; at $d = 5, 6$ its thresholds are those of the convexity criteria (3.88). Positivity in an orthogonal-polynomial basis is not attempted, and the certificates obtained this way are not small at large d .)

[Added 1 October 2026, batch 73: Questions 1–3. Manuscript 37 of Subsection 3.42ff. carries the programme of Question 1 to depths seven and eight with the largest-root criterion, and gives a practical three-anchor reduction at every depth (Theorem 3.97); manuscript 46 does depth seven independently. For Question 2, manuscript 37’s uniform cutoffs $\check{K}_d, \check{Q}_d = \Theta(d^2 e^d)$ are weaker than $\widehat{K}_d, \widehat{Q}_d$ (Remark 3.93); no improvement and no optimality. For Question 3, its tangent is Proposition 3.80, and its question on moments beyond the first (Question 3.113) restates the open part.]

3.24.3 The remaining full conjecture

Question 4. Prove that every fixed-depth rectangle has only the symmetry zeros, uniformly in depth, or exhibit the first exact counterexample. The present results show where additional zeros must be searched for at each depth, but they do not supply an induction in d . The polynomial Π_d acquires additional oscillations, which must be addressed rather than ignored. (This is Conjecture 3.19 restricted to $m \geq 5$; *update, 1 October 2026:* since Theorem 3.71, restricted to $m \geq 7$; since batch 73, Theorems 3.99 and 3.100, restricted to $m \geq 9$; Question 3.110 asks it again.)

Question 5. Can the real-zero bound (3.83) be combined with arithmetic separation to exclude integer zeros uniformly? The Mellin argument limits the number of candidates in each row, but integer nonvanishing needs a second ingredient. An effective distance-from-integers theorem for its zeros would be particularly valuable.

[Added 1 October 2026, batch 72A: the sibling report **a290268-unbounded-deficits** bears on Questions 4 and 5 without answering either. At every fixed offset $b = q - 2d$ it proves eventual nonvanishing along each ray $k/d \rightarrow \kappa$ with κ algebraic below the fold $\kappa_c \approx 0.0281$, apart from the reflection holes, and uniformly for $|b| \leq 8$, $0 \leq k \leq d/36$ (its Theorem 2.1); an Airy law at the fold (its Part II); nonvanishing at $k = 0$ on all-depth strips around $q = 2d$ (its Theorem 29.2); prime-unit families with exact valuations; and an exact example showing that root geometry alone cannot prove Conjecture 3.19 (its Part IV). None of its thresholds is numerical.]

Question 6. Establish the conjectured leading term $a(N) \sim N^2/2$ without first proving every individual nonvanishing claim. This would require bounding the total number of extra zeros across growing depths. The fixed-row bound alone does not give such a bound after summing over the whole triangular array.

[Added 1 October 2026, batch 72A: answered. The sibling report **power-tower-exponent-supports** proves $a(N) = N^2/2 + o(N^2)$ (its Corollary 13.1, with manuscript 34’s proof; the case $a = 2$ of its Theorem 3.1) by showing that only $o(N^2)$ of the candidate cells outside the positive bulk vanish — the bound across growing depths asked for — without classifying individual zeros. The proof is ineffective. The explicit lower bounds are still this report’s, the best being Theorem 3.83; Theorem 3.85 and Proposition 3.87 measure what the fixed-row bound gives. Coefficient-level results at growing depth (cosine laws, density one with $O(N^{5/3})$ exceptions on tail patches) are in **a290268-unbounded-deficits**, whose headline lower bounds this asymptotic supersedes (its Section 25).]

3.24.4 Formal proof and symbolic computation

Question 7. Formalize the polynomial factorization, bulk theorem, and integer certificate in the existing A290268 Lean development. The algebraic route has a small interface: coefficient extraction, falling factorials, the recurrence (3.61), and a finite modular nonzero checker. The branch-cut and likelihood-ratio proofs are a separate analytic layer. Keeping these layers explicit would avoid confusing a verified finite table with a verified infinite theorem. (Section 3.26.3 lists what that development has and which of its hypotheses these results would discharge. *Update, 30 September 2026:* a module-by-module algebraic blueprint for depth three is Section 3.33.2; none of its modules exists.)

Question 8. Extend the method to derivative expansions of x^{x^r} for integer $r \geq 3$, or to other exponential-logarithmic families. The correct depth generating function and the symmetry center must first be rederived. The success of the square-exponent case does not justify assuming an analogous support quasipolynomial elsewhere.

[Added 1 October 2026, batch 72A: partly answered. The sibling report **power-tower-exponent-supports**, which writes the exponent a , proves $N^2/2 + o_a(N^2)$ terms for every integer $a \geq 1$ (its Theorem 3.1, ineffective), derives the depth generating function (its equation (3.6)) and the symmetry centre $q = ad$, and classifies the first two depths (its Theorem 11.1). It also proves $N^2/2 + o(N^2)$ for $x^{\sqrt{x}}$ (its Theorem 14.1), and only finitely many zeros off the reflection line on each fixed defect large enough: from defect four for rational $a \neq 1$, from $4D + 2$ for algebraic a of degree D , from two (effectively) for irrational a (its Part III). For $a = 1/2$ and every rational $1/2 < a < 1$ there are infinite families of non-reflection zeros (its Theorem 14.3 and Corollary 16.3), so an analogous support pattern indeed cannot be assumed. Exact noncancellation at arbitrary depth stays open for every exponent.]

[Added 5 October 2026, batch 100: the sibling report **power-tower-exponent-supports** now also has a Part IV (its Sections 21–24, from bundle Report 59, an eighth note of 1 October 2026 on the same pin). For $x^{\sqrt{x}}$ it shows that on each fixed defect the scaled residual tends to a Hermite polynomial with an explicit first correction, with eventually real simple roots, so at most $O_r(\sqrt{X})$ coefficients of a fixed defect vanish through order X ; its finiteness from defect four is that report's Theorem 16.1 at $a = 1/2$, proved again with explicit integer quartics. No effective threshold is given.]

3.24.5 Suggested order of work

The most direct continuation is to formalize the bulk factorization while independently implementing certified moment bounds for depths five and six. After those bounds determine practical finite rectangles, the existing integer recurrence can generate exact certificates immediately. Only then should one search for a depth-uniform induction or arithmetic zero-separation theorem. This separates achievable finite obligations from the genuinely new ingredient needed for the complete conjecture.

3.25 Depth coordinates: conclusion, verifier and source audit

3.25.1 Conclusion of the addition

The obstruction in A290268 is not merely the existence of a few symbolic cancellations. Its coefficients combine a consecutive-root factor with an imaginary-root factor, and their tail values are signed Mellin moments of an explicitly oscillating logarithmic polynomial. Making both structures visible yields three concrete advances in the manuscript's account: positivity in the whole bulk at every depth, an explicit finite decision problem at every fixed depth, and

a complete nonvanishing classification at depths three and four, supplementing the first two depths. Of these, the first was already Theorem 3.7 of this part, and the second and third are new.

The 16,034-cell verification is logically effective because the manuscript first proves that it is exhaustive for those two infinite-parameter strata. It is not extrapolation from a finite sample. The full counting formula remains a separate target at depths five and higher, with the remaining questions and the precise proof boundary now substantially narrowed. (*Update, 1 October 2026*: depths five and six are settled by Theorem 3.71; the target is now depths seven and higher. *Update, batch 73*: depths seven and eight are settled by Theorems 3.99 and 3.100; the target is now depths nine and higher.)

3.25.2 A compact standalone finite verifier

The following standalone version verifies the two exact rectangles and six seed signs; it is shipped, byte for byte, as `code/02-depth-certificates-minimal_verify.py` (whose docstring calls this listing “Appendix A of article.tex”, the manuscript’s numbering). The fuller program `code/02-depth-certificates-verify.py` additionally checks the independent Pascal recurrence, produces the CSV witnesses, and cross-checks the original differentiation lattice.

```
def must(test):
    if not test:
        raise ArithmeticError("certificate failure")

def column(d, q, K):
    r = 2*d + 1 + q
    a = [1] + [0]*d
    for j in range(r):
        a = [(2*d-j)*a[i] + (a[i-1] if i else 0)
              for i in range(d+1)]
    out = [a[d]]
    if K == 0:
        return out
    b = [(2*d-q)*a[i] + (2*a[i-1] if i else 0)
          for i in range(d+1)]
    out.append(b[d])
    for k in range(1, K):
        c = [(2*d-q)*b[i] + (2*b[i-1] if i else 0)
              + k*(k+r)*a[i] for i in range(d+1)]
        out.append(c[d])
        a, b = b, c
    return out

for d, R, Q in [(3, 14, 38), (4, 85, 185)]:
    for q in range(Q):
        for k, value in enumerate(column(d, q, R-2)):
            hole = (q == 2*d and (k+d) % 2 == 0)
            must((value == 0) == hole)
            if not hole:
                must(value % 1009 != 0 or value % 1013 != 0)
    must(column(d, 2*d, R)[R] > 0)
    ks, ka = (d-1) % 2, d % 2
    must((-1)**Q * column(d, Q, ks)[ks] > 0)
    must((-1)**(Q+1) * column(d, Q, ka)[ka] > 0)
```

```
print("PASS")
```

The loop bounds are conclusions of the analytic reduction, not values chosen merely because the program happened to finish. Each arithmetic step is the recurrence (3.80) in the ring $\mathbb{Z}[u]/(u^{d+1})$. (The Python variable `j` in `column` is a loop index over the factors of the falling factorial, not either meaning of j in Table 3.) Thus the program is a finite, reproducible arithmetic certificate with a fully specified mathematical meaning.

3.25.3 Source audit

The manuscript’s inspected repository snapshot is `afb2d1227d8bc5df3beffc1463e958db3544960b`. At that snapshot the A290268 research note explicitly identified higher-depth negative-region nonvanishing and the general- k bulk as open parts of its analysis; the general- k bulk was in fact already Theorem 3.7 of this report, which the manuscript did not inspect. The `Main.lean` file makes the support characterization a hypothesis. The proof in the manuscript does not depend on uninspected formal claims or on the consistency of unrelated parts of the repository.

The manuscript’s scoped directory listing contains the core, closed-form, counting, structural, table, and main modules. The research note also discusses other module names and development intentions. Those descriptions are not treated as evidence that every discussed file is present or that every corresponding theorem is kernel-certified. (At the intake, the three module names that note lists beyond these six, `A290268.Hole`, `A290268.DepthOne` and `A290268.Series`, were confirmed absent from the repository’s entire history.)

External checks included the live OEIS entry, the NIST generating function, and the primary paper by Štampach, whose bibliographic data the manuscript cross-checked against a paper of R. B. Paris (bibliographic verification only, recorded in `02-depth-certificates-SOURCE_AUDIT.md`). Search results did not supply a prior published proof of the specific fixed-depth statements proved here. That limited search is not a proof of universal novelty.

3.26 Depth certificates: provenance, files and formal status

3.26.1 Provenance

Subsections 3.14–3.26 were built from **one** manuscript: batch 42, manuscript 06, archive `ProveIt_A290268_Finite_Certificates.zip` (22 pages, dated 29 September 2026), pinned to ProveIt commit `afb2d1227d8bc5df3beffc1463e958db3544960b`, and placed in this report by commit `3609d0473`. Its Section 1 is split between Subsection 3.14 (main results, novelty) and the opening of Subsection 3.15 (the conjecture, linear independence); its Sections 2–11 map one for one onto Subsections 3.15–3.24; its Section 12 and its two appendices form Subsection 3.25. Every theorem, proposition, lemma, corollary, definition, remark and question of the manuscript is printed; none was weakened or strengthened. The editorial choices were these.

- It is placed as subsections at the end of Part 3 rather than as a new part, so that no existing section, theorem or equation number moves.
- Its bulk positivity, symmetry holes, depths one and two, upper bound, depth bridge, linear independence and branch-cut formula are marked as second routes to, or re-indexings of, results of this part, which the manuscript did not read. The editorial Remarks 3.31, 3.42 and 3.46 and the parenthetical translations are additions of the intake, not of the manuscript.

- The manuscript’s letters are kept, with the renamings of Subsection 3.14.1; its reproduction commands are rewritten to the shipped file names; its archive table is replaced by the file list below; its OEIS citation is merged with this part’s.
- Its statement that the finite checks “have actually been run” is kept, with the intake’s rerun recorded beside it.

Not shipped: the manuscript’s `article.tex`, `article.pdf`, `README.md` and `SHA256SUMS.txt` (verified at placement, 10 of 10 entries); the archive itself survives in the history of commit 8315d24e3.

3.26.2 Files

The manuscript’s code and data are shipped with the prefix `02-depth-certificates-`, byte-identical to the delivery:

Shipped path (delivery name)	Purpose
<code>code/02-depth-certificates-verify.py</code> (<code>code/verify.py</code>)	Full exact verifier: both recurrences, the lattice cross-check, and data generation.
<code>code/02-depth-certificates-minimal_verify.py</code> (<code>code/minimal_verify.py</code>)	The standalone verifier of Subsection 3.25.2.
<code>code/02-depth-certificates-Makefile</code> (<code>Makefile</code>)	The delivery’s make targets; see the README before using it.
<code>data/02-depth-certificates-finite_certificate.csv</code> (<code>data/finite_certificate.csv</code>)	All 16,034 rectangle cells with signs and residues modulo 1009 and 1013.
<code>data/02-depth-certificates-sign_thresholds.csv</code> (<code>data/sign_thresholds.csv</code>)	Observed sign transitions of the reduced expectation in each tested row.
<code>data/02-depth-certificates-verification.json</code> (<code>data/verification.json</code>)	Executed-check summary and the exact seed integers.
<code>02-depth-certificates-SOURCE_AUDIT.md</code> (<code>SOURCE_AUDIT.md</code>)	Source identity, inspected scope, and limitations.

The two CSV files keep the delivered CRLF line endings.

3.26.3 Formal status

Nothing in this addition is formalized, and its placement beside a Lean development confers no formal status. The Lean development `Combinatorics/DerivativeExpansions/A290268/Lean` (library `A290268`, namespace `LeanProofs.A290268`, imported by the root `ProveIt.lean`) relates to it as follows.

- `coeff n j k` (`Core.lean`) is the recurrence (3.54) in the same coordinates; `S n` (`Count.lean`) is the conjectured support, and `card_S` proves $|S_n|$ equal to the closed form unconditionally.
- The manuscript relies on the *statements* of the three conditional theorems of `Main.lean`: `a_eq_closedForm_of_support` (hypothesis: the support of `coeff n` is `S n`), `a_eq_`

`closedForm_of_two_sided` (the structural and nonvanishing inclusions as separate hypotheses) and `a_eq_closedForm_of_support_all` (the support hypothesis at every order). It reads them, correctly, as conditional.

- `Structural.mem_S_of_coeff_ne_zero` proves the structural inclusion from a hypothesis `hhole`: vanishing on the line N odd, $-j \equiv N \pmod{4}$, $-j \geq 3$, $2(-j) - 1 \leq N$, $2k = N + 2j + 1$. With $-j = 2d + 1$ that line is exactly the hole family of Theorem 3.29, and so of Theorem 3.11; either proof would discharge `hhole` on paper. Neither is formalized, and the module `A290268.Hole` that the declaration’s docstring names as the discharge does not exist in the repository.
- Theorem 3.23 would discharge the nonvanishing inclusion only on cells of depth at most four; depths $d \geq 5$ remain, so the `Main.lean` theorems stay conditional. (Update, 1 October 2026: with Theorem 3.71, on paper, depths at most six; depths $d \geq 7$ remain. Batch 73: with Theorems 3.99 and 3.100, depths at most eight; depths $d \geq 9$ remain.)
- The only unconditional statement about the sequence itself is `Table.a_values_le_53`, $a(n)$ equal to the closed form for $n \leq 53$, a `native_decide` certificate.

3.27 A second route to depth three

What this addition is. Subsections 3.27–3.34 were written into this part on 30 September 2026 from a later manuscript, *Depth Three Is Complete for OEIS A290268: A beta-sine transform, centered moment polynomials, and exact nonvanishing* (batch 70, manuscript 03; provenance in Section 3.34). Its main theorem (Theorem 3.59) is **the case $d = 3$ of Theorem 3.23**, which the batch-42 addition had already proved in this part; it is printed here as a *second proof* of that case. Nothing in it changes the proof boundary of this part: the unproved region is still $m \geq 5$, $n \geq 2k + 2$, and the full A290268 conjecture is *not* proved. (Update, 1 October 2026: since Subsections 3.35ff., the unproved region is $m \geq 7$; since Subsections 3.42ff. (batch 73), $m \geq 9$.)

A correction to the manuscript. The manuscript says, in its abstract, its main-contribution box, its novelty paragraph, its conclusion and its source audit, that depth three was “the stated open core” of the problem at its pinned commit `a02ab3567` and that its theorem is new relative to that commit. **That is mistaken.** The pin descends from commit `0e6632abf`, which wrote Theorem 3.23 (depths one to four, every k and M) into this report on 29 September 2026; and the note the manuscript cites, `0eis/A290268/README.md`, already says at the pin, in its dated correction, that this report’s batch-42 addition “closes logarithmic deficits 3 and 4 for every n ”. The manuscript read only that note’s older line “Open core: $D \geq 3$ ”. Its sentences are printed below as the manuscript wrote them, each followed by an editorial note; none of its mathematics depends on them.

What is new relative to this part. Several results of the manuscript re-derive results printed above, by different proofs; they are printed once more, marked as second routes, with the earlier result credited:

- its main theorem and lattice corollary (Theorem 3.59, Corollary 3.60) are Theorem 3.23 at $d = 3$ and the depth-three holes of Subsection 3.20;
- its exact reduction and parameter derivative (Proposition 3.48, Lemma 3.49) are Proposition 3.24 and (3.58);
- its beta-sine transform (Theorem 3.51) is the branch-cut formula of Proposition 3.30, obtained, as in Remark 3.31, from the beta-integral mechanism of Lemma 3.15;
- its centred recurrence and vertical-line roots (Theorem 3.52, Corollary 3.53) are (3.61) and Corollary 3.27, proved here by an integration by parts and a Favard argument;
- its bulk positivity (Lemma 3.58) is Theorem 3.21 at $d = 3$, hence a case of Theorem 3.7.

New relative to this part are the harmonic factorization at the root $x = 7$ (Subsection 3.30); the sign law of the harmonic quantity E_q (Lemma 3.54), which *proves* the $q = 37/38$ sign change that Subsection 3.20 recorded only as an observation of the certificate data; the propagation recurrence (3.107) in k and the finite certificate it needs (Proposition 3.55), a different finite set from the batch-42 rectangle; the eventual signs of Corollary 3.57 for every q ; the sign phase diagram of the log-free diagonal; the beta representation of the uncentred quotient (Lemma 3.50); and the interlacing clause of Corollary 3.53. Remark 3.61 compares the two proofs and shows that their finite anchors partly coincide.

Notation. These subsections use the depth coordinates of Table 3 — k is the exponent of $\log x$, j the exponent of x , d the depth — and the batch-42 names for the objects batch 42 already named. The manuscript’s own letters clash with both, so many are renamed; Table 5 lists every renaming and is binding from here to the end of the part.

Abstract of manuscript 03. *OEIS A290268 counts the terms in the fully expanded n th derivative of x^{x^2} . The ProveIt reduction expresses every coefficient through $\gamma(k, D, M) = [w^M](2 + w)^k((1 + w)^2 \log(1 + w))^D$, and identifies the remaining support conjecture with a nonvanishing problem for these rational numbers. At the pinned repository snapshot, depths $D = 1, 2$ are solved in the accompanying research notes, whereas $D \geq 3$ is the stated open core.*

This article proves the complete zero classification at the first open depth: for every $k \geq 0$ and $M \geq 3$,

$$\gamma(k, 3, M) = 0 \iff M = k + 13 \text{ and } k \text{ is odd.}$$

Thus every depth-three coefficient is nonzero except for the symmetry-forced holes already predicted by the global conjecture.

The proof has three new ingredients. First, a general beta-sine transform represents the tail coefficients at every depth by a single logit moment with an explicit kernel $K_D(L) = ((L + i\pi)^D - (L - i\pi)^D)/(2i\pi)$. Second, a family of centered beta-moment polynomials satisfies the positive three-term recurrence $R_{N,k+1}(y) = 2yR_{N,k}(y) + k(N + k)R_{N,k-1}(y)$. Third, at $D = 3$ the problem reduces to a signed harmonic quantity whose only phase transition occurs between $q = 37$ and $q = 38$. Positivity propagation then reduces the remaining proof to a finite exact certificate over $\mathbb{Q}$. The supplied standard-library Python verifier uses `fractions.Fraction`; no floating-point assertion enters the proof.

This is a theorem relative to the definitions in the pinned ProveIt snapshot, not a claim that the full A290268 conjecture has been proved: depths $D \geq 4$ remain open. No independent peer review or Lean formalization is claimed.

Editorial notes. (i) The abstract is printed in the manuscript’s letters; below, D is d , n is N , K_D is R_d , L is τ , and $R_{N,k}$ is $\mathcal{Q}_k^{(r)}$ (Table 5). (ii) “ $D \geq 3$ is the stated open core”, “the first open depth” and “depths $D \geq 4$ remain open” are stale at the pin: depths three and four were settled by Theorem 3.23, and the open depths are $d \geq 5$ (since 1 October 2026, $d \geq 7$: Theorem 3.71; since batch 73, $d \geq 9$: Theorems 3.99 and 3.100). (iii) Of the “three new ingredients”, the first two are not new relative to this part (the bullets above); the third, the harmonic phase transition with its propagation in k , is.

Main contribution (the manuscript’s summary box). The first previously open depth in the ProveIt support program is completely settled. In the original coefficient lattice, if $d = (N + j)/2 - k = 3$ and $N \geq k + 3$, then

$$c_N(j, k) = 0 \iff j = -7, \quad k \text{ odd}, \quad N = 2k + 13.$$

Every other depth-three lattice point is nonzero.

Editorial note. “First previously open depth” is mistaken (see above): this is the depth-three content of Theorem 3.23, whose holes Subsection 3.20 lists as $q = 6$, k odd; $j = -7$, $N = 2k + 13$. In the letters of the rest of this part: $m = 3$, j odd, $n = 2j + 13$.

3.27.1 Notation: the second route

The manuscript writes the same lattice $x^{x^2+j}(\log x)^k$ as the batch-42 addition, with the same j, k, M, q and γ , so the dictionary of Table 3 applies unchanged. Its other letters collide with letters already in use in this part or in the batch-42 subsections, and are renamed as in Table 5, with no change of normalization. The first four rows are where a reader will go wrong.

Table 5: Letters of Subsections 3.27–3.34 against manuscript 03 and against the rest of Part 3. No normalization was changed.

Here	Manuscript	Meaning	Tempting false reading
N	n	derivative order	false: the manuscript's N , which is r here
r	N	$M - k$ (as in Subsection 3.15)	false: r in $n = 2r$ of Part 3
$\mathcal{Q}_k^{(r)}(y)$	$R_{N,k}(y)$	centred polynomial of Definition 3.25	false: the kernel R_d of (3.64)
$R_d(\tau)$	$K_D(L)$	kernel $\text{Im}(\tau + i\pi)^d/\pi$, (3.64)	false: the constant K_d of (3.74)
d	D	depth $(N + j)/2 - k$	$d = n - 2k - 1$ of (3.25)
τ	L	$\log(t/(1 - t))$	$L = \log x$ of Part 3
$\mathcal{P}_M(x)$	$P_M(x)$	$\prod_{\ell=1}^M (x - \ell) = (x - 1)^{\underline{M}}$	the canonical P_n ; the residual Π_d
$\mathcal{H}_{M,k}(x)$	$H_{M,k}(x)$	$\sum_{\ell} \binom{k}{\ell} \mathcal{P}_M(x + \ell) = M! \mathcal{A}_{M,k}(x - 1)$	the integer $H_{d,k,q}$ of (3.49)
$\tilde{\mathcal{Q}}_{r,k}(x)$	$Q_{N,k}(x)$	$\mathcal{Q}_k^{(r)}(x - \frac{r+1}{2})$, uncentred	$Q_{n,j}$ of Section 3.3; the constant Q_d
$\mathcal{T}_{r,k}(s)$	$S_{N,k}(s)$	$i^{-k} \mathcal{Q}_k^{(r)}(is)$	$S_{\delta}(t)$ of Subsection 3.17
$\Omega_r(x)$	$S_N(x)$	$\mathcal{P}_r(x)/(x - 7)$	$S_{\delta}(t)$
$\mathcal{I}_k(x)$	$I_k(x)$	beta moment	$I_{m,j,d}(\varepsilon)$ of (3.26)
$H_q, H_q^{(2)}$	$H_q, H_q^{(2)}$	harmonic numbers $\sum_{\ell \leq q} \ell^{-1}$, $\sum_{\ell \leq q} \ell^{-2}$	$H_{d,k,q}$; $H(u)$ of (3.15)
E_q	E_q	signed harmonic quantity (3.101)	the constant E_d ; the expectation $\mathcal{E}_{d,k,q}$
$F_k(q)$	$F_k(q)$	normalized depth-three tail value (3.102)	the Mellin transform $\mathcal{F}_{d,k}$; the falling factorial F_n
$U_{q,k}$	$U_{q,k}, U_k$	positivity functional (3.105)	the upper bound $U(n)$
V_k	V_k	$2F_k(6)$	$V_k(u)$ of (3.80)
λ_q	z	$ q - 6 /2$	$z = x^2$ of Part 3
α_q	a (tail)	$ H_q - H_6 $	the term count $a(n)$
ε	z (kernel)	a local variable near 0	—

Further changes, also without change of normalization: the manuscript's ϕ is φ of (3.48); its centre $c_N = (N + 1)/2$ is written $(r + 1)/2$, because $c_N(j, k)$ is the coefficient; its evaluation point $a = 2D + 1$ is written $2d + 1$ (it is 7 at depth three); its $B = -E_q$ is written $|E_q|$; summation and product indices written r in the manuscript are ℓ ; the hole index r of its lattice corollary is ℓ ; the index m of its count formula is ρ , as in (3.53); and its indeterminate a in Lemma 3.49 is θ . The manuscript's falling factorial P_M is the batch-42 $s^{\underline{M}}$ at $s = x - 1$, and $\mathcal{H}_{M,k}(x) = M! \mathcal{A}_{M,k}(x - 1)$, so the manuscript's x is the batch-42 s plus one: the evaluation point $x = 2d + 1$ is $s = 2d$ of (3.58). By Proposition 3.26 the manuscript's quotient polynomial $Q_{N,k}$ is the centred polynomial $\mathcal{Q}_k^{(r)}$ translated by $(r + 1)/2$, and its $R_{N,k}$ is $\mathcal{Q}_k^{(r)}$; that is why both are printed with the batch-42 letter. Its kernel K_D and the batch-42 R_d are the same polynomial, since $((\tau + i\pi)^d - (\tau - i\pi)^d)/(2i\pi) = \text{Im}(\tau + i\pi)^d/\pi$ for real τ . Finally, the two

tail values are related by

$$H_{3,k,q} = 720 (-1)^q q! F_k(q), \quad (3.89)$$

which follows from (3.49) and (3.103) below and was checked at the intake in exact arithmetic for $0 \leq k \leq 14$, $0 \leq q < 60$. In particular $\operatorname{sgn} H_{3,k,q} = (-1)^q \operatorname{sgn} F_k(q)$.

3.27.2 Problem, status and scope (manuscript Section 1)

The sequence and its support conjecture. Let

$$\left(\frac{d}{dx}\right)^N x^{x^2} = \sum_{j,k} c_N(j,k) x^{x^2+j} (\log x)^k.$$

OEIS A290268 is $a(N) = \#\{(j,k) : c_N(j,k) \neq 0\}$. The public OEIS entry still labels as conjectural the generating function

$$\sum_{N \geq 0} a(N) z^N = \frac{1 + z + 2z^2 + 2z^3 + 2z^4 + 2z^5 + 2z^6 + z^7 + 2z^8 + z^9}{(1-z)(1-z^2)(1-z^8)},$$

equivalently $a(2\rho) = 2\rho^2 + 2\rho + 1$, $a(2\rho - 1) = 2\rho^2 - \lfloor \rho/4 \rfloor$ (3.53). (Here z is a generating-function variable, local to this display.) The entry records 54 terms, through $N = 53$; the manuscript inspected it on 30 September 2026 [2].

The pinned ProveIt notes reduce the conjecture to an exact support statement. They describe a depth statistic d and a coefficient $\gamma(k, d, M)$, prove or outline the symmetry-forced holes, record complete arguments at depths $d = 1, 2$, and identify $d \geq 3$ as the open nonvanishing core [22]. The Lean module `A290268.Main` makes the final logical boundary explicit: the closed form follows as soon as the coefficient support is identified with the conjectured finite set [23].

Editorial note. The notes do say “Open core: $D \geq 3$ ” at the pin, in their section on solved strata; the same file’s dated correction (“Correction (29 September 2026)”) says that this report’s batch-42 addition closes deficits three and four for every n . The reading of `A290268.Main` is correct: its three theorems are conditional on the support characterization (Section 3.26.3).

What is proved here. The present report resolves the entire $d = 3$ stratum. It does not infer the result from a numerical scan. The infinite ranges in both k and M are proved using recurrences and positivity. A finite exact computation is used only for a compact critical strip where a signed kernel prevents a direct one-sign argument.

Proof boundary (the manuscript’s status box). The analytic and algebraic reductions are proved in the text. One finite lemma checks exact rational signs for $q \in \{1, \dots, 5\} \cup \{7, \dots, 37\}$, $0 \leq k \leq 14$, together with the centre $q = 6$. The included verifier reproduces that lemma from integer polynomial recurrences and rational harmonic numbers. Large additional boxes are checked only as cross-validation and are not needed for the theorem.

Novelty discipline. The result is new relative to repository commit `a02ab356`: the pinned A290268 status document explicitly calls $d \geq 3$ open [22]. The OEIS entry also continues to call the global formula conjectural [2]. A targeted search found no later public proof of this depth-three stratum. This statement is deliberately narrow: the report has not undergone independent refereeing, and it does not claim exhaustive priority over inaccessible or unpublished work.

Editorial note. The first sentence is mistaken: at [a02ab356](#) the repository already contained Theorem 3.23, whose $d = 3$ case is the manuscript's theorem, and the cited document's own correction points to it. The theorem is therefore *not* new relative to the pin; its proof is a second, different route. The remaining sentences — the OEIS status, the narrow search, no refereeing, no exhaustive priority — stand.

3.28 Second route to depth three: the coefficient model and a falling-factorial transform

Set $\varphi(w) = (1 + w)^2 \log(1 + w)$. The Taylor identity

$$\sum_{N \geq 0} \frac{\xi^N}{N!} \left(\frac{d}{dx} \right)^N x^{x^2} = (x + \xi)^{(x+\xi)^2}$$

becomes, after dividing by x^{x^2} and putting $w = \xi/x$,

$$x^{-x^2} (x + \xi)^{(x+\xi)^2} = \exp(x^2 w (2 + w) \log x) \exp(x^2 \varphi(w)).$$

This identity is valid formally in w over a coefficient ring containing $x, x^{-1}, \log x$; no analytic choice of logarithm is needed for coefficient extraction.

Definition 3.47 (Depth and reduced coefficient). For integers $N \geq 0$, $k \geq 0$, and an exponent shift j of the correct parity, define $d = (N + j)/2 - k$ and $M = N - k$, and, for $d, M \geq 0$, $\gamma(k, d, M) = [w^M](2 + w)^k \varphi(w)^d$.

Proposition 3.48 (Exact reduction; second route to Proposition 3.24). *Whenever $d = (N + j)/2 - k$ and $M = N - k$ are nonnegative integers,*

$$c_N(j, k) = \frac{N!}{k! d!} \gamma(k, d, M).$$

Proof. Choose the k th term from the first exponential and the d th term from the second. Their product is

$$\frac{x^{2k+2d} w^k (2 + w)^k \varphi(w)^d (\log x)^k}{k! d!}.$$

Extracting $w^N = \xi^N / x^N$ yields the power $x^{2k+2d-N} = x^j$ and the coefficient $[w^{N-k}](2 + w)^k \varphi(w)^d$. Multiplication by $N!$ gives the claim. $\square$

(The proof is the one printed for Proposition 3.24, itself the Taylor-expansion proof of Proposition 3.4.) The global support conjecture can therefore be stated entirely in terms of γ . For $d \geq 1$, the predicted zeros are exactly $M = k + 4d + 1$, $k + d \equiv 0 \pmod{2}$; at $d = 3$ this becomes $M = k + 13$, k odd.

3.28.1 A falling-factorial transform

Define $\mathcal{P}_M(x) = \prod_{\ell=1}^M (x - \ell)$, with $\mathcal{P}_0(x) = 1$. The following elementary identity is the algebraic entrance to the proof.

Lemma 3.49 (Coefficient as a parameter derivative). *For $M, d \geq 0$ and an indeterminate θ ,*

$$M! [w^M](1 + w)^\theta \log^d(1 + w) = \mathcal{P}_M^{(d)}(\theta + 1).$$

Proof. Since $[w^M](1 + w)^\theta = \binom{\theta}{M} = \mathcal{P}_M(\theta + 1)/M!$, differentiate d times with respect to θ . $\square$

(The manuscript's indeterminate is a ; it is θ here because a is the term count. With $\theta = s$ this is (3.58).) Because $2 + w = 1 + (1 + w)$, $(2 + w)^k = \sum_{\ell=0}^k \binom{k}{\ell} (1 + w)^\ell$. Taking $\theta = 2d + \ell$ in Lemma 3.49 gives

$$M! \gamma(k, d, M) = \sum_{\ell=0}^k \binom{k}{\ell} \mathcal{P}_M^{(d)}(2d + 1 + \ell). \quad (3.90)$$

Introduce $\mathcal{H}_{M,k}(x) = \sum_{\ell=0}^k \binom{k}{\ell} \mathcal{P}_M(x + \ell)$. Then

$$M! \gamma(k, d, M) = \mathcal{H}_{M,k}^{(d)}(2d + 1). \quad (3.91)$$

Suppose now that $r = M - k \geq 0$. Every summand $\mathcal{P}_M(x + \ell)$ is divisible by $\mathcal{P}_r(x)$, and direct cancellation gives

$$\mathcal{H}_{M,k}(x) = \mathcal{P}_r(x) \tilde{\mathcal{Q}}_{r,k}(x), \quad (3.92)$$

where

$$\tilde{\mathcal{Q}}_{r,k}(x) = \sum_{\ell=0}^k \binom{k}{\ell} x^{\bar{\ell}} (x - M + \ell)^{\overline{k-\ell}}. \quad (3.93)$$

Here $x^{\bar{\ell}} = x(x+1) \cdots (x+\ell-1)$. (This is the factorization (3.60) at $s = x - 1$, reached by direct cancellation instead of the formal residue substitution; so $\tilde{\mathcal{Q}}_{r,k}(x) = \mathcal{Q}_k^{(r)}(x - \frac{r+1}{2})$.)

3.29 Second route to depth three: the beta-sine transform and centred moments

The quotient polynomial has a useful integral representation.

Lemma 3.50 (Beta representation). *Let $r \geq 0$, $k \geq 0$, $M = r + k$, and $0 < \Re x < r + 1$. Then*

$$\tilde{\mathcal{Q}}_{r,k}(x) = \frac{M!}{\Gamma(x)\Gamma(r-x+1)} \int_0^1 t^{x-1} (1-t)^{r-x} (2t-1)^k dt.$$

Proof. Expand $(2t-1)^k = (t - (1-t))^k = \sum_{\ell=0}^k (-1)^{k-\ell} \binom{k}{\ell} t^\ell (1-t)^{k-\ell}$. Termwise beta integration gives $B(x+\ell, M-x-\ell+1)$. After multiplication by $M! / [\Gamma(x)\Gamma(r-x+1)]$, the first gamma ratio is $x^{\bar{\ell}}$, while the second, together with $(-1)^{k-\ell}$, is $(x-M+\ell)^{\overline{k-\ell}}$. The sum is exactly (3.93). $\square$

Using $\mathcal{P}_r(x) = \Gamma(x)/\Gamma(x-r)$ and Euler's reflection formula yields

$$\mathcal{H}_{M,k}(x) = (-1)^r \frac{M! \sin(\pi x)}{\pi} \int_0^1 t^{x-1} (1-t)^{r-x} (2t-1)^k dt. \quad (3.94)$$

(This is the mechanism of Lemma 3.15, applied to $\mathcal{H}_{M,k}$ rather than to the finite difference of that lemma.)

Define the kernel polynomial

$$R_d(\tau) = \frac{d^d}{d\varepsilon^d} \left(\frac{\sin(\pi\varepsilon)}{\pi} e^{\tau\varepsilon} \right) \Big|_{\varepsilon=0} = \frac{(\tau + i\pi)^d - (\tau - i\pi)^d}{2i\pi}. \quad (3.95)$$

Equivalently, $R_d(\tau) = \sum_{\mu \geq 0} (-1)^\mu \binom{d}{2\mu+1} \pi^{2\mu} \tau^{d-2\mu-1}$. (This is the polynomial R_d of (3.64), the manuscript's K_D ; the summation index is μ because m is the deficit.)

Theorem 3.51 (Beta-sine transform; second route to Proposition 3.30). *Let $d, k, M \geq 0$, set $r = M - k$, and assume $r \geq 2d + 1$. Then*

$$\gamma(k, d, M) = (-1)^{r+2d+1} \int_0^1 t^{2d}(1-t)^{r-2d-1}(2t-1)^k R_d\left(\log \frac{t}{1-t}\right) dt. \quad (3.96)$$

Proof. Differentiate (3.94) d times at the integer $x = 2d + 1$ and use (3.91). Near $2d + 1$, write $x = 2d + 1 + \varepsilon$ and

$$t^{x-1}(1-t)^{r-x} = t^{2d}(1-t)^{r-2d-1} \exp\left(\varepsilon \log \frac{t}{1-t}\right).$$

Also $\sin(\pi(2d + 1 + \varepsilon)) = (-1)^{2d+1} \sin(\pi\varepsilon)$. The derivative in ε is exactly $(-1)^{2d+1} R_d(\tau)$, giving (3.96). $\square$

Here $r - 2d - 1 = q$, so the sign is $(-1)^q$, and the substitution $t \mapsto 1 - t$ turns (3.96) into (3.65); the batch-42 proof deforms a Cauchy contour onto the branch cut, this one differentiates a beta integral. The first kernels are

$$R_1(\tau) = 1, \quad R_2(\tau) = 2\tau, \quad R_3(\tau) = 3\tau^2 - \pi^2, \quad R_4(\tau) = 4\tau^3 - 4\pi^2\tau.$$

Thus the transform recovers the one-sign depth-one integral and the antisymmetric logarithmic depth-two integral described in the ProveIt notes. At depth three the kernel is even but changes sign, explaining why the previous pointwise positivity strategy stops precisely there. (This is the obstruction (3.33) of Section 3.7, whose kernel is $R_3/3!$.)

3.29.1 Centred beta-moment polynomials

Fix $r \geq 1$. The manuscript writes the centred quotient $\tilde{Q}_{r,k}(\frac{r+1}{2} + y)$ as a new polynomial $R_{N,k}(y)$; by (3.60) it is $Q_k^{(r)}(y)$, and it is printed so. These polynomials contain the decisive extra structure.

Theorem 3.52 (Positive three-term recurrence; second proof of (3.61)). *For every $r \geq 1$, $Q_0^{(r)}(y) = 1$, $Q_1^{(r)}(y) = 2y$, and for $k \geq 1$*

$$Q_{k+1}^{(r)}(y) = 2y Q_k^{(r)}(y) + k(r+k) Q_{k-1}^{(r)}(y). \quad (3.97)$$

Consequently $Q_k^{(r)}$ has parity k , and every nonzero coefficient in the basis $y^k, y^{k-2}, \dots$ is a positive integer.

Proof. Let $\mathcal{I}_k(x) = \int_0^1 t^{x-1}(1-t)^{r-x}(2t-1)^k dt$. Integrate the derivative of $t^x(1-t)^{r-x+1}(2t-1)^k$. The boundary terms vanish in the beta strip. With $y = x - (r+1)/2$, the resulting identity is

$$(r+k+1) \mathcal{I}_{k+1} = 2y \mathcal{I}_k + k \mathcal{I}_{k-1}.$$

Multiplying by the factorial prefactor in Lemma 3.50 gives (3.97). The parity and coefficient positivity follow by induction. $\square$

(The batch-42 proof of (3.61) compares coefficients in a first-order differential equation for the generating function (3.59). The integer and positivity clauses sharpen that proposition's "nonnegative coefficients" only in wording: the recurrence has integer coefficients.)

Corollary 3.53 (Vertical-line roots; second route to Corollary 3.27). *Every zero of $\tilde{Q}_{r,k}$ lies on the vertical line $\Re x = (r+1)/2$. The zeros are simple, and consecutive degrees interlace after rotation to the real axis.*

Proof. Put $\mathcal{T}_{r,k}(s) = i^{-k} \mathcal{Q}_k^{(r)}(is)$. After monic normalization, (3.97) becomes a Favard recurrence with positive subdiagonal coefficient $k(r+k)/4$. Hence $\mathcal{T}_{r,k}$ has simple real zeros with the standard interlacing property. Rotating back gives the claim. $\square$

This root geometry is not required for the finite certificate, but it explains why the centred recurrence is so stable and suggests a classical orthogonal-polynomial interpretation worth developing further. (The monic polynomials of the proof are the p_k of the proof of Corollary 3.27, which reaches simplicity and the imaginary axis through a tridiagonal matrix; the interlacing clause is new to this part. The classical interpretation is (3.62).)

3.30 Second route to depth three: the harmonic phase transition

Set $d = 3$. In the tail range write $M = k + q + 7$, $q \geq 0$, $r = M - k = q + 7$. The evaluation point in (3.91) is $x = 7$, which is a root of $\mathcal{P}_r$. Factor $\mathcal{P}_r(x) = (x - 7)\Omega_r(x)$. At $x = 7$,

$$\Omega_r(7) = (-1)^q 6! q!, \quad (3.98)$$

$$\frac{\Omega'_r(7)}{\Omega_r(7)} = H_6 - H_q =: u_q, \quad (3.99)$$

$$\frac{\Omega''_r(7)}{\Omega_r(7)} = (H_6 - H_q)^2 - H_6^{(2)} - H_q^{(2)} =: 2E_q. \quad (3.100)$$

Here $H_q = \sum_{\ell=1}^q 1/\ell$, $H_q^{(2)} = \sum_{\ell=1}^q 1/\ell^2$, $H_0 = H_0^{(2)} = 0$. Thus

$$E_q = \frac{(H_q - H_6)^2 - H_q^{(2)} - H_6^{(2)}}{2}. \quad (3.101)$$

Let $y_q = 7 - \frac{r+1}{2} = \frac{6-q}{2}$. Using $\mathcal{H}_{M,k} = \mathcal{P}_r \tilde{\mathcal{Q}}_{r,k} = (x - 7)\Omega_r \tilde{\mathcal{Q}}_{r,k}$, one has $\mathcal{H}_{M,k}'''(7) = 3(\Omega_r \tilde{\mathcal{Q}}_{r,k})''(7)$. Define, with $r = q + 7$ and $\mathcal{Q} = \mathcal{Q}_k^{(r)}$,

$$F_k(q) = \frac{1}{2} (\mathcal{Q}''(y_q) + 2u_q \mathcal{Q}'(y_q) + 2E_q \mathcal{Q}(y_q)). \quad (3.102)$$

Then (3.98) gives the exact normalization

$$\boxed{\gamma(k, 3, k + q + 7) = \frac{4320(-1)^q q!}{(k + q + 7)!} F_k(q).} \quad (3.103)$$

(With (3.49) this is the dictionary (3.89), $H_{3,k,q} = 720(-1)^q q! F_k(q)$.)

3.30.1 The signed harmonic quantity

The E_q sequence has a remarkably simple first difference.

Lemma 3.54 (Unique harmonic phase transition). *For $q \geq 0$,*

$$E_{q+1} - E_q = \frac{H_q - H_6}{q + 1}. \quad (3.104)$$

Moreover, $E_0 > 0$, $E_q < 0$ for $1 \leq q \leq 37$, and $E_q > 0$ for $q \geq 38$.

Proof. Substitute $H_{q+1} = H_q + 1/(q+1)$ and $H_{q+1}^{(2)} = H_q^{(2)} + 1/(q+1)^2$ in (3.101); the square and second-order increments cancel to give (3.104).

Hence E_q strictly decreases through $q = 6$, has $E_6 = E_7$, and strictly increases thereafter. It remains only to locate the two crossings. Exact arithmetic gives

$$E_0 = \frac{203}{90} > 0, \quad E_1 = -\frac{7}{36} < 0,$$

$$E_{37} = -\frac{43927248584204635973}{2111705758564129728000} < 0, \quad E_{38} = \frac{31458788673623539948663}{1243794691794272409792000} > 0.$$

Monotonicity supplies all intervening and subsequent signs. $\square$

For $k = 0$, (3.103) and $F_0(q) = E_q$ already prove an infinite new nonvanishing diagonal. More importantly, the same phase transition controls all k through the recurrence of Theorem 3.52.

Editorial note. The $k = 0$ diagonal is not new as a nonvanishing statement: it is the $k = 0$ row of Theorem 3.23 at $d = 3$, and the formula behind it is the $D = 3$ case of the elementary-symmetric formula in the notes ((3.110) below). What the lemma adds is the exact sign law. Through (3.89) and the sign rule (3.70), the reduced expectation of the $d = 3$, $k = 0$ row has the sign of E_q ; so the lemma *proves* the observation of Subsection 3.20 that this expectation “is positive at $q = 0$, negative for $1 \leq q \leq 37$, and positive for all $q \geq 38$ ”, which the batch-42 addition read off its certificate data.

3.31 Second route to depth three: the complete tail and the full classification

3.31.1 Automatic positivity outside the critical strip

For $q < 6$, put $\lambda_q = \frac{6-q}{2} > 0$ and $\alpha_q = H_6 - H_q > 0$. For $q > 6$, put $\lambda_q = \frac{q-6}{2} > 0$ and $\alpha_q = H_q - H_6 > 0$. By parity of $\mathcal{Q}_k^{(r)}$, after removing the factor $(-1)^k$ when $q > 6$, the relevant expression is

$$U_{q,k} = \mathcal{Q}''(\lambda_q) + 2\alpha_q \mathcal{Q}'(\lambda_q) + 2E_q \mathcal{Q}(\lambda_q), \quad \mathcal{Q} = \mathcal{Q}_k^{(q+7)}. \quad (3.105)$$

More precisely,

$$2F_k(q) = \begin{cases} U_{q,k}, & q < 6, \\ (-1)^k U_{q,k}, & q > 6. \end{cases} \quad (3.106)$$

Since $\mathcal{Q}_k^{(r)}$ has positive coefficients, $\mathcal{Q}_k^{(r)}(\lambda) > 0$ and its derivatives are nonnegative for $\lambda > 0$. Therefore Lemma 3.54 immediately gives $U_{0,k} > 0$ and $U_{q,k} > 0$ for $q \geq 38$, for every k .

3.31.2 Positivity propagation in the finite strip

For $q \in \mathcal{C} := \{1, \dots, 5\} \cup \{7, \dots, 37\}$, $E_q < 0$, and with $r = q + 7$ fixed the functional reads $U_k := U_{q,k} = \mathcal{Q}_k''(\lambda_q) + 2\alpha_q \mathcal{Q}_k'(\lambda_q) - 2|E_q| \mathcal{Q}_k(\lambda_q)$, where $\mathcal{Q}_k = \mathcal{Q}_k^{(r)}$ (the manuscript's U_k , with $B = |E_q|$). Applying this differential operator to (3.97) gives

$$U_{k+1} = 2\lambda_q U_k + k(r+k)U_{k-1} + 4(\mathcal{Q}_k'(\lambda_q) + \alpha_q \mathcal{Q}_k(\lambda_q)). \quad (3.107)$$

The final term is strictly positive. Consequently, once two consecutive U_k are positive, every later U_k is positive.

The only remaining work is therefore finite.

Proposition 3.55 (Exact critical-strip certificate). *For every $q \in \mathcal{C}$:*

- (1) $U_{q,k} \neq 0$ for $0 \leq k \leq 14$;
 (2) $U_{q,13} > 0$ and $U_{q,14} > 0$.

At $q = 6$, $F_k(6) = 0$ for odd k , while $F_k(6) \neq 0$ for even $0 \leq k \leq 14$, and $F_{14}(6) > 0$.

Proof. This is the finite exact component of the proof. Starting from $Q_0 = 1$, $Q_1 = 2y$, the verifier constructs every $Q_k^{(r)}$ by (3.97) with integer coefficients. It evaluates (3.105) at the half-integer $\lambda_q = |q - 6|/2$ using `fractions.Fraction`; harmonic numbers are also exact fractions. The compressed sign results are displayed in Table 6. The supplied file `code/03-depth-three-verify_depth3.py` (delivered as `verify_depth3.py`) performs all assertions independently of any computer algebra system. $\square$

Table 6: Exact signs of $F_k(q)$. Intervals contain integers; “ $\geq r_0$ ” means all $q \geq r_0$. Rows $k = 13, 14$ are the induction anchors.

k	$F_k(q) > 0$	$F_k(q) < 0$	$F_k(q) = 0$
0	$q = 0$; $q \geq 38$	$1 \leq q \leq 37$	—
1	$0 \leq q \leq 1$; $7 \leq q \leq 34$	$2 \leq q \leq 5$; $q \geq 35$	$q = 6$
2	$0 \leq q \leq 1$; $q \geq 33$	$2 \leq q \leq 32$	—
3	$0 \leq q \leq 2$; $7 \leq q \leq 29$	$3 \leq q \leq 5$; $q \geq 30$	$q = 6$
4	$0 \leq q \leq 2$; $q \geq 28$	$3 \leq q \leq 27$	—
5	$0 \leq q \leq 3$; $7 \leq q \leq 24$	$4 \leq q \leq 5$; $q \geq 25$	$q = 6$
6	$0 \leq q \leq 3$; $q \geq 23$	$4 \leq q \leq 22$	—
7	$0 \leq q \leq 3$; $7 \leq q \leq 20$	$4 \leq q \leq 5$; $q \geq 21$	$q = 6$
8	$0 \leq q \leq 4$; $q \geq 18$	$5 \leq q \leq 17$	—
9	$0 \leq q \leq 4$; $7 \leq q \leq 15$	$q = 5$; $q \geq 16$	$q = 6$
10	$0 \leq q \leq 4$; $q \geq 13$	$5 \leq q \leq 12$	—
11	$0 \leq q \leq 5$; $7 \leq q \leq 10$	$q \geq 11$	$q = 6$
12	$0 \leq q \leq 5$; $q \geq 9$	$6 \leq q \leq 8$	—
13	$0 \leq q \leq 5$	$q \geq 7$	$q = 6$
14	all $q \geq 0$	—	—

(The manuscript writes the threshold in the caption as r ; it is r_0 here because $r = M - k$. The verifier prints these signs for $0 \leq q \leq 38$; the rows’ behaviour for $q \geq 38$ is the analytic statement $U_{q,k} > 0$ above, through (3.106).)

3.31.3 The centre $q = 6$

When $q = 6$, the evaluation point is $y = 0$, and $u_6 = 0$. By parity, $Q_k^{(13)}(0) = Q_k^{(13)'}(0) = 0$ for odd k , hence $F_k(6) = 0$. For even k , set $V_k = Q_k^{(13)''}(0) + 2E_6 Q_k^{(13)}(0) = 2F_k(6)$. At $\lambda = \alpha = 0$, (3.107) reduces, for odd k , to

$$V_{k+1} = k(13 + k)V_{k-1} + 4Q_k^{(13)'}(0).$$

The derivative term is positive. The finite certificate gives $V_{14} > 0$, so every even V_k with $k \geq 14$ is positive. The even cases below 14 are nonzero by the same certificate. Thus the centre has exactly the odd- k zeros.

Theorem 3.56 (Complete depth-three tail). *For all $k, q \geq 0$,*

$$\gamma(k, 3, k + q + 7) = 0 \iff q = 6 \text{ and } k \text{ is odd.}$$

Equivalently, in the tail $M \geq k + 7$, $\gamma(k, 3, M) = 0$ if and only if $M = k + 13$ and k is odd.

Proof. For $q = 0$ and $q \geq 38$, positivity follows directly from Lemma 3.54 and (3.106). For $q \in \mathcal{C}$, the two positive anchors in Proposition 3.55 and recurrence (3.107) imply $U_{q,k} > 0$ for all $k \geq 13$; the smaller k are nonzero by the finite certificate. The centre $q = 6$ was handled above. Finally use (3.103), whose prefactor never vanishes. $\square$

Corollary 3.57 (Stable tail signs). *For even $k \geq 14$, $F_k(q) > 0$ for every $q \geq 0$. For odd $k \geq 13$, $F_k(q) > 0$ for $q < 6$, $F_k(6) = 0$, and $F_k(q) < 0$ for $q > 6$.*

3.31.4 Bulk positivity and the full depth-three theorem

It remains to treat $3 \leq M \leq k + 6$. Multiplication by $2 + w$ gives the Pascal-type recurrence

$$\gamma(k+1, 3, M) = 2\gamma(k, 3, M) + \gamma(k, 3, M-1), \quad (3.108)$$

the case $d = 3$ of (3.63). For $k = 0$ and $3 \leq M \leq 6$, $M! \gamma(0, 3, M) = \mathcal{P}_M'''(7) > 0$, because every factor $7 - \ell$ is positive and the product-rule expansion of the third derivative has positive summands. (This is the “ $k = 0$ bulk” of the notes [22].) The tail anchor $\gamma(k, 3, k+7) > 0$ for every k follows from $q = 0$ in Theorem 3.56 together with $F_k(0) > 0$.

Lemma 3.58 (Bulk positivity; second route to Theorem 3.21 at $d = 3$). *For every $k \geq 0$ and $3 \leq M \leq k + 6$, $\gamma(k, 3, M) > 0$.*

Proof. Induct on k . The case $k = 0$ was just proved. Suppose the claim holds for k . For $3 \leq M \leq k + 6$, both terms on the right of (3.108) are nonnegative and the first is positive. At the new endpoint $M = k + 7$, the two terms are $2\gamma(k, 3, k+7) > 0$ and $\gamma(k, 3, k+6) > 0$. This proves the range $3 \leq M \leq (k+1) + 6$. $\square$

Theorem 3.59 (Complete depth-three zero classification; second proof of Theorem 3.23 at $d = 3$). *For every $k \geq 0$ and $M \geq 3$,*

$$\boxed{\gamma(k, 3, M) = 0 \iff M = k + 13 \text{ and } k \text{ is odd.}} \quad (3.109)$$

Proof. The bulk $M \leq k + 6$ is strictly positive by Lemma 3.58. The complementary range $M \geq k + 7$ is Theorem 3.56. $\square$

Corollary 3.60 (Depth three in the original coefficient lattice). *Let N, k be nonnegative integers with $N \geq k + 3$, and put $j = 2k + 6 - N$, so that $(N + j)/2 - k = 3$. Then*

$$c_N(j, k) = 0 \iff N = 2k + 13 \text{ and } k \text{ is odd.}$$

In the zero case $j = -7$. Thus the depth-three holes are exactly $c_{4\ell+15}(-7, 2\ell+1) = 0$, $\ell \geq 0$. Every other depth-three coefficient is nonzero.

Proof. Apply Proposition 3.48 and Theorem 3.59 with $M = N - k$. The positive factor $N!/(k!3!)$ does not affect vanishing. The equation $N - k = k + 13$ is $N = 2k + 13$, and substituting into $j = 2k + 6 - N$ gives $j = -7$. $\square$

This is exactly the hole predicted by the repository’s global support model at $d = 3$: the hole line is $M = k + 4d + 1$, and the parity condition $k + d$ even is k odd.

Remark 3.61 (The two proofs of depth three compared; editorial). Theorem 3.59 and the case $d = 3$ of Theorem 3.23 share the factorization (3.60) and the recurrence (3.61), and differ in how they control the tail. Batch 42 folds the branch-cut integral onto the half interval, compares likelihood ratios against a symmetric base, and needs three seed signs and the rectangle $0 \leq k \leq 12$, $0 \leq q \leq 37$ (494 integer cells, Proposition 3.41 with $(R, Q) = (14, 38)$). Manuscript 03 factors the root $x = 7$ out of $\mathcal{P}_r$, reads the tail sign off the harmonic quantity E_q , and propagates positivity in k by (3.107); it needs exact rational values for the 36 values of q in the strip $\mathcal{C}$ with $0 \leq k \leq 14$, plus the centre $q = 6$ (Proposition 3.55). The finite anchors partly coincide. By (3.89), the batch-42 central seed $H_{3,14,6} = 9707604961443840000 > 0$ is $720^2 F_{14}(6)$ with $F_{14}(6) = 18726089817600$, the manuscript's centre anchor $V_{14} > 0$; the batch-42 sum-tail seed $H_{3,0,38} > 0$ is $720 \cdot 38! E_{38} > 0$, the manuscript's harmonic anchor; and the batch-42 difference-tail seed $H_{3,1,38} < 0$ is $F_1(38) < 0$, which in the manuscript is automatic from $E_{38} > 0$. The two finite computations also agree where they overlap: through (3.89) and the sign rule (3.70), the rows $0 \leq k \leq 12$ of Table 6 reproduce every $d = 3$ row of `data/02-depth-certificates-sign_thresholds.csv` (checked at the intake). Neither proof is formalized.

3.32 Second route to depth three: further exact consequences

3.32.1 A closed harmonic formula for the log-free diagonal

For $k = 0$ and $M = q + 7$, the reciprocal multiset at the root $x = 7$ is $1, \frac{1}{2}, \dots, \frac{1}{6}, -1, -\frac{1}{2}, \dots, -\frac{1}{q}$. Its second elementary symmetric function is E_q . Therefore

$$\gamma(0, 3, q + 7) = \frac{4320(-1)^q q!}{(q + 7)!} E_q. \quad (3.110)$$

In the original derivative expansion, for $N \geq 7$,

$$c_N(6 - N, 0) = 360(-1)^{N-7}(N - 7)! \left((H_{N-7} - H_6)^2 - H_{N-7}^{(2)} - H_6^{(2)} \right). \quad (3.111)$$

The sign changes of E_q yield an exact phase diagram: $\gamma(0, 3, M) > 0$ for $3 \leq M \leq 7$; $\text{sgn } \gamma(0, 3, M) = (-1)^M$ for $8 \leq M \leq 44$; $\text{sgn } \gamma(0, 3, M) = (-1)^{M+1}$ for $M \geq 45$. The two consecutive positive values at $M = 44, 45$ mark the phase reversal.

(Formula (3.110) is the case $D = 3$ of the formula

$$\gamma(0, D, M) \frac{M!}{D!} = (-1)^q (2D)! q! e_{D-1} \left(1, \dots, \frac{1}{2D}, -1, \dots, -\frac{1}{q} \right), \quad q = M - 2D - 1,$$

of the notes [22]; the phase diagram is new. The intake rechecked (3.110) for $q \leq 123$ and the phase diagram for $M \leq 130$ in exact arithmetic.)

3.32.2 A structural explanation of the holes

At $q = 6$, the centre of the vertical root line in Corollary 3.53 is exactly the evaluation point $x = 7$. The polynomial $\mathcal{Q}_k^{(13)}$ has parity k , while the depth-three differential functional at the centre uses only even derivatives. It therefore annihilates precisely the odd degrees. The symmetry-forced hole is not an accidental cancellation of large rational numbers; it is the parity centre of the beta-moment family. (This is the parity argument of Theorem 3.29 at $d = 3$, and so the odd-falling-factorial symmetry of Section 5.4.)

3.32.3 What this does and does not imply for A290268

The theorem proves the conjectured support description on the entire $d = 3$ layer. It does not yet determine all of $a(N)$, because a coefficient of the N th derivative may have depth as large as N . To prove the OEIS formula, one still needs the corresponding nonvanishing theorem for every $d \geq 4$ (or a global argument that treats all depths simultaneously).

Editorial note. In this report the depth-four theorem is Theorem 3.23; what is still needed is every $d \geq 5$, the region $m \geq 5$, $n \geq 2k + 2$ of Section 3.10. (*Update, 1 October 2026:* depths five and six are Theorem 3.71; what is still needed is every $d \geq 7$. *Update, batch 73:* depths seven and eight are Theorems 3.99 and 3.100; what is still needed is every $d \geq 9$.)

3.33 Second route to depth three: certificate, formalization blueprint and research questions

3.33.1 Exact certificate and reproducibility

The manuscript’s artifact package contained its source and PDF, the verifier `verify_depth3.py`, the recorded output `depth3_certificate.txt`, and a README. Of these, this report ships the verifier and its recorded output, byte-identical to the delivery, as `code/03-depth-three-verify_depth3.py` and `data/03-depth-three-depth3_certificate.txt` (Section 3.34).

The verifier uses no third-party package. Its proof-critical path is:

1. construct $\mathcal{Q}_k^{(r)}$ by the integer recurrence (3.97);
2. compute $H_q, H_q^{(2)}, E_q$ with `Fraction`;
3. evaluate $F_k(q)$ and $U_{q,k}$ exactly in the finite strip;
4. assert the signs and induction anchors in Proposition 3.55.

As independent checks, the script also verifies the normalized F recurrence on $0 \leq k \leq 60$, $0 \leq q \leq 1000$; extracts $[w^M](2 + w)^k \varphi(w)^3$ directly for $0 \leq k \leq 24$, $3 \leq M \leq 180$; and confirms agreement with (3.103) and the complete zero classification throughout that box. These large-box checks are diagnostic rather than logical substitutes for the infinite proof.

The recorded certificate-body digest is

2e284dcc4a8cb79925716f96401b49b0a5f3c523437bdab8399b1ee7133bded9.

A successful run exits with status 0 and rewrites the certificate deterministically. The manuscript’s rerun command is `python3 verify_depth3.py`; with the shipped name it is

```
py code/03-depth-three-verify_depth3.py
```

run on a copy of the report directory: the program writes `depth3_certificate.txt`, under that unprefix name, into the directory that contains it, here `code/`. Its checks are Python `assert` statements, so it must not be run with `python -O`. At the intake it was rerun on a copy: exit status 0, and the rewritten certificate is byte-identical to the shipped `data/03-depth-three-depth3_certificate.txt`.

No hidden trust upgrade (the manuscript’s warning box). The exact Python certificate is substantially stronger than a floating-point scan, but it is not a kernel-checked theorem. The mathematical reduction to a finite sign problem must still be reviewed, and a future ProveIt integration should formalize both the recurrence and the finite checker in Lean or Rocq.

3.33.2 Formalization blueprint for ProveIt

The proof was designed to split cleanly along formalization boundaries. The manuscript suggests the following module structure.

1. **A290268.SeriesReduction**: formalize Proposition 3.48 in formal power series and connect it to the existing lattice recurrence.
2. **A290268.FallingFactorial**: define $\mathcal{P}_M$, prove Lemma 3.49, and derive (3.91).
3. **A290268.CenteredMoments**: define $\tilde{Q}_{r,k}$ algebraically by (3.93) and prove (3.97) without analytic integration, by polynomial identities or coefficient induction.
4. **A290268.DepthThreeHarmonic**: formalize E_q , (3.104), and the four exact anchor inequalities in Lemma 3.54.
5. **A290268.DepthThreeCertificate**: reflect the finite sign table into a Boolean checker and discharge it with `native_decide` or a kernel-reduced rational computation, according to the project’s trust policy.
6. **A290268.DepthThree**: combine positivity propagation, centre parity, bulk induction, and Theorem 3.59.

A natural public endpoint is schematically

```
theorem gamma_depth_three_eq_zero_iff
  (k M : Nat) (hM : 3 <= M) :
  gamma k 3 M = 0 <-> M = k + 13 /\ Odd k
```

followed by a bridge to the existing coefficient array:

```
theorem coeff_depth_three_ne_zero_unless_hole
  (n k : Nat) (h : k + 3 <= n) :
  coeff n (2*k + 6 - n) k = 0 <->
  n = 2*k + 13 /\ Odd k
```

The actual types will need to respect the repository’s integer exponent index and natural log-degree conventions.

Avoiding analytic overhead. The beta–sine transform is conceptually useful but not necessary for the Lean proof of depth three. The formal path can remain algebraic: $\gamma \rightarrow \mathcal{P}_M^{(3)} \rightarrow \mathcal{P}_r \tilde{Q}_{r,k} \rightarrow \mathcal{Q}_k^{(r)}$ recurrence $\rightarrow E_q, U_{q,k}$. This avoids formal integration, gamma functions, and complex analysis. The integral theorem can be formalized separately as a reusable analytic layer.

Editorial note. None of the six modules exists; the manuscript proposes them and does not claim them. The library A290268 has the modules **Core**, **ClosedForm**, **Count**, **Structural**, **Table** and **Main** (Section 3.34.3); it defines no `gamma`, and its `coeff` has type $\mathbb{N} \rightarrow \mathbb{Z} \rightarrow \mathbb{Z} \rightarrow \mathbb{Z}$, so the sketched `coeff n (2*k + 6 - n) k` needs the integer index the last sentence asks for. The same algebraic interface is Question 7 of Subsection 3.24.

3.33.3 Further research questions

The proof suggests a focused continuation program. The manuscript’s nine questions are printed as it states them; the editorial notes after seven of them record what this report already contains.

Research question 3.62 (Depth four). Can the same centred-polynomial method classify all zeros at $d = 4$? The beta–sine kernel is $R_4(\tau) = 4\tau^3 - 4\pi^2\tau$, and the root reduction involves a

third-order differential functional on $\mathcal{Q}_k^{(r)}$. The main challenge is to find a finite-dimensional positivity cone stable under the recurrence.

Editorial note: already answered as a classification. All zeros at $d = 4$ are classified by Theorem 3.23 (only $q = 8$, k even; $j = -9$, $N = 2k + 17$), which was in this report at the manuscript's pin. What the question still asks is only methodological: whether the centred-polynomial route of this addition also proves depth four, perhaps with a smaller finite certificate than the 15,540 cells of the batch-42 rectangle.

Research question 3.63 (Uniform depth induction). For general d , evaluation at the root $x = 2d + 1$ produces an order- $(d - 1)$ differential functional of $\tilde{\mathcal{Q}}_{r,k}$. Is there a uniform matrix recurrence whose invariant cone proves that only the parity-centre holes occur?

Editorial note. This is a proposed route to Question 4 of Subsection 3.24, which is open.

Research question 3.64 (Critical-strip bounds). At $d = 3$, positivity becomes automatic outside the finite harmonic strip $1 \leq q \leq 37$, and two consecutive k -anchors propagate all larger degrees. Can one bound the analogous strip and anchor degree explicitly as functions of d ?

Editorial note: partly answered. Explicit (deliberately crude) bounds K_d, Q_d as functions of d , beyond which every entry at depth d is nonzero except the holes, are Theorem 3.22 (3.74). Small or sharp bounds, for this route or that one, remain open; they are Questions 2 and 3 of Subsection 3.24. (*Update, 1 October 2026:* Theorem 3.81 makes both bounds singly exponential, $\hat{Q}_d = \lceil (2d + 1)e^d \rceil$ for the strip in q and $\hat{K}_d = 2\hat{Q}_d$ for the degree in k ; at $d = 3$ these are 141 and 282, against the strip $1 \leq q \leq 37$ and anchor degree 14 here. Sharp bounds remain open.)

Research question 3.65 (Classical identification). Identify $\mathcal{Q}_k^{(r)}$ precisely within the continuous Hahn or related orthogonal-polynomial schemes, including its measure after the rotation $y \mapsto iy$. Can known connection coefficients or total-positivity theorems replace the finite certificate?

Editorial note: the identification is answered. By (3.62), $\mathcal{Q}_k^{(r)}(y) = k! i^k P_k^{((r+1)/2)}(-iy; \pi/2)$, a Meixner–Pollaczek polynomial (a member of the Hahn class of the DLMF [20]), so its orthogonality measure after the rotation is the classical Meixner–Pollaczek weight with $\lambda = (r + 1)/2$, $\phi = \pi/2$. The second sentence, replacing the certificate, remains open.

Research question 3.66 (Eliminate the finite certificate). Find a human inequality proving the signs in Table 6, perhaps from bounds on the logarithmic derivatives of the rotated orthogonal polynomials. This would make Theorem 3.59 entirely symbolic.

Research question 3.67 (General hole theorem in the proof assistant). The repository notes give a clean $w = e^u - 1$ parity argument for $M = k + 4d + 1$. Formalizing that argument would make the structural inclusion of the conjectured support unconditional at every depth, leaving only nonvanishing.

Editorial note. On paper the hole theorem is proved three times in this report and the notes: Theorem 3.11, Theorem 3.29, and the $w = e^u - 1$ argument of the notes. The formal side is open: the hypothesis `hhole` of `Structural.mem_S_of_coeff_ne_zero` is exactly this statement, and the module `A290268.Hole` its docstring names does not exist (Section 3.26.3).

Research question 3.68 (Neighbouring derivative families). The notes report a sporadic zero in the analogous x^x family. Which functions x^{x^m} have only symmetry-forced holes, and for which m do sporadic cancellations occur? The beta–sine transform has a plausible m -parameter generalization.

Editorial note. The sporadic x^x zero is $b(8, 5) = 0$, (2.19), proved in Part 2; for x^{x^x} additional zeros provably exist (Part 4). The question for x^{x^m} , $m \geq 3$, is Question 8 of Subsection 3.24 (there written x^{x^r}); it is open. [Added 1 October 2026, batch 72A: partly answered; see the note after Question 8. The batch-72A intake scanned the recurrence of x^{x^m} for $m = 1, \dots, 5$ and $N \leq 70$ (the sibling report `power-tower-exponent-supports`, its Section 19.8): for $m = 2, \dots, 5$ every zero is structural or a reflection zero, and for $m = 1$ the only others are copies of $b(8, 5) = 0$ along one line. That is a finite observation, not a theorem. For $x^{\sqrt{x}}$ sporadic cancellations occur in two infinite families (its Theorem 14.3).]

Research question 3.69 (Asymptotic harmonic phases). For $k = 0$ and general d , the tail is governed by an elementary symmetric function of $1, \frac{1}{2}, \dots, \frac{1}{2d}, -1, -\frac{1}{2}, \dots, -\frac{1}{q}$. Determine the number and location of its sign transitions as $q \rightarrow \infty$. The $d = 3$ transition $37 \rightarrow 38$ is the first nontrivial case.

Editorial note (1 October 2026, batch 72A). This bears on the question without answering it. At every $d \geq 3$, up to the explicit factor in (3.70), the sign of the $k = 0$ tail is that of the reduced expectation $\mathcal{E}_{d,0,q}[\Pi_d(\tau^2)]$, and Theorem 3.81 shows that this expectation is positive for all $q \geq \lceil (2d+1)e^d \rceil$: every sign transition lies below that bound. The Mellin interpolant $\mathcal{F}_{d,0}$ has at most $d - 1$ real zeros (Theorem 3.44). At $d = 3$ the tail-moment formula of Remark 3.76 reproduces $2E_q$ and the transition $37 \rightarrow 38$ exactly. The precise number and location of the transitions at $d \geq 4$ remain open.

Research question 3.70 (Global A290268). Can the depthwise results be assembled into a single support theorem without proving each d separately? A successful uniform argument would immediately establish the OEIS generating function through the existing ProveIt counting reduction.

Editorial note. This is Questions 4 and 6 of Subsection 3.24; the counting reduction is Theorem 3.20 on paper and `Main.a_eq_closedForm_of_support_all` in Lean.

3.33.4 Conclusion of manuscript 03

The A290268 support problem admits more structure than a raw coefficient scan reveals. Parameter differentiation converts coefficient extraction into a falling-factorial derivative. Factoring the common roots exposes centred beta-moment polynomials with a positive three-term recurrence. At depth three, the remaining signed information collapses to one harmonic phase sequence and a recurrence-stable positivity functional. These ingredients prove the full zero classification $\gamma(k, 3, M) = 0 \iff M = k + 13$ and k odd, settling the first open depth in the pinned ProveIt program.

The result is both mathematically substantive and formalization-friendly. Its only finite component is explicit, exact, small, and independently reproducible. The next decisive test is depth four: success there would show whether the invariant-cone mechanism is a one-depth phenomenon or the start of a uniform proof of the full OEIS conjecture.

Editorial note. “Settling the first open depth” is mistaken: depth three was settled at the pin by Theorem 3.23, and this is a second proof. “The next decisive test is depth four” is a test of the *method* (Question 3.62 and its note); depth four itself is settled, and the first open depth is $d = 5$.

3.33.5 Core verifier logic (manuscript Appendix A)

The complete verifier is shipped with this report. The following excerpt shows the proof-critical recurrences; all quantities are exact fractions.


```

def E(q):
    a = H(q) - H(6)
    return (a*a - H(q, 2) - H(6, 2)) / 2

def centered_polynomials(N, max_k):
    R = [[1], [0, 2]]          # 1, 2*y
    for k in range(1, max_k):
        R.append(2*y_times(R[k]) + k*(N+k)*R[k-1])
    return R

def F_from_polynomial(k, q):
    N = q + 7
    y = Fraction(6-q, 2)
    u = H(6) - H(q)
    R = centered_polynomials(N, k)[k]
    return (R2(y) + 2*u*R1(y) + 2*E(q)*R0(y)) / 2

```

The actual source implements the polynomial operations explicitly and asserts all signs, recurrence identities, tail zeros, bulk positivity checks, and direct coefficient comparisons. (The excerpt keeps the manuscript's letters: N is r , R is $\mathcal{Q}^{(r)}$, H the harmonic numbers, and y_times , $R0$, $R1$, $R2$ stand for operations the full program spells out.)

3.33.6 Source and proof audit (manuscript Appendix B)

Item	Audit statement
Repository pin	All repository-status claims were checked against commit <code>a02ab3567fef25793b27a68248b58a7b69694342</code> . <i>Editorial: the claim that depth three was open at that commit was not; see Subsection 3.27.</i>
OEIS status	The public A290268 page was inspected on 30 September 2026; it still labels the generating function and closed form conjectural.
New theorem	Theorem 3.59 is proved in this article and is not claimed to exist in the pinned repository. <i>Editorial: it did exist there, as the case $d = 3$ of Theorem 3.23; the proof printed here is a second route.</i>
Analytic assumptions	The main depth-three proof can be read algebraically. The beta integral is used to discover and explain the centred recurrence, but the recurrence itself is a polynomial identity.
Finite computation	Only Proposition 3.55 is computer-checked. It uses exact rational arithmetic and is accompanied by source and recorded output.
Cross-checks	Larger exact boxes and direct series extraction are diagnostics, not premises of the infinite theorem.
Formal verification	No Lean/Rocq proof is claimed. A module-by-module formalization plan is given in Section 3.33.2.
Peer review	None claimed. The article should be treated as a detailed research proof requiring independent mathematical review.

3.34 Second route to depth three: provenance, files and formal status

3.34.1 Provenance

Subsections 3.27–3.34 were built from **one** manuscript: batch 70, manuscript 03, archive `ProveIt_A290268_Depth3_Article.zip` (*Depth Three Is Complete for OEIS A290268*, 17-page PDF, dated 30 September 2026), pinned to ProveIt commit `a02ab3567fef25793b27a68248b58a7b69694342`, which arrived in commit `1b3960d8a` and was placed in this report by commit `51c6943bf`. Its Section 1 and summary box form Subsection 3.27; its Sections 2–3 are Subsection 3.28, 4–5 are Subsection 3.29, 6 is Subsection 3.30, 7–8 are Subsection 3.31, 9 is Subsection 3.32; its Sections 10–13 and its two appendices form Subsection 3.33. Every theorem, proposition, lemma, corollary, definition, remark, table and research question of the manuscript is printed, with its proof, and every non-claim and limitation it states; none was weakened or strengthened. The editorial choices were these.

- It is placed as subsections at the end of Part 3, after the batch-42 addition, so that no existing section, theorem or equation number moves.
- Its main theorem, reduction, parameter derivative, beta–sine transform, recurrence, root corollary and bulk lemma are marked as second routes to results already printed in this part. Its claims that depth three was open at its pin and that its theorem is new there are kept verbatim and corrected by editorial notes, as is its statement that depth four remains open; its question “Depth four” is marked as answered by Theorem 3.23; six others carry notes relating them to this report, two of them (critical-strip bounds, classical identification) re-scoped as partly answered.
- Its letters are renamed as in Table 5, with no change of normalization; its coloured boxes are printed as quoted paragraphs; its file table is replaced by the shipped file list; its rerun command is rewritten to the shipped name; its OEIS citation is merged with this part’s. Its bibliography also lists Štampach’s paper on Bernoulli polynomials of the second kind, which its text never cites; it is the batch-42 reference [21].
- The dictionary (3.89), Remark 3.61, the editorial notes and the parenthetical translations are additions of the intake, not of the manuscript.

Not shipped: the manuscript’s source and PDF, `proveit_a290268_depth3.tex` and `proveit_a290268_depth3.pdf`; its `README.md`; and its checksum list `SHA256SUMS.txt` (verified at placement, 5 of 5 entries). The archive itself survives in the history of commit `1b3960d8a`.

3.34.2 Files

The manuscript’s code and data are shipped with the prefix `03-depth-three-`, byte-identical to the delivery:

Shipped path (delivery name)	Purpose
<code>code/03-depth-three-verify_depth3.py</code> (<code>verify_depth3.py</code>)	Exact standard-library verifier of Proposition 3.55 and the cross-checks of Section 3.33.1; writes <code>depth3_certificate.txt</code> beside itself.
<code>data/03-depth-three-depth3_certificate.txt</code> (<code>depth3_certificate.txt</code>)	Its recorded output: the anchors, the sign table for $0 \leq q \leq 38$, and the certificate-body digest.

3.34.3 Formal status

Nothing in this addition is formalized, and its placement beside a Lean development confers no formal status. The Lean library A290268 (Combinatorics/DerivativeExpansions/A290268/Lean) relates to it exactly as described in Section 3.26.3. The manuscript cites only A290268.Main, whose theorems `a_eq_closedForm_of_support`, `a_eq_closedForm_of_two_sided` and `a_eq_closedForm_of_support_all` are conditional on the support characterization; it reads them correctly. Theorem 3.59 would discharge the nonvanishing inclusion only on cells of depth three, which Theorem 3.23 already covers on paper; the Main theorems stay conditional. The six modules of Section 3.33.2 are proposals and do not exist.

3.35 Depths five and six, and count bounds

What this addition is. Subsections 3.35–3.41 were written into this part on 1 October 2026 from two later manuscripts of batch 72A, both dated 1 October 2026 and both pinned to this report’s text at commit `ca81647a9`:

- manuscript 70, *Depths Five and Six Nonvanishing for OEIS A290268* (Subsections 3.36–3.39);
- manuscript 68, *Pointwise and Summatory Bounds for OEIS A290268* (Subsection 3.40), written after manuscript 70; its pointwise bound uses 70’s Theorem 7.2, which is Theorem 3.81 here.

Provenance is in Subsection 3.41. They continue the depth coordinates of Subsections 3.14–3.26 and answer two of the research questions of Subsection 3.24. **The full A290268 conjecture is still not proved**, and neither is the leading constant $\frac{1}{2}$ of $a(N)$ in this report.

What is new relative to this part.

- *Depths five and six are classified* (Theorem 3.71): at $d = 5, 6$ the only zeros are the symmetry holes $M = k + 4d + 1$, $k + d$ even. This answers Question 1 of Subsection 3.24. The proof follows the route that question proposes: exact moment seeds for the quadratic criteria (3.88), and two finite rectangles of 343,151 and 4,785,360 cells. The open region of Conjecture 3.19 is therefore now $m \geq 7$, $n \geq 2k + 2$. [Added 1 October 2026, batch 73: depths seven and eight are classified in Subsection 3.42ff.; the open region is now $m \geq 9$.]
- *Singly exponential fixed-depth cutoffs* (Theorem 3.81): at every depth $d \geq 3$, the cutoffs $\hat{Q}_d = \lceil (2d + 1)e^d \rceil$ and $\hat{K}_d = 2\hat{Q}_d$ replace the tower-type constants (3.74) of Theorem 3.22. This answers Question 2, without a claim of optimality.
- *The largest-root affine minorant* (Proposition 3.80) replaces Lemma 3.36 at every depth, sharply among distribution-free criteria. This answers Question 3 in part: it is a first-moment criterion, not positivity in an orthogonal-polynomial basis.
- *An explicit pointwise bound* (Theorem 3.83): $a(N) \geq 3N^2/8 + (N/2)\log N - (N/2)\log \log N - 5N/4$ for every $N \geq e^{10}$. This is the first explicit bound with a gain of order $N \log N$ over the bulk; the leading constant of every explicit bound is still $\frac{3}{8}$.
- *An exact summatory bound* (Theorem 3.85), $\sum_{N \leq X} a(N) \geq 11X^3/72 + 5X^2/6 - O(X)$ for every X , with average constant $\frac{11}{24}$ (Corollary 3.86). The constant $\frac{11}{72}$ is all that the row budget of Theorem 3.44 can give (Proposition 3.87). This quantifies the last sentence of Question 6.

What is not new. The manuscripts re-derive several results printed above. Each is printed once and credited:

- manuscript 70’s Lemma 2.1 (Lemma 3.72) is Theorems 3.21 and 3.29, with the same proofs; its branch-cut formula (3.116) is Proposition 3.30, which it credits; its factorization and recurrence are Proposition 3.26, which it credits;

- its comparison Lemma 3.74 is Proposition 3.34 at $d = 5$, with the proofs of Lemmas 3.32 and 3.33;
- its two finite recurrences are (3.80) and (3.81), and its Lemma 3.73 is the criterion (3.88), which it credits;
- its Corollary 3.78 extends Corollary 3.45 by two depths, and manuscript 68’s Proposition 3.82 at the depth cap $\bar{d} = 4$ is that corollary;
- manuscript 68’s row budget is Theorem 3.44, which it credits.

A link the manuscript does not make. Manuscript 70 does not cite the second route to depth three (Subsections 3.27–3.34). Yet its tail-moment formula (3.120), taken at $d = 3$, is exactly the harmonic quantity E_q of (3.101), and it reproduces the cutoff $q \geq 38$ of that route and of Proposition 3.41. Its parameter integral in the proof of Lemma 3.75 is the beta-sine transform (3.94). Remark 3.76 records both.

Notation. The depth coordinates of Table 3 apply unchanged, with the batch-42 names for the objects batch 42 already named. Several letters of the manuscripts name *different numbers* from the same letters above, and are renamed by Table 3.A.

3.35.1 Notation: depths five and six, and count bounds

The manuscripts use the depth coordinates of Table 3: k is the exponent of $\log x$, d the depth, $M = N - k$, $q = M - k - 2d - 1$, and $H_{d,k,q}$ is the integer (3.49). The letters φ , γ , R_d , Π_d , h_d , ν_d , τ , s , y , $p = 2d$, $\delta = q - p$, S_δ , D_δ , $\mu_{d,r}$, $V_k(u)$, $J(d, k)$, $\chi_d(N)$ and $\beta(N)$ mean what they mean in Subsections 3.15–3.23. The practical cutoffs at a fixed depth are written (R, Q) , as in Proposition 3.41. Everything else that collides is renamed as in Table 3.A, with no change of normalization. The first three rows are where a reader will go wrong.

3.36 Depths five and six: the theorem and the known regions

Abstract of manuscript 70. *We complete logarithmic deficits five and six in the coefficient array for repeated derivatives of x^{x^2} . At either depth, every zero is forced by the known reflection symmetry. The inspected ProveIt report proves depths one through four and explicitly proposes Jensen criteria for the next two. We implement that proposal with certified moment cutoffs: $(R, Q) = (410, 839)$ at depth five and $(1571, 3048)$ at depth six. Two independent recurrences certify the resulting rectangles, containing 343151 and 4785360 cells. We also replace the report’s very large general fixed-depth bounds by explicit cutoffs of size $O(de^d)$, using a global affine minorant determined by the largest residual root and beta harmonic moments. This settles two specified source questions at the stated scope; the full OEIS term-count conjecture remains open in this report.*

Editorial note. “The inspected ProveIt report” is this report at commit [ca81647a9](#); “the two specified source questions” are Questions 1 and 2 of Subsection 3.24.

Proof status. This is an ordinary computer-assisted mathematical proof with explicit analytic reductions and exact finite certificates. It is not formally verified in Lean and has not been externally refereed. Novelty is assessed relative to the inspected repository source, not by an exhaustive historical-priority claim.

3.36.1 The precise theorem and the derivative interpretation

Put, as in (3.48),

$$\varphi(w) = (1 + w)^2 \log(1 + w), \quad \gamma(k, d, M) = [w^M](2 + w)^k \varphi(w)^d.$$

All indices are integers. The logarithm is expanded as a formal power series at zero.

Table 3.A: Letters of Subsections 3.35–3.41 against manuscripts 70 and 68 and against the rest of Part 3. No normalization was changed.

Here	Manuscript	Meaning	Tempting false reading
$\widehat{Q}_d, \widehat{K}_d$	Q_d, K_d (70, 68)	cutoffs $\lceil (2d+1)e^d \rceil$ and $2\widehat{Q}_d$ of Theorem 3.81	false: the constants Q_d, K_d of (3.74)
$\widehat{C}_d$	C_d (70)	$\prod_{\ell \geq 2} (\varrho_d - \omega_\ell)$	false: the constant C_d of (3.74)
ϱ_d, ω_ℓ	r_d, β_ℓ (70)	largest root and roots $\pi^2 \cot^2(\ell\pi/d)$ of Π_d	false: $r = M - k$; the bulk count $\beta(N)$
$\mathcal{Q}_k^{(r)}, \mathcal{A}_{M,k}$	$Q_k^{(r)}, A_{M,k}$	as in Subsection 3.15	$Q_{n,j}$ of Section 3.3; $A(t)$ of (3.10)
$\mathcal{B}_5(r), \mathcal{B}_6(r)$ $\mathcal{E}_{5,k,q}$	$B(r), B_6(R)$ $E_{k,q}$	base moments (3.72) expectation under the folded measure	$B(w)$ of (3.10) E_d of (3.74); E_q
$H_j, H_j^{(2)}$	$H_j, H_j^{(2)}$	harmonic numbers, as in Subsection 3.30	the integer $H_{d,k,q}$
$\Pi_5(\zeta), \Pi_6(\zeta)$ $b_5(t)$ $\Lambda_4(N)$ Ξ	$\Pi(z), \Pi_6(z)$ $b(t)$ $L_4(N)$ U	residual polynomials (3.66) $(ty)^{10}$, the base factor b_d at $d = 5$ the right side of (3.87) $1 - s^2$ in the proof of Theorem 3.81	$z = x^2$ — $\Lambda(n)$ of (3.3), which is $\beta(N)$ the bound $U(n)$
$\vartheta, c, \mathcal{G}$ ε ℓ	x, c, G (70) z (70, §1) j (70, product index)	$\pi/d, \cot \vartheta, \prod_{\ell \geq 2} (\zeta - \omega_\ell)$ a Taylor increment index of a linear factor	the variable x ; $c_N(j, k)$ $z = x^2$ the exponent j of x
$\bar{d}$ $\mathfrak{L}$ $\mathcal{N}(t)$ $\overline{\beta}(X)$ $\mathcal{S}(X)$ $\Sigma_a(X)$ $\ell_{d,k}$ $\iota(d, k)$ $i, l; \xi$	D (68) L (68) $F(t)$ (68) $B(X)$ (68) $S(X)$ (68) $A(X)$ (68) $L_{d,k}$ (68) $h(d, k)$ (68) $r, s; z$ (68)	depth cap $\log N$ $\sum_{i,l \geq 0} (t - 4i - 6l)_+$ $\sum_{N \leq X} \beta(N)$ guaranteed tail count $\sum_{N=1}^X a(N)$ $(X - 2k - 2d - 1)_+$ $\mathbf{1}\{k + d \text{ even}\}$ summation indices; generating variable	the difference $D_\delta(t)$ $L = \log x$; the old name of τ $\mathcal{F}_{d,k}$; the falling factorial F_n $B(w)$; $\mathcal{B}_d(r)$ $S_\delta(t)$ $A(t)$ of (3.10) — $h_d(\tau)$; $\chi_d(N)$ $r = M - k, s = 2t - 1; z = x^2$

Theorem 3.71 (Complete depths five and six classification). *For $d \in \{5, 6\}$, every $k \geq 0$, and every $M \geq d$,*

$$\gamma(k, d, M) = 0 \iff M = k + 4d + 1 \text{ and } k + d \text{ is even.}$$

For $0 \leq M < d$, the coefficient is zero for the separate structural reason that φ has order one at the origin.

To connect this statement with OEIS A290268, write

$$\frac{d^N}{dx^N} x^{x^2} = x^{x^2} \sum_{j \in \mathbb{Z}} \sum_{k \geq 0} c_N(j, k) x^j (\log x)^k.$$

Taylor expansion of $(x + \varepsilon)^{(x+\varepsilon)^2}$ gives the exact depth bridge

$$d = \frac{N + j}{2} - k, \quad M = N - k, \quad c_N(j, k) = \frac{N!}{k!d!} \gamma(k, d, M). \quad (3.112)$$

For example, substitute $w = \varepsilon/x$ into

$$x^{-x^2} (x + \varepsilon)^{(x+\varepsilon)^2} = \exp((2x\varepsilon + \varepsilon^2) \log x) \exp(x^2 \varphi(\varepsilon/x))$$

and extract logarithmic degree k , depth d , and ε^N . Thus the theorem completes both strata at every derivative order. At depth five the nonstructural zeros are $j = -11$, $N = 2k + 21$, k odd; at depth six they are $j = -13$, $N = 2k + 25$, k even.

Editorial note. (3.112) is Proposition 3.24, and the sketch above is its proof. In this part's letters the holes are: $m = 5$, j odd, $n = 2j + 21$; and $m = 6$, j even, $n = 2j + 25$ — in both cases $k = j + m$ even and $n = 4k - 2j + 1$, the reflection family of Theorem 3.11.

This report, Theorem 3.23 and its status summary, completes $1 \leq d \leq 4$. It also proves a terminating fixed-depth reduction with deliberately enormous bounds (Theorem 3.22), and a modular count certificate through $N = 3000$ (Proposition 3.18). Question 1 of its Section 3.24 (Subsection 3.24) already supplies the Jensen criteria used below and asks for certified practical cutoffs and the remaining finite tests. The new contributions are those cutoffs, their exact moment seeds, and the resulting all-order classifications. Depth five's rectangle lies within the earlier $N \leq 3000$ certificate; depth six's extends to $N = 6198$. The separate general cutoff answers its Question 2 with a nonoptimal singly-exponential bound. The live OEIS entry [2] still labels the full term-count formula conjectural.

3.36.2 Exact polynomial models and the known regions

Define, as in Subsection 3.15,

$$\mathcal{A}_{M,k}(s) = [w^M](2 + w)^k (1 + w)^s.$$

Then $\gamma(k, d, M) = \partial_s^d \mathcal{A}_{M,k}(2d)$, which is (3.58). For $r = M - k \geq 0$, the centered factorization from Subsection 3.15 is

$$M! \mathcal{A}_{M,k}(s) = s^r \mathcal{Q}_k^{(r)} \left(s - \frac{r-1}{2} \right), \quad (3.113)$$

where

$$\mathcal{Q}_0^{(r)} = 1, \quad \mathcal{Q}_1^{(r)} = 2y, \quad \mathcal{Q}_{k+1}^{(r)}(y) = 2y \mathcal{Q}_k^{(r)}(y) + k(k+r) \mathcal{Q}_{k-1}^{(r)}(y). \quad (3.114)$$

For completeness, its exponential generating function is

$$\sum_{k \geq 0} \frac{\mathcal{Q}_k^{(r)}(y)}{k!} \xi^k = (1 - \xi^2)^{-(r+1)/2} \exp(2y \operatorname{arctanh} \xi).$$

Expanding $(2+w)^k$ in $\mathcal{A}$, factoring out s^x and making the coefficient substitution $t = \xi/(1-\xi)$ gives this generating function and (3.113). Differentiating it proves (3.114). In particular $\mathcal{Q}_k^{(r)}$ has nonnegative coefficients and parity k .

Editorial note. (3.113), (3.114) and the generating function are (3.60), (3.61) and (3.59) (Proposition 3.26); the manuscript's sketch is the proof given there. The manuscript cites the classical Meixner–Pollaczek generating function [20, Eq. 18.23.7] for this family, as a specialization whose needed identities it proves directly; that is (3.62).

These identities give two previously established regions, recalled with their proofs to fix the boundary of the new work.

Lemma 3.72 (Bulk and symmetry). *For $d \geq 1$, one has $\gamma(k, d, M) > 0$ when $d \leq M \leq k + 2d + 1$. Also*

$$\gamma(k, d, k + 4d + 1) = 0 \quad \text{if } k + d \text{ is even.}$$

Proof. This is Theorem 3.21 together with Theorem 3.29. The manuscript recalls their proofs in condensed form — the centered factorization at $s = 2d + u$ with the Pascal induction (3.63) for $M < k$, and the odd falling factorial $u \prod_{\ell=1}^{2d} (u^2 - \ell^2)$ at $r = 4d + 1$ — and they are not printed a second time. $\square$

The remaining depth-five region is indexed by

$$q = M - k - 11 \geq 0, \quad H_{5,k,q} = \frac{(k + 11 + q)!}{5!} \gamma(k, 5, k + 11 + q).$$

The integer $H_{5,k,q}$ is the coefficient of u^5 in

$$V_k(u) = (10 + u)^{11+q} \mathcal{Q}_k^{(11+q)} \left(u + \frac{10 - q}{2} \right). \quad (3.115)$$

The known symmetry zeros are $q = 10$, k odd. The line $q = 0$ overlaps the positive bulk. (This is (3.79) at $d = 5$.)

3.37 Depths five and six: the depth-five moment certificates

3.37.1 A positive measure and a quadratic moment test

For $d \geq 1$, put $R_d(v) = \text{Im}(v + i\pi)^d / \pi$, as in (3.64). For $M = k + 2d + 1 + q$, $q \geq 0$, the branch-cut coefficient formula is

$$\gamma(k, d, M) = (-1)^{M+1} \int_0^1 t^q (1-t)^{2d} (2t-1)^k R_d \left(\log \frac{1-t}{t} \right) dt. \quad (3.116)$$

This is Proposition 3.30 of this report. To derive it, deform the Cauchy coefficient contour to the negative cut of $(1+w)^{2d} \log^d(1+w)$. Its jump is $2\pi i R_d(\log(x-1))$ at $w = -x$, $x > 1$. The substitution $t = 1/x$ gives (3.116). The small endpoint circle tends to zero as $\varepsilon^{2d+1} \log^d \varepsilon$, and the large circle tends to zero because $M > k + 2d$.

For the rest of the proof take $d = 5$, and on $1/2 < t < 1$ put

$$y = 1 - t, \quad s = 2t - 1, \quad \tau = \log(t/y), \quad b_5(t) = (ty)^{10}, \quad \delta = q - 10.$$

The residual polynomial is

$$\Pi_5(\zeta) = \zeta^2 - 2\pi^2 \zeta + \pi^4/5, \quad R_5(-\tau) = 5\Pi_5(\tau^2), \quad (3.117)$$

as in (3.68) and (3.67). Reflecting the lower half of (3.116), the integral becomes, up to a fixed nonzero signed factor,

$$\int_{1/2}^1 b_5(t) s^k \Pi_5(\tau^2) (t^\delta + (-1)^k y^\delta) dt.$$

For even k normalize the positive density $b_5 s^k S_\delta$, where $S_\delta = t^\delta + y^\delta$. For odd k and $q \neq 10$ normalize $b_5 s^k |D_\delta|$, where $D_\delta = t^\delta - y^\delta$. Call the resulting expectation $\mathcal{E}_{5,k,q}$. The omitted factor has a nonzero sign, so

$$\mathcal{E}_{5,k,q}[\Pi_5(\tau^2)] > 0 \implies H_{5,k,q} \neq 0 \quad (3.118)$$

outside the symmetry holes. (This is the folding (3.69) and the sign rule (3.70) at $d = 5$.)

Lemma 3.73 (Mean threshold nineteen). *If $Z \geq 0$ has a finite second moment and $\mathbb{E}Z \geq 19$, then $\mathbb{E}\Pi_5(Z) > 0$.*

Proof. The larger root of Π_5 is $\pi^2(1 + 2/\sqrt{5}) < 19$, using $\pi < 22/7$, hence $\pi^2 < 10$, and $2/\sqrt{5} < 9/10$. Since Π_5 is convex, $\mathbb{E}\Pi_5(Z) \geq \Pi_5(\mathbb{E}Z) > 0$. $\square$

This is the stronger quadratic criterion proposed in the source report ((3.88)), rather than a new use of Jensen's inequality. Unlike the earlier affine residuals, Π_5 is not increasing everywhere; the remaining task is to certify practical lower bounds on the mean of $Z = \tau^2$.

3.37.2 Two exact moment certificates cover the infinite tails

Let $\mu_{5,r}$ be the probability measure proportional to $b_5 s^r dt$ on $(1/2, 1)$, and write $\mathcal{B}_5(r) = \mathbb{E}_{\mu_{5,r}} \tau^2$, as in (3.72).

Lemma 3.74 (Likelihood ratio comparisons). *For nonhole integer parameters,*

$$\mathcal{E}_{5,k,q} \tau^2 \geq \begin{cases} \mathcal{B}_5(k), & k \text{ even}, \\ \mathcal{B}_5(k+1), & k \text{ odd}. \end{cases}$$

The function $\mathcal{B}_5(r)$ is nondecreasing. Within each parity class, increasing k by two increases every increasing moment of τ . Increasing q by one has the same property for $q \geq 11$.

Proof. This is Proposition 3.34 at $d = 5$. The manuscript repeats its proof: the covariance inequality of Lemma 3.32, the monotone ratios of Lemma 3.33 with the same formulas (3.71) and the same logarithmic derivative, the reflection $S_{-a} = (ty)^{-a} S_a$, $|D_{-a}| = (ty)^{-a} D_a$ for negative indices, and the factor s^2 for $k \mapsto k + 2$. It is not printed a second time. $\square$

Lemma 3.75 (The two finite moment checks). *One has $\mathcal{B}_5(410) > 19$ and $\mathcal{E}_{5,0,839} \tau^2 > 19$.*

Proof. For the first claim set $q = 10$ and define V_R by (3.115). For even R , the parameter form of the same branch-cut identity gives

$$V_R(u) = (R+21)! \frac{\sin \pi u}{\pi} \int_0^1 (t(1-t))^{10} (2t-1)^R e^{-u\tau} dt.$$

The integral is even in u by reflection. Comparing the coefficients of u and u^3 yields

$$\mathcal{B}_5(R) = \frac{\pi^2}{3} + 2 \frac{[u^3]V_R}{[u]V_R}. \quad (3.119)$$

The exact integer recurrence (3.114), truncated after degree three, verifies

$$[u]V_{410} > 0, \quad [u^3]V_{410} > 8[u]V_{410}.$$

Since $\pi > 3$, equation (3.119) gives $\mathcal{B}_5(410) > 19$.

For the second claim the reflected sum density is equivalent, for the symmetric function τ^2 , to a beta density proportional to $t^{839}(1-t)^{10}$ on $(0, 1)$. Write $H_j = \sum_{i=1}^j 1/i$ and $H_j^{(2)} = \sum_{i=1}^j 1/i^2$. Differentiating the beta integral gives

$$\mathcal{E}_{5,0,q}\tau^2 = (H_q - H_{10})^2 + \frac{\pi^2}{3} - H_q^{(2)} - H_{10}^{(2)}. \quad (3.120)$$

An exact rational summation verifies

$$(H_{839} - H_{10})^2 - H_{839}^{(2)} - H_{10}^{(2)} > 16.$$

Again $\pi > 3$ completes the claim. Both finite arithmetic checks are reproduced in the accompanying certificate; their values are approximately $[u^3]V_{410}/[u]V_{410} = 8.01181969$ and 16.00012307 , but these decimal approximations are not used to decide either sign. $\square$

Remark 3.76 (Editorial: the tail seed at depth three). The computation behind (3.120) works at every odd depth: there $k = 0$ is a sum case and $h_d = 1$, so the $k = 0$ measure is exactly the symmetrized beta law $(q + 1, 2d + 1)$, and $\mathcal{E}_{d,0,q}\tau^2 = (H_q - H_{2d})^2 + \pi^2/3 - H_q^{(2)} - H_{2d}^{(2)}$. At $d = 3$ the residual polynomial $\Pi_3(\zeta) = \zeta - \pi^2/3$ is affine, so

$$\mathcal{E}_{3,0,q}[\Pi_3(\tau^2)] = (H_q - H_6)^2 - H_q^{(2)} - H_6^{(2)} = 2\mathcal{E}_q,$$

the harmonic quantity (3.101) of the second route to depth three. Its sign law (Lemma 3.54: negative for $1 \leq q \leq 37$, positive from $q = 38$) is therefore the $d = 3$ instance of this tail seed, and the first-moment test of Proposition 3.80 at $d = 3$, whose threshold is $\varrho_3 = \pi^2/3$, gives exactly the cutoff $q \geq 38$ — the $Q = 38$ of Proposition 3.41. Likewise the parameter integral in the proof of Lemma 3.75 is the beta-sine transform (3.94) at $x = 11 + u$, $r = 21$ (the shift $x = s + 1$ of Table 5), that is, at the hole column $q = 2d$. The manuscript cites neither; the intake checked the sign change of $(H_q - H_6)^2 - H_q^{(2)} - H_6^{(2)}$ between $q = 37$ and $q = 38$ in exact arithmetic, and that 839 and 3048 are the least q for which the two depth-five and depth-six rational checks hold.

Proposition 3.77 (Practical depth five reduction). *Every nonhole entry with $k \geq 409$ or $q \geq 839$ is nonzero.*

Proof. If $k \geq 409$, even parity forces $k \geq 410$, while odd parity gives $k + 1 \geq 410$. Lemmas 3.74 and 3.75 imply $\mathcal{E}_{5,k,q}\tau^2 > 19$. For $q = 839$, the odd-parity minimum $k = 1$ also has a moment at least that of the sum measure with $k = 0$: its density ratio is

$$s \frac{D_{829}}{S_{829}} = s \tanh(829\tau/2),$$

a positive increasing function. Thus both parity minima exceed 19. Increase k by two and q by one using Lemma 3.74. Lemma 3.73 and (3.118) finish both cases. $\square$

3.37.3 The finite rectangle and completion of the proof

It remains to check

$$0 \leq k \leq 408, \quad 0 \leq q \leq 838.$$

There are $409 \cdot 839 = 343151$ cells. Their largest derivative order is $N = 2k + 11 + q = 1665$. The two exact computations are:

1. The integer jets (3.115) satisfy, in $\mathbb{Z}[u]/(u^6)$,

$$\begin{aligned} V_0 &= \prod_{\ell=0}^{10+q} (10 - \ell + u), \\ V_1 &= (2u + 10 - q)V_0, \\ V_{k+1} &= (2u + 10 - q)V_k + k(k + 11 + q)V_{k-1}. \end{aligned}$$

The coefficient $[u^5]V_k$ is computed as an exact integer and tested for zero.

2. Independently, multiplication by $(2 + w)$ gives the Pascal recurrence

$$H_{5,k+1,q} = 2H_{5,k,q+1} + (k + 12 + q)H_{5,k,q}.$$

Start with the independently computed $k = 0$ falling-factorial row through $q = 1246$, then shorten it at each step. Residues modulo 1009 and 1013 certify every nonhole except two; modulus 1019 certifies those remaining two. A nonzero residue proves integer nonvanishing. The symmetry zeros are proved analytically, rather than inferred from zero residues.

Both computations find exactly 204 zeros, namely $(k, q) = (1, 10), (3, 10), \dots, (407, 10)$, and no other zero. The integer computation reaches coefficients of at most 11042 bits. It uses only Python's standard library and runs in a few seconds in the research environment.

Editorial note. The two recurrences are (3.80) and (3.81) at $d = 5$. In the intake's two runs the integer computation took between 5 and 45 seconds. The depth-five rectangle lies inside the range $N \leq 3000$ of Proposition 3.18, so, as in Remark 3.42, its nonvanishing half re-confirms cells already certified; what that proposition does not supply are the moment seeds of Lemma 3.75, which make a finite rectangle decide every N .

The finite result, Proposition 3.77, the bulk positivity of Lemma 3.72, and the structural vanishing for $M < 5$ prove the depth-five case of Theorem 3.71. The certificates decide only the stated finite obligations; the analytic reduction is essential to the all-order conclusion.

Corollary 3.78 (A new contribution to the lower term count). *Let $a(N)$ count the nonzero derivative terms, and let $\Lambda_4(N)$ be the source report's lower bound obtained from the bulk and depths one through four, the right side of (3.87). The two new depth classifications give*

$$a(N) \geq \Lambda_4(N) + \sum_{d=5}^6 [\max(0, \lfloor N/2 \rfloor - d) - \mathbf{1}\{N \geq 4d + 1, N \equiv 2d + 1 \pmod{4}\}].$$

Proof. At depth d , outside the positive bulk, $N = 2k + 2d + 1 + q$ with $q \geq 1$ gives $\max(0, \lfloor N/2 \rfloor - d)$ cells. The sole possible hole has $q = 2d$ and $k + d$ even, exactly the displayed indicator. Depth six is proved in the next subsection. $\square$

The gain is linear in N . The leading lower-bound constant remains $3/8$, and the conjectured leading constant $1/2$ is not proved.

Editorial notes. (i) The indicator is $\chi_d(N)$ of (3.86), and the corollary is Proposition 3.82 with depth cap $\bar{d} = 6$, valid there for every N because depths five and six are now classified. For even $N \geq 12$ and odd $N \geq 25$ it adds $N - 11$, respectively $N - 13$, to the bounds of Remark 3.46. [Corrected 1 October 2026, batch 73: this note first read $N - 15$ at odd N ; at odd $N = 2\rho + 1 \geq 25$ exactly one of $\chi_5(N)$, $\chi_6(N)$ is 1, so the increment is $(\rho - 5) + (\rho - 6) - 1 = N - 13$, and the closed forms are $(3N^2 + 34N - 160)/8$ for even $N \geq 12$ and $(3N^2 + 36N - 215)/8$ for odd $N \geq 25$.] At $N = 12$ the one cell left unresolved in the example after Corollary 3.45 has depth five; it is now nonzero, and the bound equals $U(12) = 85$. (ii) “The leading lower-bound constant remains $3/8$ ” is true of every *explicit* lower bound, including Theorem 3.83 below. It is stale as a statement about the asymptotics: manuscript 34 of the same batch, placed in the new sibling report **power-tower-exponent-supports**, proves $a(N) = N^2/2 + o(N^2)$, without an effective threshold. That proof is not part of this report.

3.38 Depths five and six: the depth-six completion

For $d = 6$, the negative-region parameter is $q = M - k - 13 \geq 0$, and the residual polynomial is

$$\Pi_6(\zeta) = \zeta^2 - \frac{10}{3}\pi^2\zeta + \pi^4 = (\zeta - \pi^2/3)(\zeta - 3\pi^2).$$

Jensen’s inequality, as already proposed in the source ((3.88)), gives positive reduced expectation whenever $\mathbb{E}\tau^2 \geq 30$, since $3\pi^2 < 30$.

On the half interval the base density is now $(t(1 - t))^{12}\tau s^r$. Sum measures occur for odd k and difference measures for even k ; their symmetry holes are $q = 12$, k even. The same likelihood-ratio proof gives comparison with base exponent k in a sum case and $k + 1$ in a difference case.

Lemma 3.79 (Depth six moment seeds). *The base moment at $R = 1571$ exceeds 30, and at $Q = 3048$ both minimum parity measures have moment exceeding 30.*

Proof. At the center $q = 12$ define

$$V_R(u) = (12 + u)^{25} \mathcal{Q}_R^{(25)}(u).$$

For odd R , the parameter branch-cut integral has an odd integral factor and an overall minus sign. Expanding the integral and $\sin \pi u$ gives

$$\begin{aligned} [u^2]V_R &= (R + 25)! \int_0^1 \tau s^R (t(1 - t))^{12} dt, \\ [u^4]V_R &= \frac{(R + 25)!}{6} \int_0^1 (\tau^3 - \pi^2\tau) s^R (t(1 - t))^{12} dt. \end{aligned}$$

Therefore the τ -weighted base moment is

$$\mathcal{B}_6(R) = \pi^2 + 6 \frac{[u^4]V_R}{[u^2]V_R}.$$

Exact fourth-degree jets verify that $[u^2]V_{1571} > 0$ and $2[u^4]V_{1571} > 7[u^2]V_{1571}$. Since $\pi^2 > 9$, this proves $\mathcal{B}_6(1571) > 30$.

At $q = 3048$, the unweighted symmetrized beta measure with parameters $(q + 1, 13)$ has moment

$$(\mathbf{H}_q - \mathbf{H}_{12})^2 + \pi^2/3 - \mathbf{H}_q^{(2)} - \mathbf{H}_{12}^{(2)}.$$

Exact rational summation verifies that its rational part exceeds 27. Both minimum parity measures dominate this comparison measure: their density ratios are τs for the sum case $k = 1$, and $\tau \tanh((q - 12)\tau/2)$ for the difference case $k = 0$. These are increasing positive functions. This proves both claims. $\square$

It follows exactly as in Proposition 3.77 that all nonhole entries with $k \geq 1570$ or $q \geq 3048$ are nonzero. The remaining rectangle is

$$0 \leq k \leq 1569, \quad 0 \leq q \leq 3047.$$

It contains 4785360 cells, with maximum derivative order $N = 2k + 13 + q = 6198$. There are exactly 785 symmetry holes, namely $q = 12$ with even k .

Two independently implemented recurrences certify every nonhole by a nonzero residue modulo at least one of 1000003 and 1000033. The centered recurrence uses jets through degree six. The independent Pascal recurrence is

$$H_{6,k+1,q} = 2H_{6,k,q+1} + (k + 14 + q)H_{6,k,q},$$

initialized from the falling-factorial $k = 0$ row through $q = 4616$. Both computations find precisely the 785 prescribed holes and certify all 4784575 nonholes. Together with bulk positivity and the structural zeros this proves the depth-six case of Theorem 3.71. The exact moment seeds are also recomputed independently in the Pascal verifier.

Editorial notes. (i) The cells with $N \leq 3000$ of this rectangle are already nonzero by Proposition 3.18; the cells with $3000 < N \leq 6198$ are certified here for the first time. (ii) The program `code/08-depths-five-six-independent_pascal_depth6.py` says in its docstring that it uses the centered recurrence “only for its separate moment seed”; so the central seed is recomputed by the same recurrence as in the main certificate, and the independence of the two computations concerns the rectangle. Both depth-six computations work modulo the two primes; a nonzero residue proves nonvanishing.

3.39 Depths five and six: exponential cutoffs at every depth

3.39.1 A sharp mean criterion

The general residual polynomial (3.66) is

$$\Pi_d(\zeta) = \prod_{\ell=1}^{\nu_d} (\zeta - \omega_\ell), \quad \nu_d = \left\lfloor \frac{d-1}{2} \right\rfloor, \quad \omega_\ell = \pi^2 \cot^2 \frac{\ell\pi}{d}.$$

It is the monic factor in $R_d(-\tau) = (-1)^{d-1} d h_d(\tau) \Pi_d(\tau^2)$, where $h_d = 1$ for odd d and $h_d = \tau$ for even d ((3.67)). The root ordering is $\omega_1 > \cdots > \omega_{\nu_d} > 0$.

Proposition 3.80 (Global affine minorant at the largest root). *For $d \geq 3$, put $\varrho_d = \omega_1$ and $\widehat{C}_d = \prod_{\ell=2}^{\nu_d} (\varrho_d - \omega_\ell) > 0$, with empty product one. Then, for every $\zeta \geq 0$,*

$$\Pi_d(\zeta) \geq \widehat{C}_d(\zeta - \varrho_d). \quad (3.121)$$

Consequently, for any nonnegative random variable Z with the required finite moments, $\mathbb{E}Z > \varrho_d$ implies $\mathbb{E}\Pi_d(Z) > 0$.

Proof. For $d = 3, 4$ the polynomial is linear and the assertion is equality. For $d \geq 5$, let $\vartheta = \pi/d$ and $c = \cot \vartheta > 1$. The double-angle identity gives

$$\frac{\varrho_d}{\omega_2} = \frac{\cot^2 \vartheta}{\cot^2(2\vartheta)} = \frac{4c^4}{(c^2 - 1)^2} > 4.$$

Every other root is therefore less than $\varrho_d/4$. Put $\mathcal{G}(\zeta) = \prod_{\ell=2}^{\nu_d} (\zeta - \omega_\ell)$. For $0 \leq \zeta \leq \varrho_d$, each factor obeys $|\zeta - \omega_\ell| \leq \varrho_d - \omega_\ell$, so $\mathcal{G}(\zeta) \leq |\mathcal{G}(\zeta)| \leq \mathcal{G}(\varrho_d) = \widehat{C}_d$. Multiplication by the nonpositive number $\zeta - \varrho_d$ reverses that inequality. For $\zeta \geq \varrho_d$, every factor is positive and at least its value at ϱ_d , so $\mathcal{G}(\zeta) \geq \widehat{C}_d$. The two cases prove (3.121); taking expectations proves the final assertion. $\square$

The threshold is sharp as a uniform implication of the form $\mathbb{E}Z > c \Rightarrow \mathbb{E}\Pi_d(Z) > 0$ for arbitrary nonnegative distributions. If $c < \varrho_d$, a point mass at ϱ_d already contradicts strict positivity; a point mass sufficiently close to ϱ_d from below has negative polynomial value. This is not a claim that the coefficient distributions attain the worst case.

Editorial note. The proposition replaces the crude minorant $\Pi_d(\zeta) \geq \zeta/2 - C_d$ of Lemma 3.36. At $d = 5, 6$ its threshold ϱ_d is the largest root used in (3.88); at $d = 3, 4$ it is the root $\pi^2/3$, resp. π^2 , of the affine Π_d .

3.39.2 Singly exponential fixed-depth cutoffs

Theorem 3.81 (Singly exponential fixed depth cutoffs). *For every $d \geq 3$, take*

$$\widehat{Q}_d = \lceil (2d+1)e^d \rceil, \quad \widehat{K}_d = 2\widehat{Q}_d.$$

For $k, q \geq 0$ with $k \geq \widehat{K}_d$ or $q \geq \widehat{Q}_d$, the coefficient $\gamma(k, d, k+2d+1+q)$ is nonzero unless $q = 2d$ and $k+d$ is even. Thus the fixed-depth classification reduces to $0 \leq k < \widehat{K}_d$, $0 \leq q < \widehat{Q}_d$, a rectangle of $2\widehat{Q}_d^2 = O(d^2 e^{2d})$ cells.

The fully integer choices

$$\widehat{Q}_d = (2d+1)3^d, \quad \widehat{K}_d = 2(2d+1)3^d$$

also suffice, with exactly $2(2d+1)^2 9^d$ cells.

Proof. Put $p = 2d$ and write $Q = \widehat{Q}_d$. The general folded base density is $(t(1-t))^p h_d(\tau) s^r$, with $s = 2t - 1$. The increasing factor h_d can be omitted for a lower bound on the mean of τ^2 .

For $k \geq \widehat{K}_d$, the same base comparison as Lemma 3.74 reduces to a base exponent at least $\widehat{K}_d - 1 = 2Q - 1$. Decrease to that odd exponent and remove h_d . Under the resulting density, $\Xi = 1 - s^2$ has beta distribution with parameters $(p+1, Q)$. Since

$$\tau = -\log \Xi + 2 \log(1+s) \geq -\log \Xi,$$

Jensen's inequality and the beta logarithmic moment give

$$\mathbb{E}\tau^2 \geq (\mathbb{H}_{p+Q} - \mathbb{H}_p)^2 > \log^2 \frac{p+Q+1}{p+1} = \log^2 \left(1 + \frac{Q}{p+1}\right) > d^2.$$

Here $Q \geq (p+1)e^d$, and the strict harmonic inequality follows by integrating $1/x$ over $[p+1, p+Q+1]$.

For $q \geq Q > p$, compare the folded density with the unweighted symmetrized beta distribution $(q+1, p+1)$. The ratio is $h_d s^k$ in a sum case and $h_d s^k \tanh((q-p)\tau/2)$ in a difference case; each is increasing. The comparison moment equals

$$(\mathbb{H}_q - \mathbb{H}_p)^2 + \pi^2/3 - \mathbb{H}_q^{(2)} - \mathbb{H}_p^{(2)} > (\mathbb{H}_q - \mathbb{H}_p)^2 > \log^2 \frac{q+1}{p+1} > d^2.$$

The first strict inequality follows because both finite second-order harmonic sums are less than $\pi^2/6$.

Finally, $\varrho_d = \pi^2 \cot^2(\pi/d) < d^2$, since $\tan x > x$ for $0 < x < \pi/2$. Proposition 3.80 gives positive residual expectation, hence nonvanishing outside the symmetry holes. Since $e < 3$, the displayed integer choices also satisfy the hypotheses. The bound $e < 3$ follows directly from the exponential series and $n! \geq 2^{n-1}$ for $n \geq 2$, with strict inequality for some terms. $\square$

The rectangle size is a number of arithmetic obligations, not a bit-complexity estimate or a claim of optimality. This improves the source’s general cutoffs in response to Question 2 of its Section 3.24. The global affine minorant also extends its largest-root mean criterion to every depth, addressing the moment-bound direction of Question 3. The more economical cutoffs 410, 839 and 1571, 3048 give the explicit classifications at depths five and six.

Editorial note. For comparison: $\hat{Q}_d = 141, 492, 1633, 5245$ for $d = 3, \dots, 6$, against the practical tail cutoffs 38, 185, 839, 3048 of Proposition 3.41 and Lemmas 3.75–3.79; the constants (3.74) of Theorem 3.22 are of tower type. Theorem 3.81 also bears on the question “Critical-strip bounds” of the second route to depth three (Subsection 3.33): it gives explicit, singly exponential bounds on the region a finite certificate must cover at every depth, though not the conceptual control that question asks for.

The contribution is the certified completion of the source’s proposed depth-five/six program and the new general cutoff improvement. The source’s quadratic Jensen criteria and the classical centered-polynomial framework are credited rather than rediscovered. The full OEIS conjecture, the finite nonvanishing obligations at depths $d \geq 7$, and formal verification remain separate tasks. [Added 1 October 2026, batch 73: the obligations at depths seven and eight are discharged in Subsection 3.47ff.; those at $d \geq 9$ remain.] A bounded literature check found no further primary result resolving these exact strata; no exhaustive priority claim is made.

3.40 Pointwise and summatory count bounds (manuscript 68)

Abstract of manuscript 68. Let $a(N)$ count the terms in the fully expanded N th derivative of x^{x^2} . We derive two consequences of proved coefficient nonvanishing bounds. The exponential fixed-depth cutoff gives the pointwise refinement $a(N) \geq 3N^2/8 + (N/2) \log N - (N/2) \log \log N - 5N/4$ for every integer $N \geq e^{10}$. Separately, the existing parity-sensitive bound on zeros in a fixed row yields an exact summatory inequality and $\sum_{N=0}^X a(N) \geq 11X^3/72 + 5X^2/6 - O(X)$. Consequently the lower limit of the average of $a(N)/N^2$ is at least $11/24$. The latter is an average statement, not a pointwise bound of $11N^2/24$. Neither result proves the conjectured equivalent $a(N) \sim N^2/2$.

Status. The arguments are ordinary mathematical proofs, independently checked, and not formal Lean proofs or externally refereed results. The source-relative contributions are the growing-depth count and the summatory consequences, rather than the previously proved row zero bound. The accompanying exact arithmetic script checks the finite counting identities; no unexplored coefficient rectangle is assumed nonzero.

3.40.1 The coefficient facts used

Write

$$\frac{d^N}{dx^N} x^{x^2} = x^{x^2} \sum_{j \in \mathbb{Z}} \sum_{k \geq 0} c_N(j, k) x^j (\log x)^k, \quad a(N) = \#\{(j, k) : c_N(j, k) \neq 0\}.$$

The exact depth coordinates are

$$d = (N + j)/2 - k, \quad M = N - k, \quad \gamma(k, d, M) = [w^M](2 + w)^k ((1 + w)^2 \log(1 + w))^d.$$

Taylor expansion gives $c_N(j, k) = N! \gamma(k, d, M) / (k! d!)$. The source report proves a positive bulk of size

$$\beta(N) = \begin{cases} (3N^2 + 10N + 8)/8, & N \text{ even}, \\ (3N^2 + 12N + 1)/8, & N \text{ odd}. \end{cases} \quad (3.122)$$

Every candidate cell outside this bulk is uniquely described by

$$d \geq 1, \quad k \geq 0, \quad q \geq 1, \quad N = 2k + 2d + 1 + q. \quad (3.123)$$

Its coefficient is a positive scalar times $H_{d,k,q} = (k + 2d + 1 + q)! \gamma(k, d, k + 2d + 1 + q) / d!$. The mandatory symmetry zeros are precisely $q = 2d$ with $k + d$ even. Other zeros have not been ruled out at every depth.

Editorial note. The bridge is Proposition 3.24; (3.122) is (3.85), the lower bound $\Lambda(n)$ of Theorem 3.1.

Two proved inputs will be used in distinct ways.

1. The companion cutoff theorem (Theorem 3.81, manuscript 70's Theorem 7.2) shows that, for $d \geq 3$, every nonhole cell is nonzero when $k \geq 2\hat{Q}_d$ or $q \geq \hat{Q}_d$, where

$$\hat{Q}_d = \lceil (2d + 1)e^d \rceil. \quad (3.124)$$

Depths one and two are already classified for all parameters in the source; the companion extends the all-order classification through depth six. [Added 1 October 2026, batch 73: through depth eight since Theorems 3.99 and 3.100.]

2. The source's Mellin zero-count theorem bounds the number of zeros over all real $q > -1$ in a fixed (d, k) row by

$$J(d, k) = \begin{cases} d - 1, & k \text{ even}, \\ d, & k \text{ odd and } d \text{ odd}, \\ d - 2, & k \text{ odd and } d \text{ even}. \end{cases} \quad (3.125)$$

In particular it bounds the number of integer tail cancellations, including the symmetry hole when present.

The first input follows from a largest-root affine minorant for the folded residual polynomial and beta logarithmic moments. The second follows by counting sign changes of the integral kernel and applying the Chebyshev property of distinct real powers. They are separate statements: a per-row zero budget cannot be read as a pointwise classification along a derivative-order diagonal.

Editorial notes. (i) “Depths one and two are already classified ... in the source” understates this report: depths one to four are classified here (Theorem 3.23), and manuscript 70 adds five and six. The argument below uses the classified depths only through Proposition 3.82, where any classified depth may be counted. (ii) (3.125) is (3.83) of Theorem 3.44, quoted exactly.

3.40.2 A growing-depth pointwise bound

At a fixed derivative order, the general cutoff proves more than any fixed finite list of classified depths.

Proposition 3.82 (Exact depth count). *Let $\bar{d} \geq 3$ be an integer. If*

$$N \geq 5\hat{Q}_{\bar{d}} + 2\bar{d} - 1, \quad (3.126)$$

then

$$a(N) \geq \beta(N) + \sum_{d=1}^{\bar{d}} [\max(0, \lfloor N/2 \rfloor - d) - \chi_d(N)], \quad (3.127)$$

where

$$\chi_d(N) = \mathbf{1}\{N \geq 4d + 1, N \equiv 2d + 1 \pmod{4}\}.$$

Proof. Any cell inside the unverified rectangle has $k \leq 2\widehat{Q}_d - 1$ and $q \leq \widehat{Q}_d - 1$, and therefore, by (3.123),

$$N \leq 2(2\widehat{Q}_d - 1) + 2d + 1 + (\widehat{Q}_d - 1) = 5\widehat{Q}_d + 2d - 2.$$

The right side increases with d . Under (3.126), every cell of every depth $3 \leq d \leq \bar{d}$ lies outside its rectangle and is nonzero except for its possible symmetry hole. Depths one and two are covered by their existing classifications.

For a fixed depth, equation (3.123) gives exactly $\max(0, \lfloor N/2 \rfloor - d)$ possible k values. The unique possible hole has $N = 2k + 4d + 1$ and $k + d$ even, equivalent to the displayed indicator. The depths are disjoint and are outside the counted bulk. This proves (3.127). $\square$

Editorial note. $\chi_d(N)$ is (3.86). With $\bar{d} = 4$ and no threshold, (3.127) is Corollary 3.45; with $\bar{d} = 6$ and no threshold it is Corollary 3.78. Since depths $d \leq 6$ are classified, the threshold (3.126) is needed only for $\bar{d} \geq 7$. [Added 1 October 2026, batch 73: with $\bar{d} = 8$ and no threshold it is Corollary 3.106, and the threshold is needed only for $\bar{d} \geq 9$.]

Theorem 3.83 (A pointwise logarithmic gain). *For every integer $N \geq e^{10}$,*

$$a(N) \geq \frac{3N^2}{8} + \frac{N}{2} \log N - \frac{N}{2} \log \log N - \frac{5N}{4}. \quad (3.128)$$

In particular, the improvement over the source's quadratic bulk is of order $N \log N$, while its leading pointwise constant remains $3/8$.

Proof. Put $\mathfrak{L} = \log N$ and $\bar{d} = \lfloor \mathfrak{L} - \log \mathfrak{L} - 3 \rfloor$. Then $\bar{d} \geq 4$, $\bar{d} \leq \mathfrak{L}$, and $e^{\bar{d}} \leq N/(e^3 \mathfrak{L})$. Since $e^3 > 20$ and $\mathfrak{L} \geq 10$,

$$\widehat{Q}_{\bar{d}} \leq (2\bar{d} + 1)e^{\bar{d}} + 1 < \frac{21N}{200} + 1.$$

It follows that

$$5\widehat{Q}_{\bar{d}} + 2\bar{d} - 1 < \frac{21N}{40} + 2\mathfrak{L} + 4 < N.$$

The last inequality follows, for example, by differentiating $19e^{\mathfrak{L}}/40 - 2\mathfrak{L} - 4$ on $\mathfrak{L} \geq 10$. Proposition 3.82 applies. Using $\chi_d \leq 1$ and $\lfloor N/2 \rfloor \geq N/2 - 1/2$ gives

$$a(N) \geq \beta(N) + \frac{N\bar{d}}{2} - \frac{\bar{d}^2}{2} - 2\bar{d}.$$

Now $\bar{d} \geq \mathfrak{L} - \log \mathfrak{L} - 4$ and $\mathfrak{L}^2 + 4\mathfrak{L} \leq e^{\mathfrak{L}} = N$ for $\mathfrak{L} \geq 10$, so

$$a(N) \geq \beta(N) + \frac{N}{2}(\mathfrak{L} - \log \mathfrak{L}) - \frac{5N}{2}.$$

Finally (3.122) gives $\beta(N) \geq 3N^2/8 + 5N/4$, proving (3.128). $\square$

This proof does not declare any previously unchecked rectangle nonzero. Instead, at the particular order N , a growing number of complete depth diagonals avoid those rectangles altogether.

Remark 3.84 (Editorial: where the bound comes from, and its threshold). The gain of order $N \log N$ genuinely needs Theorem 3.81: with the tower-type constants (3.74) of Theorem 3.22 the admissible depth cap $\bar{d}$ grows far more slowly than $\log N$. For moderate N less is needed. An exact count at the intake (a scratch computation, not shipped) found that the right side of (3.87) (depths up to four) is at least the bound (3.128) for every $3 \leq N \leq 92653$, and the right side of Corollary 3.78 (depths up to six) for every $3 \leq N \leq 814895$; and that the right side of (3.127), with the largest $\bar{d}$ admitted by (3.126) or $\bar{d} = 6$, is at least (3.128) for every $3 \leq N \leq 3 \cdot 10^6$. The threshold $e^{10} \approx 22026$ of the theorem is therefore conservative; the theorem is stated here with it, as the manuscript proves it. (For $N \leq 3000$, $a(N) = U(N)$ is known in any case, by Proposition 3.18.) The manuscript's own choice of $\bar{d}$ becomes admissible, $\bar{d} \geq 3$ with (3.126), from $N = 3265$, but its estimates use $\mathfrak{L} \geq 10$.

3.40.3 An exact summatory bound from the row budgets

Let $x_+ = \max(x, 0)$ and define the finite sum

$$\mathcal{N}(t) = \sum_{i,l \geq 0} (t - 4i - 6l)_+. \quad (3.129)$$

For integer $t \leq 0$ it is zero. Put $\bar{\beta}(X) = \sum_{N=0}^X \beta(N)$.

Theorem 3.85 (Exact summatory inequality). *For every integer $X \geq 0$,*

$$\sum_{N=0}^X a(N) \geq \bar{\beta}(X) + \mathcal{N}(X-3) + 2\mathcal{N}(X-6) + \mathcal{N}(X-7). \quad (3.130)$$

Consequently

$$\sum_{N=0}^X a(N) \geq \frac{11X^3}{72} + \frac{5X^2}{6} - O(X). \quad (3.131)$$

Proof. For fixed d, k , the orders $N \leq X$ offer $\ell_{d,k} = (X - 2k - 2d - 1)_+$ integer tail values $q \geq 1$. The row budget (3.125) proves at least $(\ell_{d,k} - J(d, k))_+$ nonzero coefficients. Thus the guaranteed tail contribution is

$$\mathcal{S}(X) = \sum_{d \geq 1, k \geq 0} (X - 2k - 2d - 1 - J(d, k))_+.$$

Split k into $2i, 2i+1$ and d into $2l+1, 2l+2$. The four summands become, in order,

$$(X - 3 - 4i - 6l)_+, \quad (X - 6 - 4i - 6l)_+, \quad (X - 6 - 4i - 6l)_+, \quad (X - 7 - 4i - 6l)_+.$$

This is exactly the right-side tail in (3.130). No extra hole subtraction is appropriate: $J(d, k)$ already bounds all zeros in the row.

The generating function of this finite count is

$$\sum_{X \geq 0} \mathcal{S}(X) \xi^X = \frac{\xi^4 + 2\xi^7 + \xi^8}{(1 - \xi)^2(1 - \xi^4)(1 - \xi^6)}. \quad (3.132)$$

Its expansion at $\xi = 1$ begins

$$\frac{1}{6(1 - \xi)^4} - \frac{5}{12(1 - \xi)^3} + O((1 - \xi)^{-2}),$$

while all other poles have order at most two. Partial fractions therefore give $\mathcal{S}(X) = X^3/36 - X^2/24 + O(X)$. Direct summation of (3.122) gives

$$\bar{\beta}(X) = \begin{cases} X^3/8 + 7X^2/8 + 5X/4 + 1, & X \text{ even,} \\ X^3/8 + 7X^2/8 + 11X/8 + 5/8, & X \text{ odd.} \end{cases}$$

Adding these expressions proves (3.131). $\square$

The row-length cap is important. Subtracting the full budget d from every short terminal row would lose valid information. An equivalent coarser calculation caps possible cancellations by

$$\sum_{d,k} \min\{d, (X - 2k - 2d - 1)_+\} = \sum_{j=1}^{\lfloor (X-1)/3 \rfloor} \binom{\lfloor (X-1-3j)/2 \rfloor + 2}{2} = \frac{X^3}{72} + O(X).$$

Subtracting this from the full candidate total $X^3/6 + O(X^2)$ gives the same cubic constant $11/72$; keeping the parity-sensitive budget also gives the sharper quadratic term in (3.131).

Corollary 3.86 (Normalized average and upper limit). *One has*

$$\liminf_{X \rightarrow \infty} \frac{1}{X} \sum_{N=1}^X \frac{a(N)}{N^2} \geq \frac{11}{24}, \quad \limsup_{N \rightarrow \infty} \frac{a(N)}{N^2} \geq \frac{11}{24}.$$

Proof. Write $\Sigma_a(X) = \sum_{N=1}^X a(N)$. Summation by parts gives

$$\sum_{N=1}^X \frac{a(N)}{N^2} = \frac{\Sigma_a(X)}{X^2} + \sum_{n=1}^{X-1} \Sigma_a(n) \left(\frac{1}{n^2} - \frac{1}{(n+1)^2} \right).$$

All weights are positive. Theorem 3.85 implies $\Sigma_a(n) \geq (11/72)n^3 - O(n^2)$. Substituting this bound, and using $n^3(1/n^2 - 1/(n+1)^2) = 2 + O(1/n)$, yields

$$\sum_{N=1}^X \frac{a(N)}{N^2} \geq \frac{11X}{24} - O(\log X).$$

The first conclusion follows; the second is an immediate consequence of an average lower bound. $\square$

In particular, (3.131) is not being interpreted as a pointwise inequality with leading constant $11/24$. The conjectured leading constant $1/2$ is still not established. The source's upper bound continues to imply $\limsup a(N)/N^2 \leq 1/2$.

Editorial note. The upper bound is Theorem 3.14. The last sentence of Question 6 of Subsection 3.24 says that the fixed-row bound alone does not bound the total number of extra zeros after summing over the triangle; Theorem 3.85 computes what it does give, a summatory constant $\frac{11}{72}$ against the conjectured $\frac{1}{6}$, and Proposition 3.87 shows that this is the limit of that information. As for Corollary 3.86: an ineffective proof of $a(N) \sim N^2/2$, which would imply it, is in the sibling report named in Corollary 3.78, note (ii); it is not part of this report.

3.40.4 Scope of sharpness from integer information

The cubic constant above is optimal for a deliberately narrow collection of integer support constraints. This observation is not a sharpness claim about the actual derivative coefficients or their analytic identities.

Proposition 3.87 (An abstract support pattern). *If one retains only bulk positivity, the integer row budgets (3.125), the prescribed symmetry holes, the completed depths through six, and the cutoff exclusions (3.124), those facts alone cannot improve the summatory leading constant $11/72$.*

Proof. Retain every bulk cell and remove every prescribed symmetry hole. Put $\iota(d, k) = \mathbf{1}\{k + d \text{ is even}\}$. For $d \geq 7$ and $k < 2\widehat{Q}_d$, additionally remove the cells

$$q = 1, \dots, J(d, k) - \iota(d, k),$$

and retain all other cells. The added zeros have $q \leq d < \widehat{Q}_d$, and are disjoint from the symmetry hole $q = 2d$. Their total in each row is at most $J(d, k)$. Thus this abstract pattern respects exactly the stated constraints.

For comparison, the baseline pattern removing $1 \leq q \leq d$ has, at order N , the number of lattice points with $d \geq 1, k \geq 0$ satisfying $2k + 2d \leq N - 2$, minus the number satisfying $2k + 3d \leq N - 2$. These counts are respectively $N^2/8 + O(N)$ and $N^2/12 + O(N)$, leaving $N^2/24 + O(N)$ removed cells.

Replacing d by $J - \iota$ changes only a bounded number of possible q values per depth, hence $O(N)$ cells at an order. The mandatory holes also contribute $O(N)$. Restoring the finitely many completed depths changes $O(1)$ cells. Finally, restoring cells with $k \geq 2\widehat{Q}_d$ affects only $d = O(\log N)$, and hence $O((\log N)^2)$ cells. Since the nonstructural candidate count is $N^2/2 + O(N)$, the resulting abstract support has

$$a_{\text{model}}(N) = \frac{11N^2}{24} + O(N), \quad \sum_{N \leq X} a_{\text{model}}(N) = \frac{11X^3}{72} + O(X^2).$$

This proves the information-boundary claim. The pattern is not asserted to arise from any function or to obey the actual coefficient recurrence or Mellin identities. $\square$

[Added 1 October 2026, batch 73: depths seven and eight are now completed too (Theorems 3.99 and 3.100). Retaining them as well, and removing the extra cells only for $d \geq 9$, the same proof applies (restoring two more depths again changes $O(1)$ cells at an order), so the constant $11/72$ is still all that this information gives. This is an observation of the intake.]

Editorial note. This subsection is the manuscript's file `sharpness.tex`, which its main source loads with `\IfFileExists` as a section of its own; the proposition is unnumbered there and the label is the intake's.

3.40.5 Verification and relation to the source

The counting argument uses no large new coefficient computation. The supplied script checks direct row-budget sums, the four parity classes, the exact prefix bulk count, and the quasipolynomial generated by (3.132) for $X \leq 180$, and verifies its rational generating-function identity symbolically. These checks complement the explicit bijective/parity proofs; no finite test is extrapolated to all N .

The inspected source (this report at commit `ca81647a9`) states the fixed-row zero budget and pointwise quadratic lower bounds, but not these growing-depth or summatory consequences. The pointwise result uses the companion's new exponential cutoffs; the summatory result follows already from the source's row theorem by retaining the finite row-length restriction. No claim is made to exhaustive literature priority.

Editorial note. The growing-depth idea is Corollary 3.45 with a fixed cap $\bar{d} = 4$; what is new is letting $\bar{d}$ grow with N , which Theorem 3.81 makes possible.

3.41 Depths five and six and count bounds: provenance, files and formal status

3.41.1 Provenance

Subsections 3.35–3.41 were built from **two** manuscripts of batch 72A, which arrived in commit 1512ef835 and were placed in this report by commit d292c6765, as its additions 08 and 07. Both are dated 1 October 2026, carry the line “Research note prepared for Vladimir Reshetnikov with OpenAI”, and are pinned to ProveIt commit ca81647a9f679e96d49ffdbc54b0c0ab13d10a36.

- Manuscript 70 (addition 08), *Depths Five and Six Nonvanishing for OEIS A290268*, 9-page PDF, archive ProveIt_A290268_Depths_Five_Six.zip. Its Sections 1–2 form Subsection 3.36, 3–5 Subsection 3.37, 6 Subsection 3.38, 7 Subsection 3.39. It contributes Theorem 3.71, the moment seeds, the depth-five reduction, Corollary 3.78, Proposition 3.80 and Theorem 3.81.
- Manuscript 68 (addition 07), *Pointwise and Summatory Bounds for OEIS A290268*, 6-page PDF, archive ProveIt_A290268_Count_Bounds.zip, with its input file `sharpness.tex`. Its four sections and that input form Subsection 3.40. It contributes Proposition 3.82, Theorems 3.83 and 3.85, Corollary 3.86 and Proposition 3.87.

Every theorem, proposition, lemma and corollary of the two manuscripts is printed, with every non-claim and limitation they state; none was weakened or strengthened. The editorial choices were these.

- The additions are placed as subsections at the end of Part 3, after the batch-70 addition, so that no existing section, theorem or equation number moves; 70 comes before 68, which uses it.
- Manuscript 70’s Lemma 2.1 and Lemma 4.1 are restatements of results of this part (Theorems 3.21, 3.29 and Proposition 3.34) with the same proofs; their statements are printed and their proofs are cited, not reprinted. Its other restated formulas (bridge, factorization, branch cut, residual polynomials, folding, recurrences) are printed where the manuscript prints them, each with a reference to the result it restates.
- The letters are changed as in Table 3.A, with no change of normalization; the binding change is that manuscript 70’s C_d , K_d , Q_d and manuscript 68’s K_d , Q_d are printed $\widehat{C}_d$, $\widehat{K}_d$, $\widehat{Q}_d$, because the same letters already name the different constants (3.74). The manuscripts’ citations of the pinned source are re-targeted to this part, manuscript 68’s citation of “the companion” (its bibliography names the archive) to Theorem 3.81, and both OEIS entries and 70’s DLMF entry to this article’s bibliography.
- Manuscript 68’s `sharpness.tex` is printed as the subsubsection “Scope of sharpness from integer information”.
- Remarks 3.76 and 3.84, the notation subsection and table, and the editorial notes are additions of the intake, not of the manuscripts.

Not shipped: the two manuscripts’ sources and PDFs, 68’s `sharpness.tex`, their `README.md` files, and the two copies of the generic TeX wrapper `build_local.sh`. The archives survive in the history of commit 1512ef835. The READMEs add no mathematical claim; both say that the proofs had an “independent” mathematical check or review, which this report cannot confirm.

3.41.2 Files

The manuscripts' code and data are shipped byte-identical to the delivery, with the prefixes 08-depths-five-six- (70) and 07-count-bounds- (68).

Shipped path (delivery name)	Purpose
code/08-depths-five-six-verify_depth5.py (verify_depth5.py)	Depth-five rectangle and seeds by the centered recurrence over exact integers; writes <code>depth5_certificate.json</code> beside itself.
data/08-depths-five-six-depth5_certificate.json (depth5_certificate.json)	Its recorded output, with a digest, the exact seed ratio and a wall-clock field <code>elapsed_seconds</code> .
code/08-depths-five-six-independent_pascal_depth5.py (independent_pascal_depth5.py)	Depth-five rectangle by the Pascal recurrence modulo 1009, 1013, 1019, and both seeds; writes <code>independent_pascal_depth5.json</code> beside itself.
data/08-depths-five-six-independent_pascal_depth5.json (independent_pascal_depth5.json)	Its recorded output.
code/08-depths-five-six-verify_depth6_columns.cpp (verify_depth6_columns.cpp)	Depth-six rectangle by the centered recurrence modulo 1000003 and 1000033 (C++17); prints its JSON to standard output.
data/08-depths-five-six-depth6_column_certificate.json (depth6_column_certificate.json)	Its recorded output.
code/08-depths-five-six-independent_pascal_depth6.py (independent_pascal_depth6.py)	Depth-six rectangle by the Pascal recurrence modulo the same two primes, and both seeds; writes <code>independent_pascal_depth6.json</code> beside itself.
data/08-depths-five-six-independent_pascal_depth6.json (independent_pascal_depth6.json)	Its recorded output.
code/07-count-bounds-verify_counts.py (verify_counts.py)	Exact count checks of Subsection 3.40 for $X \leq 180$ and the generating-function identity (SymPy); prints three PASS lines and writes nothing.
data/07-count-bounds-verification.txt (verification.txt)	Its recorded output.
data/07-count-bounds-requirements.txt (requirements.txt)	Its dependency, <code>sympy==1.14.0</code> .

The commands are in the README. The JSON files and programs keep the manuscripts' letters (R, Q for the practical cutoffs, D for the depth).

3.41.3 Formal status

Nothing in this addition is formalized, and its placement beside a Lean development confers no formal status. The Lean library A290268 (`Combinatorics/DerivativeExpansions/`

A290268/Lean) relates to it as described in Section 3.26.3. Neither manuscript cites or relies on it. Theorem 3.71 would discharge the nonvanishing inclusion of that library’s support characterization on cells of depth five and six, on paper only; the theorems of A290268.Main stay conditional, and the library defines no γ .

3.42 Depths seven and eight

What this addition is. Subsections 3.42–3.51 were written into this part on 1 October 2026 from two manuscripts of batch 73, both dated 1 October 2026:

- manuscript 37, *An Eight-Depth Nonvanishing Theorem for OEIS A290268: Dominant-root tangents, single-exponential cutoffs, and exact certificates* (source 09), pinned to this report’s text at commit f18afc619, before the batch-72A Subsections 3.35–3.41 were written;
- manuscript 46, *Depth Seven Is Complete for OEIS A290268: A largest-root affine minorant, exact moment cutoffs, and a 48-million-cell certificate* (source 10), pinned to commit d3c317c09, after them, and written without knowledge of manuscript 37.

The two are not versions of each other: they share no paragraph, file or notation, and they prove the depth-seven classification by two different certificates. Provenance is in Subsection 3.51. They continue the depth coordinates of Subsections 3.14–3.26. **The full A290268 conjecture is still not proved**; the open region of Conjecture 3.19 is now deficit $m \geq 9$, $n \geq 2k + 2$.

What is new relative to this report.

- *Depth seven is classified* (Theorem 3.99), with two independent proofs: manuscript 37’s three-anchor reduction with a rectangle of 51,699,091 cells, and manuscript 46’s largest-root cutoffs with a rectangle of 48,113,025 cells replayed cell by cell by a second recurrence.
- *Depth eight is classified* (Theorem 3.100, manuscript 37 alone), with a rectangle of 552,465,284 cells. At both depths the only zeros are the symmetry holes $M = k + 4d + 1$, $k + d$ even.
- *A practical reduction at every depth* from three exact moment anchors (Theorem 3.97) and exact formulas that read every second moment off four jet coefficients (Proposition 3.98); they generalize the depth-specific seed lemmas 3.75 and 3.79.
- *An eight-depth count bound* (Corollary 3.106) and the exact depth-seven count with its rational generating function (Corollary 3.107).

What is not new. Manuscript 37 was written against the report before Theorem 3.71 and Theorem 3.81 were in it, and re-proves four results that the batch-72A addition had put here a few hours earlier. Each is printed once:

- its rows $d = 5, 6$ re-prove **Theorem 3.71** with smaller rectangles (323,158 and 4,298,240 cells against 343,151 and 4,785,360); they are printed as a *second certificate* of that theorem (Remark 3.102), not as a new classification; its rows $d = 3, 4$ are recertifications of Theorem 3.23;
- its dominant-root tangent is **Proposition 3.80**, with the same proof, and is printed by reference (Subsection 3.44); manuscript 46’s largest-root minorant is the case $d = 7$ of the same proposition, and its factored proof is printed as a second proof at $d = 7$;
- its single-exponential cutoffs $\check{K}_d, \check{Q}_d = \Theta(d^2 e^d)$ are **weaker than Theorem 3.81** ($\hat{Q}_d = \lceil (2d + 1)e^d \rceil = \Theta(de^d)$; at $d = 7$, $\check{Q}_7 = 1,927,726$ against $\hat{Q}_7 = 16,450$), and its $N \log N$ lower bound, with $-N \log \log N - C_* N$ and an ineffective constant, is **weaker than Theorem 3.83** ($-\frac{N}{2} \log \log N - \frac{5N}{4}$ for every $N \geq e^{10}$); both are kept only as remarks (Remarks 3.93 and 3.108), with their proofs, which are different from the proofs of the stronger results;

- its depth-count proposition is **Proposition 3.82** and is printed by reference; its model, factorization, bulk, holes, branch cut, folding and comparison lemmas are results of Subsections 3.15–3.18, which it credits, and are printed by reference in Subsection 3.43;
- manuscript 46 calls its largest-root minorant “new” five times, although Proposition 3.80 was in the report at its own pin; each such sentence carries an editorial note.

Notation. The depth coordinates of Table 3 and the renamings of Table 3.A apply. The manuscripts’ own letters that collide with letters already in use are renamed by Table 3.B; most important, manuscript 37’s constants K_d, Q_d are printed $\check{K}_d, \check{Q}_d$, and are neither the constants K_d, Q_d of (3.74) nor the cutoffs $\hat{K}_d, \hat{Q}_d$ of Theorem 3.81.

3.42.1 Notation: depths seven and eight

Both manuscripts use the depth coordinates of Table 3: k is the exponent of $\log x$, d the depth, $M = N - k$, $q = M - k - 2d - 1$, $p = 2d$, $r = M - k$, and $H_{d,k,q}$ is the integer (3.49). The letters $\varphi, \gamma, \mathcal{A}_{M,k}, \mathcal{Q}_k^{(r)}, R_d, \Pi_d, h_d, \nu_d, b_d(t), y, s, \tau, \delta = q - p, S_\delta, D_\delta, \mu_{d,r}, \mathcal{B}_d, \mathcal{E}_{d,k,q}, k_s, k_a, \chi_d(N)$ and $\beta(N)$ mean what they mean in Subsections 3.15–3.23, and $\varrho_d, \omega_\ell, \hat{C}_d, \hat{K}_d, \hat{Q}_d, \vartheta, H_j, \bar{d}$ what they mean in Subsections 3.35–3.41. The practical cutoffs at a fixed depth are written (R, Q) , as in Proposition 3.41; in both manuscripts, as there, the remaining rectangle is $0 \leq k \leq R - 2, 0 \leq q \leq Q - 1$. Everything else that collides is renamed as in Table 3.B, with no change of normalization. The first four rows are where a reader will go wrong.

Manuscript 46’s count $a_7(N)$ of depth-seven terms keeps its name; it is *not* the term count $a(N)$ at $N = 7$.

Abstract of manuscript 37. *OEIS A290268 counts the distinct terms in the fully collected derivatives of x^{x^2} . Its conjectured quadratic quasipolynomial is equivalent to excluding unexplained integer zeros from a three-parameter coefficient array. The maintained ProveIt report, at the repository snapshot inspected here, completes logarithmic deficits one through four and leaves deficits five and above open.*

We prove a global tangent-minorant inequality for the residual polynomial

$$\Pi_d(\zeta) = \prod_{\ell=1}^{\lfloor (d-1)/2 \rfloor} \left(\zeta - \pi^2 \cot^2 \frac{\ell\pi}{d} \right).$$

Its largest root dominates all the others sufficiently strongly that the expectation of this possibly nonmonotone polynomial is positive whenever the underlying second logarithmic moment exceeds that root. This replaces the linear-kernel restriction of the previous practical certificates. It yields explicit integer cutoffs of size $O(d^2 e^d)$, a deterministic $2^{O(d)}$ bit-operation algorithm enumerating every exceptional zero at a fixed depth, and an unconditional lower bound

$$a(N) \geq \frac{3}{8}N^2 + \frac{1}{2}N \log N - N \log \log N - C_* N \quad (N \geq N_*)$$

for absolute constants C_, N_* .*

Eighteen exact rational moment anchors and modular checks then complete every derivative order at depths five, six, seven and eight. The new finite rectangles contain 608,785,773 cells; all non-symmetry entries have nonzero residue modulo 1,000,000,007. We provide the proofs, integer recurrences, source code, recorded results, and independent finite checks. The full OEIS conjecture, and its proposed leading constant $1/2$, remain open. These are unrefereed, computer-assisted research results, not a Lean or Rocq formalization and not a claim of exhaustive historical priority.

Proof boundary (the manuscript’s box): analytic arguments cover the infinite ranges; exact programs cover explicitly bounded finite sets. Neither component alone is advertised as a proof of the complete OEIS conjecture.

Table 3.B: Letters of Subsections 3.42–3.51 against manuscripts 37 and 46 and against the rest of Part 3. No normalization was changed.

Here	Manuscript	Meaning	Tempting false reading
$\check{K}_d, \check{Q}_d, \check{b}_d, \check{E}_d$	K_d, Q_d, b_d^*, E_d^* (37)	integer constants (3.147)–(3.149), of size $\Theta(d^2 e^d)$	false: K_d, Q_d, E_d of (3.74); $\hat{K}_d, \hat{Q}_d$ of Theorem 3.81; $b_d(t)$
$\check{N}_d$	T_d (37)	order threshold $2\check{K}_d + \check{Q}_d + 2d$	false: a rational moment threshold T
ϱ_d	β_d (37), c (46)	largest root $\pi^2 \cot^2(\pi/d)$ of Π_d	false: the bulk count $\beta(N)$; $c_N(j, k)$
$\mathsf{T}, \mathsf{T}_d; \mathsf{T}'_7$	T (37); T (46)	rational threshold above ϱ_d : $\mathsf{T}_7 = 43$ (37), $\mathsf{T}'_7 = 21279/500$ (46)	$\check{N}_d$; the indeterminate θ of Subsection 3.28.1
$\hat{C}_7; \omega_2, \omega_3$	$C; b, a$ (46)	$(\varrho_7 - \omega_2)(\varrho_7 - \omega_3)$; the other roots of Π_7	C_d of (3.74); $b_d(t)$; $a(N)$
$\varpi, \tilde{\Pi}_\varpi, \mathsf{f}$	p, P_p, f (46)	π^2 as a parameter; Π_7 with π^2 replaced by ϖ ; $\tilde{\Pi}_\varpi(\varpi x)/\varpi^3$	$p = 2d$; P_n of Section 3.2
$V_{d,k,q}(u), v_i$	$F_{d,k,q}(u), f_i$ (37); $V_{k,q}, V_r$ (46)	the jet $(2d+u)^r \mathcal{Q}_k^{(r)}(u + (2d-q)/2)$ and its coefficients; $V_k(u)$ of (3.80) is its truncation	$\mathcal{F}_{d,k}$; the falling factorial F_n
$H_{7,k,q}$	$H_{k,q}$ (46)	the integer (3.49) at $d = 7$	harmonic numbers H_j
$\mathsf{I}_{d,k,q}(u)$	$I_{d,k,q}(u)$ (37)	the beta integral (3.136)	$\mathcal{I}_k$ of Table 5
$\mathsf{L}_{d,k,q}$	$L_{d,k,q}$ (37)	rational lower bound (3.157)	$L = \log x$
$\mathcal{B}_d(v), \mathcal{E}_{d,k,q}$	$B_d(v), \mathcal{E}_{d,k,q}$ (37); $\mathcal{B}_7, \mathbb{E}_{k,q}$ (46)	as in (3.72) and Subsection 3.17	$B(w)$; E_d
$H_n, H_n^{(2)}$	$H_n, H_n^{(2)}$ (46)	harmonic numbers	the integer $H_{d,k,q}$
$\beta(N)$	$\Lambda(N)$ (37)	bulk count (3.85)	—
$\eta_{\bar{d}}(N)$	$h_D(N)$ (37)	holes among depths $\leq \bar{d}$, $= \sum_{d \leq \bar{d}} \chi_d(N)$	$h_d(\tau)$
$\check{d}_N, \tilde{d}_N$	$D_N, \tilde{D}_N$ (37)	depths certified at order N	$D_\delta(t)$
$\rho; \sigma$	$r; s$ (37, §10)	$\lfloor N/2 \rfloor$; the layer $k + d$	$r = M - k$; $s = 2t - 1$
$\zeta; \xi$	z (37, 46); x (46, generating function)	variable of Π_d ; generating variable	$z = x^2$; the variable x
ϑ	θ (37); x (37, (3.158))	π/d ; an angle in $(0, \pi/2)$	the indeterminate θ ; the variable x
m	m, m_1, m_2 (37)	a modulus	the deficit m
C_*, N_*	C, N_0 (37)	unspecified absolute constants	C_d
$v; \alpha$	$x; a$ (46, §4)	$\tau/2$; an integer exponent	the variable x ; $a(N)$

Editorial notes. (i) [Added 1 October 2026, batch 73: this manuscript inspected the report at [f18afc619](#); depths five and six were classified here (Theorem 3.71) and the general cutoffs reduced to $\Theta(de^d)$ (Theorem 3.81) before it was placed. Its rows $d = 5, 6$ are a second certificate of Theorem 3.71; what is new is $d = 7, 8$. Its $O(d^2e^d)$ cutoffs and its $N \log N$ bound are weaker forms of Theorems 3.81 and 3.83.] (ii) [Added 1 October 2026, batch 73: “its proposed leading constant $1/2$ remain[s] open” is stale. The sibling report [power-tower-exponent-supports](#) proves $a(N) = N^2/2 + o(N^2)$ (its Corollary 13.1), without an effective threshold; the full OEIS conjecture is still open, and $3/8$ is still the leading constant of every explicit lower bound.] (iii) The manuscript’s 608,785,773 cells are its rectangles at depths five to eight (Table 3.E); of these, only the 604,164,375 cells at depths seven and eight certify new depths.

Abstract of manuscript 46. *OEIS A290268 counts the distinct terms in the fully expanded N -th derivative of x^{x^2} . Its conjectured rational generating function is equivalent to a complete description of the nonzero coefficients in a three-parameter array*

$$\gamma(k, d, M) = [w^M](2+w)^k((1+w)^2 \log(1+w))^d.$$

The ProveIt repository had classified the first six logarithmic depths and left $d \geq 7$ open. We settle the first open stratum. For every $k \geq 0$ and $M \geq 7$ we prove

$$\gamma(k, 7, M) = 0 \iff M = k + 29 \text{ and } k \text{ is odd.}$$

The displayed zeros are exactly the reflection-forced holes; there are no sporadic cancellations at depth seven.

The proof is computer-assisted but finite and reproducible. A branch-cut transform reduces nonvanishing to positivity of the expectation of the cubic

$$\Pi_7(\zeta) = \zeta^3 - 5\pi^2\zeta^2 + 3\pi^4\zeta - \frac{\pi^6}{7}.$$

A new global affine minorant, determined by the largest root of this cubic, shows that a first-moment inequality suffices. Exact integer and rational calculations give the cutoffs $R = 4898$ and $Q = 9825$, reducing all remaining cases to the rectangle

$$0 \leq k \leq 4896, \quad 0 \leq q \leq 9824,$$

containing 48,113,025 cells. An exact modular recurrence proves every nonhole in this rectangle nonzero; an independent centered-jet recurrence agrees cell by cell. As a consequence, the depth-seven contribution to the N -th derivative has the exact generating function

$$\sum_{N \geq 0} a_7(N) \xi^N = \frac{\xi^7}{(1-\xi)^2} - \frac{\xi^{31}}{1-\xi^4}.$$

The full A290268 conjecture remains open because depths $d \geq 8$ are not classified here.

Status and trust boundary (the manuscript’s box). This is an unrefereed, AI-assisted research manuscript. The infinite reduction is proved mathematically; the remaining finite theorem is certified by exact standard-library Python and C++17 programs shipped with this article. No floating-point decision is used. The result has not been formalized in Lean or another proof assistant, and no worldwide priority claim is made. Independent mathematical and code review is warranted before publication or OEIS adoption.

Editorial notes. (i) [Editorial note: the “new global affine minorant” is the case $d = 7$ of Proposition 3.80 (manuscript 70), which was in this report at the manuscript’s pin [d3c317c09](#); the factorization in Subsection 3.44 is a second proof.] (ii) [Added 1 October 2026, batch 73: “depths $d \geq 8$ are not classified here” is true of this manuscript; depth eight is classified in this addition by manuscript 37 (Theorem 3.100), so the open depths are $d \geq 9$.] (iii) “The ProveIt repository” is this report at [d3c317c09](#).

3.43 Depths seven and eight: the model and the known regions

The question. Both manuscripts start from the canonical expansion

$$\frac{d^N}{dx^N} x^{x^2} = x^{x^2} \sum_{j \in \mathbb{Z}} \sum_{k \geq 0} c_N(j, k) x^j (\log x)^k, \quad a(N) = \#\{(j, k) : c_N(j, k) \neq 0\},$$

the conjectured value $U(N)$ of (3.1) and (3.2), and the depth bridge $c_N(j, k) = N! \gamma(k, d, M) / (k! d!)$ with $d = (N + j)/2 - k$, $M = N - k$; so a depth is an infinite coefficient stratum, not a bound on N . Manuscript 37 notes that the formula was “still explicitly conjectural in the entry consulted on October 1, 2026”, and manuscript 46 that the entry “records a conjectured rational generating function and an equivalent quadratic quasipolynomial” [2].

Results of this report that the manuscripts restate. Both manuscripts say that these are prior results and credit them; manuscript 37 includes proofs “to fix all normalizations and to make the new certificates independently readable”. Each is printed once, at the place given, and its proof there is the one the manuscripts give unless a difference is named.

- The lattice recurrence for $c_N(j, k)$ with transitions $(j, k) \mapsto (j \pm 1, \cdot)$ of weights $2, 1, j, k$, the support window $|j| \leq N$, $j \equiv N \pmod{2}$, $0 \leq k \leq (N + j)/2$, and the linear independence of the monomials (37’s Proposition 2.1) are (3.54), Proposition 3.3 and Lemma 3.2 (37 proves independence by $x = e^t$ and the dominant exponential, the argument of Theorem 1.4).
- The bridge (37’s Proposition 2.2, 46’s Proposition 2.1) is Proposition 3.24; the structural zeros $M < d$ are immediate from $\varphi(w) = w + O(w^2)$.
- The parameter derivative, the centered factorization, its generating function, recurrence, positivity, degree and parity (37’s Proposition 2.3; 46’s §2.1) are (3.58) and Proposition 3.26 with (3.59), (3.61); the Meixner–Pollaczek identification both cite [20, Eq. 18.23.7] is (3.62). As there, “no orthogonality or zero-location theorem is needed”.
- The integer jets: with $p = 2d$, $r = p + q + 1$, put

$$V_{d,k,q}(u) = (p + u)^r \mathcal{Q}_k^{(r)}\left(\frac{p - q}{2} + u\right), \quad H_{d,k,q} = [u^d] V_{d,k,q}(u), \quad (3.133)$$

so that $c_N(-q - 1, k) = \binom{N}{k} H_{d,k,q}$. The recurrence

$$V_{d,-1,q} = 0, \quad V_{d,0,q}(u) = \prod_{i=0}^{p+q} (p + u - i), \quad V_{d,k+1,q} = (p - q + 2u) V_{d,k,q} + k(k + r) V_{d,k-1,q} \quad (3.134)$$

is (3.80) without truncation, and 37 adds the column step

$$V_{d,0,q+1}(u) = (u - q - 1) V_{d,0,q}(u). \quad (3.135)$$

Since multiplication cannot carry higher powers of u down, all of this may be evaluated in $\mathbb{Z}[u]/(u^{D+1})$ or $(\mathbb{Z}/\mathfrak{m}\mathbb{Z})[u]/(u^{D+1})$; 37 uses $D = d$ for nonvanishing tests and $D = 4$ for every moment anchor, “regardless of depth”.

- Bulk positivity $\gamma(k, d, M) > 0$ for $d \leq M \leq k + 2d + 1$, the $d = 0$ formula, and the reflection holes $H_{d,k,2d} = 0$ for $k + d$ even (37’s Propositions 3.1–3.2; 46’s Proposition 2.2) are Theorem 3.21, (3.57) and Theorem 3.29, with the same proofs. Manuscript 37: “proving that a symmetry forces some zeros is much easier than proving that symmetry accounts for *all* integer zeros.”

- The beta–sine identity (37’s Lemma 4.1)

$$\begin{aligned} V_{d,k,q}(u) &= (-1)^{M+1} M! \frac{\sin(\pi u)}{\pi} \mathfrak{l}_{d,k,q}(u), \\ \mathfrak{l}_{d,k,q}(u) &= \int_0^1 t^{q-u} (1-t)^{p+u} (2t-1)^k dt \quad (|u| < \tfrac{1}{2}), \end{aligned} \quad (3.136)$$

is Theorem 3.51 of the second route to depth three, there written with the shift of Table 5; 37 proves it from Euler’s reflection formula and the beta integral [14, 13] and the binomial sum $\sum_{\ell} \binom{k}{\ell} 2^{k-\ell} (-1)^{\ell} (1-a)^{-\ell} = (1-2a)^k (1-a)^{-k}$. Its coefficient of u^d , through $[u^d] \frac{\sin \pi u}{\pi} e^{-u\tau} = R_d(-\tau)/d!$, is the branch-cut formula (3.65) (Proposition 3.30), “without needing a contour argument” (37); manuscript 46 derives it by the contour deformation, as there.

- The residual factorization $R_d(-\tau) = (-1)^{d-1} d h_d(\tau) \Pi_d(\tau^2)$ (37’s Lemma 4.2; 46’s (3.67) at $d = 7$) is (3.67). For orientation, Π_5 and Π_6 are in (3.68) and

$$\Pi_7(\zeta) = \zeta^3 - 5\pi^2 \zeta^2 + 3\pi^4 \zeta - \frac{\pi^6}{7}, \quad R_7(-\tau) = 7\Pi_7(\tau^2). \quad (3.137)$$

Manuscript 37: already $\Pi_5'(\zeta) = 2\zeta - 2\pi^2$ is negative near the origin, so “the failure of monotonicity is real, not a technicality that may be ignored”.

- The folding onto $(1/2, 1)$, the folded measures and the exact sign rule (37’s §4.3; 46’s (3.69) at $d = 7$, base factor $(ty)^{14}$, holes on $q = 14$, k odd) are (3.69) and (3.70); in particular

$$\mathcal{E}_{d,k,q}[\Pi_d(\tau^2)] > 0 \implies H_{d,k,q} \neq 0 \quad (3.138)$$

outside the holes. Depths one and two (37’s Corollary 4.3, $\Pi_1 = \Pi_2 = 1$) are Theorems 3.16–3.17.

- The covariance inequality, the integer likelihood ratios and the comparison with the base measure (37’s Lemmas 5.1–5.2 and Proposition 5.3, with the real exponent $v \geq 0$ of $\mathcal{B}_d(v)$; 46’s Proposition 4.1 at $d = 7$) are Lemmas 3.32, 3.33 and Proposition 3.34, with the same tanh ratios. Manuscript 37: “Their correct object is an increasing function such as τ^2 , not an arbitrary residual polynomial”, and “the integer restriction matters: the behavior of S_{δ} for $0 < \delta < 1$ is different”.

Remark 3.88 (Second route to the base comparison (manuscript 46)). Manuscript 46 proves the comparison with the base measure through polynomials in $s = 2t - 1$ with nonnegative coefficients: for an integer $\alpha \geq 0$,

$$t^{\alpha} + y^{\alpha} = 2^{1-\alpha} \sum_{j \geq 0} \binom{\alpha}{2j} s^{2j}, \quad \frac{t^{\alpha} - y^{\alpha}}{s} = 2^{1-\alpha} \sum_{j \geq 0} \binom{\alpha}{2j+1} s^{2j},$$

both increasing in s , hence in τ ; negative exponents are reduced by the increasing factor $(ty)^{-\alpha}$, and $k \mapsto k + 2$ multiplies the density by s^2 . For $q \mapsto q + 1$ it uses the same tanh ratios as (3.71), in the variable $v = \tau/2$, and $\sinh(2\alpha v) \geq \alpha \sinh(2v)$. For $q > 14$ the odd parity minimum $k = 1$ dominates the even one $k = 0$ at $d = 7$, since their density ratio

$$s \frac{t^{q-14} - y^{q-14}}{t^{q-14} + y^{q-14}} = s \tanh\left(\frac{q-14}{2} \tau\right) \quad (3.139)$$

is positive and increasing; this is the argument in the proof of Proposition 3.77, at $d = 7$.

What each manuscript took to be known. Manuscript 37 compared itself with the report at [f18afc619](#):

Question or ingredient	Inspected report	Manuscript 37
Bulk positivity and reflection zeros	Established	Reproved and credited
Complete depths	$1 \leq d \leq 4$	$1 \leq d \leq 8$; new depths 5–8
Higher-degree residual sign changes	Crude affine minorant	Tangent at the largest root works globally
Universal fixed-depth cutoffs	$2^{d^{O(d)}}$ scale	$O(d^2 e^d)$ side lengths
Contribution beyond the bulk lower bound	Explicit linear improvements	Also an unconditional $N \log N$ improvement
Full conjecture $a(N) = U(N)$	Open	Still open

The manuscript’s caption: “The advance is relative to the inspected A290268 report, not a claim that classical polynomial, beta-integral, or likelihood-ratio methods are new.” It adds that the later depth-three contribution “is a second proof of a previously settled depth; we do not relabel it as a new result”.

Editorial note. [Added 1 October 2026, batch 73: the middle column was the report at [f18afc619](#). Before this manuscript was placed, the report gained depths 5, 6 (Theorem 3.71), the largest-root tangent (Proposition 3.80), cutoffs $\Theta(de^d)$ (Theorem 3.81) and an explicit $N \log N$ improvement (Theorem 3.83). Against the report today, the new rows are depths 7 and 8 only.]

Manuscript 46, pinned after those additions, records this state:

Depth	Method recorded in ProveIt	Status at pin
1	One-variable beta integral and a sign argument	complete
2	Divided differences, an entropy kernel, and antisymmetrization	complete
3, 4	Branch-cut kernels, likelihood-ratio comparisons, and finite exact certificates	complete
5, 6	Polynomial moment criteria, exact cutoffs, and independent finite recurrences	complete
7	First depth not covered by those classifications	open
≥ 8	General finite-reduction theorems exist, but no complete depth classification	open

It “deliberately continues the same framework instead of replacing it”, imports “the exact model, the bulk theorem, the branch-cut transform, and the monotone likelihood-ratio comparisons”, and lists as genuinely new: the global affine minorant for the depth-seven residual cubic; the cutoffs $R = 4898$, $Q = 9825$; the complete finite certificate at depth seven; the exact depth-seven count and generating function. “Novelty is assessed relative to the pinned repository and the inspected OEIS entry, not to an exhaustive search of all unpublished work.”

Editorial note. [Editorial note: the first item is the case $d = 7$ of Proposition 3.80 (manuscript 70), already at the manuscript’s pin; the factorization below is a second proof. The other three items are new. [Added 1 October 2026, batch 73: depth eight, “open” in the table, is classified in this addition by manuscript 37.]]

3.44 Depths seven and eight: the largest-root tangent

Manuscript 37's tangent is Proposition 3.80. Manuscript 37's Theorem 6.2 (“Dominant-root tangent for every depth”) states, for every integer $d \geq 3$ and every $\zeta \geq 0$,

$$\Pi_d(\zeta) \geq \Pi'_d(\varrho_d)(\zeta - \varrho_d), \quad (3.140)$$

and consequently, for every nonhole folded measure, $\mathcal{E}_{d,k,q}[\tau^2] > \varrho_d \Rightarrow \mathcal{E}_{d,k,q}[\Pi_d(\tau^2)] > 0$. Since $\Pi'_d(\varrho_d) = \prod_{\ell \geq 2} (\varrho_d - \omega_\ell) = \widehat{C}_d$, this is (3.121) of Proposition 3.80, and the proof is the same: for $d \geq 5$ and $\vartheta = \pi/d$, $\cot^2(2\vartheta)/\cot^2 \vartheta = \frac{1}{4}(1 - \tan^2 \vartheta)^2 < \frac{1}{4}$ (37's form of the double-angle step), so every other root is below $\varrho_d/4$, and then $|\zeta - \omega_\ell| \leq \varrho_d - \omega_\ell$ on $[0, \varrho_d]$. It is not printed a second time. Manuscript 37 states the last step as a lemma with a weaker hypothesis:

Remark 3.89 (A global supporting tangent (manuscript 37's Lemma 6.1)). Let $P(\zeta) = (\zeta - \varrho) \prod_{i=1}^m (\zeta - \omega_i)$ with $\varrho > 0$ and $0 \leq \omega_i \leq \varrho/2$ (the empty product allowed). Then $P(\zeta) \geq P'(\varrho)(\zeta - \varrho)$ for every $\zeta \geq 0$. The proof of Proposition 3.80 uses only $|\zeta - \omega_i| \leq \varrho - \omega_i$ on $[0, \varrho]$, which holds because the maximum of $|\zeta - \omega_i|$ there is attained at an endpoint and $\omega_i \leq \varrho - \omega_i$. “No convexity assertion is involved. In particular, the lemma applies even when P changes sign several times on the positive axis.” For Π_d the roots satisfy the stronger bound $\omega_\ell < \varrho_d/4$.

Manuscript 37 continues: “This is the decisive improvement over the crude high-degree minorant in the prior report. Instead of driving a moment beyond a bound involving a large power of d , one only needs to exceed the largest root, which is less than d^2 . The higher-degree oscillations remain present, but they lie above a single affine function of the right sign.”

Editorial note. [Added 1 October 2026, batch 73: the improvement over the crude minorant of Lemma 3.36 is Proposition 3.80, manuscript 70's, which was written into this report before manuscript 37 was placed; manuscript 37 found it independently.]

Remark 3.90 (Sufficient, not necessary (manuscript 37)). A positive expectation of Π_d can occur even when the second moment is below ϱ_d . The criterion is a one-sided certificate criterion, not a characterization of all signs. It proves eventual nonvanishing and confines possible exceptional zeros; it does not infer that every entry below a cutoff is zero, negative, or difficult for an intrinsic reason. (Compare the sharpness remark after Proposition 3.80.)

Manuscript 46: the case $d = 7$, by factoring. Manuscript 46 treats the cubic (3.137) with π^2 replaced by a parameter $\varpi > 0$:

$$\widetilde{\Pi}_\varpi(\zeta) = \zeta^3 - 5\varpi\zeta^2 + 3\varpi^2\zeta - \frac{\varpi^3}{7}, \quad \widetilde{\Pi}_{\pi^2} = \Pi_7. \quad (3.141)$$

Lemma 3.91 (Root geometry (manuscript 46)). $\widetilde{\Pi}_\varpi$ has three positive simple roots $\omega_3 < \omega_2 < \varrho$, and

$$0 < \omega_3 < \frac{\varpi}{3} < \omega_2 < 3\varpi < 4\varpi < \varrho < 5\varpi. \quad (3.142)$$

In particular $\varrho - \omega_2 - \omega_3 = 2\varrho - 5\varpi > 3\varpi > 0$.

Proof. Scale $\zeta = \varpi x$: $\widetilde{\Pi}_\varpi(\varpi x) = \varpi^3 f(x)$ with $f(x) = x^3 - 5x^2 + 3x - \frac{1}{7}$. Its critical points are $x = 1/3$ and $x = 3$, and

$$f(0) = -\frac{1}{7}, \quad f\left(\frac{1}{3}\right) = \frac{64}{189}, \quad f(3) = -\frac{64}{7}, \quad f(4) = -\frac{29}{7}, \quad f(5) = \frac{104}{7}.$$

The sign changes, with monotonicity between the critical points, prove (3.142). Vieta's relation $\omega_3 + \omega_2 + \varrho = 5\varpi$ and $\varrho > 4\varpi$ give the last claim. $\square$

At $\varpi = \pi^2$ the roots are $\omega_3 < \omega_2 < \varrho_7$, that is $\pi^2 \cot^2(3\pi/7) < \pi^2 \cot^2(2\pi/7) < \pi^2 \cot^2(\pi/7)$; the intake checked that f has the roots $\cot^2(\ell\pi/7)$, $\ell = 1, 2, 3$ ($\sum \cot^2 = 5$, $\prod \cot^2 = 1/7$).

Second proof of Proposition 3.80 at $d = 7$ (manuscript 46). Put $\hat{C}_7 = (\varrho_7 - \omega_2)(\varrho_7 - \omega_3) > 0$. Since $\Pi_7(\zeta) = (\zeta - \omega_3)(\zeta - \omega_2)(\zeta - \varrho_7)$,

$$\Pi_7(\zeta) - \hat{C}_7(\zeta - \varrho_7) = (\zeta - \varrho_7)((\zeta - \omega_3)(\zeta - \omega_2) - \hat{C}_7) = (\zeta - \varrho_7)^2(\zeta + \varrho_7 - \omega_2 - \omega_3). \quad (3.143)$$

For $\zeta \geq 0$, Lemma 3.91 gives $\zeta + \varrho_7 - \omega_2 - \omega_3 \geq \varrho_7 - \omega_2 - \omega_3 > 0$, so $\Pi_7(\zeta) \geq \hat{C}_7(\zeta - \varrho_7)$, with equality at $\zeta = \varrho_7$ and matching first derivative. If $Z \geq 0$ has finite third moment, $\mathbb{E}Z > \varrho_7$ implies $\mathbb{E}\Pi_7(Z) > 0$ (46's Corollary 5.3). $\square$

The identity (3.143) gives more than the inequality: the exact remainder, and contact of order two at ϱ_7 . Manuscript 46: “The importance of [the minorant] is that it replaces a cubic-moment problem by a first-moment problem. No bound on $\mathbb{E}Z^2$ or $\mathbb{E}Z^3$ is required.” It calls the observation “the key new analytic observation” and the minorant “sharp and unusually simple”.

Editorial note. [Editorial note: this is the case $d = 7$ of Proposition 3.80 (manuscript 70), in the report at the manuscript's pin; the factorization above is a second proof.]

Lemma 3.92 (A rational bound for the largest root at depth seven (manuscript 46)). *Put*

$$\pi_- = \frac{103993}{33102}, \quad \pi_+ = \frac{104348}{33215}, \quad T'_7 = \frac{21279}{500} = 42.558. \quad (3.144)$$

Then $\pi_- < \pi < \pi_+$ and $\varrho_7 < T'_7$. Consequently, for $Z \geq 0$ with finite third moment, $\mathbb{E}Z > T'_7$ implies $\mathbb{E}\Pi_7(Z) > 0$.

Proof. The bounds on π come from Machin's identity

$$\frac{\pi}{4} = 4 \arctan \frac{1}{5} - \arctan \frac{1}{239} \quad (3.145)$$

and alternating Taylor sums: twelve and thirteen terms at $1/5$, with two and three terms at $1/239$, already give a rational interval strictly inside (3.144); the checker compares in $\mathbb{Q}$. Let $\varpi_+ = \pi_+^2$. Exact arithmetic gives

$$\tilde{\Pi}_{\varpi_+}(T'_7) = \frac{97157793194275013545832051188177}{75195797154570635467617875000000} > 0, \quad (3.146)$$

and $T'_7 > 3\varpi_+$. At fixed $\zeta = T'_7$,

$$\frac{\partial \tilde{\Pi}_{\varpi}(T'_7)}{\partial \varpi} = -5T'^2_7 + 6\varpi T'_7 - \frac{3\varpi^2}{7} < 0 \quad (\pi_-^2 \leq \varpi \leq \varpi_+),$$

because even $-5T'^2_7 + 6\varpi_+ T'_7$ is negative. So $\Pi_7(T'_7) = \tilde{\Pi}_{\pi^2}(T'_7) > \tilde{\Pi}_{\varpi_+}(T'_7) > 0$. Since Π_7 is strictly increasing after its last critical point $3\pi^2$, its largest root satisfies $\varrho_7 < T'_7$. The second proof above then gives the consequence. $\square$

Editorial note. $\varrho_7 = \pi^2 \cot^2(\pi/7) \approx 42.5572$, so the margin is about 0.0008; manuscript 37 uses the cruder threshold $T_7 = 43$ (Table 3.D).

3.45 Depths seven and eight: single-exponential cutoffs, a weaker form of Theorem 3.81

Manuscript 37 proves, “without finite nonvanishing computations”, cutoffs of size $\Theta(d^2 e^d)$ at every depth. Theorem 3.81, already in this report, gives $\hat{Q}_d = \lceil (2d+1)e^d \rceil$ and $\hat{K}_d = 2\hat{Q}_d$, smaller by a factor of order d ; so the result is printed here as a remark, with its proof, which is a different route: the two-interval concentration estimate of Lemmas 3.37 and 3.38, with integer constants of size e^d in place of 2^{A_d} , instead of beta logarithmic moments. For an integer $d \geq 3$ put

$$\check{b}_d = 1 + \left\lceil \frac{(d+1)^{d(d+2)}}{d^{d(d+2)}} \right\rceil, \quad (3.147)$$

$$\check{K}_d = \check{b}_d(2d^2 + 15d + 8), \quad \check{Q}_d = 2d + 2\check{b}_d(2d^2 + 15d + 11), \quad (3.148)$$

$$\check{E}_d = 2d^2 + 13d + 5. \quad (3.149)$$

(The manuscript writes b_d^* , “the star distinguish[ing] this integer from the integration density $b_d(t)$ ”.)

Remark 3.93 (Single-exponential exceptional-zero confinement (manuscript 37’s Theorem 1.2); a weaker form of Theorem 3.81). Fix an integer $d \geq 3$. Every zero of $\gamma(k, d, M)$ with $M \geq d$ that is not a reflection hole has the form $M = k + 2d + 1 + q$ with $0 \leq k < \check{K}_d$ and $0 \leq q < \check{Q}_d$. The bounds satisfy $\check{K}_d, \check{Q}_d = \Theta(d^2 e^d)$. All exceptional zeros at depth d can be enumerated deterministically in $2^{O(d)}$ bit operations. “Here the parameter is the numerical depth d ; this is not a polynomial-time claim in its binary encoding length $\log d$.”

Write $\check{b} = \check{b}_d$. The inequalities $\log(1+x) > x - x^2/2$ and $\log(1+x) < x$ for $x > 0$ give

$$\log(\check{b} - 1) > d + \frac{3}{2} - \frac{1}{\check{d}} > d + 1, \quad \log \check{b} < d + 3, \quad (3.150)$$

since $\check{b} - 1 = \lceil (1 + 1/d)^{d(d+2)} \rceil < e^{d+2} + 1$, hence $\check{b} < e^{d+2} + 2 < e^{d+3}$; in particular $\check{b} = \Theta(e^d)$. Choose

$$t_0 = 1 - \frac{1}{\check{b}}, \quad t_1 = 1 - \frac{1}{2\check{b}}, \quad t_2 = 1 - \frac{1}{4\check{b}}.$$

On $t \geq t_0$, $\tau^2 \geq \log^2(\check{b} - 1) > (d+1)^2$. On $(1/2, t_0)$ the integral of $b_d(t)$ is at most $d+3$ (at most 1 for odd d ; for even d the extra factor τ is below $\log \check{b} < d+3$). On $[t_1, t_2]$ the length is $1/(4\check{b})$, $ty \geq 1/(8\check{b})$ and $h_d(\tau) \geq 1$, so

$$\int_{t_1}^{t_2} b_d(t) dt \geq \frac{1}{4\check{b}(8\check{b})^{2d}} > e^{-\check{E}_d}, \quad (3.151)$$

because $\log(4\check{b}(8\check{b})^{2d}) = (6d+2)\log 2 + (2d+1)\log \check{b} < \check{E}_d$, by $\log 2 < 1$ and $\log \check{b} < d+3$.

Lemma 3.94 (Manuscript 37’s Lemma 7.1). *If $v \geq \check{K}_d$, then $\mathcal{B}_d(v) > d^2$.*

Proof. Write $s_0 = 2t_0 - 1 = 1 - 2/\check{b}$ and $s_1 = 2t_1 - 1 = 1 - 1/\check{b}$. For the base density $b_d(t)s^v$, the ratio of the unnormalized mass below t_0 to the mass on $[t_1, t_2]$ is at most $(d+3)e^{\check{E}_d}(s_0/s_1)^v \leq (d+3)e^{\check{E}_d}e^{-v/\check{b}}$, since $s_0/s_1 = 1 - 1/(\check{b}-1) \leq e^{-1/\check{b}}$. As $\check{K}_d = \check{b}(\check{E}_d + 2d + 3)$ and $\log((d+3)(d-1)) < (d+3) + (d-1) = 2d+2 < 2d+3$, the ratio is less than $1/(d-1)$. So the normalized mass above t_0 exceeds $(d-1)/d$, and $\mathcal{B}_d(v) > \frac{d-1}{d}(d+1)^2 = d^2 + d - 1 - \frac{1}{d} > d^2$. $\square$

Lemma 3.95 (Manuscript 37’s Lemma 7.2). *If $q \geq \check{Q}_d$, then $\mathcal{E}_{d,k,q}[\tau^2] > d^2$ for every integer $k \geq 0$.*

Proof. By the monotonicity in each parity class of k (Proposition 3.34), it suffices to take $k = 0$ or $k = 1$. Put $\delta = q - 2d \geq 1$. Below t_0 both S_δ and D_δ are at most $2t_0^\delta$, so the unnormalized mass there is at most $2(d+3)t_0^\delta$. On $[t_1, t_2]$, $s^k \geq 1/2$ and both brackets are at least $t_1^\delta/2$ (for the difference, $D_\delta = t^\delta(1 - (y/t)^\delta)$ with $y/t < 1/2$), so the interval mass is at least $e^{-\tilde{E}_d t_1^\delta}/4$. The ratio is at most $8(d+3)e^{\tilde{E}_d}(t_0/t_1)^\delta \leq 8(d+3)e^{\tilde{E}_d}e^{-\delta/(2\check{b})}$. Since $\check{Q}_d - 2d = 2\check{b}(\check{E}_d + 2d + 6)$ and $\log(8(d+3)(d-1)) < 3 + 2d + 2 < 2d + 6$, the ratio is less than $1/(d-1)$; conclude as in Lemma 3.94. $\square$

Proof of Remark 3.93 (manuscript 37). Since $\tan x > x$ on $(0, \pi/2)$, $\varrho_d < d^2$. If $k \geq \check{K}_d$, Proposition 3.34 and Lemma 3.94 give a nonhole folded second moment above d^2 ; if $q \geq \check{Q}_d$, Lemma 3.95 does. The tangent criterion (Proposition 3.80) and the sign rule prove nonvanishing in both regions; the bulk covers $M \leq k + 2d + 1$. The scale $\Theta(d^2 e^d)$ follows from (3.150) and (3.148).

For the algorithm, construct $\check{b}, \check{K}_d, \check{Q}_d$ by integer powers and division and compute every $H_{d,k,q}$ of the rectangle exactly by (3.134)–(3.135) truncated at degree d ; ignore the proved holes and report the other zero entries. There are $\check{K}_d \check{Q}_d = 2^{O(d)}$ entries and $O(d)$ arithmetic operations per entry. With $L' = k + q + d + 1$ the coefficients have $O(L' \log L')$ bits (the initial falling product has this bound, and each step of the three-term recurrence adds $O(\log L')$), and $L' = 2^{O(d)}$ throughout, so even schoolbook arithmetic gives $2^{O(d)}$ bit operations; $\check{b}$ needs integers of only $O(d^2 \log d)$ bits. $\square$

Editorial note. The enumeration statement does not depend on the size of the cutoffs beyond $2^{O(d)}$, so it holds equally with the rectangle $2\hat{Q}_d \times \hat{Q}_d$ of Theorem 3.81, which has $O(d^2 e^{2d})$ cells. Neither manuscript nor this report claims the $2^{O(d)}$ bound is optimal.

Manuscript 37 turns the confinement into a threshold on the derivative order:

$$\check{N}_d = 2\check{K}_d + \check{Q}_d + 2d = 2\check{b}_d(4d^2 + 30d + 19) + 4d \quad (d \geq 3). \quad (3.152)$$

Remark 3.96 (Manuscript 37's Corollary 7.3). For every $d \geq 3$ and every $N \geq \check{N}_d$, all coefficients of depth d in the N th derivative have exactly the structural and reflection zeros, with no finite rectangle computation: if $k < \check{K}_d$ and $q < \check{Q}_d$ then $N = 2k + 2d + 1 + q \leq \check{N}_d - 2$.

Editorial note. The same argument with Theorem 3.81 gives the smaller threshold $N \geq 5\hat{Q}_d + 2d - 1$ of (3.126) (Proposition 3.82): at $d = 7, 8, 9$ it is 82,263, 253,400 and 769,812, against $\check{N}_d = 3,828,428, 12,738,042$ and 41,543,046 in the table below (intake arithmetic).

d	$\check{b}_d$	$\check{K}_d$	$\check{Q}_d$	$\check{N}_d$	$\hat{Q}_d$ (Thm 3.81)
3	76	5,396	11,254	22,052	141
4	213	21,300	43,886	86,494	492
5	592	78,736	161,034	318,516	1,633
6	1,636	278,120	566,068	1,122,320	5,245
7	4,504	950,344	1,927,726	3,828,428	16,450
8	12,367	3,165,952	6,406,122	12,738,042	50,677
9	33,885	10,334,925	20,873,178	41,543,046	153,959
10	92,711	33,190,538	66,937,362	133,318,458	462,556

Table 3.C: Manuscript 37's universal integer bounds (its Table 2), with the cutoff $\hat{Q}_d$ of Theorem 3.81 added by the intake for comparison. “These are proved analytically; the much smaller practical rectangles used below come from exact moment anchors. The row $d = 9$ is an eventual theorem, not a complete depth-nine classification.” The intake recomputed every entry.

3.46 Depths seven and eight: three moment anchors at every depth

“The universal rectangle is useful for complexity and growing-depth results. For concrete low depths, exact moment anchors give substantially smaller rectangles. Unlike the previous linear-kernel criterion, the following criterion applies to every $d \geq 3$.” It generalizes Proposition 3.41 (depths three and four, sign seeds) and the seed Lemmas 3.75 and 3.79 (depths five and six) to every depth and both parities.

Theorem 3.97 (Three moment anchors and a finite rectangle (manuscript 37)). *Fix $d \geq 3$. Choose an integer $R \geq 2$ with $R \equiv d - 1 \pmod{2}$, an integer $Q > 2d$, and a real threshold $\mathsf{T} > \varrho_d$. Let $k_s = (d - 1) \bmod 2$ and $k_a = d \bmod 2$. Suppose*

$$\mathcal{B}_d(R) > \mathsf{T}, \quad \mathcal{E}_{d,k_s,Q}[\tau^2] > \mathsf{T}, \quad \mathcal{E}_{d,k_a,Q}[\tau^2] > \mathsf{T}. \quad (3.153)$$

Then every nonhole entry with $k \geq R - 1$ or $q \geq Q$ is nonzero. Consequently the classification at depth d is settled by checking

$$0 \leq k \leq R - 2, \quad 0 \leq q \leq Q - 1 \quad (3.154)$$

for additional zeros.

Proof. In a sum case with $k \geq R - 1$, parity forces $k \geq R$, and the base comparison gives a moment at least $\mathcal{B}_d(R)$; in a difference case $k + 1 \geq R$, and the other base comparison gives the same. At $q = Q$ the two remaining anchors are the smallest representatives of the two parity classes; increasing k by two and q by one preserves the moment lower bounds, by Proposition 3.34, since $Q > 2d$. So every nonhole with $q \geq Q$ also has second moment above T . The tangent criterion (3.140) and the sign rule prove nonvanishing; the remaining bulk coefficients are positive. $\square$

Write $V_{d,k,q}(u) = \sum_i v_i u^i$.

Proposition 3.98 (Exact moment formulas (manuscript 37)). *For every nonhole folded measure,*

$$\mathcal{E}_{d,k,q}[\tau^2] = \begin{cases} 2\frac{v_3}{v_1} + \frac{\pi^2}{3}, & d \text{ odd}, \\ 6\frac{v_4}{v_2} + \pi^2, & d \text{ even}, \end{cases} \quad (3.155)$$

and the denominators are nonzero. If $q = 2d$ and $k \equiv d - 1 \pmod{2}$, this is exactly the base moment $\mathcal{B}_d(k)$.

Proof. Use $V(u) = C'(\sin(\pi u)/\pi)l(u)$ from (3.136), with $C' = (-1)^{M+1}M!$. For odd d , folding $l(0)$ and $l''(0)$ gives the same constant sign times the mass and the second moment of the folded density, because $h_d = 1$; in a nonhole the mass is nonzero. Comparing Taylor coefficients gives $v_1 = C'l(0)$ and $v_3 = C'(l''(0)/2 - \pi^2 l(0)/6)$. For even d , folding $l'(0)$ and $l'''(0)$ introduces the common factor $-\tau$, hence the density with $h_d = \tau$; their ratio is the second moment, $l'(0) \neq 0$, and $v_2 = C'l'(0)$, $v_4 = C'(l'''(0)/6 - \pi^2 l'(0)/6)$. At the stated centre the sum bracket is $S_0 = 2$, so the normalized measure is the base measure. $\square$

Editorial note. At $d = 5$ and $q = 2d$ this is (3.119), and at $d = 6$ the formula $\mathcal{B}_6(R) = \pi^2 + 6[u^4]V_R/[u^2]V_R$ in the proof of Lemma 3.79; manuscript 46’s centre formula (3.166) is the case $d = 7$.

Rational thresholds, with no numerical zero tests. Manuscript 37 uses

$$\frac{49}{5} < \pi^2 < 10, \quad (3.156)$$

proved by Machin’s formula below, so that the rational number

$$\mathsf{L}_{d,k,q} = \begin{cases} 2\frac{v_3}{v_1} + \frac{49}{15}, & d \text{ odd,} \\ 6\frac{v_4}{v_2} + \frac{49}{5}, & d \text{ even} \end{cases} \quad (3.157)$$

is a strict lower bound on the second moment. “All comparisons involving L are performed by cross-multiplying integers; no rounded floating-point value enters a certificate.” For $0 < \vartheta < \pi/2$,

$$\vartheta \cot \vartheta < 1 - \frac{\vartheta^2}{3} : \quad (3.158)$$

with $g(\vartheta) = (1 - \vartheta^2/3) \sin \vartheta - \vartheta \cos \vartheta$, $g(0) = 0$ and $g'(\vartheta) = \frac{\vartheta}{3}(\sin \vartheta - \vartheta \cos \vartheta) > 0$, since the derivative of the expression in parentheses is $\vartheta \sin \vartheta > 0$. For $d \geq 3$, (3.158) and (3.156) give $\pi \cot(\pi/d) < d - \pi^2/(3d)$ and so

$$\varrho_d < \left(d - \frac{49}{15d}\right)^2; \quad (3.159)$$

for $d = 3, 4$ the manuscript uses the exact $\varrho_3 = \pi^2/3 < 10/3$, $\varrho_4 = \pi^2 < 10$. For depths five to eight the root comparisons are

$$\frac{106276}{5625} < 19, \quad \frac{241081}{8100} < 30, \quad \frac{9604}{225} < 43, \quad \frac{829921}{14400} < 58.$$

Machin bounds. Both manuscripts certify π by Machin’s identity (3.145): with $a = \arctan(1/5)$, $b = \arctan(1/239)$, the double-angle formula gives $\tan(2a) = 5/12$, $\tan(4a) = 120/119$, and $\tan(4a - b) = 1$; the angle is positive and less than $4/5 < \pi/2$, so $4a - b = \pi/4$. Manuscript 37 truncates the alternating series:

$$\pi > 16\left(\frac{1}{5} - \frac{1}{375}\right) - \frac{4}{239} > \frac{157}{50}, \quad \pi < 16\left(\frac{1}{5} - \frac{1}{375} + \frac{1}{15625}\right) - 4\left(\frac{1}{239} - \frac{1}{3 \cdot 239^3}\right) < \frac{22}{7},$$

both by integer cross-multiplication, whence $\pi^2 > 24649/2500 > 49/5$ and $\pi^2 < 484/49 < 10$. Its exact verifier uses six and seven terms for $\arctan(1/5)$ and four and five for $\arctan(1/239)$ and records the rational interval; “No supplied decimal approximation of π is used to establish an anchor.” Manuscript 46’s finer interval is (3.144).

The program `code/09-eight-depths-verify_exact.py` recomputes all eighteen anchors from (3.134) in exact integers, verifies the nonzero denominators, checks the table’s strict rational bounds, and prints `data/09-eight-depths-exact_results.json`. It also verifies that each rational L is less than the next multiple of $1/1000$, “which checks the reported three-decimal interval rather than merely the positivity sign”. “The depth-three and depth-four rows are recertifications, not new depth classifications. Their slightly larger rectangles than the prior report’s are intentional: the same second-moment criterion is used uniformly across all six rows.”

Editorial note. The rows $d = 5, 6$ use cutoffs (398, 814) and (1461, 2944), smaller than manuscript 70’s (410, 839) and (1571, 3048) of Lemmas 3.75–3.79, with the same thresholds 19 and 30. In the intake’s reading the difference comes from the lower bound used for π^2 (here $49/5$; manuscript 70 uses $\pi > 3$) and, at depth six, from evaluating both parity-minimum tail measures exactly by Proposition 3.98, where manuscript 70 compares them with an unweighted beta law. Row $d = 7$ uses $\mathsf{T}_7 = 43$ and the cutoffs (5078, 10183); manuscript 46 uses the sharper $\mathsf{T}'_7 = 21279/500$ and obtains (4898, 9825) (Subsection 3.48).

d	T_d	R	Q	Base lower	Tail $k = 0$	Tail $k = 1$
3	10/3	14	39	3.517	3.408	3.627
4	10	89	192	10.119	10.033	10.101
5	19	398	814	19.037	19.002	19.024
6	30	1,461	2,944	30.014	30.002	30.010
7	43	5,078	10,183	43.006	43.003	43.005
8	58	16,613	33,257	58.003	58.002	58.003

Table 3.D: Manuscript 37’s exact anchor certificates (its Table 3). “Every printed decimal is an exact rational with denominator 1000 that is strictly below L in (3.157), and hence strictly below the true moment. The base uses $(k, q) = (R, 2d)$; the tail columns use $(k, q) = (0, Q)$ and $(1, Q)$. Every lower bound exceeds T . For all rows, $T > \varrho_d$.” Rows 3–6 recertify depths already classified in this report (Theorems 3.23 and 3.71); rows 7 and 8 are new.

3.47 Depths seven and eight: the theorems and manuscript 37’s certificates

Theorem 3.99 (Depth seven (manuscripts 37 and 46)). *For every $k \geq 0$ and every $M \geq 7$,*

$$\gamma(k, 7, M) = 0 \iff M = k + 29 \text{ and } k \text{ is odd.} \quad (3.160)$$

For $0 \leq M < 7$ the coefficient vanishes for the structural reason that $\varphi(w)^7$ has order seven at the origin.

Theorem 3.100 (Depth eight (manuscript 37)). *For every $k \geq 0$ and every $M \geq 8$,*

$$\gamma(k, 8, M) = 0 \iff M = k + 33 \text{ and } k \text{ is even.} \quad (3.161)$$

For $0 \leq M < 8$ the coefficient vanishes for the structural reason.

Both are the case $d = 7, 8$ of the statement that the only zeros at depth d are the reflection holes $M = k + 4d + 1$, $k + d$ even. In derivative coordinates (Corollary 3.105 for $d = 7$) the holes are $j = -15$, $N = 2k + 29$, k odd, and $j = -17$, $N = 2k + 33$, k even; in the letters of the rest of this part, $m = 7$, j odd, $n = 2j + 29$, and $m = 8$, j even, $n = 2j + 33$, the reflection family of Theorem 3.11. Manuscript 37 states the two theorems as part of one:

Corollary 3.101 (Eight-depth classification (manuscript 37’s Theorem 1.1; computer-assisted)). *Let $1 \leq d \leq 8$, $k \geq 0$ and $M \geq d$ be integers. Then*

$$\gamma(k, d, M) = 0 \iff M = k + 4d + 1 \text{ and } k + d \equiv 0 \pmod{2}. \quad (3.162)$$

For $M < d$ the coefficient vanishes because the series has order d at the origin.

Proof. Depths one and two are Theorems 3.16–3.17 (37: Corollary 4.3); three and four are Theorem 3.23; five and six are Theorem 3.71; seven and eight are Theorems 3.99 and 3.100. Manuscript 37 proves $d = 3, \dots, 8$ uniformly, by the proof below. $\square$

Editorial note. The manuscript says “The new part is $d = 5, 6, 7, 8$ ”. [Added 1 October 2026, batch 73: against this report the new part is $d = 7, 8$; $d = 5, 6$ were classified here by manuscript 70 before manuscript 37 was placed, and $d = 3, 4$ by manuscript 06.]

3.47.1 The integer predicate and the recurrence

For a chosen depth and rectangle the finite assertion is exactly

$$\forall 0 \leq k \leq R-2 \forall 0 \leq q < Q : H_{d,k,q} = 0 \iff q = 2d \text{ and } k+d \text{ even.} \quad (3.163)$$

The zero direction is already an algebraic theorem (Theorem 3.29). For the nonzero direction it is enough to exhibit a modulus $\mathfrak{m} \geq 2$ with $H \not\equiv 0 \pmod{\mathfrak{m}}$: “No primality assumption, random model, floating-point calculation, or reconstruction bound is required. A zero residue, in contrast, is never accepted as proof that $H = 0$.” The C++ verifier uses $\mathfrak{m}_1 = 1,000,000,007$; if a nonhole has zero residue it recomputes that cell modulo $\mathfrak{m}_2 = 1,000,000,009$, and a zero residue at both is marked *unresolved* and fails the run. “In the actual run, the second modulus was never needed.”

For a truncated jet $a = (a_0, \dots, a_d)$, multiplication by $c + bu$ is $((c + bu)a)_i = ca_i + ba_{i-1}$ ($a_{-1} = 0$), so (3.134) is, in coefficients,

$$v_{k+1,i} = (p - q)v_{k,i} + 2v_{k,i-1} + k(k + p + q + 1)v_{k-1,i}, \quad v_{k,-1} = 0. \quad (3.164)$$

For each q the outer loop stores $V_{d,0,q}$ by (3.135); the inner loop starts from $V_{d,-1,q} = 0$ and $V_{d,0,q}$ and keeps two consecutive jets. Induction on k proves that the degree- d component is $H_{d,k,q} \pmod{\mathfrak{m}}$, and induction on q that every cell, including the boundary, is reached once. Each product is reduced modulo the modulus; both operands are below $\mathfrak{m}_2$, so products are below $(\mathfrak{m}_2 - 1)^2 < 2^{64}$, the short sums below $4\mathfrak{m}_2$, and signed values such as $p - q$ are first converted to nonnegative residues. “The verifier is a specialized finite checker, not a claim that 64-bit arithmetic suffices for the arbitrary-depth decision algorithm.”

d	$R - 1$	Q	Cells	Reflection holes	Certified nonzero
3	13	39	507	6	501
4	88	192	16,896	44	16,852
5	397	814	323,158	198	322,960
6	1,460	2,944	4,298,240	730	4,297,510
7	5,077	10,183	51,699,091	2,538	51,696,553
8	16,612	33,257	552,465,284	8,306	552,456,978

Table 3.E: Manuscript 37’s complete finite rectangles (its Table 4), with $0 \leq k < R - 1$ and $0 \leq q < Q$. “Every nonhole was nonzero modulo 1,000,000,007. There were no unresolved cells and no disagreement with a predicted symmetry hole.” Rows 3–6 are recertifications; rows 7 and 8 certify new depths.

The four rectangles at depths five to eight contain 608,785,773 cells: 11,772 reflection holes and 608,774,001 certified nonzero entries. Their maximal derivative orders are 1616, 5874, 20349 and 66495. “These are depth-restricted rectangles, *not* a verification of every coefficient up to derivative order 66495. The all-order conclusion rests on [Theorem 3.97].” The source is `code/09-eight-depths-verify_rectangles.cpp`, its recorded output `data/09-eight-depths-rectangle_results.json`; “the recurrence generates the entries instead of storing hundreds of millions of residues. The certificate package is reproducible computation, not a compressed list whose omitted entries must be trusted without a generator.”

Proof of Theorem 3.100, and first proof of Theorem 3.99 (manuscript 37). For $d = 7, 8$ (and in the manuscript for each $d = 3, \dots, 8$), Table 3.D, certified by exact integer arithmetic, verifies the hypotheses of Theorem 3.97, so every nonhole outside (3.154) is nonzero. Table 3.E, certified by the modular recurrence, proves every nonhole inside it nonzero. Theorem 3.29 proves the holes, and Theorem 3.21 covers the bulk and the structural region. These regions exhaust every $k \geq 0$, $M \geq d$. $\square$

Remark 3.102 (A second certificate of Theorem 3.71). The same proof at $d = 5, 6$, with the rows of Tables 3.D and 3.E, is a second certificate of Theorem 3.71: different cutoffs ((398, 814) and (1461, 2944) against (410, 839) and (1571, 3048)), different rectangles (323,158 and 4,298,240 cells against 343,151 and 4,785,360), a different program and a different modulus. At $d = 3, 4$ it is a third certificate of Theorem 3.23 at $d = 3$ and a second at $d = 4$. Manuscript 37 was pinned before Theorem 3.71 was written; no novelty is claimed for these rows.

3.47.2 Independent finite cross-checks

The exact Python program performs two further kinds of cross-check, “neither of which is substituted for an infinite proof”. First, for $1 \leq d \leq 8$, $0 \leq q \leq 12$, $0 \leq k \leq 8$, it compares the recurrence jet with the direct sum

$$V_{d,k,q}(u) = \sum_{\ell=0}^k \binom{k}{\ell} 2^{k-\ell} \frac{M!}{(M-\ell)!} (2d+u)^{\overline{M-\ell}} : \quad (3.165)$$

all 936 comparisons agree through degree d . Second, it differentiates by the lattice recurrence (3.54) through $N = 80$, compares all 54 displayed OEIS terms through $N = 53$, and checks 10,408 depth-restricted integer coefficient identities against the bridge, “including zero cells that would disappear from a sparse representation”; the same run checks the upper and lower counting bounds on that range. “These checks provide evidence against indexing, factorial, and sign errors; they are not described as an independent formal proof.”

3.47.3 What is now known at every order

At depths five to eight the only negative-coordinate holes are:

Depth	Power j of x	Derivative order N	Parity of k
5	−11	$2k + 21$	odd
6	−13	$2k + 25$	even
7	−15	$2k + 29$	odd
8	−17	$2k + 33$	even

All other admissible coefficients at those depths are nonzero, “even at orders far outside any computed rectangle. Their signs need not be positive; in the large- k and large- q regions they are given explicitly by [(3.70)] with a positive residual expectation.” (The rows 5, 6 restate Theorem 3.71.)

3.48 Depth seven: a second, independent certificate (manuscript 46)

Manuscript 46’s proof has four ingredients: the exact coefficient model; the branch-cut folding to the expectation of $\Pi_7(\tau^2)$; the minorant of Subsection 3.44 with the rational threshold T'_7 of Lemma 3.92, so that $\mathcal{E}_{7,k,q}[\tau^2] > T'_7$ implies nonvanishing; and two independent exact recurrences for the remaining 48,113,025 cells. “The result is significant but deliberately scoped. It advances the depth frontier by one complete stratum; it does not classify depths $d \geq 8$, and therefore does not by itself prove the full conjectured generating function of A290268.” The cutoffs are $R = 4898$, $Q = 9825$: “Both are the first integers of the relevant parity or sequence satisfying the particular sufficient inequalities used below. This is not a claim that they are optimal among all possible proofs.”

The large- k seed. At the centre $q = 14$ put $V_r(u) = V_{7,r,14}(u) = (14 + u)^{29} Q_r^{(29)}(u)$. For even r the parameter form of the branch-cut identity is $V_r(u) = (r + 29)! \frac{\sin \pi u}{\pi} \int_0^1 (t(1-t))^{14} (2t-1)^r e^{-u\tau} dt$, and the integral is even in u ; comparing the coefficients of u and u^3 ,

$$\mathcal{B}_7(r) = \frac{\pi^2}{3} + 2 \frac{[u^3]V_r}{[u]V_r}. \quad (3.166)$$

The coefficients through degree three are integers, computed by the truncated recurrence $V_{r+1} = 2uV_r + r(r + 29)V_{r-1}$. The exact seed checker proves

$$\frac{[u^3]V_{4898}}{[u]V_{4898}} > \frac{\mathsf{T}'_7 - \pi^2/3}{2} = \frac{2016926119364}{102725850375}, \quad (3.167)$$

the left side being about 19.634554512765707 and the right about 19.63406593375694 (“the program decides the sign using the exact rational cross-product, not these decimals”), so $\mathcal{B}_7(4898) > \mathsf{T}'_7$. “The same sufficient inequality fails at $r = 4896$.” By Proposition 3.34, every nonhole with $k \geq 4897$ has $\mathcal{E}_{7,k,q}[\tau^2] > \mathsf{T}'_7$.

The large- q seed. For $k = 0$ and odd depth the folded measure is the symmetrized beta law, and differentiating the beta integral gives (3.120) at $d = 7$ (Remark 3.76):

$$\mathcal{E}_{7,0,q}[\tau^2] = (\mathsf{H}_q - \mathsf{H}_{14})^2 + \frac{\pi^2}{3} - \mathsf{H}_q^{(2)} - \mathsf{H}_{14}^{(2)}. \quad (3.168)$$

Exact rational summation proves

$$(\mathsf{H}_{9825} - \mathsf{H}_{14})^2 - \mathsf{H}_{9825}^{(2)} - \mathsf{H}_{14}^{(2)} > \mathsf{T}'_7 - \frac{\pi^2}{3} = \frac{4033852238728}{102725850375}; \quad (3.169)$$

the left side is about 39.268575692295734, exceeding the right by about 0.000443824781854. So $\mathcal{E}_{7,0,9825}[\tau^2] > \mathsf{T}'_7$; the same inequality fails at $q = 9824$. The odd minimum $k = 1$ dominates $k = 0$ by (3.139), and the steps $k \mapsto k + 2$, $q \mapsto q + 1$ preserve the order, so every nonhole with $q \geq 9825$ also has moment above T'_7 .

Proposition 3.103 (Infinite-to-finite reduction (manuscript 46)). *At depth seven every nonhole coefficient $\gamma(k, 7, k + 15 + q)$ is nonzero whenever $k \geq 4897$ or $q \geq 9825$. It remains to decide*

$$0 \leq k \leq 4896, \quad 0 \leq q \leq 9824, \quad (3.170)$$

which has $4897 \cdot 9825 = 48,113,025$ cells and largest derivative order $N = 2k + 15 + q \leq 19,631$.

Proof. The moment bounds above give $\mathcal{E}_{7,k,q}[\tau^2] > \mathsf{T}'_7$ in either region; apply Lemma 3.92 and (3.138). $\square$

Two independent recurrences. With $H_{7,k,q} = [u^7]V_{7,k,q}(u)$, the first checker uses the Pascal recurrence (3.81) at $d = 7$,

$$H_{7,k+1,q} = 2H_{7,k,q+1} + (k + 16 + q)H_{7,k,q}, \quad H_{7,0,q} = [u^7](14 + u)^{15+q}, \quad (3.171)$$

with the initial row extended through $q = 9824 + 4896$ and shortened by one cell after each step. The independent checker uses the centred recurrence (3.80) at $d = 7$, truncated modulo u^8 : $V_{7,0,q} = (14 + u)^{15+q}$, $V_{7,1,q} = (2u + 14 - q)V_{7,0,q}$, $V_{7,k+1,q} = (2u + 14 - q)V_{7,k,q} + k(k + 15 + q)V_{7,k-1,q}$. “The two algorithms traverse the finite set in different directions and arise

from different identities. Agreement therefore catches a broad class of indexing and recurrence errors.” The C++17 checker evaluates the Pascal recurrence modulo

$$1000000007, 1000000009, 998244353, 1004535809, 469762049; \quad (3.172)$$

one nonzero residue proves $H_{7,k,q} \neq 0$, and the analytically proved holes must, and do, vanish modulo every listed prime.

Theorem 3.104 (Finite rectangle certificate (manuscript 46)). *In the rectangle (3.170) the only zeros are $q = 14$, $k = 1, 3, 5, \dots, 4895$: 2448 holes. All 48,110,577 nonhole cells are nonzero modulo 1000000007, and the independent centred recurrence agrees with the Pascal recurrence modulo that prime on all 48,113,025 cells.*

Computer-assisted proof. `code/10-depth-seven-verify_depth7_rectangle.cpp` implements both recurrences in unsigned modular arithmetic: it builds the initial falling-product row, traverses the rectangle with (3.171), checks the hole predicate, records the first nonzero prime residue of every nonhole, then reconstructs every cell from the centred recurrence modulo the first prime and compares cell by cell; any discrepancy or unresolved nonhole fails. Its recorded output (`data/10-depth-seven-depth7_rectangle_certificate.json`):

Quantity	Certified value
Rectangle cells	48,113,025
Analytic holes	2448
Unresolved nonholes	0
Nonholes first resolved by 1000000007	48,110,577
Independent jet cells checked	48,113,025
Largest derivative order	19,631
Diagnostic transcript fingerprint	0xd571d65231d87a70

“The fingerprint is only a reproduction diagnostic; the proof-relevant data are the recurrences, the complete traversal, and the nonzero residues.” $\square$

Why the finite computation is a proof obligation, not evidence. “The certificate is not a numerical fit to the conjectured answer. Each cell stores an exact integer [(3.49)]; the recurrence identities are proved algebraically; and a nonzero residue modulo a prime logically excludes integer zero.” The remaining trust assumptions are “correctness of the short source, compiler, hardware, and execution. The independent recurrence materially reduces implementation risk, but it is not a proof-assistant kernel.”

Second proof of Theorem 3.99 (manuscript 46). Fix $k \geq 0$. If $M < 7$, vanishing is structural. For $7 \leq M \leq k+15$, nonvanishing is bulk positivity (Theorem 3.21). Otherwise $M = k+15+q$ with $q \geq 0$. If $q = 14$ and k is odd, the coefficient is zero (Theorem 3.29). Every other pair with $k \geq 4897$ or $q \geq 9825$ is nonzero by Proposition 3.103, and every other pair in (3.170) by Theorem 3.104. $\square$

Corollary 3.105 (Derivative-coordinate form (manuscript 46)). *Among the depth-seven coefficients of the N th derivative the only nonstructural zeros are $j = -15$, $N = 2k+29$, k odd; equivalently, for fixed N there is a depth-seven hole exactly when $N \geq 31$ and $N \equiv 3 \pmod{4}$.*

Proof. Substitute $d = 7$, $M = N - k$ and $M = k + 29$ into the bridge: $N = 2k + 29$ and $j = 2(k + 7) - N = -15$; k is odd. $\square$

Pseudocode and certificate details (manuscript 46’s appendices). The modular traversal is:

```

for each prime p:
    row[q] = coefficient of u^7 in falling(14+u, 15+q) mod p

for k = 0,...,4896:
    for q = 0,...,9824:
        hole = (q == 14 and k is odd)
        require all residues zero if hole
        require at least one nonzero residue otherwise
        row[q] = 2*row[q+1] + (k+16+q)*row[q] mod p

for q = 0,...,9824:
    reconstruct V_{k,q} mod the first prime using the centered jet recurrence
    compare coefficient u^7 with every stored Pascal value

```

“Every loop bound is a literal constant in the shipped source. Failures are explicit and do not depend on language-level assertions.” The seed certificate (`data/10-depth-seven-depth7_seed_certificate.json`) records depth 7, $R = 4898$, $Q = 9825$, $T'_7 = 21279/500$, the interval $103993/33102 < \pi < 104348/33215$, the centre ratio $\approx 19.634554512765707$ with margin $\approx 0.000488579008766$ over its exact target, the tail rational moment $\approx 39.268575692295734$ with margin $\approx 0.000443824781854$, and a SHA-256 digest of its own payload (the field `seed_payload_sha256`, not reproduced here); it also verifies that $R - 2$ and $Q - 1$ fail the same sufficient tests, and keeps the exact numerators and denominators. The manuscript reports that on its research run the rectangle checker took about 1.3 seconds and 190 MB of peak memory; “runtime is not part of the proof”.

Editorial note. The two certificates of depth seven agree wherever they overlap: both find the holes only at $q = 14$, k odd (2538 in manuscript 37’s larger rectangle, 2448 in manuscript 46’s), and both put the largest root below a rational threshold by Machin bounds (43 and $21279/500$). They share no code. Manuscript 46’s rectangle lies inside manuscript 37’s ($4896 \leq 5076$, $9824 < 10183$). At the intake both programs were rerun on copies (Subsection 3.51); the depth-seven rectangle checker took 18 s there, not 1.3 s.

3.49 Depths seven and eight: term counts

The upper bound and its missing converse (manuscript 37’s §10.1). Manuscript 37 re-derives $a(N) \leq U(N)$: indexing the coefficient triangle by the layer $\sigma = k + d$, $1 \leq \sigma \leq N$, $0 \leq k \leq \sigma$, it has $N(N+3)/2$ cells; at depth zero the top-logarithmic cells with $N > 2k$ remove exactly $\lfloor (N-1)/2 \rfloor$ entries; for even N there are no reflection holes, and for $N = 2\rho + 1$ a hole has $k = \rho - 2d$, $1 \leq d \leq \lfloor \rho/2 \rfloor$, $d \equiv \rho \pmod{2}$, which gives $\lfloor (\rho+1)/4 \rfloor$ holes. Subtracting gives (3.2). This is Theorem 3.14 with Lemma 3.13, which it credits; and “Equality for every N is equivalent to the absence of every additional zero at every positive depth. Completing only finitely many depths, even for all derivative orders, is not that assertion.” Its count of the bulk, $\sum_{\sigma=\rho}^{2\rho} (\sigma+1)$ for $N = 2\rho$ and $\rho + \sum_{\sigma=\rho+1}^{2\rho+1} (\sigma+1)$ for $N = 2\rho + 1$, gives $\beta(N)$ of (3.85), the $\Lambda(n)$ of Theorem 3.1.

Adding completed depths at a single order. Manuscript 37’s Proposition 10.1 is **Proposition 3.82** without its threshold, as in the note after that proposition: if at the order N every admissible cell of depth $1 \leq d \leq \bar{d}$ has no zero other than a reflection hole, then, with

$$\rho = \lfloor N/2 \rfloor,$$

$$a(N) \geq \beta(N) + \sum_{d=1}^{\bar{d}} \max(\rho - d, 0) - \eta_{\bar{d}}(N), \quad (3.173)$$

where

$$\eta_{\bar{d}}(N) = \begin{cases} 0, & N \text{ even,} \\ \#\{1 \leq d \leq \min(\bar{d}, \lfloor \rho/2 \rfloor) : d \equiv \rho \pmod{2}\}, & N \text{ odd.} \end{cases}$$

Here $\eta_{\bar{d}}(N) = \sum_{d \leq \bar{d}} \chi_d(N)$ (intake check: $N \geq 4d + 1$ is $d \leq \rho/2$, and $N \equiv 2d + 1 \pmod{4}$ is $d \equiv \rho \pmod{2}$), so (3.173) is (3.127); the manuscript's proof is the same count, and is not repeated. "The assumption is deliberately stated for the single order N . It may come either from a complete fixed-depth theorem or from the universal large-order criterion" (Remark 3.96).

Corollary 3.106 (Explicit eight-depth bounds (manuscript 37)). *Unconditionally,*

$$a(N) \geq \frac{3N^2 + 42N - 280}{8}, \quad N \geq 16 \text{ even}, \quad (3.174)$$

$$a(N) \geq \frac{3N^2 + 44N - 351}{8}, \quad N \geq 33 \text{ odd}. \quad (3.175)$$

For all smaller orders, (3.173) with $\bar{d} = 8$ remains valid.

Proof. Take $\bar{d} = 8$, valid at every N by Corollary 3.101. For even $N = 2\rho \geq 16$ the extra sum is $8\rho - 36$ and there are no holes. For odd $N = 2\rho + 1 \geq 33$ the same sum applies and exactly four of the first eight depths have the required parity, so the increment is $8\rho - 40$. Substitute $\beta(N)$. $\square$

"Relative to the prior four-depth lower bounds, this is an improvement of $2N - 26$ at even orders and $2N - 30$ at odd orders in the stated ranges. The leading coefficient remains $3/8$."

Editorial note. [Added 1 October 2026, batch 73: the gains $2N - 26$ and $2N - 30$ are measured from Remark 3.46 (depths up to four). Measured from Corollary 3.78 (depths up to six), whose closed forms are $(3N^2 + 34N - 160)/8$ for even $N \geq 12$ and $(3N^2 + 36N - 215)/8$ for odd $N \geq 25$, the gain of depths seven and eight is $N - 15$ for even $N \geq 16$ and $N - 17$ for odd $N \geq 33$: at odd $N \geq 33$ exactly one of $\chi_7(N)$, $\chi_8(N)$ is 1. The intake checked all four closed forms and both gains against a direct count of (3.127) for every $N < 2000$. The odd closed form (3.175) already holds at $N = 31$. "The leading coefficient remains $3/8$ " is true of every explicit bound; the leading constant itself is $\frac{1}{2}$, ineffectively, by the sibling report **power-tower-exponent-supports** (its Corollary 13.1).] This is Proposition 3.82 at the depth cap $\bar{d} = 8$, valid without the threshold (3.126) because depths up to eight are now classified; that threshold is needed only for $\bar{d} \geq 9$.

Corollary 3.107 (Exact depth-seven count (manuscript 46)). *Let $a_7(N)$ be the number of nonzero terms of depth seven in the N th derivative. Then*

$$a_7(N) = \begin{cases} 0, & 0 \leq N < 7, \\ N - 6 - \mathbf{1}\{N \geq 31, N \equiv 3 \pmod{4}\}, & N \geq 7, \end{cases} \quad (3.176)$$

and

$$\sum_{N \geq 0} a_7(N) \xi^N = \frac{\xi^7}{(1 - \xi)^2} - \frac{\xi^{31}}{1 - \xi^4}. \quad (3.177)$$

Proof. For fixed N and $d = 7$ the bridge gives $j = 2(k + 7) - N$, and the structural condition $M = N - k \geq 7$ is $0 \leq k \leq N - 7$: there are $N - 6$ admissible cells for $N \geq 7$ and none before. Corollary 3.105 removes exactly one cell for $N \geq 31$, $N \equiv 3 \pmod{4}$. Summing the arithmetic progression and the period-four holes gives (3.177). $\square$

Manuscript 46 calls (3.177) “an unconditional rational component of the conjectured full A290268 generating function”. The newly certified tail contribution at depth seven, beyond the bulk, is

$$\max\left(0, \left\lfloor \frac{N}{2} \right\rfloor - 7\right) - \mathbf{1}\{N \geq 29, N \equiv 3 \pmod{4}\}; \quad (3.178)$$

“the threshold $N \geq 29$ is harmless: the first integer satisfying the displayed congruence is 31, the first actual hole.” This is the summand of (3.127) at $d = 7$. “This is a linear improvement to pointwise lower bounds. It does not alter the leading quadratic constant by itself. A proof of the full OEIS formula still requires a uniform argument over all depths or an equivalent global support theorem.”

Editorial note. (3.176) counts all depth-seven terms, bulk included; the increment (3.178) counts those outside the bulk. The intake checked (3.177) against (3.176) for $N < 200$. The same argument with Theorem 3.100 gives $a_8(N) = N - 7 - \mathbf{1}\{N \geq 33, N \equiv 1 \pmod{4}\}$ for $N \geq 8$, with generating function $\xi^8/(1 - \xi)^2 - \xi^{33}/(1 - \xi^4)$; neither manuscript states it.

Logarithmically many certified depths (manuscript 37). The thresholds $\check{N}_d$ of (3.152) increase with d : $(1 + 1/d)^d$ increases, since $\frac{d}{dx}(x \log(1 + 1/x)) = \log(1 + 1/x) - \frac{1}{x+1} > 0$, its base is above 1 and the extra exponent $d + 2$ increases, so $\check{b}_d$ is nondecreasing, and the polynomial factors increase strictly. For an integer N put

$$\check{d}_N = \max(\{8\} \cup \{d \geq 9 : \check{N}_d \leq N\}). \quad (3.179)$$

At order N every depth through $\check{d}_N$ is classified (depths up to eight by Corollary 3.101, the others by Remark 3.96), so (3.173) with $\bar{d} = \check{d}_N$ is “an explicit unconditional lower bound, with no hidden asymptotic constants”.

Remark 3.108 (A growing-depth lower bound (manuscript 37’s Theorem 1.3); a weaker form of Theorem 3.83). There are absolute constants C_*, N_* such that, for every integer $N \geq N_*$,

$$a(N) \geq \frac{3}{8}N^2 + \frac{1}{2}N \log N - N \log \log N - C_*N; \quad (3.180)$$

in particular $\liminf_{N \rightarrow \infty} (a(N) - \frac{3}{8}N^2)/(N \log N) \geq \frac{1}{2}$. “This conclusion does not require the finite certificates at depths five through eight.” It “improves a lower bound below the proposed leading term, not an asymptotic expansion of $a(N)$ ”.

Proof (manuscript 37). Since $\check{N}_d = \Theta(d^2 e^d)$, $\log \check{N}_d = d + 2 \log d + O(1)$, and for large N , $\check{N}_{\check{d}_N} \leq N < \check{N}_{\check{d}_N+1}$; so $\log N = \check{d}_N + 2 \log \check{d}_N + O(1)$, first $\check{d}_N = \log N + O(\log \log N)$, and then

$$\check{d}_N = \log N - 2 \log \log N + O(1). \quad (3.181)$$

For large N , $\check{d}_N < \rho$, so $\sum_{d \leq \check{d}_N} (\rho - d) = \check{d}_N \rho - \check{d}_N(\check{d}_N + 1)/2$ and $0 \leq \eta_{\check{d}_N}(N) \leq \check{d}_N$. With $\beta(N) \geq 3N^2/8$ and $\rho \geq N/2 - 1/2$, $a(N) \geq \frac{3}{8}N^2 + \frac{N\check{d}_N}{2} - \frac{\check{d}_N^2}{2} - 2\check{d}_N$; insert (3.181) and absorb the $O((\log N)^2)$ term into C_*N . To avoid any finite certificate, use $\tilde{d}_N = \max(\{2\} \cup \{d \geq 3 : \check{N}_d \leq N\})$, depths one and two, and the thresholds for every $d \geq 3$; the calculation is unchanged. $\square$

Editorial note. Theorem 3.83 has the coefficient $-\frac{1}{2}$ of $N \log \log N$ in place of -1 , an explicit $-\frac{5N}{4}$ in place of $-C_*N$, and the explicit range $N \geq e^{10}$; it gives the same $\liminf \geq \frac{1}{2}$. Both proofs are the growing-depth count; the difference is the size of the cutoffs ($\hat{Q}_d = \Theta(de^d)$ against $\check{Q}_d = \Theta(d^2e^d)$), which moves the depth cap from $\log N - 2 \log \log N$ to about $\log N - \log \log N$. The $\liminf$ statement is also implied by the sibling report's $a(N) \sim N^2/2$.

Remark 3.109 (Why this does not establish the leading constant (manuscript 37)). The support still lies between a lower bound with leading term $3N^2/8$ and an upper bound with leading term $N^2/2$. Certifying $O(\log N)$ depths contributes $O(N \log N)$ entries, which is $o(N^2)$. To obtain $a(N) \sim N^2/2$, one needs a density statement controlling the remaining depths, not just an arbitrarily long fixed-depth list. Formula (3.180) is a lower-bound refinement, not a proposed second-order term of the sequence itself.

Editorial note. [Added 1 October 2026, batch 73: the density statement asked for here is proved, ineffectively, in the sibling report **power-tower-exponent-supports** (its Corollary 13.1): only $o(N^2)$ candidate cells vanish, so $a(N) \sim N^2/2$. The explicit lower bounds keep the leading constant $3/8$.]

3.50 Depths seven and eight: research questions

Manuscript 37 asks eight questions and manuscript 46 twelve. They are merged where they ask the same thing; each is printed as its source states it, the source is named, and dated notes mark those answered. Manuscript 37: “The following questions are motivated by proved statements above. They are not presented as settled consequences or as pre-existing named conjectures unless explicitly identified.” Manuscript 46: “The depth-seven proof exposes several concrete next problems.”

Research question 3.110 (Complete the integer zero classification (37); Global support theorem (46)). Does (3.162) hold for every positive depth d ? (46: prove for every d that the only zeros are the reflection holes; together with the existing counting reduction, this would establish the conjectured A290268 generating function and quasipolynomial.)

Manuscript 37: “This is the central coefficient form of the OEIS conjecture. The first uncompleted depth in this article is nine. The theorem now confines its possible exceptional cells to an explicit finite region, and guarantees the classification at every order $N \geq 41,543,046$. A successful approach should explain why the remaining integer lattice avoids the zero set, rather than just showing that the set has few real components.” It is Conjecture 3.19, Question 4 of Subsection 3.24, restricted to $m \geq 9$. *Editorial note.* With Theorem 3.81 the order threshold at depth nine is $5\hat{Q}_9 + 17 = 769,812$ (Remark 3.96, note), not $41,543,046$.

Research question 3.111 (Replace rectangular certificates by curved frontiers (37)). Can a rigorously monotone boundary in the (k, q) plane reduce the finite work substantially below $(R-1)Q$?

“The three-anchor method throws away most dependence between k and q . A piecewise frontier can start with finitely many certified moments and propagate each one upward in its allowed parity and tail region. Another option is to optimize the sufficient moment inequality directly. The correct deliverable is a finite cover of the complement of the tested region with explicit monotonicity hypotheses, not a plot suggesting that a boundary is monotone.”

Research question 3.112 (Sharpen the uniform cutoff and its polynomial factor (37); Sharp moment cutoffs (46)). What are the smallest useful uniform bounds for nonvanishing at depth d ? (46: the values 4898 and 9825 are minimal only for the sufficient inequalities used; find the

true smallest k and q beyond which the depth-seven expectation is positive for all remaining parameters, and derive asymptotics for those cutoffs as $d \rightarrow \infty$.)

Manuscript 37: “The present $O(d^2 e^d)$ bound uses a coarse two-interval concentration estimate. A refined endpoint analysis may reduce the factor d^2 , improve constants, or distinguish the k and q directions. Such a reduction would improve the coefficient of $N \log \log N$ in the explicit support lower bound. Proving optimality of an exponential scale for this particular second-moment criterion is a separate question from optimality for the nonvanishing problem itself.” *Editorial note*. [Added 1 October 2026, batch 73: answered in part. Theorem 3.81 reduces the factor d^2 to d , and Theorem 3.83 improves the coefficient of $N \log \log N$ from -1 to $-\frac{1}{2}$, both in this report before the manuscript was placed. Optimality, the true smallest cutoffs at depth seven and their asymptotics stay open; this is Question 2 of Subsection 3.24.]

Research question 3.113 (Exploit moments beyond the first moment of τ^2 (37)). Can a low-degree polynomial minorant, or several exact logarithmic moments, certify important regions where $\mathcal{E}[\tau^2] \leq \varrho_d$?

“The tangent is deliberately one-sided. Higher moments are also available from the beta–sine identity by extending the jet degree, so an exact optimization or sum-of-squares certificate could enlarge the positive region. It would need a proved nonnegative remainder on $[0, \infty)$; numerical positivity at sample points is insufficient.” This is the open part of Question 3 of Subsection 3.24.

Research question 3.114 (Find an arithmetic obstruction to accidental zeros (37); Arithmetic structure (46)). Can divisibility, valuation, or a sign-preserving combinatorial model prove nonvanishing without exhaustive residue checks? (46: study p -adic valuations and residue patterns of $H_{d,k,q}$.)

Manuscript 37: “The coefficients are integers with a short recurrence, but cancellations are genuine and signs vary. A productive arithmetic statement would supply one explicitly nonzero valuation-leading contribution or a uniformly available nonzero residue. The success of one modulus on the present rectangles is evidence worth studying, not a theorem that this modulus works at every depth or order.” Manuscript 46: “The first certificate prime resolves every depth-seven nonhole in the practical rectangle, an empirical fact that may reflect a stronger modular nonvanishing phenomenon.” (Compare Question 5 of Subsection 3.24.)

Research question 3.115 (Reach the missing density theorem (37); Density of difficult depths (46)). Can the number of unexplained cancellations at order N be shown to be $o(N^2)$, even without classifying each individual zero? (46: bound the number of unclassified or exceptional cells among all depths contributing to order N ; an $o(N^2)$ bound would recover the leading constant $1/2$.)

Manuscript 37: “This would suffice for the leading constant $1/2$. The all-depth finite confinement is not by itself a density bound when d grows proportionally to N . A new estimate should address the joint scaling of depth, logarithmic exponent, and tail index rather than holding one parameter fixed throughout.” *Editorial note*. [Added 1 October 2026, batch 73: answered, ineffectively, by the sibling report `power-tower-exponent-supports` (its Corollary 13.1, the case $a = 2$ of its Theorem 3.1); this is Question 6 of Subsection 3.24. An effective $o(N^2)$ bound is still open.]

Research question 3.116 (Build a proof-assistant certificate checker (37); Formal A290268 library (46)). Can the finite recurrence checker and the analytic reduction be verified in Lean or Rocq without enlarging the intended trust boundary? (46: complete the Lean formalization

of the all-depth hole theorem, then add the fixed-depth analytic interfaces and a verified modular evaluator.)

Manuscript 37: “A staged route is concrete: formalize jets as polynomials modulo u^{d+1} ; prove [(3.134)] and the coefficient bridge; verify residue certificates with a small checker; then formalize the likelihood-ratio comparison and global tangent lemma. The existing conditional support-to-count work in ProveIt is a natural integration point. The present package includes no untested Lean file and makes no kernel-verification claim.” Manuscript 46: “Depth seven is a useful stress test because the mathematics is finite but substantially larger than the published OEIS data range.” Its roadmap:

1. **Formal power-series model.** Define φ , γ , the derivative array, and prove the bridge algebraically.
2. **Centered factorization.** Formalize the factorization and recurrence, including parity and the reflection holes. “This would also discharge the currently conditional all-depth hole inclusion in the nearby A290268 Lean development.”
3. **Residual cubic.** Verify the root intervals, the identity (3.143), Machin’s formula, and all rational inequalities.
4. **Moment identities.** Formalize (3.168) and (3.166); the likelihood-ratio lemmas “can be abstracted for reuse at later depths”.
5. **Finite checker.** Prove the Pascal and jet recurrences over $\mathbb{Z}/p\mathbb{Z}$, then certify the fixed rectangle with a verified evaluator; “since the first prime resolves every nonhole, a formal certificate need store only one modular traversal plus the analytic hole theorem”.
6. **End-to-end theorem.** Combine the regions as in the second proof of Theorem 3.99 and translate through the bridge.

“The largest engineering question is whether to replay all 48 million cells inside the kernel, use a verified native evaluator, or produce a more compact algebraic certificate. The mathematical theorem does not depend on that choice.” This is Question 7 of Subsection 3.24; the library’s state is in Subsection 3.51.

Research question 3.117 (Transfer the tangent mechanism to other coefficient kernels (37)). Which derivative or generating-function problems produce residual polynomials whose largest positive root is at least twice every other nonnegative root?

“[Remark 3.89] is independent of A290268. Any application still needs its own correct integral representation, positive measure, and moment comparison. In particular, the related sequences A293239 and A281434 cannot be declared solved by analogy: their coefficient arrays and cancellation mechanisms require separate analysis.”

Research question 3.118 (Depth eight (46)). Classify all zeros of $\gamma(k, 8, M)$. The residual polynomial again has degree three in $Z = \tau^2$, but the folded parity and base measure differ from the odd case. Does a largest-root affine minorant give practical cutoffs comparable to those at depth seven?

Editorial note. [Added 1 October 2026, batch 73: answered by manuscript 37 in this addition (Theorem 3.100), with the largest-root tangent and the cutoffs $(R, Q) = (16613, 33257)$; manuscript 46 did not know manuscript 37.]

Research question 3.119 (Uniform largest-root minorants (46)). For general d , let Π_d be the residual polynomial in τ^2 . Determine when the tangent line at its largest root is a global lower

bound on $[0, \infty)$. A root-interlacing criterion could turn many higher depths into first-moment problems.

Editorial note. [Editorial note: answered for every $d \geq 3$ by Proposition 3.80 (manuscript 70), which was in this report at the manuscript’s pin; manuscript 37’s Remark 3.89 gives the root condition $\omega_\ell \leq \varrho_d/2$.]

Research question 3.120 (Invariant cones instead of rectangles (46)). Can the centered recurrence (3.61), augmented by finitely many parameter derivatives, preserve an explicit positivity cone that proves nonvanishing without a cell-by-cell certificate?

Research question 3.121 (Eliminate the finite certificate (46)). Find a human inequality or total-positivity theorem that proves the signs in (3.170) symbolically. The Meixner–Pollaczek interpretation of $\mathcal{Q}_k^{(r)}$ is a natural source of possible comparison theorems.

Research question 3.122 (Compress the certificate (46)). Although the checker is fast, the rectangle is large. Can one use block recurrences, polynomial remainder trees, or a verified divide-and-conquer argument to reduce the formal certificate from $O(RQ)$ cells to nearly linear or polylogarithmic size?

Research question 3.123 (Other exponent families (46)). Extend the coefficient model to x^{x^m} for $m \geq 3$. Which values of m have only symmetry-forced holes, and when do sporadic cancellations analogous to those in the x^x family appear?

Editorial note. [Added 1 October 2026, batch 73: answered in part. The sibling report **power-tower-exponent-supports** proves $A_a(N) = N^2/2 + o_a(N^2)$ for the term count of x^{x^a} at every positive integer a (its Theorem 3.1); which exponents have only symmetry-forced holes stays open. Here m is the exponent, not the deficit.]

Research question 3.124 (Asymptotic root geometry (46)). Determine the limiting distribution of the roots of Π_d after natural scaling by d and π^2 . Such information could predict whether affine, quadratic, or higher-moment criteria are optimal at large depth.

Editorial note. The roots are explicit, $\omega_\ell = \pi^2 \cot^2(\ell\pi/d)$, so $\omega_\ell/d^2 \rightarrow 1/\ell^2$ for each fixed ℓ as $d \rightarrow \infty$ (intake observation); the question’s content is the second sentence.

3.51 Depths seven and eight: status, provenance, files and formal status

3.51.1 The manuscripts’ statements of status

Manuscript 37: “The outcome consists of three logically distinct advances. The global supporting tangent and explicit single-exponential cutoffs are analytic results valid at every depth. The eight-depth classification adds exact finite computations and is therefore a computer-assisted theorem. The $N \log N$ lower-bound improvement follows from the uniform analytic cutoffs and does not depend on the large finite rectangles. The inspected report already proved many of the supporting identities and all classifications through depth four; those are credited and reproved here, not counted as new results. The new complete strata are five through eight. The scope of novelty is relative to that inspected maintained A290268 report. Searches of the OEIS entry, the relevant repository material, and related primary formulas are not an exhaustive priority search over unpublished or inaccessible work. No theorem in this manuscript has been checked by a Lean or Rocq kernel, and no independent referee or human proof audit has been performed. The exact checks were executed during preparation, and the included programs and JSON records distinguish finite verification from

the analytic exhaustion argument. No result has been submitted to OEIS and no repository file has been modified. The full formula [(3.1)] remains an open target.” *Editorial note.* [Added 1 October 2026, batch 73: against this report the tangent is Proposition 3.80, the cutoffs and the $N \log N$ improvement are weaker than Theorems 3.81 and 3.83, and the new strata are seven and eight.]

Manuscript 37’s audit table (its Appendix B):

Claim	Evidence in the package	Limitation
Canonical coefficient bridge	Taylor proof; 10,408 exact finite coefficient comparisons	Comparisons alone are not the all-order proof
Centered jet recurrence	Generating-function proof; 936 direct truncated-sum comparisons	No proof-assistant check
Global tangent and uniform cutoffs	Written inequalities and concentration proof	Not an integer nonvanishing classification by themselves
Eighteen moment anchors	Exact integer jets and rational comparisons	Depend on proved moment identities
Finite rectangles	Modular recurrence covering every specified cell	Compiler and runtime are trusted
Eight-depth theorem	Analytic exhaustion plus finite rectangles	Depths $d \geq 9$ not fully classified
$N \log N$ support increment	Growing-depth consequence of the universal thresholds	Does not establish leading constant $1/2$
Historical novelty	Comparison with pinned maintained report	No exhaustive global priority claim

It recommends auditing first the tangent inequality and the root-separation ratio, then the folded sign rule and the exact moment formulas (“where a missing sign or factorial would invalidate the anchors”), then the parity boundary $R - 1$ of Theorem 3.97 (“the tested maximum is $R - 2$, not $R - 1$ ”), then the two nested recurrence invariants and the modular implication; for the growing-depth result, the mass fraction $(d - 1)/d$, the integer definition of $\tilde{b}_d$ and the conversion $N = 2k + 2d + 1 + q$. It records that the report it inspected was the blob 7344f3aa4dc9344fb8648326e74e0d4bcc33eada of `article.tex` at f18afc619 (verified at the intake), and that the pinned README “states that the open region begins at depth five”.

Manuscript 46: “The first open depth in the ProveIt analysis of OEIS A290268 is now complete. At depth seven every cancellation is forced by the known reflection symmetry; there are no sporadic zeros. [...] The main remaining challenge is conceptual rather than computational: replace one-depth-at-a-time certificates by a uniform theorem over all d .” Its programs use exact integers and `fractions.Fraction`; “No random search, floating-point comparison, symbolic algebra package, online service, or presumed support formula is used to decide a theorem premise.”

3.51.2 Provenance

Subsections 3.42–3.51 were built from **two** manuscripts of batch 73 (cluster X), placed in this report by commit 26abf259b as its additions 09 and 10.

- Manuscript 37 (addition 09), *An Eight-Depth Nonvanishing Theorem for OEIS A290268: Dominant-root tangents, single-exponential cutoffs, and exact certificates*, 24-page PDF, dated October 1, 2026, “Research manuscript prepared for Vladimir Reshetnikov”, archive `ProveIt_A290268_Eight_Depths.zip`, arrived in f8c3a392a; pinned to f18afc619224ecccdf93fdf07f1709e56062b9e9 [24]. It contributes Theorem 3.100, the

first proof of Theorem 3.99, Corollary 3.101, Theorem 3.97, Proposition 3.98, Corollary 3.106, the second certificate of Theorem 3.71 (Remark 3.102), Remarks 3.93, 3.96, 3.108, 3.109, 3.89, 3.90, and eight questions.

- Manuscript 46 (addition 10), *Depth Seven Is Complete for OEIS A290268: A largest-root affine minorant, exact moment cutoffs, and a 48-million-cell certificate*, 17-page PDF, dated 1 October 2026, “AI-assisted research manuscript”, archive `ProveIt_A290268_Depth7.zip`, arrived in `3a9518c52`; pinned to `d3c317c09898410c046a4c4f74f12e903d40f287` [25]. It contributes the second proof of Theorem 3.99 (Subsection 3.48), the second proof of Proposition 3.80 at $d = 7$, Lemmas 3.91 and 3.92, Corollaries 3.105 and 3.107, Remark 3.88, the formalization roadmap and twelve questions.

Every theorem, proposition, lemma, corollary, remark, table, question and proof-boundary box of the two manuscripts is printed, with every non-claim and limitation they state; none was weakened or strengthened. The editorial choices were these.

- The addition is placed as subsections at the end of Part 3, after the batch-72A addition, so that no existing section, theorem or equation number moves; its four tables are numbered 3.B–3.E and the article-wide table counter is restored after each.
- Manuscript 37 is the base, because it proves both new depths; manuscript 46 is printed as the second, independent certificate of depth seven. The two depths are stated as two theorems, and manuscript 37’s single eight-depth theorem as Corollary 3.101.
- Results of this report that the manuscripts restate (model, bridge, factorization, bulk, holes, beta-sine identity, branch cut, folding, sign rule, comparisons; 37’s depth-count proposition and upper-bound count) are printed by reference, with the correspondence, in Subsections 3.43 and 3.49; where a proof differs it is printed (Remark 3.88). 37’s tangent theorem is Proposition 3.80 with the same proof and is printed by reference; its general lemma is a remark.
- Manuscript 37’s single-exponential confinement, its corollary on orders, and its $N \log N$ bound are weaker than Theorems 3.81 and 3.83, and are printed as remarks with their proofs; its rows $d = 3, \dots, 6$ as recertifications and a second certificate.
- The letters are changed as in Table 3.B, with no change of normalization. The binding changes: manuscript 37’s $K_d, Q_d, b_d^*, E_d^*, T_d$ are printed $\check{K}_d, \check{Q}_d, \check{b}_d, \check{E}_d, \check{N}_d$; its β_d and manuscript 46’s c are ϱ_d ; both manuscripts’ rational thresholds T are $\mathbb{T}$; 46’s $p = \pi^2$ is ϖ . The manuscripts’ citations of the pinned report and of its README are re-targeted to the results of this part; their OEIS and DLMF entries to this article’s bibliography.
- The twenty research questions are merged into fifteen where they ask the same thing (Subsection 3.50); manuscript 46’s formalization roadmap is printed under the merged formalization question.
- The notation subsection and table, the comparison column of Table 3.C, the editorial notes and all dated notes are additions of the intake, not of the manuscripts. Manuscript 46’s seed-payload digest is not reproduced (it stays in the shipped JSON).

Not shipped: the two manuscripts’ sources and PDFs, their `README.md` files and manuscript 46’s `SHA256SUMS.txt` (7/7 verified at placement); the archives survive in the history of `f8c3a392a` and `3a9518c52`. The READMEs add no mathematical claim. Manuscript 37’s README says the article “supplies an explicit threshold-based bound without asymptotic constants as well” (here (3.173) with $\bar{d} = \check{d}_N$); manuscript 46’s says the rectangle output “was independently reproduced with both GCC and Clang”, which this report cannot confirm beyond its own rerun.

3.51.3 Files

The manuscripts' code and data are shipped byte-identical to the delivery, with the prefixes 09-eight-depths- (37) and 10-depth-seven- (46).

Shipped path (delivery name)	Purpose
<code>code/09-eight-depths-verify_exact.py</code> (<code>code/verify_exact.py</code>)	The eighteen anchors of Table 3.D, the 936 direct-sum comparisons, the 54 OEIS terms and 10,408 bridge identities, the Machin interval (Python standard library); prints its JSON to standard output.
<code>data/09-eight-depths-exact_results.json</code> (<code>data/exact_results.json</code>)	Its recorded output.
<code>code/09-eight-depths-verify_rectangles.cpp</code> (<code>code/verify_rectangles.cpp</code>)	The six rectangles of Table 3.E modulo $10^9 + 7$, fallback $10^9 + 9$ (C++17); prints its JSON.
<code>data/09-eight-depths-rectangle_results.json</code> (<code>data/rectangle_results.json</code>)	Its recorded output.
<code>data/09-eight-depths-verification_environment.txt</code> (<code>data/verification_environment.txt</code>)	The producer's toolchain and the note that one full Clang run exceeded its time limit and is not used as evidence.
<code>code/09-eight-depths-Makefile</code> (<code>Makefile</code>)	The delivered build and verification targets; written for the delivery layout (do not use; see the README).
<code>code/10-depth-seven-verify_depth7_seeds.py</code> (<code>verify_depth7_seeds.py</code>)	Machin bounds, $\varrho_7 < T'_7$, both seeds and the failure of the same tests at $R - 2$, $Q - 1$, in exact rationals; writes <code>depth7_seed_certificate.json</code> beside itself.
<code>data/10-depth-seven-depth7_seed_certificate.json</code> (<code>depth7_seed_certificate.json</code>)	Its recorded output.
<code>code/10-depth-seven-verify_depth7_rectangle.cpp</code> (<code>verify_depth7_rectangle.cpp</code>)	Theorem 3.104: Pascal traversal modulo five primes and the independent jet replay (C++17); prints its JSON.
<code>data/10-depth-seven-depth7_rectangle_certificate.json</code> (<code>depth7_rectangle_certificate.json</code>)	Its recorded output.

The commands are in the README. The JSON files and programs keep the manuscripts' letters (R, Q, T). At the placement all four programs were rerun on copies under their delivery names (31 s, 65 s, 2 s and 18 s), and at this write again on a copy under the shipped names: each passed, and each output equalled the shipped file, ignoring line endings.

3.51.4 Formal status

Nothing in this addition is formalized, and its placement beside a Lean development confers no formal status. The Lean library A290268 (`Combinatorics/DerivativeExpansions/A290268/Lean`) relates to it as described in Section 3.26.3; neither manuscript relies on it, and neither

ships a Lean or Rocq file. Theorems 3.99 and 3.100 would discharge the nonvanishing inclusion of that library's support characterization on cells of depth seven and eight, on paper only; with the earlier theorems, on every depth at most eight. The theorems of A290268.Main stay conditional, and the library defines no γ . The formalization roadmap of Question 3.116 is a proposal.

4 x^{x^x} : OEIS A281434

Framework. This part instantiates Theorem 1.2(a) with the three-variable frame $u = x^x$, $v = x^{-1}$, $z = \log x$; Proposition 4.2 below is that instance, derived here independently, and it is the only one of the three whose frame is not two-dimensional. Because $u = x^x$ is not of the form x^λ with λ constant, the independence statement this part needs is *not* a special case of Theorem 1.4; the asymptotic argument that replaces it is given in full below.

Notation. In this part z **denotes** $\log x$, not x^2 ; compare Part 3. See Section 1.5. Throughout this part $a(n)$ denotes the term count for x^{x^x} .

Abstract of the original report. Let $a(n)$ be the number of nonzero monomials in the fully expanded n th derivative of x^{x^x} , after collecting identical terms. We replace repeated symbolic differentiation by a seven-transition integer recurrence on three exponents. It computes the prefix through n in $O(n^4)$ integer arithmetic operations and $O(n^3)$ live integers. A faster implementation performs the recurrence modulo a machine-size integer and verifies every possible zero by an independent exact coefficient formula; this method is deterministic, not a probabilistic zero test. We prove the explicit bounds

$$\frac{7n^3 + 39n^2 + 26n + 3\varepsilon_n(n-1)}{24} \leq a(n) \leq \frac{2n^3 + 3n^2 + 4n}{3}, \quad n \geq 1,$$

where ε_n is 1 for odd n and 0 for even n . In particular, $a(n) = \Theta(n^3)$. The upper bound counts an explicit lattice envelope; the lower bound uses nonvanishing leading coefficients and high-multiplicity roots of coefficient polynomials. We also identify an unavoidable family of logarithm-free cancellations and an infinite family with positive logarithm exponent. The stronger statement $a(n) \sim \frac{2}{3}n^3$, and the still stronger linear-deficit conjecture, are not proved here. Exact code, tests, terms through $n = 100$, and selected computations through $n = 200$ accompany this article.

4.1 The problem and the scope of the results

The sequence A281434 is defined in the OEIS as the number of terms in the fully expanded n th derivative of x^{x^x} , with offset 0 [3]. Exponentiation is right-associated. Thus the function in question is

$$f(x) = x^{(x^x)} = \exp(x^x \log x), \quad x > 0,$$

not $(x^x)^x = x^{x^2}$. Its initial term counts are

$$1, 3, 12, 30, 64, 113, 188, 285, 415, 577, 780, 1017, 1312, \dots$$

The positivity restriction supplies an unambiguous real logarithm; all subsequent coefficient manipulations are exact polynomial identities.

The phrase “fully expanded” must include collecting like terms and deleting zero coefficients. Counting product-rule branches is a different problem. Even recording every monomial that could be reached by differentiation gives an overcount: distinct contributions can cancel exactly.

For $n \geq 1$, define

$$B(n) = \frac{2n^3 + 3n^2 + 4n}{3}, \quad \Delta(n) = B(n) - a(n). \quad (4.1)$$

The polynomial $B(n)$ is A037237($n - 1$) [6]. The related OEIS entry A352697 records the differences $\Delta(n)$ and conjectures both $\Delta(n) = O(n)$ and infinitely many zero differences [5]. These observations are useful motivation, but neither is needed by our algorithms or proofs.

Theorem 4.1 (Main results). *There is a deterministic exact algorithm for A281434 requiring $O(n^4)$ integer arithmetic operations and $O(n^3)$ stored integers to compute $a(0), \dots, a(n)$. There is also a machine-modular implementation for a single $a(n)$, with exact verification of every candidate cancellation. For $n \geq 1$,*

$$L(n) := \frac{7n^3 + 39n^2 + 26n + 3\varepsilon_n(n-1)}{24} \leq a(n) \leq B(n), \quad \varepsilon_n = \frac{1 - (-1)^n}{2}. \quad (4.2)$$

Consequently

$$a(n) = \Theta(n^3), \quad \lim_{n \rightarrow \infty} \frac{\log a(n)}{\log n} = 3. \quad (4.3)$$

The distinction between a growth *order* and an asymptotic *equivalent* is important. Equation (4.3) is unconditional. An equivalent $a(n) \sim \frac{2}{3}n^3$ would additionally require $\Delta(n) = o(n^3)$; the present bounds do not establish that. The computations are consistent with a much smaller deficit, but this numerical observation will not be substituted for a nonvanishing theorem.

4.2 A canonical differential-polynomial representation

4.2.1 Three auxiliary variables

Put

$$u = x^x, \quad v = x^{-1}, \quad z = \log x, \quad A = z(z+1).$$

The three elementary derivatives and the logarithmic derivative of f are

$$u' = u(z+1), \quad v' = -v^2, \quad z' = v, \quad \frac{f'}{f} = u(A+v). \quad (4.4)$$

There is a unique polynomial $P_n \in \mathbb{Z}[u, v, z]$ such that

$$f^{(n)}(x) = f(x)P_n(x^x, x^{-1}, \log x). \quad (4.5)$$

Write

$$P_n(u, v, z) = \sum_{k,j,\ell \geq 0} c_n(k, j, \ell) u^k v^j z^\ell. \quad (4.6)$$

The corresponding expanded term of the original derivative is

$$c_n(k, j, \ell) x^{x^x + kx - j} (\log x)^\ell.$$

Therefore

$$a(n) = \#\{(k, j, \ell) : c_n(k, j, \ell) \neq 0\}. \quad (4.7)$$

In particular, a negative coefficient contributes one term, not minus one.

4.2.2 Why the representation is unambiguous

Suppose a finite linear combination of $u^k v^j z^\ell$, after substitution, vanishes for all sufficiently large positive x . Group it by k and choose the largest k having a nonzero Laurent-polynomial coefficient in x and $\log x$. Division by x^{kx} makes every smaller- k group exponentially small relative to that coefficient. Within the largest group, choose the smallest j , and then the largest ℓ with nonzero coefficient. Its asymptotic magnitude cannot be canceled by the remaining Laurent-logarithmic terms. This is a contradiction. Hence the displayed monomials are linearly independent in the required finite sense.

This argument is also a safeguard against a representation-dependent term count: we are counting a specified canonical basis, not the summands in an arbitrarily factored expression.

4.2.3 The recurrence and its proof

Proposition 4.2 (Differential recurrence). *Starting from $P_0 = 1$, the polynomials satisfy*

$$P_{n+1} = u(z^2 + z + v)P_n + u(z + 1)\partial_u P_n - v^2\partial_v P_n + v\partial_z P_n. \quad (4.8)$$

Proof. Differentiate (4.5), apply the product rule, and substitute (4.4). The result is exactly (4.8). The base case is $f^{(0)} = f$. Induction proves the identity at every order, and also proves integrality of every coefficient. $\square$

For one coefficient $c = c_n(k, j, \ell)$ the seven contributions are as follows. All contributions with the same destination must be added before zero coefficients are removed.

Destination at order $n + 1$	Contribution	Source in the derivative
$(k + 1, j, \ell + 2)$	c	f'/f : uz^2
$(k + 1, j, \ell + 1)$	c	f'/f : uz
$(k + 1, j + 1, \ell)$	c	f'/f : uv
$(k, j, \ell + 1)$	kc	differentiation of u^k
(k, j, ℓ)	kc	differentiation of u^k
$(k, j + 1, \ell)$	$-jc$	differentiation of v^j
$(k, j + 1, \ell - 1)$	ℓc	differentiation of z^ℓ

Transitions with a zero multiplier are omitted. In particular, no negative logarithm exponent is ever introduced.

4.2.4 The first cancellation

The first two polynomials have the compact forms

$$P_1 = u(A + v), \quad (4.9)$$

$$P_2 = u^2(A + v)^2 + u((z + 1)A + (3z + 2)v - v^2). \quad (4.10)$$

Expansion gives $a(1) = 3$ and $a(2) = 12$. At the next order, the coefficient polynomial of uv^2 is

$$-(z + 1) + (\partial_z - 1)(3z + 2) = -4z.$$

Its constant term vanishes. The candidate envelope has 31 cells at order 3, but the derivative has 30 terms. A Boolean reachability calculation would fail at this very early example.

4.3 The support envelope and the cubic upper bound

Definition 4.3. For $n \geq 1$, $1 \leq k \leq n$, and $0 \leq j \leq n$, put

$$m_{n,k,j} = \max(0, k - j), \quad (4.11)$$

$$U_{n,k,j} = \begin{cases} n + k, & j = 0, \\ \min(n + k - j - 1, 2n - 2j), & j \geq 1. \end{cases} \quad (4.12)$$

Let $\mathcal{E}_n$ consist of the integer triples with these ranges and $m_{n,k,j} \leq \ell \leq U_{n,k,j}$.

Proposition 4.4 (Support containment). *For $n \geq 1$, $\text{supp } P_n \subseteq \mathcal{E}_n$.*

Proof. Consider a nonzero differentiation path, and let a, b, c, d, e, r, s count its uses of the seven transitions in the displayed table, in that order. Then

$$k = a + b + c, \quad j = c + r + s, \quad \ell = 2a + b + d - s, \quad n = a + b + c + d + e + r + s.$$

In particular,

$$\begin{aligned} n - j &= a + b + d + e, \\ \ell - (k - j) &= a + d + r \geq 0, \\ 2(n - j) - \ell &= b + d + 2e + s \geq 0, \\ n + k - j - \ell &= b + c + e + s \geq 0. \end{aligned}$$

If $j > 0$, at least one transition of type c or s must occur: the transition of type r has multiplier $-j$ and cannot create the first positive j . Thus $n + k - j - \ell \geq 1$ when $j > 0$. The remaining ranges follow immediately from the transition table and the first differentiation. Every coefficient is a sum over such paths, so cancellation can remove cells but cannot add a cell outside these constraints. $\square$

Proposition 4.5 (Size of the envelope). *For $n \geq 1$, $\#\mathcal{E}_n = B(n)$.*

Proof. For $j = 0$ there are $n + 1$ logarithm exponents for each of n values of k , giving $n(n + 1)$ cells. For $j \geq 1$, set $d = n - j$, so $0 \leq d \leq n - 1$. The number of cells in this j -slice is

$$\sum_{k=1}^n (\min(d + k, 2d + 1) - \max(0, k - n + d)) = n(2d + 1) - d(d + 1).$$

Indeed, replacing each minimum by $2d + 1$ overcounts by $d(d + 1)/2$, and the sum of the subtracted maxima is another $d(d + 1)/2$. Consequently

$$\begin{aligned} \#\mathcal{E}_n &= n(n + 1) + \sum_{d=0}^{n-1} (n(2d + 1) - d(d + 1)) \\ &= n(n + 1) + n^3 - \frac{(n - 1)n(n + 1)}{3} \\ &= \frac{2n^3 + 3n^2 + 4n}{3}. \end{aligned}$$

$\square$

This proves the upper bound in (4.2). More precisely, $\Delta(n)$ is exactly the number of zero coefficients *inside* $\mathcal{E}_n$. The polynomial $B(n)$ is not being guessed by interpolation: it is the exact count of a proved finite set of candidate monomials.

4.4 An independent exact formula for every coefficient

The recurrence is already an exact algorithm. An independent formula is valuable for two other reasons: it certifies possible cancellations after a modular computation, and it proves that every (k, j) row is nonzero.

4.4.1 Notation for Stirling numbers and auxiliary polynomials

We use signed Stirling numbers $s(n, k)$ of the first kind and Stirling numbers $S(n, k)$ of the second kind, with the conventions

$$y^n = \prod_{i=0}^{n-1} (y - i) = \sum_k s(n, k) y^k, \quad y^n = \sum_k S(n, k) y^{\underline{k}}.$$

These conventions agree with DLMF §26.8 [12]. In particular, $s(n, k)$ has sign $(-1)^{n-k}$ and is nonzero for $1 \leq k \leq n$; $S(n, k) > 0$ on the same range.

For nonnegative d, p, q , define the noncentral Stirling numbers

$$T_d(p; q) = \sum_{r=0}^d \binom{d}{r} q^{d-r} S(r, p). \quad (4.13)$$

They can be computed without rational arithmetic:

$$T_0(p; q) = \mathbf{1}_{p=0}, \quad T_{d+1}(p; q) = (p + q)T_d(p; q) + T_d(p - 1; q). \quad (4.14)$$

Indices $p < 0$ or $p > d$ have value zero. The useful boundary values are

$$T_d(0; q) = q^d, \quad T_d(d; q) = 1, \quad T_d(p; 0) = S(d, p), \quad T_d(p; 1) = S(d + 1, p + 1),$$

where $0^0 = 1$. Equivalently,

$$e^{-w} \sum_{m \geq 0} (m + q)^d \frac{w^m}{m!} = \sum_{p=0}^d T_d(p; q) w^p. \quad (4.15)$$

To verify this identity, expand $(m + q)^d$, express powers of m in falling factorials, and sum $\sum_m m^p w^m / m! = w^p e^w$.

Define

$$Q_{j,r}(\beta) = \beta^{\underline{j}} (\beta + 1)^{\overline{r}} = \prod_{i=0}^{j-1} (\beta - i) \prod_{i=1}^r (\beta + i). \quad (4.16)$$

These integer polynomials are easy to build by multiplying by linear factors. Notice that $\deg Q_{j,r} = j + r$.

4.4.2 A parameterized derivative identity

For parameters α, β , Taylor expansion with the increment xt gives

$$\frac{(x + xt)^{\alpha(x+xt)+\beta}}{x^{\alpha x + \beta}} = (1 + t)^\beta \exp(\alpha x [tz + (1 + t) \log(1 + t)]). \quad (4.17)$$

Comparing coefficients of t^n and then of $(\alpha x)^d$ yields

$$\frac{d^n}{dx^n} x^{\alpha x + \beta} = x^{\alpha x + \beta - n} \sum_{d=0}^n (\alpha x)^d C_{n,d}(z, \beta), \quad (4.18)$$

where, with $j = n - d$,

$$C_{n,d}(z, \beta) = \sum_{s=0}^d \binom{n}{s} \frac{z^s}{(d - s)!} \partial_\beta^{d-s} Q_{j,d-s}(\beta). \quad (4.19)$$

For completeness, choose s factors tz from the d th power of the bracket in (4.17). The remaining factors contribute $(1+t)^{d-s} \log^{d-s}(1+t)$. Write the logarithm power as ∂_β^{d-s} acting on $(1+t)^{\beta+d-s}$ and use

$$[t^{n-s}](1+t)^{\beta+d-s} = \frac{(\beta+d-s)^{n-s}}{(n-s)!} = \frac{Q_{j,d-s}(\beta)}{(n-s)!}.$$

The factorial prefactors simplify to $\binom{n}{s}/(d-s)!$, proving the formula.

4.4.3 The coefficient oracle

Theorem 4.6 (Exact coefficient formula). *Suppose $(k, j, \ell) \in \mathcal{E}_n$, and put*

$$d = n - j, \quad h = d + k - \ell,$$

$$q_- = \max(0, k - \ell, k - d), \quad q_+ = \min(k, h, j).$$

Then

$$c_n(k, j, \ell) = \sum_{q=q_-}^{q_+} \binom{n}{\ell - k + q} \binom{h}{q} T_d(k - q; q) [\beta^h] Q_{j, h-q}(\beta). \quad (4.20)$$

An empty sum is zero. Outside the envelope, the coefficient is zero.

Proof. Expand the outer exponential:

$$f(x) = \sum_{m \geq 0} \frac{x^{mx} z^m}{m!}, \quad x^{mx} z^m = \partial_\beta^m x^{mx+\beta} \Big|_{\beta=0}.$$

Apply (4.18) with $\alpha = m$. For any polynomial $C(\beta)$,

$$\partial_\beta^m (e^{\beta z} C(\beta)) \Big|_{\beta=0} = \sum_{q=0}^m \frac{m!}{(m-q)!} z^{m-q} [\beta^q] C(\beta).$$

After division by $f = e^{uz}$, the sum over m occurring for fixed d, q is

$$\begin{aligned} e^{-uz} \sum_{m \geq q} \frac{m^d u^m z^{m-q}}{(m-q)!} &= u^q e^{-uz} \sum_{r \geq 0} (r+q)^d \frac{(uz)^r}{r!} \\ &= u^q \sum_{p=0}^d T_d(p; q) (uz)^p. \end{aligned}$$

To obtain $u^k z^\ell$, take $p = k - q$ and take $s = \ell - k + q$ from (4.19). Set $r = d - s = h - q$. Finally,

$$[\beta^q] \frac{\partial_\beta^r Q_{j,r}(\beta)}{r!} = \binom{q+r}{q} [\beta^{q+r}] Q_{j,r}(\beta),$$

and $q + r = h$. This proves (4.20); the stated bounds on q enforce $p \in [0, d]$, $s \in [0, d]$, and $q \leq j$.

The infinite series can be read as locally normally convergent analytic series for positive x near any fixed point, or as formal exponential identities followed by extraction of the finitely many relevant coefficients. $\square$

4.4.4 Three boundary checks

Useful independently checkable consequences are

$$[v^0]P_n = (z+1)^n \sum_{k=1}^n S(n, k)(uz)^k, \quad n \geq 1, \quad (4.21)$$

$$[u^n]P_n = (z(z+1) + v)^n, \quad (4.22)$$

$$[v^n]P_n = u^n. \quad (4.23)$$

The first follows either from (4.20) or from setting $v = 0$ in the recurrence. The second follows because reaching u -degree n requires every step to differentiate the outer factor f . For the third, the coefficient of the next highest v -power is obtained by multiplication by $u - n$, starting from 1.

4.5 A rigorous cubic lower bound

An upper bound alone does not establish cubic growth. We now show that a positive cubic number of the candidate cells must survive, without assuming that cancellations are rare.

4.5.1 Every row has its advertised highest coefficient

Let

$$R_{n,k,j}(z) = [u^k v^j]P_n(u, v, z).$$

Lemma 4.7 (Nonzero leading coefficient). *For every $1 \leq k \leq n$ and $0 \leq j \leq n$, the polynomial $R_{n,k,j}$ is nonzero and has degree exactly $U_{n,k,j}$.*

Proof. For $j = 0$, equation (4.21) gives

$$R_{n,k,0}(z) = S(n, k)z^k(z+1)^n.$$

Now let $j \geq 1$ and put $d = n - j$. If $k \leq d$, the proposed highest exponent is $\ell = d + k - 1$, so $h = 1$ in (4.20). Only $q = 0, 1$ contribute. The coefficient is

$$(-1)^{j-1}(j-1)! \left(\binom{n}{d-1} S(d, k) + \binom{n}{d} S(d+1, k) \right), \quad (4.24)$$

which is nonzero. Here we used $[\beta]Q_{j,0} = [\beta]Q_{j,1} = (-1)^{j-1}(j-1)!$.

If $k \geq d+1$, the highest exponent is $\ell = 2d$, and $h = k - d$. The only possible summand is $q = h$, giving

$$[z^{2d}]R_{n,k,j}(z) = \binom{n}{d} s(j, k-d). \quad (4.25)$$

It is nonzero because $1 \leq k - d \leq j$. The support envelope already rules out higher exponents, completing the proof. $\square$

4.5.2 A high-multiplicity root

Lemma 4.8 (Divisibility). *The row polynomial is divisible by*

$$z^{\max(k-j,0)}(z+1)^{M_{n,k,j}}, \quad M_{n,k,j} = \max(n-2j, k-j, 0). \quad (4.26)$$

Proof. Extracting $u^k v^j$ from (4.8) gives

$$\begin{aligned} R_{n+1,k,j} &= z(z+1)R_{n,k-1,j} + R_{n,k-1,j-1} + k(z+1)R_{n,k,j} \\ &\quad + (\partial_z - (j-1))R_{n,k,j-1}. \end{aligned} \quad (4.27)$$

Rows with out-of-range indices are zero. Begin with $R_{0,0,0} = 1$. For the factor $z+1$, the first and third terms increase the inherited multiplicity by one. The second term has inherited multiplicity at least $\max(n-2j+2, k-j, 0)$. The derivative in the last term can reduce its inherited multiplicity by at most one; before differentiation it is at least $\max(n-2j+2, k-j+1, 0)$. Each is sufficient for the target exponent $\max(n+1-2j, k-j, 0)$. This proves the $(z+1)$ assertion by induction. The same argument with the factor z proves the exponent $\max(k-j, 0)$. Since z and $z+1$ are relatively prime, both divisibilities hold together. $\square$

Lemma 4.9 (Multiplicity forces many terms). *A nonzero polynomial with a nonzero root of multiplicity at least M has at least $M+1$ nonzero coefficients.*

Proof. Suppose $p(z) = \sum_{i=1}^t b_i z^{e_i}$, with distinct nonnegative integers e_i and $b_i \neq 0$. If a root $\alpha \neq 0$ has multiplicity at least t , then $(z\partial_z)^r p(\alpha) = 0$ for $0 \leq r < t$. Thus

$$\sum_{i=1}^t e_i^r b_i \alpha^{e_i} = 0, \quad 0 \leq r < t.$$

The matrix (e_i^r) is an invertible Vandermonde matrix, contradicting $b_i \alpha^{e_i} \neq 0$. Therefore the multiplicity is at most $t-1$. $\square$

4.5.3 Summing the mandatory terms

By Lemmas 4.7–4.9, each (k, j) row contributes at least $1 + M_{n,k,j}$ nonzero monomials. Rows are disjoint in the canonical basis, so

$$a(n) \geq \sum_{k=1}^n \sum_{j=0}^n (1 + \max(n-2j, k-j, 0)). \quad (4.28)$$

For fixed j , the sum of the maxima over k is

$$\sum_{k=1}^n M_{n,k,j} = \begin{cases} n(n-2j) + \frac{j(j+1)}{2}, & 0 \leq j \leq \lfloor n/2 \rfloor, \\ \frac{(n-j)(n-j+1)}{2}, & j > \lfloor n/2 \rfloor. \end{cases} \quad (4.29)$$

In the first case, start with $n-2j$ in each row and add the excess $1 + \dots + j$. In the second case, only $k > j$ contributes. Summing these elementary progressions, and adding the $n(n+1)$ ones in (4.28), gives exactly $L(n)$ in (4.2). This completes the proof of the growth bounds in Theorem 4.1.

In particular,

$$\frac{7}{24} \leq \liminf_{n \rightarrow \infty} \frac{a(n)}{n^3} \leq \limsup_{n \rightarrow \infty} \frac{a(n)}{n^3} \leq \frac{2}{3}. \quad (4.30)$$

These constants are proved bounds, not estimates of an experimentally fitted leading coefficient.

4.6 Exact cancellation families

4.6.1 Logarithm-free coefficients and derivatives of a falling factorial

Let $F_n(y) = y^n$. In (4.20), setting $\ell = 0$ forces $q = k$, so for $d = n - j$ and $1 \leq k \leq j$,

$$c_n(k, j, 0) = \frac{k^d}{d! k!} F_n^{(d+k)}(d). \quad (4.31)$$

Although written using factorial quotients, this expression is an integer; the implementation instead uses integer polynomial coefficients and binomial coefficients. Formula (4.31) reduces every logarithm-free cancellation to an exact derivative evaluation of a single elementary polynomial.

Proposition 4.10 (Central cancellations at every odd order). *For odd $n \geq 1$,*

$$\Delta(n) \geq \left\lfloor \frac{n+1}{4} \right\rfloor. \quad (4.32)$$

Proof. Write $n = 2d + 1$ and set $j = d + 1$. Then

$$F_n(d + \beta) = \beta \prod_{r=1}^d (\beta^2 - r^2)$$

is odd. For each $1 \leq k \leq d + 1$ with $k \equiv d \pmod{2}$, its coefficient of β^{d+k} vanishes. Formula (4.31) gives a missing monomial $(k, d + 1, 0)$. There are $\lfloor (d + 1)/2 \rfloor = \lfloor (n + 1)/4 \rfloor$ such values of k . $\square$

Thus $a(n) = B(n)$ is impossible at every odd $n \geq 3$. This does not resolve whether equality occurs infinitely often at even orders. Nor does it classify all logarithm-free cancellations: additional integer derivative zeros occur away from the central symmetry.

4.6.2 An infinite family with positive logarithm exponent

It is tempting to guess that all cancellations are logarithm-free. The following family disproves that guess.

Since $f = e^g$ with $g = uz$, the u^{n-1} part of $f^{(n)}/f$ consists of the partition type having one block of size two and all other blocks of size one. Equivalently, induction in (4.8) gives

$$[u^{n-1}]P_n = \binom{n}{2} (A + v)^{n-2} ((z + 1)A + (3z + 2)v - v^2), \quad n \geq 2. \quad (4.33)$$

For $2 \leq j \leq n - 1$, put $m = n - j - 1$ and

$$a = \binom{n-2}{j}, \quad b = \binom{n-2}{j-1}, \quad c = \binom{n-2}{j-2}.$$

The coefficient of v^j , after removing the common factor $\binom{n}{2}$, is

$$z^m (z + 1)^m [-cz^2 + (a + 3b - c)z + (a + 2b)]. \quad (4.34)$$

Its coefficient of z^{2m+1} is

$$a + 3b - (m + 1)c. \quad (4.35)$$

Using

$$\frac{a}{c} = \frac{m(m+1)}{j(j-1)}, \quad \frac{b}{c} = \frac{m+1}{j-1},$$

this vanishes exactly when $m = j^2 - 4j$ in the stated range.

Proposition 4.11 (Non-logarithm-free cancellation family). *For every integer $t \geq 2$, let*

$$n = t^2 + t - 1, \quad k = n - 1, \quad j = t + 2, \quad \ell = 2t^2 - 7. \quad (4.36)$$

Then $(k, j, \ell) \in \mathcal{E}_n$, $\ell > 0$, and $c_n(k, j, \ell) = 0$.

Proof. Substitute $j = t + 2$ into $m = j^2 - 4j$: it gives $m = t^2 - 4$ and $n = m + j + 1 = t^2 + t - 1$. Then $\ell = 2m + 1$ is positive, is inside the envelope, and its coefficient vanishes by (4.35). The next higher coefficient is $-\binom{n}{2}c \neq 0$, so this is a genuine internal missing term. $\square$

The first corresponding orders are 5, 11, 19, 29, 41, 55, ..., with missing triples

$$(4, 4, 1), (10, 5, 11), (18, 6, 25), (28, 7, 43), \dots$$

These are additional to the logarithm-free central cancellations. Since all orders in (4.36) are odd, this proves $\Delta(n) \geq \lfloor (n+1)/4 \rfloor + 1$ along that infinite subsequence.

4.7 Algorithms and complexity

4.7.1 The dependency-free exact recurrence

The most direct implementation stores a dictionary from triples to integer coefficients. At each differentiation it applies the seven transitions, combines equal keys, and removes zeros. A short reference version is given in Section 4.10. It uses no symbolic-algebra system and no numerical approximation.

The supplied production version packs a triple into one integer key. For a requested maximum order N , take

$$b = N + 2, \quad c = 2N + 3, \quad \text{key}(k, j, \ell) = (kb + j)c + \ell.$$

Incrementing k, j, ℓ corresponds to adding $bc, c, 1$, respectively. The chosen bases exceed every possible exponent, so no carries can identify different monomials. The packed representation reduces tuple allocation and dictionary overhead; it does not change the mathematics.

Proposition 4.12 (Arithmetic complexity). *The exact recurrence computes the entire prefix through N in $O(N^4)$ integer arithmetic operations and $O(N^3)$ live integer coefficients. Each coefficient has $O(N \log N)$ bits.*

Proof. At order n there are at most $B(n) = O(n^3)$ terms, and each contributes to at most seven destinations. Summing through N gives $O(N^4)$ operations. Only the current and next dictionaries are needed, giving $O(N^3)$ storage.

Let H_n be the sum of the absolute values of the coefficients of P_n . The absolute sum of all multipliers leaving a term is $3 + 2k + j + \ell \leq 3 + 5n$. Therefore

$$H_{n+1} \leq (5n + 3)H_n, \quad H_0 = 1, \quad H_n \leq \prod_{r=0}^{n-1} (5r + 3) \leq 5^n n!.$$

Every individual coefficient satisfies this bound, so its bit length is $O(n \log n)$. $\square$

This is a unit-cost *integer-arithmetic* bound, not a claim that arbitrarily large integers are machine words. With elementary small-integer multiplication, a conservative bit bound is $O(N^5 \log^2 N)$ time and $O(N^4 \log N)$ coefficient-storage bits. Hashing and object overhead affect practical memory. We do not claim the algorithm is asymptotically optimal.

4.7.2 A deterministic modular sieve with exact refinement

Most coefficients need never be constructed as large integers. Fix any integer modulus $M \geq 2$, and run the same recurrence in $\mathbb{Z}/M\mathbb{Z}$. If a residue is nonzero, the original integer coefficient is certainly nonzero. Only zero residues need further attention.

The exact single-term algorithm is:

1. Compute $P_n \bmod M$ by the seven-transition recurrence.
2. Enumerate the triples in $\mathcal{E}_n$ whose residues are zero.
3. Evaluate each such coefficient with (4.20), using integer arithmetic, and count only the coefficients that are actually zero.
4. Return $B(n)$ minus that exact zero count.

A zero residue is *never* accepted as proof of a zero coefficient. Consequently this algorithm is correct for every admissible modulus, including small or composite moduli. No randomness, primality assumption, or conjecture about support is required. Tests deliberately use moduli 2, 7, and 12 to force many false modular zeros.

The supplied NumPy implementation uses signed 64-bit arrays and the modulus 1 000 000 007 by default. At order m , the array has shape $(m+1, m+1, 2m+1)$. The next array has shape $(m+2, m+2, 2m+3)$, and each transition becomes a shifted slice addition. All entries are reduced modulo M after each step. The implementation checks

$$(10n+10)M < 2^{63}$$

before starting; this conservative condition bounds the intermediate products and sums before modular reduction. Overflow would invalidate the certificate, so the check is part of correctness rather than merely an optimization detail.

The dense modular pass takes $O(n^4)$ fixed-width operations and $O(n^3)$ fixed-width words. Let $Z_M(n)$ denote the number of zero residues inside the envelope. The oracle tables $Q_{j,r}$ and $T_d(p;q)$, and Pascal's triangle for binomial coefficients, are built lazily. For a single target n , filling all potentially needed tables takes $O(n^3)$ integer arithmetic operations and storage; each candidate coefficient then requires at most $O(n)$ integer products and additions. Thus a conservative oracle bound is

$$O(n^3 + nZ_M(n)) \subseteq O(n^4).$$

In practice the lazy tables are much smaller when the modular pass leaves few candidates. Unlike the reference recurrence, the oracle may multiply two large integers, so its unit-arithmetic bound should again not be read as a bit-operation bound.

4.7.3 Which implementation to use

For a whole prefix, the dependency-free recurrence has the simplest predictable cost and automatically supplies every intermediate count. For an isolated larger index, the modular

method usually avoids nearly all large-integer coefficient work. Running a new modular pass separately for every prefix index would discard this advantage and incur repeated work.

Neither method differentiates a transcendental expression in a computer algebra system. Both return the exact OEIS term count, including signed cancellations, rather than an expression-size estimate.

4.8 Computational validation and results

4.8.1 Checks actually performed

The accompanying files record ten passing test groups. The tests compare the reference recurrence with all 43 published A281434 values through $n = 42$, and compare the generated prefix with all 80 published A352697 deficits through $n = 80$ [3, 5]. Every candidate coefficient through $n = 12$ is checked against the independent formula (4.20). The two methods are also compared through $n = 12$ for each of four moduli, including the deliberately small and composite choices described above.

Additional tests verify the envelope, its exact cardinality, the lower-bound sum, the $(z + 1)$ multiplicities, the two leading-coefficient formulas, the odd central cancellations, and initial instances of the infinite non-logarithm-free family. Such finite tests check the implementations and help detect algebraic mistakes; the universal statements are justified by the proofs, not by the tests.

The integer recurrence was run through $n = 100$. The modular implementation was separately checked against it at $n = 20, 40, 60, 80, 100$. Selected larger computations used the modular pass and exact verification of all candidate zeros. The archive contains the complete generated prefix, the tests, and the selected missing triples.

4.8.2 Selected exact values

n	$a(n)$	$B(n)$	$\Delta(n)$
0	1	–	–
1	3	3	0
2	12	12	0
3	30	31	1
5	113	115	2
11	1 017	1 023	6
20	5 760	5 760	0
40	44 320	44 320	0
60	147 680	147 680	0
80	347 838	347 840	2
100	676 800	676 800	0
101	697 178	697 203	25
150	2 272 700	2 272 700	0
151	2 318 265	2 318 303	38
200	5 373 600	5 373 600	0

For example, the two missing triples at $n = 80$ are $(33, 36, 0)$ and $(42, 45, 0)$. At $n = 101$, all 25 missing terms are supplied by the central logarithm-free symmetry in Proposition 4.10. These are exact statements about the recorded finite computations.

4.8.3 Measured performance

On the execution environment used for this report, the packed integer recurrence generated the full prefix $0 \leq n \leq 100$ in approximately 26.16 seconds. Selected single-index modular runs were:

n	Modular zero candidates	Actual zeros	Elapsed seconds
40	0	0	0.021
60	0	0	0.096
80	2	2	0.342
100	0	0	0.820
101	25	25	0.698
150	0	0	4.970
151	38	38	5.460
200	0	0	17.769

These are individual measurements, not a controlled benchmark suite or a promise of timings on other hardware. In particular, computing one target by the modular method and computing an entire prefix by the integer method are different workloads. The exact timings, interpreter information, and modulus are saved in `data/benchmarks.csv` and `data/environment.json`. The fact that every zero residue in this small selection was a genuine zero is not assumed by the algorithm. The small-modulus tests exercise the opposite situation and confirm that false modular zeros are rejected by the exact oracle.

4.9 What is and is not established asymptotically

The unconditional conclusion is cubic-order growth, with the fully explicit bounds (4.2). Consequently, $a(n) = n^{3+o(1)}$ in the logarithmic-exponent sense. The lower bound remains valid even if many more cancellations occur at orders far beyond the computed range.

There are three progressively stronger claims:

$$\begin{array}{ll}
 \text{Proved here:} & a(n) = \Theta(n^3). \\
 \text{Not proved here:} & a(n) \sim \frac{2}{3}n^3. \\
 \text{Stronger OEIS conjecture:} & a(n) = \frac{2}{3}n^3 + n^2 + O(n).
 \end{array}$$

The last statement is equivalent to $\Delta(n) = O(n)$, given the exact formula for $B(n)$, and implies the middle statement. Neither follows from agreement with $B(n)$ at many computed indices.

A proof of the middle statement needs only a subcubic bound on the number of holes; it need not classify every cancellation. In particular, the entire plane $\ell = 0$ has only $O(n^2)$ candidate cells. Thus even a complete failure to classify the logarithm-free derivative zeros would not by itself prevent proving the leading equivalent. The remaining issue is a uniform bound on cancellations with $\ell > 0$ across the three-dimensional envelope. Formula (4.20) supplies an explicit integer expression for those coefficients, and the family in Proposition 4.11 shows why a blanket nonvanishing assertion would be false.

For the stronger linear-deficit conjecture, even the logarithm-free plane requires more delicate control. Its central symmetry already forces $\Delta(n)$ to be at least a positive linear function along the odd orders, so an $o(n)$ bound is impossible. The appropriate target is an upper bound of matching order, not an assertion that the derivative eventually fills its whole envelope.

The computation problem is therefore resolved by unconditional exact algorithms, and the polynomial growth exponent is established. Determining the exact leading asymptotic constant remains a separate nonvanishing problem in this analysis.

4.10 A compact exact implementation

The following version emphasizes transparency. The archive's `code/a281434.py` additionally provides packed keys, the coefficient oracle, modular arrays, input validation, and a command-line interface.

```
from collections import defaultdict

def a281434_prefix(N):
    if not isinstance(N, int) or isinstance(N, bool) or N < 0:
        raise ValueError("N must be a nonnegative integer")
    p = {(0, 0, 0): 1}
    values = [1]
    for _ in range(N):
        q = defaultdict(int)
        for (k, j, ell), c in p.items():
            q[k+1, j, ell+2] += c
            q[k+1, j, ell+1] += c
            q[k+1, j+1, ell] += c
            if k:
                q[k, j, ell+1] += k*c
                q[k, j, ell] += k*c
            if j:
                q[k, j+1, ell] -= j*c
            if ell:
                q[k, j+1, ell-1] += ell*c
        p = {key: c for key, c in q.items() if c}
        values.append(len(p))
    return values
```

The only potentially negative transition is the one with multiplier $-j$. Deleting it, taking absolute values before accumulation, or replacing the dictionary by a set changes the sequence.

4.11 Reproducing the supplied computations

Python 3.9 or later is sufficient for the dependency-free method. The modular method additionally requires NumPy. From the archive's `code` directory, the commands below produce a single term, inspect missing monomials, create a prefix, and run the tests.

```
python a281434.py 100 --method modular
python a281434.py 80 --method modular --holes
python a281434.py 100 --prefix
python -m unittest -v test_a281434.py
python run_experiments.py --reference
python run_experiments.py 20 40 60 80 100 101 150 151 200
```

The first command returns 676800. The second returns 347838 and the two missing triples listed in the article. The full prefix and related deficits are in `data/b281434.txt` and `data/b352697.txt`. These are generated data files accompanying this report, not claims

that the OEIS has incorporated the additional terms.

The LaTeX source builds with

```
pdflatex -interaction=nonstopmode -halt-on-error article.tex
pdflatex -interaction=nonstopmode -halt-on-error article.tex
pdflatex -interaction=nonstopmode -halt-on-error article.tex
```

Repeated runs resolve the table of contents and cross-references. The bundled `build.sh` and `build.ps1` perform these commands.

5 The three answers side by side

5.1 What was proved

	Part 2: x^x A293239	Part 3: x^{x^2} A290268	Part 4: x^{x^x} A281434
Frame	$t = x^{-1}, L = \log x$ (2 variables)	$z = x^2, L = \log x$ (2 variables)	$u = x^x, v = x^{-1}, z = \log x$ (3 variables)
Clause of Thm 1.2	(a)	(b)	(a)
Growth order	$\Theta(n^2)$	$\Theta(n^2)$	$\Theta(n^3)$
Upper bound	$a(n) \leq q(n)$, the proposed value	$a(n) \leq U(n)$, the conjectured value	$a(n) \leq \frac{2n^3 + 3n^2 + 4n}{3}$ ($n \geq 1$; at $n = 0$ it reads $0 \geq 1$)
Lower bound	$n + 1 + \lfloor n^2/4 \rfloor$	$(3n^2 + 10n + 8)/8$ even, $(3n^2 + 12n + 1)/8$ odd	$\frac{7n^3 + 39n^2 + 26n + 3\varepsilon_n(n-1)}{24}$
Exact structure	$a(n) = 1 + \binom{n+1}{2} - Z(n)$, Z counting zeros of the Lehmer–Comtet triangle A008296	$a(n) = U(n) -$ (extra zeros), all predicted zeros explained	$a(n) = \mathcal{E}_n - \Delta(n)$, Δ the in-envelope cancellation count
Settled	the reduction; the Θ ; a <i>disproof</i> of the recurrence printed with range $n > 6$, at $n = 8$	the upper bound; the cancellation mechanism; the Θ ; a complete zero classification on the first two deficit diagonals	the Θ ; two exact algorithms; central cancellations at every odd order $n \geq 3$
Not settled	the principal closed formula: needs Conjecture 2.9	the conjecture itself, and the constant $\frac{1}{2}$: needs Conjecture 3.19	the constant: $a(n) \sim \frac{2}{3}n^3$ would need $\Delta(n) = o(n^3)$

Update (29 September 2026). For x^{x^2} the table records Part 3 as first written. Its later Sections 3.14–3.26 add to the settled column the complete zero classification on the third and fourth deficit diagonals, and improve the lower bound to $(3n^2 + 26n - 72)/8$ for even $n \geq 8$ and $(3n^2 + 28n - 111)/8$ for odd $n \geq 17$ (Remark 3.46); the leading constant $\frac{1}{2}$ is still not settled, and Conjecture 3.19 is still needed, now only on deficits $m \geq 5$.

Update (1 October 2026, batch 72A). For x^x , the later Sections 2.11–2.15 add to the settled column complete nonvanishing on every offset $\mu(p-1)$ of the Lehmer–Comtet triangle for primes $p > \mu$ under explicit conditions (for every prime when $\mu = 2, 4$, every odd prime when $\mu = 3, 5$), finite candidate lists for the offsets $(p-1)^2$ and $p^2 - 1$, and exact p -adic valuations on divisibility progressions. Conjecture 2.9 is still needed. For x^{x^2} , Sections 3.35–3.41 add the complete zero classification on the fifth and sixth deficit diagonals and the explicit lower bound $3n^2/8 + (n/2)\log n - (n/2)\log \log n - 5n/4$ for $n \geq e^{10}$ (Theorem 3.83); the leading constant of every explicit lower bound is still $3/8$, and Conjecture 3.19 is now needed only on deficits $m \geq 7$.

Update (1 October 2026, batch 73). For x^{x^2} , Sections 3.42–3.51 add the complete zero classification on the seventh and eighth deficit diagonals (Theorems 3.99 and 3.100) and the lower bound $(3n^2 + 42n - 280)/8$ for even $n \geq 16$, $(3n^2 + 44n - 351)/8$ for odd $n \geq 33$ (Corollary 3.106); for $n \geq e^{10}$ Theorem 3.83 remains the best explicit bound eventually. Conjecture 3.19 is now needed only on deficits $m \geq 9$.

[Added 1 October 2026, batch 72A: in the row “Growth order” both quadratic entries are

sharpened to $a(n) \sim n^2/2$ by the sibling report **power-tower-exponent-supports** (its Corollaries 12.1 and 13.1, the cases $a = 1, 2$ of its Theorem 3.1), without an effective threshold: for x^x , $Z(n) = o(n^2)$; for x^{x^2} , the constant $\frac{1}{2}$ of the row “Not settled” is settled in the limit. The row “Lower bound” and the updates above still give the best explicit bounds, and the closed formulas still need Conjectures 2.9 and 3.19.]

5.2 Why the orders differ

The growth orders line up with the number of variables in the frame, and Part 1 explains why the upper half of that observation must hold. In a frame of m variables, each exponent in $\text{supp } P_n$ is bounded by a constant multiple of n , because each application of the recurrence raises any one exponent by a bounded amount. The envelope therefore lies in a box of side $O(n)$ in m dimensions, so $|\mathcal{E}_n| = O(n^m)$; with Corollary 1.5 for the two frames it covers, and Part 4’s own independence argument for the third, Proposition 1.6 then gives $a(n) = O(n^m)$. With $m = 2$ for x^x and x^{x^2} and $m = 3$ for x^{x^x} , the observed orders n^2 , n^2 , n^3 are exactly the envelope dimensions.

Two cautions are worth stating, because the pattern invites more than it supports.

First, the matching *lower* bounds are not a consequence of the frame at all. A frame bounds the support from outside; it says nothing about how much of the envelope survives cancellation, and in principle almost all of it could vanish. Each part proves its own lower bound by exhibiting coefficients that are nonzero, and those arguments have nothing in common. Part 2 counts nonzero entries along each diagonal of its triangle, bounding the zeros there by the degree of a polynomial that cannot vanish identically. Part 3 proves strict *positivity* on a region, through a substitution that turns its coefficients into an integral with a sign-definite kernel. Part 4 forces a root of high multiplicity and then uses the fact that a polynomial with a nonzero root of multiplicity M has at least $M + 1$ nonzero coefficients. Three mechanisms, no common method; the framework supplies none of them.

Second, the frame is a choice, not an invariant. Nothing stops one from embedding any of these problems in a larger frame and obtaining a worse upper bound; what the three parts exhibit is the smallest frame each author found in which the derivative closes. The claim “the order is the dimension” is therefore a statement about these three constructions, not a theorem about power towers, and it should not be extrapolated to $x^{x^{x^x}}$ without doing the work.

5.3 What the shared framework buys, and what it does not

It buys four things, with two qualifications that should be stated rather than glossed. The count is an invariant of the function rather than of a differentiation algorithm — by Corollary 1.5 for Parts 2 and 3, and by Part 4’s own argument for the third frame, which that corollary does not cover. The coefficients are integers, so the computations are exact and a nonzero residue is a proof of nonvanishing (Proposition 1.8). The *envelope* bound is free once the envelope is described (Proposition 1.6) — though only Part 4’s advertised upper bound is the bare envelope count; the bounds listed for the other two in Section 5.1 are the envelope minus proved cancellation families, and those families are real theorems, not framework. And the problem is reduced from “count the terms” to “find the zeros”, which is the form all three parts actually work in.

It buys nothing at all about the zeros. That is where each of the three problems becomes its own problem, and where all three still have something open. In Part 2 the zeros are the vanishing entries of a classical triangle and the missing step is their complete classification. In Part 3 the zeros are forced by a reflection symmetry and the missing step is that there are no

others. In Part 4 the zeros are not even conjecturally classified, and the open question is the weaker one of their density. These three statements are not special cases of one another, and no argument in any part transfers to any other.

5.4 What the three do share, beyond the framework

There is more common structure than the framework alone, and it is worth setting down, because it is the one thing a reader could take from these three parts that none of them states on its own.

Every cancellation family that any of the three parts *proves* comes from the same source: the falling factorial $y^n = y(y-1)\cdots(y-n+1)$ is symmetric about its centre, so that $(n-1-y)^n = (-1)^n y^n$, and a polynomial that is odd about a point has no even coefficient there. Each part reaches that fact in its own coordinates:

- Part 2, Proposition 2.8, centres at $r = 2h$ to get $F_{4h+1}(2h+u) = u \prod_{s=1}^{2h} (u^2 - s^2)$, an odd polynomial, killing the coefficient of u^{2h} ;
- Part 3, Lemma 3.10, uses $F_n(n-1-u) = (-1)^n F_n(u)$ directly, transported through a 2-step difference;
- Part 4, Proposition 4.10, gets $F_n(d+\beta) = \beta \prod_{r=1}^d (\beta^2 - r^2)$, again odd.

So all three counting problems are, at bottom, questions about the integer zeros of derivatives and finite differences of one classical polynomial, and every zero any of them can currently explain is a parity zero of that polynomial about its centre. What is *not* shared is everything past that point: the triangles the parity acts on are different, the difference operators are different, and — this is the crux — in no case is the parity family known to be the whole zero set, which is exactly what each part’s open question asks. [Added 1 October 2026, batch 72A: the sibling report **power-tower-exponent-supports** shows that this does not extend to x^a in general: for $a = 1/2$ and every rational $1/2 < a < 1$ infinitely many cancellations lie off the reflection line (its Theorem 14.3 and Corollary 16.3). For the three functions here the statement stands.]

Given that, a caution about the two $\Theta(n^2)$ results. Parts 2 and 3 both bound a two-dimensional envelope and subtract cancellations, and both leave the leading constant open, and it would be easy to read them as one result twice. They are not: the arrays are different, one is normalized by x^{-n} and the other is not, and the open statements are statements about different objects. What can be said honestly is the metamathematical thing: no argument given in either part carries over to the other. Whether either open conjecture *implies* the other is not something this article can settle, and it does not claim to. [Added 1 October 2026, batch 72A: both leading constants are now proved, ineffectively, by one contour argument for every integer exponent, in the sibling report **power-tower-exponent-supports**; that argument is not part of this article, and the point above, that no argument given here carries over between the two parts, stands.]

5.5 What would settle each one

- x^x (**Part 2**). Conjecture 2.9: that for $1 \leq r \leq m$ the only vanishing first-kind Lehmer–Comtet entries are the sporadic $b(8, 5)$ and the symmetry family $b(4h+1, 2h)$, $h \geq 1$. The sporadic entry and the whole family are both *proved* to vanish in Part 2 — and the sporadic one is genuinely sporadic, not the first term of a second family, which is part of what makes the converse hard; the open direction is that there are no others. A proof

would give the closed formula for $a(n)$ and, with it, the corrected recurrence for the actual derivative count on its true range $n \geq 14$.

- x^{x^2} (**Part 3**). The nonvanishing statement isolated in Conjecture 3.19: that no coefficient vanishes outside the *two* explained families — the degree-forced top diagonal ($j = k$ with $n > 2k$) and the reflection family (k even, $n = 4k - 2j + 1$). Coefficients do vanish in bulk outside the reflection family alone; it is the union that is conjecturally complete. Theorem 3.20 shows the statement is exactly equivalent to the OEIS conjecture, so nothing weaker will do. Since the later addition to that part it is proved on deficits $m \leq 4$ (Theorem 3.23); what remains is $m \geq 5$. (*Update, 1 October 2026*: also on $m = 5, 6$, Theorem 3.71; what remains is $m \geq 7$. *Update, batch 73*: also on $m = 7, 8$, Theorems 3.99 and 3.100; what remains is $m \geq 9$.)
- x^{x^x} (**Part 4**). A uniform bound $\Delta(n) = o(n^3)$ would give the equivalent $a(n) \sim \frac{2}{3}n^3$; the stronger $\Delta(n) = O(n)$, conjectured not in A281434 itself but in the companion entry A352697 which records the deficits, needs control even on the logarithm-free plane, where only $O(n^2)$ cells are candidates. Whether equality $\Delta(n) = 0$ recurs infinitely often at even orders is open.

The three obstructions do not even have the same shape. In Parts 2 and 3 the open statement is a nonvanishing statement: no coefficient outside the explained families is zero. In Part 4 it cannot be, because additional zeros provably exist — Proposition 4.11 exhibits an infinite family of them with positive logarithm exponent, and Proposition 4.10 forces more at every odd order $n \geq 3$. There the open question is only *how many*: a density statement, $\Delta(n) = o(n^3)$, not a nonvanishing one. That difference is the sharpest thing the comparison has to offer, and it is the reason the three investigations are printed here as three.

References

- [1] V. Reshetnikov and contributors, *A293239: Number of terms in the fully expanded n th derivative of x^x* , The On-Line Encyclopedia of Integer Sequences, 2017. Conjectured formulas attributed in the entry to C. Barker and V. Kotešovec. Consulted 20 September 2026. <https://oeis.org/A293239>.
- [2] The OEIS Foundation Inc., *A290268: Number of terms in the fully expanded n -th derivative of x^{x^2}* , contributed by Vladimir Reshetnikov, with formula comments by Peter Luschny, October 2017. Accessed September 20, 2026. <https://oeis.org/A290268>.
- [3] Vladimir Reshetnikov, *A281434: Number of terms in the fully expanded n -th derivative of $x^{(x^x)}$* , The On-Line Encyclopedia of Integer Sequences, entry created October 5, 2017. <https://oeis.org/search?q=id:A281434&fmt=text>. Accessed September 20, 2026.
- [4] N. J. A. Sloane and contributors, *A008296: Triangle of Lehmer–Comtet numbers of the first kind*, The On-Line Encyclopedia of Integer Sequences. Consulted 20 September 2026. <https://oeis.org/A008296>.
- [5] Vladimir Reshetnikov, *A352697: $a(n) = A037237(n-1) - A281434(n)$* , The On-Line Encyclopedia of Integer Sequences, March 29, 2022. <https://oeis.org/A352697>. Accessed September 20, 2026.
- [6] N. J. A. Sloane and contributors, *A037237: Expansion of $(3+x^2)/(1-x)^4$* , The On-Line Encyclopedia of Integer Sequences. <https://oeis.org/A037237>. Accessed September 20, 2026.
- [7] L. Comtet, *Advanced Combinatorics: The Art of Finite and Infinite Expansions*, D. Reidel, Dordrecht, 1974, p. 139.
- [8] D. H. Lehmer, *Numbers associated with Stirling numbers and x^x* , Rocky Mountain Journal of Mathematics **15**(2) (1985), 461–479. <https://projecteuclid.org/euclid.rmjm/1250127221>.
- [9] H. W. Gould, *A set of polynomials associated with the higher derivatives of $y = x^x$* , Rocky Mountain Journal of Mathematics **26**(2) (1996), 615–625. <https://doi.org/10.1216/rmjm/1181072076>.
- [10] S. Jeong, *On a question of Lehmer concerning the Comtet numbers*, arXiv:2607.26613v1, 2026. In particular, Proposition 3.10(iii) records the symmetry zero family. The recurrence question answered there concerns the numbers of the *second* kind, not the complete first-kind zero classification used here. <https://arxiv.org/abs/2607.26613>.
- [11] Steven S. Poon, *Higher Derivatives of the Falling Factorial and Related Generalizations of the Stirling and Harmonic Numbers*, arXiv:1401.2737, 2014. <https://arxiv.org/abs/1401.2737>.
- [12] NIST Digital Library of Mathematical Functions, §26.8: *Set Partitions—Stirling Numbers*. <https://dlmf.nist.gov/26.8>. Accessed September 20, 2026.
- [13] NIST Digital Library of Mathematical Functions, §5.5(ii), especially Eq. 5.5.3, Euler’s reflection formula. Accessed September 20, 2026. <https://dlmf.nist.gov/5.5.E3>.
- [14] NIST Digital Library of Mathematical Functions, §5.12, especially Eq. 5.12.1, Euler’s beta integral. Accessed September 20, 2026. <https://dlmf.nist.gov/5.12.E1>.

- [15] D. M. Davis, *Coefficients in powers of the log series*, Integers **11** (2011), A16; preprint arXiv:1001.3068 (2010), Theorem 1.1 and Section 3. <https://arxiv.org/abs/1001.3068>.
- [16] M. Ravichandran, *Principal submatrices, restricted invertibility and a quantitative Gauss–Lucas theorem*, International Mathematics Research Notices **2020** (15), 4809–4832. Theorem 1.4 in arXiv version 2, attributed there to Marcus. <https://arxiv.org/abs/1609.04187>.
- [17] A. Adelberg and T. Lengyel, *New results on the p -adic valuation of Stirling numbers*, Integers **22** (2022), A82. <https://math.colgate.edu/~integers/w82/w82.pdf>.
- [18] V. Reshetnikov, *ProveIt: A290268 investigation notes*, Oeis/A290268/README.md, snapshot afb2d1227d8bc5df3beffc1463e958db3544960b, accessed 29 September 2026. <https://github.com/VladimirReshetnikov/ProveIt/blob/afb2d1227d8bc5df3beffc1463e958db3544960b/Oeis/A290268/README.md>.
- [19] V. Reshetnikov, *ProveIt: A290268 support-to-count reduction*, Combinatorics/DerivativeExpansions/A290268/Lean/A290268/Main.lean, same snapshot, accessed 29 September 2026. <https://github.com/VladimirReshetnikov/ProveIt/blob/afb2d1227d8bc5df3beffc1463e958db3544960b/Combinatorics/DerivativeExpansions/A290268/Lean/A290268/Main.lean>.
- [20] NIST Digital Library of Mathematical Functions, *Hahn class: generating functions*, Section 18.23, Eq. 18.23.7 (Meixner–Pollaczek generating function), accessed 29 September 2026. <https://dlmf.nist.gov/18.23.E7>.
- [21] F. Štampach, *Asymptotic behavior and zeros of the Bernoulli polynomials of the second kind*, Journal of Approximation Theory **262** (2021), article 105517. Preprint arXiv:2011.13808 (2020). <https://arxiv.org/abs/2011.13808>.
- [22] V. Reshetnikov, *A290268: number of terms in the expanded n -th derivative of x^{x^2}* , ProveIt repository, file Oeis/A290268/README.md, commit a02ab3567fef25793b27a68248b58a7b69694342, 2026. <https://github.com/VladimirReshetnikov/ProveIt/blob/a02ab3567fef25793b27a68248b58a7b69694342/Oeis/A290268/README.md>.
- [23] V. Reshetnikov, *A290268: reduction of the conjecture to the support characterization*, ProveIt repository, file Combinatorics/DerivativeExpansions/A290268/Lean/A290268/Main.lean, same commit, 2026. <https://github.com/VladimirReshetnikov/ProveIt/blob/a02ab3567fef25793b27a68248b58a7b69694342/Combinatorics/DerivativeExpansions/A290268/Lean/A290268/Main.lean>.
- [24] ProveIt research corpus, *Term counts in the derivatives of power towers: OEIS A293239, A290268, A281434*, article.tex and README.md of this report at commit f18afc619224ecccdf93fdf07f1709e56062b9e9 (manuscript 37’s pinned source), consulted 1 October 2026. <https://github.com/VladimirReshetnikov/ProveIt/tree/f18afc619224ecccdf93fdf07f1709e56062b9e9>.
- [25] V. Reshetnikov, *ProveIt repository*, commit d3c317c09898410c046a4c4f74f12e903d40f287, especially Oeis/A290268 and this report (manuscript 46’s pinned source), inspected 1 October 2026. <https://github.com/VladimirReshetnikov/ProveIt/tree/d3c317c09898410c046a4c4f74f12e903d40f287>.