

A manifest of docs/reports

One hundred and eighty independent mathematical research packages,
classified by subject

Vladimir Reshetnikov

2 October 2026

1 Scope and conventions

This document catalogues the one hundred and eighty research packages stored under `docs/reports`. The collection originally held one hundred and four reports; after two consolidations it contributes one hundred and two maintained reports. The other seventy-eight came through ProveIt’s incoming-reports procedure (`docs/incoming`) — six in batch 36, one each in batches 40, 41 and 44, two in batch 60, one each in batches 62 and 70, three in batch 72, sixteen in batch 73, two in batch 74, three in batch 75, two in batch 76, thirty-four in batch 77, three in batch 78 and two in batch 79 — whose archives remain in the repository’s history. Later batches also extended existing reports; their entries name the batch. The next paragraphs describe the historical collection before those two consolidations. On 29 September 2026 the two Lipparini Problem 6.2 reports were unified, and the finite-height transfinite-word report was integrated with its guarded-block companion. Both proof routes and all distinct extensions are retained; old directories contain redirects, not extra catalogue entries. The seven surreal-number reports that this catalogue used to carry were moved out to a repository of their own, with their history, and are no longer listed here. They arrived as one hundred and twenty-five ZIP archives, in seven deliveries, which have been unpacked and grouped into the subject hierarchy described in Section 2. The archives themselves are no longer kept in the repository.

One hundred and twenty-five archives yield one hundred and three reports for two different reasons. Twenty archives duplicated another: sixteen pairs, three three-way clusters and one four-way cluster. Two more were folded into a neighbour not because they repeated it but because they shared a subject — the three reports on term counts in the derivatives of power towers are now one article with a common framework and three parts. That consolidation is editorial, and the distinction matters: those three were put through the duplicate sweep and cleared it, and their three theorems are three different theorems. The seventh delivery of eighteen contained no duplicate at all, which is worth recording: a sweep that finds nothing is evidence about the collection, not a sweep that failed. Each pair has been merged into one report that proves the shared theorem once and keeps everything both originals built on top of it. Where the two reached a result by genuinely different routes, both proofs are kept and marked as alternatives, with a sentence on what each one buys; the merged shuffle report keeps three independent finite certifications of one theorem, and the merged report on surreal option graphs kept two different five-vertex counterexamples for the same reason before it moved out with the other surreal-number reports. A merged entry below names both of its source archives and says what was doubled.

Not every pair aimed at one problem is a duplicate. The two Gonshor reports both prove his product-birthday bound, but on domains that are incomparable in both directions, so neither theorem contains the other and they are catalogued separately; the same holds for reports answering different numbered problems of one paper, noted where it occurs.

Every package is self-contained and follows the same shape: a typeset article (`*.pdf`) with its complete L^AT_EX source, a `README.md` summarising the result, usually a `code/` directory with exact-arithmetic verification scripts, a `data/` or `results/` directory holding the recorded output of those scripts, and often `STATUS.md`, `sources.md` or `notes/` files auditing the sources consulted and the precise limits of the claim.

Two cautions apply uniformly and are therefore not repeated in the individual entries. First, these are AI-assisted research drafts: none has been independently refereed, and none has been checked in a proof assistant. Several continue a Lean or Rocq project elsewhere in ProveIt; that confers no formal status on them, and their entries say which project they continue. Second, the descriptions below record *what each report sets out to settle and what it claims to prove*. Compiling this manifest involved reading each package’s statement of its problem and result; it did not involve checking any proof. Where a report itself flags a limitation — a prior disproof, an imported theorem, a correction to the source, or an exception left open — that is noted, because it changes what the report is.

The naming of the directories is editorial. Several distinct reports attack the same published problem, and some arrived under colliding archive names, so each directory has been given a name describing its own angle on the problem; the merged directories usually name their combined scope. The September 29 consolidations retain the base directory names to preserve existing links. The original archive names are recorded with every entry.

2 The directory tree

Category	Subcategory	Reports
enumerative-combinatorics	—	25
ordinals-and-order-types	ordinal-arithmetic	3
	wqo-powersets-and-statures	4
	friendly-order-types	3
	games-on-ordinals	3
	transfinite-words	1
	(top level)	6
generating-functions-and-asymptotics	oeis-sequence-asymptotics	59
	binary-product-coefficient-counts	3
	(top level)	5
log-concavity-and-unimodality	—	14
hankel-determinants	catalan-and-ballot	4
	somos-and-elliptic	3
	growth-and-runs	2
	(top level)	1
congruences-and-valuations	supercongruences	4
	iterated-series	3
	(top level)	4
automata-and-formal-languages	—	8
tetration-and-digit-stabilization	—	6
graph-theory	—	3
quaternionic-analysis	—	1
jacobian-conjecture	—	4
galois-theory-and-radicals	—	2
hilbert-tenth-problem	—	9
Total		180

More than a quarter of the collection consists of counterexamples and corrections rather than proofs. Ordinal Chomp, open-query games, compositional tree series, the secant numbers, the decimal-length bound for tetration stabilization, power-tower stabilization, iterated-power digit stabilization, cluster variables, chromatic coefficients of cycles, products of q -integers, domination roots, Hardinian array constants, the mixed-base counterexample, recursive exponents, Gregory sign thresholds and binary substitution discrepancy all refute, or replace with a proved statement, an assertion published or catalogued elsewhere. Several reports whose main outcome is a proof still correct the statement they prove: the Somos-8 certificates, the shifted Catalan Hankel polynomials and the Rueppel determinants record a sign error, a missing one-unit shift and a numbering ambiguity in their sources, and the independence-system thresholds report refines its target after finding that the original conjecture had already been disproved by someone else.

3 Ordinals and order types

3.1 Friendly order types

Three reports on Isa Vialard’s *friendly order type* $f(P)$, the rank of the tree of open-ended bad sequences in which every chosen point has an incomparable partner in the current residual. All three prove the same finite formula; they differ in what they build on top of it, and in how they

prove it — one by a forest-rank monovariant that survives moves deleting a whole upset, one through a lemma bounding the cut maxima. A fourth report proved the formula by the first report’s argument almost verbatim and has been merged into it.

1. Friendly Order Types Through Incomparability Components

ordinals-and-order-types/friendly-order-types/
incomparability-components-and-infinite-tails/friendly_order_types.pdf
from friendly_order_type_research (1).zip

The same component formula, together with an exact friendly rank for connected infinite well-partial-orders: writing $o(C) = \lambda + n$ with λ the limit part and $T(C)$ the finite set of points with finite principal upsets. Includes an exhaustive finite verification and a proof audit.

2. Friendly Order Type Through Incomparability Components

ordinals-and-order-types/friendly-order-types/
matroid-rank-and-ordinal-sums/article.pdf
from friendly_order_type_research (3).zip

Identifies $|P| - c(P)$ with the graphic-matroid rank and the incidence-matrix rank of the incomparability graph. Adds a substitution formula, a finite Cartesian-product formula, and additivity over arbitrary well-ordered ordinal sums; evaluates every well-partial-order whose incomparability components are finite, and gives countable counterexamples delimiting stronger infinite extrapolations.

3. Friendly Order Types of Finite Posets and Ordinal Products

ordinals-and-order-types/friendly-order-types/
components-forests-and-ordinal-products/friendly_order_types.pdf
from friendly_order_types_research.zip + friendly_order_type_research.zip

Proves $f(P) = |P| - c(\text{Inc}(P))$ for a finite poset P , where c counts connected components of the incomparability graph (isolated vertices included), and classifies the witnesses: every maximum friendly subset is P minus one minimal root per component, counted by $\prod |\text{Min}(C_i)|$. With Vialard’s width-transfer theorem this gives ordinal width $\omega^{f(P)}$ for the Dershowitz–Manna multiset ordering. On the transfinite side it proves $f(\alpha_1 \times \cdots \times \alpha_r \times B) = \Theta$, the natural product, except that it is the predecessor of Θ when every α_i is a successor and B has a greatest element, and classifies arbitrary finite ordinal boxes. Also: spanning-forest certificates, exact monotonicity under added comparisons, a bivariate generating function, and explicit four-point factors with equal (o, h, w, f) whose products differ. *Merged; the Cartesian-product formula is proved twice, once by corner deletion and once by a chain-grid spanning-subgraph argument.*

3.2 Well-quasi-ordered powersets and statures

Four reports on the open questions in Section 6 of Abriola, Halfon, Lopez, Schmitz, Schnoebelen and Vialard, *Measuring well-quasi-ordered finitary powersets* (arXiv:2312.14587v2), and on the corresponding topological invariant.

4. Cofinal strata and ordinal absorption

ordinals-and-order-types/wqo-powersets-and-statures/
cofinal-strata-of-finitary-powersets/article.pdf
from cofinal_strata_ordinal_powersets.zip; batch 42: ProveIt_Ordinal_Heights_Matchings.zip;
batch 36: ProveIt_Nonuniform_Powerset_Heights.zip, ProveIt_Persistent_Frontier_Heights.zip,
persistent_frontier_heights.zip, ProveIt_Ordinal_Heights_Research.zip

For $S_\alpha(Q)$, a finite poset Q with each point replaced by a chain $\alpha = \omega^\rho$, proves $o(K(S_\alpha(Q))) = \sum_k \omega^{\rho \otimes k} c_k(Q)$ and $h(K(S_\alpha(Q))) = \alpha \cdot \text{height}(M(Q))$, where c_k counts size- k maximal antichains not dominated by a larger one. The maximal-order-type theorem allows nonuniform pure well-partial-order fibers and is proved by a finite cofinal-stratification argument. Parts II and III, merged in batch 36 from four manuscripts that each prove it, supply the nonuniform height formula the original left open. For positive pure fibers $\omega^{\rho q}$, $h(K)$ is the largest ordinary ordinal sum, along a chain of maximal frontiers, of the largest fiber leaving the frontier at each step

plus the largest fiber of the last frontier; after Cantor-normal-form refinement the same formula covers nonzero limit fibers. For arbitrary ordinal fibers, finite and successor ones included, the height is given instead by a finite dynamic program over downsets with fallback weight 1, and the frontier compression is not valid there. Adds exact point ranks, NP membership of the threshold problem for the explicit pure-block encoding, and a separate part for well-partial-order fibers with $h(K(P_q))$ a positive pure power. Part IV, added in batch 42, shows that for height-two skeletons with uniform fibers the height is $\alpha(\nu_{\text{ur}} + 1)$, with ν_{ur} the maximum uniquely restricted matching number, so deciding the height is NP-complete already when every fiber is ω .

5. Intrinsic ordinal approximations for finitary powersets

ordinals-and-order-types/wqo-powersets-and-statures/
intrinsic-ordinal-approximations/article.pdf
from intrinsic_ordinal_approximations.zip

Studies the intrinsic invariant $a_o(A) = \sup\{o(B) + 1 : \text{Idl}(B) \hookrightarrow A\}$. Proves the universal bound $a_o(A) \leq h(P_f(A))$ with equality exactly when every finitely generated downset of A is an algebraic dcpo, gives the closed formula $h(P_f(\prod_i \alpha_i)) = \sup(\bigotimes_i (1 + a_i) + 1)$, and shows the universal identity fails on $\lambda + F$.

6. Ordinal Order Maps and Finitary Powersets of Ordinal–Finite Grids

ordinals-and-order-types/wqo-powersets-and-statures/
ordinal-order-maps-and-grids/ordinal_order_maps.pdf
from ordinal_order_maps_research.zip

Exact maximal order types and heights for finite Hoare powersets of $\beta \times Q$ with Q a finite poset: for $\beta = \omega^{\gamma_0} + \dots + \omega^{\gamma_{m-1}}$, $o(\text{Mon}(Q, \beta)) = \bigoplus_{s \text{ isotone}} \omega^{\bigoplus_q \gamma_s(q)}$.

7. Sharp ordinal bounds for Hoare powerspaces

ordinals-and-order-types/wqo-powersets-and-statures/
hoare-powerspace-statures/article.pdf
from hoare_statures_research.zip

For all Noetherian spaces, $1 + \|X\| \leq \|HX\| \leq 2^{\|X\|}$, with witnesses attaining both endpoints at every ordinal and T_1 upper witnesses; the older ω^α bound is attainable at infinite input stature α exactly when $\omega \cdot \alpha = \alpha$; the complete spectrum at $\omega + n$ is $\omega \cdot j(U) + (j(Q) - j(U))$. Extends the known well-quasi-order results of Abriola et al. to arbitrary Noetherian spaces.

3.3 Transfinite words

One unified report on Harry Altman’s finite-alphabet sharpness question (arXiv:2409.03199v2, Example 3.15), combining three source manuscripts. The common height-two formula is stated once. The finite-height induction and the more detailed guarded-tail investigation are retained as distinct routes.

8. Finite-alphabet transfinite words: exact order types and guarded periodic blocks

ordinals-and-order-types/transfinite-words/order-types-below-omega-squared/article.pdf
from omega2_order_types_proposed_solution.zip + ordinal_words_research.zip +
finite_alphabet_transfinite_words.zip

Claims $o(V_k(P)) = \omega^{\omega^{N_k(P)-1}}$ for $|P| \geq 2$ and each positive finite k , using canonical finite atom posets with $N_{k+1} = |P| + J(I_k(P)) - 1$. At height two this is $\omega^{\omega^{n+j(P)-2}}$; empty and singleton alphabets are explicit exceptions. Retains the selected-tail-family classification, its universal-only exception, chain/antichain extremality, exact-length strata, and canonical linear-time token comparison. The guarded proof has both natural-product and elementary amplification routes; the finite-height proof uses protected separators. Two independent implementations and their separate audit records are preserved. Neither an all-height union formula nor a higher-cutoff selected-family classification is claimed.

3.4 Ordinal arithmetic

Three reports on ordinal and infinitary arithmetic, including a consolidation of two independent proposed solutions to Problem 6.2 of Paolo Lipparini, *Monotone infinitary operations on ordinals* (arXiv:2505.00424v2): define and study the least weakly monotone, e -special strictly increasing operation on countable sequences of ordinals.

9. An Explicit Formula for the Minimal e -Monotone Infinitary Ordinal Sum

[ordinals-and-order-types/ordinal-arithmetic/lipparini-minimal-infinitary-sum/article/explicit_ordinal_sum.pdf](#)

from `lipparini_problem_6_2_proposed_solution.zip` + `ordinal_operation_research.zip`

The same three-case formula for the least admissible operation is proved by two retained routes: profile ranks with arithmetic absorption and finite-product lower bounds, and corrected blocks with threshold interpolation and finite-part restoration. The latter avoids the product-rank lemma; both import Lipparini's characterization of the weaker S . Retains the multiset quotient, bounded-poset heights, finite-cap ranks with alternative proofs, equality on comparable inputs, cardinality preservation, and counterexamples to grouping and continuity. The correction is zero, a finite increment, or one *natural* summand ω . Both implementations and their separate verification records are preserved.

10. Jacobsthal Products All at Once

[ordinals-and-order-types/ordinal-arithmetic/jacobsthal-products-all-at-once/article.pdf](#)

from `Jacobsthal_Products_All_at_Once.zip`

Question 2.16 of Altman, *Intermediate arithmetic operations on ordinal numbers* (MLQ 63 (2017) 228–242), asks for an all-at-once order-theoretic model of finite and infinitary Jacobsthal products, rather than one built from parenthesized binary multiplication or a limit. Gives a constructive affirmative answer: a Cantor-normal-form comparison rule on actual tuples or finite-support functions, invariant under order isomorphisms and extending the componentwise order, whose definition never evaluates a Jacobsthal product. Adds a restricted maximality characterisation and a finite-tail reduction. The associativity identity and the binary normal-form formula are classical and are not claimed here; the construction is not claimed to be preserved by arbitrary order embeddings.

11. Pairwise Coherence Does Not Guarantee Completion

[ordinals-and-order-types/ordinal-arithmetic/](#)

[pairwise-coherence-of-infinitary-products/ordinal_product_coherence.pdf](#)

from `ordinal_product_coherence.zip`

Answers the pairwise-coherence sufficiency question of Lipparini, *Noncommutative infinitary semigroups* (arXiv:2605.28413v1, Remark 7.1(b)), negatively. On the two-element group C_2 , retaining every finite product and declaring every infinite all- e product to be e gives a partial system satisfying full pairwise regrouping, even for arbitrary partitions, yet with no injective extension defining all finite and ω -indexed products; two elements is minimal. Adds: for any monoid M the canonical partial domain $P(M)$ is coherent, and for directly finite M it completes exactly when the unit group is trivial; and for every r a system on $3r + 4$ elements coherent through nesting depth r but conflicting at depth $r + 1$, so no fixed finite depth of the coherence test characterises completability.

3.5 Games on ordinals

12. An ordinal Chomp transition at 2

[ordinals-and-order-types/games-on-ordinals/](#)

[ordinal-chomp-transition-at-two/ordinal_chomp_counterexample.pdf](#)

from `ordinal_chomp_transition_counterexample.zip` + `ordinal_chomp_counterexample.zip`

For a numerical semigroup S ordered by $x \leq_S y$ iff $y - x \in S$, and the ordinal extension ordered by ordinary ordinal subtraction (so S^2 is the *lexicographic*, not coordinatewise, square), Chomp on S

is a second-player win while Chomp on S^σ is a first-player win for every $\sigma \geq 2$, and $\omega \cdot 4 + 4$ is the *least* winning opening. Hence $\text{ch}(\langle 4, 5, 6 \rangle) = \text{ch}(\langle 4, 6, 9 \rangle) = 2$, a negative answer to Question 4.6 of Rivero Herrera (arXiv:2504.07317v3) and a refutation of Conjecture 4.1 of v1, already for three small natural-number generators. The proof reduces the infinite game to a finite spectral statement via a finite-window recurrence and an exact finite-state certificate, then transfers it by an explicit response procedure. *Merged; the shared $\langle 4, 6, 9 \rangle$ certificates were verified to hold the same 253 rows entry for entry, and the two reports' poisoned and unpoisoned conventions are reconciled by an explicit dictionary.*

13. Every ordinal as a point-separating game value

[ordinals-and-order-types/games-on-ordinals/
point-separating-game-values/ordinal_separation.pdf](#)
from ordinal_separation_research.zip

For every ordinal λ , orders $\{0, 1\}^\lambda$ lexicographically and gives it the topology of all downward-closed sets, and the refinement generated by downward-closed sets minus finite sets. The point-separating game value is exactly λ for both: the first space is compact T_0 , the second compact T_1 , and for infinite λ neither is Hausdorff. Taking $\lambda = \omega^2 + 1$ supplies a compact T_1 witness of cardinality continuum for Problem 1.2 of Chiozini, Csernák and Soukup (arXiv:2510.05754v3), with no CH, forcing or large-cardinal hypothesis. Hausdorff witnesses are not claimed. This is a different problem of that paper from the one the open-query games report answers.

14. Open-query games on ordinal spaces

[ordinals-and-order-types/games-on-ordinals/open-query-membership-games/article.pdf](#)
from ordinal_membership_games.zip; batch 52: ProveIt_Finite_Output_Open_Query_Games.zip;
batch 56: ProveIt_Filter_Profiles_Open_Query_Complexity.zip; batch 65:
ProveIt_Closure_Word_Duality.zip; batch 68: ProveIt_Universal_Word_Profiles.zip

A negative answer to Problem 1.3 of Chiozini, Csernák and Soukup (arXiv:2510.05754v3): two copies of $[0, \omega]$ each have set-membership number 1, but their topological sum has value 2. The general theorem computes the invariant for every nonempty Hausdorff space of finite Cantor–Bendixson height h as $\lceil \log_2(h + 1) \rceil$ when the last nonempty derivative is a singleton and $\lceil \log_2(h + 2) \rceil$ otherwise. Part II, added in batch 52, replaces the membership bit by one of q output labels: the optimal depth is $\lceil \log_2 L \rceil$ for the least number L of ordered layers with open suffixes; on Hausdorff spaces of finite height $L \leq (q - 1)h + s$, sharply, with s the number of labels on the last derivative; a topological sum of components of depth at most d has depth at most $d + \lceil \log_2 q \rceil$, sharply; and computing the optimal depth is polynomial for two labels but NP-complete for three, even on disjoint chains. At $q = 2$ it reproduces Part I. Part III, added in batch 56, answers Part II's question of when that bound is attained, for spaces whose derived set is finite and nonempty: a coloring with q_c colors needs exactly $q_c + \tau(G_c)$ layers, τ the directed feedback vertex number of a color digraph; the bound is attained exactly under a bin-partition condition on the partition capacities of the neighborhood filters; and at three labels a sum of three one-point free-ultrafilter spaces has value 2 while a sum of three convergent sequences has value 3. Part IV, added in batch 65, drops the separation and finiteness hypotheses for a fixed coloring: a layer word is admissible exactly when it contains as subsequences all reduced words whose nested color closure is nonempty. It classifies the finite-depth behaviors, shows that the longest such word has the length of the least Cantor–Bendixson height of a Hausdorff representative, gives $q_c + \tau(G_c)$ layers whenever the second derivative is empty, bounds finite-poset obstructions by $mq(q - 1)^{m-1}$ points, and proves the product law $\rho(X \times Y) = \rho(X) + \rho(Y) - 1$ for that longest length ρ . Part V, added in batch 68, computes the worst case over colorings at every finite height on locally compact or sequential (more generally rank-accessible) Hausdorff spaces: $\max_c L = h(q - 1) + \min(q, t)$ with t the size of the last nonempty derivative, also when t is infinite, attained by one coloring constant off a countable set; it adds product and sum laws and an inverse theory of layer and query-depth spectra. Local data determining an arbitrary fixed coloring at height two, and a characterization of the spaces attaining that worst case (rank accessibility is sufficient, not necessary), remain open.

3.6 Order-theoretic invariants without a subcategory

Six reports whose objects sit outside every subcategory above: the minor order on finite graphs, two on sites of filters on ω , congruences of finite lattices, a translation-invariant ideal, and the lexicographic order of the well-orderings of the reals.

15. Ordinal complexity of graph minors

[ordinals-and-order-types/graph-minor-ordinals/graph_minor_ordinals.pdf](#)

from [graph_minor_ordinals.zip](#)

For finite simple graphs of maximum degree at most two under minor containment, computes maximal order type, ordinal antichain-tree width and descending-tree height: $o(D) = w(D) = \omega^{\omega \cdot 2}$ and $h(D) = \omega$, with exact values ω^{b+k} and ω^{b+k-1} for the subfamilies bounded by path size b and cyclic-component count k . The general theorem is that a wqo C of connected graphs with $o(C) = \alpha < \varepsilon_0$ has finite disjoint-union closure of type ω^α , proved by a component-splitting argument, since minor containment is not multiset embedding. A corollary is that continuity fails for increasing unions of minor-closed classes. The report is explicit that it does not compute the maximal order type of the minor order on all finite graphs and settles no named conjecture.

16. A two-valued obstruction to gluing Fréchet powers

[ordinals-and-order-types/frechet-two-valued-obstruction/article.pdf](#)

from [frechet_two_valued_obstruction.zip](#)

For the presheaf $F \mapsto A^\omega / \sim_F$ on proper filters on ω containing the cofinite filter, with the dense Grothendieck topology, proves in ZF that it is a sheaf if and only if $|A| \leq 1$. The counterexample is an explicit matching family of constant sections using two values, on a covering sieve with countably many generators each of countable base; the distinction between the number of sieve generators and the size of the full sieve is essential to it. The question is Massas, *A Semi-Constructive Approach to the Hyperreal Line* (2023), p. 517 after Fact 5.10, repeated in the 2024 dissertation *Duality and Infinity*, p. 249.

17. Bounded-character Fréchet filter sites

[ordinals-and-order-types/frechet-bounded-character-sites/article.pdf](#)

from [filter_complexity_research.zip](#), [Frechet_Cardinal_Boundaries.zip](#)

Continues the previous entry's non-claim about other filter sites. For the site P_κ of free filters on ω generated by fewer than κ sets modulo the cofinite filter, represents the Booleanization C_κ as a regular-open algebra on the free ultrafilters and proves that $P(\omega)/\text{Fin}$ is order dense in it exactly for $\kappa \leq \mathfrak{p}$; above $\mathfrak{p}$ no complete Boolean homomorphism preserving the modulo-finite events exists in either direction. The atoms are the ultrafilters of character below κ , so C_κ is atomless exactly for $\kappa \leq \mathfrak{u}$; $\mathfrak{h}$ is the first size of a family of events that cannot be decided jointly on a dense set of regions. Computes the associated sheaf by Boolean mixtures, with full Boolean-valued transfer, countable saturation of ultrafilter specializations, and sizes $\mathfrak{c}$, $2^\mathfrak{c}$, $2^{2^\mathfrak{c}}$ for the raw power and the two sheafifications. A three-stage example uses Millán's forcing model as an imported input. Merged from two manuscripts whose indexings differ by a successor shift; the second one's closure-based results are stated at successor stages only.

18. A sharp cover bound for congruences of finite lattices

[ordinals-and-order-types/lattice-congruence-cover-bound/sharp_cover_bound.pdf](#)

from [lattice_cover_bound_research.zip](#)

If a finite n -element lattice has an element with at least $k \geq 3$ upper covers then $|\text{Con}(L)| \leq 2^{n-k-1}$, with equality exactly for the glued sums $C_s \oplus M_k \oplus C_t$. This replaces the multicolour Ramsey threshold $R_{k-1}(k)$ of the source's Lemma 4.4 (arXiv:2603.11454) by k itself. Adds a sharp nonextremal gap, a fan-label ideal refinement, and a bound on the size of Czédli's skeleton in terms of congruence density. All congruence machinery, including Freese's maximum, is proved from scratch. The target is an informal suggestion in the source, not a numbered conjecture.

19. A non-Baire translation-invariant ideal

[ordinals-and-order-types/non-baire-translation-invariant-ideal/article.pdf](#)

from `non_baire_ideal_research.zip`

An affirmative construction for Question 1 of Filipów and Tryba, *Densities for sets of natural numbers vanishing on a given family* (J. Number Theory 211 (2020) 371–382; arXiv:2309.00982, p. 7): a translation-invariant ideal that is not Baire and carries a perfect translation-almost-disjoint family.

20. Lexicographic orders of the well-orderings of the reals

[ordinals-and-order-types/lexicographic-well-orderings-of-reals/article.pdf](#)

from batch 76: `lexicographic_well_orderings_of_reals.zip` + `Lexicographic_Well_Orderings_of_Reals (1).zip` + `lexicographic_well_orderings_research.zip`

With $\kappa = 2^{\aleph_0}$ and $\mathcal{W}$ the order of all well-orderings of $\mathbb{R}$, as enumerations compared at their first unequal real entry, proves in ZFC that $|\mathcal{W}| = 2^\kappa$, that every linear order of size at most κ embeds, even into each fixed-type slice, and that an ordinal embeds exactly when it is below κ^+ ; yet the binary lexicographic coding length and the real representation rank of $\mathcal{W}$ are both κ^+ , so $\mathcal{W}$ embeds into no bounded union of strata, in particular not into the slice $\mathcal{W}_\kappa$, which is equimorphic with the cube 2^κ in lexicographic order and with the Kanovei–Shelah ultrafilter index. Also: each stratum $\mathcal{W}_{\lambda+n}$, λ a limit of cardinality κ , is equimorphic with $2^\lambda \times n!$; $\mathcal{W}_{\text{lex}}^\gamma$ is equimorphic with $\mathcal{W}$ exactly for $0 < \gamma < \kappa^+$; adjacency and the $n!$ blocks; the exact point-character and gap spectra (every gap has a countable side; exactly 2^κ gaps), a (cf κ , cf κ) gap of $\mathcal{W}_\kappa$, a dense finite-condensation quotient equimorphic with $\mathcal{W}$, bounded surreal sign orders and an ultrafilter index built from well-orderings. The cube hierarchy re-proves a corollary of [point-separating-game-values](#), cited first; real-power strictness is Kuhlmann’s theorem; no priority is claimed, and no intrinsic embedding criterion or classification of the gaps of the longer strata. *Merged from three independent manuscripts proving the same core: each shared theorem printed once, with the other arguments as second routes. Each answers a question of another: the ZFC non-embedding into $\mathcal{W}_\kappa$, which one had only under $2^{<\kappa} < 2^\kappa$, and the cut and character spectra and the quotient another had asked about.*

4 Automata and formal languages

Eight reports on state complexity, synchronization, language hierarchies, crossover closures, additive complexity and the asymptotic enumeration of automata. Three are merges: the shuffle bound arrived four times, the DFAO reversal bound twice, and the last report joins a manuscript on accessible automata with one on strongly connected automata that adapts its method.

21. The six-state shuffle bound

[automata-and-formal-languages/shuffle-six-state-bound/shuffle_six_state.pdf](#)

from `shuffle_six_state_research (1).zip` + `shuffle_six_state_research.zip` + `Shuffle_Six_State_Theorem.zip` + `six_state_shuffle_research.zip`

For complete DFAs with $m, n \geq 2$ and $\min(m, n) \leq 6$, the worst-case deterministic state complexity of the shuffle is $F(m, n) = 2^{mn-1} + 2^{(m-1)(n-1)}(2^{m-1} - 1)(2^{n-1} - 1)$, over a finite alphabet allowed to depend on m and n ; the unrestricted conjecture stays open. The route is a reachability theorem in the universal shuffle automaton — a subset is reachable exactly when it meets the initial row and the initial column — proved by induction on $(m + n, |S|)$ down to a finite certificate obligation. *Merged from four reports. Two shipped the same 10,642 witnesses; a third supplies an independent database of 10,619 witnesses sharing no witness with them, so the finite step is certified twice over; a fourth adds a fifth analytic reduction that cuts the obligation to 453 certificates and replaces the solver search with a deterministic enumeration. All three routes are kept.*

22. A coloring obstruction to maximal binary DFAO reversal

[automata-and-formal-languages/dfao-reversal-coloring-obstruction/article.pdf](#)

from dfao_reversal_research.zip + binary_dfao_reversal_research.zip; batch 39:

ProveIt_Three_Output_Reversal.zip; batches 40–41: ProveIt_DFAO_Eventual_Optimality.zip,

ProveIt_Binary_DFAO_Reversal.zip, ProveIt_Binary_Reversal_Exactness.zip; batch 43:

ProveIt_Four_Output_Reversal.zip, ProveIt_Four_Output_Reversal (1).zip; batch 67:

ProveIt_Near_Linear_DFAO_Reversal.zip, ProveIt_DFAO_Spectral_Cancellation.zip

For transformations a, b on an n -element set and an output map to k labels, proves $|\tau\langle a, b \rangle| \leq k^n - k! + g(k) < k^n$ for $3 \leq k \leq n$, with $g(k)$ the largest order of a permutation of k symbols — the binary-alphabet nonattainment question of Davies, arXiv:1705.07150v2, Section 5. The mechanism is a collision-orbit graph, a rightmost singular factorization, and a cyclic relabeling bound. Adds an abelian-permutation extension, a structural universal bound matching the published lower construction on all 372 pairs with $7 \leq n \leq 30$, and polynomial-time missing-coloring certificates. *Merged; the two originals ran the same three-case argument, and both a qualitative short proof and the counting proof are kept.* Part II, added in batch 39, settles three outputs exactly: $R_2(n, 3) = 3^n - 3(2^A + 2^B - 2) + B$ for every $n \geq 7$, with (A, B) the closest coprime split of n , together with rigidity of the extremizers; the formula matches all twenty-four $k = 3$ rows of the finite table. Part III, merged in batches 40–41 from three manuscripts, settles every $k \geq 4$ past an explicit threshold: $R_2(n, k) = k^n - \text{Bip}_k(a_n, b_n) + a_n b_n$ once $n \geq N(k) \leq 3k^4$, and for all $n \geq \max(7, k + 1)$ when $k = 4, \dots, 7$ (computer-assisted below $N(k)$), with the lower construction imported from Davies; a non-effective eventual route for every $k \geq 3$, 919 finite certificates and a second proof of the $k = 3$ formula are kept. Explicit exactness for $k \geq 8$ below $\min(N(k), T(k))$, with $T(k)$ the threshold of Part V, stays open. Part IV, merged in batch 43 from two manuscripts on $k = 4$, re-derives that case with an analytic threshold $n \geq 26$ (against 212), and adds stability, the missing-orbit inventory, the extremal output maps and a minimal order-15 recurrence. Part V, added in batch 67, proves the formula for every $k \geq 4$ once $n \geq T(k) = 4k \log k + O(k)$ instead of $n \geq 3k^4$ (so for all k up to $cn/\log n$, $c < 1/4$), with a theta-function limit of the optimal deficit when n/k^2 tends to a constant and a Gaussian law in between. Part VI, added in batch 67, determines the minimal recurrences of the explicit witness deficit for every fixed k : order 9 at three outputs (against Part II’s 12), a product-12 cancellation on the odd residues but not in the full sequence at nineteen outputs, the only product-12 cancellation at any k , and explicit tail numerators.

23. Synchronizing Dzyga’s automata by a cyclic retract

[automata-and-formal-languages/dzyga-synchronizing-retract/article.pdf](#)

from Dzyga_Automata_Cyclic_Retract_Research.zip

For the Figure-3-complete automaton A_k of Conjecture 8 in Szykuła’s 2026 survey, exhibits an explicit word $R_k = p(ec^{r-1})^{r-2}e$ with $r = k + 4$ that sends every state to q_2 for every $k \geq 1$, of expanded length $8k^2 + 54k + 92 = \frac{8}{9}n^2 + \frac{22}{3}n + 16$ for $n = 3k + 6$. The proof is a cyclic retraction with two projected reflections whose discrepancy supplies one adjacent merger, and does not depend on the finite computations; shortening the reflected tail by one state gives a nonsynchronizing automaton with a dihedral permutation quotient. This settles the synchronization clause of the stated model, not the numerical threshold conjecture, and the universal lower bound $\tau(q_0) \geq 4k + 8$ is not proved.

24. Gaps in bounded-shuffle hierarchies

[automata-and-formal-languages/insertion-degree-spectra/article.pdf](#)

from shuffle_spectra_research.zip; batch 41: ProveIt_Binary_Insertion_Spectra.zip; batch 66:

ProveIt_Sharp_Insertion_Isolation.zip

A counterexample to the insertion-degree No-Gap Conjecture of Hughes, arXiv:2608.27755v1, Section 11, followed by positive results: every finite set of positive integers containing 1 is the insertion-degree spectrum of two nonempty finite languages over a suitable alphabet, and any prescribed finite list of degrees above 1 is realizable with every other output of degree 1. A ternary pair has spectrum $\{1, 3\}$ with sole difficult output **abcba**, whose length 5 is the minimum possible

for a gap witness; a binary pair has spectrum $\{1, 2, 4\}$ while all 18 singleton source-pair spectra are initial intervals, so source-pair ambiguity can erase a degree. The separate iteration-depth no-gap conjecture is proved for arbitrary languages. Part II, added in batch 41, fixes the alphabet: every finite set containing 1 is the spectrum of finite *binary* languages with the single inserted word a^m , so two letters always suffice; its m^2 length bound holds in its full-coverage class only. Part III, added in batch 66, determines that class exactly: an isolated degree- m output with g gaps between its a 's (the two end gaps included) and T letters a exists exactly when $g \geq m$ and $T \geq \max(g(m-1) + 1, \tau_m)$, with $\tau_3 = 12$ and $\tau_m = 5m - 2$ for $m \geq 4$, so the minimum lengths for $m = 3, 4, 5$ are 14, 21 and 27. It characterizes the extremal centers and the mandatory targets, and at $m = 3$ classifies the shortest target languages: $3^9 = 19,683$ of them, of minimum size 37, attained by 512.

25. Constrained crossover closure

[automata-and-formal-languages/constrained-crossover-closure/article.pdf](#)

from ProveIt_Constrained_Crossover.zip (batch 44); batch 66: ProveIt_Crossover_Stabilization.zip, ProveIt_Crossover_Classification.zip; batch 71: quadratic-finite-rank-package.zip; batch 73: rational-rank-slopes-package.zip, dfa-crossover-package.zip, common-path-rank-package.zip (the earlier editions dfa-crossover-package (1).zip and common-path-rank-package (1).zip superseded, not printed)

Answers the question Hughes (arXiv:2608.27755v1, Section 4) leaves open on the complexity of deciding $L \otimes_c L = L$ for the constrained crossover of a context-free language: it is co-r.e.-complete. Adds an aligned-rank no-gap theorem for iterated crossover and effectivity results in which Presburger quantifier elimination is used only as an algorithmic step. Related to, but using nothing from, the insertion-degree report above. Part II, added in batch 66, shows finite stabilization of regular seeds PSPACE-complete for NFA input, already over two letters; the upper bound also covers DFA input, improving Part I's EXPSPACE bound (Part VI settles DFA input); in the stabilizing case the rank is at most quadratic in the number of states. Part III, added in batch 66, decides regularity, inclusion and equality of the full closures of context-free seeds, classifies the finitely stabilizing sparse one-marker seeds into regular, linear non-regular and non-context-free closures, and shows that thresholding the exact stabilization depth is coNP-complete for succinct ray input. Part IV, added in batch 71, shows that the quadratic order is attained for NFAs: the largest finite rank $F(s)$ of a binary language recognized by an s -state NFA satisfies $(s-3)^2 \leq F(s) \leq (2s+5)^2$ for $s \geq 4$, the lower bound replacing Part II's linear one. Parts V–VII, added in batch 73, answer the questions Parts II and IV left. Part V: on every live residue the largest rank of a length- n hull word is $\sigma_r n + O(1)$ with a rational, computable slope σ_r , and $F(s) \leq 2(s-1)^2 + 3$. Part VI: finite stabilization of an explicit complete binary DFA is coNP-complete, and the largest finite rank of an s -state complete binary DFA satisfies $(s-7)^2 + 2 \leq F_{\text{DFA}}(s) \leq s^2 + 3$ for $s \geq 8$. Part VII: $F(s) \leq s^2 + 8s$ for binary NFAs ($s^2 + \max(8, 2q-4)s$ over q letters), so $F(s)/s^2 \rightarrow 1$ and $F_{\text{DFA}}(s)/s^2 \rightarrow 1$, with an exact algorithm and $F(1), \dots, F(4) = 1, 2, 5, 7$, the last three computer-assisted. The exact values beyond four states and the lower-order terms stay open. Lemmas the new Parts re-prove are cited, not reprinted, and two lemmas shared by Parts VI and VII are printed once.

26. A cut-sensitive attack on the first open MIX grammar case

[automata-and-formal-languages/mix-grammar-cut-sensitive/mix_grammar_report.pdf](#)

from mix_grammar_research.zip

The selected question — whether $L_3 = \text{MIX}$ for the canonical tuple grammars of Yuyama, *On the Kanazawa–Salvati Conjecture* (arXiv:2609.13871v1) — is *not* settled here, and the report says so at the front. What it does contain, with complete written proofs, is a cut-sensitive membership recurrence, a broad three-component positive family, an infinite family in $L_3 \setminus L_2$, and an exhaustive computation through length 21.

27. Natural densities of the Tribonacci word's additive complexities

[automata-and-formal-languages/tribonacci-additive-complexity/article.pdf](#)

from [tribonacci_additive_density_research.zip](#)

Answers Remark 19 of Popoli, Shallit and Stipulanti (arXiv:2410.02409). With β the Tribonacci constant, the additive complexity of the Tribonacci word takes values in $\{3, 4, 5\}$ with natural densities $d_3 = (21\beta^2 - 38\beta + 5)/22$, $d_4 = (-55\beta^2 + 72\beta + 67)/22$ and $d_5 = (17\beta^2 - 17\beta - 25)/11$, all positive and summing to 1, with a power-saving error $O(N^\theta)$, $\theta = \log(7/5)/\log \beta$, at *every* integer cutoff rather than only along Tribonacci cutoffs. The 76-state output automaton is reconstructed independently rather than copied, and certified by evaluating its degree-76 characteristic polynomial at the adjacency matrix.

28. Accessible and strongly connected deterministic automata, to every fixed order

[automata-and-formal-languages/accessible-and-strong-automata/article.pdf](#)

from batch 77: [accessible-automata-reproducibility.zip](#) + [strong-automata-reproducibility.zip](#)

For a fixed alphabet of $k \geq 2$ letters, with $t = k(1 - e^{-t})$, $\rho = e^{-t}$, $v = t/k$ and $c = 1 - k\rho$, Part I proves that the number a_n of rooted accessible complete deterministic transition structures on n states (OEIS A006689, A006690) satisfies $a_n/S(kn + 1, n) = \sum_{j \leq J} c_j n^{-j} + O(n^{-J-1})$ for every J , with $c_0 = c$, $c_1 = k\rho v/(2c)$ and the small-state counts entering first at c_k . The route is an exact first-failure renewal with two boundaries and a finite coefficient algorithm; c_1, c_2, c_3 are tabulated. Part II adapts the method, with a strict checkpoint, to strongly connected structures (A027834/A027835, A006691/A006692, A304312/A304313), transfers the expansion through the classical formal logarithm, confirms the identification that the A304312/A304313 records leave conjectural, and gives exact cyclic automorphism sectors with a uniform divisor-tail bound and a self-contained proof of Liskovets's existing unrooted formula. The leading equivalents (Korshunov, Bassino–Nicaud, Lebensztayn) are classical, and priority of the refinement is left unresolved: Korshunov's 1978 survey was not obtained. The Lambert- W inverses and threshold brackets are instances of the transseries volume. *Merged from two manuscripts; the passages Part II repeats are printed once, in Part I.* Minimal DFAs of finite languages, a different object, are counted in the report [a082161-airy-amplitudes](#).

5 Hankel determinants and continued fractions

5.1 Somos sequences and elliptic curves

29. A proof of Barry's Hankel–Somos-6 conjecture

[hankel-determinants/somos-and-elliptic/barry-somos-6-hankel-recurrence/article.pdf](#)

from [barry_somos6_research.zip](#)

Conjecture 7 of Paul Barry, arXiv:2211.12637: for the Hankel determinants H_N of a four-parameter Jacobi-type generating function, $H_N H_{N-6} = \alpha H_{N-1} H_{N-5} + \gamma H_{N-3}^2$ with $\alpha = t^2(r + 2)^2$ and $\gamma = t^3((s + t)(r + s + t + 2)^2 + t(r + 2)^3)$. Also proves a homogeneous four-parameter form and a seven-parameter polynomial Hankel theorem making exceptional specialisations rigorous; the verifier never divides by a determinant.

30. Finite Hankel Certificates for Three Somos-8 Conjectures

[hankel-determinants/somos-and-elliptic/somos-8-hankel-certificates/article.pdf](#)

from [somos_hankel_proofs.zip](#)

Conjectures 11, 12 and 13 of the same Barry paper, by importing Hone's genus-two Hankel theorem (arXiv:1907.05204v3, Thm. 5.4) and supplying a binomial translation, a polynomial-minor specialisation valid on the degenerate locus, and a rank-bound lemma that turns four symbolic rows into an all-index identity. Identifies a sign error in the source's Catalan expression for Conjecture 13.

31. Elliptic addition and Jacobi continued fractions

[hankel-determinants/somos-and-elliptic/elliptic-addition-and-jacobi-fractions/article.pdf](#)
from [elliptic_continued_fraction_research.zip](#)

A consistently indexed, non-torsion version of Barry's 2023 Conjecture 4, with the coordinate formulas of Conjecture 3: every continued-fraction tail is given explicitly and removing one layer is shown to be elliptic translation. Hypotheses: characteristic zero, $Y^2 + aXY + bY = X^3 + cX^2 + dX$ nonsingular, $b \neq 0$, and $P = (0, 0)$ of infinite order; order-three and order-four examples show the regular fraction genuinely needs the last one.

5.2 Catalan and ballot moments

32. Shifted Catalan Hankel Polynomials

[hankel-determinants/catalan-and-ballot/shifted-catalan-hankel-polynomials/article.pdf](#)
from [catalan_hankel_conjecture_solution.zip](#); batch 39: [ProveIt_Catalan_Root_Collisions.zip](#);
batch 42: [ProveIt_Cyclotomic_Catalan_Recurrences.zip](#); batch 53:
[Two_Endpoint_Catalan_Confluence.zip](#); batch 55: [ProveIt_Interior_Collision_Research.zip](#); batch 57:
[ProveIt_Catalan_Growing_Shifts.zip](#)

For $D_N^{(m)}(a, b) = \det(aC_{i+j+m} + bC_{i+j+m+1})$, proves the denominator-exponent assertion on page 4 of Barry, arXiv:2011.10827v1, and shows the separately printed Conjecture 2 needs an explicit one-unit shift correction. The minimal characteristic polynomial is $(X^2 - (a+2b)X + b^2)^{m(m-1)/2+1}$, so the minimal order is $m(m-1) + 2$; all exceptional reduced denominators, the exact coefficient formula, numerator degree and reciprocal symmetry are supplied. Part II, added in batch 39, replaces $aC + bC'$ by an arbitrary polynomial multiplier and computes the root collisions: exact sector degrees with nonzero leading constants, the minimal recurrence order off resonance, a secondary-Hankel factorization and a certified root-free algorithm for the recurrence order. Part III, added in batch 42, gives every characteristic multiplicity for one repeated root at a cyclotomic resonance, and shows that the question it answers needed one more parameter: the sign of z^L , not only t and $\text{ord}(z^2)$. Part IV, added in batch 53, answers the endpoint half of Part II's question on uniform limits: for fixed-size clusters of roots approaching the endpoints 0 and 4 (d roots in all) the normalized determinant is a convergent series in even powers of $1/(N + d/2)$, locally uniform through every collision inside a cluster, with a factorized leading profile, an explicit first interaction term, and ratio to the fixed-endpoint determinant tending to 1 exactly when the outward displacements are $o(N^{-2})$. Part V, added in batch 55, answers the other half: for fixed-size clusters of roots colliding on the scale $1/N$ at fixed bases other than 0 and 4, the determinant is an exact finite sum of exponential sectors whose amplitudes are convergent series in $1/(N + d/2)$, uniform through every collision, with an explicit first correction (this expansion is not even), an explicit remainder bound, and factorized amplitudes for several separated clusters. For two equal-multiplicity roots inside the spectrum the limit is an oscillatory Jacobi Hankel determinant, the first correction oscillates as $\sin(2(N+t)\theta)$, and simple zeros move by a convergent phase-dependent expansion. Limits uniform up to the endpoints remain open. Part VI, added in batch 57, lets the shift grow with the determinant size: for real $a \geq 0$ the ratio $D_N^{(m)}(a, 1)/D_N^{(m)}(0, 1)$ tends to 1 exactly when $aN(N+m+1)/(4m+2) \rightarrow 0$, at any relative growth of N and m ; the size-biased inverse Jacobi zeros follow a shifted Marchenko–Pastur (Catalan-moment) law, and three critical scales, an algebraic free energy and a uniform saddle prefactor describe the crossover.

33. Product formulas for ballot-polynomial Hankel determinants

[hankel-determinants/catalan-and-ballot/ballot-polynomial-hankel-determinants/article.pdf](#)
from [Cigler_Hankel_Conjectures_13_15.zip](#)

Conjectures 13, 14 and 15 in Section 5 of Johann Cigler, arXiv:2111.14492v3. An explicit coefficient product gives the sign and exact support of the normalised shift- k Hankel determinant of the ballot moments and positivity of every residual coefficient for $n \geq 1$; the same products give the

conjectured generating-function denominators, exact numerator degrees and reciprocal numerator identities. Adds strict log-concavity of the residual coefficients and their limiting binomial profile.

34. Rectangular Schur polynomials and Cigler's Conjecture 8

[hankel-determinants/catalan-and-ballot/cigler-conjecture-8-schur/article.pdf](#)
 from [cigler_conjecture8_research.zip](#)

A complete proof of Conjecture 8 in Cigler, arXiv:2111.14492v3, Section 4 (printed pp. 16–17), for the Hankel determinants of $b_m(t) = \sum_r \binom{\lfloor m/2 \rfloor}{r} \binom{\lceil m/2 \rceil}{r} t^r$, including the numerator-polynomial degree and reciprocity assertions. A different conjecture of that paper from the ballot-polynomial entry (Conjectures 13–15) and from the parity entry (Conjecture 16).

35. A parity factorization for Cigler's Conjecture 16

[hankel-determinants/catalan-and-ballot/cigler-conjecture-16-parity/article.pdf](#)
 from [cigler_conjecture16_research.zip](#); batch 56: [ProveIt_Cigler_Schur_Recurrences.zip](#); batch 66: [ProveIt_Cyclotomic_Pole_Collapse.zip](#); batch 70: [ProveIt_Cigler_Conjecture17_Resolution.zip](#)

A complete proof of Conjecture 16 in Cigler, arXiv:2111.14492v3, Section 6 (pp. 21–22), on the Hankel polynomials, together with a sharp stabilization statement: the report also determines the first coefficient at which the conjectured stabilization fails. Part II, added in batch 56, proves the report's experimental rectangular-Schur formula: with $a = \lfloor k/2 \rfloor$ and $b = \lfloor (k-1)/2 \rfloor$ the normalized determinant is $s_{(n^a)}(1^a, t, (t^2)^b)$ for all $k \geq 1$ and $n \geq 0$. It determines the exact minimal recurrence at each shift, with reduced denominator $\prod_{q=0}^{k-1} (1 - t^q z)^{1 + \lfloor q(k-1-q)/2 \rfloor}$ over $\mathbb{Q}(t)$ and at every real $t \neq 0, \pm 1$, proves Cigler's denominators (84) and (86) minimal, and shows that the positive reciprocity sign printed in his (85), part of Conjecture 18, is false when $k = 2a$ with a even. Part III, added in batch 66, determines the reduced denominator and the minimal recurrence at every complex parameter: at a primitive root of unity of order $\ell \geq 3$ at most one residue class loses pole order, decided by an explicit parity test, and it loses one order except for a loss of three at primitive fourth roots when $k \equiv 1 \pmod{4}$. Part IV, added in batch 70, proves Cigler's Conjecture 17 in sign-corrected form: once his denominator $(1-t)^a(1-t^2)^{ab}$ is cleared, the determinant splits into k blocks, polynomials in t whose coefficients are integer-valued polynomials in n , of exact degrees $q(k-1-q)$ in t and $\lfloor q(k-1-q)/2 \rfloor$ in n , with explicit leading terms in n ; their reciprocity sign is -1 exactly when $4 \mid k$. The sign printed in Cigler's (83) fails for every $k \equiv 1 \pmod{4}$ from 5 on (his displayed block for $k = 5$ is palindromic), so the conjecture as printed is false. Its own README explains why this is a different problem from the other Cigler entries in this collection.

5.3 Arithmetic Hankel determinants

The numerator formula recorded as a conjecture in OEIS A122251, with A122249 as its binary special case. Two reports attacked it and proved it the same way, sharing even their worked counterexamples; they are merged here into one. The merged report sits at the category's top level.

36. A Proof of an OEIS Hankel-Numerator Conjecture

[hankel-determinants/a122251-numerators-and-denominators/report.pdf](#)
 from [hankel_numerator_research.zip](#) + [hankel_oeis_conjecture_proof.zip](#)

For the Hankel determinant

$$H_n(m, a) = \det(1/(a + m(i+j)))_{0 \leq i, j \leq n},$$

the reduced numerator is $\prod_{p|m} p^{n(n+1)v_p(m)+2 \sum_{k \leq n} v_p(k!)}$: independent of the coprime offset a , a perfect square, and supported on the primes dividing m . Taking $m = p$ and $a = 1$ settles the OEIS A122251 conjecture, with A122249 as the binary case. Also gives a closed form for the reduced denominator and all its local valuations, a sharp gcd theorem over the allowed offsets realised at just two explicitly constructed shifts, the extension to unequal row and column steps $\det(1/(a + m_1 i + m_2 j))$, recurrences, digit-sum asymptotics, and a Lambert-series generating

function with a natural boundary. *Merged; the determinant is evaluated three independent ways, and a second proof of the numerator formula runs through an explicit inverse matrix and MacMahon's boxed plane partitions.*

5.4 Growth laws and run statistics

37. A proof of the Apéry Hankel growth conjecture

[hankel-determinants/growth-and-runs/](#)

[apery-hankel-determinant-growth/apery_hankel_growth.pdf](#)

from [apery_hankel_research.zip](#); batch 73: [apery_hankel_research.zip](#) (a different manuscript under the same archive name); batch 77: [apery-shifted-release.zip](#)

For $A_k = \sum_j \binom{k}{j}^2 \binom{k+j}{j}^2$ and $D_n = \det(A_{i+j})$, proves $\log D_n = n(n+1) \log \frac{17+12\sqrt{2}}{4} + O(n)$ — the limit conjectured in the September 2026 revision of OEIS A228143 — using Edgar's moment-density theorem. Also an explicit law for $\det(A_{m(i+j)+r_n})$ with $r_n/n \rightarrow \rho$, and non-D-finiteness of the formal generating series of D_n . Part II, added in batch 73, determines what Part I left open: by sharp Szegő asymptotics, $h_n/\Lambda^{2n} \rightarrow K = (\pi C/2) \exp\langle \log \varphi \rangle \approx 0.6334$, an arcsine average of Edgar's density (a quadrature value, not interval-certified), so $D_n/D_{n-1} \sim K \Lambda^{2n}$ and $\log D_n = n(n+1) \log \Lambda + (n+1) \log K + o(n)$, with $C = 17 + 12\sqrt{2}$ and $\Lambda = C/4$; it refines Part I's stride theorem at $\rho = 0$. Its re-proof of Part I's main theorem is printed as a corollary, not a new result. Part III, added in batch 77, proves for $H_N(r) = \det(A_{r+i+j})_{0 \leq i,j < N}$ and $s = r/N$ in compact subsets of (s_*, ∞) , $s_* = 3\sqrt{2}/4 - 1$, the full equivalent $H_N(r) = M(s)N^{-1/24}e^{N^2A(s)+NB(s)}(1+c(s)/N+O(N^{-2}))$ with an exact Selberg normalization, by adapting Charlier–Gharakhloo's determinant-ratio analysis to a logarithmic potential; also that the floor-shifted sequence is eventually increasing, and rational-residue inverse thresholds, whose inversion cites the transseries volume. Its quadratic term $A(s)$ is Part I's stride theorem at $m = 1$ and is not claimed as new; strides $m \geq 2$, $s \leq s_*$ and the prefactor of D_n stay open.

38. Binary Runs in Rueppel Hankel Determinants

[hankel-determinants/growth-and-runs/rueppel-binary-run-determinants/rueppel_hankel.pdf](#)

from [rueppel_hankel_research.zip](#)

For the Rueppel series $r(x) = \sum_{j \geq 0} x^{2^j-1}$, the polynomial identity $H_n(tx + 1/r(x^2)) = (-1)^{1+\varepsilon+\lfloor R(n)/2 \rfloor}(\varepsilon + (R(n) - \varepsilon)t^2)$, where $R(n)$ counts runs of equal bits in the binary expansion of n and ε is the low bit of the Gray code $n \oplus \lfloor n/2 \rfloor$. Settles statements 8, 10, 13 and 23 of Barry's 2021 paper (not the identically numbered ones elsewhere), with a correction.

6 Congruences and p -adic valuations

6.1 Supercongruences

39. Congruences for OEIS A348410 at every prime

[congruences-and-valuations/supercongruences/a348410-all-prime-congruences/article.pdf](#)

from [a348410_supercongruence.zip](#) + [A348410_Supercongruence_Research.zip](#)

For $A_{\alpha,\beta}(n) = [x^n](1-x)^{-\alpha n}(1+x)^{-\beta n}$ proves the odd-prime cubic supercongruence $v_p(A(Np) - A(N)) \geq 3v_p(N) - \varepsilon_p$ with $\varepsilon_3 = 1$, $\varepsilon_p = 0$ for $p \geq 5$, and no coprimality condition; $(\alpha, \beta) = (2, 1)$ is Bala's A348410 conjecture. Separately proves an exact binary leading-defect formula, $A(2^r) - A(2^{r-1}) \equiv 2^{3r-3} \alpha \beta \binom{(\alpha+\beta+1)m-1}{m-1} \pmod{2^{3r-2}}$, hence the all-prime bound $3r - v_p(12)$ and sharp denominators 12 and 24. An appendix audits arXiv:2104.10754v1, whose framing conclusion and weighted harmonic assertion fail on its own displayed examples. *Merged; the odd-prime and binary halves run on different mechanisms — a doubling automorphism, and reflection with formal integration by parts — and both are kept.*

Not a duplicate of signed-binomial-supercongruences, and checked. That report's $a_{r,s,t}(n) = [x^{tn}](1+x)^{rn}(1-x)^{sn}$ specializes at $t = 1$, $(r, s) = (-\beta, -\alpha)$ to the coefficient studied

here, and the two odd-prime statements then agree. Neither contains the other. That one is general in the extraction step t , which is fixed at 1 here; this one adds the prime 2 with an exact defect and its sharpness, every prime at once, minimal common denominators, the primitive-cyclic-word interpretation, the Lambert/Euler product, the algebraic generating functions, a spacing-three counterexample, and a sufficient criterion beyond the family. The proofs are different arguments: that one imports Jacobsthal's congruence, this one uses none and pairs a quadratic-tail estimate with a coefficient estimate.

40. Signed binomial supercongruences and OEIS A352373

[congruences-and-valuations/supercongruences/a352373-signed-binomial-supercongruences/signed_binomial_supercongruence.pdf](#)
from OEIS_A352373_supercongruence.zip

For $a(n) = [x^{tn}](1+x)^{rn}(1-x)^{sn}$ proves $a(mp^h) \equiv a(mp^{h-1}) \pmod{p^{3h}}$ for $p \geq 5$ and $a(m3^h) \equiv a(m3^{h-1}) \pmod{3^{3h-1}}$, with m not required coprime to p . Covers A352373, A348410, A351856 and A351857 as specialisations; the only non-elementary import is the classical Jacobsthal binomial congruence.

Not a duplicate of *all-prime-congruences*, and checked. Setting $t = 1$ and $(r, s) = (-\beta, -\alpha)$ turns the theorem below into that report's odd-prime tower, and the moduli agree after reindexing the multiplier. Neither contains the other: this one is general in the extraction step t , that one is fixed at $t = 1$ but proves the prime 2 case, the all-prime bound, the sharp denominators and much else. The two proofs are genuinely different — this one imports Jacobsthal's congruence, that one does not use it at all.

41. Apéry numbers under exponentiation and series reversion

[congruences-and-valuations/supercongruences/a267220-apery-transform-supercongruences/article.pdf](#)
from Apery_OEIS_Supercongruences.zip, apery_oeis_conjectures(1).zip

Proves both supercongruence conjectures contributed by Peter Bala to OEIS A267220 on 17 October 2024, for every integer parameter including negative and zero. With $A(x) = \exp \sum_{n \geq 1} a_n x^n / n$ for the Apéry numbers a_n , $F(x) = x / \text{Rev}(xA(x))$, $u_m(n) = [x^n]A(x)^{mn}$ and $v_m(n) = [x^n]F(x)^{mn}$: for odd p and $R = \nu_p(N)$ both families satisfy $\nu_p(u_m(N) - u_m(N/p)) \geq 2R + \nu_p(m)$, the unsigned statement for v_m holds at *every* prime including 2, and a *signed* refinement $(-1)^{mN}u_m(N) \equiv (-1)^{mN/p}u_m(N/p)$ recovers full precision at $p = 2$, where the unsigned bound genuinely loses one power. Merged from two reports that share a seed lemma and a Möbius/dilogarithm layer but diverge at the crux; both routes are kept, and the merged text is careful that the signed and unsigned 2-adic statements are different statements, not a sharper and a weaker version of one.

42. Four Motzkin congruences

[congruences-and-valuations/supercongruences/a001006-motzkin-congruences/article.pdf](#)
from motzkin_congruences.zip

Proves all four conjectures in Peter Bala's comment of 10 February 2022 on OEIS A001006, with sharp ranges. The engine is a prime-power block-transfer theorem: for odd p and $q = p^{k-s+1}$, $h = mp^{s-1}$, the Motzkin numbers satisfy $M_{mp^k+d} \equiv T_h M_d$ for $0 \leq d \leq q - 3$, with explicit corrections at $d = q - 2$ and $d = q - 1$; the boundary case gives $M_{np^{r-2}} \equiv \frac{\varepsilon_p^r - 1}{2} T_{n-1} - U_{n-1} \pmod{p}$. Adds the exact first failures beyond the two initial-block ranges, arbitrary-multiplier and all-prime-power boundary extensions, a prime-modulus digit algorithm, and examples showing why unweighted repetition modulo a prime cube is generally false. Discusses Burns, Kohen and Pan-Sun rather than claiming novelty.

6.2 Congruences for iterated and compositional series

43. Arithmetic of a Self-Iterating Generating Function

[congruences-and-valuations/iterated-series/a396807-self-iterating-series/article.pdf](#)
from [A396807_congruences_and_prime_power_lifts.zip](#)

Both conjectures in OEIS A396807 for $A(x) = x + A^{[5]}(x)A^{[6]}(x)$ (functional iteration): $A(x) \equiv x/(1-x)$ and $A^{[k]}(x) \equiv x/(1-kx)$ modulo 10 for every integer k , negative iterates included. Adds a sharp modulus theorem for $F = x + cF^{[r]}F^{[s]}$, rational lifts at every selected prime power, and minimal period 46860 for A396807 modulo 100 starting at index 1.

44. Prime-power congruences for compositional tree series

[congruences-and-valuations/iterated-series/compositional-tree-series-congruences/article.pdf](#)
from [compositional_tree_congruences.zip](#)

For $A_l(x) = x \exp(A_l^{ol})$, shows the OEIS A396805 claim $a_5(n) \equiv n \pmod{5}$ is *false*: first failure $n = 23$, with exception set $\{20j + 3, 20j + 4 : j \geq 1\}$ and residue 2 throughout. Proves eleven other conjectural assertions in A396803–A396806 via a general reduction of coefficients mod p^α to typed rooted trees on fewer than p^α labels, yielding eventual periods, rational residue generating functions, and an exact classification of $a_l(n) \equiv n \pmod{p}$.

45. Lacunary iterates modulo four

[congruences-and-valuations/iterated-series/a168362-lacunary-iterates-mod4/article.pdf](#)
from batch 73: [ProveIt_A168362_Mod4_Theorem.zip](#)

For $F(x) = \sum_{j \geq 0} x^{2^j}$ and its compositional iterates $G_m = F^{\circ m}$, OEIS A168362 is $a(n) = [x^n]G_n(x)$, and the entry records a sparse 2026 pattern for $a(n) \pmod{4}$. Proves $[x^N]G_m(x) \equiv 0 \pmod{4}$ for all $m, N \geq 1$ whenever N has at least three binary ones, gives exact bit formulas at powers of two and sums of two powers of two, and so classifies the diagonal completely: apart from 2, 3, 4, 6, $a(n) \equiv 2 \pmod{4}$ exactly when $n = 2^r + 2^s$, $r > s \geq 1$, $r \geq 3$, and $(r-s) \pmod{2^s}$ lies in an explicitly computable residue set R_s . This confirms every exception printed in the entry; their number up to X is $\kappa \log_2 X + O((\log \log X)^3)$ with $\kappa = \sum_s |R_s|/2^s \approx 2.32393$. The ratio conjecture $a(n+1)/a(n) \sim en$ stays open.

6.3 Valuations, divisibility and periodicity

46. Binary Carries and the Divisibility of Two-Stack-Sortable Permutation Numbers

[congruences-and-valuations/a000139-binary-carries-parity/research.pdf](#)
from [a000139_carries_research.zip](#)

Proves Peter Bala’s parity conjecture for OEIS A000139, $a(n) = 2(3n)!/((n+1)!(2n+1)!)$: $a(n)$ is odd exactly when n is odd and its binary expansion contains no 11. The full valuation is $v_2(a(n)) = s(n) + s(n+1) - s(3n)$ in binary digit sums. Two independent parity proofs, plus exact valuation distributions, recurrences, asymptotics, extremal formulas and limit laws.

47. Polynomial divisibility of height-one factorial ratios

[congruences-and-valuations/factorial-ratio-polynomial-divisibility/article.pdf](#)
from [bala_factorial_divisibility_research.zip](#); batch 74: [OEIS_Fractional_Factorial_Theorems.zip](#)

A general rational-root criterion for uniform polynomial divisibility of balanced integral factorial ratios, applied to Peter Bala’s uniform-product conjectures for OEIS A211417 and A295431: explicit multipliers for every product length, with the least multipliers certified in eight particular examples. Includes an atlas of the 52 sporadic height-one ratios with recurrences, generating functions and asymptotic expansions. Part II, added in batch 74, proves integrality and algebraic generating functions for A347854–A347858 through a rational-dilation theorem for height-one factorial ratios with composite denominators, settling the Gamma-interpolated values Part I left open for these five entries; it also proves $6(6n+1)(12n-1) \mid A295432(n)$ and gives finite-monodromy certificates and moment laws. Its re-proof of Part I’s A295431 theorem, with larger constants and no optimality, is printed only as a marked second route.

48. Integrality of the Brent–Glasser–Guttmann asymptotic coefficients

[congruences-and-valuations/a321941-asymptotic-coefficient-integrality/article.pdf](#)

from batch 73: [A321941_Arithmetic_Conjectures.zip](#)

Brent, Glasser and Guttmann introduced rational coefficients d_k in the all-orders asymptotic expansion of a Hadamard product involving the exponential integral, conjectured that $r_k = 64^k d_k$ (OEIS A321941) is always an integer, and observed $r_k \equiv \binom{2k}{k} \pmod{32}$ for $k \geq 3$. Proves both, by keeping a quadratic invariant of the underlying second-order recurrence rather than only its symmetric-square recurrence, and extends them to polynomials $P_k(t) \in \mathbb{Z}[t]$ with $P_k(1) = r_k$, even for $k > 0$, with a coefficientwise congruence modulo 32. Consequences: the scale 64 is minimal, denominator bounds are sharp at infinitely many indices and exact when the index has at most four binary ones, and the inverse expansion has no denominator primes other than 3. The analytic expansion itself is the published theorem; the negativity conjecture stays open.

49. When the Secant Numbers Become Periodic

[congruences-and-valuations/secant-number-periodicity/article.pdf](#)

from [secant_periodicity_research.zip](#)

Refutes the OEIS A000364 conjecture that the secant numbers from $n = 1$ are purely periodic modulo every m : at $m = 27$, $a(1) \equiv 1$ but $a(19) \equiv 10$. The exact replacement is that pure periodicity from index one holds if and only if no odd prime cube divides m ; the least eventual period always divides $\varphi(m)$. Determines the least eventual period and exact onset for every modulus, with densities and a certified mean onset.

7 Tetration and digit stabilization

50. A Counterexample to a Decimal-Length Bound for Tetration Stabilization

[tetration-and-digit-stabilization/congruence-speed-decimal-length-bound/article.pdf](#)

from [tetration_counterexample.zip](#)

An explicit 2544-digit base $A = 1 + qu$ ($q = 5^{2547}$) for which $\nu_{10}(T_{n+1} - T_n) = \min(1 + 2548n, 2547(n + 1))$ at every height, so the congruence speed at height $\text{len}(A) + 2$ is 2548 while the permanent value is 2547. This contradicts the decimal-length form of Conjecture 1 of Marco Ripà, *The congruence speed formula* (NNTDM 27(4), 2021). The report audits the later Ripà–Onnis confirmation: their weaker valuation bound is compatible with the counterexample, the decimal-length bound is not.

51. Tetration, perfect powers, and frozen decimal digits

[tetration-and-digit-stabilization/perfect-power-minima-for-frozen-digits/article.pdf](#)

from [tetration_perfect_power_minima.zip](#)

Determines $\rho(n)$, the smallest n th perfect power $x^n > 1$ whose eventual number of newly frozen terminal decimal digits is exactly n : $\rho(1) = 2$, $\rho(2) = 49$, and $\rho(n) = B(s)^n$ for $n \geq 3$ with $s = n - v_2(n)$ and $B(s) \in \{2^s \pm 1, 3 \cdot 2^s \pm 1\}$ according to $s \pmod{4}$. All other roots are ruled out by a deterministic quadratic size barrier plus small exact certificates.

52. Phase-shift classification for integer tetration

[tetration-and-digit-stabilization/](#)

[phase-shift-classification/tetration_phase_classification.pdf](#)

from [tetration_phase_classification.zip](#)

Proves the 21-word phase-shift conjecture in OEIS A376842, the 14-word even-base restriction from the appendix of Ripà’s 2025 paper, and the 28-word prescribed-anchor conjecture in OEIS A376446. The mechanism compares the 2-adic and 5-adic valuations of $D_b = T_{b+1} - T_b$ at the first permanently constant-speed height: strict 2-adic dominance gives phase 5, strict 5-adic dominance a multiplicative orbit in $\mathbb{F}_5^\times$, and balanced valuations exactly four further anchored words.

53. A Sharp Convergence Theorem for the q -Newton Series of Subcritical Tetration

[tetration-and-digit-stabilization/q-newton-series-convergence/article/](#)

[tetration_qnewton.pdf](#)

from [tetration_qnewton_research.zip](#)

Proves convergence of the explicit q -binomial interpolation series of MathOverflow question 259278 for every real base $1 < a < e^{1/e}$, with $\log a = qe^{-q}$: the grouped series converges absolutely and locally uniformly for $\operatorname{Re} z > -2$, converges conditionally on $\operatorname{Re} z = -2$ except at $z = -2 + 2\pi ik/(-\log q)$, and diverges for $\operatorname{Re} z < -2$. Adds strict complete monotonicity and uniqueness under an eventual log-convexity condition.

54. Stabilization of integer power towers

[tetration-and-digit-stabilization/power-tower-stabilization/article.pdf](#)

from [tetration_stabilization_research.zip](#) + [tetration_exact_digits.zip](#)

Refutes the claim that every prime base ending in 9 reaches its permanent decimal congruence speed by height 2 (Ripà, Conjecture 2; arXiv:2208.02622v1, Conjecture 4.1): 2749 is the smallest prime counterexample, with $H(2749) = 3$. Proves exact formulas for $C_a(h)$, $V(a)$ and $H(a)$, an exact natural density for onsets, and infinitely many primes at each prescribed onset via CRT and Dirichlet. Also resolves the three conjectural comments in OEIS A324017 — two proved, the third disproved, with $3 \uparrow\uparrow 2 \equiv 27$ and $3 \uparrow\uparrow 3 \equiv 59 \pmod{64}$ — via $T_{h+r} - T_h \equiv -2q^h \pmod{q^{h+1}}$ for even $q \geq 4$, so exactly one radix- q digit freezes per height. Adds a factorization-free digit-lifting algorithm, modular saturation across Knuth arrow ranks, and a sign-sensitive local p -adic Lambert W formula. *Merged; the shared one-digit-per-height theorem was proved identically in both originals and is proved once here.*

55. Delayed digit stabilization: a counterexample and sharp bounds

[tetration-and-digit-stabilization/](#)

[iterated-power-digit-stabilization/delayed_digit_stabilization.pdf](#)

from [digit_stabilization_research.zip](#)

For $S_a(b) = v_{10}(a^{10^{b+1}} - a^{10^b})$ and $D_a(b) = S_a(b) - S_a(b-1)$, answers negatively the question whether $D_a(b) = 1$ for all $b \geq 3$: the smallest counterexample is $a = 3 \cdot 2^{99} + 1$, with $S_a(2) = 100$, $S_a(3) = 102$, $D_a(3) = 2$. Proves minimality, an exact onset formula, the smallest counterexample at every stage, the natural density of exceptions, and arbitrary-radix extensions. The article is explicit that a^{10^b} is not genuine tetration and derives the tower formulas separately.

8 Log-concavity, unimodality and positivity

56. Single curves do not force log-concavity

[log-concavity-and-unimodality/cluster-variable-log-concavity-counterexample/article.pdf](#)

from [annular_single_lamination_counterexample.zip](#)

An unpunctured annulus with one marked point on each boundary and a single unit-weight simple lamination curve gives, after mutations 2, 1, 2 with the two mutable variables specialised to 1, the polynomial $1 + 3q^2 + 2q^3 + 4q^4 + 2q^5 + q^6$; the consecutive coefficients 3, 2, 4 form a strict valley, refuting Conjecture 1 of arXiv:2508.04396v3 and conflicting with its general unpunctured-surface unimodality statement. The geometric realisation is proved, not merely the matrix calculation, and a fixed seed gives infinitely many such examples.

57. Cycles refute infinite log-concavity for chromatic coefficients

[log-concavity-and-unimodality/infinite-log-concavity-of-cycles/article.pdf](#)

from [chromatic_cycle_logconcavity.zip](#)

For the cycle C_n with absolute chromatic coefficients under the iterated operator $L(a)_k = a_k^2 - a_{k-1}a_{k+1}$: infinitely log-concave for $3 \leq n \leq 11$; first failure at depth 5 for $n = 12$, depth 4 for $13 \leq n \leq 16$, and depth 3 for $n \geq 17$. So C_{12} is the smallest cycle counterexample (no minimum over all graphs is claimed), with an explicit third-iterate witness polynomial valid for all $n \geq 3$.

58. Oscillating Log-Concavity in Geometric Generating Functions

[log-concavity-and-unimodality/oscillating-log-concavity-geometric-transform/article.pdf](#)
from [oscillating_log_concavity_research.zip](#)

Challenges 2 and 3 in Section 3 of Heim and Neuhauser, arXiv:2302.13327v1, for the *geometric* transform $1/(1 - uG(z))$. For divisor-sum weights, every real exponent and every $u > 0$ give infinitely many strict log-concavity and log-convexity indices with eventually bounded gaps; likewise for integer power weights with exponent ≥ 2 , the exponents 0 and 1 being explicitly exceptional. For square weights at $u = 1$, every four consecutive indices carry both signs (sharp), each with density $1/2$ in every arithmetic progression. A subdominant-pole theorem is the common mechanism.

59. A counterexample and a sharp spacing-three criterion for products of q -integers

[log-concavity-and-unimodality/q-integer-product-unimodality/report.pdf](#)
from [q_unimodality_attack.zip](#)

The necessity half of Conjecture 5.4 of Connelly, Ito, Martinez, Shevchenko and Yang (arXiv:2605.12822v1) is false: $(1 + q)^6(1 + q^3)$ is symmetric and weakly unimodal although no a_i is divisible by 3 and $b = 2$ exceeds the asserted bound, and the example lies inside the source's claimed range. The report also proves the proposed sufficient condition for every $r \geq 2$, a necessary degree bound in the absence of an r -divisible ordinary factor, and a complete classification of spacings two and three.

60. Sharp size thresholds for failures of log-concavity in independence systems

[log-concavity-and-unimodality/independence-system-thresholds/article.pdf](#)
from [sharp_log_concavity_thresholds.zip](#)

For each rank $r \geq 4$, determines the smallest ambient ground-set size admitting failure of ordinary, ordered and ultra log-concavity, and shows the three thresholds coincide across all independence systems, those satisfying Xie–Xu's augmentation bound, and those with additive augmentation parameter 3 (values $7/6/5$, $7/7/6$, $8/7/7$, and $r + 1$ throughout for $r \geq 7$). The initial target, Xie–Xu Conjecture 4.8 (arXiv:2408.09152v1), had already been disproved by Alex Chengyu Li in September 2026; the report states this and refines the target to the rankwise minimum-size question.

61. A sharp order-five threshold for higher-order Stirling subset log-concavity

[log-concavity-and-unimodality/higher-order-stirling-rows/article.pdf](#)
from [stirling_log_concavity.zip](#)

For $S_{n,k}^{(r)}$, the number of partitions of $n + (r - 1)k$ labels into k blocks of size $\geq r$, proves that all rows are log-concave if and only if $1 \leq r \leq 5$, with strict inequalities in the interior — the log-concavity assertion of Deb–Sokal Conjecture 1.4(c) (arXiv:2507.18959v1). Also proves the cycle cases $r = 3, 4$; the fifth-order *cycle* conjecture is explicitly left unresolved, and the included operator counterexample is flagged as not being a counterexample to it.

62. A sum-of-squares proof of Ulas's real-zero conjecture

[log-concavity-and-unimodality/stern-polynomial-positivity/stern_conjecture_note.pdf](#)
from [stern_conjecture_research.zip](#)

Conjecture 4.4 of Maciej Ulas, arXiv:1909.10844v1, for the Stern polynomials indexed by $a_n = (4^n - 1)/3$ and $h_n = 2(4^n - 1)(2 \cdot 4^n + 1)/3 + 1$: for every $n \geq 0$ and every real t , $W_n(t) = (A_{n+1}(t) - tA_n(t)/2)^2 + 3t^2A_n(t)^2/4 > 0$. The adjacent-polynomial quadratic identity behind it was already in Ulas's paper and is re-derived, not claimed as new.

63. Two-coin counterexamples to generalized Shepp–Olkin thresholds

[log-concavity-and-unimodality/bernoulli-entropy-quasiconcavity/article.pdf](#)
from [bernoulli_entropy_research.zip](#); batch 43: [proveit_product_entropy.zip](#); batch 57: [ProveIt_Alphabet_Entropy.zip](#)

For every finite real $q > 1$, both the Rényi- q and the Tsallis- q entropy of a sum of independent Bernoulli variables fail even *quasiconcavity* as a function of the parameter vector, already on the open square $(0, 1)^2$ with two coins. This disproves the proposed thresholds — Rényi 2 and Tsallis

3.65986... — adjoined to Conjecture 4.2 of arXiv:1503.01570 and restated in arXiv:1510.05390, Section 8. Part II, added in batch 43, settles the subunit joint-concavity question for the joint bit vector only: product Rényi entropy is concave in every dimension for $0 < q \leq 2$, and product Tsallis entropy is concave for every $0 < q < 1$ exactly when $n \leq 9$. Part III, added in batch 57, answers Part II's question on larger alphabets for full categorical simplices: for $0 < q < 1$ product Tsallis entropy on $\Delta_{k_1} \times \cdots \times \Delta_{k_n}$ is concave exactly when $\sum_i \Theta_{k_i}(q) \leq 1$ for an explicit alphabet profile Θ_k ; at $q = 1/2$ three factors with a common alphabet of k symbols are concave exactly when $k \leq 17$; concavity for every choice of n alphabets holds exactly when $nq \leq 1$; and for $1 < q < 2$ concavity reduces to positive semidefiniteness of an $n \times n$ compression matrix. For the ordinary sum Part I's question stays open; an external claim about it is recorded as unverified.

64. Nonreal roots survive in polynomial iterative equations

[log-concavity-and-unimodality/nonreal-roots-in-iterative-equations/article.pdf](#)

from nonreal_roots_research.zip; batch 41: [ProveIt_Increasing_Spectral_Core.zip](#); batch 42:

[ProveIt_Orientation_Reversing_Spectra.zip](#); batch 54: [ProveIt_Exact_Algebraic_Degree.zip](#)

Here $P[f] = \sum_j a_j f^{oj}$ means a combination of *iterates*, not a polynomial in pointwise values. Gives a negative answer to Problem 6.1 of Draga–Morawiec (arXiv:1503.00570), which proposed deleting nonreal maximal-modulus characteristic roots: an explicit odd, strictly increasing, globally bi-Lipschitz homeomorphism realises a cubic whose nonreal root cannot be discarded. Adds a classification of the circular root spectra that occur. Part II, added in batch 41, classifies the minimal iterative polynomials of increasing homeomorphisms with roots on several circles, through a universal spectral core and an affine rigidity theorem. Part III, added in batch 42, does the same for decreasing homeomorphisms with $P(0) \neq 0$: two-cycle and one-sided spectra, a decreasing core with one or two witnesses, and a finite-smoothness classification. Part IV, added in batch 54, answers Part III's question on algebraic degree: for a polynomial coordinate H and $f = H \circ (a \text{ id}) \circ H^{-1}$ the degree of f over $\mathbb{R}(x)$ is $\deg H / \deg W$, where W is the largest normalized right compositional factor of H with $W(at) = a^{\deg W} W(t)$, and it is exactly $\deg H$ for every coordinate of Part III's finite-smoothness theorem. For every odd $n \geq 5$ it gives real-analytic involutions of algebraic degree n , with distortion arbitrarily close to 1, whose graph extension has Galois group S_n , so they are not expressible by radicals.

65. Canonical binomial decomposition does not preserve real roots

[log-concavity-and-unimodality/binomial-decomposition-real-roots/article.pdf](#)

from mu_welker_recursive_decomposition.zip + recursive_binomial_counterexample.zip

A negative answer to Question 3.10 of Mu and Welker (arXiv:2503.24076v1) for their canonical decomposition $F(t) = G(t) + tH(t)$ built from the integer binomial expansion of every coefficient: real-rootedness of F does not pass to the parts. Two cubic witnesses are carried, one with a repeated root and one with simple roots, each with its own digit table, discriminant pair and exact nonreal zeros; the report also gives the minimal witness within cubics, two all-degree thresholds neither of which implies the other, and a two-sided limit. *Merged; the two originals were two write-ups of one proof program, each already containing both witnesses with the headline and the co-star swapped.*

66. A first-triple criterion for log-concavity of MDS weight distributions

[log-concavity-and-unimodality/mds-weight-log-concavity/article.pdf](#)

from mds_log_concavity_research.zip

For $d = n - k + 1 \geq 2$, $k \geq 3$ and $q > d$, the compressed formal MDS weight distribution $(1, A_d, \dots, A_n)$ is log-concave if and only if the single inequality $A_{d+1}^2 \geq A_d A_{d+2}$ holds — the whole sequence is controlled by its first triple.

67. A uniform Hankel obstruction for higher-order Stirling subset polynomials

[log-concavity-and-unimodality/stirling-hankel-obstruction/article.pdf](#)

from stirling_hankel_obstruction.zip

Proves the higher-order negative assertion of Deb–Sokal Conjecture 1.4(d), with a stronger statement at every positive shift: for $r \geq 3$ and $a \geq 1$ the 3×3 Hankel determinant $\det(s_{r,a+i+j}(x))$

of the row-generating polynomials fails the required positivity. Clause (d) of that conjecture; clause (c) is the subject of the higher-order Stirling rows entry.

68. Normalization by the matching number for weighted matching-support polynomials

[log-concavity-and-unimodality/matching-rank-normalization/article.pdf](#)

from batch 72: [ProveIt_Weighted_Rank_Five_and_Two_Shore_Covers.zip](#),
[strictness_equality_package.zip](#), [ProveIt_Weighted_Rank_Four_Stability.zip](#),
[ProveIt_Sharp_Full_Hall_Classification.zip](#),
[ProveIt_Unit_Weight_Rank_Normalization_Counterexample.zip](#), [ProveIt_One_Shore_Counterexamples.zip](#),
[ProveIt_Sharp_Rank_38_Hall_Endpoint.zip](#), [ProveIt_Sharp_Rank_38_Full_Hall_ULC.zip](#),
[ProveIt_Last_Hall_Gap_Separation.zip](#), [ProveIt_Rank_Two_Matrix_Blocks_and_Conditioning.zip](#),
[ProveIt_Three_Vertex_Marginals_and_Star_Cores.zip](#),
[ProveIt_Rooted_Hall_Sectors_and_Cubic_Obstruction.zip](#), [ProveIt_Rank_Six_Finite_Last_Gap.zip](#),
[ProveIt_Rank_Six_Second_Newton_Inequality.zip](#), [ProveIt_Rank_Six_Common_Exterior_Theorem.zip](#),
[ProveIt_Rank_Six_Nested_and_Conditional_Results.zip](#),
[ProveIt_Rank_Six_Conditioning_for_Every_Core.zip](#), [ProveIt_Rank_Six_Complete_Core_Cubic.zip](#),
[ProveIt_Rank_Six_Matching_Hole_Cores.zip](#), [ProveIt_Rank_Six_Two_Complete_Rows.zip](#),
[ProveIt_Rank_Six_Two_Complete_Columns.zip](#), [ProveIt_Rank_Six_At_Least_Six_Core_Edges.zip](#),
[ProveIt_Eventual_One_Shore_Scaling_Through_Rank_Seven.zip](#) (with a second manuscript nested inside it), [ProveIt_Eventual_Five_Plus_Three_Covers.zip](#), [ProveIt_Four_Plus_Four_Top_Gap.zip](#),
[ProveIt_Eventual_One_Shore_Scaling_Through_Rank_Eight.zip](#); batch 73:
[matching-rank-four-package.zip](#), [all-two-by-three-cores-result.zip](#),
[two-element-matroid-lorentzian-result.zip](#), [sharp-weighted-rank-boundary-result.zip](#)

Continues Parts XI–XIII of the preorder root-polytope report on whether the bipartite matching-support polynomial p_G , with positive vertex activities, is ultra-log-concave of the order of the matching number $\nu(G)$; merged from twenty-seven manuscripts, one of them delivered only inside another’s archive. Part I proves it for arbitrary positive activities on both shores whenever some minimum vertex cover meets one shore in at most two vertices, through an explicit transversal-matroid lift, so for every graph with $\nu(G) \leq 5$; with that report’s rank-six Hall counterexample the least matching number of a weighted failure is exactly six. Equality in a normalized inequality occurs only for balanced star forests, covers with two vertices on each shore give real-rooted support polynomials, and the full Hall family is ultra-log-concave for all activities exactly when one core has at most two vertices. Part II answers the unit-weight and one-shore versions negatively: a connected all-unit graph with 6916 vertices and $\nu = 3450$ fails, as does a 125-vertex graph with $\nu = 50$ weighted on one shore only, and for one common activity on the right core of a full Hall graph the least failing matching number is exactly 38. Part III, at $\nu = 6$ with all left activities one, proves the second and last of the five inequalities for every graph (the first holds in general), and all five for several core families. Part IV proves eventual ultra-log-concavity, after a common scaling of the right part of a minimum cover, for every graph with $\nu(G) \leq 8$ and unit left activities, and locates the least matching number of a persistent failure in $[9, 38]$. Results that the preorder report already proves are cited there, not reprinted, and theorems proved by several manuscripts are printed once with the other proofs as second routes. Part V, added in batch 73 from four manuscripts written in parallel with batch 72, shows that at unit weights and matching number three a nonreal zero needs fifteen vertices, attained by a unique graph; proves a two-element Lorentzian deformation of every matroid basis polynomial and an all-rank first-layer positive-semidefiniteness theorem (which proves Part III’s three-vertex marginal theorem without its 5,339-monomial certificate); gives a stable seed for two-neighbour star cores; and shows by three 13-vertex examples that the two-tail lifts do not extend to three tails. What those manuscripts re-prove of Parts I and III is cited or kept as second routes; six further archives of the batch (two earlier editions, four re-proofs or special cases) were not placed.

69. Rank ultra-log-concavity of preorder gamma and directed support polynomials at the actual degree

[log-concavity-and-unimodality/preorder-gamma-rank-ulc/article.pdf](#)

from batch 73: preorder-degree-four-part-1-of-4.zip, preorder-degree-four-part-2-of-4.zip, preorder-degree-four-part-3-of-4.zip, preorder-degree-four-part-4-of-4.zip, preorder-gamma-degree-three-package.zip, template26-structural-supplement.zip, role-cover-package.zip, three-core-rayleigh-result.zip, physical-cover-three-result.zip, bipartite-and-pendant-support-result.zip, independent-role-degree-three-package.zip, universal-sink-last-gap-package.zip, universal-sink-all-degrees-package.zip, universal-sink-five-core-package.zip, universal-sink-five-core-boundary-package-part-1.zip, universal-sink-five-core-boundary-package-part-2.zip; superseded, not printed: weighted-degree-three-package.zip, universal-sink-degree-four-package.zip; batch 77: universal-sink-five-core-cubic-package.zip

Answers Research question 96 of the preorder root-polytope report: is the gamma polynomial $\Gamma_D = \sum_k \gamma_k t^k$ of a finite preorder, with γ_k counting ordered pairs of disjoint k -sets joined along arcs, ultra-log-concave after normalization by its *actual* degree? Part I proves it for every preorder of actual degree at most four with unit activities: degree three by a Gallai–Edmonds reduction and exact integer-population certificates (a nonreal-rooted cubic needs fifteen vertices), degree four by bounded cores, 76 four-attachment templates and 50 global binomial-square identities, with a smaller second route for the largest certificate. Part II treats independent tail and head activities on arbitrary loopless directed relations: real stability of the signed monomer polynomial under every two-by-two role cover (generalizing the $2 + 2$ theorem of the matching-rank report), coefficientwise Rayleigh inequalities for a three-vertex shore, rank ultra-log-concavity under every physical vertex cover of size three and on bipartite physical graphs, a pendant-head stability criterion, and the full statement for independently weighted preorders of actual degree at most three. Part III adds a cloud of universal sinks to an arbitrary directed core: a sharp last-coefficient bound for every core size, and the statement at every actual degree for four-vertex cores and at actual degrees four and five for five-vertex cores. The degree-three and degree-four theorems are computer-assisted. Merged from eleven manuscripts and a six-page supplement, from eighteen archives (three of them handed over from the matching-support cluster); two earlier versions are superseded and theorems proved more than once are printed once with second routes. Section 14 of Part III, added in batch 77 from a further manuscript, proves $\gamma_2^2 \geq 3\gamma_1\gamma_3$ on the face $\gamma_4 = 0$ for every loopless directed five-vertex core with universal sinks and independent activities, with the constant 3 sharp even for preorder cores, by a degree-drop reduction to a computer-assisted two-sink lemma with 7,901 rational square certificates; so five-vertex cores are rank ultra-log-concave at every actual degree. For preorder cores it is a case of Part II’s degree-three theorem, so the new content is the non-transitive core; Section 12’s counterexample to the constant 3 has $\gamma_4 > 0$. Some certificate data over 2 MB, and Section 14’s 95.1 MB of certificate shards, which cannot be regenerated, stay in the arrival commits, and the report says how to retrieve or rebuild them.

9 Quaternionic analysis

One report, first merged from ten manuscripts and since extended by two more. The subject is new to this collection, and the delivery was unusual: one expository survey and nine independent attempts at the same follow-up problem, all written on the same day. The nine agree, which is the substance of the entry.

70. Slice regularity and the global inverse Fueter problem

[quaternionic-analysis/slice-regularity-and-fueter-inversion/article.pdf](#)

from quaternionic_analysis.zip and nine companion archives; batch 39:

ProveIt_Fueter_Polynomial_Periods.zip; batch 64: ProveIt_Fueter_Global_Inversion.zip

One article, first assembled from ten manuscripts. Part I is the delivered expository survey: slice regularity and Cauchy–Fueter regularity side by side, with Cauchy formulas, Taylor and Laurent expansions, the regular product and the geometry of zeros, the argument principle, Rouché, open

mappings, Schwarz–Pick and an intrinsic Riemann mapping theorem. Parts II–XI answer the follow-up question the other nine manuscripts all attacked independently: for an axially monogenic $g = P + IQ$, when is there a single-valued slice-regular f with $\Delta_4 f = g$? The answer is a two-period obstruction. The closed one-forms $\omega_g = \frac{1}{2}(-P dx + Q dy)$ and $\theta_g = \frac{1}{2}((xP + yQ)dx + (yP - xQ)dy)$ have vanishing periods exactly when a global primitive exists, and then $f(x + Iy) = (x + Iy)C + V$ with $dC = \omega_g$, $dV = \theta_g$, unique modulo $q \mapsto qa + b$; the monodromy is affine, and the cokernel of the Fueter map has real dimension $8m$. The point the nine exist to make is that θ_g is computed from the target, with no first primitive chosen. **Nine independent derivations reaching the same two forms, the same $8m$ and the same logarithmic energy constant is the evidence the entry rests on.** Where they differed it was almost always notation — thirteen conflicts are recorded and resolved in `MERGE_NOTES.md` — but two were mathematical and would have produced a false statement if the merge had been mechanical: the size law at a critical singularity has genuinely different sharp constants in two norms that four manuscripts both called M_g , and the conserved flux recovering the slope residue needs the weight $3(\operatorname{Re} q - u) + \operatorname{Im} q$ rather than $q - u$, the coefficient 3 being exactly what makes it right-Fueter-regular. All ten source manuscripts and all ten verification programs are kept; the deduplication is in the article. Parts XII–XIII, added in batch 39 from an eleventh manuscript, answer the report’s open problem on higher dimensions in large part: in ambient dimension $2h + 2$ one polynomial closed form of degree $2h - 1$ (at $h = 1$ the form $\theta_g + t\omega_g$) gives a complete obstruction, with a constructive inverse, a cokernel of dimension $2h d_V m$, an exact residue filtration and sharp size and energy constants. Part XIV, added in batch 64 from a twelfth manuscript, carries this across the real axis and to planar domains of arbitrary, also infinite, connectivity: every admissible polynomial period character is realized, a continuous linear right inverse of the period map exists exactly when the admissible period rank is finite, and a continuous nonlinear section exists always.

10 Graph theory

71. Mutual-visibility sets in Andrásfai graphs

`graph-theory/andrasfai-mutual-visibility/article.pdf`
from `andrasfai_visibility_research.zip`

Proves the generating function attributed to Andrew Howroyd in OEIS A391632, $\sum a(n)x^n = x(4 - 19x + 33x^2 - 40x^3 + 8x^4)/(1 - 10x + 29x^2 - 32x^3 + 8x^4)$, by an exact weight-preserving bijection between mutual-visibility sets of the n -Andrásfai graph and anchored closed walks of length $3n - 1$ in a four-state labelled graph. Adds the full bivariate generating function, an exact binomial formula for every cardinality, maximum-set classification, eventual polynomiality at each fixed deficit, exponential asymptotics, a central limit theorem, and linear-time recognition and exact uniform sampling.

72. A bicyclic 30-vertex graph with domination root -4

`graph-theory/domination-root-minus-four-order-30/domination_root_30.pdf`
from `domination_root_30_research.zip`; batch 67: `ProveIt_Domination_Cycle_Obstructions.zip`

An explicit connected planar graph on 30 vertices and 31 edges, treewidth 2 and domination number 10, with $D(G, x) = x^{10}(x + 4)Q_{19}(x)$ — a negative answer to question 2 of Section 4 in Alikhani and Griswold, arXiv:2608.00109v1, which asks whether 33 is the minimum order. Their 33-vertex example had already disproved the original integer-root conjecture; no claim of global minimality at 30 is made. Three independent exact polynomial computations agree. Part II, added in batch 67, proves $1 \leq |D(G, -1)| \leq 3^\beta$ sharply for cycle rank β , a simultaneous divisibility budget for integer roots and their multiplicities, and the exact value sets of $|D(G, -1)|$: $\{1, 3, 7, 9\}$ for $\beta \leq 2$, so no root -6 at cycle rank at most two, and a ten-element set for $\beta \leq 3$ (computer-assisted). Part I’s zero- C four-cycle search forces $|D(G, -1)| = 3$, so it can produce no integer root below -2 other than a simple -4 ; no root -6 or -8 is constructed.

73. A marked-cycle proof of the trapezohedral domination conjecture

[graph-theory/trapezohedral-minimal-dominating-sets/article.pdf](#)

from [trapezohedral_domination_research.zip](#), [A381190_proof_package.zip](#)

Proves the generating function labelled conjectural in OEIS A381190: for the n -trapezohedral graph, the number of dominating sets that are inclusion-minimal among all dominating sets and induce a connected subgraph is $a_n = 2n(2p_{n-2} + p_{n-3})$ with $p_m = p_{m-2} + p_{m-3}$ (Padovan). The proof classifies such sets by a cyclic arrangement of gaps of length 2 and 3 with a restricted marked edge. Adds the size distribution, strict log-concavity, an unbiased sampler, a Binet formula and a central limit theorem.

11 Enumerative combinatorics

74. A Twelve-Value Proof of the A181280 Formula

[enumerative-combinatorics/a181280-binary-matrix-formula/article.pdf](#)

from [A181280_research_package.zip](#)

The binary-matrix formula recorded as Conjecture 20 by Kauers and Koutschan (2023) and still marked conjectural in OEIS A181280: an explicit spectral recurrence bound and a finite certificate resting on twelve decisive counts, together with further consequences. `STATUS.md` records that general C-finiteness was already proved in Kauers's December 2025 lecture slides.

75. Esthetic Numbers, Folded Paths, and Algebraic Diagonals

[enumerative-combinatorics/esthetic-numbers-and-folded-paths/article.pdf](#)

from [esthetic_walks_research.zip](#)

The diagonal conjecture $T(q, q-2) = A182555(q-2)$ in OEIS A377000, where $T(q, k)$ counts base- q positive integers of length k whose adjacent digits differ by one. Proves it exactly, proves the entry's other two conjectures as retrieved, and derives formulas for every fixed-offset diagonal $q = k + d$.

76. Diagonal Hardinian Arrays: A Fixed-Parameter Asymptotic Formula

[enumerative-combinatorics/hardinian-array-diagonal-asymptotics/article.pdf](#)

from [hardinian_arrays_research.zip](#)

For every fixed $r \geq 0$, $H_r(n, n) \sim C_r 4^{rn} n^{-r(r-1)/2}$ with an explicit C_r in Gamma values. The formula confirms the constants tabulated by Dougherty-Bliss and Kauers (2024) but refutes their proposed restriction to powers of 2, 3 and π : $C_7 = 2^{18} 5^2 / (3^{45} \pi^3)$. Adds an exact finite determinant at fixed array size, an exact enumeration algorithm, and a limiting negative-binomial law for total deficits.

77. An explicit formula for lucky parking spots

[enumerative-combinatorics/lucky-parking-spots/article.pdf](#)

from [lucky_parking_spots_research.zip](#)

Proves Conjecture 13 of Butler, Hadaway, Lenius, Martens and Moats (JIS 29 (2026), Article 26.1.1): a closed form for $C(n, j)$, the number of parking functions in which *spot* j is lucky. Adds an exact reflection law, all fixed right-edge columns, a closed form for OEIS A374533, Poisson edge limits, uniform bulk convergence, polynomial reciprocity and asymptotic expansions.

78. A hexagonal-lattice proof of the A195806 conjecture

[enumerative-combinatorics/a195806-hexagonal-lattice/article.pdf](#)

from [A195806_hexagonal_proof.zip](#)

An elementary derivation of the six residue-class formulas and the constant-coefficient recurrence recorded as conjectural in OEIS A195806, starting from the original triangular-array constraints rather than from interpolation. The steps are an integral six-parameter bijection, four independent translation intervals, a dihedral chamber reduction, and an exact lattice count.

79. Leaves of large type in the numerical-semigroup tree

[enumerative-combinatorics/numerical-semigroup-leaf-types/article.pdf](#)

from [leaf_type_counterexample.zip](#)

Disproves Conjecture 4.4 of Chappelon, Ramírez Alfonsín and Stamate, *Numerical semigroup tree: type-representation* (arXiv:2507.15006v2), with an explicit genus-34 counterexample, and establishes a sharp asymptotic order for the type of leaves in that tree.

80. Strict growth and the Catalan limit for adjacency-bounded 132-avoiding permutations

[enumerative-combinatorics/adjacency-bounded-132-avoiders/article.pdf](#)

from [strict_growth_132.zip](#); batch 38: [Closing_Catalan_Growth_Gap.zip](#); batch 43:

[ProveIt_Largest_Jump_Research.zip](#); batch 52: [ProveIt_Macroscopic_Jump.zip](#),

[ProveIt_Macroscopic_Jump_Research.zip](#); batch 64: [ProveIt_Critical_Jump_Moments.zip](#); batch 68:

[ProveIt_Polynomial_Rarity.zip](#); batch 71: [132_Avoider_Crossover_Package.zip](#); batch 72:

[General_132_Avoider_Crossovers_Package.zip](#), [Uniform_132_Avoider_Spectral_Bridge_Package.zip](#),

[Relative_132_Avoider_Asymptotic_Package.zip](#), [Theta_132_Avoider_Boundary_Package.zip](#),

[Integer_Count_132_Avoider_Asymptotic_Package.zip](#), [Uniform_132_Avoider_Saddle_Sum_Package.zip](#),

[Growing_Macro_132_Avoider_Counts_Package.zip](#)

For 132-avoiding permutations with every adjacent difference at most m and growth constant α_m : proves $\alpha_{m+1} > \alpha_m$ for all $m \geq 1$ — the remaining strict-inequality part of Nadler’s Conjecture 2, which A. C. Li announced publicly on 8 September 2026, eleven days before Part I’s date (a dated note in Part I records the announcement; Part I’s proof does not use it, and it is not a retraction) — and the refinement $4 - \alpha_m = (2 \log m + 4 \log \log m + O(1))/m$, with $2 \log \pi - 4 \log 2 \leq \liminf E_m \leq \limsup E_m \leq 2 \log \pi$. Part II, added in batch 38, closes that window: a Catalan-weighted supersolution and a growing cutoff give $E_m \rightarrow 2 \log \pi$, with an all-orders inverse-log expansion. Its numerics at $m = 10^6$ (near 3.15) are far from the limit and are not evidence for it. The all- m component ordering is still not claimed. Part III, added in batch 43, describes the joint limit near the boundary $m = n - d$: the distance $n - M_n$ of the largest jump converges in total variation at rate $n^{-1/2}$ to an explicit algebraic law, uniformly in $d = o(n)$. Part IV, merged in batch 52 from two manuscripts, treats the bulk regime $n - m \asymp n$: $\sqrt{n} P(n - M_n > un) \rightarrow (3/\sqrt{\pi})(1 - 2u)/\sqrt{u(1 - u)}$ for $0 < u < 1/2$ (and $\rightarrow 0$ for $u \geq 1/2$), so for $1/2 < \theta < 1$ the fraction of 132-avoiders of length n with every adjacent difference at most $\lfloor \theta n \rfloor$ is asymptotic to $3(2\theta - 1)/\sqrt{\pi\theta(1 - \theta)n}$, and the equivalent $3/\sqrt{\pi(1 - \theta)n}$, which Part III declined to assert, is false. It also gives the exact constant of every moment $E(n - M_n)^p$, $p > 1/2$ (for instance $E(n - M_n) \sim 3\sqrt{n/\pi}$); for $\theta \leq 1/2$ it proves only $o(n^{-1/2})$ and polynomial lower bounds. Part V, added in batch 64, settles the critical moment, $E\sqrt{n - M_n} = \frac{3}{2\sqrt{\pi}} \log n + c_\star + o(1)$ with an explicit $c_\star \approx -0.9421$, and gives the first correction to every subcritical moment. Part VI, added in batch 68, gives the order of Part IV’s fraction for every fixed $0 < \theta < 1$: it is $\Theta(n^{-\lfloor 1/\theta \rfloor/2})$, with an order-crossover scale $n^{1-1/(2(q-1))}$ near each $\theta = 1/q$ (so the fraction is $\Theta(1/n)$ at $\theta = 1/2$). Part VII, added in batch 71, finds the first amplitude and crossover function: n times the fraction at $\lfloor n/2 + t\sqrt{n} \rfloor$ tends to $6 + 12t_+/\sqrt{\pi}$, locally uniformly in real t , so the fraction at $\lfloor n/2 \rfloor$ is asymptotic to $6/n$; and for $1/3 < \theta \leq 1/2$ the fraction at $\lfloor \theta n \rfloor$ is asymptotic to $I_3(\theta)/(2\pi n)$, with an explicit two-dimensional integral I_3 and $I_3(1/2) = 12\pi$. Part VIII, added in batch 72, finds every remaining fixed amplitude and reciprocal crossover: for $1/p < \theta < 1/(p - 1)$ the fraction at $\lfloor \theta n \rfloor$ is asymptotic to $(p + 1)/(2^p \pi^{(p-1)/2}) I_p(\theta) n^{-(p-1)/2}$, with I_p an explicit integral over a capped simplex, so every fixed $0 < \theta < 1$ now has an equivalent; and near every $\theta = 1/r$ the fraction, multiplied by $n^{r/2}$, tends to an explicit constant plus a multiple of t_+^{r-1} in the scaled window variable t (at $1/3$: $5/\sqrt{2\pi} + (729\sqrt{3}/(4\pi)) t_+^2$). Part IX, merged in batch 72 from six manuscripts, lets the cap grow with $n/m \rightarrow \infty$: a logarithmic bridge to the fixed-cap growth rate uniformly for $n \geq cm \log m$, one saddle sum on that whole range, explicit Gamma-sum equivalents near $n \asymp m \log m$ and for $1 \ll n/m \ll \log m$, and a theta-function modulation for $n/(m(\log m)^{3/2}) \rightarrow \infty$ that tends to one only when $n \gg m(\log m)^2$; the equivalents pass to the actual class when $n(\log m)^2/m^2 \rightarrow 0$.

81. The first pattern failure in 123-avoiding permutations

[enumerative-combinatorics/a273821-first-pattern-failure/article.pdf](#)

from batch 74: A273821_First_Pattern_Failure.zip

For a uniform 123-avoiding permutation of $[n]$, let K_n be the largest number of its largest entries whose induced permutation still avoids 132. Proves the bivariate generating function of the joint counts $T(n, k)$ marked conjectural in OEIS A273821, by inserting successive minima and telescoping a Catalan convolution to $T(n, k) = 2^{k-1} \binom{2n-2k}{n-k-1} - \binom{2n-k-1}{n-k-1}$; every column $k \geq 2$ is hypergeometric with growth rate 4, answering the entry's growth question. Also an explicit discrete limit law for K_n with exponentially weighted total-variation error $O(n^{-1})$, a crossover at $k \asymp \sqrt{n}$, two exactly separated exponential sectors for k proportional to n , and, under the weight y^{K_n} , a transition at $y = 2$ with critical law $K_n/n \rightarrow \text{Beta}(1, \frac{1}{2})$.

82. Throwback Dynamics with Repeated Entries

[enumerative-combinatorics/throwback-queue-dynamics/throwback_research.pdf](#)

from throwback_research.zip

For the queue process that removes the head and reinserts it at the zero-based position equal to its weight: every token recurs if and only if at most m input tokens have weight at most m , for every $m \geq 1$, with an explicit description of the recurrent tokens when that fails. Adds exact cycle periods and frequencies, a coprimality criterion for perfect mixing, Catalan and prime-parking counts for recurrent cores, and the largest labelled period of a core of size r .

83. Reciprocity and Matrix Duality for Preorder Polytopes

[enumerative-combinatorics/preorder-polytope-reciprocity/article.pdf](#)

from preorder_reciprocity.zip

Proves Conjectures 7.2, 4.2 and 4.4 of Athanasiadis and Chapoton, *Polytopes and posets associated to preorders* (arXiv:2605.26916v1), and the ordinary $q = 1$ case of their Conjecture 4.9 (not its full q -refinement). Two mechanisms: Ehrhart–Macdonald reciprocity after subtracting the all-ones vector, extending the double identity to independently weighted capacities; and duplication of both colour classes of the bipartite graph followed by Postnikov's lattice-point formula, giving the conjectured matrix transpose.

84. Support reciprocity and the q -zeta polynomial of preorder lattice-point posets

[enumerative-combinatorics/preorder-q-zeta-reciprocity/article.pdf](#)

from preorder_qzeta_reciprocity.zip + preorder_q_zeta_reciprocity.zip +
preorder_q_reciprocity_research.zip

Proves the full q -refinement of Athanasiadis–Chapoton Conjecture 4.9, equation (21): for a preorder τ on $[n]$, with P_τ the nonnegative integer vectors satisfying $a(I) \leq |I|$ on every order ideal and B_τ its maximal elements, $Z_q(P_\tau, [-1]_q) = (-1)^n \sum_{b \in B_\tau} q^{-|\text{supp}(b)|}$, together with the stronger identity at each exact support. The $q = 1$ case was already covered by the preorder-polytope reciprocity report and is cited, not reclaimed. *Merged from three reports that reached the same two theorems by three architecturally independent routes — an incidence inversion with Boolean localization, the same with a Hall-based privatization and a self-contained Euler identity, and a capacity-enumerator reciprocity with a transfer involution — and all three survive.*

85. An explicit shelling for preorder lattice-point posets

[enumerative-combinatorics/preorder-shellability/article.pdf](#)

from preorder_shellability_research.zip

An affirmative solution to Question 4.6 of Athanasiadis and Chapoton (arXiv:2605.26916v1, p. 13): an explicit pure shelling of the order complex of P_τ with its minimum removed, and a coning that shells the order complex of P_τ itself. Both complexes are therefore Cohen–Macaulay over the integers and over every field. A different question of that paper from the three other preorder entries.

86. Multiple-chain preorder polytopes

[enumerative-combinatorics/multiple-chain-exponential-formula/article.pdf](#)

from `multiple_chain_exponential_formula_research.zip` + `multiple_chains_research.zip`

Proves the exponential generating-function identity proposed in Section 9.5 of Athanasiadis and Chapoton (arXiv:2605.26916v1) for lattice points of multiple-chain preorder polytopes counted by number of nonzero coordinates, in a rectangular generalization with independent block width and height. Adds a duality corollary, an Ehrhart evaluation, a degree- 2^d root theorem with a quartic elimination certificate, two treatments of the γ -vector, a Stieltjes moment representation, the boundary constant in two forms, and a central limit theorem with exact variance. *Merged; the identity is proved twice, once through a subset encoding with a bounded symbol and once through slack meanders in a Laurent ring, and the eleven deliberate redundancies are listed in the text.*

87. Rationality and growth of stabilized permutation-weight series

[enumerative-combinatorics/stabilized-permutation-weights/article.pdf](#)

from `stabilized_permutation_weights.zip`

Studies the full stabilized generating functions of the permutation-weight q -Eulerian polynomials, beyond their initial connection with OEIS A256193: an all- d finite-spectrum theorem, rationality, a universal integer denominator, the exact radius of convergence and residue-class asymptotics. The two-descent sequence gets an explicit formula in powers of 3, powers of 2 and Fibonacci numbers.

88. A Reflexive Root-Polytope Model for Preorder h -Polynomials

[enumerative-combinatorics/preorder-root-polytopes/article.pdf](#)

from `preorder_root_polytopes.zip`; batch 39: `ProveIt_Height_Two_Preorder_Research.zip`; batch 42: `ProveIt_Cactus_Rigidity.zip`; batch 44: `ProveIt_Preorder_Gamma.zip`, `ProveIt_Preorder_Gamma_Research.zip`; batch 55: `ProveIt_Lorentzian_Support.zip`, `ProveIt_Support_Polynomials_Research.zip`, `ProveIt_ADE_Stability.zip`, `ProveIt_Matroid_Lifts.zip`; batch 63: `ProveIt_Marked_Tree_Stability.zip`; batch 64: `ProveIt_Two_Flow_Lattice_Sampling.zip`; batch 66: `ProveIt_Random_Tree_Stability.zip`; batch 68: `ProveIt_Hall_Bottlenecks.zip`; batch 70: `ProveIt_Matching_Rank_Research.zip`; batch 71: `ProveIt_Weighted_Rank_Three_Leaf_Compression.zip`

Proves Conjecture 5.1(c) of Athanasiadis and Chapoton (arXiv:2605.26916v1): for a preorder τ on $[n]$ the directed root polytope $C_\tau = \text{conv}(\{\pm e_i\} \cup \{e_i - e_j : j \leq_\tau i\})$ is reflexive with $h^*(C_\tau, t) = h_\tau(t)$, and any pulling triangulation of its boundary is combinatorially the boundary of a simplicial polytope with that h -polynomial. Palindromicity and unimodality, Conjecture 5.1(a,b), follow, as do the full g -theorem conditions. The higher-dimensional augmented bipartite identity and preorder duality are credited to Dai, Hou, Liu, Thawinrak and Wang (arXiv:2608.16037v2). Part II, added in batch 39, treats height-two preorders through matching polynomials: γ -positivity in height two (Conjecture 5.2 there), a family showing that real-rootedness (Conjecture 5.3) fails — its first counterexample is credited to a MathDB posting by Shivam Patel — and crown obstructions: an induced crown forces large minimal nonfaces in every lattice triangulation of C_τ , so the boundary complex is not flag when the crown has at least six elements. Flag realizability in general (Conjecture 5.1(d)) remains open. Part III, added in batch 42, shows that exactly the cacti admit a faithful unit-phase minor model, a property strictly stronger than real stability. Part IV, merged in batch 44 from two manuscripts, proves Conjecture 5.2 for every finite preorder: γ_k counts ordered pairs of disjoint k -sets joined by a τ -increasing bijection, and a reflexive relation gives a palindromic polynomial exactly when it is transitive. Part V, merged in batch 55 from two manuscripts, shows that for a bipartite graph with shores of sizes m and n every support enumerator divided by $\binom{\min(m,n)}{k}$ is log-concave (the unweighted inequality is Theorem A of Röhrle and Ulirsch, arXiv:2402.15317, carried through the support-counting bridge), so every h_τ and every height-two γ -vector is ultra-log-concave; it classifies the palindromic support polynomials of all bipartite graphs, gives FPRASEs for partition functions and for single coefficients, and shows exact counting #P-complete already in height two. Part VI, added in batch 55 with the stability sections of a further manuscript, shows that stability of the signed support polynomial is block-local, holds for the incidence-apex graph of a tree exactly when the tree is a finite or affine ADE tree, and is classified for books and bipartite multi-theta graphs; the books show it

strictly stronger than real-rootedness. Part VII, added in batch 55, shows that the transversal matroid of a preorder determines the preorder up to isomorphism and that height-two preorder matroids generate exactly the gammoids under minors. Part VIII, added in batch 63, classifies stability for one apex adjacent to an arbitrary set of vertices of a tree (a corollary of Part VI, also proved directly), with exact $O(n^5)$ enumeration, optimization and repair of the stable apex neighbourhoods, a sharp diameter window, and real-rootedness of their size-counting polynomial exactly for paths. Part IX, added in batch 64, gives explicit polynomial-time two-flow bijections between the demand vectors of a bipartite graph with prescribed support and the supplier sets matchable to it (the correspondence itself is Oh's), and with them an almost-uniform sampler of actual lattice points of every finite preorder polytope. Part X, added in batch 66, answers Part VIII's question for a uniform random labelled tree with every vertex marked independently with probability p : the probability that the marking is stable is asymptotic to $\frac{9}{8}\sqrt{2\pi}p^3n^{5/2}e^{-pn}$ whenever $p\sqrt{n} \rightarrow \infty$, an explicit entire function governs the window $p\sqrt{n} \rightarrow \lambda$, and below it stability is asymptotically the event of at most two marks; for a fixed host the diameter determines the probability up to a polynomial factor, separating the typical rate $\log(1-p)$ from the annealed rate $-p$. Part XI, added in batch 68, answers Part V's question on normalization by the matching number negatively for positive vertex weights on both shores: a graph with 30 vertices, 104 edges, matching number eight and weights in $\{1, 500\}$ has a weighted support polynomial that is not ultra-log-concave of order eight. A Hall-bottleneck family gives a rank-six phase diagram, the endpoint threshold for every core size and a lower bound asymptotic to $\frac{4}{3}\nu$ for any normalization depending on the matching number ν alone; the first rank-normalized inequality holds for every positively weighted graph, with a defect identity. Part XII, added in batch 70, answers the unit-weight version affirmatively for matching number at most three, by a sum-of-squares identity, and for a connected family of matching number four and disjoint unions, leaving unit weights from matching number four on and the one-shore version open. It also proves the first rank-normalized inequality for weighted directed relations, gives a 9×9 rank-three bimatroid obstruction, exact neighbourhood-type kernels with an exact support sampler, and for every finite preorder the first gamma-vector inequality $\gamma_1^2 \geq \frac{2d}{d-1}\gamma_2$, with d the gamma degree, which also bounds the cost of computing the gamma vector. Part XIII, added in batch 71, merges leaves with a common single neighbour into one leaf of summed weight without changing the support polynomial, and so proves normalization by the matching number for arbitrary positive weights on both shores whenever a minimum vertex cover is split $1 + (\nu - 1)$ between the shores, hence for every weighted graph of matching number at most three. That left a positively weighted counterexample matching number four, five or six, with every minimum vertex cover meeting both shores in at least two vertices, and at matching number four, with or without weights, only covers split $2 + 2$. Batch 72 continued this line in a separate report, *matching-rank-normalization* (log-concavity section), to which dated notes here now point: the unit-weight and one-shore versions fail (at matching numbers 3450 and 50), and the least matching number of a weighted failure is exactly six. Batch 73 opened a second separate report, *preorder-gamma-rank-ulc* (log-concavity section), which answers this report's Research question 96, rank ultra-log-concavity of the gamma polynomial at its actual degree, for unit activities through degree four and for independent activities through degree three. Conjectures 7.2, 4.2 and 4.4 of the same paper are proved in the preorder-polytope reciprocity report.

89. Largest Simplex Faces under Ordinal Sums

enumerative-combinatorics/

simplex-faces-of-order-and-chain-polytopes/Largest_Simplex_Faces.pdf

from Series_Parallel_Simplex_Faces_Research.zip

Answers the concluding open question of Freij-Hollanti and Lundström, *Simplex inequalities of order and chain polytopes of recursively defined posets* (Order 43 (2026), article 31). For $s_O(P)$ and $s_C(P)$, the largest dimensions of simplex *faces* of the order and chain polytopes: both are computable in $O(n)$ on any series-parallel decomposition, the order case carrying eight normalized endpoint states. For an ordinal sum of antichains, $s_O = r + \nu$ with ν a maximum b -matching of

the layer path under capacities $\min(a_i - 1, 2)$, and $s_C = r + \max(0, k - 1)$. The extremal gap is sharp: $0 \leq s_C - s_O \leq \lfloor (n - 2)/3 \rfloor$, attained for every $n \geq 2$.

90. Skew partitions, continued fractions and connected components

[enumerative-combinatorics/skew-partition-continued-fractions/article.pdf](#)
from [A225114_proof_bundle.zip](#)

Proves the repeated-power continued-fraction formula recorded as conjectural in OEIS A225114, which counts skew partition diagrams with no empty row or column. Unique concatenation gives $A(q) = 1/(1 - P(q))$ for the edge-connected diagrams, and a boundary-path encoding followed by a weighted two-coloured Motzkin-path decomposition gives $P(q) = q/(1 - 2q - q^3/(1 - 2q^2 - q^5/(1 - 2q^3 - \dots)))$, whence $A(q) = 1/(2 - C(q))$ with $C = 1 + P$. Adds a four-variable refinement, a formula by height, width and component count, a sharp truncation bound, exponential asymptotics and a central limit theorem for the number of components. **Explicitly not a novelty claim:** a prior-art check found Pasechnik’s MathOverflow sketch of ten September 2024, and the entry credits Kurkov; the report says so first and credits both.

91. From mesh avoidance to Catalan inflation

[enumerative-combinatorics/mesh-avoidance-catalan-inflation/article.pdf](#)
from [A289587_proof.zip](#), [OEIS_A289587_proof.zip](#)

Proves Scheuerle’s generating-function conjecture in OEIS A289587 for permutations avoiding 321 and the mesh pattern $(12, 174)$, and the equal count for $(12, 234)$ by inversion: with $D = x^4 - 2x^3 - 5x^2 - 2x + 1$, $P = x^4 + 4x^3 + 11x^2 + 10x + 3$ and $Q = x^2 + 5x + 3$, the generating function is $(P - Q\sqrt{D})/(8x(1 + x)^2)$. The headline structural result is the exact *occurrence count* $\text{occ} = \binom{f}{2} + \sum_U \binom{|U|}{2}$, of which avoidance is the case $\text{occ} = 0$. Merged from two reports that reached the identical generating function by the identical run-contraction substitution $u \mapsto u/(1 + xu)$; the merge keeps *both* refinements, because they refine the same sequence by different statistics — excedances and records — and are genuinely different theorems, agreeing only in total. The two local predicates they are built on disagree on 31 of the 321-avoiders of length at most six, the smallest being 12, and the merged article proves where they coincide rather than assuming it.

92. Prescribed coefficient-lattice minima

[enumerative-combinatorics/coefficient-lattice-minima/article.pdf](#)
from [coefficient_lattice_minima_research.zip](#)

Settles the unnumbered realization conjecture after Lemma 1 of O’Bryant, *On Nathanson’s Triangular Number Phenomenon* (arXiv:2506.20836): every nondecreasing list of even integers is the list of successive L^1 -minima of the coefficient lattice of some finite integer set. The witness is the mixed-radix set $A = \{0, 1, h_1, h_1h_2, \dots\}$, whose minima are exactly $(2h_1, \dots, 2h_r)$ with $r + 2$ distinct elements, and the radices may be given in any order.

93. Coefficientwise magic positivity through dimension ten

[enumerative-combinatorics/magic-positivity-parking-polytopes/article.pdf](#)
from [magic_positivity_research.zip](#)

A polytope is *magic positive* when its Ehrhart polynomial has nonnegative coefficients in the basis $t^j(t + 1)^{n-j}$. Proves, in exact arithmetic, that the generalized parking-function polytope $X_n(b)$ is magic positive for every dimension $3 \leq n \leq 10$ and *every* positive integer parameter vector, uniformly in b rather than on a sampled box — a partial solution of Conjecture 8.1 of Hill, Luo, Trinh and Vindas-Meléndez (arXiv:2607.15503v1). Dimension three is proved by printed algebraic formulas; the first-coefficient harmonic-number formula and the endpoint identities hold in every dimension. The conjecture in unrestricted dimension remains open.

94. Monotone blocks and reverse replies

[enumerative-combinatorics/reverse-reply-avoidance-game/article.pdf](#)
from [reverse_reply_solution.zip](#)

Proves Conjecture 2.9 of Ulfarsson (arXiv:2603.16004): for every $k \geq 3$ and every $n \geq R(k) = k^4 - 2k^3 - 3k^2 + 5k + 5$, the second player wins the normal-play permutation-avoidance game on

S_n with pattern length k by always replying with the reversal, so the Sprague–Grundy value is 0. The proof is elementary, with no computer search: a minimal-support lemma isolates one pattern inside a family built from singletons plus one inflatable monotone interval, and inflation restores the original length so the reply is legal. $R(k)$ is asserted sufficient, not minimal.

95. Nonintegral tropical degree from two horizontal segments

[enumerative-combinatorics/tropical-hilbert-nonintegral-degree/article.pdf](#)

from [tropical_hilbert_counterexample.zip](#)

Computes the strict-witness tropical Hilbert function of Grigoriev (arXiv:2404.06440v2) exactly for a two-segment set: $\text{TH}_{V_d}(k) = k + \min\{k, \lceil k/d \rceil\} + 1$. Taking $d = 3$ gives tropical degree $4/3$ — not an integer — and a quasipolynomial of least period 3, so eventual polynomiality fails. Every rational in $(1, 2]$ is realized, denominators are unbounded, and for $d = p/q > 1$ in lowest terms the least eventual period is exactly p . Both bounds are proved by hand.

96. Finite-prefix corrections for polyomino growth

[enumerative-combinatorics/polyomino-growth-finite-prefix-corrections/article.pdf](#)

from [ProveIt_Polyomino_Research.zip](#); batch 40: [ProveIt_Spectral_Saturation.zip](#)

Continues the formal project [Combinatorics/Polyominoes/KlarnerConstant](#), whose Lean-verified bound is $\lambda \leq 9047/2000 = 4.5235$ for Klarner’s constant. Proves a finite-prefix correction theorem for proper positive polynomial recurrences and applies it to Bui’s seventeen-neighbourhood system with an exact marked-occurrence table through size 18, giving the computer-assisted, unformalized bound $\lambda \leq 2249/500 = 4.498$ and $A_n \leq (927884613/10^9)(2249/500)^n$. A second theorem gives balanced integer-monomial divergence certificates, complete for productive, strongly connected, nonlinear systems, and yields $4.52349 \leq \mu_{\text{original}} \leq 4.5235$ for the original recurrence — a statement about the recurrence, not a lower bound on λ . At intake, a fresh build of the delivered enumerator (the same program, not an independent one) regenerated the size-18 table and matched it exactly. Part II, added in batch 40, shows where the method stops: exact spectral saturation in the irreducible critical case, a square-root law, and an all-prefix barrier $\mu_N \geq 4.3149$ for Bui’s map — a limit of this method, not a lower bound on λ and not an improvement on 4.498.

97. Three consecutive square-pyramidal numbers: exact Frobenius formulas

[enumerative-combinatorics/a069762-pyramidal-frobenius/article.pdf](#)

from batch 75: [OEIS_Pyramidal_Frobenius.zip](#)

For $s_n = n(n+1)(2n+1)/6$, gives the Frobenius number F_n of $\langle s_n, s_{n+1}, s_{n+2} \rangle$ (OEIS A069762, which had no formula) exactly: a quintic quasipolynomial of least eventual period six, $F_n = n^5/9 + 8n^4/9 + O(n^3)$, valid for every $n \geq 2$ apart from six exceptions, the last at $n = 23$. The proof is an explicit Apéry normal form after two gcd reductions. Also the exact genus, quasipolynomial for every gap-power sum, a triangular limit law $2(1-x)$ for a uniform gap divided by F_n , the minimal generating-function denominator with an order-18 eventual recurrence, and a convergent Puiseux inverse. Eventual quasipolynomiality itself is Rouné–Woods’ and Shen’s theorem; the claim is the explicit formula. The inversion instantiates the transseries volume’s Lagrange–Bürmann and staircase theorems, which the report cites.

98. A pole hierarchy for valley-monotone compositions

[enumerative-combinatorics/valley-monotone-bargraphs/article.pdf](#)

from batch 75: [OEIS_Valley_Monotone_Bargraphs.zip](#)

Proves Conjecture 2.3 of Flórez, Ramírez and Villamizar (JCTA 2024): the number $d(n)$ of compositions of n with weakly increasing interior valley heights (non-decreasing bargraphs by area) satisfies $d(n) = C\lambda^n(1 + O(\theta^n))$ with $\theta < 1$, $\lambda = 1.97642067794583\dots$ and $C = 0.60390151906189\dots$ certified by rational interval arithmetic; the digits printed in 2024 are the finite ratio at $n = 100$. The input is the stacks generating function of OEIS A001523; $d(n)$ itself has no verified OEIS number. Also infinitely many simple real poles increasing to 1, so the generating function is not D-finite, a finite-pole expansion to every radius below 1, a central limit theorem for bottom-level valleys, a terminal-component law and the asymptotics of the pole accumulation. Whether the real poles exhaust the spectrum is left open.

12 Generating functions and asymptotics

12.1 Coefficient counts of binary products

Three reports on counting coefficients of $\prod_i (1 + x^{G_i})$. The first two answer, independently, the universal-rationality part of Richard Stanley’s MathOverflow question 431075 (“A conjectured rational generating function”, 2022), which had no posted answer when checked on 19 September 2026. A third answer used the same mixed-base construction and the same enumerative argument as the first, and has been merged into it.

99. A Mixed-Base Counterexample to Stanley’s Rationality Question

generating-functions-and-asymptotics/binary-product-coefficient-counts/
mixed-base-counterexample/article.pdf

from stanley_mixed_base_counterexample.zip + Stanley_Rationality_Counterexample.zip

With $G_{2j+1} = 2^j$ and $G_{2j+2} = 3^j$ the exponent generating function is rational, while $\nu(2n) = 2^{n - \lfloor (n+1)\alpha \rfloor}$ for $\alpha = \log_3 2$ counts the coefficients equal to one in $\prod_{i \leq N} (1 + x^{G_i})$ and its generating function is not rational, not D-finite, not P-recursive, and has a natural boundary. Classifies rationality in the family interleaving a^j and q^j with digit polynomial $1 + y + \dots + y^{a-1}$ for all $2 \leq a < q$ — Stanley’s general $r = a - 1$ — as holding exactly when $q^u = a^v$. Adds a finite-quotient rigidity theorem, an abstract natural-boundary theorem for $\sum q^{\lfloor n\theta \rfloor} t^n$ at every irrational θ , the exact limit set of a_n/λ^n , and a streaming recurrence making the point that non-P-recursive does not mean hard to compute. *Merged; the enumerative half was the same argument in both originals, but the two routes to non-rationality differ and both are kept — an elementary finite-state reduction mod p , and the rigidity route that reaches the stronger transcendence conclusions.*

100. Rational exponents, nonrational coefficient counts

generating-functions-and-asymptotics/binary-product-coefficient-counts/
recursive-exponent-construction/research.pdf

from stanley_counterexample.zip

A different negative answer, from the recursion $a_0 = 2$, $a_1 = 1$, $a_n = a_{n-1} - 2a_{n-2}$, with $b_n = T_{n-1} + 2^n - 1 + a_n$, $T_n = T_{n-1} + b_n$, $G_{2n-1} = 2^{n-1}$ and $G_{2n} = b_n$: the counting series has a natural boundary on the unit circle, hence is neither rational, algebraic nor D-finite.

101. Coefficient residues in generalized Fibonacci products

generating-functions-and-asymptotics/binary-product-coefficient-counts/
kbonacci-coefficient-residues/article.pdf

from stanley_kbonacci_research.zip + quasifibonacci_parity_research.zip

Settles Conjectures 6.2 and 6.3 of Stanley, *Theorems and Conjectures on Some Rational Generating Functions* (arXiv:2101.02131v3, pp. 23–24). For k -bonacci seeds each exceeding the sum of their predecessors, the number of odd coefficients of $\prod_{i \leq n} (1 + x^{w_i})$ has generating function $(1 + 2z^k)/(1 - 2z + 2z^k - 2z^{k+1})$, independently of the seeds; the proof builds periodic signs of block length $k + 1$ and block product -1 and shows every signed finite product is flat. *Merged; the Conjecture 6.3 halves proved the same statement on the same domain, and the surviving report adds Conjecture 6.2 and the asymptotic analysis.*

12.2 Single-sequence OEIS studies

Fifty-nine reports on the growth or the closed form of a single OEIS sequence. One article counts the terms in the derivatives of three power towers — x^{x^x} , x^{x^2} and x^x — whose three parts reach three different answers by three different recurrences, and two reports added in batch 72 sit beside it: one proves the leading constant $\frac{1}{2}$ for x^{x^a} at every integer a , hence for x^x and x^{x^2} , and the other studies the x^{x^2} coefficients at unbounded logarithmic deficit. Two more reports share a Bell-scale saddle-point method on unrelated sequences. Shared technique is not shared theorem, and none of these duplicates another. Thirteen reports came in batch 73, one in batch 74, one in batch 75 and thirty-three in batch 77; they are the last forty-eight entries below, the batch-77 ones

grouped by the cluster that placed them (ascent sequences; iterated Bell numbers, cone walks and divisor weights; trees and networks; words, permutations, partitions and placements; moments, renewals and moving parameters). Four batch-73 reports merge two manuscripts that proved the same theorems independently on the same day (A139217, A022629, A215561, A279619): the shared theorems are printed once and the other proofs kept as marked second routes. Batch 75 brought a third and a fourth independent proof of the A022629 theorems, and a manuscript about two thirds of which re-proves results of the A215561 report; those duplicates are tabulated in the reports, not reprinted, and only their new results were added. Batch 77 did the same with a fifth independent proof of the A022629 theorems, a fourth of the A215561 ones, a third of the A279619 constant and a second of the A126764 corrections. Several of its new reports join a manuscript with addenda that nest it byte for byte; each paper is printed once. The four partition reports of batches 73–74 (A022629, A033552, A097356, A238016) study different products in different saddle regimes and share no proposition, nor do the batch-77 partition reports (A174065, A291698, A292507, A301746, A301981), which share at most a method with them; the A189281 and A330266 reports share a method but no theorem. Many of these manuscripts invert their asymptotics by Lambert W and rounding; those steps are instances of the inversion theorems of the transseries volume in *Analysis/Transseries*, which the reports cite rather than claim.

102. Bell-scale growth of OEIS A088714

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a088714-bell-scale-growth/article.pdf](#)

from A088714_asymptotic_growth.zip; batch 73: golden_moment_laws.zip; batch 76: A088714_Bell_Local_Estimates.zip

For the unique formal series with $A(x) = 1 + xA(x)^2A(xA(x))$, proves $\log a_n = n \log n - n \log \log n - n + O(n \log \log n / \log n)$, so $(e \log n / n) a_n^{1/n} \rightarrow 1$ and the ordinary generating function has radius of convergence zero while the exponential one is entire. The lower bound comes from a single explicit path through an exact triangular recurrence, the upper from a Stirling-positivity comparison with derivatives of Touchard polynomials. The multiplicative equivalent is *not* claimed, and the finer Bell-number form is labelled a conjecture with a sufficient ratio condition isolated. Part II, added in batch 73, treats A088714 and its companion A088713 as moment sequences: determinate moment measures, strict Hankel positivity, golden-ratio endpoint laws with exponents $(3 - \sqrt{5})/2$ and $(\sqrt{5} - 1)/2$, the ratio law $a_{n+1}/a_n \sim n/\log n$ and the far tail. Its re-proof of Part I's theorem is replaced by a pointer; the finer Bell conjecture stays open. Part III, added in batch 76 from a continuation manuscript: Part I's formal endpoint coefficient expansion, proved uniformly for $m \leq C \log n$; the one-sided correction $\liminf(r_n - n/W(n)) \geq 3/2$ for $r_n = a_n/a_{n-1}$ (the conjecture needs 2); and a reduction of the finer Bell conjecture to one scalar defect estimate $E_n = O(W(n)^q/n^2)$, sufficient and in some form necessary, with a tilted-variance criterion, the complementary-endpoint bound $5/(r_n r_{n-1})$ and exact numerics to $n = 1600$. Its re-derivation of Part I's summability step is a pointer; $W(n)$ enters only as the Bell saddle of the transseries volume, which it cites; the conjecture and its constant stay open.

103. The exponential growth constant of OEIS A158415

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a158415-growth-constant/article.pdf](#)

from A158415_asymptotic_growth.zip

For the number of distinct values expressible with n symbols from $E ::= 1 \mid \sqrt{E} \mid (E + E)$, proves a computable $\gamma > 1$ exists with $a(n)^{1/n} \rightarrow \gamma$ and $1.86204 < \gamma < 1.9208$, gives an effective characterisation as a supremum of algebraic finite-data bounds, and proves the boundary identity $\sum a(n)\gamma^{-n} = \gamma^2$, which rules out a pure $C\gamma^n$ prefactor. The rigorous lower bounds rest on a uniform injectivity theorem: any finite alphabet of representable values can serve as nested-radical digits, via an odd valuation at a prime above 2. No closed form for γ is claimed, and terms 22–28 get certified intervals rather than exact values.

104. Dyadic scaling rigidity of 3AP-free permutations

generating-functions-and-asymptotics/oeis-sequence-asymptotics/
a003407-dyadic-scaling-rigidity/article.pdf

from ProveIt_Dyadic_Scaling_Rigidity.zip (batch 40); batch 43: ProveIt_3AP_Nonholonomicity.zip

For $\theta(n)$, the number of permutations of $[n]$ with no three-term arithmetic progression (OEIS A003407), studies the dyadic scaling of $\theta(n)$: a continuous log-periodic growth profile with its exact fundamental period, rigidity of the profile under real rescaling, explicit losses and gains of the 2:1 gluing, and the empirical and sampling laws; the adjacent ratios $\theta(n+1)/\theta(n)$ have no limit. The parity recurrences and Sharma's Theorem 2.8, which the manuscript imports, are already proved in Lean in `Combinatorics/Ramsey` (the table $\theta(1..20)$ there rests on a `native_decide` certificate); the report's own theorems are not formalized. Part II, added in batch 43, proves θ is not P-recursive and its generating functions are not D-finite (the ordinary one is transcendental), using the published counts $\theta(64)$, $\theta(75)$, which nothing in ProveIt certifies; whether the generating function has a natural boundary stays open.

105. An exact coefficient formula for Ramanujan's theta function

generating-functions-and-asymptotics/oeis-sequence-asymptotics/
a260306-formula-proof/a260306_proof.pdf

from A260306_formula_proof.zip

Proves both finite formulas recorded in OEIS A260306, for every coefficient index, for the expansion $\theta(N) \sim \sum c_n N^{-n}$ defined by $\sum_{k < N} N^k/k! + \theta(N)N^N/N! = e^N/2$. The route is an exact Laplace-integral difference, an analytic change of variable with $F(\psi(w)) = w^2/2$, and Lagrange inversion reducing each c_n to one coefficient extraction, then a finite change of basis into the entry's double and triple sums. Includes the analytic error estimate the identification needs, rather than only checking the two sums agree, and identifies A065973 as the reduced denominators.

106. The maximal positive coefficient of a Hermite polynomial

generating-functions-and-asymptotics/oeis-sequence-asymptotics/
a277280-hermite-maximum/article.pdf

from A277280_asymptotics.zip

An exact closed form for OEIS A277280, the largest coefficient of the physicists' Hermite polynomial H_n with signs retained: with $k_n = \lfloor (2n+6 - \lfloor \sqrt{8n+21} \rfloor)/8 \rfloor$ and $d_n = n - 4k_n$, the exponent d_n is the unique maximizer and $a(n) = n! 2^{d_n} / ((2k_n)! d_n!)$. The distinction from the largest coefficient in absolute value is kept explicit throughout.

107. Asymptotics of OEIS A277364

generating-functions-and-asymptotics/oeis-sequence-asymptotics/
a277364-bell-asymptotics/A277364_asymptotics.pdf

from A277364_asymptotics.zip

For $a(n) = \sum_{k \leq n/2} S(n, k)$, partitions of an n -set into at most $\lfloor n/2 \rfloor$ blocks, proves $a(n) \sim B_n$ with an explicit first correction: writing $r = W_0(n)$, $a(n) = \exp\{n(r-1+1/r)-1\}(1+r)^{-1/2}(1-P(r)/n+O(r^2/n^2))$ for an explicit rational P . Also gives a parity-split equivalent for the omitted tail $B_n - a(n)$.

108. Term counts in the derivatives of power towers

generating-functions-and-asymptotics/oeis-sequence-asymptotics/
power-tower-derivative-term-counts/article.pdf

from A293239_research_report.zip, A290268_partial_results.zip,
A281434_exact_algorithm_and_cubic_growth.zip; batch 42: ProveIt_A290268_Finite_Certificates.zip;
batch 70: ProveIt_A290268_Depth3_Article.zip; batch 72:
ProveIt_Lehmer_Cmtet_Prime_Multiple_Nonvanishing.zip,
ProveIt_Lehmer_Cmtet_Stirling_Residues_and_Candidate_Lists.zip,
ProveIt_Lehmer_Cmtet_Uniform_Padic_Nonvanishing.zip, ProveIt_A290268_Depths_Five_Six.zip,
ProveIt_A290268_Count_Bounds.zip; batch 73: ProveIt_A290268_Eight_Depths.zip,
ProveIt_A290268_Depth7.zip

One article on three OEIS sequences that ask the same question of three functions: how many distinct monomials are in the fully collected n th derivative of x^x (A293239), x^{x^2} (A290268)

and x^{x^x} (A281434)? Part I proves once the apparatus all three had established separately — the canonical differential-polynomial form $f^{(n)} = fP_n$ or $fx^{-n}P_n$ with its first-order integer recurrence, the independence of logarithmic monomials that makes the count an invariant of f rather than of a differentiation algorithm, the identity $a(n) = |\text{supp } P_n|$, and the envelope bound. Two normalizations are needed, because neither clause covers all three in the frames the parts use. **None of the three headline questions is fully settled, and the article says so in its abstract:** for x^x the count reduces exactly to zeros of the first-kind Lehmer–Comtet triangle A008296 and $a(n) = \Theta(n^2)$, but the closed formula needs an isolated nonvanishing step — and separately the recurrence printed in the entry for $n > 6$ is *disproved* at $n = 8$; for x^{x^2} the conjectured expression is proved to be an upper bound and $a(n) = \Theta(n^2)$, but the article does not prove the constant $\frac{1}{2}$ (since 1 October 2026 the sibling report *power-tower-exponent-supports* does, for x^x as well); for x^{x^x} explicit bounds give $a(n) = \Theta(n^3)$ but not $a(n) \sim \frac{2}{3}n^3$, and there the obstruction is not a nonvanishing statement but a density one, since further zeros provably exist. A closing part sets the three side by side and records what they share beyond the framework: every cancellation any of them can prove is a parity zero of the same falling factorial. This is a thematic consolidation, not a deduplication: the three were checked and are not duplicates, and all fifty numbered results of the three originals are preserved unchanged. Batch 42 added to the $x^{(x^2)}$ part depth certificates that close the nonvanishing problem at logarithmic deficits 3 and 4 for every n and improve its lower bound on the count; the open region was then deficit at least 5. Where that manuscript re-derived results already in the report, it is printed as a second route. Batch 70 added a second proof of the deficit-three case (no zeros beyond the forced holes), by a harmonic sign law and positivity propagation in k with its own finite certificate; that sign law proves a sign change the batch-42 data had only observed. Its manuscript’s claim that deficit three was still open is corrected in the text. Batch 72 added three manuscripts to the x^x part: Lehmer–Comtet nonvanishing on the prime-multiple offsets $\mu(p-1)$, $p > \mu$, under explicit conditions (in full for $\mu = 2, 4$, and for $\mu = 3, 5$ at odd p), Stirling residues that leave one candidate zero on $(p-1)^2$, and exact p -adic valuations on two families; the closed formula stays open. It added two to the x^{x^2} part: deficits five and six are classified by exact moment seeds, so the open region was then deficit at least 7, every fixed deficit reduces to a singly exponential rectangle, and $a(n) \geq \frac{3}{8}n^2 + \frac{n}{2} \log n - \frac{n}{2} \log \log n - \frac{5}{4}n$ for $n \geq e^{10}$, still the best explicit bound. Re-derivations of earlier results are printed once and credited. The two sibling reports of the same batch, *power-tower-exponent-supports* and *a290268-unbounded-deficits*, are not part of this article; dated notes in it point to them. Batch 73 added two independent manuscripts to the x^{x^2} part: deficit seven is classified twice, by three-anchor certificates of 51,699,091 and 48,113,025 cells, and deficit eight once (552,465,284 cells); at both only the reflection holes vanish, so the open region is now deficit at least 9. Their re-derivations of the deficit-five and -six classification, the largest-root minorant and weaker cutoffs and count bounds are printed as second routes or remarks.

109. Derivative supports of x^{x^a}

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/power-tower-exponent-supports/article.pdf](#)

from batch 72: ProveIt_All_Integer_Exponent_Derivative_Supports.zip,
 ProveIt_A293239_Leading_Asymptotic.zip, ProveIt_A290268_Leading_Asymptotic.zip,
 ProveIt_Half_Exponent_Derivative_Supports.zip,
 ProveIt_Algebraic_Exponents_and_Cancellation_Frequencies.zip (with a second manuscript nested inside it), ProveIt_Irrational_Exponents_Effective_Cancellation_Bounds.zip

Let $A_a(N)$ count the nonzero monomials $x^{a\ell-N}(\log x)^k$ in the N th derivative of x^{x^a} divided by x^{x^a} . For every fixed positive integer a , proves $A_a(N) = N^2/2 + o(N^2)$, by saddle contours in the coordinate $\tanh(t/2)$, a convex-envelope coverage argument and an integer-triangle estimate; for $a = 1$ this is $a(N) \sim N^2/2$ for OEIS A293239 (density zero of the zeros of the Lehmer–Comtet triangle), and for $a = 2$ it is $\lim a(N)/N^2 = \frac{1}{2}$ for OEIS A290268, the constant the power-tower term-count report leaves open. The same leading asymptotic holds for $a = 1/2$, with two infinite families of cancellations off the reflection line. On a fixed factor defect r it proves finiteness of the cancellations off an explicit reflection line for rational $a \neq 1$ ($r \geq 4$) and for algebraic a of degree

D ($r \geq 4D + 2$), at most $(2\lfloor r/2 \rfloor)^2 - 1$ cancellation cells, effectively, for irrational a and none for transcendental a , and exact square-root counting laws for rational $1/2 < a < 1$ on defects two and three. Every asymptotic statement is ineffective. Merged from seven manuscripts: the integer theorem was proved three times, and the separate proofs at $a = 1$ and $a = 2$, more explicit at those exponents, are printed as second routes.

110. OEIS A290268 at unbounded logarithmic deficit

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a290268-unbounded-deficits/article.pdf](#)

from batch 72: ProveIt_A290268_Explicit_Finite_Ratio.zip, ProveIt_A290268_Airy_Fold.zip, ProveIt_A290268_Arithmetic_Phase.zip, ProveIt_A290268_Sparse_Cancellations.zip, ProveIt_A290268_Pointwise_Three_Halves.zip, ProveIt_A290268_Strict_Quadratic_Gain.zip, ProveIt_A290268_Direct_Contour.zip, ProveIt_A290268_Global_049_Bound.zip, ProveIt_A290268_Growing_Logarithmic_Degrees.zip (superseded, not printed)

Studies the integer coefficients $H_{d,k,q}$ behind the term count of the N th derivative of x^{x^2} when the logarithmic deficit d grows with N . A regular saddle branch ending at a fold $\kappa_c \approx 0.0281461$ carries a uniform phase law that proves eventual nonvanishing at every algebraic limiting ratio $k/d \rightarrow \kappa < \kappa_c$, with possible zeros only near explicit transcendental rays; an additive Airy law holds in the window $|k - \kappa_c d| = O(d^{1/3})$. Two-dimensional cosine laws with explicit phase curvatures give density one with $O(N^{5/3})$ exceptions on tail patches and positivity on the triangle $40k + 420q \leq 91d$; block congruences modulo primes, prime-unit families with exact valuations, all-depth central strips and an example showing that root geometry alone cannot prove the support conjecture complete it. The manuscripts' headline lower bounds $((3/8 + c)N^2, 3N^2/8 + CN^{3/2}, \liminf a(N)/N^2 \geq 28176/57247)$ are kept in a table as superseded by $a(N) = N^2/2 + o(N^2)$ from the companion report on derivative supports; the phase laws behind them are not implied by it. Merged from eight manuscripts with shared material printed once; a ninth is wholly implied by the base and is described, not printed. The support conjecture stays open.

111. Iterated sums and products from one

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a352969-correctness-and-growth/article.pdf](#)

from A352969_correctness_and_growth.zip

For $S_0 = \{1\}$, $S_{n+1} = (S_n + S_n) \cup (S_n \cdot S_n)$ and $a_n = |S_n|$ (OEIS A352969), proves correctness of the listed terms and the doubly exponential law $2^{2^{n-r_{n-4}}} \leq a_n \leq 2^{2^{n-1}}$, hence $\log \log a_n = n + O(\sqrt{\log n})$. The step is simultaneous, not repeated closure — a distinction the report is careful about.

112. Distinct third derivatives of a power tower

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a199085-exponentiation-derivative-values/article.pdf](#)

from A199085_generating_function.zip

For OEIS A199085, the number of distinct values taken by the third derivative at $x = 1$ over all fully parenthesized exponentiations of n copies of x , proves $a(1) = a(2) = 1$ and $a(n) = \lfloor (n^2 - 2)/3 \rfloor$ for $n \geq 3$, with the rational generating function $z/(1 - z) + z^3/(1 - z)^3 - z^7/((1 - z)^2(1 - z^3))$. The proof is for all n , not an extrapolation: every attainable pair of second and third derivatives is classified and a witness constructed for each admissible pair. Despite the shared vocabulary this is a different question from *power-tower-derivative-term-counts*, which fixes the function and varies the derivative order; here the order is fixed at three and the parenthesization varies.

113. Finite-fringe rigidity in congruence-restricted greedy dissociated sequences

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a139217-greedy-dissociated-fringes/article.pdf](#)

from batch 73: finite_fringe_oeis_A139217_A139218.zip + ProveIt_OEIS_A139217_A139218.zip

OEIS A139217 and A139218 choose greedily the least positive integer congruent to 1, respectively 2, modulo 3 that keeps the set dissociated (all subset sums distinct). After three terms the signed subset-sum difference set is an interval with a fixed finite fringe of holes, $\{3\}$ for A139217

and $\{1, 3, 6, 11\}$ for A139218, and a general propagation theorem keeps it so forever. Hence exact closed forms, for instance $a(n) = \frac{3}{4}2^n + 1$ for $n \equiv 0, 1$ and $\frac{3}{4}2^n - 2$ for $n \equiv 2 \pmod{3}$ for A139217 ($n \geq 3$), rational generating functions and inverse expansions; both satisfy $a(n) = a(n-1) + a(n-2) + 2a(n-3)$, for A139217 only from $n = 6$, so the entry's range " $n > 4$ " fails at $n = 5$. *Merged from two manuscripts that proved the same theorems independently on the same day; each shared theorem is printed once and credited to both, the second proof kept as a second route, and the cross-identities, remainder bound and tables that only one has are added.*

114. Five OEIS asymptotic conjectures for distinct-partition norms

generating-functions-and-asymptotics/oeis-sequence-asymptotics/
a022629-distinct-partition-norms/article.pdf

from batch 73: OEIS_Distinct_Partition_Norms.zip + A022629_Research_Package.zip; batch 75: A022629_research_package.zip, OEIS_Norm_Weighted_Partitions.zip; batch 77: oeis-fixed-power-asymptotics-result.zip

For $a_\alpha(n) = \sum N(\lambda)^\alpha$ over partitions λ of n into distinct parts, $N(\lambda)$ the product of the parts, proves the logarithmic growth formulas marked conjectural in A022629, A092484 and A265840–A265842, and with $K = \sqrt{2n}$, $r = \log K$, $c = \pi^2/(6\alpha^2)$ the expansion $\log a_\alpha(n) = \alpha K(r - 1 + c/r + c/r^2 + \dots)$ to every order; a separate local-limit argument gives multiplicatively accurate saddle asymptotics with all Edgeworth orders, since exponentiating a truncated logarithm does not. Also a Lambert- W inverse, a Gumbel law for the largest part, asymptotic to $(1+1/\alpha)\sqrt{2n}$, and the modulus of the first Fourier resonances. *Merged from two independent manuscripts with identical coefficients (one treats A022629, the other's $\alpha = 1$): shared results printed once, with the second proofs, an explicit differential-operator derivation and the results only the A022629 manuscript has.* Two batch-75 manuscripts, the third and fourth independent proofs of these theorems with identical coefficients, are recorded in one table rather than reprinted; their new results form Sections 15–19: sharp signed cumulant scales, a saddle remainder of every fixed order with first correction $\sim -3/(8\alpha Kr)$, localization of the first index attaining a large value to within $1 + o(1)$ of an explicit inverse, eventual strict log-concavity and fixed-degree Jensen hyperbolicity; and the convolution powers $\prod(1 + k^\alpha q^k)^\mu$ (A297321, A022630, A022631) with μ -independent coefficients, two further inverse coefficients and an exact check of the eighth forward coefficient. A batch-77 manuscript is a fifth independent proof of the core theorems (all logarithmic orders for $\alpha = 1$), again tabulated, not reprinted. Its new results form Section 20: the slot family $\prod(1 + k^\alpha q^k)^{k^\beta}$, $\alpha > 0$, $\beta \geq 0$ an integer, with two logarithmic orders, an exact-saddle expansion to every fixed order, a threshold inverse and strict monotonicity, which for $\alpha = \beta = 1$ proves Kotěšovec's conjectured leading term for OEIS A266891; and resummed "thermal" forms of the logarithm and the inverse, no sharper than the two-term truncations.

115. Three exceptions before every square

generating-functions-and-asymptotics/oeis-sequence-asymptotics/
a097356-sqrt-restricted-partitions/article.pdf

from batch 73: OEIS_SquareRoot_Partition_Research.zip; batch 74: A097356_Sharp_Envelopes_Transseries_Inversion.zip

OEIS A097356 counts partitions of N into parts at most $\lfloor \sqrt{N} \rfloor$. Proves that the posted lower constant 0.0881548837... is exactly the limit inferior, while the literal eventual lower bound fails at precisely three indices in every square block, the three integers before the next square. The engine is a uniform all-orders expansion of $p_m(\alpha m^2 + \sigma m + \tau)$ with a finite Gaussian coefficient operator; for A097356 it carries a phase factor $\rho^{\lfloor \sqrt{N} \rfloor}$, giving the cluster set, limiting distribution, summatory law and an inverse with $O(1)$ index error once a piecewise-linear phase map is added to the Lambert- W_{-1} core. A206227 and A206240 are covered too; the leading asymptotic is Szekeres's. A second derivation from batch 74, written independently with equal constants, is merged: its proofs are second routes, and its own results (exact shell-polynomial degrees, second-order envelope constants, the exact phase loss, a Fourier identity and an $O(N^{-1/2})$ rate for the limiting law) form an added section.

116. Partitions into Catalan numbers

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a033552-catalan-partitions/article.pdf](#)

from batch 73: Catalan_Partitions_Research.zip

For OEIS A033552, partitions whose parts are distinct Catalan numbers, a complete expansion in inverse powers of a logarithmic size parameter with a nonconstant periodic amplitude, an explicit first correction and an inverse expansion for the threshold at which $p(n)$ reaches a given value, from a contour-controlled saddle estimate, a Barnes G product identity and a Fourier series in $\Gamma(s)\zeta(1+s)$; also a phase-dependent limit law for the largest part index. The entry poses no asymptotic conjecture, and the methods are classical.

117. Central Poupard numbers

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a125054-central-poupard-numbers/article.pdf](#)

from batch 73: Central_Poupard_OEIS.zip

For $A_n(c) = \sum_k \binom{n}{k} c^{n-k} T_k$ with T_k the tangent numbers, $A_n(1)$ is OEIS A125054. Proves the continued fraction recorded there as a conjecture in December 2025 and $A_n(1) \equiv 3 \pmod{9}$ for $n \geq 1$, by formal continued fractions. Analytically: an exact convergent decomposition into odd-pole exponential sectors, a uniform all-order endpoint-saddle expansion for $c = \tau n^2$ with a first-order transition at $\tau_* \approx 0.2624665$ and a displaced finite-size window, mixture and Gaussian laws, located polynomial zeros, the same transition for moments of any density with tail $Cy^\beta e^{-\alpha y}$, and a Lambert- W inverse.

118. Theta laws for high-power ballot sums

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a357825-theta-ballot-power-sums/article.pdf](#)

from batch 73: OEIS_Theta_Ballot_Sums.zip

For $a_n = \sum_h B(n, h)^n$, with $B(n, h)$ the ballot paths of length n ending at height h (OEIS A357825), the normalized ratio whose non-convergence the entry records is a Jacobi theta function at a drifting parity-sensitive phase times $e^{-5/6}$. Proves a uniform all-orders expansion in theta derivatives, the exact lower and upper limiting constants, the cluster interval and phase law, ratio and Turán asymptotics and eventual strict log-convexity; the multiset counterpart differs by an explicit exponential sector of relative order $n^{3/2-n}$, and for $\sum_h B(n, h)^k$ transitions occur at $k \asymp n$ and $k \asymp n^{3/2}$. Bala's supercongruence stays open.

119. Fixed-composition excursions

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a215561-fixed-composition-excursions/article.pdf](#)

from batch 73: oeis_balanced_excursions.zip + late_growing_excursions.zip; batch 75: OEIS_Fixed_Content_Ballot_Words.zip; batch 77: oeis-ballot-asymptotics-20261001.zip

For words with n copies of each of r letters whose every prefix sum is at most its average (OEIS A215561), proves the fixed-row conjecture $A_r(n) \sim \kappa_r (rn)! / (rn (n!)^r)$ with κ_r a positive algebraic number given by a finite kernel-root product, a complete expansion in inverse powers of n , D-finiteness of every row with a transcendental ordinary generating function for $r \geq 3$ (the specific A215570 recurrence is proved in Part IV, batch 75), a contact law and a Lambert- W inversion. Part II holds the second manuscript's own results: the leading term for every finite labelled step set, a second all-orders route, second-order terms for $r = 2, 3$, a giant primitive component, eventual log-convexity and a rounding bound. *Merged from two manuscripts that proved the main theorems independently, with constants agreeing to the last digit; shared results printed once and credited to both, the different routes kept as second proofs.* Part IV, added in batch 75 from a third manuscript, proves Kauers–Koutschan's Conjecture 15, the order-three recurrence for A215570, by a positive Legendre-coefficient formula and a rational telescoping certificate printed in full, and computes the coefficients $\alpha_{5,2}, \dots, \alpha_{5,8}$ of the five-letter row exactly. About two thirds of that manuscript re-proves results of Parts I–II, its row theorem only with relative error $O(n^{-1/5})$; those results are listed in one comparison table, not reprinted. Part V, added in batch 77 from a

fourth manuscript written without knowledge of this report, re-proves the main theorems with the same constants, sextic, Hessian and $\alpha_{5,1}$; these are tabulated, not reprinted. It adds the first correction $\alpha_{4,1} = -\frac{37}{200}\varrho^3 + \frac{11}{40}\varrho^2 + \frac{13}{40}\varrho - \frac{259}{400} \approx -0.5097846849$ of the four-letter row A215562, with $\varrho = \varphi - \sqrt{\varphi}$ and φ the golden ratio, the Hessian H_4 , an operator-norm second route to the global-arc bound and a nested-logarithm inverse for every row; $\alpha_{4,2}$, $\alpha_{4,3}$ and the rows $r = 6, 7, 8$ stay open.

120. An exact gamma constant for an Apéry-like sequence

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a279619-level-seven-gamma-constant/article.pdf](#)

from batch 73: A279619_Gamma_Asymptotics.zip + A279619_exact_asymptotics.zip; batch 77: level7-asymptotics-replay.zip

Proves the asymptotic displayed in OEIS A279619 with the constant identified: $a_n \sim \sqrt{3\pi} 27^n / (\Gamma(\frac{1}{7})\Gamma(\frac{2}{7})\Gamma(\frac{4}{7})n^{3/2})$, with $\sum a_n/27^n = 3\Gamma(\frac{1}{7})\Gamma(\frac{2}{7})\Gamma(\frac{4}{7})/(8\pi^2)$, a second exact sum at $z = 1/125$, rational corrections to every fixed order with error bounds, ratio, root and critical-tail expansions, and a continued-fraction companion with a recessive alternating solution. The Lucas-congruence conjecture of the entry is not established. *Merged from two manuscripts that proved the constant independently on the same day by different routes — a hypergeometric pullback with Ramanujan’s level-seven series, and an eta product at $i/\sqrt{7}$ evaluated by Chowla–Selberg — both kept.* A third manuscript, added in batch 77, proves the constant once more; its route is printed as Section 16: the Fricke involution of the weight-one theta series gives the singular coefficient directly at $\tau = i/\sqrt{7}$, the level-seven case of Guillera–Zudilin’s modular limit formula, and an Epstein zeta function gives the eta value. It adds no theorem, but an explicit generator for the corrections of every convolution power, with coefficients in a polynomial ring over $\mathbb{Q}$, and rounding brackets for every fixed weight, whose inversion cites the transseries volume; its claim to answer O’Brien’s Question 10.2 is not carried.

121. Spectral collisions and boundary asymptotics for zig-zag Eulerian numbers

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a205497-zigzag-eulerian-spectra/article.pdf](#)

from batch 73: zigzag_oeis_research.zip

For the descent counts $z(n, k)$ of linear extensions of a fence (OEIS A205497), determines the exact reduced denominator of every fixed-column generating function; its degree, the minimal recurrence order, is $r_k = \frac{40}{27\pi^2}(k+1)^3 + O(k^2 \log k)$, about $80/(9\pi^2)$ of the usual unreduced order. Also a uniform relative-error bound, a Gumbel crossover $\exp(-e^{-s})$ when $n/(k + \frac{3}{2}) - \log n \rightarrow s$, and strict log-concavity near both edges for $k \leq cn/\log n$; full-row log-concavity and real-rootedness are not proved. Two OEIS presentation corrections.

122. Winding sectors and the four-displacement restricted-permutation problem

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a000382-winding-correction/article.pdf](#)

from batch 73: ProveIt_A000382_winding_correction.zip

For the permutations of $\mathbb{Z}/n$ with every displacement in $\{0, 1, 2, 3\}$, a bijective proof by winding number of $\Phi_n = 2(L_n + 1)$, L_n the Tribonacci–Lucas numbers; the formula itself is already printed in OEIS A004306. OEIS identifies Φ_n with A000496 and with four times A000382, which first disagree at $n = 8$: the recurrence marked conjectural in A000382 is proved, but for the sequence produced by Mendelsohn’s 1961 formula, not for the permutation count, and the exact difference has generating function $z^4/(1 - 2z + z^4)$. Also an inverse transseries.

123. Fixed-multiplicity shuffles and balanced Smirnov words

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a330266-balanced-smirnov-poisson/article.pdf](#)

from batch 73: ProveIt_Balanced_Smirnov_OEIS.zip; batch 77: balanced-smirnov-repair-reproducibility.zip

Proves Kirgizov’s conjecture in OEIS A330266 for every fixed $k \geq 2$: the fraction of shuffles of n ranks of k cards with no two equal neighbours tends to $e^{-(k-1)}$, and the number of equal

adjacencies is asymptotically $\text{Poisson}(k-1)$, with an exact Laguerre-kernel generating integral and explicit corrections, for $k = 4$ $e^{-3}(1 - \frac{9}{8n} - \frac{3}{128n^2} + \frac{189}{1024n^3} + O(n^{-4}))$. Part I marks one tail estimate, on which the uniform expansion and the corrections rest, as a sketch; the limit law does not depend on it. Part II, added in batch 77 from a second manuscript, proves it for every fixed k : the factorial-moment domination $E(X)_M \leq (k-1)^M$, an explicit uniform remainder with a Touchard-polynomial constant, an all-orders expansion in integer powers, a degree bound giving factorial cumulants $O(n^{1-r})$, and a sharper inverse along the count sequence, which cites the transseries volume. Every coefficient Part I displays is re-derived unchanged and cited, not reprinted; the second manuscript's contraction, ratio and local derivations are kept as second proofs.

124. Unconditional all-orders asymptotics for two OEIS permutation sequences

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a189281-path-forest-expansions/article.pdf](#)

from batch 73: `oeis_path_forests.zip`; batch 74: `ProveIt_Fixed_Gap_Asymptotics.zip`

OEIS A189281 and A110128 count permutations avoiding a prescribed signed or absolute difference between entries two positions apart. Proves the displayed higher expansions without the conjectured recurrences, through a stabilization theorem for tilings of long path forests: for any fixed offsets the violation count has an all-orders expansion $e^{\theta u}$ times polynomials, with a finite exact formula for every coefficient, extended through order sixteen. The recurrences and the integrality suggested by A189281 are not claimed; a rational-function conjecture is proposed. Part II, added in batch 74 from a manuscript that re-derives Part I's theorems independently (its coefficients through order twelve agree and are printed only as a check), adds shape-uniform remainders, comparison bounds with an $n = 12$ example, a closed $c_3(r, s)$ and integrality for oriented one-path families.

125. Two Fourier peaks and a moving lattice maximum

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a039831-two-fourier-peaks/article.pdf](#)

from batch 73: `A039831_Two_Fourier_Peaks.zip`

Proves Kotešovec's 2023 conjecture $M_n \sim 3(n/e)^n$ in OEIS A039831, for the largest coefficient of $\prod_{j \leq n} (1 + q + q^3 + \dots + q^{2j-1})$, with explicit corrections through relative order n^{-4} : rounding produces a log-periodic term at order n^{-3} , and a second Fourier neighbourhood at $q = -1$ a parity term at order n^{-4} , which moves the maximizing exponent infinitely often. Also a zero-multiplicity law under Bernoulli smoothing and a Lambert- W inverse. A039824 is not resolved.

126. The cubic boundary for restricted partitions

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a238016-restricted-partitions-cubic-boundary/article.pdf](#)

from batch 74: `OEIS_Restricted_Partitions_Cubic_Boundary.zip`

For partitions of N into parts at most m , proves $p_m(N) \sim N^{m-1}/(m!(m-1)!)$ if and only if $N/m^3 \rightarrow \infty$, a condition still called conjectural in OEIS A238016 and A258670 (sufficiency is Erdős-Lehner's; necessity is new here), with ratio $e^{1/(4c)}$ at $N \sim cm^3$, a uniform centred expansion for $N \geq c_0 m^3$, five corrections for A238608, exponential sectors at $N = q^m$, inverse charts and a Poisson law for equal adjacent parts. The cubic scale is the $\alpha \rightarrow \infty$ end of the A097356 report's saddle question, not an answer to it.

127. L-convex polyominoes and a colored-partition comparison

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a126764-lconvex-polyominoes/article.pdf](#)

from batch 75: `A126764_Lconvex_Research.zip`; batch 77: `lconvex-asymptotics-reproducibility.zip`

Proves the Guttman-Kotešovec asymptotic recorded as a conjecture in OEIS A126764 for fixed L-convex polyominoes of area n , $a_n \sim \frac{13\sqrt{2}}{768} n^{-3/2} e^{\pi\sqrt{13n/6}}$, with three correction terms in closed form and relative error $O(n^{-7/4})$. The leading term and first correction already follow from a squeeze $0 \leq p_n - 4a_n \leq 3(p_n - 2p_{n-1} + p_{n-2})$ ($n \geq 3$) against the coefficients p_n of an eta quotient counting coloured partitions; a finite-cut identity and a full-circle estimate give the other

two. Also an all-orders radial expansion of the generating function at $q = e^{-t}$ (not a coefficient expansion) and a W_{-1} inverse, an instance of the transseries volume's Lambert and staircase theorems. The area generating function is a cited input; Part I's manuscript metadata says it was prepared with ChatGPT. Part II, added in batch 77 from an independent second manuscript, proves the all-orders coefficient expansion that Part I left open: from an exact decomposition $A = PB^2 + R$, with P Part I's eta quotient, B a partial theta function and R exponentially smaller, $a_n = \frac{13\sqrt{2}}{768}N^{-3/2}e^{\pi\sqrt{13N/6}}(\sum_{r < M} c_r N^{-r/2} + O(N^{-M/2}))$ with $N = n - 1/6$, for every M , re-proving Part I's three corrections. Also eventual strict log-concavity, monotonicity by an injection, and a shifted W_{-1} inverse with integer brackets of every order, again an instance of the transseries volume; Part I's boundary series equals B . Part II states that Guttmann and Kotěšovec print the reciprocal of the amplitude, which the report does not check.

128. The logarithmic deficit of 120-avoiding ascent sequences

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a202061-ascent-120-deficit/article.pdf](#)

from batch 77: [oeis-a202061-report.zip](#) + [oeis-a202061-sharp-deficit.zip](#) + [oeis-a202061-second-order-report.zip](#) + [oeis-a202061-third-order-report.zip](#) + [oeis-a202061-allorders-report.zip](#)

For ascent sequences avoiding 120 (OEIS A202061), with $\mu \approx 7.2959$ the largest root of $\mu^3 - 8\mu^2 + 5\mu + 1$, $D_n = n \log \mu - \log a_n$, $F = n^{1/3}(\log n)^{2/3}$ and $L = \log n$, five Parts prove, each refining the last: $D_n = \Theta(F)$ from an exact positive height-walk representation, which excludes every equivalent $C_0 \mu^n e^{-dn^\gamma} n^g$ with $\gamma \geq 1/3$, among them the stretch $n^{3/8}$ that Conway, Conway, Elvey Price and Guttmann estimated numerically, and that the generating function is not D-finite (Part I); $D_n/F \rightarrow C_\star \approx 2.2326$, explicit (II); the term $(7/3)C_\star F \log L/L$ (III); the next constant $\kappa \approx -1.9401$, which proves Part III's conjecture (IV); and, for every fixed J , an expansion of $D_n/(C_\star F)$ in powers of $1/L$ with universal polynomials $P_J(\kappa + (7/3) \log L)$ (V); each with its inverse. No multiplicative equivalent, amplitude or effective threshold is claimed, priority is not established, and the inverses are instances of the transseries volume. The A202062 report gives a second route to the growth rate μ . *Merged from five manuscripts whose archives nest their predecessors byte for byte; each paper is printed once, in chain order, and one manuscript's restatements of Part I are cited.*

129. Ascent sequences avoiding 000

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a202058-ascent-000-growth/article.pdf](#)

from batch 77: [oeis-a202058-report.zip](#) + [oeis-a202058-fine-report.zip](#) + [a202058-lognormal-report.zip](#) + [a202058-ratio-report.zip](#)

For ascent sequences avoiding 000 (OEIS A202058), $T = 3\pi^2/8$ and $c_n = a_n T^n / n!$, Part I proves the factorial growth constant $(a_n / n!)^{1/n} \rightarrow 8/(3\pi^2)$ conjectured by Conway, Conway, Elvey Price and Guttmann, through the exact radius T of the exponential generating function, with $\log c_n = O(n^{9/13}(\log n)^3)$. Part II proves $-O(n^{2/3}(\log n)^{5/3}) \leq \log c_n \leq O((\log n)^2)$, which excludes a factor e^{cn^σ} with $c, \sigma > 0$, gives exact counts through 400, shows that $a_n / n!$ is not PF₃, and conjectures $\log c_n \sim (2/3)(\log n)^2$. Part III proves that law in limsup and cumulative form, with $\log A(t) \sim (2/3) \log^2(1/(T - t))$; Part IV proves the ratio limit $a_{n+1}/((n+1)a_n) \rightarrow 8/(3\pi^2)$ with error $O(n^{-1/4}(\log n)^{3/4})$ and the lower bound $\log c_n \geq -O(\sqrt{n}(\log n)^{3/2})$. The pointwise law, log-concavity and a multiplicative equivalent stay open; the Lambert- W inverses are instances of the transseries volume. *Merged from four manuscripts in chain order; Part III's re-proof of Part II's bounds is kept as a second route, Part IV's uncredited restatement of a Part II identity is credited, and statements a later Part supersedes carry dated notes.*

130. Exact enumeration of 201-avoiding ascent sequences

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a202062-ascent-201-enumeration/article.pdf](#)

from batch 77: [oeis-a202062-report.zip](#)

For ascent sequences avoiding 201 (OEIS A202062), proves the cubic generating function convec-

tured by Guttman and Kotěšovec, by an exact two-label generating tree whose two-catalytic kernel equation is solved by two rational invariants, one from an order-seven kernel orbit; the report finds their conjectured cubic equivalent to its own and a sign error in their printed final combination. For every M , $a_n = C\mu^n n^{-9/2}(\sum_{m \leq M} c_m n^{-m} + O(n^{-M-1}))$, with $\mu \approx 7.2959$ the largest root of $\mu^3 - 8\mu^2 + 5\mu + 1$, $C \approx 13.4300$ in closed form and $c_m \in \mathbb{Q}(\mu)$, the first three exact; also a W_{-1} inverse with ceiling brackets, an instance of the transseries volume's Lambert theorems. With Conway, Conway, Elvey Price and Guttman's equal-growth theorem it gives $\lim b_n^{1/n} = \mu$ for A202061, a second route to Part I of the A202061 report. Novelty rests on a scoped literature check, and the envelope constants are not explicit.

131. Asymptotic enumeration and inversion for A122399

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a122399-surjection-diagonal/article.pdf](#)

from batch 77: [oeis-a122399-research.zip](#)

For $a_n = \sum_k k^n k! S(n, k)$ (OEIS A122399), the diagonal of $1/(1 - ue^y(e^x - 1))$: an exact, absolutely convergent vertical-contour formula for the two-size numbers $A_{m,n}$ with a nonasymptotic tail bound; an all-orders saddle expansion uniform for m/n in compact subsets of $(0, \infty)$; and $a_n = CD^n(n!)^2 n^{-1/2}(1 - 0.0600744 \dots /n + \dots)$ through n^{-4} , with $D \approx 3.1610887$ (A317855) and the first correction in closed form. Also a controlled inverse with integer-threshold envelopes, whose initializer is an instance of the transseries volume's inverse-Gamma core; a central limit theorem for the number of occupied blocks; and $a_{n+\varphi(p^r)} \equiv a_n \pmod{p^r}$ for $n \geq r$, which proves Bala's conjecture that $p-1$ is a period modulo p ; no least period is claimed. The leading equivalent is Kotěšovec's (OEIS, 2018), the methods are standard saddle methods, and no global novelty is claimed.

132. Asymptotics of iterated Bell diagonals (OEIS A139383, A261280)

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a139383-iterated-bell-diagonals/article.pdf](#)

from batch 77: [oeis-iterated-bell-report-attribution-revised.zip](#) + [iterated-bell-proportional-addendum.zip](#) (the first edition [oeis-iterated-bell-report.zip](#) superseded, not printed)

For the iterated Bell numbers $H(n, m) = n! [z^n] (e^z - 1)^{\circ m}$, with OEIS A139383 = $H(n, n)$ and A261280 = $H(n, n+1)$, Part I proves the leading equivalents those entries label conjectural, and for each fixed integer k an expansion $H(n, n+k) = e^k C n^{2n-5/6} 2^{1-n} e^{-n} (1 + \sum_j P_{j,k} (\log n) n^{-j})$ to every fixed order, with an outward-rounded interval certificate that $2 < C < 4$ (the digits $C \approx 2.86540798$ are not certified), an exact coefficient generator, and smooth inverses with a safe integer bracket. Part II, from an addendum conditional on Part I, extends the expansion to proportional depth $m = \lambda n + \delta$, uniformly for λ in compact subsets of $(0, \infty)$, with an amplitude $D(\lambda)$, $D(1) = 2C$, proved strictly positive for every $\lambda > 0$, a floor-depth phase $e^{-\{\lambda n\}/\lambda}$, eventual monotonicity and inverses; the endpoint regimes stay open. Prellberg's 2002 bivariate formula and parabolic-coordinate method already give the leading equivalent: no priority is claimed for it, for the method or for the coefficients. The inverses are instances of the transseries volume. The superseded first edition differs only in that attribution; the check that Prellberg's broad printed theorem fails for $f(z) = z/(1-z)$ is shared with the Takeuchi report.

133. Corner polyhedra and Schnyder labelings: polynomial exponents and limiting amplitudes

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a377922-corner-polyhedra-schnyder/article.pdf](#)

from batch 77: [oeis-bimodal-cone-research.zip](#) + [oeis-bimodal-full-asymptotics.zip](#)

Fusy, Narmanli and Schaeffer encode corner polyhedra (OEIS A377922) and 3-connected Schnyder labelings (A377920) as quadrant walks with a two-state parity and conjecture equivalents $\kappa \mu^n n^{-\alpha}$ with irrational exponents (their Conjecture 25). Part I proves $p_n = \Theta((9/2)^n n^{-\alpha_P})$ and $s_n = \Theta((16/3)^n n^{-\alpha_S})$ with the conjectured $\alpha_P = 1 + \pi / \arccos(9/16)$ and $\alpha_S = 1 + \pi / \arccos(22/27)$, by a one-unit cycle buffer, a random-clock transfer and a Fourier local limit; since both exponents

are irrational, the generating functions of A377922, A377920 and A377921 are not D-finite; and first-threshold inverses with $O(1)$ error. Part II, from an addendum, proves the full equivalents with finite positive amplitudes κ_P, κ_S (completing Conjecture 25), the coefficientwise equivalent $\tilde{s}_n \sim (16/19)^3 \kappa_S (16/3)^n n^{-\alpha_S}$ for A377921, and a Lambert- W_{-1} inverse with a vanishing-width ceiling bracket, an instance of the transseries volume. No digits of κ_P, κ_S , no rates and no correction terms are claimed; A377923 is outside both theorems; the iid cone theorems are Denisov–Wachtel’s, and neither manuscript claims priority.

134. Precise asymptotics of the powered Catalan numbers (OEIS A113227)

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a113227-powered-catalan/article.pdf](#)

from batch 77: [oeis-powered-catalan-report.zip](#) + [oeis-powered-catalan-fourier-addendum.zip](#)

The powered Catalan numbers a_n (OEIS A113227) count permutations avoiding the vincular pattern 1-23-4. From the known continued fraction Part I derives a positive discrete moment representation $a_n = \sum_{\lambda} \omega_{\lambda} \lambda^n$ whose transform is a Bessel quotient, locates its large spectral nodes, and proves $a_n \sim e^{e-2} (2\pi)^{-1/2} r^{-3/2} B_n(e)$, with B_n the Touchard polynomial and $r = n/W(n/e)$, answering the question of precise asymptotics that Elizalde left open in 2006; also an expansion to every fixed order with explicit c_1, c_2 , a canonical continuous inverse and an exponential-generating-function corollary. Part II, from an addendum, proves a zero-free complex wedge for the Bessel weight, replaces the spectral sum by a lattice sum with relative error $O(e^{-dn})$ for every $d < \log 2$, and splits a_n , for each fixed J , into $J + 1$ exact Fourier sectors, each with an all-orders expansion. No convergence, growing number of sectors, Borel sum or transseries is claimed, and the inverse is an instance of the transseries volume. The recurrence, models and continued fraction are prior art (Callan, Beaton et al., Gladkovskii); the sibling pattern 12-34 (A113226) has its own report.

135. Growing powers of Mahonian coefficients

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a380274-mahonian-growing-powers/article.pdf](#)

from batch 77: [oeis-mahonian-crossover-result.zip](#)

For $S_n(r) = \sum_k M_n(k) r^k$, with $M_n(k)$ the number of permutations of n with k inversions, proves that for every $r_n \rightarrow \infty$ the ratio $S_n(r_n)/C_n^{r_n}$, C_n the central Mahonian number, is asymptotic to a theta function $\Theta_{\delta}(r_n/V_n)$, where $V_n = n(n-1)(2n+5)/72$ is the inversion variance and $\delta \in \{0, \frac{1}{2}\}$ depends on $n \bmod 4$: a Gaussian integral below the critical scale $r \asymp n^3$, a discrete Gaussian partition function on it and $1 + 2\delta$ above it, with the matching limit laws for the common inversion count of colliding random permutations. Also an explicit fourth-order correction, uniform expansions to every fixed order on the transition scale, an unnormalized multiplicative equivalent, and a Lambert- W_0 index inverse on a fixed critical path, an instance of the transseries volume. The fixed-power equivalent with three corrections, including OEIS A380274 ($r = 3$) and A380275 ($r = 4$), is in Wang’s 2026 manuscript and is not claimed; priority for the growing-power extension is not established. A dated note relates the crossover to the critical theorem of the theta-ballot report, whose general lattice-peak question stays open.

136. Divisor-weighted partition products and assemblies (OEIS A301746 and A294363)

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a301746-divisor-weighted-asymptotics/article.pdf](#)

from batch 77: [oeis-divisor-square-asymptotics-result.zip](#) + [oeis-divisor-assembly-result.zip](#)

Two manuscripts on generating functions weighted by the divisor function $d(k)$. Part I treats OEIS A301746, the coefficients of $\prod_k (1 + q^k)^{d(k)^2}$, and proves the conjecture recorded there. From the fourth-order Mellin pole at $s = 1$ it obtains $\log a_n \sim \sqrt{n} (\log n)^{3/2} / (2\sqrt{6})$, Kotěšovec’s 2018 conjecture, with three further inverse-logarithmic corrections and the matching inverse. Part I also gives a relative expansion to every order in exact saddle cumulants, strict monotonicity and shrinking brackets for the integer threshold; the contributions of zeta zeros are left open. Part II treats A294363, $n![z^n] \exp(\sum_k d(k) z^k)$, whose Mellin kernel has a double pole: an explicit

multiplicative equivalent through a shifted Lambert- W saddle, an all-orders expansion, the OEIS logarithmic statement, and a factorial-core inverse. Its coefficients are rational in the Lambert parameter, and the inverse's absolute error tends to zero. The two Parts share a method template, also used by the batch-77 source of the A022629 report, but no theorem. No priority is claimed for the methods, and the Lambert, reversion and bracket steps are instances of the transseries volume.

137. Ascent sequences with bounded multiplicity (OEIS A294220)

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a294220-ascent-multiplicity-caps/article.pdf](#)

from batch 77: `oeis-bounded-multiplicity-family-report.zip + oeis-fixed-sector-addendum.zip`

For ascent sequences of length n in which every value occurs at most b times (the array OEIS A294220; cap three is A317784), Part I proves for every fixed $b \geq 3$ that $(a_n^{(b)}/n!)^{1/n} \rightarrow \mu_b = 1/T_b$, with $T_b = \int_0^\infty \log E_b(v)/(E_b(v) - 1) dv$ and E_b the truncated exponential, with error $O_b(n/\log n)$ in $\log a_n^{(b)}$; also a Lambert- W inverse in the length, the large-cap law $T_b = \pi^2/6 + (b+1)(\zeta(b+2) - 1) + O(b^{5/2}(4/9)^b)$ with explicit constants, and a W_{-1} inverse in the cap with counterexamples to naive rounding. The cap-two value $T_2 = 3\pi^2/8$, the constant $8/(3\pi^2)$ conjectured by Conway, Conway, Elvey Price and Guttmann, is taken from the A202058 report, not re-proved. Part II, from an addendum, writes $T_b - \pi^2/6$ as a convergent sum of positive Taylor sectors, proves that the remainder after any fixed number of sectors is asymptotic to the next sector, and expands each fixed sector to all orders in $1/b$; the remainder above is $\sim \frac{8}{9\sqrt{\pi}}\sqrt{b}(4/9)^b$. No prefactor or all-orders expansion in n , no joint regime $b = b(n)$ and no priority is claimed; the inverses are instances of the transseries volume.

138. Galled trees by gall count: the cube-root crossover, positive density and the near-maximal endpoint

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a399421-galled-trees/article.pdf](#)

from batch 77: `galled-density-reproducibility.zip + galled-crossover-reproducibility.zip + galled-endpoint-report.zip` (the earlier edition `galled-crossover-reproducibility (1).zip` superseded, not printed)

For $g(n, k)$, the rooted binary unlabeled simplex time-consistent galled trees with n leaves and k galls (OEIS A399421, row sums A397952), proves expansions to every fixed order in three disjoint regimes. Part I: for $k \asymp n^{1/3}$ the fixed- k asymptotic acquires the factor $\exp(-2\gamma_0 k^{3/2}/\sqrt{n})$, $\gamma_0 = 1.1300337163\dots$, with a series in $n^{-1/3}$ and an explicit first correction. Part II: for k/n in compact subsets of $(0, 1/2)$, a saddle expansion in $1/n$, a central limit theorem with mean $0.19741431n$ and variance $0.05548588n$, local and large-deviation laws and integer inverse brackets; it records that the inspected versions of Agranat-Tamir et al. print $\gamma \approx 0.3846$ where their defining equation gives $0.3835758866\dots$, the value credited on OEIS to Kotěšovec. Part III: for $n - 2k - 1 \asymp \sqrt{n}$, an expansion in $n^{-1/2}$ whose factor two comes from the parity of two moving square-root singularities. The generating function and the fixed- k and scalar leading terms are prior work, the cube-root mechanism is Fuchs and Hao Yu's for leaf-labeled classes, matching the regimes stays open, and the inverses are instances of the transseries volume. *Merged from three manuscripts of one workspace, one per regime, with no theorem proved twice; the common model is printed once.* Five heavy triangle files are not shipped; the README regenerates them.

139. Airy amplitudes for relaxed trees, compacted trees and finite-language automata

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a082161-airy-amplitudes/article.pdf](#)

from batch 77: `relaxed-binary-trees-reproducibility.zip + compacted-binary-trees-reproducibility.zip + fixed-arity-airy-package.zip + ternary-airy-amplitudes-result.zip + larger-alphabet-automata-report.zip + oeis-dfa-finite-language-report.zip`

Relaxed trees, compacted trees and minimal automata of finite languages satisfy one triangular recurrence family indexed by the arity k ; their growth orders are due to Elvey Price, Fang and

Wallner and to Ghosh Dastidar and Wallner, and the existence of a limiting amplitude was open. Proves positive amplitudes and Poincaré expansions to every fixed order in $n^{-1/3}$. Part I (A082161): $a_n = \gamma n! 4^n e^{3zn^{1/3}} n (1 + \sum_k c_k n^{-k/3})$, z the largest Airy zero, with three explicit logarithmic corrections. Part II (A254789): the same with $n^{3/4}$, exact positive renewal models, and cancellation of the $n^{-1/3}$ term in the logarithm of the compacted-to-relaxed ratio. Part III: every fixed $k \geq 3$ and all three models at scale $(n!)^{k-1} (k^k / (k-1)^{k-1})^n e^{\gamma n^{1/3}} n^{(2k-1)/3}$, with common first two corrections, first model-dependent defects of order $n^{-(k-2)}$, eventual decrease of the normalized ratios and a cancellation identity; its ternary section, the base proof, adds the third coefficient, and its comparison section bounds the automaton-to-relaxed ratio below by $\prod_{j=2}^n (1 - 1/(2j^{k-1})) \geq (2\sqrt{2}/\pi) \sin(\pi/\sqrt{2})$, uniformly in k and n . Part IV (A331120): binary automata, with 8^n and $n^{7/8}$. No amplitude is evaluated, the expansions are not uniform in k , the Lambert- W inverses are instances of the transseries volume, and the sources' claim that Ghosh Dastidar–Wallner use a false bound $U \leq q$ was not checked against the printed paper. *Merged from six manuscripts: the ternary one is the $k = 3$ case and base proof of the arity one, the comparison one the earliest form of its comparison step, and the automaton one repeats the compacted one's method and Jacobi appendix, printed once.*

140. Airy asymptotics for tree-child networks

generating-functions-and-asymptotics/oeis-sequence-asymptotics/
a213863-tree-child-networks/article.pdf

from batch 77: d-combining-asymptotics-reproducibility.zip +
tree-child-asymptotics-reproducibility.zip

For maximally reticulated d -combining tree-child networks and their word diagonal a_n (OEIS A213863 at $d = 2$), proves for every fixed $d \geq 2$ that $a_n = \gamma_d (n!)^{d-1} \Gamma^n e^{Bn^{1/3}} n^\rho (1 + \sum_r C_{d,r} n^{-r/3})$ to every finite order, with $\gamma_d > 0$ defined by convergent limits, $\Gamma = 4(d+1)^{d-1}/(d-1)!$, $B = 3z((d-1)/(d+1))^{2/3}$, z the largest Airy zero, and ρ explicit, by a contractive Jacobi reduction and polynomial–Airy profiles; ternary coefficients are given. Part II, for $d = 2$: explicit coefficients; every order for the total number of binary tree-child networks, $TC_n \sim (\sqrt{e} \gamma/12) (n!)^2 12^n e^{Bn^{1/3}} n^{-5/3}$, through the Pons–Batle identity proved by Lin, Liu, Liu, Liu and Xin; weighted all-orders expansions of the reticulation deficit around Poisson(1/2); an exact eventual total-variation identity with sharp rate $|B|n^{-2/3}/(48\sqrt{e})$; explicit inverses with integer brackets, instances of the transseries volume. For $d \geq 3$ the total count has only its leading term, and no amplitude is evaluated or certified. *Merged from two manuscripts with one proof skeleton: the binary manuscript's own proof of the case $d = 2$ is kept as a second route, and four identical displays are printed once.*

141. Historic tree asymptotics and a high-order obstruction

generating-functions-and-asymptotics/oeis-sequence-asymptotics/
a333497-historic-trees/article.pdf

from batch 77: historic-tree-report.zip

For reduced 5-historic trees (OEIS A333497), the exponential coefficients h_n of the solution of $H''' = H^2$, $H(0) = H'(0) = H''(0) = 1$, proves a unique positive blowup point ρ , $180^{1/4} \leq \rho \leq 60^{1/3}$, and an expansion of $h_n/(30(n+2)!\rho^{-n-3})$ to every fixed grade with exact Gamma ratios, oscillating in $\log n$ through the exponent $\lambda = (13 + i\sqrt{71})/2$: the relative error is $2|K|n^{-13/2} \cos(\arg K - \frac{1}{2}\sqrt{71} \log n) + O(n^{-15/2})$ with $K \neq 0$, so it changes sign infinitely often. The same for A336009 ($G'''' = G^2$), with the exact real-mode coefficient $-1/18052070400$ from a conserved energy. Main result: the factorial-pole asymptotic of Burghart and Wagner's Conjecture 1 fails for every ODE order $r \geq 30$ (conjecture parameter $m = r - 1 \geq 29$), through an unstable conjugate pair (an exact rational bridge for $30 \leq r \leq 37$), a sign-variation cone, strict sign-regularity and a generalized stable manifold. The orders $4 \leq m \leq 28$ and Conjecture 2 stay open, no replacement asymptotic is given, and the fitted constants are not certified; the manuscript also reads the journal's Table 1 as listing the radius under its reciprocal's label. The Lambert- W start and two-ceiling window of the inverse are instances of the transseries volume.

142. Asymptotic expansions of the Takeuchi numbers

generating-functions-and-asymptotics/oeis-sequence-asymptotics/
a000651-takeuchi-numbers/article.pdf

from batch 77: takeuchi-asymptotics-research.zip

For the Takeuchi numbers T_n (OEIS A000651) and $w = W(n)$, proves $\log T_n = e^w(w^2 - w + 1) + w^2/2 - \frac{1}{2}\log(1+w) + \log C + p_1(w)/n + p_2(w)/n^2 + O((1+w)^8/n^3)$ with explicit rational p_1, p_2 and $0 < C < \infty$, without assuming the leading equivalent, which is Prellberg's (announced 2002); then every fixed order, each p_j by a finite exact algorithm and one convergent quadrature, and an explicit $p_3 \sim w^4/72$, so $p_j = O(w^j)$ fails at $j = 3$. The transfer is a quantitative stability lemma for descending Markov chains with jumps close to $1 + \text{Pois}(W(r))$. Also the Takeuchi-minus-Bell coefficients, and real and integer inverses that are instances of the transseries volume. The manuscript states that the general theorem printed in Prellberg's 2002 slides omits a restriction ($f = z/(1-z)$, $a = z$, $b = 1$ gives B_{n-1} , not a multiple of B_n), without effect on the Takeuchi case; a139383-iterated-bell-diagonals makes the same check, printed only here. C is not enclosed, and effective constants and optimal truncation are open.

143. Rooted identity trees with fixed maximum outdegree

generating-functions-and-asymptotics/oeis-sequence-asymptotics/
a116379-bounded-identity-trees/article.pdf

from batch 77: rooted-identity-trees-source.zip

For rooted unlabeled identity trees with at most d children at each vertex (OEIS A116379 for $d = 3$, A116380 for $d = 4$, whose leading asymptotic OEIS records as unknown), proves for every fixed $d \geq 2$ the square-root law $a_{d,n} = C\rho^{-n}n^{-3/2}(\sum_{j \leq R} D_j n^{-j} + O(n^{-R-1}))$ to every fixed order, with $C > 0$ and explicit D_1, D_2 , from the signed root equation: $1/4 \leq \rho \leq 9/20$, a transversality bound with the constant $1 - 810/(319\sqrt{19})$ independent of d , a characteristic point and a unique dominant singularity. Tabulates ρ, C, D_1, D_2 for $d = 3, 4$ to 30 uncertified decimals ($\rho_3 = 0.40077450047\dots$, $C_3 = 0.41431687066\dots$) and inverts a specified finite model by W_{-1} , an instance of the transseries volume. Nothing is uniform in d ; the derivative cancellation is Bell, Burris and Yeats's, and no priority is claimed, since Meir-Moon-Mycielski (1983) and Haigh-Kennedy-Quintas (1984) were not read in full.

144. Tournament score sequences and weighted component asymptotics

generating-functions-and-asymptotics/oeis-sequence-asymptotics/
a000571-tournament-score-sequences/article.pdf

from batch 77: tournament-score-sequences-source.zip

Refines Kolesnik's leading asymptotics for the score sequences of tournaments (OEIS A000571) and of strong tournaments (A351822), from the generating function $\exp A(z)$ of Claesson-Dukes-Franklín-Stefánsson: to every fixed order, $T_n^\pm = (e^{\pm\lambda}/(2\sqrt{\pi}))4^n n^{-5/2}(1 + c_1^\pm/n + \dots)$ with $\lambda = A(1/4) = 0.33023754398\dots$, $c_1^\pm = -1/8 \pm 5\mu/2$, $\mu = A'(1/4)/4 = 0.67395260185\dots$ and explicit $c_2^\pm$. For the count weighted by u^m , m the number of strong blocks: fixed subcritical weights; at the critical weight an expansion in $n^{-1/2}$ with no n^{-1} term; uniformly on a logarithmic window above it a half-power expansion with Kummer-function coefficients; and at $\tau = \frac{1}{2}\log n + 2\log\log n + \log(\mu\sqrt{\pi}/2) + s$ a coexistence profile proportional to $1 + e^{-s}$, which is not claimed to be a mixture law. Inverses of specified models are instances of the transseries volume. Nothing is claimed for $\tau < 0$ or for growing orders, and the renewal mechanisms are credited to Blanchet and Glynn.

145. Fixed forbidden distances in involutions

generating-functions-and-asymptotics/oeis-sequence-asymptotics/
a239144-forbidden-distance-involutions/article.pdf

from batch 77: fixed-forbidden-distance-involutions.zip +
involution-refinements-attribution-revision.zip (its earlier edition
involution-refinements-reproducibility.zip superseded, not printed)

For involutions of $[n]$ avoiding the transpositions of a forbidden graph of fixed maximum degree, proves a uniform complex probability-generating function expansion to every fixed order in $n^{-1/2}$,

with coefficients evaluated at the graph's own matching-root moments, and an exact split of the avoidance count into two Gaussian sectors of relative weights $e^{\pm\sqrt{n}}$, each with its own expansion. For a fixed finite set of forbidden distances the local statistics are eventually exactly affine in $1/n$, so the columns of OEIS A239144 have constant coefficients: $A_{n,r}/I_n = e^{-r}(1 + r/\sqrt{n} - r/(2n) - r(8r^2 - 12r + 3)/(24n^{3/2}) + O(n^{-2}))$. Section 9 holds the second manuscript's case $r = 1$: the all-orders law of the number of adjacent transpositions (A217876; zero column A170941), its total-variation distance from Poisson(1), a second proof by growing moments and a Touchard transform, a Chebyshev sector decomposition with sector ratio $(-1)^n e^{-2\sqrt{n}}(1 - 31/(12\sqrt{n}) + O(1/n))$, and coefficients through order eight. *Merged: the second manuscript proves the case $r = 1$ with the same coefficients $31/24$, $-55/48$, $1717/1920$; its inverse statements are given by pointer.* The Gaussian matching identity and the root-moment method are prior work (Godsil, Godsil-McKay, McLeod), the zero-count limit is Ganjtabesh-Steyaert's, the Poisson law Li's and $31/24$ Kotěšovec's. The inverses are instances of the transseries volumes.

146. All fixed order asymptotics for distinct parabolic double cosets

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a260700-parabolic-double-cosets/article.pdf](#)

from batch 77: [double-coset-report.zip](#)

Let p_n count the distinct subsets of S_n that are double cosets of standard parabolic subgroups (OEIS A260700). Starting from Browning's exact enumeration, proves $p_n = Kn! \rho^{-2n} (\sum_{m \leq M} B_m(\rho) n^{-m} + O(n^{-M-1}))$ for every fixed M , with $\rho = \log 2$, $K = e^{-\rho^2/2}/(4\rho^2)$ and $B_1(r) = r^2/2 - r^3 - 11r^4/12 + r^5/4$; B_2, B_3, B_4 are explicit. The first correction proves Browning's higher-order conjecture with $c = -KB_1(\rho) = 0.10819705289 \dots > 0$, the sign proved without a numerical test. The tools are uniform generalized-Fubini pole extraction, summable bounds on the signed inner defects and a joint bound for the omitted long-cycle configurations of the outer Stirling transform, with a finite exact coefficient algorithm. The leading equivalent is Browning's theorem, novelty rests on a bounded search, and five questions stay open. Smooth Lambert- W inverses and an eventual two-ceiling bracket for the integer threshold are instances of the transseries volume; the report keeps the counterexample $y = p_{25} + 1$ to the naive rounding $N(y) = \lceil x_1(y) \rceil$. A companion of the A260952 report below; neither uses the other.

147. Full support signed permutations and their asymptotic coefficients

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a260952-full-support-signed-permutations/article.pdf](#)

from batch 77: [Coxeter_Full_Support_Asymptotics.zip](#)

Proves Kotěšovec's 2015 conjecture recorded in OEIS A260952, $a_m \sim -2^{m+1}m!/(9(\log 3)^{m+1})$, with the corrections $1 - 2\tau/m - 9\tau^2/(4(m)_2) - 27\tau^3/(2(m)_3) - 1755\tau^4/(16(m)_4) + O(m^{-5})$, $\tau = \log 3$, every further fixed order by a finite formula, and $a_m < 0$ for all $m \geq 1$. The a_m are the asymptotic coefficients of the full-support elements of the Weyl groups of types B and D (A109253, A112225); they are written as a positive Stirling transform whose late expansion holds to every fixed order on the scale $\log(1+q)$, uniformly in q . The two exact endpoint sums of the signed-permutation model have separate all-order expansions for $1/4 < |q| < 4$ (not extended to $q \geq 4$), which give an all-order sparse-sign crossover: the number of negative entries tends to a Poisson law conditioned to be positive. The models are Bergeron-Hohlweg-Zabrocki's and the Stirling-transform construction Martin-Kearney's. Lambert- W growth-index inverses with a stated rounding width are instances of the transseries volume. A companion of the A260700 report above; neither uses the other.

148. Ordered long-jump rook paths

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a227578-ordered-rook-paths/article.pdf](#)

from batch 77: [ordered-rook-reproducibility.zip](#)

Let $A(n, k)$ count the paths from $(n, \dots, n)$ to the origin that lower one coordinate by any positive amount at each step and stay weakly ordered (OEIS A227578; columns A059231, A227580, A227583, A227596). For every $k \geq 2$ the report proves an exact coefficient identity with a squared

Vandermonde numerator, so each column is D-finite (no posted recurrence is certified), and, by the Raichev–Wilson smooth-point theorem, $A(n, k) = C_k(k+1)^{kn}n^{-(k^2-1)/2}(\sum_{j < M} d_j(k)n^{-j} + O(n^{-M}))$ for every M , with C_k in closed form through a Mehta-type Gaussian integral, $d_1(k) = -(k-1)(k+1)(2k^4 + 8k^3 + 9k^2 + 6k + 12)/(12k(k+2)^2)$ and $d_2(k)$ in closed form; every $d_j(k)$ comes from a finite generator and is a rational function of k with poles only at $0, -1, -2$. This proves the general fixed- k growth that OEIS records as a conjecture; the leading constants for $k \leq 5$ were already recorded there by Kotěšovec, and the methods are classical. The W_{-1} inverse and the two-ceiling threshold enclosure are instances of the transseries volume. “Rook” means a move here, not a rook polynomial.

149. Regular cyclic word covers

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a108242-regular-cyclic-word-covers/article.pdf](#)

from batch 77: [regular-cyclic-word-covers-reproducibility.zip](#)

Counts sets and multisets of rotation classes of words of fixed length $\ell \geq 3$ over an n -letter alphabet in which every letter occurs exactly r times. An exact slot-partition identity gives, for fixed degree, the ratio to the carrier $(rn)!/(\ell^{rn/\ell}(rn/\ell)!(r!)^n)$ to every order in $1/n$, with rational coefficients and absolute remainder control. For $\ell = 3$ it identifies OEIS A108242, A110105, A110104 and A110106 (following the examples and Mishna’s source where OEIS titles disagree with the parameters), with corrections through n^{-5} , such as $1 + 8/(9n) + 104/(81n^2) + \dots$, and the corrections each posted carrier induces. At the critical degree $r = \lambda n^{(\ell-2)/2}$ it proves an expansion to every fixed order in $n^{-1/2}$, uniform on compact positive λ -intervals, with multiplier $\exp(s\lambda^2/(2\ell) + \mathbf{1}_{2|\ell} \lambda/\ell)$, $s = \pm 1$; the even-length term comes from half-turn symmetry. The leading equivalents were already posted by Kotěšovec, and the literature comparison is an applicability screen, not a priority claim. A W_0 inverse with a lattice threshold enclosure is an instance of the transseries volume. The open cyclic-words question of the A330266 report concerns other objects and is not answered.

150. Unitary divisor sum colored partitions and their asymptotics

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a301981-unitary-divisor-partitions/article.pdf](#)

from batch 77: [unitary-partition-report.zip](#)

For the products $\prod(1 - z^m)^{-\sigma^*(m)}$ (OEIS A301981) and $\prod(1 + z^m)^{\sigma^*(m)}$ (A301982), σ^* the unitary divisor sum, proves that the stretched-exponential equivalents conjectured in both OEIS entries are false: the ratio of coefficient to model does not eventually stay in any compact subinterval of $(0, \infty)$, by a least-height zeta-zero noncancellation (the Mellin kernel has denominator $\zeta(2s-1)$) and an Abelian transfer. Unconditionally $\log a_n = cn^{2/3} + o(n^{1/3})$ with $c = 3(A/4)^{1/3}$, $A = \pi^2/6$ for A301981 and $\pi^2/8$ for A301982, and the error is not $O(n^{1/4-\varepsilon})$; for each sequence separately, an error $O_\varepsilon(n^{1/4+\varepsilon})$ for every $\varepsilon > 0$ is equivalent to the Riemann hypothesis, which is neither proved nor disproved. Also a corrected equivalent at the explicit point $(2A/n)^{1/3}$ (without a rate), all-fixed-order exact-saddle and off-center Hermite expansions, and two-ceiling threshold brackets whose integer step is an instance of the transseries volume. Which one-sided divergence occurs stays open. The refutation concerns OEIS text, not a repository claim, and no OEIS submission is implied.

151. Log-periodic factors in radix-layer partition asymptotics

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a174065-radix-layer-partitions/article.pdf](#)

from batch 77: [radix-partition-reproducibility.zip](#)

For $b \geq 2$ let $a_b(n) = [z^n] \prod_{j \geq 0, i \geq 1} (1 + z^{ib^j})$; for $b = 2^m$ it counts partitions into parts k with $v_2(k) \equiv 0 \pmod{m}$: ordinary partitions at $b = 2$, OEIS A174065 at $b = 4$, A393565 at $b = 8$. The report proves an expansion of $a_b(n)$ to every fixed order against $KN^{-\kappa}e^{2\sqrt{AN}}$, with explicit differential operators acting on a positive analytic 1-periodic function H of $\log_b \sqrt{N/A}$, and shows by the zero-free line of ζ that H is nonconstant for every $b > 2$. Hence the equivalents recorded in A174065 and A393565 have correct smooth constants but omit a persistent log-periodic factor; a

separate Abelian argument excludes every constant-only equivalent, and the subsequential ratios fill exactly $[\min H, \max H]$, with $(\min H)(\max H) = 1$ from an exact radial reciprocity. Also eventual strict increase and log-concavity, and a two-ceiling threshold enclosure whose integer step is an instance of the transseries volume. The methods are classical Mellin analysis (Flajolet–Gourdon–Dumas, Madritsch–Wagner). The correction concerns OEIS text, not a repository claim, and no OEIS edit was made. The mechanism, but no theorem, is shared with the A033552 report.

152. Full asymptotic expansions for partition sums of acyclic orientations

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/](#)

[a372395-acyclic-orientation-partitions/article.pdf](#)

from batch 77: [orientation-partition-asymptotics.zip](#)

OEIS A372395 and A370613 sum the numbers of acyclic orientations of the complete multipartite graphs K_λ over the partitions λ of n , unrestricted or into distinct parts. The report proves $B_\pm(n) = A_\pm n! n^{-p_\pm} e^{C_\pm \sqrt{n}} (\sum_{j \leq J} a_{\pm,j} n^{-j/2} + O(n^{-(J+1)/2}))$ for every J , with $p_+ = 1$, $p_- = 3/4$, explicit amplitudes including a tilted Gamma-integral factor, and $a_{+,1} = -0.30756533\dots$, $a_{-,1} = 0.23095021\dots$; this answers the question of Sun (arXiv:2605.04006), to whom the exact Gamma representation and the constants $C_\pm$ are due. A uniform quadratic majorant controls the signed representation; a finite Gaussian-expectation algorithm gives every coefficient, and a second short route gives $a_{\pm,1}$. A corollary extends the expansion to the sums of $(-1)^{|V|} \chi_{K_\lambda}(-v)$ for fixed $v > 0$, with $\Gamma(n+v)/\Gamma(v)$ in place of $n!$. The expansions are Poincaré expansions, not transseries, and four questions stay open. The Lambert- W inverse with its integer-threshold envelope is an instance of the transseries volumes.

153. Sparse nonattacking chess placements

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/](#)

[a201513-sparse-chess-placements/article.pdf](#)

from batch 77: [Sparse_Chess_Asymptotics.zip](#)

For n nonattacking kings (OEIS A201513) or knights (A201540) on an $n \times n$ board, measured against Kotěšovec’s common leading equivalent $B_n = e^{-9/2} n^{2n}/n!$, proves rational expansions to every fixed order, four corrections explicit: $K_n/B_n = 1 + 7/(3n) + 119/(9n^2) + 11327/(810n^3) + 90475/(1944n^4) + O(n^{-5})$ and $N_n/B_n = 1 + 37/(3n) + 914/(9n^2) + 330917/(810n^3) + 966583/(1944n^4) + O(n^{-5})$. For any fixed finite symmetric move set S the count is $(n^{2n}/n!) e^{-(|S|+1)/2}$ times an all-orders series, also uniformly for k pieces with k/n in a compact positive interval, and the first correction is $c_1 = B + 1/3 - \tau$ (boundary loss B , triangle density τ), which is why kings and knights share the leading term but not the next; wazir (A201511) and fers (A201861) coefficients through order four follow. The cluster-expansion method is credited to Davies–Jenssen–Perkins and Pulvirenti–Tsagkarogiannis. The coefficients depend on the carrier: against the Stirling-expanded carrier the first corrections are $9/4$ and $49/4$. The Lambert- W growth-index inverse and its rounding qualification are instances of the transseries volume.

154. Asymptotic enumeration of tied football seasons

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/](#)

[a380592-tied-football-seasons/article.pdf](#)

from batch 77: [tied-football-report.zip](#)

In a double round-robin league of n labelled teams with $3/1/0$ scoring, $a(n)$ (OEIS A380592) counts the seasons in which every team finishes on the same score. The report proves $a(n) \sim B_n \Theta(c_n)$ with $B_n = 3^{n(n-1)} n^{-1/2} (9/(56\pi n))^{(n-1)/2} e^{-9829/65856}$, $c_n = 8(n-1)/3 - 13/84$ and the positive 1-periodic theta function $\Theta(c) = \sum_j e^{-4\pi^2 j^2/9} e^{2\pi i j c}$, so the multiplier depends on $n \bmod 3$ and cannot be replaced by 1. The proof uses mean-matching exponential tilts and a sharp global Fourier localization: a negative quartic gives an exact Gaussian majorant and an outlier bound handles the complement. On each residue class the conditioned common score tends to a discrete Gaussian centred at c_n , which is not its conditional mean. After checking the hypotheses of Isaev’s complex cumulant theorem (arXiv:2508.16952), the external input, it gives every fixed order in $1/n$ with a finite exact algorithm and the first correction. The residue-aware inverse

threshold enclosures need no Lambert function, the phase being quadratic, and are instances of the transseries volume's reversion and staircase theorems. The model and its exact counts are Jahn–Habermann–Lavrov's.

155. All-orders asymptotics and asymptotic inversion for A292507

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a292507-binomial-partition-transform/article.pdf](#)

from batch 77: A292507_report_and_code.zip

For $a_n = \sum_k \binom{n}{k} b(k)$, where $b(k) = p(k) - p(k-1)$ counts the partitions of k without parts 1 (OEIS A292507), proves an all-orders Poincaré expansion $a_n = C 2^n n^{-3/2} e^{\beta\sqrt{n}} (\sum_{j \leq M} c_j n^{-j/2} + O(n^{-(M+1)/2}))$ with $\beta = \pi/\sqrt{3}$ and $C = (\pi/6)e^{\pi^2/24}$, an additive form of Kotěšovec's logarithmic conjecture in the entry; c_1, c_2, c_3 are in closed form, and a finite Gaussian-moment algorithm gives every c_j ($c_1, \dots, c_6$ computed). Also an all-orders reversion of the real model in $X^{-1/2}$ and $\log X$ with a residual bound, and shrinking integer-threshold brackets with existence constants only, not certified thresholds; these are instances of the transseries volume's Lambert, perturbed-inversion and staircase theorems and are cited, not claimed. The only analytic input is Nemes' explicit partition-remainder estimate. The leading binomial-smoothing mechanism is prior art (A218481), the coefficient formula is Gutkovskiy's, and no publication priority is claimed; nor is convergence of the correction series or a complete transseries.

156. Permutations avoiding the vincular pattern 12–34 (A113226)

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a113226-vincular-avoiders/article.pdf](#)

from batch 77: a113226-asymptotics.zip + a113226-refined-addendum.zip

For the permutations avoiding the vincular pattern 12–34 (OEIS A113226), equivalently Bevan–Cheon–Kitaev's $\{3, 2+2\}$ -free naturally labelled posets, Part I derives the closed EGF $H(z) = \exp\{z/2 + q(3 \arcsin q - \pi/2)/\sqrt{1-q^2}\}$, $q = e^{z/2}/2$, identifies the cumulants as A136127 shifted, and proves the stretched-exponential form of Bevan–Cheon–Kitaev's Conjectures 13 and 14 with exact constants and to all orders: $A_n \sim D n! \rho^{-n} e^{3cn^{1/3}} n^{-5/6} (1 + \sum_j a_j n^{-j/3})$ with $\rho = \log 4$, $c = (\pi^2/(4\rho))^{1/3}$ and $D = 2e^{-3}\sqrt{c/(3\pi)} \approx 0.0357$, where their fit gave about 0.032; $a_1, \dots, a_4$ are exact. Its Lambert- W inverse is an instance of the transseries volume. Part II, an addendum, marks the distinct strict downsets and the isolated elements: a record-to-cycle bijection explaining the exponential transform, a complex-uniform saddle expansion, all-orders local large deviations for interior densities, Gaussian and local limit laws, and a conditional Poisson law of isolated elements with total-variation error $O(n^{-2/3})$. Endpoint densities, exponentially small sectors and certified constants are not claimed; Part II answers Part I's questions 1–2 in part, and 3–4 stay open. The two heavy exact tables are not shipped; the README regenerates them.

157. Asymptotic expansions and inversion for A124380

[generating-functions-and-asymptotics/oeis-sequence-asymptotics/a124380-signed-moment-asymptotics/article.pdf](#)

from batch 77: a124380-reproducibility.zip

For $\sum_n a_n z^n = \sum_k z^k \prod_{j \leq k} (1 + jz)$ (OEIS A124380), proves an exact decomposition $a_n = I_n + (-1)^n J_n$ into a positive and a signed moment integral, the multiplicative equivalent $a_n \sim \frac{1}{2} e^{1/2} x \exp\{x^2(2L-1) + x(L+1) + L^2/8\}$ with $x = \sqrt{n/2}$ and $L = \log x$, which is more than Kotěšovec's logarithmic conjecture in the entry, and every fixed-order relative expansion with coefficients polynomial in L . The signed sector is exponentially smaller and has its own all-orders oscillatory expansion, with the error measured against its amplitude so that it stays valid near zeros of the sine, and coefficients $q_j^-(L) = (-1)^j q_j(L - i\pi)$. Also eventual strict log-convexity, an all-orders inverse in $X = \sqrt{y/W(y/e)}$, an instance of the transseries volume's inverse-Gamma core with no novelty claimed, and a conditional certification rule for the integer threshold; $T(y) = \lceil \nu(y) \rceil$ is not asserted. The combinatorial models are Chen–Fan–Zhao's and Cerbai–Claesson–Sagan's, no priority is claimed, and four research questions stay open.

158. Complete increasing subsequences in multiset permutations (A047909 and A268485)

generating-functions-and-asymptotics/oeis-sequence-asymptotics/
a047909-beta-renewal-subsequences/article.pdf
from batch 77: beta-renewal-asymptotics-reproducibility.zip +
beta-renewal-uniform-reproducibility.zip

Studies the probability $p_m(k) = H(m, k)(m!)^k/(mk)!$ that a word with m copies of each of $1, \dots, k$ contains $12 \cdots k$ (OEIS A047909, diagonal A268485). Its representation as the probability that k independent $\text{Beta}(1, m)$ increments sum to at most 1, and the Gaussian limit, are El Maazouz–Pitman’s and are credited, not claimed. Part I proves an expansion to every fixed order in the window $k = m + x\sqrt{m}$ with $P_1, \dots, P_4$ explicit ($P_1 = (x^2 + 8)/6$); on the diagonal only odd half-integer powers occur beyond the leading $1/2$, with four corrections computed. It adds smooth quantile and Lambert- W count inverses with qualified integer brackets and all-orders rare-tail expansions for k/m in compact subsets of $(0, 1)$ or $(1, \infty)$. Part II, an addendum, joins the regimes: relative error $O(m^{-(R+1)/2})$ for both tails, uniformly on compact ranges of k/m and through $k = m + 1$, recovering Part I’s coefficients by a second route, and both real Lambert- W branches for tail thresholds. The inverses are instances of the transseries volume and the tilting method is classical (Esscher, Cramér, Daniels). Effective constants, growing orders, ratios tending to 0 or ∞ , and certified rounding remain open.

159. Uniform asymptotics and moving zeros for the level-15 family of Cooper

generating-functions-and-asymptotics/oeis-sequence-asymptotics/
cooper-level-15-moving-zeros/article.pdf
from batch 77: modular-crossover-replay.zip

For Cooper’s level-15 polynomials $T_\varepsilon(n)$, of degree n in the parameter ε , proves a two-singularity expansion with rates $\varepsilon + 11$ and $\varepsilon - 1$, uniform to every fixed order on the disc $|\varepsilon + 5| \leq 1/10$, with an additive error that stays valid where the two terms cancel; in the window $\varepsilon = -5 + u/n$ the profile is $e^{u/6} + 10(-1)^n e^{-u/6}$. Hence for every large odd n there is exactly one zero with $|u - 3 \log 10| < 1$, real and simple, at $\varepsilon_n = -5 + 3 \log(10)/n + (162/125 - \frac{9}{2} \log 10)/n^2 + O(n^{-3})$, and no real zero in a bounded window for large even n . Explicit path, sheet and connection proofs give all ten constants of Cooper’s Table 8 (levels 11, 14, 15 and 24), labelled conjectured and determined numerically there, including two negative-axis connections with exact eta multipliers. The formulas are Cooper’s, the level-11 equivalent is Kotěšovec’s (A284756) and the positive modular radial principle Guillera–Zudilin’s. Also a finite generator for every correction order and a qualified, non-effective threshold inverse, an instance of the transseries volume. No historical priority and no global location of the zeros are claimed; the OEIS numbers of five of the sequences were not verified.

160. Moving fugacity asymptotics for distinct partitions

generating-functions-and-asymptotics/oeis-sequence-asymptotics/
a291698-moving-fugacity-partitions/article.pdf
from batch 77: moving-fugacity-reproducibility.zip

For $a_n(\alpha) = [q^n] \prod_{k \geq 1} (1 + n^\alpha q^k)$, distinct partitions with one size-dependent weight $u = n^\alpha$ per part, proves, uniformly for α in compact subsets of $(0, \infty)$, an all-orders principal Bessel expansion with a finite generator for its coefficients and, through an exact Jacobi decomposition, finitely many conjugate resonance sectors with separated local integrals and sectorwise remainders. This yields the equivalents conjectured by Kotěšovec for A291698 ($\alpha = 1$) and A292304 ($\alpha = 2$), and shows that the elementary Sommerfeld form is relatively valid exactly when $\alpha \geq 1/2$. Also a specified monotone real continuation with a Lambert- W inverse and its first correction, a formal reversion generator and an integer-threshold statement, instances of the transseries volume with no novelty claimed. The onset is slow: at $n = 20,000$ the principal term for $\alpha = 2$ is still off by about 5.6%. The sector count and depth are fixed; no growing-sector theorem, convergence or explicit constants are claimed, nor priority, and four questions stay open.

12.3 Asymptotics, records and discrepancy

161. The Exponential Scale of Gregory Sign Thresholds

generating-functions-and-asymptotics/
gregory-coefficient-sign-thresholds/gregory_sign_threshold.pdf
from gregory_sign_threshold_research.zip

For $(x/\log(1+x))^m = \sum_n G_n^{(m)} x^n$ and $\text{Gr}(m)$ the least N beyond which $(-1)^{n-1} G_n^{(m)} > 0$, proves $\log \text{Gr}(m) = m + 1 - \gamma - (\zeta(2) + \zeta(3))/m + C_2/m^2 + O(m^{-3})$ with C_2 explicit, plus a finite algorithm for every coefficient of the complete expansion; in particular $\text{Gr}(m)/e^m \rightarrow e^{1-\gamma}$ and $\text{Gr}(m)/3^m \rightarrow 0$. This replaces Conjecture 6.1 of Xu and Zhao (arXiv:2609.11072v1), whose printed lower bound already conflicts with its own reported $\text{Gr}(5) = 113$; that value is independently certified here.

162. Exact Records in the Square-Root-of-Three Walk

generating-functions-and-asymptotics/sqrt-three-walk-records/article.pdf
from sqrt3_record_recurrence.zip

Proves the four-phase record recurrence for $S(N) = \sum_{n \leq N} (-1)^{\lfloor n\sqrt{3} \rfloor}$ in the corrected form on page 7 of Bruin and Fokkink, *On the records and zeros of a deterministic random walk* (arXiv:2503.11734v2), where it is experimental and attributed to Arias de Reyna and van de Lune. Includes minimality of every claimed first passage, a rational generating function, closed forms, sharp logarithmic extrema, and an extension to every even integer trace $D \geq 4$.

163. Sharp discrepancy for two families of binary substitutions

generating-functions-and-asymptotics/binary-substitution-discrepancy/article.pdf
from oeis_discrepancy_counterexamples.zip; batch 73:
OEIS_Quadratic_Morphic_Words_Research_Package.zip

For the fixed word of $0 \mapsto 1, 1 \mapsto 101010$, the position sequences A284365 and A284366 both carry a conjectured upper position error below 2, and both conjectures are false: the first counterexamples are at ranks 2,977,771 and 8,933,313, with common error $(2977771\sqrt{21} - 13645857)/2 \approx 2.00487$ and sharp common upper bound $(6 + \sqrt{21})/5 \approx 2.11652$. Proves complete discrepancy limit intervals for every substitution $0 \mapsto 1, 1 \mapsto (10)^m$, with infinite counterexample families and exact recurrences for their indices. Part II, added in batch 73, treats the disjoint family $0 \mapsto 1, 1 \mapsto 10^a 1^b$: the prefix discrepancy is bounded exactly when $a \leq b + 1$, with sharp, unattained position-error envelopes from a four-state Bellman system, and sharp bounds for A284368–A284371. The density question stays open in both Parts.

164. Exact eventual maximizers of Rudin–Shapiro autocorrelations

generating-functions-and-asymptotics/
rudin-shapiro-autocorrelation-maximizers/article.pdf
from rudin_shapiro_maximizers.zip

For the aperiodic autocorrelation $C_m(k)$ of the level- m Rudin–Shapiro sequence, classifies the maximizing shift exactly for all large m : $k_m^* = (2^{m+1} + (-1)^m)/3$, with $A_m = b_{m-2}$, settling the discrete uniqueness-and-location conjecture of Tarnu (J. Approx. Theory 287 (2023) 105866), repeated as Conjecture 1.5 of Choi–Tarnu (2025), including the limit $k_m^*/2^m \rightarrow 2/3$. The certificate is a rational polytope argument with exact arithmetic.

165. Apéry arrays and conjectural zeta accelerations

generating-functions-and-asymptotics/apery-array-zeta-accelerations/article.pdf
from Apery_OEIS_Conjectures_Proofs.zip, apery_oeis_conjectures.zip

Proves the entire shifted-diagonal $\zeta(3)$ family recorded as conjectural in OEIS A143007, and both the super- and subdiagonal $\zeta(2)$ families of A108625, at every nonnegative offset: for all $k \geq 0$, $\zeta(3) = H_k^{(3)} + \sum_{n \geq 1} (2n+k)(3n^2+3nk+k^2)/(n^3(n+k)^3 T_{n-1,n+k-1} T_{n,n+k})$. The main diagonals are the classical Apéry sequences A005259 and A005258. Proofs are finite telescoping identities with rational lattice potentials and explicit limit estimates; no computer test or unproved

conjecture enters. Adds rational error bounds, numerator and denominator recurrences, a three-term recurrence along every shifted diagonal, and sharp fixed-offset asymptotic constants. Merged from two reports whose boxed identities were character-identical up to macro names.

13 The Jacobian conjecture and Keller maps

Four reports continuing the formal project `Algebra/JacobianConjecture`, which verifies in Lean and Rocq Alpöge’s three-variable counterexample and a degree-six stable representative in five variables (its stabilization to all higher dimensions in Lean only); its further degree reductions are SymPy certificates. None of the four reports’ own theorems is formalized; each README names the project declarations it builds on. Three study the project’s three-variable map, one of them its iteration; the fourth studies a different map, Gao’s five-dimensional F_6 .

166. The fiber geometry of Gao’s Keller map F_6

[jacobian-conjecture/gao-f6-fiber-geometry/article.pdf](#)

from `ProveIt_Six_Sheets_Research.zip`, `ProveIt_Keller_Fibers_Research.zip`

For Gao’s five-dimensional six-sheeted Keller map F_6 (arXiv:2608.00222v1, §4.5.1), classifies every geometric fiber: the cardinalities are exactly 0, 1, 2, 3, 4, 6, all seven root-multiplicity patterns occur, and off the hyperplane $c = 0$ the inverse points are the simple roots of an explicit sextic. The omitted set is a smooth closed surface isomorphic to $\mathbb{G}_m \times \mathbb{A}^1$, of codimension three, with the single effective omitted parameter (21/32, 5/32, 25/128). Gives an explicit finite normalization of rank six, the two-component nonproperness locus, the discriminant’s normalization and local analytic types, global monodromy S_6 with two arithmetic certificates, no proper intermediate field, real fiber cardinalities $\{0, 1, 2, 3, 4\}$, and an obstruction to equivalence with a stabilization of a map in at most three variables. The map is Gao’s; the fiber theory is the reports’. Merged from two manuscripts proving the same core in identical normalizations; the second one’s different proofs are kept as marked second routes. Not to be confused with the project’s formally verified dimension-five stable map.

167. Arithmetic local–global dichotomies for the three-variable Keller map

[jacobian-conjecture/arithmetic-local-global-fibers/article.pdf](#)

from `ProveIt_Arithmetic_Local_Global.zip`; batch 58: `ProveIt_Dense_Integral_Hasse.zip`; batch 64: `ProveIt_Beyond_Split_Fibers.zip`

For the project’s three-variable Keller map, gives an explicit infinite family of integral targets with a preimage in every $\mathbb{Z}_p^3$ but none in $\mathbb{Z}^3$ (each fiber is three rational points, so these are not rational Hasse failures); classifies the split integer-root plane $C = 2$ by a gcd condition, with locally soluble parameter density $27/(4\pi^2)$ surviving the removal of S -integral fibers; proves the exact integral 2-adic fiber law (21, 7, 3, 1)/32 for fiber sizes 0, 1, 2, 3, determined modulo 8; and contrasts it with a rational Hasse principle over every number field. The complex stratification is credited to Gao and the finite-field histograms to an external note. An intake remark records an integral triple collision over the project’s own collision value. Not a new disproof of the Jacobian conjecture. Part II, added in batch 58, answers Part I’s density question: such failures are Zariski dense in every plane $C = c \neq 0$, hence in $\mathbb{A}^3$, and none lies on $C = 0$. It gives a complete local criterion for completely split fibers, finds finitely many integral split fibers on each plane, and proves $\#\mathcal{H}_c^{\text{sp}}(T) \sim \kappa(c)T^{2/3}$ for the completely split failures in square boxes $|A|, |B| \leq T$, with $\kappa(c)$ an explicit Gamma factor times an Euler product. Part III, added in batch 64, gives a complete local-integral certificate for every target, nonsplit fibers included, and on $C = 2$ counts the integral Hasse failures in the same boxes: $\frac{27}{2\pi^2}T \log T + O(T)$, almost all of them nonsplit.

168. All-degree rigidity of a weighted Keller class

[jacobian-conjecture/weighted-keller-rigidity/article.pdf](#)

from `ProveIt_Weighted_Keller_Rigidity.zip`; batch 44: `ProveIt_Affine_Weighted_Keller_Classification.zip`

For polynomial maps of weights $(-1, 1, 2)$ written through $t = xy$ and $v = x^2z$, classifies the Keller maps in every degree over any field of characteristic zero: those with p, q affine in v and r affine

in (t, v) are an explicit tame family and a two-parameter diagonal orbit of the known map, so every noninvertible member has coordinate degrees $(7, 6, 4)$ and sixteen monomials, with no input degree bound. Extends to $r = R(t) + \gamma v$, where the possible maximum degrees of the noninvertible members are exactly 7 and every even integer at least 8, and shows that with $r = 1 + t + v$ the fixed degree-seven coefficient scheme is the double point $\mathbb{Q}[\varepsilon]/(\varepsilon^2)$, with an explicitly obstructed tangent (a torus times it when the two coefficients of r are freed). The classification does not cover unrestricted weighted maps. Sharpens to all degrees the project's degree-seven sparsity search; it does not show degree seven minimal in dimension three. Part II, added in batch 44, removes the restriction on the coefficient of v in r : whenever p, q, r are affine in v that coefficient is constant, so the classification covers the whole affine-in- z class; it adds dual-number slopes of every degree and a universal second-order obstruction.

169. Dynamical degrees of the three-variable Keller map

[jacobian-conjecture/keller-map-dynamical-degrees/article.pdf](#)

from `ProveIt_Keller_Dynamics.zip` (batch 70)

Studies the project's three-variable Keller map F under iteration, which neither the project nor the other reports touch. Over any field of characteristic other than three, $\deg F^n$ satisfies $d_{n+2} = 6d_{n+1} + d_n$ with $d_0 = 1, d_1 = 7$ (1, 7, 43, 265, ...), so the first dynamical degree is $3 + \sqrt{10}$; every positive weighted degree of every iterate is explicit, and every coordinate's downward-completed Newton polyhedron has exactly two vertices. Outside characteristic three, composing with a shear $z \mapsto z + y^2 h(xy)$, $h \neq 0$ of degree m , gives first dynamical degree $(2m + 7 + \sqrt{4m^2 + 32m + 61})/2$; in characteristic zero these maps are all Keller, noninjective, of generic degree three and in one tame left-right equivalence class, yet their dynamical degrees are unbounded. In characteristic three the second coordinate is y exactly and the degrees follow separate closed formulas (first dynamical degree 4 or $m + 3$); in characteristic two the map is purely inseparable of degree two. A monomial-dominance escape theorem gives a forward-invariant complex escape region, and integral points there have maximal arithmetic degree and are Zariski dense. No Lean or Rocq source is supplied; the formalization route is a plan.

14 Galois theory and radicals

Two reports continuing the formal project `Algebra/PolynomialFormulas` (Lean and Rocq root formulas through degree four, Abel–Ruffini, and Lazard's quintic solver); none of its theorems is formalized.

170. Specialization-safe radical solvers

[galois-theory-and-radicals/specialization-safe-radical-solvers/article.pdf](#)

from `ProveIt_Radical_Solvers_Research.zip`; batch 52: `ProveIt_Optimal_Kummer_Atlases.zip`

Gives an exact cyclotomic-resultant criterion for the characteristics in which equal-cardinality subset sums of the roots of an irreducible prime-degree polynomial can collide, and evaluates it: the bad characteristics are $\{5\}$ for quintic pairs, $\{2, 7\}$ for septic pairs and $\{2, 7, 29\}$ for septic triples. Classifies the collisions completely in characteristic 29 (exactly the cyclic polynomials $(X - \eta)^7 - b$) and in characteristic 2 (translated $Z^7 + aZ^3 + bZ + d$, realizing the nonsolvable $\text{GL}_3(\mathbb{F}_2)$). Constructs separating subset invariants in every characteristic, reconstructs all roots from one coherent p -th radical on each Fourier chart once the intermediate-field invariant data are given (generalizing Lazard's pivot-one scheme for quintics; a coefficient-only solver is left as further work), realizes every Fourier support by rational polynomials with full affine Galois group, and shows the Picard group of the cyclic configuration quotient is $\mathbb{Z}/p$, so no single global Kummer coordinate exists. The Picard result has a written proof only. Part II, added in batch 52, answers Part I's question on the least number of charts: $\text{CH}^*(\text{Conf}_p(\mathbb{A}^d)/C_p) = \mathbb{Z}[t]/(pt, t^{d(p-1)})$, so every cover by regular Kummer charts, even with nonlinear character sections and different characters on different charts, needs exactly $d(p-1)$ charts ($p-1$ for polynomials: four for quintics, six for

septics), and the Fourier charts attain it. This too is a written proof, by integral intersection theory; its checker verifies only finite identities.

171. Finite separating ranges for sextic block resolvents

[galois-theory-and-radicals/sextic-block-resolvent-separators/article.pdf](#)

from `ProveIt_Finite_Sextic_Separators.zip` (batch 41); batch 57:

`ProveIt_Three_Plus_One_Resolvents.zip`; batch 70: `ProveIt_Zero_Parameter_Sextics.zip`

The project finds parameters separating the sextic pair and triple block-resolvent descriptors by an unbounded `Nat.find` search. This report proves in writing that on the curve $(t, 2t^2)$ one of $t = 1, \dots, 196$ always separates both families, sharply for arbitrary complex test sets, by orbit counting. The bound is not formalized; formalizing it would make that search bounded, and two further unbounded searches of the project remain. Part II, added in batch 57, shows that deciding radical solvability of an irreducible sextic needs no separating parameter: the sextic is solvable exactly when its triple resolvent has a base-field root at one parameter chosen in advance, or its matching resolvent has one at each of three distinct parameters, because a base-field matching root of a nonsolvable sextic comes from a Galois-invariant pentad, of which there is at most one, and occurs at no more than two parameters. Part III, added in batch 70, sharpens this: one compressed matching test at $t = 0$ together with one triple test decides, by an integer certificate that the five values of a pentad are never all equal and by reciprocal transport; and since a pentad's points span an affine space of dimension at least two, the polynomial of false-positive parameters has degree at most one and does not vanish at zero, so any two distinct matching parameters suffice. It adds a denominator-free Pfaffian of order twenty for the constant triple resolvent and a cubic 15×15 matrix polynomial whose determinant is the cube of the constant matching resolvent.

15 Hilbert's tenth problem

Nine reports on Diophantine representations of computation that control their witnesses, beside the formal project `Computability/HilbertTenthProblem`, which proves MRDP in both directions in Lean, the universal pair $(58, 4)$ and the single-fold unary exponential representation of Jones and Matiyasevich. The first three (canonical certificates; probabilistic, quantum and continuous computation; liveness beyond halting) continue that project. The other six, added in batches 76, 78 and 79 (groups as substrates, signal machines, stochastic and thermal exactness, quadratic orthant certificates, polynomial witness histories, smooth finalizers), treat further substrates or witness domains and use MRDP at most as an imported theorem. Batch 79 also extended five of the older reports. None of the reports' own theorems is formalized; each README names the project declarations it builds on or relates to.

172. Canonical Diophantine certificates

[hilbert-tenth-problem/canonical-diophantine-certificates/article.pdf](#)

from `Diophantine_Causal_Computation.zip`, `canonical_diophantine_certificates.zip`, `ProveIt_Linear_Size_Diophantine_Certificates.zip`, `Witness_Faithful_Diophantine_Compilation.zip`, `canonical_trace_polytopes.zip`, `diophantine_counter_programs.zip` (batch 60); batch 61: `Causal_Diophantine_Queue_Compilation.zip`; batch 62: `Names_to_Numbers_Diophantine_Heaps.zip`, `order_free_diophantine.zip`, `ProveIt_Priority_Pumping_Diophantine.zip`, `one_shared_fallback_research.zip`; batch 63: `Diophantine_Certificates_Without_Histories.zip`; batch 78: `Exact_Wiring_Diophantine_Report.zip`, `Topology_Is_Not_Free_Interaction_Nets.zip`, `no_ghost_wires.zip`, `sandpile_diophantine_research.zip`; batch 79: `Positive_Spectrum_Diophantine.zip`, `Spectral_Guards_Without_Time_Expansion.zip`, `no_borrowed_firings.zip`, `Compressed_Queue_Diophantine_Research.zip`, `Eager_Tree_Calculus_Research_Package.zip`

Diophantine representations that control their witnesses, not only their existence. For Petri nets and interval-guarded translations, a convex quadratic integer polynomial whose natural zeros are in bijection with the commutation classes of enabled executions of length T (its real zero set is a rational polytope with exactly those lattice points), with quartic second routes, a Cartier–Foata

variant for bounded causal height, exact commutation tests, #P-completeness of counting and an exponential lower bound for normal-form automata. Single-fold quartics for multisets and nested repetitions of counter actions independent of the repetition counts, for “ $p(i) \geq 0$ for all $i \leq T$ ” with $O((d+1)^3)$ witnesses independent of T , for memory logs with $24L$ witnesses, and for FRACTRAN and scheduled SKI runs. For cyclic-tag and deletion-tag systems and fixed-read FIFO networks, certificates with one coordinate per consumed symbol and one guard slack, with no intermediate queue quantified; faithful word memory with polynomial insertion at both ends needs two natural coordinates. Proves that a single-fold (finite-fold) universal halting polynomial exists exactly when every enumerable set has such a representation, an open problem not solved here. Parts X–XIII, added in batch 62 from four manuscripts: quartic certificates for dynamic heaps and local graph programs, with birth-order normalization of fresh names, two-stack universality and a proof that perpetual memory safety has no existential representation; for polynomials nonnegative on the real orthant, degrees two and three represent exactly the semilinear sets and degree four every enumerable set, with single-fold quadratics for semilinear sets and order-free self-assembly compilers; for first-enabled-rule multiset and FRACTRAN programs the sharp pumping threshold $2H + 1$, a uniform single-fold quartic and c.e.-completeness of eventual periodicity; and conservative priority reaction networks with one shared fallback, with single-fold trace and minimum-fuel quartics and a least-reservoir relation whose single-fold (finite-fold) representability is again equivalent to that of every c.e. set. Part XIV (batch 63): one quartic per run-length routing topology with exactly $4n + 4m + s$ witnesses and a unique root iff the run terminates, and a fixed polynomial-time rank function whose single-fold (finite-fold) graph is equivalent to the corresponding principle for all c.e. sets. Part XV, added in batch 78 from three manuscripts: exact port wiring for Lafont’s interaction combinators. If $M \neq N$ are perfect matchings of $2n$ ports and Q is a uniformly random one, closing by Q gives different parities of cyclic wires with probability between $2/5$ and $2/3$, both bounds sharp and the equality cases classified; bounded six-rule compilers with $O(C_0 + F + T)$ natural coordinates, degree four and one witness per legal schedule (scheduled histories, not commutation classes), built on a minimal ten-state boundary calculus, a ten-pattern annihilation kernel and memory certificates that are ledger variants of Part V’s $24L$ (whose product lemma two of the manuscripts re-prove; no novelty is claimed for it); and a canonical permutation-orbit certificate with an implemented fixed-schedule compiler. This answers Part X’s question on an interaction-combinator frontend for bounded scheduled histories; quotienting by independent schedules stays open. Part XVI (batch 78): for abelian sandpiles on a finite undirected multigraph with a sink, an exact support criterion for the odometer and cubic integer polynomials with exactly one natural zero for every input, carrying the odometer, the stable endpoint and canonical burning ranks instead of a toppling history, with $10n + 6E$ coordinates in the compact form (a directed counterexample fixes the scope); on $\mathbb{Z}^d$ with a stable periodic background, an exterior collar that certifies global halting exactly, with a unique canonical radius that by Cairns’s imported universality theorem has no computable bound; and the exact minimal eventual period of stabilization along an input ray. It answers Part XIV’s question on infinite-background abelian computation for undirected graphs, and its single-fold equivalence for Cairns’s relation is an instance of Part IX’s. Batch 79 added a second route inside Part XVI, written independently: $12n + 16E$ quadratic residuals in $11n + 12E$ natural witnesses with exactly one natural zero, carrying the odometer, whose sum of squares is a quartic with an integral rejection gap, and on $\mathbb{Z}^3$ one fixed radius-one normal form with 47 finite-deviation fields whose field solution exists, uniquely, exactly when the input stabilizes (neither ledger improves the other). Part XVII (batch 79, from two manuscripts proving one theorem by two routes): complete discrete sign charts of positive-base exponential polynomials along a loop, verified at run endpoints only (a row of order m has at most $2m - 1$ sign runs, sharp), with one witness per input in arithmetic augmented by power atoms $y = b^n$ ($O(D^3)$ witnesses for fixed bases, $O(D^4)$ with the bases as inputs) or, for an externally fixed bit bound, in an ordinary quartic; the complete sign-chart graph of $2^n - y$ has a single-fold (finite-fold) representation exactly when every c.e. set has one, which sharpens a proposition of Part IX; and $\Re((3 + 4i)^n)$ changes sign with positive density on

every residue class. Part XVIII (batch 79): for reliable FIFO actions along a straight-line trace grammar, a quartic with exactly $6g + 3c + 1$ natural witnesses and one zero exactly when the trace connects the supplied words, quantifying no intermediate queue; exact macro repetition counts, infinite repeatability as a word equation, a Fine–Wilf pumping threshold and unique-witness quartics for supplied infinite loops; its re-proofs of Parts I and VI in one coordinate are printed as pointers. Part XIX (batch 79): for Barry Jay’s Tree Calculus under eager evaluation and every externally fixed bound N on distinct calls, a quartic with $3N^2 + 19N$ natural witnesses whose zeros project to the terminating applications (memoized proof DAGs, made well-founded by additive call costs), a canonical refinement with exactly one witness, a strict bracket-abstraction compiler from weak call-by-value λ -calculus, and a literal universal tree of 175 constructor-DAG nodes with an r.e.-complete halting domain, whose code of about $4.5 \cdot 10^{10}$ bits is charged by 175 quadratic equations and never materialized. Merged from twenty-one manuscripts; not a Lean formalization.

173. Diophantine laws of probabilistic, quantum and continuous computation

[hilbert-tenth-problem/probabilistic-quantum-and-continuous-computation/article.pdf](#)

from ProveIt_Diophantine_Quantum_Normalization.zip, quantum_diophantine_research.zip, ProveIt_Diophantine_Barriers.zip (batch 60); batch 62: Quartic_Probability_Landscapes.zip, ProveIt_Reversible_Equilibria.zip, Diophantine_Certificates_for_Coherent_Computation.zip, proveit_rational_rotations_research.zip, dense_rotations_research.zip, strict_contractions_research.zip, Diophantine_Contracting_2Adic.zip, Precision_Is_Memory_Diophantine_Neural_Computation.zip; batch 64: Rational_Tubes_Integer_Zeros.zip; batch 76: Infinite_Quantum_Runs_Diophantine_Certificates.zip

One fixed quartic represents the output law of every effective probabilistic or quantum program as masses of projected roots; exact normalization budgets with a unique-witness quartic certificate; normalized two-outcome probabilities are exactly the weakly computable reals, and conditional comparisons are Σ_2^0 -complete. Deciding whether a seven-outcome rational quantum instrument in dimension four has an impossible outcome word is undecidable (via Neary’s matrix-mortality theorem), with bijective quartics for words of fixed length. One rational piecewise-affine map of the cube realizes every disjoint pair of enumerable sets as exact acceptance and robust rejection, with quartic safety certificates for a computable critical-noise radius. Parts V–VII, added in batch 62 from eight manuscripts: Boolean circuits compile to coercive integer quartics of height at most 4 whose real zeros are Boolean, with uniform Hessian and rounding constants, so that halting probabilities of fair-coin programs are root counts, component counts or Euler characteristics; birth–death chains with defect environments have the sharp spectral gap $(3 - 2\sqrt{2})/4$ and a root mass that is rational exactly for eventually periodic environments; unique-witness quartics for coherent contraction trees over $\mathbb{Z}[e^{i\pi/4}]$, and dense rational $\text{SO}(4)$ gate groups from the rotations $(3 \pm 4i)/5$, $(3 \pm 4j)/5$ with c.e.-complete state transfer or c.e.-complete membership; a fixed $1/2$ -Lipschitz rational piecewise-linear map of $\mathbb{R}^4$ and a $1/2$ -contraction of $\mathbb{Z}_2$ with Σ_1^0 -complete exact half-space reachability and exact extinction; and unique-witness quadratics for rational clipped neural networks, whose tied four-layer synthesis is equivalent to Hilbert’s tenth problem over $\mathbb{Q}$. Part VIII (batch 64): complete integer tube certificates, sums of squares of quadratics with at most one witness, for reaching open rational targets under rational polynomial flows, and global mass-action normal forms that carry them to chemical reachability. Part IX, added in batch 76 from one manuscript: for time-homogeneous rational quantum loops (completely positive, trace-nonincreasing), the group inverse and Cesàro projection as the unique solution of four linear equations with no rank choice, all stopping moments by one finite recurrence certified by a Faddeev–LeVerrier circuit, unique seven-coordinate natural wires extending Part VI’s quartic compilation to rational inputs, an exact weak-clock resolvent with an exponential limit law, a geometric clock on the Eisert–Müller–Gogolin instruments giving every device the law $(16/25)(9/25)^n$ while forbidden-record existence stays undecidable (also in dimension four, conditional on Part II), a two-state counterexample to the unrestricted reading of Lardizabal’s rationality conjecture, and no uniform existential Diophantine graph for computably scheduled one-dimensional loops. Its three arguments repeated from Parts I, II and V are pointers. The report now points to the sibling reports

stochastic-and-thermal-exactness, which answers Part VII’s structural-subclass question negatively in part (dated notes), and **signal-machine-collision-certificates**, which shares no theorem. A dated note (batch 79) applies the finalizer of **smooth-diophantine-finalizers** to Part V’s Boolean-circuit quartics, whose zero sets are singular at every real zero. Merged from thirteen manuscripts; not a Lean formalization.

174. Beyond halting: recurrence, progress deadlines and Diophantine liveness

[hilbert-tenth-problem/liveness-beyond-halting/article.pdf](#)

from `Beyond_Halting_Diophantine_Liveness.zip`, `Quadratic_Diophantine_Dynamics.zip` (batch 62);
batch 63: `quadratic_infinite_computation.zip`; batch 79: `clock_spectra_research.zip`

For computable finitely branching systems, existence of a recurrent run with a computable progress deadline is Σ_3^0 -complete and equivalent to a nonempty effectively closed class of accepting schedules, and to an accepting schedule of hyperimmune-free degree; fixed deadlines are Π_1^0 , bounded gaps Σ_2^0 , unrestricted recurrence Σ_1^1 -complete, with strict separations and every c.e. degree as an exact deadline spectrum. An auxiliary-free degree-two polynomial with coefficients at most eight gives the transition relation of counter networks (degree two is minimal); a second, incoming-edge encoding has height six, a sum-of-squares quartic companion and an exact planar realization; and a two-stack quartic certifies finite runs meeting prescribed deadlines. Part VI (batch 79) answers in part the report’s question on deadline spectra beyond c.e. cones: clock problems are, up to uniform computable transformations, the upward effectively closed mass problems of Baire space; every Δ_2^0 degree, and every degree of a unique path through a recursive tree (among them $\mathbf{0}^{(r)}$ and $\mathbf{0}^{(\omega)}$), is a least clock degree of an input of a fixed two-stack system; unions of incomparable cones have no least degree, and a least clock degree is hyperarithmetical and contains a self-modulus. Its degree-two two-stack certificate is a second route, at lower degree, to the report’s quartic compiler. Merged from four manuscripts; not a Lean formalization.

175. Groups as Diophantine substrates

[hilbert-tenth-problem/group-theoretic-substrates/article.pdf](#)

from batch 76: `arithmetic_van_kampen.zip` + `three_commutative_phases_research.zip`; batch 78:
`Order_Is_Not_a_Moment.zip` + `ProveIt_Affine_Matrix_Diophantine_Research.zip`

Low-degree certificates for computation in groups, all through the integral Heisenberg group H . Part I: with Sanov’s chart (the free group of rank two as the integral points of $x + t + 4xt - yz = 0$), quartics whose zeros are in bijection with padded factorizations and that accept exactly the words of van Kampen area at most m , with $(2s + 13)m - 4$ witnesses, so that minimal arity tracks the Dehn function; proof-DAG compression, and exactly $k + \ell$ product gates for $[a^{2^k}, b^{2^\ell}]$, optimal in the stated calculus; a height bound rests on an imported Cornulier–Tessera lemma. Part II: three injective commuting phase homomorphisms into $H^N \times \mathbb{Z}^m$ whose ordered product realizes every integral quadratic system fibre-bijectively, hence, with MRDP, three fixed free abelian subgroups of a Heisenberg power with c.e.-complete product membership, a proposed, unrefereed negative answer to the three-subgroup question of König–Lohrey–Zetzsche and Roman’kov; two-phase decidability, known for integer exponents and credited, is new only for natural and mixed exponents. Part III (batch 78): for three letters the realizable pair counts K_{AB} at fixed K_{AC}, K_{BC} form a gapless interval, with a 24-witness quartic single-fold over complete profiles for $c = 2$, holes for four letters, and decidable three-generator submonoid membership in H^r given one multiplicity, a partial answer to a question of the Hilbert’s-tenth programme. Part IV (batch 78): the hit set of an affine unimodular curve in any subgroup of $SL_d(\mathbb{Z})$ is finite (at most $d - 1$ points) or exactly periodic, with sharp periods and no uniform algorithm; input degree two, via the repository’s quadratic loader (reconstructed and attributed), is the least that loads a noncomputable c.e. set; and an affine abelian compiler into $SL_{6g+2}(\mathbb{Z})$. Built from four manuscripts that share no theorem, the Heisenberg setup printed once; one Mihailova step is a second route to the probabilistic-quantum report’s. Not a Lean formalization.

176. Collision geometry is linear: Diophantine certificates for rational signal machines and conservative particle dynamics

[hilbert-tenth-problem/signal-machine-collision-certificates/article.pdf](#)

from batch 78: Collision_Geometry_Diophantine_Signal_Machines.zip +

Signal_Machine_Diophantine_Certificates.zip; batch 79:

Conservative_Signal_Frontend_Corrected.zip, Sparse_Lattice_Diophantine_Certificates.zip;

superseded, not printed: Conservative_Signal_Diophantine_Frontend.zip

A rational signal machine computes by collisions of finitely many kinds of particles moving at fixed rational speeds on a line. Once a complete finite collision history is fixed, its realizations are exactly the natural zeros of a sum of squares of integer affine residuals, of degree two, with one witness tuple each; typed endpoint conditions on maximal adjacency intervals exclude unrecorded and omitted simultaneous collisions, and causal depth controls the denominators. Part I eliminates every event coordinate: the realizing initial gaps form a relatively open rational polyhedral cone cut out by at most $3(n - 1) + (3m + 1)E$ conditions, which gives small integral realizations, an NP upper bound for bounded synthesis, a canonical first-hit quartic, exact quasi-polynomial seed counts, an exactly solved first-collision tournament and a four-speed accumulating clock with denominators $2 \cdot 3^k$. Part II keeps event times and positions as witnesses on a rational grid, adds halting, proves full column rank (strict convexity in the witnesses) and bounds the canonical witnesses by $60^K \max(1, M_0)$ for the four-speed interface of Durand-Lose’s universal machine, which is not transcribed. Part I’s convex-quadratic barrier is a special case of Part XI of [canonical-diophantine-certificates](#), printed as a second route. No fixed-arity polynomial for unbounded halting is claimed. Part III (batch 79) writes out a fixed reversible, number-conserving universal signal machine with 114 meta-signals, 445 two-to-two rules and exactly 18 live signals on every encoded input, compiled from Morita’s 15-state, 6-symbol reversible universal Turing machine through Durand-Lose’s conservative stacks; its event dynamics is an 18-dimensional partial piecewise homogeneous integer-linear map, and every fixed horizon has an explicit canonical quadratic certificate (4,196,998 auxiliaries per step for the literal machine). This answers Part I’s question on an audited universal frontend, except for an ordinary-integer loader; the manuscript is the corrected edition of an earlier delivery, which is superseded. Part IV (batch 79) compiles a fixed horizon of one-dimensional partitioned number-conserving lattice dynamics into one quartic sum of squares with a unique natural witness, corrects a sign in Morita and Imai’s construction, and proves decidability and effective semilinearity at total mass at most two. Four large packet and fixture files of Parts III and IV are regenerated rather than stored. *Parts I and II are two independent manuscripts proving one theorem by two routes, both printed in full; Parts III and IV come from two further manuscripts; not a Lean formalization.*

177. Stochastic and thermal exactness: exact events in uniformly convergent systems

[hilbert-tenth-problem/stochastic-and-thermal-exactness/article.pdf](#)

from batch 78: ProveIt_Exact_Erasure_Research.zip + ProveIt_Exact_Convergence_Research.zip +

Thermal_Arithmetic_Diophantine_Hamiltonians.zip; batch 79: Mixing_Does_Not_Remove_Arithmetic.zip,

Coercive_Green_Diophantine.zip + Well_Conditioned_Diophantine_Computation.zip

One pattern in five settings: an approximation converges uniformly at a known rate, while an exact event of the same system is undecidable. Part I lifts integer matrix mortality to strictly positive rational column-stochastic matrices with no common invariant complex line, entries in any band around $1/n$ and total-variation contraction η^T ; a product has rank one exactly when the source product vanishes, so exact erasure is undecidable for at most eight positive 7×7 matrices (from an imported mortality bound) and many-one complete for a nine-state family with a variable alphabet, with horizon- T quartics having one witness per erasing word. Part II builds a fixed turn-based stochastic game with discount $1/2$, reward $1/4$ and full-support dyadic rows: value iteration converges at rate 2^{-n} , yet exact attainment of the known fixed point $\frac{1}{2}\mathbf{1}$ from a dyadic terminal payoff is c.e.-complete; the one-player case stays open. Part III turns natural arithmetic circuits into diagonal Hamiltonians $H = (1 + \sum_i n_i)Q$ of degree at most five whose zero states biject with the witnesses; compact resolvent and finite thermal trace are equivalent to finite witness multiplicity and undecidable on a padded universal family, and a uniformly computable critical

fugacity satisfies $t_* < 1$ exactly on halting inputs. The quadratization, no-cutoff and finite-fold steps are those of **canonical-diophantine-certificates**, printed as pointers; defects that the research tree found in three shipped programs are disclosed, not patched, and Part II's two largest certificates are regenerated by its code rather than stored. Part IV (batch 79): every nonzero rational polynomial P of translation rank $r(P)$ is realized by $s = r(P) + 1$ states (optimal) with commuting, invertible, strictly positive doubly stochastic rational matrices in any band around $1/s$, acceptance $1/s + \varepsilon q^{|n|} P(n)$ and total-variation contraction θ^L ; on a fixed alphabet, with the input on a rational quadratic Bézier curve or a line segment, the exact hit of the equilibrium value is c.e.-complete, with no computable horizon. It reuses Part I's lift, printed as a pointer. Part V (batch 79, two manuscripts proving one theorem by two routes): uniformly coercive operators $\alpha I - A$ on computable path forests, and on a connected graph of maximum degree three, with polynomial-time approximable inverses, whose unique normalized integer Green certificates exist exactly on halting inputs; the rational Green values are classified by a Pell equation, finite support and attainment of a strongly convex energy are undecidable, and membership over $\mathbb{Z}[S^{-1}]$ is decidable for every finite prime set S . *Merged from six manuscripts; Parts I–IV share no theorem; a sibling of probabilistic-quantum-and-continuous-computation*, not a Lean formalization.

178. Quadratic orthant certificates: maximal parallelism, timed irreversible races, a fixed Waterfall universal machine, active membranes and reset Petri nets

hilbert-tenth-problem/quadratic-orthant-certificates/article.pdf

from batch 78: Maximal_Parallel_Diophantine.zip + Total_Quadratic_Diophantine_Semantics.zip + Waterfall_Diophantine_Certificates.zip; batch 79: Membrane_Motif_Research_Package.zip, Universal_Membrane_Research_Package.zip, Reset_Petri_Net_Certificates.zip

Parts I–III, V and VI compile fixed finite systems into integer polynomials of degree two, nonnegative on the real orthant, whose natural zeros biject with the semantic objects and whose unsquared products express a condition that balance equations cannot; Part IV uses degree four. Parts I–III sit at the level $D_2^+ = \text{SL}$ of the degree classification in Part XI of **canonical-diophantine-certificates**. Part I certifies one maximal parallel round of a multiset-rewriting system with $d + 2m + 4H$ natural auxiliaries, and also histories and exact first halting; the resource-cover rank $\rho(A)$ gives the sharp counting exponent $m - \rho(A)$, fixed-horizon counts are eventually quasipolynomial, counting is $\#P$ -complete, the deadlock of $A + B \rightarrow C$ has no convex quadratic, and a program-uniform certificate needs degree four. Part II gives irreversible exclusive-site systems with positive integer delays a total semantics: one quadratic per catalogue has exactly one natural zero for every delay vector, recording the earliest-completion outcome and certified permanent absence, which has no computable grounding bound. Part III proves the exact macrostep behaviour of Iijil's fixed 46-clock Waterfall program for the Neary–Woods universal machine $U_{15,2}$ and compiles k steps into one quadratic with $35k$ witnesses and an empty or singleton fibre; its arity grows with k . Re-proofs and generalizations of Part XI's results are printed as second routes; the research tree's reduction to 107 witnesses at $k = 7$ and its independent finding of the same Neary–Woods erratum are credited. Iijil's two machine data files are third-party and not under MIT-0. Parts IV–VI were added in batch 79. Part IV certifies one maximally parallel step of polarizationless noncooperative active membranes with division and dissolution by a sum of squares of quadratic residuals (degree four, one polynomial per finite schema), with exact inclusion maximality, an exact variable and residual ledger, the sharp next-population bound $2^n - 1$ and a four-rule system that defeats identical-subtree sharing. Part V gives two literal universal active-membrane frontends for the Neary–Woods machine (affine: 528 three-counter instructions and 2,544 rules; prime: 8,408 two-counter instructions and 34,605 rules), both c.e.-complete, the transcendental density of the prime language, of Turing degree $0'$, and outcome quadratics whose products are strong selector gates, so that every zero over the real orthant is natural and every fibre is empty or a singleton. Part VI builds a unit-weight reset Petri net (539 places, 771 transitions, 233 reset arcs from three places) with c.e.-complete exact-target reachability, exactly one accepting word at each length of an arithmetic progression and a canonical certificate with $2813h$ witnesses, natural over the nonnegative reals; it re-derives

Part V’s program layer without citing it, which is printed once. Seventeen heavy schemas and traces of Parts V and VI are regenerated rather than stored. *Merged from six manuscripts: the first three share no theorem, and Parts IV and V are complementary*; not a Lean formalization.

179. Polynomial witness histories: unique linear certificates with finite polynomial unknowns

[hilbert-tenth-problem/polynomial-witness-histories/article.pdf](#)

from batch 79: `unique_polynomial_histories.zip` + `one_coordinate_certificates.zip` + `Linear_Boundary_Transport_Research.zip`

Certificates whose unknowns are a fixed number of finite polynomials rather than integers: identities linear in them (in Part II of degree two, every product through one shared stride), a clock that makes the whole witness unique with 0/1 coefficients, and no computable input-only bound on its degree or support. Part I compiles cellular computation over $\mathbb{N}[X, Y]$ with $s^3 + s + 5$ polynomial unknowns and $3s + 3$ affine-linear rows (or $3\lceil \log_2 s \rceil + 6$ unit-coefficient rows, or nine weighted rows), with a unique Boolean-coefficient witness exactly at a first singleton acceptance, exact degrees, a Σ_1^0 -complete fixed matrix and a single quadratic. Part II answers Part I’s question on one coordinate negatively: coefficient-bounded univariate linear systems over $\mathbb{N}[X]$ are decided by a finite-state procedure, so no universal univariate linear compiler has computably bounded witness heights; eleven degree-two identities with the self-calibrating stride $S = X^{m+2h+2}$ keep the full unique Boolean history, in bijection with Part I’s witnesses. Part III, independent of both: linear boundary transport uniform over every nonzero commutative ring, a compiler with $q + d$ linear identities in $q + 2d$ sorted polynomial unknowns, a row-tagged single equation, an exact integer/rational energy gap and a Hilbert-space range theorem. A fixed number of polynomial unknowns is not a fixed number of integer unknowns: no ordinary fixed-arity representation is claimed, the no-bound results are the bounded-search argument of Part IX of **canonical-diophantine-certificates**, and undecidability of linear equations over polynomial semirings is older work (Narendran). Three expanded-polynomial exports of 1.0–1.6 MB are regenerated by the shipped code rather than stored. *Merged from three manuscripts, one answering another*; not a Lean formalization.

180. Smooth everywhere, universal on the integers: smooth Diophantine finalizers

[hilbert-tenth-problem/smooth-diophantine-finalizers/article.pdf](#)

from batch 79: `Smooth_Quartic_Diophantine_Certificates.zip`

For integer polynomials f_i of degree at most two with homogeneous quadratic lifts $q_i(x, t)$, the finalizer $F = \sum_i q_i(x, t)^2 + (tu - 1)^2 + g(z_1) + g(z_2) + 2g(z_3)$, $g(z) = 2z^2 - z$, is a quartic in five more variables whose natural zeros are in bijection with those of the system and whose integer zeros are exactly two sign-related copies of its integer zeros, while its affine scheme is smooth over $\mathbb{Z}$ with geometrically integral fibres, certified by an explicit integral Bézout identity $AF + \sum_v B_v F_v = 1$ of product degree at most seven. It has a nonnegative point over $\mathbb{Z}[1/2]$, points over every $\mathbb{Z}_p$ and modulo every integer, rational points dense in its real locus and exactly two real path components, so no local obstruction remains. With first weight one, universal smoothness holds exactly for total guard weight 4 or 16, and three guards (five auxiliaries) are optimal within positive weight-four guard families; also a bounded counter-program frontend with a unique natural witness, c.e.-completeness on the compiled class with no computable bound on the least integer witness, and power-of-two padding to higher degree. General smooth-variety undecidability is Poonen’s; the counter adapter and the quartic normalization are those of **canonical-diophantine-certificates**, credited; no finite-fold or single-fold MRDP theorem is claimed. A dated note in **probabilistic-quantum-and-continuous-computation** applies the finalizer to that report’s Boolean-circuit quartics. From one manuscript; not a Lean formalization.