

Iterated Euler diagonals (OEIS A290353 and A290354)

Kotěšovec's asymptotic form, the first logarithmic correction,
and the logarithmic height crossover

Part IV: a common density for parabolic iteration

Three research reports of one external research session
(bundle Reports 225, 226 and 228), merged into one ProveIt report

Part IV: research companion prepared for Vladimir Reshetnikov,
developed with OpenAI ChatGPT

Parts I–III: 5 October 2026 Part IV: 8 October 2026 merged: 6 and 9 October 2026

Abstract

Let $F_0(z) = z$ and $1 + F_{m+1}(z) = \prod_{d \geq 1} (1 - z^d)^{-[z^d]F_m(z)}$, so that $A(n, m) = [z^n]F_m(z)$ counts unlabeled rooted trees with n leaves, all at level m (OEIS A290353), and $a_n = A(n, n)$ is OEIS A290354. This report prints three manuscripts as Parts I–III, in the order in which each depends on the one before. Part I proves $a_n \sim c(2n/\pi)^n n^{-4/3}$ with $c = T^{1/3}e^{-B}h(1) > 0$, $T = \pi/2$, $B = \frac{1}{2} - \frac{\pi}{6} - \frac{\pi^2}{16} - \frac{\log 2}{6}$, where h is the smooth, strictly positive density of a positive measure whose Laplace transform is the entire parametrization P of $w \mapsto e^w - 1$ with normalized Fatou coordinate $-2/w + \frac{1}{3} \log w + O(w)$; this proves the form of the asymptotic that Václav Kotěšovec stated as a conjecture on A290354 (August 14, 2017), while the digits of $c \approx 4.4923$ remain uncertified. Part I also proves proportional-height asymptotics for $A(n, m)$ with n/m in a compact subset of $(0, \infty)$, fixed and square-root height shifts with the floor phase, and a Lambert- W inverse at exact sequence values. Part II proves the first logarithmic correction $1 + (-\frac{1}{18} \log^2 n + A \log n + C)/n + O(n^{-7/5})$, with A and C exact in $h(1), h'(1), h''(1)$ and a new clock constant κ in closed form in $\pi, \log 2$ and Catalan's constant, the corresponding uniform proportional-height correction and a sharpened inverse. Part III proves $A(n, m)/(2T^{-n}m^{n-1}) \rightarrow e^{-1/(3\lambda)}$ when $m/(n \log n) \rightarrow \lambda \in (0, \infty)$, with every fixed inverse-logarithm order, through an exact Hankel representation of h near zero, $h(0) = 2$, a smooth germ $t^{1/3}h(t)$ and rational-plus-zeta coefficients; it also gives Kaneiwa's normalization and smooth height prescriptions. Validated digits of c , higher corrections, effective constants, the regime $n/m \rightarrow \infty$ and a conjectured identity linking h with the amplitude of the labelled iterated Bell numbers (a neighbouring report) remain open. The report is unrefereed, and nothing in it is formalized.

Contents

Guide to this report	4
What is proved, and where	5
The OEIS statements, and the inverses	6
Notation across the three Parts	7

Provenance and merge decisions	9
What was checked, and what was not	10
Relation to neighbouring reports and formal projects	11
I Diagonal and proportional height asymptotics for iterated Euler transforms	12
1 Definition and statement	12
2 Subsidiary terms and the nonautonomous clock	14
3 Matching and the positive real orbit	15
4 The repelling coordinate with uniform sectors	17
5 The local limit and angular tails	19
6 The positive measure and its density	21
7 Coefficient extraction and the amplitude	22
8 Proportional height and fixed shifts	23
9 An inverse consequence	24
10 Exact computation and numerical diagnostics	25
11 Further questions and quantitative extensions	27
II First logarithmic correction for iterated Euler diagonals	29
12 Definition and main results	29
13 Analytic foundation from Report 225	31
14 Subsidiary forcing through fifth scaled order	33
15 Two further nonautonomous clock terms	34
16 Weighted defect and accumulated clock error	36
17 Finite Fatou expansion and forced matching	37
18 Uniform local limit on a growing strip	38
19 Derivative extraction and classical Fourier inversion	40
20 Uniform proportional height correction	42
21 A sharpened inverse formula	42
22 Exact computation and numerical diagnostics	43
23 Further questions and the boundary of the theorem	45

III	Logarithmic height crossover for iterated Euler transforms	47
24	The array and the crossover	47
25	Imported analytic interfaces and the scope of the result	49
26	Finite Fatou expansions with sectorial remainders	52
27	The endpoint value by direct Fourier inversion	52
28	An exact Hankel representation	53
29	A smooth endpoint germ and the logarithmic transseries	54
30	The rational and zeta coefficient structure	56
31	Transfer to shrinking degree to height ratios	57
32	Exact logarithmic cancellation and the crossover proof	58
33	Kaneiwa normalization and its first correction	59
34	Smooth suppression ratio inverses	59
35	Exact finite comparisons and reproducibility	62
36	Further questions and limits	63
37	Further questions and research (all three Parts)	65
IV	A common density for parabolic iteration	69
38	Provenance, inputs and notation of Part IV	69
39	Question, result and scope	73
40	Four analytic inputs and exact conventions	75
41	Positive coefficient measures and their limit	76
42	Identification, analyticity and endpoint consequences	78
43	A bound valid for every iteration depth	80
44	Summing the Bell local limit	81
45	Consequences for chain-length constants	83
46	Computation and reproducibility	83
47	Literature, provenance and research boundaries	85
48	Further research questions	86
A	A short proof checklist for independent review	87

Guide to this report

The three Parts are three manuscripts of the session bundle of Reports 1–243 (arrival commit 60f54ea06), placed in this directory by commit 47fc7a069 (batch 106). They study one object, the iterated Euler transforms F_m of $1 + x$ and their coefficient array $A(n, m)$:

Part	Bundle report	Delivered title	Labels
I	225 (base)	Diagonal and proportional height asymptotics for iterated Euler transforms	<code>ied:lead:</code>
II	226	First logarithmic correction for iterated Euler diagonals	<code>ied:corr:</code>
III	228	Logarithmic height crossover for iterated Euler transforms	<code>ied:log:</code>
IV	—	A Common Density for Parabolic Iteration (added 9 October 2026)	<code>ied:par:</code>

Arrangement. The base of the merge is Report 225, whose `report225.tex` was staged as this `article.tex`. It proves the leading theorems and constructs the analytic foundation (the nonautonomous clock, the sectorial Fatou coordinate, the positive Laplace density h), so it is printed first, as Part I, with its section, statement and equation numbers unchanged. Report 226 uses Part I’s analytic interface and refines its clock and its coefficient extraction to the first correction; it is Part II. Report 228 uses both, Part I’s coordinate and density and Part II’s absolute derivative estimates, and pushes the transfer to degree-to-height ratios shrinking like a power of $1/\log m$; it is Part III. The three Parts thus appear in dependency order, which is also the order of writing.

Embedded copies. Report 226’s archive carries Report 225’s source and PDF under `supporting/`, and Report 228’s carries those of Reports 225 and 226; all six files are byte-identical to the standalone deliveries (SHA-256 at placement, `cmp` again at the write). They are printed once each, as Parts I and II, and are not shipped. Report 226’s archive also carries byte copies of eight code and data files of Report 225 (its ledger says “copied unchanged”); they are shipped once, under Report 225’s prefix.

Restatements. Part II’s Section 13 and Part III’s Section 25 state, in their own words, the results they import (from Part I, and for Part III also from Part II). They are printed in full, with notes naming the Part I or Part II statement behind each item, because their displays are cited by the later proofs, because Part II states as interface items several estimates that Part I proves only inside its proofs (the inverse-orbit bound and the disk and neighbourhood forms of the tail and outer bounds), and because Part III’s contains material that is not a restatement (its L^1 consequence (III.16) and its comparison with the iterated Bell report). (Wording corrected after the independent check of 6 October 2026; the first wording read “because they contain material that is not a restatement (Part II’s bound on inverse orbits, Part III’s L^1 consequence ...)”). Part II’s bound on inverse orbits is the estimate $|G^j(u)| \geq c(|u| + j)$ of Part I’s proof of Lemma 4.1.)

Numbering. Sections are numbered continuously and statements within sections, so a statement number moves with its section only. Part I keeps Report 225’s Sections 1–11. Report 226’s Section k is Section $k + 11$ here (Sections 12–23): its Theorems 1.1 and 1.2 are Theorems 12.1 and 12.2, its Corollary 10.1 is Corollary 21.1. Report 228’s Section k is Section $k + 23$ here (Sections 24–36): its Theorem 1.1 is Theorem 24.1, its Lemma 3.1, Proposition 4.1, Theorems 5.1, 6.1, 8.1 are 26.1, 27.1, 28.1, 29.1, 31.1, and its Proposition 11.1 and Theorems 11.2–11.3 are 34.1, 34.2 and 34.3. The manuscripts number equations consecutively through each document; each Part keeps its

manuscript's equation numbers, prefixed in Parts II and III: Report 226's equation (k) is (II. k) here and Report 228's is (III. k). Section 37 was added in the write; no delivered number moved.

Notes added when the manuscripts were merged are set in small type and tagged **[write]**, with their date (6 October 2026). They give cross-references between the Parts, mark statements that a later Part extends or answers, mark results proved twice, and record what the write checked. Superseded statements and answered questions are printed as delivered, with a dated note. No delivered statement was changed, and no delivered symbol was renamed. Each Part has a section of open questions near its end: Part I its Section 11, Part II its Section 23, Part III its Section 36; Section 37 gathers them for the whole report under Vladimir's standing rule of 4 October 2026, with their sources, sketches and what is missing, and adds the question shared with the iterated Bell report.

[write] Part IV, added 9 October 2026 (second write). Since its second write the report has four Parts. Parts I–III, the front matter and Section 37 are the report as merged on 6 October 2026 and independently checked; the abstract and the front matter describe them alone, and nothing of them changed except this note, the Part IV row of the table above, dated notes at the ends of the front-matter sections “What is proved, and where”, “Provenance and merge decisions” and “Relation to neighbouring reports and formal projects”, a dated note at the end of Section 37, and the title, author line, date and PDF fields. Part IV (Sections 38–48 and Appendix A, labels **ied:par:**) is the manuscript *A Common Density for Parabolic Iteration* (8 October 2026, “Research companion prepared for Vladimir Reshetnikov / Developed with OpenAI ChatGPT”). From four analytic inputs proved in Part I here and in the iterated Bell report it proves the identity $I(t) = t 2^{t/3} h(t)$ for every $t > 0$ (Theorem 39.1), which answers item 9 of Section 37, and the holomorphy of h on $\Re t > 0$, which answers the positive-axis part of item 6; the germ at zero stays open. It also proves the geometric depth aggregate (Theorem 39.2), the strict-chain amplitude $C(q)$ holomorphic on $\Re q > 0$, Lengyel's constant $C = \frac{1}{2}(2 \log 2)^{(\log 2)/3} h(\log 2)$ (Corollary 39.3, (39.12)), and the endpoint expansion of I (Corollary 42.2). These are implications from the four inputs, which are unreferenced results of this report and of **a139383-iterated-bell-diagonals**; Section 38 prints that trust boundary, the provenance, the checks, the non-claims and the letters (L is $\log(1 + 1/q)$ in Part IV, $\log n$ in Part III; $H(n, m)$ is the labelled iterated Bell array, not a matching function).

What is proved, and where

Here $T = \pi/2$, $B = \frac{1}{2} - \frac{\pi}{6} - \frac{\pi^2}{16} - \frac{\log 2}{6}$, $f(w) = e^w - 1$, P is the entire solution of $P(s+1) = e^{P(s)} - 1$ whose inverse is the normalized Fatou coordinate $\Phi(w) = -2/w + \frac{1}{3} \log w + O(w)$, h is the density with $P(s) = \int_0^\infty e^{st} h(t) dt$, and $\beta = n/m$.

Statement	Where
P entire, the Laplace transform of a finite positive measure with every exponential moment, no atom at zero, support $[0, \infty)$; its density h is C^∞ and strictly positive on $(0, \infty)$	Part I, Theorem 1.1, Lemma 4.1, Proposition 6.1, Corollary 6.2
$a_n \sim c(n/T)^n n^{-4/3}$, $c = T^{1/3} e^{-B} h(1) > 0$, with an absolutely convergent Fourier formula for $h(1)$ (Kotěšovec's conjectured form on A290354)	Part I, Theorem 1.1
$A(n, m) \sim h(\beta) m^{-1} (m/T)^n m^{-\beta/3} T^{\beta/3} e^{-\beta B}$ uniformly for β in a compact subset of $(0, \infty)$; ratio e^k for fixed shifts; floor phase $e^{-\{\lambda n\}/\lambda}$; Gaussian-shaped profile for $ k \leq C\sqrt{n}$	Part I, Theorem 8.1, Corollaries 8.2 and 8.3
Lambert- W inverse	Part I, Corollary 9.1; sharpened in Part II,
$n = X + (\frac{4}{3} \log X - \log c)/q + o(1/\log X)$, nearest-integer recovery at exact sequence values (no onset)	Corollary 21.1

Statement	Where
Subsidiary forcing through fifth scaled order; two further clock terms M, N ; the clock constant $\kappa = \frac{\pi^3}{96} + \frac{\pi^2}{12} - \frac{13\pi}{48} - \frac{11}{36} - \frac{G}{6} - \frac{\pi \log 2}{24}$ (G Catalan)	Part II, Lemma 14.1, Section 15, (II.3)
$a_n = c(n/T)^n n^{-4/3} \{1 + (-\frac{1}{18} \log^2 n + A \log n + C)/n + O(n^{-7/5})\}$, A, C exact in $h(1), h'(1), h''(1), \kappa$	Part II, Theorem 12.1
The uniform first correction $\mathcal{Q}_\beta(d_m)/m$ for β in a compact subset of $(0, \infty)$	Part II, Theorem 12.2
Finite Fatou expansions to every order with rational a_j and differentiated sectorial remainders	Part II, Lemma 17.1 (to w^2); Part III, Lemma 26.1 (every order)
$h(0) = 2$; exact Hankel representation of h near zero; $t^{1/3}h(t)$ has a one-sided C^∞ germ; power-log expansion of h at zero (h is not C^1 at 0); $\log(h/2)$ coefficients	Part III, Proposition 27.1, Theorems 28.1 and 29.1, Corollary 29.2, Proposition 30.1
$c_j = \rho_j - \zeta(j)/(j3^j)$, $\rho_j \in \mathbb{Q}$	
Transfer $mA(n, m)r_m^n = h(\beta) + O(\log m/(m\beta^{30}))$ for all n ; relative equivalent on $(\log m)^{-D} \leq \beta \leq B_0$	Part III, Theorem 31.1
$A(n, m)/(2T^{-n}m^{n-1}) \rightarrow e^{-1/(3\lambda)}$ for $m/(n \log n) \rightarrow \lambda$, every fixed order in $1/(\lambda \log n)$, $c_2 = (9 - \pi^2)/108$; Kaneiwa's normalization shifts K to $K + 1$	Part III, Theorem 24.1, Section 33
Smooth suppression-ratio inverses, rounded height prescriptions, observed-ratio inverses	Part III, Proposition 34.1, Theorems 34.2, 34.3
<i>Open in all three Parts</i>	
Validated digits of $c, h(1), h'(1), h''(1)$; corrections beyond the first and an all-orders theorem; effective constants, onsets and certified thresholds; the regime $n/m \rightarrow \infty$ and the large-argument behaviour of h ; real analyticity of h on $(0, \infty)$ and analyticity of the germ at zero; polynomially small ratios and the transition to Kaneiwa's fixed-degree regime; other arrays and initial functions; the conjectured identity $I(\beta) = \beta 2^{\beta/3} h(\beta)$ with the iterated Bell amplitude. See Section 37.	

The finite computations of the three companions (exact integer recurrences, exact rational formal coordinates, symbolic clock identities) check identities, normalizations and coefficient algebra; no asymptotic statement rests on them, and every decimal value of $c, h(1), h'(1), h''(1)$ in the report is an uncertified floating diagnostic.

[write] **Part IV and this table, 9 October 2026.** Two entries of the last row are now proved in Part IV, conditionally on its four inputs (Section 40): the identity $I(\beta) = \beta 2^{\beta/3} h(\beta)$ with the iterated Bell amplitude, for every $\beta > 0$ (Theorem 39.1), and real analyticity of h on $(0, \infty)$, indeed holomorphy on $\Re t > 0$ ((39.6)). Analyticity of the germ at zero, and every other entry of the row, stay open. Part IV adds: the geometric depth aggregate $\sum_m e^{-Lm} H(n, m)$ of the labelled array (Theorem 39.2), the strict-chain amplitude $C(q)$ on $\Re q > 0$ and Lengyel's constant in terms of $h(\log 2)$ (Corollary 39.3), the endpoint expansion of I (Corollary 42.2) and density formulas for the constants K_1, K_2 of the chain-length cumulants (Corollary 45.1).

The OEIS statements, and the inverses

The entries as read by the write on 5 October 2026 (A290354 revision #26 of June 21, 2018; A290353 revision #28 of September 3, 2026):

- **A290354** (Alois P. Heinz, July 28, 2017): “a(n) is the n-th term of the n-th Euler transform of the sequence with g.f. $1+x$ ”, also “the number of unlabeled rooted trees with exactly n leaves, all in level n ”; formula “a(n) = A290353(n,n)” and the formula line, verbatim,

Conjecture: $a(n) \sim c * 2^n * n^{(n-4/3)} / \pi^n$, where $c = 4.4923\dots$ - _Vaclav
Kotesovec_, Aug 14 2017

Since $2^n n^{n-4/3} / \pi^n = (n/T)^n n^{-4/3}$, Theorem 1.1 proves this form, with $c = T^{1/3} e^{-B} h(1) > 0$ identified analytically. It does not certify the digits: the value $c \approx 4.4923299$ (Part I, Section 10; Part II's diagnostics give the same) is a floating-point evaluation without error control, and agrees with the OEIS's 4.4923... only numerically, as Part I says. The entry's b-file covers $n = 0..414$; the manuscripts compare only the 22 displayed terms ($n = 0..21$) and an independently generated fixture through 414, as they state. The entry cross-references A139383, the diagonal of the labelled iterated Bell numbers (see "Relation to neighbouring reports").

- **A290353** (Alois P. Heinz, July 28, 2017): "Square array $A(n,k)$, $n \geq 0$, $k \geq 0$, read by antidiagonals, where column k is the k -th Euler transform of the sequence with g.f. $1+x$ "; " $A(n,k)$ is the number of unlabeled rooted trees with exactly n leaves, all in level k "; generating function of column $k > 0$: $\prod_{j>0} 1/(1-x^j)^{A(j,k-1)}$. Its k is the manuscripts' height m (Part II writes $b_{n,m}$). The entry's cross-references name rows 2, 3, 4 as A001477, A000217, A000330, that is $A(2,m) = m$, $A(3,m) = m(m+1)/2$, $A(4,m) = m(m+1)(2m+1)/6$; these are the coefficients $[q^2]F_h$, $[q^3]F_h$, $[q^4]F_h$ computed in Part II's Lemma 14.1 (checked by the write against the exact recurrence for $m \leq 12$). The entry carries no asymptotic statement.

The inverses. Part I's Corollary 9.1 and Part II's Corollary 21.1 start from $X \log(X/T) = Y$. This is the factorial core **p0:prop:factorial-core** of the transseries volume [11] with $\kappa = 1$, its d equal to $-\log T$ and $L = Y$: its solution $\exp(W_0(Y/T) + \log T) = T e^{W_0(Y/T)}$ equals $Y/W_0(Y/T) = X$; after taking logarithms it is the Lambert core **p0:thm:lambert-core** with $a = b = 1$ and $L = \log(Y/T)$ in the variable $\log X - \log T$ (L named after the independent check of 6 October 2026). The corrections $X + \delta_0$ (Part I) and $X + \delta_0 + \delta_1/X$ (Part II) are proved directly, by Taylor expansion of an explicit smooth model and the mean-value theorem at the true integer; they parallel the first terms of **p0:thm:perturbed-inversion** but are not instances of it, because the perturbation $-\frac{4}{3} \log x + \log c + Q(\log x)/x$ is not a small analytic perturbation in that theorem's sense. The nearest-integer recovery at exact values $y = a_n$ resembles part (3) of **p0:thm:staircase**, but is not an instance either: its smooth root $x(y)$ solves a model equation and is not an admissible interpolation of (a_n) , and the manuscripts state no threshold for a general y between two terms. Part III's inverses (Section 34) involve no Lambert W : the model $\tau = \chi_n \beta - C(\beta)$ has the linear core $\chi_n \beta = \tau$ (the case $b = 0$ of **p0:thm:lambert-core**), perturbed by a germ C known to be C^∞ but not known to be analytic, so its finite reversion (Theorem 34.2) is proved by Taylor's theorem and the mean-value theorem, again an analogue rather than an instance of **p0:thm:perturbed-inversion**; its rounded height prescriptions are not instances of **p0:thm:staircase** (no monotonicity of $R(n, m)$ in m is available). None of the manuscripts claims novelty for the inversion mechanics.

Notation across the three Parts

Common to all Parts: F_h (the height index is h in Parts I–II and m in Part III's recurrence), $f(w) = e^w - 1$, $T = \pi/2$, B , Φ , P , the density h and its measure ν , $\beta = n/m$, the radius r , and the contour parameter s . Part II's $b_{n,m}$ is Parts I and III's $A(n, m)$. Part II's $d_\star = \frac{1}{3} \log T - B$, so its $D_n = n - \frac{1}{3} \log n + d_\star$ is Part I's D_n , and Part III's d_m is Part II's d_m . The symbols below change meaning between the Parts or inside one; each Part's text fixes its meaning there, and each Part opens with a short table of its own readings. No symbol was renamed.

Symbol	Meanings
h	The density of ν (all Parts) and the iteration height in F_h, R_h, E_h (Parts I, II; Part II's $t = hz$). <i>Tempting false reading:</i> in $F_h(z)$ and $[q^2]F_h = h$ the letter is the height, not the density.
K	Parts I, II: an integer entrance index ($M_0 \leq K \leq n$ in Lemma 3.3; $K = \lfloor \sqrt{n} \rfloor$ in Proposition 5.1 and (II.38); $K = \lceil At \rceil$ in Proposition 5.2); Part II also K_0 . Part III: the constant $K = \frac{1-\gamma+\log(T/2)}{3} - B$ (III.2), shifted to $K + 1$ by Kaneiwa's normalization; R_K is Kaneiwa's ratio. <i>False reading:</i> Part III's K is not an index.
κ	Parts II, III: the clock constant (II.3), $\kappa \approx -0.2543$. Part I: a small fixed $\kappa > 0$ bounding $K \leq \kappa n$ (before (23)).
ρ_j	Part II: $\rho_1 = h'(1)/h(1)$, $\rho_2 = h''(1)/h(1)$ and $\rho_j(\beta) = h^{(j)}(\beta)/h(\beta)$ (II.6). Part III: the rational numbers ρ_j , $j \geq 2$, of (III.6). <i>False reading:</i> Part III's $\rho_2 = 1/12$ is not Part II's $\rho_2 \approx -0.153$.
$c, C, \mathcal{C}$	$c = T^{1/3}e^{-B}h(1)$ the amplitude (Parts I, II). Part I: $C_z(w)$ the clock (10), C_λ (32), C_0 a fixed bound (Lemma 2.1). Part II: C the constant term of the correction (II.8), $\mathcal{C}_z(w)$ the extended clock (II.32), C_0 a real-width bound, C_{30} . Part III: $C_0 = 1 - \gamma - \log 2$, $\mathcal{C}$ Catalan's constant (Part II's G), $C(\beta)$ the smooth germ (III.43); c_j the log-density coefficients (III.6). All Parts: c_k the coefficients in the exact recurrence; C, c generic constants. The iterated Bell report's $C \approx 2.8654$ is a different amplitude.
$G, \mathcal{G}, g$	Part I: $G(u) = 2/\log(1 + 2/u)$ (Lemma 4.1), Part III's $\mathcal{G}(u)$; $G_n(s) = F_n(T/(D_n - s))$ (24); $g(w) = \log(1 + w)$ (also Part II). Part II: G Catalan's constant; $G_n(s) = F_n(re^{s/n})$ (exponential, not Part I's rational parametrization); $G_3(y)$. Part III: $G_m(s) = F_m(r_me^{s/m})$; $g(t) = t^{1/3}h(t)$ the germ.
L	Parts I, II: the clock function $L(y)$ (9), and a fixed integer L in Propositions 5.1 and 18.1; Part I also $L(x) = x \log(x/T)$ in the proof of Corollary 9.1. Part III: $L = \log n$.
W	The positive Lambert function (Parts I, II). Part III: $W(x) = P(-x)$ (proof of Theorem 28.1); Part III uses no Lambert function.
λ	Part I: $m = \lfloor \lambda n \rfloor$ (Corollary 8.2), so $\lambda \approx m/n$. Part III: $m/(n \log n) \rightarrow \lambda$. <i>False reading:</i> Part III's λ is not a slope m/n . The iterated Bell report's λ is m/n .
α, β	$\beta = n/m$ in all Parts; Part I also uses $0 < \alpha < \beta < \pi$ for angles (proof of Lemma 4.1), β_0, β_1 interval ends, $\alpha_n = \{\lambda n\}$. Part II: $\alpha = 4/3$ (Section 21), $\alpha = n - H(z) - \kappa z$ (proof of Proposition 18.1). Part III: B_0 the upper end.
H	Parts I, II: the matching function $H(z) = T/z + \frac{1}{3} \log(1/z) + B$. Part III: $H = H_{\theta,a}$ a Hankel contour, $H_j(X)$ the log-polynomials of (III.27). The iterated Bell report's $H(n, m)$ is its array.
a, b	a_n the diagonal (all). Part II: $a(y) = (1 + y^2)/2$ (Part I's $A(y)$), $b(y, t)$, u, v step coefficients, $b_{n,m}$ the array. Parts I–III: $b_j = \sum_{d j} d[z^d]F_h$. Part III: a_j the rational Fatou coefficients (Part I's Section 10 lists the first three), b_j the germ's Taylor coefficients. Part I: b a small constant (Proposition 5.2).
A	$A(n, m)$ the array (Parts I, III). Part II: A the log n coefficient (II.8). Part I: $A(y)$, and A a large constant (Proposition 5.2). Part III: $\mathcal{A}(w)$ (III.20).
d, D	Part I: $d = n - h$ (Lemma 3.2), D_n . Part II: $d = -\frac{1}{3} \log n + d_\star$, $d_m, d_y = 1 + y^2$, $D_n = n + d$. Part III: $d_m, D_N(u), D_t(v)$, D the band exponent (III.33).
R, r	Parts I, II: $R_h(z) = \sum_{j \geq 2} F_h(z^j)/j$; Part II also $R_n(s)$. Part III: the ratio $R(n, m)$ (III.3), R_2, R_K , the interval $\mathcal{R}$, and a radius R . r, r_m, r_n radii (all); Part III also a target ratio r .
Q	Part I: $Q(u) = u + \frac{1}{3} \log u$ (proof of Lemma 4.1), $Q_m(t)$ (proof of Theorem 8.1). Part II: $Q_2, Q_3, Q_\beta(d), Q_n(s), Q(u) = -u^2/18 + Au + C$. Part III: $Q_m(s)$, Part II's Q_n at height m .
M, N	Part II: the clock functions $M(y), N(y)$; also $M = n^{1/20}$, M_n, M_0 . Part I: constants M, M_0 . Part III: $M = m^{1/20}$; N a truncation order.
q	Part I: the entrance constant q , $q = \log(X/T) + 1$, q_m . Part II: the derivative order q ($= 30$), q_X . Part III: a truncation degree q .
E	Parts I, II: E_h the clock error. Part III: $E(m, \beta)$ the transfer error (III.32).

Symbol	Meanings
δ, τ	Part I: δ the tail range, τ a first crossing time (Lemma 3.2). Part II: δ_0, δ_1 inverse coefficients. Part III: $\tau = -\log r$.
X, y	Parts I, II: $X = Y/W(Y/T)$; $y = w/z$ the clock ratio; $y = a_n$ in the inverses. Part III: $X = \log t$; $y = m/n$, y_* , y_J height variables.
ℓ	Part II: $\ell = \log c$. Part III: $\ell_n = E_{n-1}/(n-1)!$ Kaneiwa’s coefficient.
I, Ψ	Part II: I a compact interval of ratios. Part I: Ψ the reciprocal Abel coordinate; Part III also $\Psi_{\text{IBD}} = -\Phi + \frac{1}{3} \log 2$, the iterated Bell report’s coordinate, and $I(\beta)$ that report’s amplitude (Section 25).

Provenance and merge decisions

Three manuscripts of the session bundle of Reports 1–243. Their author lines read “Report 225”, “Report 226” and “Report 228” (PDF author field “Research report”); none names a person, a tool or an addressee, says it is AI-assisted, or carries “prepared for private review” wording. All three are dated 5 October 2026. Report 225 cites the iterated Bell report at commit 65969409c and Report 228 at commit 5fdc0d68a; both commits exist in the repository, and that report’s `article.tex` is unchanged at HEAD (blob ab5014ec), so their section numbers refer to its current text. Report 226 pins no commit. They arrived in commit 60f54ea06 and were placed by 47fc7a069 (batch 106).

- **Part I**, bundle Report 225, archive `Report225.zip` (`src/report225.tex`, 639 lines, 16 pages); the base. Contributes the subsidiary-forcing estimate, the nonautonomous clock and its matching constant B , the noncircular real entrance, the sectorial Fatou coordinate and entire parametrization, the local limit and angular tails, the positive measure and smooth strictly positive density, the diagonal equivalent, proportional height, fixed and square-root shifts, the floor phase, the Lambert inverse, and Kaneiwa’s and Bechtloff Weising’s antecedents.
- **Part II**, bundle Report 226, archive `Report226.zip` (`src/report226.tex`, 715 lines, 17 pages). Its `supporting/` holds Report 225’s source and PDF byte for byte, and eight of its code and data files are byte copies of Report 225’s. Contributes fifth-order forcing, the clock functions M and N , the closed form of κ , the weighted fourth-power defect, the finite Fatou expansion and forced matching, the growing-strip local limit, thirtieth-derivative extraction, the first correction on the diagonal and uniformly in compact ratios, and the sharpened inverse.
- **Part III**, bundle Report 228, archive `Report228.zip` (`src/report228.tex`, 729 lines, 17 pages). Its `supporting/` holds the sources and PDFs of Reports 225 and 226 byte for byte. Contributes the L^1 form of Part II’s estimates, finite Fatou expansions to every order, $h(0) = 2$, the Hankel representation, the endpoint germ and its rational-plus-zeta coefficients, the transfer to logarithmic bands, the crossover theorem, Kaneiwa’s normalization, the smooth inverses, and the local identification with the iterated Bell coordinate together with the conjectured amplitude identity.

Where the merge had to choose:

1. *Order and base*. Dependency order I–II–III, with the base first (see the Guide). Statements that a later Part extends stay as delivered, with dated notes.
2. *Restatements*. Part II’s Section 13 restates Part I’s Lemma 2.1, the clock and entrance (Sections 2–3, (23)), the coordinate (Lemma 4.1), the tail and outer bounds (Proposition 5.2, Lemma 5.3) and the density (Proposition 6.1), and re-derives Corollary 6.2. Part III’s Section 25 restates Part I’s coordinate and density and Part II’s absolute estimates, then

derives a new L^1 bound. The intake dossier suggested replacing these sections by pointers, as the iterated Bell report did for its Part II; the write printed them in full, as the batch-105 model report did, because nearly every display in them is cited by the later proofs, because Part II's states as items estimates that Part I proves only inside proofs, and because Part III's contains material found nowhere else in the merge ((III.16) and the iterated Bell comparison). (Wording corrected after the independent check of 6 October 2026; the first wording read "and because each contains material found nowhere else in the merge".) No passage of one manuscript repeats another verbatim.

3. *Front matter and bibliography.* The three title blocks and abstracts are printed at the heads of their Parts; one table of contents replaces three. The three bibliographies are merged into one, keeping every detail of every entry; where two manuscripts describe an entry differently both descriptions are kept. The citation keys were renamed (Report 228's `oeis` is A290353, Reports 225 and 226's `oeis` is A290354); Reports 226 and 228's entries for Reports 225 and 226 now point to Parts I and II. The write added the entry for the transseries volume [11].
4. *Part III's table.* Report 228's `\input{crossover_table.tex}` is replaced by the table's text, byte for byte apart from its two leading comment lines, which are kept as L^AT_EX comments; the generated file is shipped as `data/228-logheight-results-crossover_table.tex`.
5. *Build-specific lines.* The manuscripts' `\pdfmapfile` lines (font-map additions for their TeX Live build) and Report 228's table-of-contents format tweak are not carried into the merged preamble; they affect no text.
6. *Write additions.* This front matter, the reading-conventions tables of the Parts, Section 37 with Remarks 37.1 and 37.2, the dated notes, labels for unlabelled sections, and the label prefixes. Everything else is delivered text.

[write] **A stale blob, 9 October 2026.** The statement above that the iterated Bell report's `article.tex` is "unchanged at HEAD (blob `ab5014ec`)" was true at this report's first write (`84fc1aaed`) and is stale since the same day: the reciprocal note proposed there was added to that report by commit `3f9fc9d4f`, whose blob `2b1d8fbe81` is that report's `article.tex` at HEAD and at Part IV's pin. The change is an added note; the section numbers the manuscripts cite are unchanged. Part IV (Section 38) is the fourth manuscript; its pin is `ProveIt 58175ca455`, at which this report's text is Parts I–III as merged and checked.

What was checked, and what was not

The report is unrefereed and not formalized. At intake (dossier of batch 106, 5 October 2026) the three manuscripts were read in full and no step was found false. Checked independently there: the first twelve terms of A290354 and the normalized ratios $a_n(T/n)^n n^{4/3} = 4.6013, 4.5529, 4.5178$ at $n = 40, 80, 200$ that Part I quotes, from an own implementation of the recurrence; the closed form of κ against a direct quadrature of the defining equation (II.23), -0.2543481698511 against -0.2543481698560 (difference from the truncated tail); that with Part II's diagnostic constants the corrected prediction tracks the exact a_n far better than the leading term (residual 1.4×10^{-4} against 2.5×10^{-2} at $n = 200$); and $2^{1/3}h(1)$ against the iterated Bell amplitude at $\beta = 1$ (Remark 37.1). The delivered suites pass on copies (README). The write checked the rows $A(2, m)$, $A(3, m)$, $A(4, m)$ against Part II's Lemma 14.1, the reversion coefficients (III.46)–(III.47), c_2 and c_3 from $S(t)$ and the Gamma expansion, the linear coefficient $C_0/3$ and K in (III.36), the reduction of Theorem 12.2 to Theorem 12.1 at $\beta = 1$, the floor amplitude C_λ of (32), the two orders of the inverse (II.53), and recomputed $c = 4.4923298968$ from the shipped diagnostic value of $h(1)$ (added after the independent check of 6 October 2026: an independent evaluation of $h(1)$ gives $c = 4.4923298971$; Remark 37.1).

What was *not* done: the analytic estimates (the clocks and their defects, the sector bootstraps, the tails, the Hankel deformations) were read and their displayed algebra spot-checked, but they are the manuscripts’ proofs, corroborated numerically, not an independent verification. The write did not read Kaneiwa (1980), Bechtloff Weising, Dudko–Sauzin, Prellberg–Mishna or Nagaev–Vakhtel; those citations and their page and theorem numbers are as the manuscripts and their source ledgers give them.

Relation to neighbouring reports and formal projects

The labelled analogue. SetTheory/Cardinals/docs/reports/generating-functions-and-asymptotics/oeis-sequence-asymptotics/a139383-iterated-bell-diagonals (labels ibd:) treats $H(n, m) = n![z^n](e^z - 1)^{\circ m}$, OEIS A139383 and A261280: a fixed-shift diagonal theorem to every order (its Part I) and proportional depth $m = \lambda n + \delta$ for λ in compact subsets of $(0, \infty)$ (its Part II), by the same parabolic map $w \mapsto e^w - 1$ without subsidiary forcing. No theorem is shared: the arrays differ, and Part I here says the labelled theorem “cannot be directly transferred” (Section 1.1), because the forcing changes the leading scale ($(n/T)^n n^{-4/3}$ here, $n^{2n-5/6}/(2^{n-1}e^n)$ there) and the clock constant. The two reports share the normalized orbit: Part III records $\Psi_{\text{IBD}}(w) = -\Phi(w) + \frac{1}{3} \log 2$ and the unproved relation $I(\beta) = \beta 2^{\beta/3} h(\beta)$ between that report’s amplitude I and the density h here (Section 25). The values agree to eight significant digits at $\beta = 1$ (Remark 37.1), and the relation, if proved, would bear on that report’s open endpoint $\lambda \rightarrow \infty$ (Section 37, item 9, and Remark 37.2). That report answers no question asked here, and this one answers none asked there; a reciprocal note is proposed for it.

Other neighbours. No other report of the collection treats A290353 or A290354 (repository search, 6 October 2026; A290354 occurs elsewhere only in a cross-reference line of the iterated Bell report’s OEIS data). The batch-106 reports a252782-diagonal-euler-transforms (one Euler transform with weights j^n) and a005121-strict-partition-chains (the depth sum $\sum_m 2^{-m-1} H(n, m)$ of the labelled array) share only words with this one. The transseries volume supplies the factorial core of which the leading inverses are instances (“The OEIS statements, and the inverses”). Placement in the collection confers no formal status: no Lean or Rocq file in the repository treats these sequences, and no statement of this report is formalized.

[write] **Part IV and the neighbours, 9 October 2026.** The two sentences above that no theorem is shared with the iterated Bell report and that the relation $I(\beta) = \beta 2^{\beta/3} h(\beta)$ is unproved describe Parts I–III. Part IV proves that relation for every $\beta > 0$ from inputs of both reports (Theorem 39.1), and gives the leading amplitude of a005121-strict-partition-chains (Lengyel’s constant) and its marked amplitude on $\Re q > 0$ (Corollary 39.3). Notes for those two reports, recording what Part IV answers there (the endpoint amplitude of I and the exploratory values of $I(1)$ in the iterated Bell report; the heuristic of `spc:q:depth` and, in part, `spc:q:disk` in the chain report), are proposed separately; they are not applied by this write.

Part I

Diagonal and proportional height asymptotics for iterated Euler transforms

Bundle Report 225, archive `Report225.zip`, delivered as `src/report225.tex` (16 pages), the base of the merge. Delivered title block “Diagonal and proportional height asymptotics for iterated Euler transforms / Report 225 / 5 October 2026”. Printed as delivered, with `[write]` notes; section, statement and equation numbers are those of the delivery.

Reading conventions of Part I	Elsewhere in the report
$A(n, m) = [z^n]F_m$; $a_n = A(n, n)$	Part II: $b_{n,m}$ for $A(n, m)$
K an integer entrance index; κ a small constant in $K \leq \kappa n$	Part III: K a constant; Parts II, III: κ the clock constant
$L(y)$ the clock function; $C_z(w)$ the clock; E_h	Part II: $C_z(w)$; Part III: $L = \log n$, $E(m, \beta)$
$G(u)$, $g(w) = \log(1 + w)$,	Part II: G Catalan, $G_n(s) = F_n(re^{s/n})$; Part III:
$G_n(s) = F_n(T/(D_n - s))$	$\mathcal{G}(u)$, $g(t) = t^{t/3}h(t)$
$m = \lfloor \lambda n \rfloor$ (Corollary 8.2)	Part III: $m/(n \log n) \rightarrow \lambda$

Abstract of this Part (as delivered). We study the number a_n of unlabeled rooted trees with n leaves, all at level n , equivalently the diagonal of the iterated Euler transform in OEIS A290354. We prove

$$a_n \sim c \left(\frac{2n}{\pi} \right)^n n^{-4/3}, \quad c > 0,$$

and identify c by a normalized repelling parametrization of $w \mapsto e^w - 1$. The main estimate is a uniform intermediate-angle bound, obtained from an explicit nonautonomous clock, a positive-coefficient Cauchy entrance, and a forced sector bootstrap. A positive Laplace measure identifies the coefficient amplitude and proves its strict positivity without assuming a density expansion at zero. Uniform proportional-height asymptotics for the array A290353, fixed and square-root height shifts, and a Lambert W inverse corollary are included. Numerical diagnostics give $c \approx 4.4923$, agreeing with Kotesovec’s conjecture; the decimal computation is not a validated enclosure. This is an unrefereed mathematical research report, not a proof-assistant formalization.

1 Definition and statement

Let

$$F_0(z) = z, \quad F_{h+1}(z) = \exp \left(\sum_{j \geq 1} \frac{F_h(z^j)}{j} \right) - 1. \quad (1)$$

Each F_h is analytic in $|z| < 1$, has nonnegative coefficients, vanishes at zero, and has linear coefficient one. These facts follow by induction; the sum over j converges normally on compact subdisks because $F_h(z) = O(z)$ at zero. For $n \geq 1$ put $a_n = [z^n]F_n(z)$ and set $a_0 = 1$ separately. The constant-term exception matches the OEIS convention and has no asymptotic effect.

For clarity, write $A(n, m) = [z^n]F_m(z)$ for $n \geq 1$, and put $A(0, m) = 1$ by convention. This is the square array A290353 [1], with degree n and height m ; the sequence A290354 is its main diagonal. A tree whose leaves all have level $m + 1$ is a root carrying a nonempty unordered multiset

of trees whose leaves all have level m . If there are $[z^d]F_m(z)$ types of a subtree with d leaves, the ordinary multiset generating function is

$$1 + F_{m+1}(z) = \prod_{d \geq 1} (1 - z^d)^{-[z^d]F_m(z)}. \quad (2)$$

Taking its formal logarithm gives (1). This also proves integrality and explains why there is no factorial normalization. The first diagonal values are

$$1, 1, 2, 6, 30, 170, 1337, 12166, 133476, 1676364.$$

The separate constant row is a generating-function convention, rather than a claim about a nonempty tree with no leaves.

Set

$$T = \frac{\pi}{2}, \quad B = \frac{1}{2} - \frac{\pi}{6} - \frac{\pi^2}{16} - \frac{\log 2}{6}, \quad D_n = n - \frac{1}{3} \log n + \frac{1}{3} \log T - B. \quad (3)$$

The logarithms used near the positive axis have their real determination there.

Theorem 1.1. *There is an entire function P satisfying*

$$P(s+1) = e^{P(s)} - 1, \quad (4)$$

whose inverse coordinate near zero is normalized by

$$\Phi(w) = -\frac{2}{w} + \frac{1}{3} \log w + O(w), \quad \Phi(e^w - 1) = \Phi(w) + 1.$$

It is the Laplace transform of a finite positive measure ν on $[0, \infty)$ with every exponential moment. This measure has a smooth, strictly positive density h on $(0, \infty)$. Moreover,

$$a_n \sim c(n/T)^n n^{-4/3}, \quad c = T^{1/3} e^{-B} h(1) > 0. \quad (5)$$

For every real σ ,

$$h(1) = \frac{e^{-\sigma}}{2\pi} \int_{-\infty}^{\infty} P'(\sigma + it) e^{-it} dt, \quad (6)$$

and this integral is absolutely convergent.

The normalization of Φ is essential: translating a Fatou coordinate translates P and changes the density amplitude. Formula (3) is tied to the zero constant term in the displayed normalization.

1.1 Prior work and the scope of the result

The OEIS entry attributes (5), with $c = 4.4923\dots$, to Václav Kotěšovec on 14 August 2017 and labels it a conjecture [2]. Kaneiwa's 1980 paper [3], Propositions 1 and 2 on printed page 19, proves that the unlabeled r -fold partition number $p(r; n)$ is a polynomial of degree $n - 1$ in r and identifies its leading coefficient through $\tan x + \sec x$. His indexing gives

$$p(r; n) = [z^n]F_{r+1}(z), \quad a_n = p(n - 1; n) \quad (n \geq 1).$$

Thus polynomiality, the tangent leading profile, and its $\pi/2$ pole are prior results. Fixed- n polynomial information does not alone justify the growing- n diagonal limit. Kaneiwa credits Bell's

1938 polynomiality result for the separately defined *labeled* function $\tilde{p}(r; n)$ on page 18; that sentence should not be read as attributing his unlabeled Proposition 1 to Bell.

Bechtloff Weising's higher-order Bell symmetric functions [4], Definition 2.1 and Proposition 2.12, specialize in one variable to $1 + F_{m+1}$; in particular $a_n = [m_{(n)}]B_n^{(n-1)}$. This is an algebraic and species connection, not a diagonal asymptotic theorem in that source. The related iterated-Bell report [5] concerns $n![z^n](e^z - 1)^{om}$ and cannot be directly transferred: the subsidiary forcing in (1) changes both the leading scale and the clock constant. No general priority claim for parabolic-coordinate methods is made here.

[write] **The OEIS statement and the neighbouring report, 6 October 2026.** The front matter quotes A290354's formula line verbatim as read on 5 October 2026: "Conjecture: $a(n) \sim c * 2^n * n^{(n-4/3)} / \Pi^n$, where $c = 4.4923\dots$ - _Vaclav Kotesovec_, Aug 14 2017". Theorem 1.1 proves this form with c identified; the digits 4.4923 are matched only by an uncertified evaluation (Subsection 10.2). Kaneiwa's indexing $p(r; n) = [z^n]F_{r+1}$ is used again in Part III, Section 33, which compares his leading coefficient with the crossover. The iterated Bell report [5] is the labelled analogue; Part III, Section 25, records the common normalized orbit and a conjectured identity between its amplitude and the density h here (Section 37, item 9).

2 Subsidiary terms and the nonautonomous clock

Write

$$R_h(z) = \sum_{j \geq 2} \frac{F_h(z^j)}{j}, \quad f(w) = e^w - 1.$$

The subsidiary estimate below is independent of the size of the main value $F_h(z)$.

Lemma 2.1. *Uniformly for $h|q|$ sufficiently small and $|q|$ small,*

$$F_h(q) = q + hq^2 + O((h + h^2)|q|^3). \quad (7)$$

Consequently, for each fixed C_0 , uniformly for $h \leq C_0/|z|$ as $z \rightarrow 0$,

$$R_h(z) = \frac{z^2}{2} + \frac{z^3}{3} + \frac{hz^4}{2} + O(|z|^4). \quad (8)$$

Proof. For real $0 < q < 1$, coefficient positivity and the absence of a constant term give $F_h(q^j) \leq q^{j-1}F_h(q)$. Hence the exponent defining $F_{h+1}(q)$ is at most $F_h(q)[- \log(1 - q)]/q$. While $F_h(q) \leq 2q$, the one-step increment is at most $CF_h(q)^2$. The elementary reciprocal comparison, or induction against $q/(1 - C'hq)$ with C' fixed large, proves $F_h(q) \leq 2q$ for $hq \leq c$ and c sufficiently small. Positivity gives the corresponding complex bound.

Summing one-step $O(|q|^2)$ increments gives $F_h(q) = q + O(h|q|^2)$. Substituting this back into the recurrence gives the sharper increment $F_{h+1}(q) - F_h(q) = q^2 + O((h + 1)|q|^3)$, because the quadratic main term and the leading subsidiary term each contribute $q^2/2$. Summation proves (7). Apply it at $q = z^j$ for $j \geq 2$. The $j = 2$ error is $O(h^2|z|^6) = O(|z|^4)$; the baseline tail $j \geq 4$ and the remaining corrections are bounded by geometric series. This proves (8). $\square$

[write] **Extended in Part II, 6 October 2026.** Part II's Lemma 14.1 extends (8) to fifth scaled order, from the exact coefficients $[q^2]F_h = h$, $[q^3]F_h = h(h + 1)/2$, $[q^4]F_h = h(h + 1)(2h + 1)/6$, which are the rows $n = 2, 3, 4$ of A290353 (checked by the write against the exact recurrence for $h \leq 12$).

Define, in a narrow sector about the positive real axis,

$$J(y) = 2 \arctan y - \frac{\pi}{2}, \quad L(1) = 0, \quad L'(y) = \frac{y^3/3 - y - 4/3 - 2J(y)}{(1+y^2)^2}. \quad (9)$$

Put

$$C_z(w) = \frac{J(w/z)}{z} + L(w/z). \quad (10)$$

The clock domain will be $|\arg(w/z)| \leq \eta$, $|w/z| \geq 1/2$, with a slightly wider fixed sector available for Taylor segments. In particular $C_z(z) = 0$.

Lemma 2.2. *Suppose $h \leq C_0/|z|$, $w_h = F_h(z)$ remains sufficiently small, and w_h/z remains in the stated clock domain through time k . Then, with $E_h = C_z(w_h) - h$,*

$$E_{h+1} - E_h = \frac{z^2(L(w_h/z) - E_h)}{1 + (w_h/z)^2} + O(|w_h|^2 + |z|^2), \quad (11)$$

with uniform constants. Consequently

$$|E_k| \leq C \left(\sum_{h < k} |F_h(z)|^2 + k|z|^2 \right). \quad (12)$$

Proof. By Lemma 2.1,

$$w_{h+1} = w_h + \frac{w_h^2 + z^2}{2} + \frac{w_h^3}{6} + \frac{w_h z^2}{2} + \frac{z^3}{3} + \frac{h z^4}{2} + O((|w_h| + |z|)^4). \quad (13)$$

Write $y = w_h/z$, $A(y) = (1+y^2)/2$ and $B_1(y, hz) = y^3/6 + y/2 + 1/3 + hz/2$. The increment of y is $zA + z^2B_1 + O((|w_h| + |z|)^4/|z|)$. Since $J' = 1/A$, the coefficient of z in the clock increment is

$$\frac{B_1}{A} - \frac{y}{2} + AL'(y) = \frac{hz - J(y)}{1 + y^2}.$$

But $hz - J(y) = z(L(y) - E_h)$, proving the cancellation in (11).

For completeness the remainder is analytic and uniform. Derivatives of $J(w/z)/z$ of order $r \geq 1$ are $O((|w| + |z|)^{-r-1})$, and derivatives of $L(w/z)$ are $O((|w| + |z|)^{-r})$. The displacement in (13), divided by $|w|$, is $O(|w| + |z|)$ because $|w/z|$ is bounded below. Nested clock cones therefore keep the full Taylor segment away from $w = \pm iz$. The omitted recurrence terms, the mixed quadratic Taylor terms, and the cubic Taylor remainder all contribute $O((|w| + |z|)^2)$ to the clock increment.

Finally $L(y)/(1+y^2)$ is bounded on the clock cone and $|1+y^2|$ is bounded below. The multiplicative discrete Gronwall factor is at most $\exp(Ck|z|^2)$, uniformly bounded in the stated range. Since $E_0 = 0$, this gives (12). $\square$

3 Matching and the positive real orbit

Lemma 3.1. *Uniformly in a fixed narrow sector about the positive real axis,*

$$L(y) = \frac{1}{3} \log y + B + O(y^{-2}) \quad (|y| \rightarrow \infty). \quad (14)$$

Consequently, when $|w/z|$ is sufficiently large in that sector,

$$C_z(w) = H(z) - \frac{2}{w} + \frac{1}{3} \log w + O\left(\frac{|z|^2}{|w|^3} + \frac{|z|^2}{|w|^2}\right), \quad H(z) = \frac{T}{z} + \frac{1}{3} \log \frac{1}{z} + B. \quad (15)$$

For real $0 < x \leq w \leq \epsilon$, with ϵ small fixed,

$$H(x) - C_x(w) \asymp \frac{1}{w}, \quad C'_x(w) > 0. \quad (16)$$

Proof. The derivative in (9) is $1/(3y) + O(y^{-3})$; in particular there is no y^{-2} term. To compute its constant, integrate the part $(y^3/3 - y)/(1 + y^2)^2$ first. After subtraction of $(\log y)/3$ its contribution from 1 to infinity is $-(\log 2)/6 - 1/3$. In the rest set $u = \arctan y$. Its contribution is

$$\int_{\pi/4}^{\pi/2} (\pi - 4/3 - 4u) \cos^2 u \, du = \frac{5}{6} - \frac{\pi}{6} - \frac{\pi^2}{16}.$$

Their sum is B in (3). The arctangent expansion gives (15).

For real $w \geq x$, the arctangent identity gives

$$H(x) - C_x(w) = \frac{2}{x} \arctan \frac{x}{w} - \frac{1}{3} \log w + O((x/w)^2).$$

The first term is between fixed positive multiples of $1/w$; the remaining terms are smaller for small w , uniformly for $x/w \leq 1$. The expansion of L may be bounded on all $y \geq 1$ by enlarging its constant on a compact interval. Finally the main derivative $2/(x^2 + w^2)$ dominates the $O(1/(x + w))$ derivative of the correction for small ϵ . This proves (16). $\square$

Lemma 3.2. Fix $\sigma \in \mathbb{R}$ and let $r = T/(D_n - \sigma)$. There are constants $M_0, c_0, c, C > 0$ such that, for all sufficiently large n , if $d = n - h \geq M_0$ and $x > 0$ satisfies $|x - r| \leq c_0 d/n^2$, then every real iterate through h stays in the small-clock range and

$$\frac{c}{d} \leq F_h(x) \leq \frac{C}{d}. \quad (17)$$

Proof. We have $H(r) = n - \sigma + O(\log n/n)$ and $H'(x) = O(n^2)$ when $x \asymp n^{-1}$. Make c_0 small, then M_0 large compared with $|\sigma|$. Uniformly in the displayed range,

$$d/2 \leq H(x) - h \leq 2d. \quad (18)$$

It remains to justify that the small-clock hypotheses are not circular. Let τ be the first time $F_\tau(x) \geq \epsilon$, if it occurs by time h . Start with a slightly larger admissible clock radius, so that one-step overshoot gives $F_\tau(x) \leq 2\epsilon$. Positivity gives

$$F_{j+1}(x) - F_j(x) \geq \frac{1}{2} F_j(x)^2, \quad \sum_{j < \tau} F_j(x)^2 \leq 2(F_\tau(x) - x).$$

Lemma 2.2 therefore gives $E_\tau = O(\epsilon + \tau x^2)$; here $\tau \leq h \leq n$ and $x \asymp n^{-1}$. Lemma 3.1 and monotonicity of C_x imply $\tau \geq H(x) - C/\epsilon$. If M_0 is large enough, this contradicts (18). Thus no such first crossing occurs. Apply (12) again through h and use (16); the clock error is $O(F_h(x) + nx^2)$ and can be absorbed into the $1/F_h(x)$ comparison. Equation (18) now gives (17). $\square$

Lemma 3.3. *With r as in Lemma 3.2, there is $q > 0$ such that for every integer $M_0 \leq K \leq n$ and every $|z - r| \leq qK/n^2$, the orbit through $n - K$ lies in the clock cone, satisfies $|F_h(z)| \asymp (n - h)^{-1}$, and obeys*

$$C_z(F_{n-K}(z)) = n - K + O(K^{-1} + n^{-1}). \quad (19)$$

Proof. At height $h \leq n - K$, use the disk about r of radius $R_h = c_0(n - h)/n^2$. Positivity and Lemma 3.2 bound its supremum by $F_h(r + R_h) \leq C/(n - h)$. On its inner half, Cauchy's estimate gives $|F'_h| \leq Cn^2/(n - h)^2$. Hence

$$|F_h(z) - F_h(r)| \leq C|z - r| \frac{n^2}{(n - h)^2}.$$

Divide by the positive lower bound in (17). Taking $q \leq c_0/2$ and then decreasing it makes the relative error at most $CqK/(n - h) \leq Cq$. Also $|z - r|/r \leq CqK/n$. Thus $F_h(z)/z$ lies in the required fixed narrow cone, with modulus at least $1/2$, for every $h \leq n - K$. The two-sided comparison follows. Finally $\sum_{h < n-K} |F_h(z)|^2 \leq C/K$, so Lemma 2.2 gives (19). $\square$

4 The repelling coordinate with uniform sectors

The needed coordinate is constructed here to make the sector and branch choices explicit.

Lemma 4.1. *For each closed angular sector strictly contained in $|\arg w| < \pi$, there is a sufficiently small radius on which an analytic function Φ satisfies*

$$\Phi(w) = -2/w + \frac{1}{3} \log w + O(w), \quad \Phi'(w) = 2/w^2 + O(1/w), \quad \Phi(f(w)) = \Phi(w) + 1, \quad (20)$$

whenever both arguments lie in the chosen domain. These functions agree on common smaller sectors. There is a local inverse on every sufficiently distant closed subsector of $|\arg(-s)| < \pi$, with

$$P_{\text{loc}}(s) = -2/s + O(\log |s|/|s|^2), \quad P'_{\text{loc}}(s) = 2/s^2 + O(\log |s|/|s|^3). \quad (21)$$

For small w in a smaller sector, $P_{\text{loc}}(\Phi(w)) = w$. The local inverse extends to the entire P in Theorem 1.1 and agrees with it on its sector of definition.

Proof. Let $g(w) = \log(1 + w)$ and conjugate by $u = 2/w$. The resulting map satisfies

$$G(u) = \frac{2}{\log(1 + 2/u)} = u + 1 - \frac{1}{3u} + O(u^{-2}).$$

For $Q(u) = u + (\log u)/3$,

$$Q(G(u)) - Q(u) = 1 + O(u^{-2}). \quad (22)$$

We give the uniform geometric estimate behind summation of this defect. Fix $0 < \alpha < \beta < \pi$. If $|\arg u| \leq \alpha$, then

$$|u + j| \geq \cos(\alpha/2)(|u| + j), \quad j \geq 0,$$

and the straight ray has distance at least $c_{\alpha, \beta}(|u| + j)$ from the complement of $|\arg v| < \beta$. Distinguish an inner analyticity radius R_0 from a larger starting threshold R_1 . Bootstrap

$$|G^j(u) - (u + j)| \leq a(|u| + j),$$

where a is less than half the preceding radial and angular margins. This gives $|G^j(u)| \geq c(|u|+j) > R_0$ if R_1 is large enough. Summing the one-step $O(1/|G^j(u)|)$ error improves the bound to

$$|G^j(u) - (u + j)| \leq C \left(\frac{1}{|u|} + \log(1 + j/|u|) \right).$$

Divided by $|u| + j$, this is uniformly $O(R_1^{-1})$ because $\log(1+x)/(1+x)$ is bounded. Increase R_1 to obtain a strict improvement and $cR_1 > 2R_0$. This closes the induction, including times when the straight ray decreases in modulus. Only the angular sector is invariant; the starting-radius boundary need not be.

The defects in (22) are now absolutely summable with total $O(1/|u|)$. Therefore

$$\Psi(u) = \lim_{m \rightarrow \infty} (Q(G^m(u)) - m) = Q(u) + O(1/u)$$

exists locally uniformly, is analytic, and satisfies $\Psi(G(u)) = \Psi(u) + 1$ wherever both sides are defined. Nested sectors and thresholds justify shifting the limit. Cauchy estimates on smaller sectors give derivative bounds. Define

$$\Phi(w) = -\Psi(2/w) + \frac{1}{3} \log 2.$$

This proves (20); $g \circ f$ is the identity near zero.

For the inverse, solve $\Psi(u) = -s + (\log 2)/3$. A disk of radius $C \log |s|$ centered at the right side stays inside a slightly larger allowed sector for large $|s|$. On its boundary the logarithmic correction to the identity map is smaller than the radius for C large. Rouché's theorem gives exactly one root, and $\Psi'(u) = 1 + O(1/u)$ is nonzero. The roots depend analytically on s , agree on overlaps by uniqueness, and give (21). If a small w already satisfies $\Phi(w) = s$ in a smaller sector, then $u = 2/w = -s + O(\log |s|)$ lies in the same uniqueness disk. Thus $P_{\text{loc}}(\Phi(w)) = w$; global injectivity of Φ is neither assumed nor needed.

First restrict P_{loc} to a strict left half-plane $\text{Re } s < -R_2$ so that $f(P_{\text{loc}}(s)) = P_{\text{loc}}(s + 1)$ holds when both parameters lie there. On $\text{Re } s < m - R_2$, define $P(s) = f^{\circ m}(P_{\text{loc}}(s - m))$. Successive definitions agree on overlaps, so they produce an entire function obeying (4). Agreement with P_{loc} on every connected larger inverse sector follows from the identity theorem, since there is agreement on a nonempty open part of the left half-plane. Positivity first far to the left and then forward iteration show $P(s) > 0$ for real s . $\square$

[write] Finite expansions of Φ elsewhere in the report, 6 October 2026. Part II's Lemma 17.1 proves $\Phi(w) = -2/w + \frac{1}{3} \log w - w/36 + w^2/540 + O(w^3)$ with differentiated sectorial remainders, and Part III's Lemma 26.1 the expansion to every order with rational coefficients a_j , both by summing the Abel defect along the inverse orbits constructed in this proof. The coefficients displayed in Subsection 10.2 are those a_1, a_2, a_3 .

Combining Lemmas 3.1, 3.3, and 4.1 gives the crucial entrance phase. Restrict $M_0 \leq K \leq \kappa n$ with $\kappa > 0$ small fixed, so that $|F_{n-K}(z)/z| \asymp n/K$ lies in the large-ratio matching range. Uniformly on the entrance disk,

$$\Phi(F_{n-K}(z)) = n - K - H(z) + O(K^{-1} + n^{-1} + K^3/n^2). \quad (23)$$

Both matching scales below satisfy this restriction.

5 The local limit and angular tails

Proposition 5.1. *Locally uniformly for $s \in \mathbb{C}$,*

$$G_n(s) := F_n\left(\frac{T}{D_n - s}\right) \longrightarrow P(s). \quad (24)$$

Here G_n is defined on each fixed compact set for all sufficiently large n .

Proof. Fix a compact set S and take $K = \lfloor \sqrt{n} \rfloor$. The point $z = T/(D_n - s)$ differs from the real reference T/D_n by $O_S(n^{-2})$, so Lemma 3.3 applies. Exactly,

$$H\left(\frac{T}{D_n - s}\right) = n - s + \frac{1}{3} \log \frac{D_n - s}{n}.$$

Thus (23) gives entrance phase $s - K + o(1)$ uniformly on S .

Choose a fixed integer L larger than the inverse-sector threshold and $2 \max_{s \in S}(|s| + 1)$. For $0 \leq j \leq K - L$, the prototype $s - K + j$ lies in a fixed interior left sector, has modulus comparable to $K - j$, and has real part bounded away from zero. Bootstrap phase error at most one. By (8), the actual w -step differs from $f(w)$ by $O(n^{-2})$. The disk of this radius about the unforced image lies in the coordinate domain: the image has distance at least $c/(K - j)$ from its angular boundary and zero. On that disk $|\Phi'| \leq C(K - j)^2$. Lemma 4.1 identifies the perturbed point with the same inverse branch. Thus the phase error at step j is at most $C(K - j)^2/n^2$, and the total is $O(K^3/n^2) = o(1)$. Together with the initial uniform error this strictly improves the bootstrap for large n .

At time $n - L$ the value tends uniformly to $P(s - L)$. In the remaining fixed L steps the subsidiary forcing is still $O(n^{-2})$. Finite compositions of the entire function f preserve convergence on these compact images, proving (24). $\square$

Proposition 5.2. *Fix $\sigma \in \mathbb{R}$ and put $r = T/(D_n - \sigma)$. There are $M, \delta, C > 0$ such that, for all large n and $M \leq t \leq \delta n$,*

$$|F_n(re^{it/n})| \leq C/t, \quad \left| \frac{re^{it/n}}{n} F'_n(re^{it/n}) \right| \leq C/t^2. \quad (25)$$

The same holds for negative t with $|t|$ in place of t .

Proof. Choose A large and $K = \lceil At \rceil$. Let $z_0 = re^{it/n}$. Since $|z_0 - r| \leq Ct/n^2$, choosing A after the entrance constant q and then choosing $b > 0$ small places

$$|z - z_0| \leq bt/n^2 \quad \text{inside} \quad |z - r| \leq qK/n^2.$$

Choose δ small enough that $K/n \leq \kappa$ and $K \leq n/2$. Expansion of H , using $H'(z) = O(n^2)$ on this disk and $H(r) = n - \sigma + O(\log n/n)$, gives

$$s_0 := \Phi(F_{n-K}(z)) = -K + it + O(1 + bt + t^2/n + t \log n/n + K^3/n^2). \quad (26)$$

The bounded term includes fixed σ . First choose b and δ to control proportional errors, then M to control fixed constants and the local inverse radius, and finally n to control $\log n/n$. The error in (26) is at most $t/10$.

Use the fixed prototype $p_j = -K + j + it$, $0 \leq j \leq K$, and bootstrap $|s_j - p_j| \leq t/4$. Then

$$\operatorname{Im} s_j \geq 3t/4, \quad \operatorname{Re} s_j \leq t/4, \quad |\arg(-s_j)| \leq \pi/2 + \arctan(1/3) < 2\pi/3, \quad ct \leq |s_j| \leq CK.$$

After increasing M , the inverse estimate gives

$$|\arg P_{\text{loc}}(s_j)| \leq 2\pi/3, \quad c/K \leq |P_{\text{loc}}(s_j)| \leq C/t \leq \epsilon/4.$$

Construct Φ beforehand on the wider sector $|\arg w| < 3\pi/4$, $|w| < \epsilon$, with further harmless room for nested-sector estimates. The same bounds hold for $s_j + 1$, with the next prototype index. The unforced image $f(P_{\text{loc}}(s_j)) = P_{\text{loc}}(s_j + 1)$ has distance at least c'/K from zero and the angular boundary, and at least $\epsilon/2$ from the outer circle.

The actual forcing is $O(n^{-2})$ by Lemma 2.1. Its full perturbation disk lies inside the wider coordinate domain for large n , since $K \leq n/2$. On this disk $|\Phi'| \leq CK^2$, so one step changes the ideal unit phase increment by at most CK^2/n^2 . The inverse-identification clause of Lemma 4.1, with a fixed slight angular enlargement, ensures that the actual perturbed value is $P_{\text{loc}}(s_{j+1})$. It does not switch branches.

The total additional phase error is at most $CK^3/n^2 \leq C'A^3\delta^2t$. Make this at most $t/20$. Combined with the initial $t/10$ error it is at most $3t/20$, strictly improving the $t/4$ bootstrap. Therefore $|F_n(z)| \leq C/t$ throughout the disk about z_0 . Cauchy's derivative estimate there yields $|F'_n(z_0)| \leq Cn^2/t^2$, proving (25). Conjugation proves the negative-angle case.

The noncircular order of constants is: fixed nested sectors and clock cones; small ϵ and inner analyticity radii; c_0, q ; large A ; small b ; small δ ; large M ; large n . All may depend on fixed σ , but none depends on n, t once chosen. $\square$

Lemma 5.3. *For each fixed $\delta > 0$, uniformly for $\delta \leq |\theta| \leq \pi$ on the same moving circle,*

$$F_n(re^{i\theta}) = O(n^{-1}), \quad F'_n(re^{i\theta}) = O(1). \quad (27)$$

Proof. Put $z = \zeta/n$ and $v_h = nF_h(z)$. On a bounded v -domain, the subsidiary estimate gives

$$v_{h+1} = v_h + \frac{v_h^2 + \zeta^2}{2n} + O(n^{-2}), \quad v_0 = \zeta.$$

The limiting ordinary differential equation has solution

$$v(u) = \zeta \tan(\zeta u/2 + \pi/4), \quad 0 \leq u \leq 1.$$

A pole at positive real $u \leq 1$ requires $\zeta = (\pi/2 + 2k\pi)/u$. On $|\zeta| = T$ only $\zeta = T$ causes a pole; the negative pole family has modulus at least $3T$. Thus the closed arc away from T has compact complex neighborhoods on which the full ODE trajectory is bounded.

Choose a bounded disk strictly containing those trajectories. Its vector field is uniformly Lipschitz, and the local discretization error is $O(n^{-2})$. Discrete Gronwall gives $O(n^{-1})$ error while the numerical orbit stays in the disk; this estimate prevents a first exit for large n . Hence $F_n = O(n^{-1})$ uniformly on the complex neighborhoods. Cauchy in the ζ variable gives $F'_n = O(1)$. The moving arc is contained in those neighborhoods because $nr \rightarrow T$. $\square$

6 The positive measure and its density

Proposition 6.1. *The entire function P is the Laplace transform of a finite positive measure ν on $[0, \infty)$ with every exponential moment, no atom at zero, and support $[0, \infty)$. Its restriction to $(0, \infty)$ has a smooth strictly positive density h .*

Proof. Put $r_n = T/D_n$ and define the finite positive measures

$$\nu_n = \sum_{m \geq 1} [z^m] F_n(z) r_n^m \delta_{m/n}.$$

Their actual Laplace transforms are $F_n(r_n e^{s/n})$. The rational parameter in Proposition 5.1 is

$$D_n - \frac{T}{r_n e^{s/n}} = D_n(1 - e^{-s/n}) = s + O_S(\log n/n)$$

on every compact S . Hence these transforms converge locally uniformly to $P(s)$.

The masses tend to $P(0) > 0$. Lévy's continuity theorem applied to the normalized characteristic functions gives a finite weak limit ν , supported on $[0, \infty)$. For every real u , the transforms at $u + 1$ are uniformly bounded. On $t > R$, the integral of e^{ut} is bounded by e^{-R} times the integral of $e^{(u+1)t}$. This proves exponential uniform integrability, so $P(u) = \int e^{ut} \nu(dt)$. Analyticity and the exponential moments extend this identity to every complex u .

By Lemma 4.1, $P(-x) \sim 2/x$. Thus $\nu(\{0\}) = 0$. Its support has positive points arbitrarily close to zero: a positive gap would imply exponential decay of $P(-x)$, contradicting $2/x$.

For fixed real σ , Lemma 4.1 also gives $P'(\sigma + it) = O(t^{-2})$ as $|t| \rightarrow \infty$, since a vertical line eventually lies in an interior inverse sector. This is the Fourier transform of the finite measure $ue^{\sigma u} \nu(du)$ and is integrable. Fourier inversion gives this weighted measure a continuous density. Dividing by $ue^{\sigma u}$ shows that $\nu(du) = h(u) du$ on $(0, \infty)$, with h continuous and nonnegative.

The functional equation (4) and uniqueness of finite measures give

$$(e^u - 1)\nu = \sum_{k \geq 2} \frac{\nu^{*k}}{k!}. \quad (28)$$

Both sides are finite: the left side has mass $P(1) - P(0)$, and the positive series on the right has finite total mass. In particular $\nu * \nu \leq 2(e^u - 1)\nu$, so its closed support S is closed under addition. Positive support points tending to zero have integer multiples approaching any prescribed positive number. Therefore $S = [0, \infty)$.

Fix $t > 0$. Full support and continuity yield an interval $I \subset (0, t)$ on which $h > 0$. Full support also yields a point $b \in t - I$ with $h(b) > 0$. Then $h(t - b) > 0$, and continuity implies $(h * h)(t) > 0$. The convolution is finite and continuous on $(0, \infty)$: near any fixed t , split the defining integral into two small endpoint pieces and a compact interior. On an endpoint piece one factor is bounded away from zero's neighborhood while the other has arbitrarily small L^1 mass near zero; the interior follows by dominated convergence. Thus the density inequality from (28), initially almost everywhere, holds everywhere by continuity:

$$(e^t - 1)h(t) \geq \frac{1}{2}(h * h)(t) > 0.$$

This proves strict positivity, without a pointwise density expansion at zero. $\square$

[write] **The density at zero, 6 October 2026.** Part III supplies the expansion this proof does without: $h(0) = 2$ (Proposition 27.1), an exact Hankel representation near zero (Theorem 28.1), a C^∞ germ $t^{t/3}h(t)$ (Theorem 29.1) and the power-log expansion (III.28); in particular h is continuous at 0 but not C^1 there.

Corollary 6.2. *The density h is C^∞ on $(0, \infty)$.*

Proof. Fix real σ . Cauchy's estimates in the inverse sectors of Lemma 4.1, applied on disks of radius proportional to $|s|$, give

$$P^{(q)}(\sigma + it) = O_q(|t|^{-q-1}) \quad (|t| \rightarrow \infty)$$

for every integer $q \geq 1$. This is the Fourier transform of the finite measure $u^q e^{\sigma u} \nu(du)$. Given $k \geq 0$, choose $q > k$. Then $|t|^k P^{(q)}(\sigma + it)$ is integrable. Differentiation under Fourier inversion gives a C^k density for this weighted measure. On $(0, \infty)$ it agrees with $u^q e^{\sigma u} h(u)$ by uniqueness and continuity. Divide by the positive smooth weight on every compact subinterval of $(0, \infty)$. Since k was arbitrary, h is smooth there. $\square$

[write] **Proved twice, 6 October 2026.** Part II restates this corollary with the same argument after its interface list (Section 13, around (II.16)).

7 Coefficient extraction and the amplitude

Proof of Theorem 1.1. All structural claims about P and ν were proved above. Fix real σ and put $r = T/(D_n - \sigma)$. Differentiating the Cauchy coefficient formula, or integrating it by parts over the full circle, gives exactly

$$na_n r^n = \frac{1}{2\pi} \int_{-\pi n}^{\pi n} \left[\frac{z}{n} F'_n(z) \right] e^{-it} dt, \quad z = r e^{it/n}. \quad (29)$$

For the derivative normalization, at $z = T/(D_n - s)$ one has

$$\frac{z}{n} F'_n(z) = \frac{D_n - s}{n} G'_n(s).$$

Locally uniform holomorphic convergence in Proposition 5.1 gives $G'_n \rightarrow P'$. On the circle, the actual parameter is $s_n = D_n - (D_n - \sigma)e^{-it/n} \rightarrow \sigma + it$, uniformly for bounded t . The central integrand therefore tends to $P'(\sigma + it)e^{-it}$.

Proposition 5.2 bounds the intermediate tails by $C \int_M^\infty t^{-2} dt = O(M^{-1})$, uniformly in n . Lemma 5.3 bounds $z F'_n(z)/n$ by $O(n^{-2})$ on outer arcs, whose t -length is $O(n)$, so their contribution is $O(n^{-1})$. First let $n \rightarrow \infty$ and then $M \rightarrow \infty$ in (29). Fourier inversion of the density of $u e^{\sigma u} \nu(du)$ at $u = 1$ gives

$$na_n r^n \longrightarrow \frac{1}{2\pi} \int_{\mathbb{R}} P'(\sigma + it) e^{-it} dt = e^\sigma h(1) > 0. \quad (30)$$

This also proves (6).

Finally, expansion of (3) yields

$$r^{-n} = (n/T)^n n^{-1/3} T^{1/3} e^{-B-\sigma} (1 + o(1)).$$

The factor $e^{-\sigma}$ from the contour radius cancels the factor e^σ in (30). Division by n proves (5) with the stated positive constant. $\square$

[write] **Refined in Part II, 6 October 2026.** Part II's Theorem 12.1 adds the first relative correction, with remainder $O(n^{-7/5})$, by extracting coefficients from the thirtieth derivative (Section 19) instead of the first.

8 Proportional height and fixed shifts

Theorem 8.1. *Let $0 < \beta_0 < \beta_1 < \infty$ be fixed. Uniformly over positive integer pairs (n, m) with $m \rightarrow \infty$ and $\beta = n/m \in [\beta_0, \beta_1]$,*

$$A(n, m) \sim \frac{h(\beta)}{m} \left(\frac{m}{T}\right)^n m^{-\beta/3} T^{\beta/3} e^{-\beta B}. \quad (31)$$

The exact integer height m and exact ratio $\beta = n/m$ occur in this formula.

Proof. Fix real σ , let $r_m = T/(D_m - \sigma)$, and extend

$$Q_m(t) = \frac{z}{m} F'_m(z), \quad z = r_m e^{it/m}, \quad |t| \leq \pi m,$$

by zero elsewhere. Proposition 5.1 gives convergence to $Q(t) = P'(\sigma + it)$ uniformly on each compact interval. Proposition 5.2 gives $|Q_m(t)| \leq C/t^2$ in the intermediate range; Lemma 5.3 gives $O(m^{-2})$ in the outer range. The limiting derivative also has an integrable $O(t^{-2})$ tail. Splitting the integral at a fixed M , then letting $m \rightarrow \infty$ and $M \rightarrow \infty$, proves

$$\|Q_m - Q\|_{L^1(\mathbb{R})} \rightarrow 0.$$

Their Fourier transforms consequently converge uniformly for all real frequencies, by the L^1 norm bound. At the exact frequency $\beta = n/m$, the coefficient identity is

$$nA(n, m)r_m^n = \frac{1}{2\pi} \int_{\mathbb{R}} Q_m(t) e^{-i\beta t} dt = \beta e^{\beta\sigma} h(\beta) + o(1).$$

Fourier inversion applies to the continuous density of $ue^{\sigma u}\nu(du)$. On the chosen compact interval, β is bounded away from zero and h has a strictly positive minimum. The absolute error therefore gives the uniform relative estimate

$$A(n, m) = \frac{e^{\beta\sigma} h(\beta)}{m} r_m^{-n} (1 + o(1)).$$

Write $q_m = -(\log m)/3 + (\log T)/3 - B - \sigma$. Since $n = \beta m$ exactly,

$$\begin{aligned} r_m^{-n} &= (m/T)^n \exp(\beta m \log(1 + q_m/m)) \\ &= (m/T)^n m^{-\beta/3} T^{\beta/3} e^{-\beta B - \beta\sigma} \left(1 + O((\log m)^2/m)\right), \end{aligned}$$

uniformly for these β . Cancellation of the σ factors proves the theorem. $\square$

Corollary 8.2. *For every fixed integer k , as $n \rightarrow \infty$,*

$$\frac{[z^n]F_{n+k}(z)}{a_n} \rightarrow e^k.$$

For fixed real $\lambda > 0$, put $\alpha_n = \{\lambda n\}$ and $m = \lfloor \lambda n \rfloor$. Then

$$A(n, m) \sim C_\lambda e^{-\alpha_n/\lambda} (\lambda n/T)^n n^{-1-1/(3\lambda)}, \quad C_\lambda = h(1/\lambda) \lambda^{-1-1/(3\lambda)} T^{1/(3\lambda)} e^{-B/\lambda}. \quad (32)$$

Proof. For $m = n + k$, continuity gives $h(n/m) \rightarrow h(1)$, while $(m/n)^n \rightarrow e^k$. All remaining ratios of prefactors in (31) and (5) tend to one. For the second assertion, $m = \lambda n - \alpha_n$ gives

$$(m/(\lambda n))^n = e^{-\alpha_n/\lambda}(1 + O(n^{-1})), \quad n/m = 1/\lambda + O(n^{-1}).$$

Substitution into (31), using continuity of h and $O(\log n/n)$ for changes in the logarithmic powers, proves (32). $\square$

The oscillating floor factor need not tend to one. Dropping it changes the leading equivalent. Neither theorem asserts uniformity as $\beta \downarrow 0$ or $\beta \rightarrow \infty$.

[write] **Extended in Parts II and III, 6 October 2026.** Part II's Theorem 12.2 adds the first correction on the same compact ranges. Part III's Theorem 31.1 extends the leading equivalent of Theorem 8.1 (the same formula, since $e^{\beta d_m} = m^{-\beta/3} T^{\beta/3} e^{-\beta B}$) toward $\beta \downarrow 0$, uniformly on every band $(\log m)^{-D} \leq \beta \leq B_0$, with an explicit relative error; this yields the crossover of Theorem 24.1. Uniformity as $\beta \rightarrow \infty$ remains open (Section 37, item 5).

Corollary 8.3. *For each fixed $C > 0$, uniformly over integers $|k| \leq C\sqrt{n}$,*

$$e^{-k} \frac{[z^n] F_{n+k}(z)}{a_n} = \exp\left(-\frac{k^2}{2n}\right) (1 + o(1)).$$

Proof. Put $m = n + k$ in Theorem 8.1. Uniformly in this range, $\beta = n/m = 1 + O(n^{-1/2})$, so $h(\beta)/h(1) \rightarrow 1$. The other prefactor ratios tend uniformly to one: their logarithms, aside from the continuous density ratio, are $O((1 + |k|) \log n/n)$. The exponential ratio is determined by

$$n \log(1 + k/n) = k - \frac{k^2}{2n} + O_C(n^{-1/2}).$$

Substitution and division by the diagonal equivalent prove the claim. $\square$

This is a Gaussian-shaped asymptotic profile for height shifts. It is not a probability central limit theorem; no probability distribution on heights has been introduced.

9 An inverse consequence

Corollary 9.1. *Let $y = a_n$ with $n \rightarrow \infty$, put $Y = \log y$, and let W denote the positive real Lambert function. Put*

$$X = \frac{Y}{W(Y/T)}, \quad q = \log(X/T) + 1.$$

Then

$$n = X + \frac{\frac{4}{3} \log X - \log c}{q} + o(1/\log X). \quad (33)$$

In particular $n - X \rightarrow 4/3$. The explicit corrected expression in (33), and the large real root $x(y)$ of

$$\log y = x \log(x/T) - \frac{4}{3} \log x + \log c,$$

recover n by nearest-integer rounding for all sufficiently large exact sequence values $y = a_n$.

Proof. Let $L(x) = x \log(x/T)$. Then $L(X) = Y$ and $L'(X) = q$. Theorem 1.1 gives $L(n) - L(X) = \frac{4}{3} \log n - \log c + o(1)$. First comparison gives $n/X \rightarrow 1$, and the mean-value theorem then gives $n - X = O(1)$. Thus

$$L(n) - L(X) = q(n - X) + O(X^{-1}), \quad \log n = \log X + O(X^{-1}).$$

Division by q proves (33). Applying the same mean-value argument to the full leading model gives $x(a_n) = n + o(1/\log n)$; the rounding assertions follow. $\square$

No effective onset is asserted. This is recovery from exact sequence values, not a universal threshold-rounding rule for arbitrary y between them. The leading equivalent alone does not provide a higher-order error term.

[write] **Sharpened in Part II; the transseries volume, 6 October 2026.** Part II's Corollary 21.1 sharpens (33) to an error $O(X^{-7/5}/\log X)$; its $\delta_0 = (\frac{4}{3} \log X - \log c)/q_X$ is the correction here. $X = Y/W(Y/T)$ solves $X \log(X/T) = Y$, an instance of the factorial core `p0:prop:factorial-core` of [11] ($\kappa = 1$, its $d = -\log T$). The correction and the rounding at exact values are proved directly by the mean-value theorem; they parallel `p0:thm:perturbed-inversion` and part (3) of `p0:thm:staircase` but are not instances of them (front matter, “The OEIS statements, and the inverses”).

10 Exact computation and numerical diagnostics

10.1 The coefficient algorithm

Logarithmically differentiating (2) gives

$$\frac{zF'_{h+1}(z)}{1 + F_{h+1}(z)} = \sum_{j \geq 1} b_j z^j, \quad b_j = \sum_{d|j} d[z^d]F_h(z).$$

Therefore, writing $1 + F_{h+1}(z) = \sum_{k \geq 0} c_k z^k$, one obtains

$$c_0 = 1, \quad c_k = \frac{1}{k} \sum_{j=1}^k b_j c_{k-j} \quad (k \geq 1). \quad (34)$$

Truncation at degree N is exact for every coefficient retained. The public standard-library program checks divisibility before each integer division. It stores F_h with constant term zero and restores the convention $A(0, h) = 1$ only when reporting the array. Degree and height are each capped at 640; this is an implementation and testing limit, not a limitation of the theorems. The diagonal calculation uses $O(N^3)$ integer arithmetic operations and $O(N)$ stored integers; operation costs grow with the bit lengths.

An independent product implementation expands each factor using

$$(1 - z^d)^{-a} = \sum_{k \geq 0} \binom{a + k - 1}{k} z^{dk} \quad (a \geq 1)$$

and multiplies the finite truncations. It checks every entry through degree and height 12. A separate generated exact fixture contains diagonal values through 414. The public tests also compare the 22 displayed OEIS values, $n = 0, \dots, 21$. The full OEIS b-file was not successfully retrieved in the

source review; the 415-entry fixture is therefore explicitly described as generated mathematical data, not as a 415-term external comparison.

All public pass/fail checks use explicit exceptions, so Python optimization does not remove them. Local chunked decimal conversion works with Python’s minimum configurable integer-string digit limit of 640 without modifying that process-wide limit. This matters because the largest exact integers exceed that many digits.

10.2 The amplitude computation is not certified

The normalized ratios $a_n(T/n)^n n^{4/3}$ decrease slowly over the tested range: approximately 4.60133 at $n = 40$, 4.55287 at $n = 80$, 4.52400 at $n = 160$, 4.51777 at $n = 200$, and 4.50437 at $n = 414$. These finite values alone neither prove a limit nor certify its decimal digits.

A separate optional numerical script evaluates the repelling parametrization. The first terms of its normalized formal coordinate are

$$\Phi(w) = -2/w + \frac{1}{3} \log w - \frac{w}{36} + \frac{w^2}{540} + \frac{w^3}{7776} + \cdots.$$

The script uses nine correction coefficients, solves the truncated inverse equation near $s - M$ by Newton iteration, applies M forward iterates of f , and propagates the derivative by multiplication by e^w . Subtracting $2/(1-s)^2$ from the derivative before quadrature improves the observed tail: its known contribution to $h(1)$ is $2/e$ for $\sigma < 1$. Tests varying the forward depth, vertical contour, cutoff, and mesh give values near 4.4923299, consistent with the conjectured 4.4923... Floating-point agreement is not an interval enclosure. There is no certified digit count in this report, and the analytic theorem does not depend on these calculations.

[write] **Checks, 6 October 2026.** The shipped diagnostic `data/225-leading-data-amplitude_diagnostic.json` records $h(1) = 1.8146095596784502$ and $c = 4.492329896751906$; the write recomputed $T^{1/3}e^{-B}h(1) = 4.4923298968$ from that $h(1)$. Part II’s derivative diagnostics give $h(1) \approx 1.81460956$ as well (Subsection 22.2). The coefficients above agree with Part III’s $a_1 = -1/36$, $a_2 = 1/540$, $a_3 = 1/7776$, and the write’s rerun of `coordinate_series.py--order9` also gives $a_4 = -71/435456$. All values remain uncertified.

10.3 Reproducibility and limits

The accompanying archive contains the complete editable TeX, exact programs and tests, numerical diagnostic script, exact fixture, source ledger, and a deterministic builder. The builder verifies an input manifest, runs explicit tests, creates a private pdfL^AT_EX format, compiles three times without shell escape, rejects layout and reference warnings, and packages sorted members with fixed timestamps. It preserves the source tree and writes outputs only to a new separate directory. Ordinary and optimized cap-640 runs are compared by full ZIP bytes and by each member. Byte identity is a claim about the same toolchain; arbitrary TeX, font, or numerical-library versions need not reproduce identical bytes.

[write] **The companion here, 6 October 2026.** Report 225’s companion is shipped as `code/225-leading-*.py`, `data/225-leading-*` and `225-leading-sources-SOURCE_LEDGER.md`; its delivered README was replaced by the report README, and its PDF and `MANIFEST.sha256` are not shipped. The builder and tests run only in the delivered layout; the report README gives the rerun route from the arrival commit.

11 Further questions and quantitative extensions

11.1 Validated evaluation of the positive constant

The exact expression (6) is a useful starting point for a certificate, but a validated computation requires more than increased precision. One must bound the omitted Fatou-coordinate terms on a stated sector, enclose the inverse branch, control the forward propagation of both value and derivative, validate the finite-interval quadrature, and give an explicit integrable bound for the remaining vertical tail. The sector proof above supplies qualitative constants; it does not extract numerical values for all of them. A practical goal is a reproducible interval for c narrow enough to validate the conjectured initial digits, with each of these error sources accounted for separately.

The rational subtraction used in the diagnostic script could be extended to several terms derived from the large- s inverse expansion. This can reduce cancellation and the cutoff length, but every subtracted term and its Fourier contribution must be explicit. Mesh or precision stability cannot replace bounds for truncation of the asymptotic coordinate and the infinite integral.

[write] **Still open, 6 October 2026.** Not settled by Parts II or III; Section 37, item 1. Part II’s closed form of κ removes one constant from the first correction, which still needs $h(1)$, $h'(1)$, $h''(1)$.

11.2 Higher corrections to the diagonal

The logarithmic center D_n is determined only to the precision needed for a leading equivalent. Higher corrections require retaining more subsidiary terms in $R_h(z)$, including their dependence on h , and constructing a corresponding hierarchy of nonautonomous clocks. Their matching constants interact with the inverse-coordinate expansion. It is natural to investigate corrections involving powers of $1/n$ and polynomials in $\log n$, but no complete expansion of this form is asserted here.

A formal expansion at bounded s is insufficient for a coefficient theorem. The error must be controlled in an integrable norm on the growing contour, with the intermediate-angle and outer-angle arguments strengthened to the same order. The present choice $K = \lfloor \sqrt{n} \rfloor$ balances errors only for convergence; a higher-order theorem would need a new quantified matching analysis. This identifies a concrete obstruction to promoting formal series directly into an all-orders claim.

[write] **Partly answered by Part II, 6 October 2026.** Part II proves the first correction, $(-\frac{1}{18} \log^2 n + A \log n + C)/n$ with remainder $O(n^{-7/5})$ (Theorem 12.1), by two further clock terms and a quantified growing-strip matching with $K = \lfloor n^{1/2} \rfloor$; it confirms the expected shape, a polynomial in $\log n$ of degree two over n . Corrections beyond the first and an all-orders theorem remain open (Section 37, item 2).

11.3 Effective bounds and inverse thresholds

Making the constants in the clock, stopping, and sector estimates explicit would give finite- n bounds and a computable onset for the leading equivalent. Such bounds would in turn permit certified recovery of integer thresholds for arbitrary input sizes. Corollary 9.1 currently supplies eventual rounding at exact sequence values; it does not say how large “eventually” is, nor does it control the location of a general input between adjacent terms. A rigorous threshold algorithm should combine exact small-index computation with explicit upper and lower bounds, rather than treat a floating-point evaluation of c as exact.

One can also seek inverse statements for $A(n, \lfloor \lambda n \rfloor)$. Formula (32) shows why the bounded floor phase must enter such statements. A single smooth approximation that discards this phase can

have a nonvanishing multiplicative error.

[write] **Still open, 6 October 2026.** Part II's Corollary 21.1 sharpens the inverse at exact sequence values, still without onset; Part III's inverses concern ratios in the window $m \asymp n \log n$, not $A(n, \lfloor \lambda n \rfloor)$. Section 37, items 3 and 4.

11.4 Other height scales and the density

Theorem 8.1 treats n/m in a fixed compact subset of $(0, \infty)$. Extending it toward zero or infinity requires information about $h(t)$ at the corresponding endpoint and renewed uniformity estimates. In particular, the strict-positivity proof does not give an expansion of $h(t)$ at zero. The measure identity

$$(e^t - 1)\nu = \sum_{k \geq 2} \nu^{*k}/k!$$

is a possible starting point for endpoint analysis and stronger regularity, such as real analyticity. Smoothness away zero is already established in Corollary 6.2, but differentiation of this convolution identity or exchange of limits with its series needs separate justification.

Kaneiwa's fixed-leaf polynomial regime, proportional height, and fixed-height partition regimes are genuinely different limiting questions. A uniform transition theory between them would explain how the polynomial and parabolic descriptions fit together. It may also clarify which parts of this proof extend to other multiset recurrences or other initial generating functions. Such a generalization must recompute the subsidiary forcing and the clock constant; the unforced map alone does not determine the answer.

[write] **Partly answered by Part III, 6 October 2026.** At the endpoint zero Part III proves the expansion asked for here: $h(0) = 2$, the Hankel representation and the power-log series (Proposition 27.1, Theorem 28.1, Corollary 29.2), and it extends the height theorem toward $n/m \rightarrow 0$ on logarithmic bands (Theorem 31.1), with Kaneiwa's normalization compared in Section 33. Theorem 28.1 also gives real analyticity of h on some interval $(0, \epsilon)$. Open: $n/m \rightarrow \infty$, real analyticity on all of $(0, \infty)$, the transition to the fixed-degree regime, other recurrences (Section 37, items 5–8).

Part II

First logarithmic correction for iterated Euler diagonals

Bundle Report 226, archive `Report226.zip`, delivered as `src/report226.tex` (17 pages). Delivered title block “First logarithmic correction for iterated Euler diagonals / Report 226 / 5 October 2026”. Printed as delivered, with `[write]` notes. Delivered Section k is Section $k + 11$ here, its statement $k.j$ is $(k + 11).j$, and its equation (k) is (II. k).

Reading conventions of Part II	Elsewhere in the report
“Report 225”, [6]	Part I
$b_{n,m}$ the array; d_* , d , d_m	Parts I, III: $A(n, m)$; Part I: D_n , $d = n - h$
$\rho_1, \rho_2 = h'(1)/h(1), h''(1)/h(1)$; $\rho_j(\beta)$	Part III: ρ_j rational
A, C the correction constants; G Catalan’s constant	Parts I, III: $A(n, m)$; Part III: $\mathcal{C}$ Catalan, C_0 , $C(\beta)$
$M(y)$, $N(y)$ clock functions; $M = n^{1/20}$; $\mathcal{C}_z(w)$	Part III: $M = m^{1/20}$, N an order
$q = 30$ the derivative order; $\alpha = 4/3$, $\ell = \log c$	Part I: q entrance constant; Part III: ℓ_n Kaneiwa

Abstract of this Part (as delivered). For the iterated Euler diagonal a_n in OEIS A290354, we prove

$$a_n = c(2n/\pi)^n n^{-4/3} \left\{ 1 + \frac{-\log^2 n/18 + A \log n + C}{n} + O(n^{-7/5}) \right\}.$$

The constants A and C are exact expressions in derivatives of the positive density associated with the normalized repelling parametrization of $w \mapsto e^w - 1$. A new nonautonomous clock constant is evaluated in terms of π , $\log 2$, and Catalan’s constant. The proof constructs two further clock terms, obtains a weighted fourth-power defect, and matches a forced Fatou correction on a growing vertical strip. Thirtieth-derivative coefficient extraction turns these estimates into the stated relative remainder. A uniform correction for compact positive degree-to-height ratios and a sharpened Lambert W inverse are included. This companion uses the analytic foundation of the preceding unrefereed Report 225, whose unchanged PDF and TeX accompany the archive. It is an unrefereed research proof, not a proof-assistant formalization. Floating-point diagnostics are not certified decimal enclosures; no all-orders theorem is claimed.

12 Definition and main results

Define

$$F_0(z) = z, \quad F_{h+1}(z) = \exp \left(\sum_{j \geq 1} \frac{F_h(z^j)}{j} \right) - 1. \quad (\text{II.1})$$

For $n \geq 1$ and $m \geq 0$, put $b_{n,m} = [z^n]F_m(z)$ and $a_n = b_{n,n}$; set $a_0 = 1$ and $b_{0,m} = 1$ separately when reporting the integer arrays. Thus a_n is A290354 and $b_{n,m}$ is A290353 [2, 1]. The analytic generating functions themselves have constant term zero. They count unlabeled rooted trees whose leaves all have the stated height: a new root carries a nonempty unordered multiset of trees at the previous height. Equivalently,

$$1 + F_{h+1}(z) = \prod_{d \geq 1} (1 - z^d)^{-[z^d]F_h(z)}.$$

There is no factorial normalization. The diagonal begins

$$1, 1, 2, 6, 30, 170, 1337, 12166, 133476, 1676364.$$

The exact constants and functions used throughout are

$$T = \frac{\pi}{2}, \quad B = \frac{1}{2} - \frac{\pi}{6} - \frac{\pi^2}{16} - \frac{\log 2}{6}, \quad d_\star = \frac{1}{3} \log T - B, \quad (\text{II.2})$$

$$\kappa = \frac{\pi^3}{96} + \frac{\pi^2}{12} - \frac{13\pi}{48} - \frac{11}{36} - \frac{G}{6} - \frac{\pi \log 2}{24}, \quad (\text{II.3})$$

where $G = \sum_{j \geq 0} (-1)^j / (2j+1)^2$ is Catalan's constant. Let P denote the entire repelling parametrization normalized by

$$P(s+1) = e^{P(s)} - 1, \quad \Phi(w) = -\frac{2}{w} + \frac{1}{3} \log w + O(w), \quad \Phi(P(s)) = s \quad (\text{II.4})$$

in the inverse sector near negative real s . Report 225 [6] constructs this normalization and proves

$$P(s) = \int_0^\infty e^{sx} h(x) dx, \quad h \in C^\infty(0, \infty), \quad h(x) > 0 \quad (x > 0). \quad (\text{II.5})$$

The density has every exponential moment. Set

$$c = T^{1/3} e^{-B} h(1), \quad \rho_1 = \frac{h'(1)}{h(1)}, \quad \rho_2 = \frac{h''(1)}{h(1)}. \quad (\text{II.6})$$

The zero additive constant in (II.4) is essential: a translation of the Fatou coordinate would change P and h .

Theorem 12.1. *As $n \rightarrow \infty$,*

$$a_n = c(n/T)^n n^{-4/3} \left(1 + \frac{-\log^2 n / 18 + A \log n + C}{n} + O(n^{-7/5}) \right), \quad (\text{II.7})$$

where $c > 0$ and

$$A = \frac{d_\star}{3} + \frac{4}{9} + \frac{\rho_1}{3}, \quad (\text{II.8})$$

$$C = -\frac{d_\star^2}{2} - d_\star \left(\frac{4}{3} + \rho_1 \right) - T\kappa - \frac{1}{3} - \frac{4\rho_1}{3} - \frac{\rho_2}{2}.$$

The error is relative to the displayed leading expression. Its exponent is a convenient nonoptimal choice.

With $d = -\frac{1}{3} \log n + d_\star$, the numerator of the correction can equivalently be written

$$-\frac{d^2}{2} - d \left(\frac{4}{3} + \rho_1 \right) - T\kappa - \frac{1}{3} - \frac{4\rho_1}{3} - \frac{\rho_2}{2}. \quad (\text{II.9})$$

In particular the squared-logarithm coefficient is exactly $-1/18$. Retaining only a more accurate contour radius would miss some terms in A and C .

Theorem 12.2. *Let $I = [\beta_0, \beta_1] \subset (0, \infty)$ be fixed. Uniformly for integers $m \rightarrow \infty$ and $n \geq 1$ with $\beta = n/m \in I$, put $d_m = -\frac{1}{3} \log m + d_\star$ and $\rho_j(\beta) = h^{(j)}(\beta)/h(\beta)$ for $j = 1, 2$. Then*

$$b_{n,m} = h(\beta) T^{\beta/3} e^{-\beta B} (m/T)^n m^{-1-\beta/3} \left(1 + \frac{\mathcal{Q}_\beta(d_m)}{m} + O_I(m^{-7/5}) \right), \quad (\text{II.10})$$

$$\begin{aligned} \mathcal{Q}_\beta(d) = & -\frac{\beta d^2}{2} - d \left(1 + \frac{\beta}{3} + \beta \rho_1(\beta) \right) - \beta \kappa T \\ & - \frac{1}{3} - \left(1 + \frac{\beta}{3} \right) \rho_1(\beta) - \frac{\beta}{2} \rho_2(\beta). \end{aligned}$$

The exact height and exact ratio occur in the formula; no additional factor of β belongs in its leading amplitude.

No uniform conclusion is asserted as β approaches zero or infinity. In particular replacing an integer height by an unfloored real height without retaining its phase is not justified.

[write] **Toward $\beta \rightarrow 0$ in Part III, 6 October 2026.** Part III extends the *leading* term of (II.10) to the bands $(\log m)^{-D} \leq \beta \leq B_0$ (Theorem 31.1); no Part extends the correction $\mathcal{Q}_\beta$ there, and $\beta \rightarrow \infty$ is open (Section 37, item 5).

12.1 Dependence on the preceding report and prior work

Report 225 proves the leading equivalent and its analytic foundation. The present report extends its clock and coefficient analysis; it does not independently reprove all entrance and angular-tail estimates. Section 13 records the full interface used, and the accompanying archive includes the unchanged complete proof of Report 225. Neither report has undergone journal refereeing.

The OEIS attributes the leading equivalent with numerical amplitude 4.4923... to Václav Kotěšovec, dated 14 August 2017 [2]. The fixed-degree polynomiality and tangent leading profile have earlier antecedents explained and sourced in Report 225. The correction proved here is not inferred merely from that fixed-degree information. We make no worldwide priority claim for the result or for parabolic-coordinate methods.

[write] **Reading Part II against Part I, 6 October 2026.** “Report 225” and [6] are Part I of this report; its Sections 2–7 are Part I’s Sections 2–7, with the same numbers. The included copy is byte-identical to Part I and is not shipped. At leading order Theorem 12.1 is Part I’s Theorem 1.1 and Theorem 12.2 is Part I’s Theorem 8.1. Here $b_{n,m}$ is Part I’s $A(n, m)$, and $d_\star = \frac{1}{3} \log T - B$, so D_n is Part I’s D_n . Part II’s ρ_j are not Part III’s rational ρ_j .

13 Analytic foundation from Report 225

All branches agree with real values on the positive axis. Constants in local analytic estimates may depend on fixed sectors or fixed real-width bounds, but not on the growing indices. The following are the precise results imported from Report 225, Sections 2–7 [6].

- (i) Each F_h is analytic in $|z| < 1$, with nonnegative coefficients and linear coefficient one. If $(h+1)|q|$ is sufficiently small, then $|F_h(q)| \leq C|q|$ uniformly. This follows by positive-coefficient comparison on the real radius and a reciprocal recurrence bound; it does not assume that the main value $F_h(z)$ is small.

(ii) Define

$$J(y) = 2 \arctan y - T, \quad L(1) = 0, \quad L'(y) = \frac{y^3/3 - y - 4/3 - 2J(y)}{(1+y^2)^2},$$

$$H(z) = \frac{T}{z} + \frac{1}{3} \log \frac{1}{z} + B. \quad (\text{II.11})$$

The old clock $J(w/z)/z + L(w/z)$ gives a noncircular positive-real entrance bound and its Cauchy extension. More precisely, let $D_n = n - \frac{1}{3} \log n + d_*$ and $r = T/D_n$. For suitable fixed $q > 0$, K_0 , and $\epsilon > 0$, if $K_0 \leq K \leq \epsilon n$, then on $|z - r| \leq qK/n^2$ the orbit through $n - K$ is small and

$$|F_h(z)| \asymp (n - h)^{-1} \quad (0 \leq h \leq n - K), \quad F_h(z)/z \text{ lies in a fixed narrow positive cone.} \quad (\text{II.12})$$

The cone has a slightly larger fixed cone available for Taylor segments. At entrance the old phase estimate is

$$\Phi(F_{n-K}(z)) = n - K - H(z) + O(K^{-1} + n^{-1} + K^3/n^2). \quad (\text{II.13})$$

(iii) The sector coordinate Φ in (II.4) satisfies the Abel equation $\Phi(f(w)) = \Phi(w) + 1$, $f(w) = e^w - 1$. Its inverse is uniquely selected in fixed interior sectors about the closed left half-plane for s of large modulus, and

$$P(s) = -\frac{2}{s} + O\left(\frac{\log |s|}{|s|^2}\right). \quad (\text{II.14})$$

A small point w in a smaller coordinate sector satisfying $\Phi(w) = s$ is the same inverse value $P(s)$; global injectivity is unnecessary. The inverse iterates $g^{\circ j}(w)$, $g(w) = \log(1 + w)$, obey $|g^{\circ j}(w)| \leq C/(|w|^{-1} + j)$ in nested sectors. Extension by finitely many iterates of f gives the entire P .

(iv) There are fixed $b, \delta, M_0, C > 0$ such that for $M_0 \leq |t| \leq \delta n$,

$$|z - re^{it/n}| \leq b|t|/n^2 \implies |F_n(z)| \leq C/|t|. \quad (\text{II.15})$$

For the outer angles $\delta \leq |\theta| \leq \pi$, the compact arc $\zeta \approx Te^{i\theta}$ has fixed complex neighborhoods on which $F_n(\zeta/n) = O(n^{-1})$. These neighborhoods follow from the bounded ODE profile $\zeta \tan(\zeta u/2 + \pi/4)$ for $0 \leq u \leq 1$ away from its positive pole. Both are bounds on two-dimensional analytic neighborhoods, not merely on one-dimensional arcs.

(v) The positive-measure construction yields (II.5), with no atom at zero, support $[0, \infty)$, and strict positivity of h on $(0, \infty)$. Its leading coefficient consequence is $a_n \sim c(n/T)^n n^{-4/3}$.

[write] Where Part I proves each item, 6 October 2026. (i) Lemma 2.1 and the first step of its proof. (ii) (9)–(10), Lemmas 2.2–3.3 and (23) (Part I writes $K \leq \kappa n$ for the $K \leq \epsilon n$ here; this κ is not the clock constant). (iii) Lemma 4.1, (20)–(21); the bound on inverse iterates is the estimate $|G^j(u)| \geq c(|u| + j)$ of its proof, read in $w = 2/u$. (iv) Proposition 5.2, whose proof (not its statement) bounds F_n on the whole disk about z_0 , and Lemma 5.3, whose proof (not its statement, which concerns the moving circle) gives the bound on complex neighbourhoods of the arc. (Extended after the independent check of 6 October 2026; the first wording read “Proposition 5.2, whose proof bounds F_n on the whole disk about z_0 , and Lemma 5.3”.) (v) Proposition 6.1 and Theorem 1.1. The paragraph below repeats Part I’s Corollary 6.2 with its proof. The section is printed in full because its displays are cited below.

For clarity, the derivative regularity needed below also follows directly from this interface. Sector Cauchy estimates in (II.14) give, for each fixed $q \geq 1$,

$$P^{(q)}(\sigma + it) = O_q(|t|^{-q-1}) \quad (\text{II.16})$$

uniformly for σ in a bounded interval. This is the Fourier transform of $x^q e^{\sigma x} h(x)$. If $q > k$, its product with t^k is integrable, so Fourier differentiation gives a C^k weighted density. Dividing by the nonzero smooth weight on compact positive intervals proves $h \in C^\infty(0, \infty)$. No boundary behavior of h or its derivatives at zero is used.

14 Subsidiary forcing through fifth scaled order

Write

$$R_h(z) = \sum_{j \geq 2} F_h(z^j)/j, \quad w = F_h(z), \quad y = w/z, \quad t = hz.$$

The map is $w_{h+1} = f(w_h + R_h(z))$.

Lemma 14.1. *Uniformly for $h \leq C_0/|z|$, as $z \rightarrow 0$,*

$$\begin{aligned} R_h(z) = & \frac{z^2}{2} + z^3 \left(\frac{1}{3} + \frac{t}{2} \right) + z^4 \left(\frac{1}{4} + \frac{t^2}{4} \right) \\ & + z^5 \left(\frac{1}{5} + \frac{7t}{12} + \frac{t^3}{6} \right) + O(|z|^6). \end{aligned} \quad (\text{II.17})$$

Proof. Coefficient extraction in one Euler step gives

$$[q^2]F_h = h, \quad [q^3]F_h = \frac{h(h+1)}{2}, \quad [q^4]F_h = \frac{h(h+1)(2h+1)}{6}.$$

Indeed, if the current coefficients are a, b, c , their increments are respectively 1, $a+1$, and $b+a^2/2+3a/2+1$; the latter becomes $(h+1)^2$. All three start at zero.

Take $\varrho = c_0/(h+1)$ with c_0 small. The small-subsidiary bound gives $\sup_{|q| \leq \varrho} |F_h(q)| \leq C\varrho$. Cauchy and a geometric sum therefore give, for $|q| \leq \varrho/2$,

$$F_h(q) = q + hq^2 + \frac{h(h+1)}{2}q^3 + \frac{h(h+1)(2h+1)}{6}q^4 + O((h+1)^4|q|^5). \quad (\text{II.18})$$

For small z , every $q = z^j$, $j \geq 2$, lies in this disk. The $j = 2$ remainder is $O((h+1)^4|z|^{10}) = O(|z|^6)$, and $j \geq 6$ contributes $O(|z|^6)$. At order z^5 , the baseline is $1/5$; the linear- h part of $[q^3]F_h$ at $j = 2$ contributes $t/4$, $[q^2]F_h$ at $j = 3$ contributes $t/3$, and the cubic- h part of $[q^4]F_h$ at $j = 2$ contributes $t^3/6$. This proves (II.17), including the otherwise easily lost $7t/12$ term. $\square$

[write] **Extends Part I, 6 October 2026.** This extends Part I's (8), whose $hz^4/2$ is the term $z^3t/2$ here. The three coefficients $[q^2]F_h$, $[q^3]F_h$, $[q^4]F_h$ are the rows $n = 2, 3, 4$ of A290353 (its cross-references name A001477, A000217, A000330); the write checked them against the exact recurrence for $h \leq 12$.

For the clock algebra write $d_y = 1 + y^2$ and $a(y) = d_y/2$. Expansion of the exponential gives

$$\Delta y = za + z^2b(y, t) + z^3v(y, t) + z^4u(y, t) + O(|z|^5(1 + |y|)^6), \quad (\text{II.19})$$

where

$$\begin{aligned} b(y, t) &= y^3/6 + y/2 + 1/3 + t/2, \\ v(y, t) &= y^4/24 + y^2/4 + y/3 + yt/2 + 3/8 + t^2/4, \\ u(y, t) &= y^5/120 + y^3/12 + y^2/6 + y^2t/4 + 3y/8 + yt^2/4 \\ &\quad + 11/30 + 5t/6 + t^3/6. \end{aligned} \quad (\text{II.20})$$

The estimate is uniform for bounded t when $|w| + |z|$ is small. These are finite Taylor identities with analytic remainders, not extrapolations from numerical values.

15 Two further nonautonomous clock terms

Define $M(1) = N(1) = 0$ on the same simply connected cone as J, L . First set

$$Q_2(y, t) = J'v + J''ab + \frac{J'''}{6}a^3 + L'b + \frac{L''}{2}a^2. \quad (\text{II.21})$$

The required equation for M is

$$aM' = -Q_2(y, J) - \frac{L}{d_y}. \quad (\text{II.22})$$

Its expanded form is

$$M' = -\frac{1}{36d_y^3} \{ 36J^2(y^2 - 1) + 12J(5y^3 - 3y - 8) + 72Ld_y + y^6 + 9y^4 + 40y^3 - 21y^2 - 24y - 29 \}. \quad (\text{II.23})$$

The feedback term L/d_y is necessary. Setting $hz = J$ before retaining the self-consistency terms would give a wrong next clock.

Next define

$$Q_3(y, t) = J'u + J''(av + b^2/2) + \frac{J'''}{2}a^2b + \frac{J'''}{24}a^4 + L'v + L''ab + \frac{L'''}{6}a^3 + M'b + \frac{M''}{2}a^2, \quad (\text{II.24})$$

$$G_3(y) = Q_3(y, J) + L \partial_t Q_2(y, J) + M/d_y, \quad N' = -G_3/a. \quad (\text{II.25})$$

Here $\partial_t Q_2(y, J) = J/d_y + L'/2$. Equations (II.22) and (II.25), with the zero values at one, are exact recursive definitions involving only differentiation, integration, and rational operations. They specify the functions without a fitted constant.

15.1 Large ratio expansions

Uniformly on each closed smaller cone,

$$J = T - 2/y + 2/(3y^3) + O(y^{-5}),$$

$$L = \frac{1}{3} \log y + B + \frac{5}{6y^2} + \frac{4/9 + 2T/3}{y^3} - \frac{7}{4y^4} + O(y^{-5}). \quad (\text{II.26})$$

Substitution in (II.23) gives

$$M' = -\frac{1}{36} - \frac{1}{6y^2} - \frac{5T/3 + 10/9}{y^3} + \frac{-(2/3) \log y - 2B - T^2 + 9/2}{y^4} + O(y^{-5}), \quad (\text{II.27})$$

and integration yields

$$M = -\frac{y}{36} + \kappa + \frac{1}{6y} + \frac{5T/6 + 5/9}{y^2} + \frac{(2/9)\log y + 2B/3 + T^2/3 - 77/54}{y^3} + O(y^{-4}). \quad (\text{II.28})$$

Thus M has neither a divergent $\log y$ nor a divergent $\log^2 y$ term. The constant is the convergent integral

$$\kappa = \frac{1}{36} + \int_1^\infty \left(M'(y) + \frac{1}{36} \right) dy. \quad (\text{II.29})$$

For the next clock, direct expansion in (II.25) gives

$$\begin{aligned} G_3 &= -\frac{y^3}{540} - \frac{y}{360} + \frac{T}{12} + \frac{1}{18} \\ &\quad + \frac{(5/18)\log y + 5B/6 + 5T^2/12 - 3/20}{y} + O(\log y/y^2), \\ N' &= \frac{y}{270} + \frac{1}{540y} - \frac{T/6 + 1/9}{y^2} \\ &\quad + \frac{-(5/9)\log y - 5B/3 - 5T^2/6 + 161/540}{y^3} + O(\log y/y^4), \\ N &= \frac{y^2}{540} + \frac{\log y}{540} + n_0 + \frac{T/6 + 1/9}{y} \\ &\quad + \frac{(5/18)\log y + 5B/6 + 5T^2/12 - 11/1080}{y^2} + O(\log y/y^3). \end{aligned} \quad (\text{II.30})$$

The finite n_0 is fixed by $N(1) = 0$ and is not needed at the present order. Unlike M , N contains a divergent logarithm. Its clock contribution is $z^2 \log y$, below the first correction but relevant to future orders.

15.2 Exact evaluation of the new clock constant

Put $u = \arctan y$. An exact primitive, checked by differentiation and its value at $u = T/2$, is

$$\begin{aligned} L(\tan u) &= -\frac{1}{3} \log \cos u + \frac{1}{3} - \frac{1}{6} \cos 2u + (T - 2/3)u - u^2 \\ &\quad + (T/2 - 1/3 - u) \sin 2u - \frac{1}{6} \log 2 - \frac{1}{4} T^2 + T/3. \end{aligned} \quad (\text{II.31})$$

Substituting it and $J = 2u - T$ in (II.23) gives

$$(M' + 1/36) dy = \left(\frac{2}{3} \cos^2 u \log \cos u + \mathcal{P}(u) \right) du,$$

where the finite trigonometric polynomial is

$$\mathcal{P} = p_0 + p_2 \cos 2u + p_4 \cos 4u + q_2 \sin 2u + q_4 \sin 4u$$

and

$$\begin{aligned} p_0 &= 2u^2 - 2Tu + T^2/2 + 8u/3 - 4T/3 + 1/12 + \log 2/6, \\ p_2 &= 3u^2 - 3Tu + 3T^2/4 + 10u/3 - 5T/3 + 1/3 + \log 2/6, \\ p_4 &= u^2 - Tu + T^2/4 + 2u/3 - T/3 + 1/12, \\ q_2 &= 2u/3 - T/3 + 2/9, \quad q_4 = 7u/6 - 7T/12 + 7/18. \end{aligned}$$

Elementary termwise integration yields

$$\int_{T/2}^T \mathcal{P}(u) du = -\frac{5\pi}{16} - \frac{5}{12} - \frac{\log 2}{12} + \frac{\pi \log 2}{24} + \frac{\pi^3}{96} + \frac{\pi^2}{12}.$$

The two logarithmic integrals are

$$\begin{aligned} \int_{\pi/4}^{\pi/2} \log \cos u du &= -\frac{\pi}{4} \log 2 - \frac{G}{2}, \\ \int_{\pi/4}^{\pi/2} \cos 2u \log \cos u du &= \frac{\log 2}{4} + \frac{\pi}{8} + \frac{1}{4}. \end{aligned}$$

The first follows by combining the half-period log-cosine integral with $G = -\int_0^{\pi/4} \log \tan u du$; the latter identity also follows from the Fourier series of the logarithm. For the second, integration by parts leaves the boundary contribution $\log 2/4$ and $\int_{\pi/4}^{\pi/2} \sin^2 u du = \pi/8 + 1/4$. The other endpoint term tends to zero. Using $\cos^2 u = (1 + \cos 2u)/2$ in (II.29), including its $1/36$ normalization, proves exactly (II.3).

[write] **Checks, 6 October 2026.** At intake the closed form was compared with a direct quadrature of (II.29), using the primitive (II.31) (itself checked against quadrature of (9)): -0.2543481698511 against -0.2543481698560 , the difference coming from the truncated tail. The delivered symbolic suite (`code/226-firstcorr-code-clock_checks.py`) reproduced `data/226-firstcorr-data-symbolic_checks.json` on a copy at the write.

16 Weighted defect and accumulated clock error

Define

$$\mathcal{C}_z(w) = J(w/z)/z + L(w/z) + zM(w/z) + z^2N(w/z), \quad E_h = \mathcal{C}_z(F_h(z)) - h. \quad (\text{II.32})$$

All four functions vanish at ratio one, so $E_0 = 0$.

Lemma 16.1. *Suppose $h \leq C_0/|z|$, the orbit is in the small-clock range, and $y_h = F_h(z)/z$ remains in the fixed cone. Then*

$$|E_{h+1} - E_h| \leq \frac{C|z|^2|E_h|}{(1 + |y_h|)^2} + C(|F_h(z)|^4 + |z|^4). \quad (\text{II.33})$$

Consequently, on the entrance disks of (II.12),

$$\mathcal{C}_z(F_{n-K}(z)) = n - K + O(K^{-3} + n^{-3}). \quad (\text{II.34})$$

Proof. Put $Y = 1 + |y|$ on a slightly wider cone. There $|d_y| \asymp Y^2$ and $|w| + |z| \asymp |z|Y$. The definitions and sector expansions give, for the fixed derivative orders required below,

$$\begin{aligned} J^{(r)} &= O(Y^{-r-1}), & L^{(r)} &= O(Y^{-r}), \\ M^{(r)} &= O(Y^{1-r}), & N^{(r)} &= O(Y^{2-r}) \quad (r \geq 1), \end{aligned}$$

and $J = O(1)$, $L = O(1 + \log Y)$, $M = O(Y)$, $N = O(Y^2)$. Since $\Delta y = O(|z|Y^2)$, its relative size is $O(|w| + |z|)$; all Taylor segments stay in the larger cone for a sufficiently small clock radius.

Temporarily replace R_h by its polynomial through degree five, keeping a bounded independent t . The discarded $O(z^6)$ forcing changes the clock by $O(z^6/(|w| + |z|)^2) = O(z^4)$. Expand J/z to fourth order in the displacement, L to third, zM to second, and z^2N to first. Their omitted terms are bounded respectively by

$$|z|^{-1}Y^{-6}(|z|Y^2)^5, \quad Y^{-4}(|z|Y^2)^4, \quad |z|Y^{-2}(|z|Y^2)^3, \quad |z|^2(|z|Y^2)^2.$$

Each is $O(|z|^4Y^4)$. The step remainder $O(|z|^5Y^6)$ in (II.19), multiplied by the leading ratio derivative $O(|z|^{-1}Y^{-2})$, has the same bound.

The retained clock increment is exactly

$$1 + z \frac{t - J}{d_y} + z^2\{Q_2(y, t) + aM'\} + z^3\{Q_3(y, t) + aN'\} + O(|z|^4Y^4).$$

Set $t_0 = z\mathcal{C}_z(w) = J + zL + z^2M + z^3N$. It is bounded, and $t_0 - J = O(|z|(1 + \log Y))$ because $|z|Y$ is small. Expansion in t about J gives the terms

$$z^2\{L/d_y + Q_2(y, J) + aM'\} + z^3\{M/d_y + L\partial_t Q_2(y, J) + Q_3(y, J) + aN'\}.$$

They vanish by (II.22) and (II.25). All discarded t -Taylor terms are $O(|z|^4Y^4)$: representative largest bounds are $|z|^4Y^3$ and $|z|^4(1 + \log Y)^2Y^2$, both admissible.

The actual variable satisfies $t = hz = t_0 - zE_h$ exactly. Both t and t_0 are bounded before any bound on E_h is used, so their connecting segment is bounded. On it $\partial_t R_{\text{polynomial}} = O(z^3)$ and $\partial_w \mathcal{C}_z = O((|w| + |z|)^{-2})$. The derivative of the increment with respect to t is therefore $O(|z|/Y^2)$. Multiplication by $-zE_h$ gives the feedback term in (II.33); $|z|^4Y^4 \asymp |w|^4 + |z|^4$ gives the remainder.

Discrete Gronwall has multiplier at most $\exp(Cn|z|^2) = O(1)$. The already established entrance orbit gives

$$\sum_{h < n-K} |F_h(z)|^4 = O(K^{-3}), \quad n|z|^4 = O(n^{-3}).$$

Together with $E_0 = 0$ this proves (II.34). The orbit bounds come from the old clock, so the improved defect is not being used to bootstrap its own hypotheses. $\square$

17 Finite Fatou expansion and forced matching

Lemma 17.1. *In fixed smaller coordinate sectors,*

$$\Phi(w) = -\frac{2}{w} + \frac{1}{3} \log w - \frac{w}{36} + \frac{w^2}{540} + O(w^3). \quad (\text{II.35})$$

For

$$U_0(w) = \frac{2}{3w^3} + \frac{5}{6w^2} + \frac{1}{6w}, \quad (\text{II.36})$$

one has $U_0(f(w)) - U_0(w) = -\Phi'(w)/2 + O(w)$.

Proof. Let Φ_2 be the four displayed terms in (II.35). Direct series division gives

$$\Phi_2(f(w)) - \Phi_2(w) - 1 = -w^4/5184 + O(w^5).$$

Sum this defect along the inverse orbit from Section 13. Its absolute total is bounded by $C \sum_{j \geq 0} (|w|^{-1} + j)^{-4} = O(w^3)$. This constructs a normalized Abel coordinate with (II.35). Its

difference from the original normalized coordinate is inverse-map invariant and tends to zero along inverse iterates, hence vanishes. Sector Cauchy estimates justify differentiated remainders. A second finite calculation gives

$$U_0(f(w)) - U_0(w) + \Phi'_2(w)/2 = -w/1080 + O(w^2).$$

Replacing Φ'_2 by Φ' changes this by $O(w^2)$. Both lower pole terms in U_0 are needed, including for cancellation of the constant term. $\square$

[write] **Generalized in Part III, 6 October 2026.** Part III's Lemma 26.1 proves the expansion to every order, with rational a_j and differentiated remainders, by the same summation of the Abel defect along inverse orbits; its $a_1 = -1/36$, $a_2 = 1/540$ are the coefficients here.

Combining the expansions of J, L, M, N gives, when $|w/z|$ is large in the clock cone,

$$\begin{aligned} \mathcal{C}_z(w) &= H(z) + \kappa z + \Phi(w) + z^2 U_0(w) \\ &\quad + O\left(|w|^3 + |z|^2(1 + \log |w/z|) + \frac{|z|^3}{|w|^3} + \frac{|z|^4}{|w|^5}\right). \end{aligned} \quad (\text{II.37})$$

For example, the z^2/w^3 , z^2/w^2 , and z^2/w terms come respectively from J , L , and zM and assemble exactly into $z^2 U_0$. Lower M, N tails are absorbed by the displayed remainder in the small-clock range.

Take

$$K = \lfloor n^{1/2} \rfloor, \quad z \asymp n^{-1}, \quad w = F_{n-K}(z) \asymp K^{-1}. \quad (\text{II.38})$$

The four matching errors in (II.37) have orders at most K^{-3} , $n^{-2} \log n$, $n^{-3} K^3$, and $n^{-4} K^5$; each is $O(n^{-3/2})$. The clock error in (II.34) has the same order. Thus, for $\Theta_z = \Phi + z^2 U_0$,

$$\Theta_z(F_{n-K}(z)) = n - K - H(z) - \kappa z + O(n^{-3/2}). \quad (\text{II.39})$$

The sign of the new phase constant is negative because $+\kappa z$ is moved to the other side of the clock identity.

For an actual late step, use the Abel equation before Taylor expansion: $\Phi(w_{h+1}) = \Phi(w_h + R_h) + 1$. Since $R_h = z^2/2 + O(z^3)$, Lemma 17.1 yields

$$\Theta_z(w_{h+1}) - \Theta_z(w_h) - 1 = O\left(|z|^2 |w_h| + \frac{|z|^3}{|w_h|^2} + \frac{|z|^4}{|w_h|^4}\right). \quad (\text{II.40})$$

Indeed the $z^2 \Phi'/2$ term cancels the leading U_0 difference. The second-order Taylor error for Φ is $O(z^4/w_h^3)$, absorbed by the last term, and perturbing the argument in U_0 gives $O(z^4/w_h^4)$.

18 Uniform local limit on a growing strip

Proposition 18.1. *For every fixed real-width bound C_0 , uniformly at*

$$z = \frac{T}{D_n - s}, \quad |\operatorname{Re} s| \leq C_0, \quad |\operatorname{Im} s| \leq n^{1/20} + 2,$$

one has

$$F_n(z) = P(n - H(z) - \kappa z) + O(n^{-3/2}). \quad (\text{II.41})$$

Proof. Write $M_n = n^{1/20} + 2$ and use K in (II.38). The distance to $r = T/D_n$ is $O(M_n/n^2)$, so the whole parameter set lies in the old entrance disk of radius qK/n^2 . All entrance and new matching estimates are uniform there.

Let $\alpha = n - H(z) - \kappa z$. The exact identity

$$H\left(\frac{T}{D_n - s}\right) = n - s + \frac{1}{3} \log \frac{D_n - s}{n}$$

gives $\alpha = s + O((\log n + M_n)/n)$; its real part stays bounded and its imaginary part differs from $\text{Im } s$ by $o(1)$.

Choose one fixed integer L large relative to C_0 and the inverse-sector threshold. It does not depend on M_n . For $0 \leq j \leq K - L$, set $m = K - j$ and use the prototype phase $\alpha - m$. Its real part is at most $C_0 + 1 - m < 0$ and

$$|\alpha - m| \asymp m + |\text{Im } s|.$$

It lies in a fixed interior inverse sector about the closed left half-plane. A bootstrap phase error at most one preserves fixed angular and radial margins. The corresponding inverse values satisfy

$$|w_j| \asymp (m + |\text{Im } s|)^{-1}. \quad (\text{II.42})$$

The actual $O(n^{-2})$ forcing changes the uncorrected Φ phase by at most $Cn^{-2}(m + M_n)^2$. Its total is

$$O\left(n^{-2} \sum_{m=L}^K (m + M_n)^2\right) = O(K^3/n^2) = o(1),$$

because $M_n = o(K)$. The old entry error in (II.13) is also $o(1)$. These strictly improve the bootstrap for large n . The inverse-identification clause ensures every perturbed point uses the same branch. There is no new injectivity assumption.

The bounds just obtained imply the sums needed for the corrected phase:

$$\sum |w_j| = O(\log K), \quad \sum |w_j|^{-2} = O(K^3), \quad \sum |w_j|^{-4} = O(K^5).$$

Summing (II.40) to time $n - L$ gives

$$O(n^{-2} \log n + n^{-3} K^3 + n^{-4} K^5) = O(n^{-3/2}),$$

of the same order as the error in (II.39). Removing $z^2 U_0$ at time $n - L$ costs $O(n^{-2}(1 + M_n)^3) = O(n^{-37/20})$. Hence

$$\Phi(F_{n-L}(z)) = \alpha - L + O(n^{-3/2}).$$

The inverse derivative is uniformly bounded on the shifted strip, by analyticity on its bounded part and the sector estimate on its unbounded part. We obtain $F_{n-L}(z) = P(\alpha - L) + O(n^{-3/2})$.

The values $P(\alpha - L)$ form a bounded set and tend to zero at large imaginary height. Its closure is compact. The fixed entire composition $f^{\circ L}$ and its derivatives are bounded on a compact enlargement, so the remaining L iterates preserve the error; their subsidiary forcing contributes only $O(n^{-2})$. This proves (II.41). Large imaginary parts remain in the inverse sector, which is why one fixed L suffices. $\square$

18.1 Exponential parameter and explicit remainders

Set $d = -\frac{1}{3} \log n + d_*$, $D_n = n + d$, $r = T/D_n$, and

$$G_n(s) = F_n(re^{s/n}), \quad Q_n(s) = (d + 1/3)s - s^2/2 - d/3 - \kappa T.$$

The exact phase identity is

$$\begin{aligned} n - H(re^{s/n}) - \kappa re^{s/n} &= D_n(1 - e^{-s/n}) - \frac{1}{3} \log(1 + d/n) \\ &\quad + s/(3n) - \kappa T e^{s/n}/(n + d). \end{aligned} \quad (\text{II.43})$$

It implies

$$n - H(re^{s/n}) - \kappa re^{s/n} = s + Q_n(s)/n + R_n(s), \quad |R_n(s)| \leq Cn^{-2}(1 + |d| + |s|)^3. \quad (\text{II.44})$$

The rational parameter for $re^{s/n}$ is $D_n(1 - e^{-s/n})$; this preserves a slightly enlarged fixed-width strip and increases its height by $o(1)$. Both P' and P'' are uniformly bounded on such strips. Taylor expansion in (II.44) has additional error bounded by $Cn^{-2}(1 + |d| + |s|)^4$. On $|\operatorname{Im} s| \leq n^{1/20} + 2$ it is smaller than $n^{-3/2}$. Proposition 18.1 therefore gives

$$G_n(s) = P(s) + \frac{Q_n(s)P'(s)}{n} + O(n^{-3/2}). \quad (\text{II.45})$$

Cauchy's estimate on fixed-radius disks inside the enlarged strip gives the same error for every fixed derivative, in particular for derivative thirty. These degree-three and degree-four bounds avoid an unspecified polynomial whose degree could compromise the growing-strip argument.

19 Derivative extraction and classical Fourier inversion

For every nonnegative integer q , termwise coefficient extraction gives exactly

$$na_n r^n = \frac{1}{2\pi} \int_{-\pi n}^{\pi n} G_n^{(q)}(it) e^{-it} dt. \quad (\text{II.46})$$

Indeed a term of degree k acquires $(k/n)^q$, and orthogonality over this full period selects $k = n$, whose factor is one. Analyticity in a neighborhood of the circle justifies the termwise step. No integration by parts or endpoint assumption is needed.

19.1 Transport of analytic disks

The estimates imported from Report 225 are disk estimates, which permit arbitrary fixed derivative order. Choose a sufficiently small fixed $a > 0$. For $|t| \leq \delta n$,

$$|s - it| \leq a|t| \implies |re^{s/n} - re^{it/n}| \leq re^{a\delta} a|t|/n \leq b|t|/n^2.$$

Thus (II.15) and Cauchy give

$$|G_n^{(30)}(it)| \leq C_{30}|t|^{-31} \quad (M_0 \leq |t| \leq \delta n). \quad (\text{II.47})$$

For the outer angles, the fixed neighborhoods in $\zeta = nz$ admit s disks of radius $a_0 n$ for one fixed small $a_0 > 0$. The bound $G_n = O(n^{-1})$ there gives $G_n^{(30)} = O(n^{-31})$. Disks crossing the endpoints $\pm \pi n$ cause no problem because the exponential parametrization is periodic.

Put $M = n^{1/20}$. The finite-height intermediate and outer integrals outside $[-M, M]$ are

$$O(M^{-30} + n^{-30}) = O(n^{-3/2}).$$

The central integral of the differentiated error in (II.45) is

$$O(Mn^{-3/2}) = O(n^{-29/20}).$$

For the limiting transforms, (II.16) and Leibniz give

$$(Q_n P')^{(30)}(it) = O(|t|^{-30} + \log n |t|^{-31})$$

for large $|t|$. Its omitted tail, divided by n , is smaller still, and that of $P^{(30)}$ is $O(M^{-30})$. Both central integrals can consequently be replaced by full-line integrals with error $O(n^{-7/5})$. The margin between $29/20$ and $7/5$ is deliberate.

19.2 Signs and derivative weights in inversion

Write $\mathcal{I}[V](x) = (2\pi)^{-1} \int_{\mathbb{R}} V(it)e^{-itx} dt$ whenever the integral is absolutely convergent. We only apply it to the differentiated transforms with $q = 30$. At positive x , Fourier inversion and its justified ordinary derivatives give

$$\begin{aligned} \mathcal{I}[P^{(q)}](x) &= x^q h(x), \\ \mathcal{I}[(P')^{(q)}](x) &= x^{q+1} h(x), \\ \mathcal{I}[(sP')^{(q)}](x) &= -(x^{q+1} h)'(x) + qx^q h(x), \\ \mathcal{I}[(s^2 P')^{(q)}](x) &= (x^{q+1} h)''(x) - 2q(x^q h)'(x) + q(q-1)x^{q-1} h(x). \end{aligned} \tag{II.48}$$

For example, $(sP')^{(q)} = sP^{(q+1)} + qP^{(q)}$ and $(s^2 P')^{(q)} = s^2 P^{(q+1)} + 2qsP^{(q)} + q(q-1)P^{(q-1)}$. The t - and t^2 -weighted integrals are absolutely convergent by (II.16); multiplication by s corresponds to minus ordinary differentiation in x for this Laplace sign convention. These identities invoke no endpoint distributions and no regularity at zero.

At $x = 1$, the four inversions in (II.48) are respectively

$$h(1), \quad h(1), \quad -h(1) - h'(1), \quad 2h'(1) + h''(1).$$

In particular the $h(1)$ coefficient in the last expression cancels. Substituting into (II.46) and (II.45) gives

$$\begin{aligned} na_n r^n &= h(1) + \frac{1}{n} \{(-d/3 - \kappa T)h(1) \\ &\quad - (d + 1/3)(h(1) + h'(1)) - h'(1) - h''(1)/2\} + O(n^{-7/5}). \end{aligned} \tag{II.49}$$

Finally,

$$\begin{aligned} r^{-n} &= (n/T)^n e^d \left(1 - \frac{d^2}{2n} + O(\log^4 n/n^2)\right), \\ e^d &= n^{-1/3} T^{1/3} e^{-B}. \end{aligned}$$

Multiplication, division by the positive $h(1)$, and substitution of d give (II.9), hence (II.7) and (II.8). Cross-products and the radius remainder are $O(\log^4 n/n^2)$ and are smaller than the stated error. This completes the proof of Theorem 12.1.

20 Uniform proportional height correction

Proof of Theorem 12.2. Use the height-scaled variables $r = T/(m + d_m)$ and $G_m(s) = F_m(re^{s/m})$. The exact period identity at $\beta = n/m$ is

$$m\beta^q b_{n,m} r^n = \frac{1}{2\pi} \int_{-\pi m}^{\pi m} G_m^{(q)}(it) e^{-i\beta t} dt, \quad q = 30. \quad (\text{II.50})$$

A degree- k term acquires $(k/m)^q$, and the full period selects $k = n$, yielding exactly β^q . All growing-strip and tail estimates depend on height m , not on the target degree. Changing the oscillatory factor has modulus one for real β ; division by β^q is uniformly harmless on I .

Evaluate (II.48) at $x = \beta$ and divide by β^q . The first result is $h(\beta)$; the correction terms become

$$\beta h(\beta), \quad -h(\beta) - \beta h'(\beta), \quad 2h'(\beta) + \beta h''(\beta).$$

For example the third identity simplifies to $-h - xh'$ after division by x^q , and in the fourth the coefficient of h cancels identically, leaving $2h' + xh''$. Hence

$$\begin{aligned} mb_{n,m} r^n &= h(\beta) + \frac{J_\beta(d_m)}{m} + O_I(m^{-7/5}), \\ J_\beta(d) &= (-d/3 - \kappa T)\beta h(\beta) - (d + 1/3)(h(\beta) + \beta h'(\beta)) \\ &\quad - h'(\beta) - \beta h''(\beta)/2. \end{aligned}$$

Since $n = \beta m$ exactly,

$$r^{-n} = (m/T)^n e^{\beta d_m} \left(1 - \frac{\beta d_m^2}{2m} + O_I(\log^4 m/m^2) \right), \quad e^{\beta d_m} = m^{-\beta/3} T^{\beta/3} e^{-\beta B}.$$

Multiplication gives $\mathcal{Q}_\beta = J_\beta/h - \beta d_m^2/2$ in (II.10). Continuity and positivity imply $\min_I h > 0$; the derivatives are bounded on I . This converts the uniform absolute coefficient error into the asserted relative error. At $\beta = 1$ the expression reduces exactly to Theorem 12.1. $\square$

[write] **Checked, 6 October 2026.** The write checked the reduction: $\mathcal{Q}_1(d)$ is (II.9). The theorem refines Part I's Theorem 8.1, whose leading term it contains.

The exact-height formula can be used directly for $m = n + k$, for square-root height windows, or for $m = \lfloor \lambda n \rfloor$ with fixed $\lambda > 0$. In the last case the leading floor phase remains $\exp(-\{\lambda n\}/\lambda)$, as in Report 225. Formula (II.10) retains all such dependence by keeping the exact m and n/m ; replacing them prematurely by their limits can discard contributions as large as the correction itself.

21 A sharpened inverse formula

Set $\alpha = 4/3$, $\ell = \log c$, and

$$Q(u) = -u^2/18 + Au + C.$$

Theorem 12.1 implies

$$\log a_n = n \log(n/T) - \alpha \log n + \ell + \frac{Q(\log n)}{n} + O(n^{-7/5}), \quad (\text{II.51})$$

because the square of the first relative correction is $O(\log^4 n/n^2)$.

Corollary 21.1. *For an exact sequence value $y = a_n$, let $Y = \log y$ and, for large Y , define*

$$X = \frac{Y}{W(Y/T)}, \quad U = \log X, \quad q_X = \log(X/T) + 1, \quad (\text{II.52})$$

where W is the positive real Lambert branch. Put

$$\delta_0 = \frac{\alpha U - \ell}{q_X}, \quad \delta_1 = -\frac{\delta_0^2/2 - \alpha\delta_0 + Q(U)}{q_X}. \quad (\text{II.53})$$

Then

$$n = X + \delta_0 + \frac{\delta_1}{X} + O\left(\frac{X^{-7/5}}{\log X}\right). \quad (\text{II.54})$$

Proof. The definition of W gives $X \log(X/T) = Y$. The leading equivalent first gives $n = X + O(1)$. Define the explicit smooth model

$$F(x) = x \log(x/T) - \alpha \log x + \ell + Q(\log x)/x.$$

For bounded δ , Taylor expansion about X gives uniformly

$$\begin{aligned} F(X + \delta) - Y &= q_X \delta - \alpha U + \ell \\ &\quad + \frac{\delta^2/2 - \alpha\delta + Q(U)}{X} + O((1 + U^2)/X^2). \end{aligned}$$

Now $\delta_0 = O(1)$ and $\delta_1 = O(U)$, so substituting $\delta = \delta_0 + \delta_1/X$ cancels both retained orders. The residual is $O((1 + U^2)/X^2) = O(X^{-7/5})$. At the true integer n , (II.51) gives $F(n) - Y = O(n^{-7/5})$. On the bounded comparison interval $F'(x) \sim \log X$. The mean value theorem proves (II.54). Only F is differentiated; the asymptotic remainder is used solely at the true integer. $\square$

The constants δ_0, δ_1 in this inverse depend on X and are distinct from the fixed clock constant $d_\star$. The large real solution of $F(x) = \log y$ has the same error at $y = a_n$. With exact constants this yields eventual nearest-integer recovery, but neither an effective onset nor a certificate for arbitrary thresholds between consecutive sequence values. A fixed numerical error in $\log c$ creates an index error of order $1/\log X$, much larger than the fine rate in (II.54); the diagnostic decimal must not be substituted as if exact.

[write] **Relation to Part I and to the transseries volume, 6 October 2026.** δ_0 is the correction of Part I's (33); the write rechecked the two orders (II.53) by expanding $F(X + \delta_0 + \delta_1/X) - Y$. As in Part I, X is an instance of `p0:prop:factorial-core` of [11]; the two Newton-type corrections are proved directly and are not instances of `p0:thm:perturbed-inversion` (front matter).

22 Exact computation and numerical diagnostics

22.1 Exact recurrence and finite checks

Writing $1 + F_{h+1}(z) = \sum_{k \geq 0} c_k z^k$ and logarithmically differentiating the Euler product gives

$$b_j = \sum_{d|j} d[z^d]F_h(z), \quad c_0 = 1, \quad c_k = \frac{1}{k} \sum_{j=1}^k b_j c_{k-j}. \quad (\text{II.55})$$

Degree truncation is exact. The public code checks divisibility before every integer division. An independent finite-product implementation uses the binomial expansion of $(1 - z^d)^{-a}$ and agrees through degree and height twelve. A separately generated exact fixture contains all diagonal values through $n = 414$; only the first twenty-two displayed OEIS terms are an external sequence comparison. The fixture is not presented as a retrieved OEIS b-file.

The code also verifies finite formal-coordinate identities and the new clock algebra exactly over rational symbolic expressions. Such checks protect against algebraic transcription errors; they do not prove analytic remainder estimates, sector entrance, or the all-orders convergence of any formal series. All pass/fail checks use explicit exceptions and remain active under Python optimization. Local chunked decimal serialization handles long exact integers under `PYTHONINTMAXSTRDIGITS=640` without changing the global limit. The implementation caps and tested ranges are stated in the archive README.

22.2 Uncertified amplitude and derivative experiments

The accompanying optional floating-point programs solve a truncated Fatou inverse, take finitely many forward iterates, propagate derivatives, and numerically invert the transform. For example, P''' is the transform of $x^3h(x)$. Its absolutely convergent inversion and its first two ordinary x derivatives give $h(1)$, $h'(1)$, and $h''(1)$. Subtracting $12/(1-s)^4$ improves the observed tail; its inverse is $2x^3e^{-x}$, with values and derivatives at one equal to $2/e$, $4/e$, and $2/e$. After subtraction and quadrature, the derivative weights are undone by

$$h' = (x^3h)'|_1 - 3h(1), \quad h'' = (x^3h)''|_1 - 6h'(1) - 6h(1).$$

These are diagnostic computations with no interval error control for the truncated coordinate, inverse Newton step, forward propagation, finite contour, or quadrature.

Representative numerical values, not certified digit claims, are

$$\begin{aligned} h(1) &\approx 1.81460956, & \rho_1 &\approx -0.43688195, & \rho_2 &\approx -0.15328008, \\ \kappa &\approx -0.25434817, & A &\approx 0.60098418, & C &\approx -0.49816124. \end{aligned}$$

The associated leading amplitude is approximately 4.4923299. At $n = 414$, floating-point normalization of the exact integer gives

$$a_n(T/n)^n n^{4/3} \approx 4.50436880933,$$

while the unfitted first-correction prediction gives approximately 4.50433122939, a difference about 3.76×10^{-5} . No coefficient was fitted to these finite sequence values. Agreement and parameter sensitivity are useful diagnostics, not proofs of a remainder or of displayed digits.

[write] Checks and a comparison, 6 October 2026. At intake, with these constants the corrected prediction matched the exact a_n with residual 1.4×10^{-4} at $n = 200$, against 2.5×10^{-2} for the leading term. With Part I's diagnostic $h(1)$, $2^{1/3}h(1) = 2.2862647816$ agrees to eight significant digits with the iterated Bell report's exploratory amplitude $I(1)$ (Remark 37.1); both values are uncertified.

22.3 Reproducible archive

The archive contains this complete editable TeX, exact programs and tests, optional diagnostic programs and recorded examples, the exact fixture, source provenance, and a deterministic source-preserving builder. It also includes the unchanged Report 225 PDF and TeX as a clearly labeled supporting reference. The new report does not modify that earlier deliverable.

The builder validates its source manifest, runs the exact checks, creates a private pdfL^AT_EX format, compiles without shell escape, rejects overfull boxes, missing glyphs, and unresolved references, checks that source bytes were preserved, and writes a new PDF and sorted fixed-timestamp ZIP to a separate new directory. Ordinary and optimized runs under the restricted digit limit are compared as actual complete ZIPs and member bytes. Identical-toolchain byte reproducibility is distinct from mathematical correctness; different TeX or font versions may change PDF bytes. Every rendered page is also inspected before release.

[write] **The companion here, 6 October 2026.** Report 226’s own files are shipped as `code/226-firstcorr-*.py`, `data/226-firstcorr-*` and `226-firstcorr-sources-SOURCE_LEDGER.md`. Its eight byte copies of Report 225’s files and its `supporting/` copies of Report 225 are shipped once, as Report 225’s files and as Part I. Its builder and guard tests expect those files beside them; the report README gives the rerun route from the arrival commit.

23 Further questions and the boundary of the theorem

23.1 A genuine all orders expansion

The natural formal expectation is

$$\frac{a_n}{c(n/T)^n n^{-4/3}} \sim 1 + \sum_{r \geq 1} \frac{P_r(\log n)}{n^r}, \quad \deg P_r \leq 2r.$$

Only $P_1(X) = -X^2/18 + AX + C$ is proved here. If an all-orders construction establishes the stated degree bound and proves that no further contribution reaches degree $2r$, the leading coefficient would be

$$[X^{2r}]P_r(X) = \frac{(-1/18)^r}{r!}.$$

In particular the formal second-order highest term would be $\log^4 n / (648n^2)$. This conditional inference is not an all-orders theorem and does not follow from the first correction alone.

A proof needs an induction controlling arbitrary fixed scaled subsidiary order, large-ratio clock expansions and resonant logarithms, forced Fatou cohomological corrections, and growing-strip errors with extraction derivative order chosen for each target precision. The $\log y/540$ in N already shows why clocks cannot be treated as a plain power series in z : a $z^2 \log(1/z)$ contribution can enter the global phase. Exponentiating the radius alone misses this structure.

[write] **Still open, 6 October 2026.** Not settled by Part III, which works at one polynomial rate in m (Subsection 36.4). The conditional coefficient $(-1/18)^r/r!$ is recorded as unproved in Section 37, item 2.

23.2 Effective constants and certified inversion

Validated values of $h(1)$ and its derivatives require explicit sector remainder bounds, an enclosed inverse branch, controlled forward derivative propagation, interval quadrature, and an integrable certified tail. The present proof supplies qualitative constants and no finite onset. The exact formula for κ is elementary apart from Catalan’s constant, but certifying the full correction still requires the density derivative calculations.

An effective relative-error bound could turn the scalar inverse model into a certified threshold algorithm by combining exact small-index counts with rigorous upper and lower bounds. Theo-

rem 12.1 alone does not decide where a general input lies between a_n and a_{n+1} . One should separate this computational threshold problem from eventual rounding at exact sequence values.

[write] **Still open, 6 October 2026.** Section 37, items 1 and 3.

23.3 Endpoint height regimes and density structure

The compact-positive ratio theorem leaves open $n/m \rightarrow 0$ and $n/m \rightarrow \infty$. These require endpoint information about h and renewed uniform contour estimates. Smooth strict positivity away from zero supplies neither an expansion at zero nor a large-argument asymptotic. The positive-measure identity from Report 225,

$$(e^x - 1)\nu = \sum_{k \geq 2} \nu^{*k}/k!, \quad \nu(dx) = h(x) dx,$$

is a possible starting point, but exchanging endpoint limits or many derivatives with this convolution series requires separate arguments.

A transition theory connecting fixed-degree polynomiality, proportional height, and fixed-height partition asymptotics would also be useful. The forcing calculations here demonstrate that another initial generating function or multiset recurrence cannot simply inherit the same clock constants from the autonomous map. Its subsidiary terms and matching normalization must be recomputed.

[write] **Partly answered by Part III, 6 October 2026.** Part III treats $n/m \rightarrow 0$ at leading order on logarithmic bands (Theorem 31.1) and proves an expansion of h at zero (Proposition 27.1, Theorem 29.1, Corollary 29.2), so “an expansion at zero” is no longer missing. Open: $n/m \rightarrow \infty$ and a large-argument asymptotic of h , the correction on shrinking bands, the transition to fixed degree, other recurrences (Section 37, items 5, 7, 8).

Part III

Logarithmic height crossover for iterated Euler transforms

Bundle Report 228, archive `Report228.zip`, delivered as `src/report228.tex` (17 pages). Delivered title block “Logarithmic height crossover for iterated Euler transforms / Report 228 / 5 October 2026”. Printed as delivered, with `[write]` notes. Delivered Section k is Section $k + 23$ here, its statement $k.j$ is $(k + 23).j$, and its equation (k) is (III. k). Section 37, which closes the report, is added in the write.

Reading conventions of Part III	Elsewhere in the report
“Reports 225 and 226”, [6], [7]	Parts I and II
$m/(n \log n) \rightarrow \lambda$; $L = \log n$; $R(n, m)$ the ratio	Part I: $m = \lfloor \lambda n \rfloor$; $L(y)$ the clock; $R_h(z)$
K a constant; $C_0 = 1 - \gamma - \log 2$; $\mathcal{C}$ Catalan; $C(\beta)$	Parts I, II: K an index; C_0 bounds; Part II: G
the germ	Catalan
$\rho_j \in \mathbb{Q}$, c_j log-density coefficients; a_j Fatou	Part II: ρ_j ratios of derivatives of h ; all: c_k, b_j in
coefficients; b_j germ coefficients	the exact recurrence
$g(t) = t^{t/3}h(t)$; $W(x) = P(-x)$; $\mathcal{G}(u)$	Parts I, II: $g(w) = \log(1 + w)$; W Lambert; Part I:
	$G(u)$
$\tau = -\log r$; χ_n ; $y = m/n$	Part I: τ a time; Parts I, II: $y = w/z$

Abstract of this Part (as delivered). For the iterated Euler-transform array $A(n, m)$ in OEIS A290353, we prove

$$\frac{A(n, m)}{2(2/\pi)^n m^{n-1}} \rightarrow \exp\left(-\frac{1}{3\lambda}\right) \quad \text{if} \quad \frac{m}{n \log n} \rightarrow \lambda \in (0, \infty).$$

The result extends the positive fixed-slope analysis of Reports 225 and 226 to degree-to-height ratios decreasing as any fixed inverse power of the logarithm. We prove an exact small-argument Hankel representation for their normalized positive density h , show that $t^{t/3}h(t)$ has a one-sided smooth germ, and obtain every fixed logarithmic endpoint order with differentiated remainders. A rational coefficient generator isolates a single zeta value at each order of the logarithmic density. Exact cancellation of two logarithms yields every fixed inverse-logarithm order in the crossover window, without $\log \log n$ terms. We also invert the smooth suppression model and quantify rounded integer-height prescriptions and observed-ratio inverses. The exact Kaneiwa leading-monomial normalization changes the first correction constant by +1.

This unrefereed research proof uses explicitly stated analytic interfaces from the two preceding reports, included unchanged in the archive. It does not assert convergence of an infinite endpoint series, an all-orders expansion in $1/m$, an effective onset, discrete monotonicity, or exact integer-height recovery. Numerical comparisons are unvalidated floating diagnostics built from exact finite coefficients.

24 The array and the crossover

Define

$$F_0(z) = z, \quad F_{m+1}(z) = \exp\left(\sum_{j \geq 1} \frac{F_m(z^j)}{j}\right) - 1, \quad A(n, m) = [z^n]F_m(z) \quad (n \geq 1). \quad (\text{III.1})$$

This is the height-indexed array in OEIS A290353 [1]. It counts unlabeled rooted trees whose leaves all have the prescribed height. Equivalently, the new root carries a nonempty unordered multiset of trees from the preceding height, and

$$1 + F_{m+1}(z) = \prod_{d \geq 1} (1 - z^d)^{-A(d,m)}.$$

The generating functions have constant term zero; the tabular convention $A(0, m) = 1$ is imposed separately. There is no factorial normalization.

Put

$$\begin{aligned} T &= \frac{\pi}{2}, & B &= \frac{1}{2} - \frac{\pi}{6} - \frac{\pi^2}{16} - \frac{\log 2}{6}, \\ C_0 &= 1 - \gamma - \log 2, & K &= \frac{1 - \gamma + \log(T/2)}{3} - B, \end{aligned} \quad (\text{III.2})$$

where γ is Euler's constant. Our primary normalization is

$$R(n, m) = \frac{A(n, m)}{2T^{-n}m^{n-1}}. \quad (\text{III.3})$$

All asymptotics with two indices below take $n, m \rightarrow \infty$ in their specified ranges.

Theorem 24.1. *If $m/(n \log n) \rightarrow \lambda \in (0, \infty)$, then*

$$R(n, m) \longrightarrow e^{-1/(3\lambda)}. \quad (\text{III.4})$$

More precisely, let $\Lambda \subset (0, \infty)$ be compact, put $L = \log n$, and set $m = \lfloor \lambda n L \rfloor$. For every fixed $N \geq 1$, uniformly for $\lambda \in \Lambda$,

$$\log R(n, m) = -\frac{1}{3\lambda} + \frac{K}{\lambda L} + \sum_{j=2}^N \frac{c_j}{(\lambda L)^j} + O_{\Lambda, N}(L^{-N-1}), \quad (\text{III.5})$$

where the sum is empty for $N = 1$, and

$$c_j = \rho_j - \frac{\zeta(j)}{j3^j}, \quad \rho_j \in \mathbb{Q} \quad (j \geq 2). \quad (\text{III.6})$$

A finite coefficient algorithm determines every ρ_j . In particular,

$$c_2 = \frac{9 - \pi^2}{108}, \quad c_3 = \frac{17}{1080} - \frac{\zeta(3)}{81}.$$

For example, exponentiation gives

$$R(n, m) = e^{-1/(3\lambda)} \left\{ 1 + \frac{K}{\lambda L} + \frac{K^2/2 + (9 - \pi^2)/108}{\lambda^2 L^2} + O_{\Lambda}(L^{-3}) \right\}. \quad (\text{III.7})$$

Every order here means a fixed finite truncation followed by its asymptotic limit. It says neither that the infinite series converges nor that there is an all-orders expansion in powers of m^{-1} . One polynomial rate in the coefficient-transfer error will suffice for all fixed inverse-logarithm orders.

The smooth suppression-ratio inverses are proved in Section 34. With $\tau = -\log r$ and $\chi_n = (\log n)/3 - K$, their first height prescription is

$$\frac{m}{n} \approx \frac{\chi_n}{\tau} - \frac{c_2}{\chi_n} - \frac{c_3\tau}{\chi_n^2} - \frac{c_2^2\tau + c_4\tau^2}{\chi_n^3}. \quad (\text{III.8})$$

The inverse concerns a smooth asymptotic ratio; no unique discrete solution, least-height threshold, or exact recovery by rounding is claimed.

[write] Reading Part III against Parts I and II, 6 October 2026. “Report 225”, [6] and “Report 226”, [7] are Parts I and II; their included copies are byte-identical and not shipped. Here λ measures $m/(n \log n)$, not Part I’s slope m/n of Corollary 8.2; K is a constant, not Parts I and II’s entrance index; ρ_j are rational numbers, not Part II’s $h^{(j)}(1)/h(1)$; $L = \log n$, not the clock function.

25 Imported analytic interfaces and the scope of the result

The analytic foundation comes from Reports 225 and 226 [6, 7]. Their unchanged complete TeX and PDF files accompany this report, with hashes in the source ledger. We state exactly what is used; the present report supplies the endpoint, shrinking-ratio, and inverse arguments in full.

25.1 The normalized coordinate and density

Report 225, Sections 4 and 6, constructs an entire function P with

$$P(s+1) = e^{P(s)} - 1, \quad \Phi(w) = -\frac{2}{w} + \frac{1}{3} \text{Log } w + O(w). \quad (\text{III.9})$$

The coordinate Φ is analytic on every sufficiently small closed proper subsector of $|\arg w| < \pi$, with compatible branches on overlaps. It has zero additive constant in (III.9). The local inverse agrees with P on every sufficiently distant proper subsector of $|\arg(-s)| < \pi$; in particular

$$P(\Phi(w)) = w, \quad P'(s) = \frac{2}{s^2} + O\left(\frac{\log |s|}{|s|^3}\right) \quad (\text{III.10})$$

there. For the reciprocal inverse iteration

$$\mathcal{G}(u) = \frac{2}{\text{Log}(1 + 2/u)} = u + 1 - \frac{1}{3u} + O(u^{-2}), \quad (\text{III.11})$$

the normalized coordinate satisfies

$$\Psi(\mathcal{G}(u)) = \Psi(u) + 1, \quad \Psi(u) = u + \frac{1}{3} \text{Log } u + O(u^{-1}), \quad \Phi(w) = -\Psi(2/w) + \frac{1}{3} \log 2.$$

On nested proper sectors its inverse orbits satisfy $|\mathcal{G}^{\circ j}(u)| \geq c(|u| + j)$, and the normalized orbit limit is available on these sectors. These sector estimates, rather than a solely real-axis expansion, are used below.

The same report proves the positive Laplace representation

$$P(s) = \int_0^\infty e^{st} h(t) dt, \quad h \in C^\infty(0, \infty), \quad h(t) > 0 \ (t > 0). \quad (\text{III.12})$$

The density has every exponential moment. Fourier inversion applies to P' on vertical lines and to the fixed higher derivatives used below. We do not import endpoint smoothness or an endpoint expansion for h .

[write] **Where Part I proves this, 6 October 2026.** “Report 225, Sections 4 and 6” are Part I’s Sections 4 and 6: Lemma 4.1 (whose proof constructs $G = \mathcal{G}$, Ψ and $\Phi = -\Psi(2/w) + \frac{1}{3}\log 2$, with the orbit bound $|G^j(u)| \geq c(|u| + j)$), Proposition 6.1 and Corollary 6.2; Fourier inversion of P' is (30).

25.2 The absolute derivative estimates before division

For large m , set

$$\begin{aligned} d_m &= -\frac{1}{3}\log m + \frac{1}{3}\log T - B, & r_m &= \frac{T}{m + d_m}, \\ G_m(s) &= F_m(r_m e^{s/m}), & Q_m(s) &= (d_m + \frac{1}{3})s - \frac{1}{2}s^2 - \frac{1}{3}d_m - \kappa T. \end{aligned} \quad (\text{III.13})$$

Here κ is the fixed real constant of Report 226. Its value is immaterial for the transfer in this report; for a fully specified interface it is

$$\kappa = \frac{\pi^3}{96} + \frac{\pi^2}{12} - \frac{13\pi}{48} - \frac{11}{36} - \frac{\mathcal{C}}{6} - \frac{\pi \log 2}{24}, \quad \mathcal{C} = \sum_{j \geq 0} \frac{(-1)^j}{(2j+1)^2}.$$

Report 226, Sections “Derivative extraction and classical Fourier inversion” and “Uniform proportional height correction,” proves the following absolute estimates, all depending on height m , not on the target degree. With $M = m^{1/20}$, the thirtieth derivative of

$$G_m(s) - P(s) - m^{-1}Q_m(s)P'(s)$$

is $O(m^{-3/2})$ on $s = iu$, $|u| \leq M$, by analytic disk estimates on an enlarged strip. The finite-height intermediate derivative is $O(|u|^{-31})$, and the outer derivative is $O(m^{-31})$ on an interval of length $O(m)$. Their absolute tails are

$$O(M^{-30} + m^{-30}) = O(m^{-3/2}). \quad (\text{III.14})$$

The limiting derivative has $P^{(30)}(iu) = O(|u|^{-31})$, and

$$(Q_m P')^{(30)}(iu) = O(|u|^{-30} + \log m |u|^{-31}) \quad (\text{III.15})$$

for large $|u|$. No degree-to-height division has yet occurred.

Write $\tilde{G}_{m,30}(u)$ for $G_m^{(30)}(iu)$ on $[-\pi m, \pi m]$ and zero elsewhere. This notation means extension of the derivative’s values, not differentiation of a zero extension. Taking absolute values before integrating the preceding bounds proves

$$\left\| \tilde{G}_{m,30} - P^{(30)}(i \cdot) - \frac{(Q_m P')^{(30)}(i \cdot)}{m} \right\|_{L^1(\mathbb{R})} = O(m^{-7/5}). \quad (\text{III.16})$$

Indeed the central error is $O(Mm^{-3/2}) = O(m^{-29/20})$; (III.14) and the limiting P tail are $O(m^{-3/2})$; the correction tail after division by m is

$$O(m^{-1}(M^{-29} + \log m M^{-30})),$$

which is smaller still. These estimates even give the exponent 29/20 here; the weaker 7/5 is convenient and already sufficient. Equation (III.16) is an absolute L^1 consequence of the proved estimates, not an extrapolation of Report 226’s compact-slope relative theorem.

[write] **Where Part II proves this, 6 October 2026.** Report 226’s sections “Derivative extraction and classical Fourier inversion” and “Uniform proportional height correction” are Part II’s Sections 19 and 20: (II.45), (II.47), the outer-angle bound and the tail estimates following it, there at height n and here at height m (Part II’s proof of Theorem 12.2 makes the same change). d_m and Q_m are Part II’s d_m and Q_n with n replaced by m . The L^1 bound (III.16) itself is derived here.

25.3 Classical background and attribution

The endpoint calculation belongs to the universal unforced parabolic map $w \mapsto e^w - 1$. Finite sectorial Fatou expansions, their rational formal recursion, and Hankel inversion are classical methods. Dudko and Sauzin [8] give general sectorial asymptotics, including nonzero resiter; the present reciprocal germ has resiter 1/3. We give a direct finite-defect proof adapted to the precise normalization above.

The public iterated Bell report [5] already contains all-finite-order inverse-coordinate machinery and a related contour amplitude. Its inspected proportional-depth transfer fixes n/m in a positive compact interval and explicitly does not assert endpoint uniformity. In its notation, renamed here to avoid collisions, the small-variable coordinate is

$$\Psi_{\text{IBD}}(w) = -\Phi(w) + \frac{1}{3} \log 2.$$

This local identity follows from the common normalized reciprocal orbit limit. A formal contour match suggests an amplitude relation $I(\beta) = \beta 2^{\beta/3} h(\beta)$, but that identification requires its own analytic contour correspondence and is not used here. We neither attribute an endpoint theorem to that source nor claim a new array-specific origin for the universal coefficient calculus.

Prellberg’s 2002 seminar, as summarized by Mishna [9], already presents the parabolic logarithmic mechanism and its 1/3 specialization. Nagaev and Vakhel [10] give an important unforced antecedent: their critical branching large-deviation correction has the form $\exp[-\gamma u \log u/m]$, with $\gamma = 1/3$ in the Poisson case, in an appropriate moderate-deviation range. The heuristic high-moment saddle $u \asymp n$ places the logarithmic crossover in that range when $m/n \rightarrow \infty$. Passing from those probability estimates to a full moment equivalent needs tail domination, which is not supplied by this comparison; they do not furnish the forced Euler coefficient transfer proved here. The new scope relative to Report 225 is the quantitative transfer as n/m decreases to logarithmic scales, using the forced-array contour estimates and its constants T, B . No worldwide priority claim is made for the parabolic mechanism or universal endpoint calculus.

[write] **The iterated Bell report, 6 October 2026.** [5] is `a139383-iterated-bell-diagonals` in this repository; its article is unchanged since the pin, so the sections named are its current Sections 2, 4 (Part I there) and 12, 13, 17 (Part II there). Its Ψ is Ψ_{IBD} here and its $I(\beta)$ is the amplitude of its Theorem `ibd:pd:thm:main`. The relation $I(\beta) = \beta 2^{\beta/3} h(\beta)$ is unproved; at $\beta = 1$ the two sides agree to eight significant digits (Remark 37.1), and Section 37, item 9, records it as a question shared with that report, with a conditional consequence for its open endpoint (Remark 37.2). That report’s own literature section places the Nagaev–Vakhel range outside its compact-slope regime $m \asymp n$; here $m/n \rightarrow \infty$, which is the range named above.

26 Finite Fatou expansions with sectorial remainders

Lemma 26.1. *For every integer $N \geq 0$ there are unique rational numbers $a_1, \dots, a_N$ such that*

$$\Phi(w) = -\frac{2}{w} + \frac{1}{3} \operatorname{Log} w + \sum_{j=1}^N a_j w^j + O(w^{N+1}) \quad (\text{III.17})$$

uniformly on each sufficiently small closed proper sector. For every fixed derivative order k , the k th derivative of the remainder is $O(|w|^{N+1-k})$ on smaller closed sectors. The first values are

$$a_1 = -\frac{1}{36}, \quad a_2 = \frac{1}{540}, \quad a_3 = \frac{1}{7776}, \quad a_4 = -\frac{71}{435456}.$$

Proof. Choose a finite reciprocal model

$$V_N(u) = u + \frac{1}{3} \operatorname{Log} u + \sum_{j=1}^N v_j u^{-j}.$$

The expansion of (III.11) has rational coefficients. In $V_N(\mathcal{G}(u)) - V_N(u) - 1$, the new coefficient v_j first contributes $-j v_j u^{-j-1}$. Cancellation is therefore triangular, unique and rational, and gives

$$D_N(u) := V_N(\mathcal{G}(u)) - V_N(u) - 1 = O(u^{-N-2}).$$

Along an inverse orbit, $V_N(u) - u - (\operatorname{Log} u)/3 \rightarrow 0$, so the normalized orbit limit is the same Ψ as in Section 25. Telescoping gives the sign-sensitive identity

$$\Psi(u) - V_N(u) = \sum_{j \geq 0} D_N(\mathcal{G}^{\circ j}(u)). \quad (\text{III.18})$$

The sign is positive: the finite sum equals $V_N(\mathcal{G}^{\circ J}(u)) - J - V_N(u)$ before taking $J \rightarrow \infty$. The orbit bound gives absolute convergence and

$$\sum_{j \geq 0} |D_N(\mathcal{G}^{\circ j}(u))| \leq C \sum_{j \geq 0} (|u| + j)^{-N-2} = O(|u|^{-N-1}).$$

Transforming by $\Phi(w) = -\Psi(2/w) + (\log 2)/3$ proves (III.17); thus $a_j = -v_j/2^j$. Cauchy's derivative estimates on disks of radius proportional to $|w|$ inside nested sectors give every fixed differentiated remainder. Direct triangular substitution gives the stated coefficients. No convergence of the infinite formal coordinate series is required. $\square$

27 The endpoint value by direct Fourier inversion

The leading endpoint value follows already from the first-order interface in Report 225.

Proposition 27.1. *The density extends continuously to zero with*

$$h(t) = 2 + O(t \log(1/t)), \quad h(0) = 2. \quad (\text{III.19})$$

Proof. Fourier inversion of P' on the line $\operatorname{Re} s = -1/t$ gives

$$th(t) = \frac{e}{2\pi} \int_{\mathbb{R}} P'(-1/t + iu) e^{-iut} du.$$

The entire line is in the fixed inverse sector $|\arg(-s)| \leq \pi/2$. The leading term $2/s^2$ in (III.10) inverts to $2t$. The absolute error is bounded by a constant times

$$\int_{\mathbb{R}} \frac{\log \sqrt{t^{-2} + u^2}}{(t^{-2} + u^2)^{3/2}} du = O(t^2 \log(1/t)),$$

by $u = v/t$. Division by t proves the assertion. $\square$

[write] **Answers Parts I and II, 6 October 2026.** This answers, at the endpoint zero, the remark closing the proof of Part I's Proposition 6.1 and the questions of Subsections 11.4 and 23.3 that ask for an expansion of h at zero.

28 An exact Hankel representation

Set

$$\mathcal{A}(w) = \Phi(w) + 2/w - \frac{1}{3} \operatorname{Log} w = O(w). \quad (\text{III.20})$$

Fix $\pi/2 < \theta < \pi$ and $a > 0$. The contour $H = H_{\theta,a}$ starts at infinity on argument $-\theta$, travels inward to $ae^{-i\theta}$, follows the circle $|v| = a$ counterclockwise through the positive axis to $ae^{i\theta}$, and leaves along argument θ . All logarithms in the contour formula use the principal branch.

Theorem 28.1. *For sufficiently small positive t ,*

$$h(t) = \frac{2(2t)^{-t/3}}{2\pi i} \int_H e^v v^{t/3-2} \exp[-t\mathcal{A}(2t/v)] dv. \quad (\text{III.21})$$

The integral is absolutely convergent and is an exact identity. It supplies analytic continuation to a sufficiently small punctured sector about positive t .

Proof. Let $W(x) = P(-x)$. With an upward vertical contour, the absolutely convergent inverse of P' is

$$th(t) = -\frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} W'(x) e^{tx} dx, \quad c > 0. \quad (\text{III.22})$$

To check the sign, replacing s by $-x$ reverses the vertical orientation and $P'(-x) = -W'(x)$. Equivalently, change u to $-u$ in the absolutely convergent Fourier formula. We take $c = 1/t$.

Choose $\theta < \theta_1 < \pi$. For small t , the line and scaled contour $x = v/t$ lie beyond the inverse threshold in the sector $|\arg x| \leq \theta_1$, apart from no problematic compact portion since $|v| \geq a$. Truncate both at common large x -radius R . The circular connectors from the vertical endpoints to the rays lie to the left of the vertical line, so $|e^{tx}| \leq e$ there. The estimate $W'(x) = O(R^{-2})$ and connector length $O(R)$ give integral $O(R^{-1})$. Thus these connectors vanish as $R \rightarrow \infty$. The vertical integral converges absolutely by $W' = O(|x|^{-2})$, and the ray integrals by $\cos \theta < 0$. Cauchy's theorem deforms the upward line to the stated lower-inward, upper-outward Hankel orientation.

For the second deformation put

$$\begin{aligned} D_t(v) &= \frac{1}{3}(\text{Log } v - \log(2t)) - \mathcal{A}(2t/v), \\ x_\ell(v) &= v/t + \ell D_t(v), \quad 0 \leq \ell \leq 1. \end{aligned}$$

Uniformly for $|v| \geq a$ on H ,

$$|D_t(v)| \leq C(1 + |\log |v|| + |\log t|), \quad \sup_{v \in H} |tD_t(v)/v| \longrightarrow 0.$$

Consequently this homotopy stays in the larger proper inverse sector and satisfies $|x_\ell(v)| \geq |v|/(2t)$ for small t . Its endpoint at $\ell = 1$ is exactly

$$x(v) = -\Phi(2t/v) = v/t + \frac{1}{3}(\text{Log } v - \log(2t)) - \mathcal{A}(2t/v). \quad (\text{III.23})$$

The compatible principal branches are valid since $\arg(2t/v) = -\arg v$ stays strictly inside $(-\pi, \pi)$. In particular no negative-real branch cut is crossed.

At ray endpoints $v = Re^{\pm i\theta}$ the homotopy connectors have length $O(\log R + |\log t|)$ and

$$\text{Re}(tx_\ell(v)) = R \cos \theta + O(t \log R + t |\log t|).$$

Their integrals tend to zero exponentially as $R \rightarrow \infty$, for fixed small t . On each finite truncated homotopy the integrand $W'(x)e^{tx}$ is entire, so Cauchy's theorem applies. This is a deformation of paths; it does not require global univalence of the map $v \mapsto x(v)$.

By the imported inverse identification,

$$W(x(v)) = P(\Phi(2t/v)) = 2t/v, \quad W'(x(v)) dx = -2tv^{-2} dv.$$

Substitution into (III.22), retaining its minus sign, gives (III.21), including the factor 2. As an orientation check, $(2\pi i)^{-1} \int_H e^v v^{-2} dv = 1$, which gives the positive endpoint value.

Finally choose $\eta > 0$ with $\eta + \theta < \pi$. For complex t with $0 < |t| < \epsilon$ and $|\arg t| < \eta$, every $2t/v$ stays in a fixed coordinate sector. Use the corresponding branch of $\text{Log } t$ in (III.21). Uniform convergence on compact subsets follows because $e^{\text{Re } v}$ on the rays dominates all fixed polynomial and logarithmic factors. The resulting holomorphic function agrees with the real density for positive t by the identity already proved. $\square$

[write] **A consequence, 6 October 2026.** Since the continuation is holomorphic on a punctured sector containing $(0, \epsilon)$ and agrees with h there, h is real analytic on $(0, \epsilon)$. Real analyticity on all of $(0, \infty)$, asked in Part I's Subsection 11.4, is not shown (Section 37, item 6).

29 A smooth endpoint germ and the logarithmic transseries

Here “endpoint transseries” refers only to finite asymptotic expansions in powers $t^j(\log t)^r$ and their differentiated remainders. No summability, exponentially small completion, or convergence assertion is attached to the term.

Theorem 29.1. *Let $g(t) = t^{t/3}h(t)$ for $t > 0$. There are real coefficients b_j , with $b_0 = 2$, such that for every fixed $N, k \geq 0$,*

$$\frac{d^k}{dt^k} \left(g(t) - \sum_{j=0}^N b_j t^j \right) = O(t^{N+1-k}) \quad (t \downarrow 0). \quad (\text{III.24})$$

Consequently g has a one-sided C^∞ extension to $[0, \epsilon)$ and $b_j = g^{(j)}(0)/j!$. This does not assert analyticity at zero.

Proof. Cancel the explicit $t^{-t/3}$ in (III.21) to obtain

$$g(t) = \frac{2 \cdot 2^{-t/3}}{2\pi i} \int_H e^v v^{-2} \exp \left[\frac{t}{3} \operatorname{Log} v - t\mathcal{A}(2t/v) \right] dv. \quad (\text{III.25})$$

Use Lemma 26.1 to any required finite order. On H , $|v| \geq a$, and the remainder after retaining $\mathcal{A}$ through degree q is uniformly $O(|t|^{q+1}|v|^{-q-1})$. Multiplication by t improves its order once more. The remaining factors $2^{-t/3}$ and $v^{t/3}$ are entire in t . Their finite Taylor remainders are bounded by fixed powers of $1 + |\operatorname{Log} v|$ times a small fixed power of $|v|$. Along the rays these are integrably dominated by $e^{\operatorname{Re} v}$; the inner arc is compact and avoids zero. Finite expansion and integration thus yield, uniformly in a smaller punctured t -sector,

$$g(t) = \sum_{j=0}^N b_j t^j + O(|t|^{N+1}).$$

Apply Cauchy's estimate to the actual holomorphic remainder on disks of radius δt centered at positive t . This proves (III.24), rather than formally differentiating an unsupported pointwise expansion. For a prescribed derivative order choose $N \geq k$; its limit is $k!b_k$. Choosing one additional order establishes compatibility of these endpoint derivatives. The constant follows from Proposition 27.1, or directly from the orientation check in Theorem 28.1. $\square$

The classical reciprocal-Gamma Hankel identity, on the orientation specified above, is

$$\frac{1}{2\pi i} \int_H e^v v^{-z} dv = \frac{1}{\Gamma(z)}.$$

For a finite nonnegative multi-index $\mathbf{r} = (r_1, r_2, \dots)$, let $R_{\mathbf{r}} = \sum r_j$ and $J_{\mathbf{r}} = \sum j r_j$. Expanding the last exponential in (III.25) gives the finite-order generator

$$g(t) \sim 2 \cdot 2^{-t/3} \sum_{\mathbf{r}} \frac{(-1)^{R_{\mathbf{r}}} 2^{J_{\mathbf{r}}}}{\Gamma(2 + J_{\mathbf{r}} - t/3)} \left(\prod_j \frac{a_j^{r_j}}{r_j!} \right) t^{R_{\mathbf{r}} + J_{\mathbf{r}}}. \quad (\text{III.26})$$

At any specified degree only finitely many multi-indices occur. Equation (III.26) is interpreted by truncation; Theorem 29.1 proves its remainder, and no convergence of the full multi-index sum is used.

Corollary 29.2. *Writing $X = \log t$, for every fixed $N \geq 0$,*

$$h(t) = \sum_{j=0}^N t^j H_j(X) + O(t^{N+1}(1 + |X|)^{N+1}), \quad H_j(X) = \sum_{r=0}^j b_{j-r} \frac{(-X/3)^r}{r!}. \quad (\text{III.27})$$

The k th derivative of its remainder is $O(t^{N+1-k}(1 + |\log t|)^{N+1})$ for every fixed k . In particular H_j has degree exactly j , with leading coefficient $2(-1/3)^j/j!$.

Proof. Use $h(t) = \exp[-t \operatorname{Log} t/3]g(t)$ in the punctured sector and multiply finite expansions. The exponential remainder obeys the stated power-log bound. Cauchy estimates on the same proportional disks give the differentiated bound; equivalently one may differentiate the finite product with its controlled remainder. Since $b_0 = 2$, its highest logarithmic coefficient is the stated nonzero number. $\square$

In particular,

$$h(t) = 2 + \frac{2}{3}t(C_0 - \log t) + \frac{t^2}{9} \left((\log t - C_0)^2 + \frac{3}{2} - \frac{\pi^2}{6} \right) + O(t^3(1 + |\log t|)^3). \quad (\text{III.28})$$

The differentiated estimates imply

$$h'(t) = -\frac{2}{3}(\log t + \gamma + \log 2) + O(t \log^2 t), \quad h''(t) = -\frac{2}{3t} + O(\log^2 t).$$

Thus the original density h is not C^1 at zero, although the explicitly renormalized germ g is smooth.

30 The rational and zeta coefficient structure

Because $g(t) = 2 + O(t)$ on a sector, a branch of $\log(g(t)/2)$ exists near the endpoint, approaching zero. Theorem 29.1 gives ordinary finite Taylor remainders for this logarithm and each fixed derivative. Therefore

$$\log \frac{h(t)}{2} = -\frac{t}{3} \log t + \frac{C_0}{3}t + \sum_{j=2}^N c_j t^j + O(t^{N+1}). \quad (\text{III.29})$$

After removing the single explicit $t \log t$ term, every k th derivative of the remainder is $O(t^{N+1-k})$.

Proposition 30.1. *The coefficients in (III.29) have the form (III.6). Their first rational parts are*

$$\begin{aligned} \rho_2 &= \frac{1}{12}, & \rho_3 &= \frac{17}{1080}, & \rho_4 &= \frac{13}{3888}, \\ \rho_5 &= \frac{2071}{2449440}, & \rho_6 &= \frac{5131261}{22044960000}, & \rho_7 &= \frac{261791507}{3968092800000}. \end{aligned}$$

Proof. Factor $1/\Gamma(2 - t/3)$ out of (III.26). The remaining formal ordinary series $S(t)$ has rational coefficients and constant term one, because each Gamma ratio is exactly

$$\frac{\Gamma(2 - t/3)}{\Gamma(2 + J - t/3)} = \prod_{k=2}^{J+1} (k - t/3)^{-1},$$

with empty product one when $J = 0$. Thus

$$\frac{g(t)}{2} \sim \frac{2^{-t/3}}{\Gamma(2 - t/3)} S(t), \quad \log S(t) \in t^2 \mathbb{Q}[[t]]. \quad (\text{III.30})$$

At each finite order this is justified by the already proved expansion. The familiar local Gamma expansion yields

$$-\log \Gamma(2 - t/3) = \frac{1 - \gamma}{3}t + \sum_{j \geq 2} \frac{1 - \zeta(j)}{j 3^j} t^j.$$

Combining it with (III.30) proves the rational/zeta form and the linear coefficient $C_0/3$. For example,

$$S(t) = 1 + \frac{t^2}{36} + \frac{11t^3}{3240} + O(t^4),$$

which immediately gives c_2 and c_3 . Finite multi-index enumeration with the rational a_j gives the further displayed ρ_j . The archive supplies exact-arithmetic reproduction of these identities, including a separate finite reversion check. The analytic remainders follow from the preceding proofs, not from symbolic checking. $\square$

31 Transfer to shrinking degree to height ratios

Theorem 31.1. *Uniformly over positive integer degrees n as $m \rightarrow \infty$, writing $\beta = n/m$, one has the absolute estimate*

$$mA(n, m)r_m^n = h(\beta) + O\left(\frac{\log m}{m\beta^{30}}\right). \quad (\text{III.31})$$

For every fixed $B_0 > 0$, uniformly for $0 < \beta \leq B_0$,

$$\begin{aligned} A(n, m) &= \frac{h(\beta)}{m} (m/T)^n e^{\beta d_m} \{1 + O(E(m, \beta))\}, \\ E(m, \beta) &= \frac{\log m}{m\beta^{30}} + \frac{\beta \log^2 m}{m}. \end{aligned} \quad (\text{III.32})$$

The relative expression is an asymptotic equivalent whenever its stated error tends to zero. In particular this holds uniformly on every fixed logarithmic band

$$(\log m)^{-D} \leq \beta \leq B_0, \quad D > 0 \text{ fixed}. \quad (\text{III.33})$$

Proof. Cauchy's estimates for the inverse-sector expansion of P give $P^{(q)}(iu) = O(|u|^{-q-1})$ at infinity for every fixed $q \geq 1$. Leibniz' rule, the quadratic form of Q_m , and $d_m = O(\log m)$ imply

$$\|(Q_m P')^{(30)}(i \cdot)\|_1 = O(\log m).$$

The bounded interval uses analyticity; on the tails (III.15) is integrable. Hence the pre-division estimate (III.16) implies

$$\|\tilde{G}_{m,30} - P^{(30)}(i \cdot)\|_1 = O(\log m/m). \quad (\text{III.34})$$

For every positive integer n , termwise full-period extraction is exact:

$$m\beta^{30} A(n, m)r_m^n = \frac{1}{2\pi} \int_{-\pi m}^{\pi m} G_m^{(30)}(iu) e^{-i\beta u} du. \quad (\text{III.35})$$

Indeed a monomial of degree k acquires $(k/m)^{30}$; orthogonality over a period selects $k = n$, leaving β^{30} . The contour is inside the disk of analyticity for large m . No integration by parts or endpoint matching assumption is needed.

Fourier inversion of $P^{(30)}$ gives $\beta^{30}h(\beta)$ for each $\beta > 0$. The oscillatory factor has absolute value one, so (III.34) and division by β^{30} give (III.31). The constant before that division is independent of the target degree. The displayed β^{-30} loss is therefore genuine and has not been discarded. Only this absolute assertion is globally uniform in β .

By Proposition 27.1, strict positivity, and continuity, $\min_{[0, B_0]} h > 0$. Thus division by $h(\beta)$ is uniform on this interval. Furthermore

$$\begin{aligned} r_m^{-n} &= (m/T)^n \exp\{n \log(1 + d_m/m)\} \\ &= (m/T)^n e^{\beta d_m} \left\{1 + O\left(\frac{\beta \log^2 m}{m}\right)\right\} \end{aligned}$$

uniformly for $\beta \leq B_0$. This proves (III.32). On (III.33),

$$E(m, \beta) = O\left(\frac{(\log m)^{30D+1}}{m} + \frac{\log^2 m}{m}\right) = o(1).$$

The constants may depend on B_0, D , and the fixed derivative order. There is no uniform relative theorem for arbitrary $\beta \downarrow 0$ here. $\square$

[write] **Relation to Parts I and II, 6 October 2026.** On a fixed compact β -range (III.32) is Part I's (31) with the rate $O(\log m/m + \log^2 m/m)$ in place of $o(1)$; Part II's Theorem 12.2 is sharper there. The new content is the uniformity down to $(\log m)^{-D}$.

32 Exact logarithmic cancellation and the crossover proof

Combining (III.3) and (III.32) gives

$$R(n, m) = \frac{h(\beta)}{2} e^{\beta d_m} \{1 + O(E(m, \beta))\}.$$

In a logarithmic band with sufficiently small upper endpoint, its logarithm is legitimate because $E = o(1)$ and $h > 0$. Equation (III.29) then gives, for every fixed $N \geq 1$,

$$\log R(n, m) = -\frac{\beta}{3} \log n + K\beta + \sum_{j=2}^N c_j \beta^j + O(\beta^{N+1} + E(m, \beta)). \quad (\text{III.36})$$

The identity behind this formula is exact:

$$-\frac{\beta}{3} \log \beta - \frac{\beta}{3} \log m = -\frac{\beta}{3} \log n, \quad \log \beta + \log m = \log n.$$

It removes potential $\log \log n$ terms before any asymptotic substitution.

Proof of Theorem 24.1. If $m/(n \log n) \rightarrow \lambda \in (0, \infty)$, then $\beta \rightarrow 0$, $\beta \log n \rightarrow 1/\lambda$, and $\log m \sim \log n$. The ratio belongs, for example, to a fixed logarithmic band with $D = 2$ for all sufficiently large n , and $E \rightarrow 0$. Equation (III.36) gives (III.4).

For the uniform fixed-order assertion let $m = \lfloor \lambda n L \rfloor$, $L = \log n$, and $\lambda \in \Lambda$. Then

$$\beta = \frac{1}{\lambda L} + O_\Lambda \left(\frac{1}{n L^2} \right), \quad E(m, \beta) = O_\Lambda(L^{30}/n).$$

Integer rounding changes βL by $O_\Lambda(1/(nL))$. These errors are smaller than every fixed inverse power of L . Substitute the finite expansion (III.36); its endpoint remainder becomes $O_{\Lambda, N}(L^{-N-1})$. This proves (III.5) and, by ordinary finite exponentiation, (III.7). No estimate of arbitrarily high order in m^{-1} was used. $\square$

More generally, whenever $\beta \rightarrow 0$ inside a fixed logarithmic band,

$$R(n, m) = \exp \left[-\frac{n}{3m} \log n \right] \{1 + O(\beta + E(m, \beta))\}. \quad (\text{III.37})$$

Thus $R \rightarrow 1$ if $m/(n \log n) \rightarrow \infty$ and m/n grows at most as a fixed power of $\log n$. Also $R \rightarrow 0$ if $m/n \rightarrow \infty$ while $m/(n \log n) \rightarrow 0$; this range lies in a logarithmic band because $m/n = o(\log n)$. Fixed degree followed by height tending to infinity is a different limit and is not covered by this transfer.

33 Kaneiwa normalization and its first correction

Kaneiwa [3] proves that his $p(r; n)$ is a degree- $(n - 1)$ polynomial in r with leading coefficient

$$\ell_n = \frac{E_{n-1}}{(n-1)!} = [x^{n-1}](\tan x + \sec x). \quad (\text{III.38})$$

His indexing is $p(r; n) = A(n, r + 1)$, so $r = m - 1$. His fixed-degree polynomial theorem alone provides no uniform error as n and r grow together.

The nearest pole of $\tan x + \sec x$ is $T = \pi/2$, with principal part $2/(T - x)$. The apparent singularity at $-T$ is removable; the next pole is $-3T$. Subtract the principal part and use Cauchy's coefficient estimate on any fixed circle of radius strictly between T and $3T$. For every fixed $1 < c < 3$ this gives

$$\frac{\ell_n}{2T^{-n}} = 1 + O(c^{-n}). \quad (\text{III.39})$$

If $n \rightarrow \infty$ and $\beta = n/m \rightarrow 0$, then

$$(n - 1) \log(1 - 1/m) = -\beta + O(1/m + n/m^2) \rightarrow 0.$$

Consequently the crossover limit is unchanged when (III.3) is replaced by

$$R_K(n, m) = \frac{A(n, m)}{\ell_n(m - 1)^{n-1}}. \quad (\text{III.40})$$

For higher corrections the height shift must be retained. Directly,

$$\log R_K(n, m) = \log R(n, m) + \beta + O(1/m + n/m^2 + c^{-n}). \quad (\text{III.41})$$

In the crossover window these residual errors are smaller than every fixed inverse power of $\log n$. Thus Theorem 24.1 and its exponential version hold for R_K with K replaced everywhere by $K + 1$, while every c_j for $j \geq 2$ is unchanged. In particular the second-order exponential coefficient becomes $((K + 1)^2/2 + c_2)/(\lambda^2 L^2)$. If one instead normalizes by $\ell_n m^{n-1}$, K stays unchanged.

[write] **Kaneiwa in Part I, 6 October 2026.** The indexing $p(r; n) = A(n, r + 1)$ is Part I's $p(r; n) = [z^n]F_{r+1}$ (Subsection 1.1), where the same Propositions 1 and 2 of Kaneiwa are credited.

34 Smooth suppression ratio inverses

Fix a compact interval $\mathcal{R} = [r_0, r_1] \subset (0, 1)$ and write

$$\tau = -\log r \in [\tau_0, \tau_1] = [-\log r_1, -\log r_0], \quad \chi_n = \frac{\log n}{3} - K. \quad (\text{III.42})$$

The suppression logarithm τ is distinct from the density argument. Define the actual smooth germ

$$C(\beta) = \log \frac{g(\beta)}{2} - \frac{C_0}{3}\beta. \quad (\text{III.43})$$

It has $C(0) = C'(0) = 0$ and finite Taylor expansions $C(\beta) = \sum_{j=2}^N c_j \beta^j + O(\beta^{N+1})$, with all fixed differentiated remainders. This definition uses the actual density, not an assumed convergent formal sum.

34.1 The unique small model root

Proposition 34.1. *There is $\beta_0 > 0$ such that for every sufficiently large n and every $r \in \mathcal{R}$ the equation*

$$\tau = \chi_n \beta_* - C(\beta_*) \quad (\text{III.44})$$

has exactly one solution $\beta_(n, r) \in (0, \beta_0)$. Uniformly in r , $\beta_* \sim \tau/\chi_n$.*

Proof. Choose β_0 so that $g > 0$ and C is smooth on $[0, \beta_0]$. For large n , the derivative of the right side is $\chi_n - C'(\beta) \geq \chi_n/2$ throughout that interval. Its values at the endpoints are zero and a quantity tending to infinity. The intermediate value theorem and strict increase prove existence and uniqueness in this endpoint interval. With $C(\beta) = O(\beta^2)$ and τ in a fixed positive compact interval, first $\beta_* = O(\chi_n^{-1})$, and then $\beta_* = \tau/\chi_n + O(\chi_n^{-3})$. No uniqueness outside the stated interval is asserted. $\square$

34.2 Every fixed reversion order

Put $x = 1/\chi_n$ and $u = \beta_*/x$. Equation (III.44) becomes

$$u = \tau + C(xu). \quad (\text{III.45})$$

Finite substitution determines every coefficient recursively. The first terms are

$$\beta_* = \frac{\tau}{\chi_n} + \frac{c_2 \tau^2}{\chi_n^3} + \frac{c_3 \tau^3}{\chi_n^4} + \frac{2c_2^2 \tau^3 + c_4 \tau^4}{\chi_n^5} + O_{\mathcal{R}}(\chi_n^{-6}), \quad (\text{III.46})$$

$$y_* := \frac{1}{\beta_*} = \frac{\chi_n}{\tau} - \frac{c_2}{\chi_n} - \frac{c_3 \tau}{\chi_n^2} - \frac{c_2^2 \tau + c_4 \tau^2}{\chi_n^3} + O_{\mathcal{R}}(\chi_n^{-4}). \quad (\text{III.47})$$

The absent χ_n^{-2} term in (III.46) and absent constant term in (III.47) are consequences of $C'(0) = 0$. Since $c_2 < 0$, the first displayed correction to the height model is positive.

Theorem 34.2. *For every fixed $J \geq 0$ there are explicitly generated polynomials $d_j(\tau)$ such that*

$$y_J(n, r) = \frac{\chi_n}{\tau} + \sum_{j=1}^J d_j(\tau) \chi_n^{-j}, \quad (\text{III.48})$$

$$y_*(n, r) - y_J(n, r) = O_{\mathcal{R}, J}(\chi_n^{-J-1}).$$

Here $d_1 = -c_2$, $d_2 = -c_3 \tau$, and $d_3 = -c_2^2 \tau - c_4 \tau^2$. The coefficient d_J requires only $c_2, \dots, c_{J+1}$.

Proof. Take a finite formal truncation $U_M(\tau, x)$ of (III.45) through degree M . Taylor's theorem for the smooth C gives a residual

$$U_M - \tau - C(xU_M) = O_{\mathcal{R}, M}(x^{M+1}).$$

Both the true u and this truncation stay in a common positive compact interval, uniformly in τ . There

$$\frac{\partial}{\partial u} \{u - \tau - C(xu)\} = 1 - xC'(xu) \geq \frac{1}{2}$$

for small x . The real mean-value theorem bounds $u - U_M$ by twice the residual. Since u stays bounded away from zero, finite reciprocal expansion then gives every claimed order for $y_* = 1/(xu)$.

Write $\beta_* = \tau x V$. The equation becomes $V = 1 + \sum_{j \geq 2} c_j \tau^{j-1} x^j V^j$ at each finite order. Every nonconstant coefficient of V is therefore a polynomial divisible by τ . The same holds for $1/V - 1$, so division by τx introduces no negative powers of τ beyond the leading $1/(\tau x)$. Formal recursion shows that the coefficient of x^j in y_* first involves c_{j+1} , proving polynomiality and the stated dependency. In particular direct substitution gives (III.46) and (III.47) with the displayed signs.

Equivalently one can solve the truncated polynomial model $\tau = \chi_n \beta - \sum_{j=2}^{J+1} c_j \beta^j$ in its unique small-root interval. Its omitted C remainder is $O(\beta^{J+2})$; division by the model derivative costs χ_n^{-1} and reciprocation costs χ_n^2 . Thus its height differs from y_* by $O(\chi_n^{-J-1})$, and it has the same y_J . Other polynomial roots are irrelevant. No convergence of an infinite reversion series is invoked. $\square$

34.3 Rounded height prescriptions and observed ratios

Theorem 34.3. *Choose any integer $m_J(n, r)$ with $|m_J - ny_J(n, r)| \leq 1$. Then, uniformly for $r \in \mathcal{R}$,*

$$\log \frac{R(n, m_J)}{r} = O_{\mathcal{R}, J} \left(\chi_n^{-J-2} + \frac{\chi_n^{30}}{n} \right). \quad (\text{III.49})$$

Rounding the exact smooth model ny_ instead gives error $O_{\mathcal{R}}(\chi_n^{30}/n)$ in this logarithmic ratio.*

Conversely, suppose integer heights have $m/n \asymp \log n$ and the actual observed ratio $r_n = R(n, m)$ lies in $\mathcal{R}$. Then

$$\frac{m}{n} = y_J(n, r_n) + O_{\mathcal{R}, J} \left(\chi_n^{-J-1} + \frac{\chi_n^{31}}{n} \right), \quad (\text{III.50})$$

with constants also depending on the fixed comparison bounds in $m/n \asymp \log n$.

Proof. Using the exact smooth germ in (III.32), rather than a Taylor truncation, gives on this height window

$$\log R(n, m) = -\chi_n \beta + C(\beta) + O(\chi_n^{30}/n). \quad (\text{III.51})$$

Indeed $m \asymp n\chi_n$, $\beta \asymp \chi_n^{-1}$, and $E(m, \beta) = O(\chi_n^{30}/n)$. As a function of $y = m/n$, the smooth part is

$$F_n(y) = -\chi_n/y + C(1/y), \quad F'_n(y) = \frac{\chi_n - C'(1/y)}{y^2} \asymp \chi_n^{-1}$$

uniformly on any common crossover-height window. The error $y_J - y_* = O(\chi_n^{-J-1})$ therefore costs $O(\chi_n^{-J-2})$ in the log ratio. Integer rounding costs $O(1/(n\chi_n))$, which is absorbed by χ_n^{30}/n . Since $F_n(y_*) = -\tau = \log r$, this proves (III.49) and the exact-model statement.

For the converse, take $\tau_n = -\log r_n$ at the actual observed ratio. Equation (III.51) and the lower derivative bound show that $m/n - y_*(n, r_n) = O(\chi_n^{31}/n)$. Add Theorem 34.2 to obtain (III.50). Inverting the ratio error costs precisely one extra factor χ_n . $\square$

These formulas use a specified target r , or the actual r_n at each n . Replacing r_n by its mere limiting value generally loses higher-order information unless an appropriate convergence rate is known separately. The available coefficient theorem gives no discrete monotonicity, exact ratio solvability, least integer height, or effective onset. Thus the estimates are smooth prescriptions and asymptotic observed-ratio inverses, not threshold or exact-integer-recovery theorems.

For the Kaneiwa normalization (III.40), replace K by $K + 1$ throughout this section, hence replace χ_n by $(\log n)/3 - K - 1$. The residual shift in (III.41) is smaller than χ_n^{30}/n on the window. The coefficients c_j , reversion signs, and stated orders remain unchanged.

[write] **No Lambert function here, 6 October 2026.** Unlike Parts I and II, these inverses involve no Lambert W : the core of (III.44) is linear (the case $b = 0$ of `p0:thm:lambert-core` in [11]), and the reversion with the smooth germ C is proved by Taylor’s theorem; it is an analogue, not an instance, of `p0:thm:perturbed-inversion`, whose hypotheses ask for analyticity, and the rounded prescriptions are not instances of `p0:thm:staircase`. The write rechecked (III.46)–(III.47) by substitution.

35 Exact finite comparisons and reproducibility

35.1 An exact finite coefficient algorithm

Logarithmic differentiation of the Euler product gives, with the preceding-height coefficients $f_d = A(d, m)$,

$$b_j = \sum_{d|j} df_d, \quad c_0 = 1, \quad c_k = \frac{1}{k} \sum_{j=1}^k b_j c_{k-j}, \quad A(k, m+1) = c_k \quad (k \geq 1). \quad (\text{III.52})$$

Truncation at degree N is exact for all coefficients retained. The public implementation uses integers and checks divisibility at each division. A separate finite-product implementation checks small tables using $(1 - z^d)^{-a} = \sum_{k \geq 0} \binom{a+k-1}{k} z^{dk}$. The working limits, degree and height at most 640, are explicit implementation caps, not theorem restrictions. Computing height M through degree N uses $O(MN^2)$ integer operations and $O(N)$ live coefficients apart from saved outputs; bit costs grow with coefficient size.

The code uses explicit exceptions for all pass/fail conditions, so checks remain active under Python’s `-O`. Chunked integer decimal conversion works with the minimum configurable integer-string digit limit 640 and does not change any process-wide limit. A guard test exercises a value exceeding that digit count. The exact coefficient generation is independent of any decimal normalization or asymptotic theorem.

35.2 Finite crossover diagnostics

Table 1 uses exact recurrence values at $n = 20, 40, 80$ and $m = \lfloor \lambda n \log n \rfloor$ for $\lambda = 1/2, 1, 3/2$. All heights are below 640. For finite-height comparison define

$$R_2(n, \lambda) = e^{-1/(3\lambda)} \left(1 + \frac{K}{\lambda \log n} + \frac{K^2/2 + c_2}{(\lambda \log n)^2} \right). \quad (\text{III.53})$$

This is the explicit second-order ratio expansion, rather than the exponential of a truncated logarithm. The exact coefficient uses the displayed integer height. The nominal floor choices agree at 80 and 160 decimal digits; this is a computational check, not an interval certificate for the transcendental floor. Values of R and R_2 are rounded floating evaluations, not interval enclosures; only $A(n, m)$ is computed exactly. The column $e^{-1/(3\lambda)}$ is the limiting ratio. These finite examples illustrate the size of the correction and do not establish an effective error bound or a certified asymptotic onset. The accompanying JSON records exact decimal coefficients, the chosen integer heights, normalization formulas, and diagnostic precision. The coefficient generator separately verifies the rational values through ρ_7 and the finite inverse identities. Symbolic and exact-integer checks are reproducible algebra; floating agreement is not an analytic remainder certificate.

[write] **The labelled counterpart, 6 October 2026.** Remark 37.2 gives the same comparison for the labelled array $H(n, m)$ at the same heights, against a conditional prediction.

n	λ	m	$R(n, m)$	$e^{-1/(3\lambda)}$	$R_2(n, \lambda)$
20	0.5	29	0.89828117	0.51341712	0.86765860
20	1	59	0.94696428	0.71653131	0.93775982
20	1.5	89	0.96417635	0.80073740	0.95910776
40	0.5	73	0.80413195	0.51341712	0.78974074
40	1	147	0.89724904	0.71653131	0.89222870
40	1.5	221	0.93039388	0.80073740	0.92738192
80	0.5	175	0.74764498	0.51341712	0.73949701
80	1	350	0.86485251	0.71653131	0.86215689
80	1.5	525	0.90778645	0.80073740	0.90621689

Table 1: Unvalidated floating comparisons built from exact finite coefficients. The two displayed approximations have different purposes: the limit records the crossover theorem, and R_2 retains the first two crossover ratio corrections.

35.3 Reproducing the archive

The archive includes editable TeX, exact and symbolic programs, generated comparisons, the source ledger, and unchanged Reports 225 and 226. Its builder verifies a source manifest, reruns the public checks, copies inputs to a new external output directory, creates a private pdfL^AT_EX format, compiles without shell escape, and rejects unresolved references, missing characters, or overfull boxes. Sources are preserved. Sorted ZIP members have fixed timestamps and permissions. Normal and optimized runs, including actual extraction and rebuilding of the ZIP, are compared by complete archive bytes and member hashes. Byte identity is a same-toolchain claim; different TeX, font, Python, or numerical-library versions can change output bytes.

[write] **The companion here, 6 October 2026.** Report 228’s companion is shipped as `code/228-logheight-*.py`, `data/228-logheight-*`, `228-logheight-code-README.md` and `228-logheight-sources-SOURCE_LEDGER.md`; its `supporting/` copies are Parts I and II. The delivered `code/reproduce.py--check` regenerated both result files byte for byte on a copy at the write; the report README gives the rerun route.

36 Further questions and limits

36.1 Polynomially small ratios

The explicit β^{-30} loss identifies an obstacle to much smaller ratios. The present theorem is organized around fixed logarithmic bands and the height window $m \asymp n \log n$. A broad transfer theory for polynomially small β , especially one approaching the fixed-degree polynomial regime, is not proved here. A narrow range may be tested against the displayed absolute error, but that observation alone does not supply a sharp uniform transition theorem, optimal range, or a high-height error matching Kaneiwa’s polynomial expansion. A useful next step is to exploit lower derivative orders or the retained first correction in the absolute Fourier norm, with explicitly balanced endpoint errors. Such a result would need its own statement and proof rather than a claim of arbitrary-small- β uniformity.

[write] **Still open, 6 October 2026.** Section 37, item 7.

36.2 Effective bounds and validated evaluation

Our remainder constants and onset are qualitative. A numerical certificate would require explicit bounds for the sectorial coordinate remainder, its inverse identification, contour tails, the finite-height derivative errors, and floating evaluation of the chosen model. Higher numerical precision or repeated mesh agreement cannot replace these bounds. An effective counterpart of (III.49) could produce rigorously enclosed height intervals for target ratios. Determining a least height would additionally require suitable discrete information about $R(n, m)$; no monotonicity result is available in this report.

[write] **Still open, 6 October 2026.** Section 37, items 1 and 3.

36.3 Stronger endpoint structure

The smooth germ g has an all-fixed-order Taylor asymptotic expansion on a sector, but analyticity at zero and convergence of its Taylor series have not been shown. Classical resurgence theory suggests studying sharper coefficient growth and summability questions, with attention to the nonzero resiter. These would concern the universal map, not a new Euler-specific density. Likewise a complete analytic comparison with the public IBD contour amplitude would clarify the global contour normalization beyond the local coordinate identity recorded above.

[write] **Still open; shared with the iterated Bell report, 6 October 2026.** Section 37, items 6 and 9.

36.4 Other arrays and other height windows

The constants T and B arise from the subsidiary Euler forcing. For another multiset recurrence, or another initial function, the universal endpoint calculation may persist while the clock and transfer change. Extending this method requires proving the new forced entrance and angular-tail estimates. At the opposite endpoint $\beta \rightarrow \infty$, a different density regime is needed. Finally, every fixed inverse-logarithm order here coexists with only one polynomial-in-height transfer rate; an all-orders $1/m$ theorem would require a further hierarchy of quantitative forced corrections and is not a consequence of the present proof.

[write] **Still open, 6 October 2026.** Section 37, items 2, 5 and 8.

37 Further questions and research (all three Parts)

[write] **Added in the write, 6 October 2026.** This section gathers the open questions of the three Parts under Vladimir’s standing rule of 4 October 2026: every claim that a manuscript states without proof, every limitation it records as a question, and the question shared with the iterated Bell report. Each item names its sources, the sketch they give, and what is missing. Questions answered inside the merge are marked at the questions themselves. No claim of the three manuscripts was found to be wrong, and none is refuted here.

1. *Validated constants.* Sources: Part I, Subsection 11.1; Part II, Subsection 23.2; Part III, Subsection 36.2. The amplitude c has the absolutely convergent formula (6); the first correction needs $h(1)$, $h'(1)$, $h''(1)$, since κ is now in closed form (II.3). Sketch: interval enclosures of the truncated Fatou coordinate on a stated sector (Lemma 26.1 gives the rational coefficients to every order), of the inverse branch, of the forward propagation of values and derivatives, of the finite quadrature and of the vertical tail. Missing: explicit numerical values for every constant of the sector proofs. Current values ($c \approx 4.4923299$, $h(1) \approx 1.81460956$) are floating diagnostics only.
2. *Corrections beyond the first, and an all-orders theorem.* Sources: Part I, Subsection 11.2; Part II, Subsection 23.1; Part III, Subsection 36.4 (an all-orders theorem in $1/m$). The expected form is $1 + \sum_r P_r(\log n)/n^r$ with $\deg P_r \leq 2r$; Part II proves P_1 . Part II’s conditional inference, that $[X^{2r}]P_r = (-1/18)^r/r!$ (so $\log^4 n/(648n^2)$ at second order) *if* an all-orders construction gives the degree bound and no other contribution reaches degree $2r$, is unproved. Missing: an induction over subsidiary orders, clock hierarchies with resonant logarithms (Part II’s N already has $\frac{1}{540} \log y$), forced Fatou cohomological corrections, and growing-strip errors with an extraction order chosen per precision.
3. *Effective constants, onsets and certified thresholds.* Sources: Part I, Subsection 11.3; Part II, Subsection 23.2; Part III, Subsection 36.2. All three Parts give qualitative constants only; their inverses recover n at exact sequence values eventually, without onset, and Part III’s prescriptions are smooth. Missing: explicit clock, stopping and sector constants; for least heights in Part III, discrete information (monotonicity) about $R(n, m)$, which no Part supplies.
4. *Inverse statements for $A(n, \lfloor \lambda n \rfloor)$.* Source: Part I, Subsection 11.3, second paragraph. The floor phase $e^{-\{\lambda n\}/\lambda}$ of (32) must enter; a smooth approximation that drops it has a non-vanishing multiplicative error. Not treated by any Part (Part III’s inverses concern $m \asymp n \log n$).
5. *The regime $n/m \rightarrow \infty$, and corrections near $n/m \rightarrow 0$.* Sources: Part I, Subsection 11.4; Part II, Subsection 23.3; Part III, Subsection 36.4. The side $n/m \rightarrow 0$ is answered at leading order for logarithmic bands by Part III (Theorem 31.1); the large-argument behaviour of h and any transfer as $\beta \rightarrow \infty$ are open, and no Part proves a first correction of Part II’s kind on a shrinking band.
6. *Analytic structure of h .* Sources: Part I, Subsection 11.4 (“stronger regularity, such as real analyticity”); Part III, Subsection 36.3 (analyticity of the germ g at zero, convergence of its Taylor series, coefficient growth and summability, with the nonzero resiter $1/3$). Known: $h \in C^\infty(0, \infty)$ (Corollary 6.2); h continues holomorphically to a small punctured sector about $(0, \epsilon)$, so h is real analytic on $(0, \epsilon)$ (Theorem 28.1); g has a one-sided C^∞ germ (Theorem 29.1). Missing: real analyticity on all of $(0, \infty)$ (the measure identity (28) is the suggested route; item 9 would give it), and any statement on the Taylor series of g .
7. *Polynomially small ratios and Kaneiwa’s regime.* Sources: Part III, Subsection 36.1; Part I, Subsection 11.4, second paragraph; Part II, Subsection 23.3, second paragraph. Part III’s transfer loses β^{-30} and stops at logarithmic bands; a uniform theory linking proportional height, the

crossover and Kaneiwa's fixed-degree polynomials (Section 33) is open. Part III's suggested step: lower derivative orders or the retained first correction in the absolute Fourier norm, with balanced endpoint errors.

8. *Other arrays and initial functions.* Sources: Part I, Subsection 11.4; Part II, Subsection 23.3; Part III, Subsection 36.4. The forcing and clock constants (T, B, κ) must be recomputed for another multiset recurrence or initial function; the endpoint calculus of the unforced map may persist. Missing: the forced entrance and tail estimates in each new case.
9. *The amplitude identity with the iterated Bell report* (a question shared with `a139383-iterated-bell-diagonals`). Source: Part III, Section 25 ("a formal contour match suggests an amplitude relation $I(\beta) = \beta 2^{\beta/3} h(\beta)$, but that identification requires its own analytic contour correspondence and is not used here") and Subsection 36.3 ("a complete analytic comparison with the public IBD contour amplitude"). Here I is the iterated Bell report's Part II amplitude, $H(n, m) \sim (n-1)! 2^{-n} (\lambda n)^n (\lambda n)^{-1/(3\lambda)} I(1/\lambda)$ for $m = \lambda n$ (its `ibd:pd:eq:factorial`), defined there by a fixed-radius contour functional. Sketch: both reports use the same normalized reciprocal orbit, and $\Psi_{\text{IBD}} = -\Phi + \frac{1}{3} \log 2$ holds locally; the iterated Bell report's own literature section derives $I(\beta) = \beta J_P(\beta)$ from Prellberg's contour integral under the same kind of unproved contour correspondence. Evidence: agreement to eight significant digits at $\beta = 1$ (Remark 37.1). Missing: an analytic identification of that report's contour functional with the Laplace inversion of P (orientation, normalization, vanishing endpoint terms). Consequences if proved: (a) h is real analytic on $(0, \infty)$, because that report proves I real analytic there (item 6); (b) Part III's endpoint results ($h(0) = 2$, Corollary 29.2, (III.29)) give the behaviour of $I(\beta)$ as $\beta \rightarrow 0$, that is, the amplitude half of the endpoint $\lambda \rightarrow \infty$ that the iterated Bell report leaves open; a labelled analogue of Theorem 31.1 would still be needed for a coefficient theorem (Remark 37.2).

Remark 37.1 ([**write**] The identity at $\beta = 1$, numerically, 6 October 2026). The iterated Bell report's shipped exploratory runs (`data/53-iterated-bell-exploratory-amplitude.json` there) give $I(1) = 2.2862647982, 2.2862647847, 2.2862647863$ (its $C = \sqrt{2\pi} I(1)/2$ between 2.865407976 and 2.865407993). Part I's shipped diagnostic $h(1) = 1.8146095596784502$ (`data/225-leading-data-amplitude_diagnostic.json`) gives $2^{1/3} h(1) = 2.2862647816$. The two sides agree to eight significant digits (2.2862648); their differences, 3×10^{-9} to 1.7×10^{-8} , are of the size of the spread of the three iterated Bell runs (1.4×10^{-8}). (Added after the independent check of 6 October 2026: an independent evaluation of (6) by the verification (four values of σ , Gauss–Legendre quadrature to $|u| = 4 \cdot 10^4$ with an integration-by-parts tail; uncertified) gives $h(1) = 1.814609559835$ and $2^{1/3} h(1) = 2.2862647818$, so the shipped diagnostic is low by 1.6×10^{-10} , from its truncation of the vertical tail at $|u| = 3000$; the comparison is unchanged, with differences 2.9×10^{-9} to 1.65×10^{-8} and agreement to eight significant digits.) Both sides are floating values without error control; this is evidence for item 9, not a proof of it, and it says nothing about $\beta \neq 1$.

Remark 37.2 ([**write**] A conditional consequence for the labelled array, 6 October 2026). Let $H(n, m) = n! [z^n] (e^z - 1)^{\circ m}$ and $R_{\text{lab}}(n, m) = H(n, m) / (n! m^{n-1} 2^{1-n})$. Suppose, for some $\beta_0 > 0$ and $D > 1$, (a) $I(\beta) = \beta 2^{\beta/3} h(\beta)$ for $0 < \beta \leq \beta_0$, and (b) $H(n, m) = (n-1)! 2^{-n} m^n m^{-\beta/3} I(\beta) \{1 + \epsilon(n, m)\}$ uniformly for $(\log m)^{-D} \leq \beta = n/m \leq \beta_0$, with $\epsilon \rightarrow 0$ (the iterated Bell report's leading form `ibd:pd:eq:factorial` with $\lambda n = m$, extended to logarithmic bands). If $m/(n \log n) \rightarrow \lambda \in (0, \infty)$, then $R_{\text{lab}}(n, m) \rightarrow e^{-1/(3\lambda)}$, the same limit as Part III's Theorem 24.1; and if $\epsilon = O(L^{-N-1})$ with $L = \log n$ and $m = \lfloor \lambda n L \rfloor$, then

$$\log R_{\text{lab}}(n, m) = -\frac{1}{3\lambda} + \frac{K_{\text{lab}}}{\lambda L} + \sum_{j=2}^N \frac{c_j}{(\lambda L)^j} + O(L^{-N-1}), \quad K_{\text{lab}} = \frac{1-\gamma}{3},$$

with the same c_j as (III.6).

Proof. By (a) and (b), since $(n-1)!m^n\beta = n!m^{n-1}$,

$$R_{\text{lab}}(n, m) = 2^{\beta/3} m^{-\beta/3} \frac{h(\beta)}{2} \{1 + \epsilon(n, m)\}.$$

By (III.29), $\log(h(\beta)/2) = -\frac{\beta}{3} \log \beta + \frac{C_0}{3} \beta + \sum_{j=2}^N c_j \beta^j + O(\beta^{N+1})$ with $C_0 = 1 - \gamma - \log 2$. Hence, using $\log \beta + \log m = \log n$ exactly as in (III.36),

$$\log R_{\text{lab}} = -\frac{\beta}{3} \log n + \frac{\log 2 + C_0}{3} \beta + \sum_{j=2}^N c_j \beta^j + O(\beta^{N+1}) + \log(1 + \epsilon),$$

and $\log 2 + C_0 = 1 - \gamma$. If $m/(n \log n) \rightarrow \lambda$, then $\beta \rightarrow 0$, $\beta \log n \rightarrow 1/\lambda$, and β eventually lies in the band of (b): $\beta \leq \beta_0$, and $\beta \geq 1/(2\lambda \log n) \geq (\log m)^{-D}$ because $D > 1$ and $\log m \sim \log n$. So the right side tends to $-1/(3\lambda)$. For $m = \lfloor \lambda n L \rfloor$, $\beta = 1/(\lambda L) + O(1/(nL^2))$, whose effect is smaller than every power of L , as in the proof of Theorem 24.1; substituting gives the display. $\square$

Hypothesis (b) is not proved anywhere: the iterated Bell report's Theorem (its `ibd:pd:thm:main`) is uniform only for $\lambda = m/n$ in compact subsets of $(0, \infty)$, and it records the endpoints $\lambda \rightarrow 0, \infty$ as open. A consistency check, not a proof: with $h(0) = 2$ the right side of (b) is $n!m^{n-1}2^{1-n}(1 + o(1))$ when $\beta \log m \rightarrow 0$, the leading term of the fixed- n asymptotic $H(n, r+1) \sim n!r^{n-1}/2^{n-1}$ of Skau and Kristensen that the iterated Bell report cites. Exact values of $H(n, m)$, computed by the write from $H(n, m+1) = \sum_j S(n, j)H(j, m)$ with Stirling numbers $S(n, j)$ and evaluated in 30-digit floating arithmetic (uncertified), behave as the remark predicts, with the heights $m = \lfloor \lambda n \log n \rfloor$ of Part III's Table 1:

n	λ	m	$R_{\text{lab}}(n, m)$	$e^{-1/(3\lambda)}$	$e^{-1/(3\lambda)}(1 + K_{\text{lab}}/(\lambda \log n))$
20	0.5	29	0.56666331	0.51341712	0.56172244
20	1	59	0.75860855	0.71653131	0.75023906
20	1.5	89	0.83316752	0.80073740	0.82585011
40	0.5	73	0.55659049	0.51341712	0.55264578
40	1	147	0.74875697	0.71653131	0.74390532
40	1.5	221	0.82522543	0.80073740	0.82113138
80	0.5	175	0.54955621	0.51341712	0.54644061
80	1	350	0.74202367	0.71653131	0.73957531
80	1.5	525	0.81979276	0.80073740	0.81790547

At $\lambda = 1$ the distance to the limit is 0.042, 0.032, 0.025 at $n = 20, 40, 80$, and after the K_{lab} term 0.0084, 0.0049, 0.0024. The Euler constant $K \approx 0.816$ of Part III and $K_{\text{lab}} \approx 0.141$ differ, as the subsidiary forcing leads one to expect. These finite values neither prove the limit nor certify any digit.

[write] **Independent check of the write, 6 October 2026.** An adversarial check made by the intake after the write (commit `84fc1aead`) re-pulled A290354 (live revision #26 of 21 June 2018) and A290353 (#28) and found every quoted line character for character, including the formula line in the front matter, in the note at the start of Part I and in the bibliography. The entry's history places the conjecture in Kotěšovec's revision #23 of 14 August 2017, so the date "Aug 14 2017" that the write gives is the date of the conjecture, not of the revision read, and it is correct. The check recomputed a_n as the n -th Euler transform of $1 + x$ in integer arithmetic (equal to the b-file for $n \leq 40$; $a_{40}(T/40)^{40}40^{4/3} = 4.6013347$), confirmed that Theorem 1.1 is exactly the conjectured form, and the rows $A(2, m)$, $A(3, m)$, $A(4, m)$ for $m \leq 12$. It

re-derived Remark 37.2 from (a), (b) and (III.29), found it labelled conditional wherever it appears (title, hypotheses, the paragraph after the proof, the README) and not listed as proved in the status table, and reproduced its table to every printed digit from an own exact computation of $H(n, m)$; extended to $n = 160$ it gives $m = 812$, $R_{\text{lab}} = 0.73782182$ at $\lambda = 1$ (distance 0.0213 to the limit, 0.0014 after the K_{lab} term) and $m = 406$, $R_{\text{lab}} = 0.54374893$ at $\lambda = 0.5$ (0.0303 and 0.0018). It evaluated the closed form (II.3) to 35 digits, $\kappa = -0.25434816985107480385506489650373339$, and matched it against an independent quadrature of the defining equation (II.23) to 5×10^{-28} . It recomputed $h(1)$ independently (Remark 37.1, dated note) and confirmed the classification of the inverses in the front matter: the factorial core and the Lambert core with $a = b = 1$ are instances; the corrections, the rounding at exact values and Part III's reversion and prescriptions are not. Two wordings are corrected, each with a dated note keeping the first wording: the front matter's justification of the printed restatements (twice) overstated Part II's novelty, since its bound on inverse orbits is in Part I's proof of Lemma 4.1; and the note after Part II's Section 13 now flags the neighbourhood form of the outer bound, as it already flagged the disk form of the tail bound, as proved only inside Part I's proofs. Added: the name $L = \log(Y/T)$ in the front matter, and the independent values of $h(1)$ and $c = 4.4923298971$. No mathematical error was found.

[write] **Part IV and these questions, 9 October 2026.** Part IV (Section 38) answers part of these questions; the earlier notes, written before it arrived, are kept. *Item 9* (the amplitude identity): **answered**, conditionally on Part IV's four inputs (Section 40): $I(t) = t^{2^{t/3}}h(t)$ for every $t > 0$ (Theorem 39.1). The proof identifies the limits of the positive measures $\mu_m = \sum_n [z^n] f^{\text{om}}(z) P(-m)^n \delta_{n/m}$, whose masses are exactly $P(0)$; it needs no global comparison of contours, the route item 9 calls missing. Both consequences listed there follow: (a) h is real analytic on $(0, \infty)$ (below); (b) the endpoint behaviour of I as $\beta \rightarrow 0$ is Corollary 42.2, the amplitude half of the iterated Bell report's endpoint $\lambda \rightarrow \infty$; a uniform coefficient transfer toward that endpoint stays open (Part IV's Question 1). *Item 6*: **answered on the positive axis**: h extends holomorphically to $\Re t > 0$ (39.6). Analyticity of the germ g at zero and its Taylor series stay open (Part IV's Question 7). *Remark 37.2*: its hypothesis (a) is now Theorem 39.1 (for all $\beta > 0$, under the same inputs); hypothesis (b) is still unproved, so the remark remains conditional. *Remark 37.1*: the identity now forces $I(1) = 2^{1/3}h(1)$. With Part IV's exploratory density ($h(1) = 1.81460955983\dots$, within 10^{-12} of the verification's independent 1.814609559835) the forced value is 2.2862647817...; the iterated Bell report's three exploratory runs exceed it by 2.9×10^{-9} , 4.5×10^{-9} and 1.6×10^{-8} . Every value involved is an uncertified floating evaluation, and the spread of the three runs is 1.4×10^{-8} , so this records an offset of the runs, not a refutation of anything; the remark's "eight significant digits" stays true. *Items 1 and 5* reappear as Part IV's Questions 5 (enclosures of $h(\log 2)$, $h'(\log 2)$, $h''(\log 2)$) and 6 (large-argument asymptotics of h). *An observation* [write], uncertified: from exact chain counts (the Newton form $\sum_m e^{-Lm} H(n, m) = \sum_k (N^k)_{n1} e^{-Lk} (1 - e^{-L})^{-k-1}$, with N the strictly triangular part of the Stirling matrix) and Part IV's exploratory h , the ratio of $S_n(L)$ to the main term of Theorem 39.2 satisfies $n(\text{ratio} - 1) = 0.11556$ at $L = \log 2$ and 0.16669 at $L = 1$ for $n = 320$ (monotone in n from $n = 25$), against $L/6 = 0.11552$ and 0.16667: the first correction looks like $L/(6n)$ with no logarithmic term, as the chain report proves at $q = 1$ ($z_n/f_n = C(1 + L/(6n) + O(n^{-2}))$). At $L = 1$ ($q = 1/(e - 1)$, outside that report's disk) this is evidence on Part IV's Question 2, not a proof.

Part IV

A common density for parabolic iteration

Manuscript 02 of the archive `ProveIt_Research_2026-10-08 (1).zip`, delivered as `02_parabolic_amplitudes/article.tex` (16 pages). Delivered title block “A Common Density for Parabolic Iteration / Exact Bell–Euler amplitude identification and weighted partition-chain constants / Research companion prepared for Vladimir Reshetnikov / Developed with OpenAI ChatGPT / 8 October 2026”. Printed as delivered after Section 38, which the write added, with **[write]** notes; the manuscript’s Section k is Section $k + 38$ here, its statements and equations $(k.m)$ are $(k+38.m)$, and its Appendix A keeps its letter. Its labels carry the prefix `ied:par:.` “[12]” in this Part is Parts I–III of this report, “[5]” the iterated Bell report, “[13]” the strict partition-chain report.

Abstract of this Part (as delivered). Two recent ProveIt reports study different arrays governed by the same parabolic map $f(z) = e^z - 1$. One constructs a positive density h whose Laplace transform is the normalized entire Fatou orbit P ; the other obtains an iterated-Bell amplitude I from a fixed contour. Their precise relation was left as an explicit open question. We prove

$$I(t) = t 2^{t/3} h(t) \quad (t > 0)$$

by identifying two limits of positive coefficient measures. This avoids a global comparison of the two contours and proves that h extends holomorphically to $\Re t > 0$. A coefficientwise rational majorant then controls every iteration depth in a geometric aggregate. It gives the exact Lengyel-constant identity and, more generally, a holomorphic formula on $\Re q > 0$ for the leading amplitude of strict partition chains marked by q . Complete proofs, source pins, exact finite checks and independent numerical diagnostics are supplied. The analytic inputs from the earlier reports are identified explicitly; no worldwide priority claim or new leading-equivalent claim is made.

38 Provenance, inputs and notation of Part IV

[Added 9 October 2026.] Part IV was added in a second write of this report, after Parts I–III had been merged (`84fc1aaed`) and independently checked (`a3a2d5582`). It is the manuscript *A Common Density for Parabolic Iteration*, which starts from item 9 of Section 37 (it cites it as “Section 37, item 9”) and answers it. Everything of the manuscript is printed in this Part except its title page (recorded in the source block above) and its reference list (merged into this report’s bibliography): its abstract above, its ten sections as Sections 39–48 and its Appendix A. Its letters are kept (Section 38.6). This section and the notes marked **[write]** are the write’s; all other text of Sections 39–48 and of the appendix is the manuscript’s.

38.1 The manuscript

The archive `ProveIt_Research_2026-10-08 (1).zip` (1,394,668 bytes; 45 files in one wrapper directory `ProveIt_Research_2026-10-08/`, 1,636,954 bytes unpacked) arrived in commit `b28d0850b` (8 October 2026, 08:39 PDT). It holds three manuscripts; this is its manuscript 02, `02_parabolic_amplitudes/` (12 files). Its 9 supporting files were placed in this directory by commit `803f4d937` (9 October 2026) with the prefix `04-parabolic-`, the Part number: the three programs as `code/04-parabolic-*.py`, the four data files, `provenance.json` and `requirements.txt` as `data/04-parabolic-*`; all 9 are byte-identical to the delivery (confirmed at the write). Not

shipped, and retrievable from the arrival commit: the manuscript `article.tex`, its PDF and its `README.md`. The archive is not retired: its manuscripts 01 and 03 and its package-level files (among them `REVIEW_NOTES.md`, `THEOREM_LEDGER.json`, `RESEARCH_QUESTIONS.json`, `reproduce.py`) belong to their own placements.

The manuscript is `article.tex` (1,079 lines, dated 8 October 2026); a build of the delivered text gives 16 pages, as delivered, with no warnings. Its title page reads “Research companion prepared for Vladimir Reshetnikov / Developed with OpenAI ChatGPT” (PDF author field the same). Its pin (Section 47.2, `data/04-parabolic-provenance.json` and its three bibliography entries) is ProveIt commit 58175ca45563d9ee29875374dd77942f069268a9 (7 October 2026, 20:30 PDT), with the blobs `b5af1ca30b` of this report’s `article.tex` (Parts I–III as merged and checked, unchanged until this write), `2b1d8fbe81` of the iterated Bell report and `da8cac4780` of the strict partition-chain report, both unchanged at HEAD.

How the manuscript was merged. Its 56 labels, and 6 labels the write gave to its unlabelled Sections 1, 4, 7, 9, 10 and Appendix A, carry the prefix `ied:par::`; its 51 references (41 `\eqref`, 10 `\ref`) were updated. Its citation keys are entries of the merged bibliography, marked [Part IV], except two: `BellReport` is IBD (4 citations re-keyed; Part IV’s pin and description added) and `Prellberg` is the entry of Parts I–III (Part IV’s description added); its key `Euler` is this report at the pin, that is, Parts I–III, and its `OEIS` (A005121, listed but not cited) is A005121. Its macros `\coef` and `\Repin` were added to the preamble; its `\C`, `\R`, `\N`, `\dd`, `\ii`, `\file`, `\Log` are those of Parts I–III; its `\Oh` (a calligraphic $\mathcal{O}$, an upright O in Parts I–III) is switched at the start of this Part. From this Part on, equations are numbered within sections, as delivered. The delivered font (Palatino), running heads and paragraph skip are not reproduced. No other text was changed. The write labelled the nine items of its Section 48. The numbering, against a build of the delivered text: its Sections 1–10 are Sections 39–48 here, its Theorems 1.1 and 1.2 are Theorems 39.1 and 39.2, Corollary 1.3 is Corollary 39.3, Corollary 4.2 is Corollary 42.2, Proposition 4.3 is Proposition 42.3, Corollary 7.1 is Corollary 45.1; every statement and equation $k.m$ is $(k+38).m$; Appendix A is unchanged (all 56 labels compared).

What the manuscript says about the repository. Its account of the three reports is correct: item 9 (“Section 37, item 9”) asks for the identity; the iterated Bell report records the same question (its dated note at the end of `ibd:sec:literature`); the labels it names as resolved are `ied:q:ibd`, the positive-axis part of `ied:q:analytic`, the leading-amplitude part of `spc:q:depth`, and in part `spc:q:disk`, as the table of Section 38.3 states. The chain report’s `spc:q:depth` carries the heuristic $C \stackrel{?}{=} L^{L/3-1} I(L)/2$ of that report’s write, with $L = \log 2$; (39.12) proves it.

38.2 The trust boundary: four inputs

Every theorem of this Part is an implication. Its main theorem uses exactly the four statements of Section 40, each proved in an unrefereed, unformalized report of this collection:

- *Input F :* the normalized Fatou coordinate and the entire orbit, Part I’s Lemma 4.1 and (20);
- *Input M :* the positive Laplace measure with continuous, strictly positive density, Part I’s Proposition 6.1;
- *Input L :* the absolute local coefficient transfer `ibd:pd:eq:absolute` of the iterated Bell report, uniform for n/m in compact subsets of $(0, \infty)$; that report’s Part II is itself conditional on its Part I, whose outer arcs rest on an interval certificate of common entry (Remark 40.1 says so);
- *Input A :* the holomorphy part of `ibd:pd:prop:positive` there (not its numerical positivity seed $I(1) > 1.64$).

The corollaries use more: Corollary 42.2 uses Part III’s (III.29); Corollary 39.3’s agreement near $q = 1$ and Corollary 45.1 use the strict partition-chain report’s marked expansion and its cumulant expansions (45.1)–(45.2). The manuscript states this boundary itself (“The present proof checks the implication from their stated analytic interfaces”), and its package record (`REVIEW_NOTES.md`, not shipped here) lists among its remaining trust boundaries that “The parabolic source reports are unrefereed” and that its inverse-Laplace calculations “are not interval-certified”. Nothing here makes the inputs more certain than their sources: the theorems of this Part have exactly the status of Parts I–III and of the iterated Bell report together. Prellberg’s 2002 seminar, in Mishna’s summary [9], is credited for the bivariate parabolic mechanism and an aggregate contour constant; no new-equivalent or priority claim is made.

38.3 What Part IV answers

Question	After Part IV
Item 9 (the amplitude identity)	Answered (Theorem 39.1), by limits of positive coefficient measures, without a comparison of contours.
Item 6 (analytic structure of h)	Answered on the positive axis: h holomorphic on $\Re t > 0$ (39.6). Open: the germ g at zero and its Taylor series.
Remark 37.2	Hypothesis (a) proved; (b) open; the remark stays conditional.
Remark 37.1	$I(1) = 2^{1/3}h(1)$ is now forced; the iterated Bell report’s uncertified runs sit 2.9×10^{-9} to 1.6×10^{-8} above the forced value (a record, not a refutation).
Iterated Bell report, endpoint $\lambda \rightarrow \infty$	The amplitude half: Corollary 42.2; the uniform transfer stays open (Question 1).
Chain report, <code>spc:q:depth</code>	The leading-amplitude part: its heuristic $C = L^{L/3-1}I(L)/2$ is (39.12), a theorem; all-order summation and the logarithmic cancellation stay open (Question 2).
Chain report, <code>spc:q:disk</code>	In part: $C(q)$ holomorphic on $\Re q > 0$, positive on $q > 0$; the zero set and global uniform asymptotics stay open (Questions 3, 4).
Items 1–8 otherwise	Untouched; items 1 and 5 reappear as Questions 5 and 6.

Dated notes at the end of Section 37 and of the front matter say the same; the notes for the two neighbouring reports are proposed separately and not applied.

What Part IV proves again: nothing of Parts I–III; its Lemma 41.1 re-derives the radius from Input F, and its Lemma 44.1 the chain report’s exact mixture $z_n = \sum_m 2^{-m-1}H(n, m)$ at $q = 1$. *What is new to the report:* everything else.

38.4 What the write read and checked

At placement (the intake dossier, 9 October 2026; own code): from the delivered exploratory $h(\log 2)$, $C(1) = (2L)^{L/3}h(L)/2 = 1.09868580552518740\dots$ against Lengyel’s constant, OEIS A086053, 1.09868580552518701\dots (agreement to 15 digits); $I(\log 2) = 1.65770061282$, consistent with the chain report’s finite- n extrapolations; the exact counts against A005121 and A139383; and $2^{1/3}h(1) = 2.28626478178$ against the iterated Bell report’s runs.

At the write (9 October 2026; Python 3.14.4, SymPy 1.14.0, mpmath 1.3.0; own code, kept with the intake records and not shipped). Every proof of Sections 41–45 was read line by line, with no

error found; and the four inputs were read at their sources (Part I's Lemma 4.1 and Proposition 6.1; `ibd:pd:eq:absolute`, stated there for $\eta = n/m \in [b, B]$ with remainder $\mathcal{O}(m^{-1}(1 + \log m)^D)$, and the holomorphy part of `ibd:pd:prop:positive`), where they say what Section 40 quotes. Rechecked by hand or with SymPy in particular: the three sensitive normalizations after the proof of Theorem 39.1; the main term of Theorem 39.2 from $(n-1)!2^{-n}(n/L)^{-L/3}I(L)$ times $n!L^{-n-1}$; (39.12) and (39.11); the endpoint expansion (42.4) and its exponentiation (42.5); the chain rule behind Corollary 45.1 and both expanded forms; the radius model of Lemma 41.1 (numerically, residual $O(\log m/m)$ at $m = 10^3, 10^6$). With exact integer and rational arithmetic: the Newton form $H(n, m) = \sum_k \binom{m}{k} (N^k)_{n1}$ against the recurrence (40.2); the chain counts against A005121; the majorant (43.2) for $n \leq 30$, $m \leq 60$; the mixture (44.3) for $n \leq 25$ at five rational q , through the closed form $\sum_m a^m H(n, m) = \sum_k (N^k)_{n1} a^k (1-a)^{-k-1}$. Numerically (uncertified): Theorem 39.2 at $L = \log 2$ and $L = 1$ for $n \leq 320$, with ratios to the main term 1.00036 and 1.00052 at $n = 320$ (note at the end of Section 37); the iterated Bell report's three runs against the forced $I(1)$. It did not rerun the shipped programs. *Not read by the write*: Prellberg's summary, Lengyel (1984) and Skau–Kristensen; the manuscript's accounts of them are reported, not confirmed.

38.5 What is not claimed

From the manuscript: the theorems are implications from the four inputs, whose sources “are unrefereed and not proof-assistant verified”, and the proof “does not promote their earlier proofs or interval software to formal verification”; no numerical seed is needed for $I(1)$, but other computational dependencies of the Bell foundation remain (Remark 40.1); the leading equivalent and the existence of Lengyel's constant are classical, and Prellberg's seminar is a direct antecedent; the mean, variance and central limit theorem of chain lengths are not new claims; the continuation of $C(q)$ does not give uniform complex asymptotics or zero-freeness; the endpoint series does not give the uniform transfer, and the germ is not shown convergent; Proposition 42.3 is a transfer principle under stated hypotheses; the inverse-Laplace values are diagnostics, not certified; “A literature search and source inspection do not establish that no equivalent theorem has appeared elsewhere”. The write adds: its checks are exact finite computations, hand or SymPy algebra and floating evaluations; its observation on the first correction of the aggregate is numerical; nothing was submitted anywhere, and the notes for the two neighbouring reports are not applied.

38.6 Reading conventions

Part IV keeps its letters. Where they clash with Parts I–III, read them as follows.

Symbol	Part IV	Parts I–III
$H(n, m), f$	$n![z^n]f^{\circ m}$, the labelled iterated Bell array; $f(z) = e^z - 1$	$H(z)$ the matching function, H_j , $H_{\theta,a}$; $f(w) = e^w - 1$
$I(t), h, P, \Phi$	the iterated Bell amplitude; the density; the orbit; the coordinate (Parts I–III's)	I (Section 25); I a compact interval (II)
t, β	$t = n/m$	$\beta = n/m$
$L, L(q)$	$L > 0$ the tilt of $S_n(L)$; $L(q) = \log(1 + 1/q)$; $L = \log 2$ at $q = 1$	$L(y)$ a clock, $L = \log n$ (III)
$C, C(q), K_1, K_2, K$	Lengyel's constant, the marked amplitude; cumulant constants; $K = \max(D, 2)$	c the amplitude; C, K constants of Parts II–III

Symbol	Part IV	Parts I–III
$S_n(L), Z_n(q),$ $T_n, \mathcal{H}_n$	the geometric aggregate; marked chain counts; $T_n = \sum_m a^m H(n, m)$; the number of strict transitions	$T = \pi/2$
r_m, u_m, μ_m, s_m	$P(-m); 2/r_m$; the coefficient measures; $m + \Phi(r_m e^{s/m})$	r, r_m radii; s a contour parameter
$A(x), a, \rho$	$\log h(x) + (x/3) \log(2x)$; $a = q/(q+1)$ or the a of Proposition 42.3; ρ the log coefficient there	$A(n, m)$ the Euler array; $a_n; \rho_j$
$\mathcal{W}_n(L), w_n$ c_j, D	the central depth window; $\sqrt{n}(\log n)^2$ Part III's log-density coefficients; the logarithmic degree of Input L	$c_j; D$ a band exponent (III)

39 Question, result and scope

Let $f(z) = e^z - 1$, and let f^{om} denote m iterations of f , with $f^{o0}(z) = z$. Define

$$H(n, m) = n![z^n]f^{om}(z), \quad n \geq 1, \quad m \geq 0. \quad (39.1)$$

The array H is the iterated exponential Bell array. The principal objects here are functions that describe its asymptotic coefficients, rather than a new interpretation of these integers.

The report on iterated Euler transforms [12] constructs an entire function P satisfying

$$P(s+1) = e^{P(s)} - 1, \quad P(s) = \int_0^\infty e^{st} h(t) dt. \quad (39.2)$$

Its additive normalization is fixed by the inverse coordinate

$$\Phi(w) = -\frac{2}{w} + \frac{1}{3} \log w + \mathcal{O}(w), \quad \Phi(P(s)) = s \quad (39.3)$$

near the positive real w -axis, equivalently for sufficiently negative real s . The density h is continuous and strictly positive on $(0, \infty)$, and the measure in (39.2) has every exponential moment.

The iterated-Bell report [5] defines a second function I by a fixed-radius contour and proves the local coefficient formula

$$H(n, m) \sim (n-1)! 2^{-n} m^n m^{-n/(3m)} I(n/m) \quad (39.4)$$

uniformly when n/m ranges over compact subsets of $(0, \infty)$. Section 37, item 9, of [12] explicitly asks whether

$$I(t) = t 2^{t/3} h(t). \quad (39.5)$$

The same question is recorded in the iterated-Bell report. Both sources observe agreement of numerical values, but leave the orientation, normalization and endpoint terms of a proposed contour comparison unresolved.

Theorem 39.1 (Amplitude identification). *Under the four analytic input statements recorded in Section 40, identity (39.5) holds for every real $t > 0$. The density therefore has the unique holomorphic extension*

$$h(z) = \frac{2^{-z/3}}{z} I(z), \quad \Re z > 0. \quad (39.6)$$

In particular, h is real analytic on all of $(0, \infty)$.

The formulation records ordinary mathematical dependencies: the four statements are proved in the cited repository reports. Those sources are unrefereed and not proof-assistant verified. The present proof checks the implication from their stated analytic interfaces; it does not promote their earlier proofs or interval software to formal verification.

Our second main result concerns the geometric aggregate

$$S_n(L) = \sum_{m \geq 0} e^{-Lm} H(n, m), \quad L > 0.$$

The sum converges for every n . The following theorem justifies the missing depth summation and identifies its normalization.

Theorem 39.2 (Geometric aggregation). *For every $L > 0$,*

$$S_n(L) \sim (n!)^2 (2L)^{-n} n^{-1-L/3} L^{L/3-1} I(L). \quad (39.7)$$

The equivalent is uniform for L in any fixed compact subinterval of $(0, \infty)$.

For strict partition chains, put

$$Z_1(q) = 1, \quad Z_n(q) = q \sum_{j=1}^{n-1} \left\{ \begin{matrix} n \\ j \end{matrix} \right\} Z_j(q) \quad (n \geq 2),$$

where the braces denote Stirling numbers of the second kind. Thus q marks the number of strict transitions, not the number of partitions in the chain. Write

$$L(q) = \log(1 + 1/q) \quad (q > 0). \quad (39.8)$$

Corollary 39.3 (Marked chain amplitude). *For every $q > 0$,*

$$Z_n(q) \sim C(q) (n!)^2 [2L(q)]^{-n} n^{-1-L(q)/3}, \quad (39.9)$$

uniformly on compact positive q -intervals, with

$$C(q) = \frac{L(q)^{L(q)/3-1} I(L(q))}{q+1} \quad (39.10)$$

$$= \frac{[2L(q)]^{L(q)/3} h(L(q))}{q+1}. \quad (39.11)$$

This $C(q)$ is positive for real $q > 0$ and extends holomorphically to the half-plane $\Re q > 0$.

At $q = 1$, $Z_n(1)$ is OEIS A005121, whose normalized leading constant is Lengyel's constant C . Thus

$$C = \frac{1}{2} (\log 2)^{(\log 2)/3-1} I(\log 2) = \frac{1}{2} (2 \log 2)^{(\log 2)/3} h(\log 2). \quad (39.12)$$

The leading equivalent and the existence of this constant are classical [14, 9]; our contribution is the rigorous identification with the precise functions and normalization used by the current repository reports.

39.1 What changes in the research record

Question or statement	Status in this companion
Euler–Bell amplitude identity, [12], item 9	Proved by positive coefficient measures.
Real analyticity of h on $(0, \infty)$	Proved; holomorphic extension to $\Re z > 0$.
Strict-chain amplitude from the depth theorem	Proved with a uniform bound for every depth.
Marked amplitude beyond a neighborhood of $q = 1$	Holomorphic on $\Re q > 0$; positive on the positive axis.
Bell proportional-depth leading form	An input from [5], with older antecedents credited.
All fixed-order marked-chain expansions near $q = 1$	Prior repository result [13]; not reproved here.
Complex zero-freeness and global uniform complex expansions	Remain open; amplitude continuation alone does not imply them.

40 Four analytic inputs and exact conventions

We separate the new argument from the analytic foundations it uses. This permits an audit without reconstructing the long Euler-transform clock estimates or the iterated-Bell contour certificate.

Input F: normalized Fatou coordinate and orbit

Lemma `ied:lead:lem:fatou` and equation `ied:lead:eq:fatou` of [12] give an analytic Φ on a sufficiently small sector about the positive real axis with (39.3). Its local inverse P_{loc} extends to an entire function P , agrees with that inverse in its sector, and satisfies the functional equation in (39.2). In particular, for all sufficiently large integers m ,

$$r_m = P(-m) > 0, \quad \Phi(r_m) = -m.$$

Only this inverse statement near the negative real s -axis and the entire functional equation will be needed.

Input M: positive Laplace measure

Proposition `ied:lead:prop:measure` of [12] proves that P is the Laplace transform of a finite positive measure on $[0, \infty)$. It has no atom at zero, every exponential moment, and a continuous strictly positive density h on $(0, \infty)$. The identification needs only finiteness, positivity, the Laplace formula and continuity. Strict positivity is used for later relative aggregation. Smoothness, proved in the source’s Corollary `ied:lead:cor:smooth`, is not needed as an input.

Input L: absolute local coefficient transfer

Equation `ibd:pd:eq:absolute` of [5] states that, for each fixed $0 < a < b < \infty$, there is a finite D such that

$$\frac{H(n, m)}{(n-1)! 2^{-n} m^n m^{-n/(3m)}} = I(n/m) + \mathcal{O}\left(\frac{(1 + \log m)^D}{m}\right) \quad (40.1)$$

uniformly for integers $n, m \geq 1$ with $a \leq n/m \leq b$. Here $m \rightarrow \infty$, and the implicit constant may depend on a, b . This is an *absolute* estimate; it appears before the source's positivity proof and involves no division by I .

Input A: holomorphy of the contour amplitude

The first part of the proof of Proposition `ibd:pd:prop:positive` of [5] shows that I is holomorphic on $\Re t > 0$. It uses a complex-linear contour functional and locally uniform domination. The later numerical seed for strict positivity is not needed for holomorphy, and is not used here.

Remark 40.1 (Dependency and numerical boundaries). The amplitude identification gives a route from positivity of h to positivity of I without a numerical enclosure for $I(1)$. This does not erase other computational dependencies of the earlier Bell foundation. That report uses an interval certificate for common entry on a fixed semicircle. We import its resulting absolute transfer statement rather than independently formalizing that certificate.

40.1 Nonnegative coefficients

Every coefficient of f is nonnegative and $f(0) = 0$, so every coefficient of $f^{\circ m}$ is nonnegative. The exact recurrence is

$$H(n, m+1) = \sum_{j=1}^n \left\{ \begin{matrix} n \\ j \end{matrix} \right\} H(j, m), \quad H(n, 0) = \mathbf{1}_{\{n=1\}}. \quad (40.2)$$

Indeed, iterates commute, and

$$\frac{f(z)^j}{j!} = \sum_{n \geq j} \left\{ \begin{matrix} n \\ j \end{matrix} \right\} \frac{z^n}{n!}.$$

These facts ensure positivity of the measures below. The use of $(n-1)!$ rather than $n!$ in (40.1) is essential: after coefficient extraction it supplies $1/n$, ultimately producing the factor t in (39.5).

41 Positive coefficient measures and their limit

Lemma 41.1 (The exact radius and its expansion). *Let $r_m = P(-m)$ for sufficiently large integers m . Then*

$$\frac{2}{r_m} = m - \frac{1}{3} \log m + \frac{1}{3} \log 2 + \mathcal{O}\left(\frac{\log m}{m}\right), \quad (41.1)$$

$$\log \frac{mr_m}{2} = \frac{\log m - \log 2}{3m} + \mathcal{O}\left(\frac{(\log m)^2}{m^2}\right). \quad (41.2)$$

Proof. Put $u_m = 2/r_m$. Since $\Phi(r_m) = -m$ and $\log r_m = \log 2 - \log u_m$, Input F gives

$$u_m + \frac{1}{3} \log u_m = m + \frac{1}{3} \log 2 + \mathcal{O}(1/u_m).$$

It first follows that $u_m/m \rightarrow 1$: the logarithmic term is $o(u_m)$, and the identity forces $u_m \rightarrow \infty$. Consequently $u_m = m + \mathcal{O}(\log m)$ and $\log u_m = \log m + \mathcal{O}(\log m/m)$. Substitution proves (41.1). Taylor expansion of $-\log(1 + (u_m - m)/m)$ proves (41.2). $\square$

For every sufficiently large integer m , define

$$\mu_m = \sum_{n \geq 1} ([z^n] f^{\circ m}(z)) r_m^n \delta_{n/m}. \quad (41.3)$$

This is a finite positive measure because $f^{\circ m}$ is entire. The functional equation gives its exact normalization:

$$\mu_m([0, \infty)) = f^{\circ m}(r_m) = f^{\circ m}(P(-m)) = P(0). \quad (41.4)$$

Thus no asymptotic renormalization of these measures is needed.

Lemma 41.2 (Laplace convergence). *For each fixed real s ,*

$$\int_0^\infty e^{st} d\mu_m(t) = f^{\circ m}(r_m e^{s/m}) \longrightarrow P(s). \quad (41.5)$$

The convergence of the right-hand side is locally uniform for complex s .

Proof. For s in a fixed compact subset of $\mathbb{C}$ and m large, $z_m = r_m e^{s/m}$ lies in the local inverse sector. Set $s_m = m + \Phi(z_m)$. The local identity $P(\Phi(z_m)) = z_m$ and iteration of (39.2) give exactly

$$f^{\circ m}(z_m) = P(s_m).$$

Subtracting the two coordinate expansions and writing $u_m = 2/r_m$,

$$\begin{aligned} s_m &= \Phi(r_m e^{s/m}) - \Phi(r_m) \\ &= u_m(1 - e^{-s/m}) + \frac{s}{3m} + \mathcal{O}(1/m) \\ &= s + \mathcal{O}\left(\frac{1 + \log m}{m}\right). \end{aligned}$$

The estimates are uniform on the chosen compact set. The last line uses $u_m = m + \mathcal{O}(\log m)$. Entire analyticity of P proves convergence. The Laplace equality follows from the positive power series in (41.3). $\square$

Lemma 41.3 (Compact test convergence). *For every continuous function φ with compact support in $(0, \infty)$,*

$$\int \varphi(t) d\mu_m(t) \longrightarrow \int_0^\infty \varphi(t) h(t) dt. \quad (41.6)$$

Proof. Push μ_m and $h(t) dt$ forward under $x = e^{-t}$ to finite measures on $[0, 1]$, giving them zero mass at $x = 0$. Their total masses are $P(0)$. For every integer $k \geq 0$, Lemma 41.2 at $s = -k$ gives convergence of their k th moments. Hence their integrals converge for each polynomial.

Uniform polynomial approximation on $[0, 1]$ gives convergence for every continuous function there: an approximation error at most ϵ contributes at most $2P(0)\epsilon$, uniformly in m . The function

$$g(x) = \varphi(-\log x) \quad (0 < x \leq 1), \quad g(0) = 0$$

is continuous on $[0, 1]$ because φ has compact support inside $(0, \infty)$. Applying the conclusion to g proves (41.6). $\square$

Remark 41.4. A compact asymptotic formula for individual coefficients alone need not identify a global transform limit. Here the masses are finite and fixed, and their Laplace transforms determine polynomial moments after compactification. The same measures can then be tested in the local coefficient regime. No positive-measure interpretation of the signed Bell contour functional is required.

42 Identification, analyticity and endpoint consequences

Proof of Theorem 39.1. Fix $0 < a < b < \infty$ and put $t = n/m$. Dividing (40.1) by $n!$ and multiplying by r_m^n gives, uniformly for $a \leq t \leq b$,

$$([z^n]f^{\circ m}(z))r_m^n = \frac{1}{n} \left(\frac{mr_m}{2} \right)^n m^{-t/3} \left[I(t) + \mathcal{O}\left(\frac{(1 + \log m)^D}{m} \right) \right]. \quad (42.1)$$

By Lemma 41.1,

$$n \log \frac{mr_m}{2} - \frac{t}{3} \log m = -\frac{t}{3} \log 2 + \mathcal{O}\left(\frac{(1 + \log m)^2}{m} \right).$$

Therefore

$$([z^n]f^{\circ m}(z))r_m^n = \frac{1}{m} \left[\frac{2^{-t/3} I(t)}{t} + \mathcal{O}\left(\frac{(1 + \log m)^K}{m} \right) \right], \quad (42.2)$$

where $K = \max\{D, 2\}$ suffices. Only boundedness of I on $[a, b]$ and $t \geq a$ have been used, not positivity of I .

Let φ have compact support in (a, b) . Summing (42.2) against $\varphi(n/m)$ is a Riemann sum. There are $\mathcal{O}(m)$ relevant atoms, so their accumulated error is $\mathcal{O}((1 + \log m)^K/m) = o(1)$. Thus

$$\int \varphi d\mu_m \longrightarrow \int_a^b \varphi(t) \frac{2^{-t/3} I(t)}{t} dt.$$

Lemma 41.3 gives another expression for the same limit. The continuous function $h(t) - 2^{-t/3} I(t)/t$ has zero integral against every such test function and therefore vanishes pointwise. Arbitrariness of a, b proves (39.5) on $(0, \infty)$.

Input A makes the right-hand side of (39.6) holomorphic on $\Re z > 0$. It agrees with h on the positive axis. Uniqueness follows from the identity theorem. $\square$

There are three sensitive normalizations. The denominator in (41.1) is $m - \frac{1}{3} \log m + \frac{1}{3} \log 2$. The factor $1/n$ in (42.1) comes from $(n-1)!/n!$. The exponential tilt cancels $m^{-t/3}$ but leaves $2^{-t/3}$. Their combination gives exactly the stated identity.

Equality was established on each positive interval by test functions. It was not inferred from matching finitely many endpoint coefficients or decimal values. The proof also never takes $n/m \rightarrow 0$ in a theorem whose hypotheses require a fixed positive compact ratio range.

Corollary 42.1 (Analytic positivity). *$I(t) > 0$ for every $t > 0$.*

Proof. Input M gives $h(t) > 0$, and (39.5) has a positive multiplier. $\square$

42.1 The small-parameter amplitude

The Euler report supplies a separate endpoint input: for every fixed $N \geq 2$,

$$\log \frac{h(t)}{2} = -\frac{t}{3} \log t + \frac{1 - \gamma - \log 2}{3} t + \sum_{j=2}^N c_j t^j + \mathcal{O}(t^{N+1}), \quad t \downarrow 0, \quad (42.3)$$

with fixed differentiated remainders and $c_2 = (9 - \pi^2)/108$. This is equation `ied:log:eq:logdensity` and its coefficient discussion in [12]; it is not required for the main theorem.

Corollary 42.2 (Endpoint transseries for I). *For every fixed $N \geq 2$,*

$$\log \frac{I(t)}{2t} = -\frac{t}{3} \log t + \frac{1-\gamma}{3}t + \sum_{j=2}^N c_j t^j + \mathcal{O}(t^{N+1}). \quad (42.4)$$

The fixed differentiated remainder statements transfer unchanged. In particular,

$$\begin{aligned} I(t) = 2t & \left\{ 1 + \frac{t}{3}(1 - \gamma - \log t) \right. \\ & \left. + t^2 \left[\frac{(1 - \gamma - \log t)^2}{18} + \frac{9 - \pi^2}{108} \right] + \mathcal{O}(t^3(1 + |\log t|)^3) \right\}. \end{aligned} \quad (42.5)$$

Proof. Take logarithms of (39.5) and substitute (42.3). The $\log 2$ term cancels exactly. Exponentiation through degree two gives (42.5). The logarithms differ by an exactly linear function, so the differentiated remainders introduce no new error. $\square$

This solves the amplitude part of the Bell large-depth endpoint question. Coefficient asymptotics when depths grow like $n \log n$ still need a uniform transfer as n/m decreases. Neither the endpoint series nor holomorphic continuation supplies that uniformity. The smooth renormalized germ at zero is not shown here to have a convergent Taylor series.

42.2 A reusable identification principle

Proposition 42.3 (General positive-measure matching). *Suppose a positive-coefficient iterated family has an entire orbit P and local inverse coordinate*

$$\Phi(w) = -\frac{1}{aw} + \rho \log w + \mathcal{O}(w), \quad a > 0, \quad \rho \in \mathbb{R},$$

with the analogues of Inputs F and M . Suppose its absolute local transfer for $t = n/m$ in positive compact sets is

$$[z^n] f^{\circ m}(z) = \frac{1}{n} (am)^n m^{-\rho t} (J(t) + o(1)),$$

where J is continuous and the error uniform. Then

$$J(t) = t a^{-\rho t} h(t) \quad (t > 0).$$

Proof. Set $r_m = P(-m)$ and $u_m = 1/(ar_m)$. The inverse identity yields $u_m = m - \rho \log m - \rho \log a + o(1)$, and therefore

$$m^{-\rho t} (amr_m)^n \longrightarrow a^{\rho t}$$

uniformly on positive compact ratio ranges. The Laplace convergence of the coefficient measures follows as before, since $u_m(1 - e^{-s/m}) \rightarrow s$. Their local density is $a^{\rho t} J(t)/t$, which compact test convergence identifies with h . $\square$

This is a transfer principle with stated hypotheses, not a claim that every parabolic map has a positive Laplace orbit or a local coefficient theorem. The exponential map has $a = 1/2$ and $\rho = 1/3$.

43 A bound valid for every iteration depth

For power series with real coefficients, write $A \preceq B$ when every coefficient of $B - A$ is nonnegative. If the series have nonnegative coefficients and zero constant term, this order is preserved by composition.

Lemma 43.1 (Rational composition majorant). *For every integer $m \geq 0$,*

$$f^{\circ m}(z) \preceq \frac{z}{1 - mz/2}. \quad (43.1)$$

In particular, for $m, n \geq 1$,

$$0 \leq H(n, m) \leq n! \left(\frac{m}{2}\right)^{n-1}. \quad (43.2)$$

Proof. For every $j \geq 1$, $j! \geq 2^{j-1}$: each factor $2, \dots, j$ is at least 2. Hence

$$f(z) = \sum_{j \geq 1} \frac{z^j}{j!} \preceq g(z) := \frac{z}{1 - z/2}.$$

Composition and induction give $f^{\circ m} \preceq g^{\circ m}$. Direct substitution shows $g^{\circ m}(z) = z/(1 - mz/2)$. Extract coefficients and multiply by $n!$. $\square$

This elementary inequality supplies what compact-ratio asymptotics omit: a bound at depths much smaller or larger than n . Near the main saddle it loses a factor of order $n^{L/3}$, but exponential tail decay dominates that polynomial loss. We give the needed summation estimates explicitly.

Lemma 43.2 (Discrete gamma sums). *Let L range over a fixed compact interval $[L_0, L_1] \subset (0, \infty)$. Uniformly as $n \rightarrow \infty$,*

$$\sum_{m \geq 1} e^{-Lm} m^n = \frac{\Gamma(n+1)}{L^{n+1}} (1 + \mathcal{O}(n^{-1/2})). \quad (43.3)$$

Put $w_n = \sqrt{n}(\log n)^2$. For $p = n$ or $p = n - 1$, there are constants $c, C > 0$ such that

$$\sum_{\substack{m \geq 1 \\ |m - n/L| > w_n}} e^{-Lm} m^p \leq C \frac{\Gamma(p+1)}{L^{p+1}} e^{-c(\log n)^4} \quad (43.4)$$

for all sufficiently large n , uniformly in L .

Proof. For a nonnegative unimodal integrable function v on $[0, \infty)$, comparison with unit intervals on each side of its maximum gives

$$\left| \sum_{m \geq 1} v(m) - \int_0^\infty v(x) dx \right| \leq 2 \sup_{x \geq 0} v(x).$$

Take $v(x) = e^{-Lx} x^p$. Its integral is $\Gamma(p+1)L^{-p-1}$ and its maximum is $(p/(eL))^p$. Stirling's formula shows that their ratio is $\mathcal{O}(p^{-1/2})$ uniformly on the stated L -interval. This proves (43.3) and bounds the complete sum by a uniform constant times its integral.

For the upper tail, set $M = n/L + w_n$ and $\theta = L^2 w_n / (4n)$. For large n , $L \pm \theta$ lie in a fixed positive compact interval. Exponential tilting gives

$$\begin{aligned} \sum_{m \geq M} e^{-Lm} m^p &\leq e^{-\theta M} \sum_{m \geq 1} e^{-(L-\theta)m} m^p \\ &\leq CT(p+1)L^{-p-1} e^{-\theta M} \left(\frac{L}{L-\theta} \right)^{p+1}. \end{aligned}$$

Let $x = \theta/L$. Because $p+1 = n + \mathcal{O}(1)$ and $-\log(1-x) \leq x + x^2$ for $0 \leq x \leq 1/2$, the logarithm of the last factor is at most

$$-\theta w_n + \mathcal{O}(\theta) + (n + \mathcal{O}(1))\theta^2/L^2.$$

Its principal terms are $-L^2 w_n^2 / (4n) + L^2 w_n^2 / (16n)$. Thus it is bounded above by $-c w_n^2 / n$ uniformly for large n .

For the lower tail, when $m \leq n/L - w_n$,

$$e^{-Lm} \leq e^{\theta(n/L - w_n)} e^{-(L+\theta)m}.$$

The same complete-sum bound leads to the logarithm

$$\theta(n/L - w_n) - (p+1) \log(1 + \theta/L).$$

Using $\log(1+x) \geq x - x^2/2$ gives the same negative bound. Finally $w_n^2/n = (\log n)^4$. Rounding the real cutoffs to integers only introduces factors $e^{\mathcal{O}(\theta)}$, which do not affect the estimates. $\square$

44 Summing the Bell local limit

Proof of Theorem 39.2. Fix a compact positive L -interval and use

$$\mathcal{W}_n(L) = \{m \in \mathbb{N} : |m - n/L| \leq w_n\}, \quad w_n = \sqrt{n}(\log n)^2.$$

Throughout this window, uniformly,

$$\frac{n}{m} = L + \mathcal{O}\left(\frac{(\log n)^2}{\sqrt{n}}\right), \quad \log m = \log(n/L) + \mathcal{O}\left(\frac{(\log n)^2}{\sqrt{n}}\right).$$

By the amplitude identification, I is continuous and strictly positive. It has a positive lower bound on the compact range of these ratios. Therefore

$$m^{-n/(3m)} I(n/m) = (n/L)^{-L/3} I(L)(1 + o(1)) \quad (44.1)$$

uniformly on the window and in L . In particular, the difference between the logarithm of the power on the left and its central value is $\mathcal{O}((\log n)^3/\sqrt{n}) = o(1)$. Its dependence on n does not obstruct this replacement.

Input L now applies within one fixed compact ratio range and gives

$$\begin{aligned} \sum_{m \in \mathcal{W}_n(L)} e^{-Lm} H(n, m) &= (n-1)! 2^{-n} (n/L)^{-L/3} I(L)(1 + o(1)) \\ &\quad \times \sum_{m \in \mathcal{W}_n(L)} e^{-Lm} m^n. \end{aligned} \quad (44.2)$$

The absolute transfer error is a relative $o(1)$ here because $m \asymp n$ and I has a positive lower bound. Lemma 43.2 shows that the last sum equals $n!L^{-n-1}(1+o(1))$. Substitution gives the main term in (39.7).

For the omitted depths, both preceding lemmas give

$$\begin{aligned} \sum_{m \notin \mathcal{W}_n(L)} e^{-Lm} H(n, m) &\leq n! 2^{1-n} \sum_{m \notin \mathcal{W}_n(L)} e^{-Lm} m^{n-1} \\ &\leq Cn! 2^{1-n} (n-1)! L^{-n} e^{-c(\log n)^4}. \end{aligned}$$

Dividing by the main scale gives at most a uniform constant times $n^{L_1/3} e^{-c(\log n)^4} \rightarrow 0$. The positive amplitude $L^{L/3-1} I(L)$ is bounded away from zero on the fixed interval. The $m=0$ term is zero for $n > 1$. All estimates were uniform, proving the theorem. $\square$

44.1 An exact mixture for strict chains

Lemma 44.1. *For $q > 0$ and $a = q/(q+1)$, for every $n \geq 1$,*

$$Z_n(q) = (1-a) \sum_{m \geq 0} a^m H(n, m) = \frac{1}{q+1} S_n(L(q)). \quad (44.3)$$

Proof. The majorant proves absolute convergence. Put $T_n = \sum_{m \geq 0} a^m H(n, m)$ and sum (40.2):

$$T_n = \mathbf{1}_{\{n=1\}} + a \sum_{j=1}^n \left\{ \begin{matrix} n \\ j \end{matrix} \right\} T_j.$$

The diagonal Stirling number equals 1, so $T_1 = (1-a)^{-1}$ and

$$T_n = \frac{a}{1-a} \sum_{j < n} \left\{ \begin{matrix} n \\ j \end{matrix} \right\} T_j \quad (n \geq 2).$$

Multiplying by $1-a$ and using $a/(1-a) = q$ gives the defining recurrence and initial value of $Z_n(q)$. Uniqueness by induction proves the result. $\square$

Proof of Corollary 39.3. Apply the aggregation theorem to (44.3) and then the amplitude identification. This proves (39.9), (39.10) and (39.11), their positivity and compact real uniformity.

For the complex continuation define $L(q) = \text{Log}(1+1/q)$ on $\Re q > 0$, using the principal logarithm. There $\Re(1+1/q) > 1$, so L is holomorphic and $\Re L(q) = \log|1+1/q| > 0$. In particular L is nonzero and $\text{Log } L$ is holomorphic as well. The function

$$\frac{1}{q+1} \exp \left[\left(\frac{L(q)}{3} - 1 \right) \text{Log } L(q) \right] I(L(q))$$

is holomorphic throughout $\Re q > 0$ and agrees with the identified positive-real amplitude. On a common neighborhood of $q = 1$ it agrees, by the identity theorem, with the prior local marked amplitude constructed in [13]. $\square$

The continuation concerns the amplitude. A uniform expansion for $Z_n(q)$ on a complex region additionally needs control of cancellation and of its remainders; it does not follow from amplitude continuation.

45 Consequences for chain-length constants

The prior local marked-chain expansion [13] gives, with $L = \log 2$ and $\mathcal{H}_n$ the number of strict transitions in a uniformly chosen chain,

$$\mathbb{E} \mathcal{H}_n = \frac{n}{2L} + \frac{\log n}{6} + K_1 - \frac{1}{12n} + \mathcal{O}(n^{-2}), \quad (45.1)$$

$$\text{Var}(\mathcal{H}_n) = \frac{1-L}{4L^2}n - \frac{\log n}{12} + K_2 + \frac{1}{24n} + \mathcal{O}(n^{-2}). \quad (45.2)$$

That source defines the constant terms by

$$K_1 = \left. \frac{d}{dt} \log C(e^t) \right|_{t=0}, \quad K_2 = \left. \frac{d^2}{dt^2} \log C(e^t) \right|_{t=0}.$$

The new identity expresses these in the common density and its derivatives.

Corollary 45.1 (Density formulas for constant cumulant terms). *Put $A(x) = \log h(x) + (x/3) \log(2x)$ for $x > 0$. Then*

$$K_1 = -\frac{1}{2} - \frac{1}{2}A'(L), \quad K_2 = -\frac{1}{4} + \frac{1}{4}(A''(L) + A'(L)). \quad (45.3)$$

Equivalently,

$$K_1 = -\frac{1}{2} - \frac{1}{2} \left[\frac{h'(L)}{h(L)} + \frac{\log(2L) + 1}{3} \right], \quad (45.4)$$

$$K_2 = -\frac{1}{4} + \frac{1}{4} \left[\frac{h''(L)}{h(L)} - \left(\frac{h'(L)}{h(L)} \right)^2 + \frac{h'(L)}{h(L)} + \frac{1}{3L} + \frac{\log(2L) + 1}{3} \right]. \quad (45.5)$$

Proof. Near $t = 0$, (39.11) gives

$$\log C(e^t) = -\log(1 + e^t) + A(\log(1 + e^{-t})).$$

The inner function has first derivative $-1/2$ and second derivative $1/4$ at zero; the first two derivatives of $-\log(1 + e^t)$ there are $-1/2$ and $-1/4$. The chain rule yields (45.3). Differentiating A gives the expanded expressions. $\square$

The mean, variance and central limit theorem are not new claims of this note. These formulas identify their amplitude derivatives in the normalization used by the three reports. Counting partitions instead of strict transitions would add one to the mean and to K_1 ; we retain the source's convention.

46 Computation and reproducibility

No analytic proof above uses a measured constant or a finite experiment. The accompanying programs check exact normalizations and compare independent numerical constructions.

46.1 Exact finite checks

The script `code/verify_exact.py` uses exact integer and rational arithmetic for:

- 960 instances of (43.2), for $1 \leq n \leq 24$, $1 \leq m \leq 40$;
- 144 evaluations of the depth polynomial in the Newton basis;
- 96 rational instances of the geometric mixture, at $q = 1/2, 1, 2, 3/2$;
- a symbolic check that the four-term Fatou coordinate has zero Abel defect through degree five.

All checks passed. The Newton-basis check uses that the Stirling matrix on degrees at most n is $I + N$ with N strictly triangular:

$$H(n, m) = \sum_{k=0}^{n-1} \binom{m}{k} (N^k)_{n1}.$$

The coefficient $(N^k)_{n1}$ counts strict chains of length k , independently checking the convention in (44.3). These verifications concern finite algebra, not bounds for asymptotic remainders.

46.2 Independent Laplace inversion

The script `code/explore_density.py` approximates $P(-p)$ directly. It solves a truncated inverse-coordinate equation far along the negative direction and iterates $w \mapsto e^w - 1$ forward. The four coefficients are

$$\Phi(w) = -\frac{2}{w} + \frac{1}{3} \log w - \frac{w}{36} + \frac{w^2}{540} + \frac{w^3}{7776} - \frac{71w^4}{435456} + \mathcal{O}(w^5).$$

The exact script checks those coefficients symbolically. Numerical inverse Laplace transformation then estimates $h(t)$. This route neither uses the Bell fixed-contour amplitude nor estimates the density from Bell counts.

Two runs used 160 and 320 forward steps and inversion degrees 48 and 64. Their values agree to at least twelve decimal places at the four displayed arguments. Because no interval bound for truncation or quadrature is supplied, that agreement is a diagnostic, not a certified precision claim.

t	Exploratory $h(t)$	$t2^{t/3}h(t)$
1/2	2.14193747174	1.20212176094
$\log 2$	2.03764226105	1.65770061282
1	1.81460955983	2.28626478178
2	1.02595377154	3.25720019242

The script `code/compare_counts.py` computes Bell counts by exact Stirling recurrence through degree 160 and depth 320. It compares normalized counts at $n/m = 1/2, 1, 2$ with the independently inverted density. It also computes strict-chain counts through $n = 320$ and compares their normalized constant, after the prior report's four known corrections, with (39.12). The complete outputs are in `data/count_comparison.json`. Finite agreement is never used as a proof premise.

46.3 Replay

From the companion directory run:

```
python code/verify_exact.py --output data/exact_checks.json
python code/explore_density.py --steps 160 --degree 48 \
    --output data/density_160.json
python code/explore_density.py --steps 320 --degree 64 \
    --output data/density_320.json
python code/compare_counts.py --density data/density_320.json \
    --output data/count_comparison.json
pdflatex -halt-on-error article.tex
pdflatex -halt-on-error article.tex
```

Counting uses the Python standard library; the symbolic coordinate check uses SymPy, and the optional decimal calculations use mpmath. No script accesses the network. Outputs distinguish exact checks from floating diagnostics.

[write] **The package here, 9 October 2026.** In this report the three programs are `code/04-parabolic-verify_exact.py`, `code/04-parabolic-explore_density.py` and `code/04-parabolic-compare_count_s.py`, and the data files named above are `data/04-parabolic-*`; they read and write the delivered names, so the replay runs only on a copy with the delivered layout (the README gives the recipe). The write did not rerun them. Its own exact checks of the majorant, the mixture and the Newton basis, and its numerical check of the aggregate, are recorded in Section 38.4; the table’s $t^{2^{t/3}}h(t)$ column is the inverse-Laplace h times an exact factor, so its digits are as uncertified as those of h .

47 Literature, provenance and research boundaries

47.1 Prior analytic work

Prellberg’s 2002 seminar, summarized by Mishna in 2005 [9], applies parabolic iteration and complex analysis to linear composition recurrences. Its general setting is

$$X(z) = a(z)X(f(z)) + b(z), \quad f(z) = z + cz^2 + dz^3 + \dots$$

The summary contains a two-index coefficient expression before summing over depth and a final partition-chain equivalent with an inverse-coordinate contour amplitude. For $f = e^z - 1$, the quantity $1 - d/c^2$ is $1/3$. These are direct antecedents. The leading scale and the use of Fatou coordinates are not discoveries of this note.

The four-page primary seminar summary was opened and read for this work. The separate FPSAC slide URL cited by the Bell report could not be retrieved in this session and is not represented as independently read. The repository distinguishes its detailed compact-uniform transfer theorem from the older outline.

Lengyel’s 1984 recurrence paper is the historical source identified by the summary [14]. The fixed-degree, large-iteration leading coefficient of the Bell array is also discussed by Skau and Kristensen [15]; its abstract was inspected here. That information alone does not justify a stronger claim about the full scope of the paper. Our arguments do not use it as a theorem.

47.2 Repository pin and exact questions

The inspected ProveIt revision is

58175ca45563d9ee29875374dd77942f069268a9.

The source reports are under

SetTheory/Cardinals/docs/reports/
generating-functions-and-asymptotics/oeis-sequence-asymptotics/.

Their exact source identifiers are:

Report	Git blob of <code>article.tex</code>
a290354-iterated-euler-diagonals	b5af1ca30b21cd89d90488258d8722aa8a155b0e
a139383-iterated-bell-diagonals	2b1d8fbe811b11925a0879552b7319500f16cfd2
a005121-strict-partition-chains	da8cac4780378f0a0a1af35df4ffebdee4b014d2

The precise open-question labels resolved here are `ied:q:ibd`, the all-positive-axis portion of `ied:q:analytic`, and the leading-amplitude portion of `spc:q:depth`. The marked continuation also bears on `spc:q:disk`, but does not settle the complex zero set or global uniform asymptotic estimates. The mean-constant identities identify amplitude derivatives requested beside the depth question.

[write] **The pin and the labels, 9 October 2026.** At the pin the three blobs are as stated, and the reports’ texts at HEAD are the same, except that this report now has Part IV. The labels named in the preceding paragraph exist with the meanings given: `ied:q:ibd` and `ied:q:analytic` are items 9 and 6 of Section 37, and `spc:q:depth`, `spc:q:disk` are items of the strict partition-chain report’s “Further questions and research”. What this Part answers in each is stated in Section 38.3; “the leading-amplitude portion of `spc:q:depth`” is that report’s heuristic $C \stackrel{?}{=} L^{L/3-1}I(L)/2$, which (39.12) proves.

The new work proves connections between existing normalizations and supplies an explicit tail argument. Its repository-relative novelty is verifiable at the pin. A literature search and source inspection do not establish that no equivalent theorem has appeared elsewhere, and this note makes no such claim. The connection is useful because previously separate amplitude functions become one positive analytic density, allowing parameter transfer without guessing contour constants.

48 Further research questions

The following problems are concrete continuations of this work.

1. **Uniform Bell transfer toward the endpoint.** Prove a version of (40.1) for n/m decreasing on logarithmic and then polynomial scales. The endpoint amplitude is now known. The remaining task is a uniform coefficient remainder, especially for the long-delayed parts of the fixed cut.
2. **Summation of every correction.** The Bell expansion has polynomials in $\log n$ at each inverse power of n , whereas the chain recurrence expansion has no such logarithms. The factor $m^{-n/(3m)}$ slightly displaces the central depth saddle. Carry its expansion to every fixed order and prove cancellation of its logarithms against the Bell corrections. Theorem 39.2 proves only the leading summation.
3. **Complex zeros of the marked amplitude.** Determine whether (39.10) has zeros on $\Re q > 0$; equivalently, study zeros of I or h on the image of $L(q)$. Real positivity gives no complex zero-free theorem.
4. **A common positive-parameter asymptotic theory.** Extend the recurrence proof of every fixed order from a small disk about $q = 1$ to neighborhoods of every positive compact interval.

Compare its limiting constant directly with (39.11). Continuation of the amplitude alone is insufficient.

5. **Certified density and cumulant constants.** Give interval enclosures for $h(\log 2)$, $h'(\log 2)$ and $h''(\log 2)$ from the normalized orbit and explicit inversion tails. Equations (39.12), (45.4) and (45.5) then yield certified C, K_1, K_2 . The present decimal calculations stop before that certification.
6. **Large-argument density asymptotics.** Determine $h(t)$ as $t \rightarrow \infty$. This also determines the Bell amplitude at small proportional depth and, through $L(q) \rightarrow \infty$ as $q \downarrow 0$, the marked-chain amplitude at small transition weight. Compact-parameter results do not control this endpoint.
7. **Convergence or summability of the endpoint germ.** The renormalization $t^{t/3}h(t)$ has finite-order Taylor asymptotics. Decide convergence, or determine coefficient growth and summability. Holomorphy on $\Re t > 0$ does not decide behavior at its boundary point zero.
8. **Other positive parabolic maps.** Apply Proposition 42.3 to a family for which both the positive Laplace orbit and local coefficient theorem can be proved. The main difficulty is establishing those hypotheses, rather than identifying the measures once they hold.
9. **Formal verification of the bridge.** Formalize the implication from the four explicit analytic inputs: radius asymptotics, finite positive measures, polynomial approximation and Riemann sums. This gives a small auditable theorem boundary before the much larger earlier contour foundations are formalized.

[write] **Further questions of Part IV, 9 October 2026.** Following Vladimir Reshetnikov's standing rule of 4 October 2026, every claim of the manuscript that is not proved in full is stated as an open question; these nine items are the manuscript's own, as delivered, and the write labelled them. No claim of the manuscript was found wrong; nothing is moved or refuted. Relations to Section 37 and to the neighbouring reports: Question 1 is the coefficient half of the iterated Bell report's endpoint $\lambda \rightarrow \infty$ (the amplitude half is Corollary 42.2) and hypothesis (b) of Remark 37.2; Question 2 is the open part of the chain report's `spc:q:depth` (the write's numerical observation at the end of Section 37 suggests a first correction $L/(6n)$ without logarithm); Questions 3 and 4 are the open part of `spc:q:disk`; Question 5 contains item 1 at $\log 2$; Question 6 is the $\beta \rightarrow \infty$ half of item 5; Question 7 is the open part of item 6.

A A short proof checklist for independent review

A reviewer can assess the new reasoning by checking this chain:

1. $r_m = P(-m)$ is positive and gives the exact mass $f^{\circ m}(r_m) = P(0)$.
2. The additive constant of Φ in (39.3) is zero. It gives the $\log 2/3$ term in (41.1).
3. The inverse identity is applied only near r_m , where $r_m e^{s/m}$ stays in the stated sector.
4. Negative integer Laplace values and equal finite masses give compact test convergence after $x = e^{-t}$.
5. Input L is absolute and uniform on each fixed positive compact interval; such an interval contains $\mathcal{O}(m)$ atoms.
6. Continuity converts equality of the tested densities into a pointwise identity.
7. The central aggregation window lies in one compact ratio region; all outside depths obey the rational majorant.
8. The majorant's loss $n^{L/3}$ is retained before its tail is declared negligible.

9. For complex continuation, $L(q)$ stays in $\Re L > 0$ on $\Re q > 0$, with every logarithm branch specified. None of these steps uses the decimal values. The earlier analytic interfaces remain the inherited dependency.

References

- [1] The OEIS Foundation Inc., *A290353*, square array of iterated Euler transforms, entry by Alois P. Heinz, 28 July 2017. Retrieved 5 October 2026. <https://oeis.org/A290353>.
[As cited in Report 228: *A290353*, iterated Euler-transform array, entry by Alois P. Heinz, 28 July 2017. Source checked 5 October 2026. Report 226: Source checked for Report 225 on 5 October 2026.]
- [2] The OEIS Foundation Inc., *A290354*, entry by Alois P. Heinz, 28 July 2017; asymptotic conjecture by Václav Kotěšovec, 14 August 2017. Retrieved 5 October 2026. <https://oeis.org/A290354>.
[As cited in Report 226: leading asymptotic conjecture by Václav Kotěšovec, 14 August 2017. Source checked for Report 225 on 5 October 2026.]
- [3] R. Kaneiwa, *On r -fold Partitions and a Certain Form of Infinite Products*, Otaru University of Commerce, Humanities Research **61** (1980), 11–26. <https://barrel.repo.nii.ac.jp/records/2331>; persistent identifier <https://hdl.handle.net/10252/2056>.
[Report 228 adds: Propositions 1 and 2, printed p. 19.]
- [4] M. Bechtloff Weising, *Higher Order Bell Symmetric Functions*, arXiv:2505.18504v2. <https://arxiv.org/abs/2505.18504v2>. Definitions and statements checked in the versioned HTML on 5 October 2026.
- [5] *Asymptotics of Iterated Bell Diagonals (OEIS A139383, A261280)*, unrefereed research report in VladimirReshetnikov/ProveIt, combined fixed-shift and proportional-depth manuscripts, 2 October 2026. Catalogue checked 5 October 2026; indexed commit 65969409cfc007047182e9303e1d2d0106324dea. [Repository report](#).
[As cited in Report 228: unrefereed report in VladimirReshetnikov/ProveIt, 2 October 2026. Snapshot at main commit 5fdc0d68a062c815100ac27bd173a8c4b901c08d, checked 5 October 2026. [Pinned public source](#). Sections 4, 12, 13, and 17.] **[[write]]** In this repository: `SetTheory/Cardinals/docs/reports/generating-functions-and-asymptotics/oeis-sequence-asymptotics/a139383-iterated-bell-diagonals`; its `article.tex` is the same blob at both pins and at HEAD, and its Sections 4, 12, 13 and 17 are the sections Report 228 names.]
[Part IV (key `BellReport`): ProveIt research report *Iterated Bell diagonals*, with the proportional-depth addendum; pinned revision 58175ca45563d9ee29875374dd77942f069268a9, blob 2b1d8f8e81.]
- [6] *Diagonal and proportional height asymptotics for iterated Euler transforms*, Report 225, 5 October 2026. **[[write]]** Printed in this report as Part I.] As cited in Report 226: preceding unrefereed research report; unchanged PDF and editable TeX included in the archive’s `supporting` directory; exact provenance in `sources/SOURCE_LEDGER.md`. As cited in Report 228: unrefereed research report, 5 October 2026; unchanged complete TeX and PDF in the accompanying archive; provenance in `sources/SOURCE_LEDGER.md`. **[[write]]** The included copies are byte-identical to Part I and are not shipped; the ledgers are shipped as `226-firstcorr-sources-SOURCE_LEDGER.md` and `228-logheight-sources-SOURCE_LEDGER.md`.]
- [7] *First logarithmic correction for iterated Euler diagonals*, Report 226, unrefereed research report, 5 October 2026. Unchanged complete TeX and PDF in the accompanying archive; provenance in `sources/SOURCE_LEDGER.md`. **[[write]]** Printed in this report as Part II; the included copy is byte-identical to it and is not shipped.]
- [8] A. Dudko and D. Sauzin, *The resurgent character of the Fatou coordinates of a simple parabolic germ*, C. R. Acad. Sci. Paris I **352** (2014), 255–261. <https://doi.org/10.1016/j.crma.2013.12.005>.

- [9] T. Prellberg, *On the Asymptotic Analysis of a Class of Linear Recurrences*, September 2002 seminar, summary by M. Mishna, Algorithms Seminar 2002–2004 (2005), 47–50. Sections 2.2 and 3.1. <https://algo.inria.fr/seminars/summary/Prellberg2002a.pdf>.
[Part IV: seminar of 23 September 2002, summary by M. Mishna, in F. Chyzak (ed.), *Algorithms Seminar 2002–2004*, INRIA, 2005, pp. 47–50; “The four-page primary seminar summary was opened and read”.]
- [10] S. V. Nagaev and V. I. Vakhtel, *Limit theorems for probabilities of large deviations of a Galton–Watson process*, Discrete Math. Appl. **13**(1) (2003), 1–26. Theorems 1–2 and their Poisson specialization; the source-comparison limitations are recorded in the ledger. <https://doi.org/10.1515/156939203321669537>; <https://www.mathnet.ru/eng/dm183>.
- [11] [[write]] *Transseries and Inversion*, the transseries volume of this repository, Analysis/Transseries/docs/series-and-transseries/Transseries_And_Inversion/transseries_and_inversion.tex: Proposition p0:prop:factorial-core, Theorems p0:thm:lambert-core, p0:thm:perturbed-inversion and p0:thm:staircase.
- [12] [Part IV] ProveIt research report, *Diagonal and proportional height asymptotics for iterated Euler transforms*, with Parts II–III and editorial notes, a290354-iterated-euler-diagonals/article.tex. Pinned revision 58175ca45563d9ee29875374dd77942f069268a9 (blob b5af1ca30b). [This report’s Parts I–III, unchanged at the pin.]
- [13] [Part IV] ProveIt research report, *Strict partition chains and Lengyel numbers*, a005121-strict-partition-chains/article.tex. Pinned revision 58175ca45563d9ee29875374dd77942f069268a9 (blob da8cac4780).
- [14] [Part IV] T. Lengyel, *On a recurrence involving Stirling numbers*, European Journal of Combinatorics **5** (1984), 313–321. Bibliographic attribution checked in [9]; the original paper was not independently reread for the companion.
- [15] [Part IV] I. H. Skau and K. F. Kristensen, *An asymptotic formula for the iterated exponential Bell numbers*, arXiv:1903.07979, 2019. <https://arxiv.org/abs/1903.07979>.
- [16] [Part IV; listed, not cited in its text] The OEIS Foundation Inc., *A005121*, <https://oeis.org/A005121>. Accessed 8 October 2026. Sequence metadata are context; the checks independently recompute every count.